

# Supplement to Prediction of Infectious Disease Epidemics via Feature-Weighted Density Ensembles

*Evan L Ray, Krzysztof Sakrejda, Nicholas G Reich*

*March 2017*

In this supplement, we include additional figures and results.

## Component Model Log Scores and Weighting Features

Supplemental Figures 1, 2, and 3 illustrate the relationship between log scores and weighting features for predictions from the three component models made during the training phase in weeks before the season onset (for predictions of onset timing) or the season peak (for predictions of peak timing or peak incidence).

## Estimated Mean Model Performance

Supplemental Figure 4 displays estimated mean log scores for each model and ensemble method in predictions made before the season onset or season peak. Supplemental Figures 5, 6, and 7 show estimates for the difference in mean log scores between the model with the highest estimated mean log score for each target and each other model. The point and 95% interval estimates are obtained from a mixed effects model with a separate fixed effect mean for the interaction of model and prediction target; random effects for each combination of region, season, model, and prediction target; and lag 1 autocorrelation nested within each combination of region, season, model, and prediction target. For predictions of onset timing, the only difference in model performance that is statistically significant is the difference between **CW** and **KDE**. For predictions of peak timing, the only difference that is statistically significant is between **SARIMA** and **KDE**. For predictions of peak incidence, none of the differences between the models are statistically significant.




Figure 1: Log scores achieved by each component model in each week of the season, summarizing across all seasons in both the training phase when all three component models produced predictions. The thick line is a smoothed estimate of mean log score at each week in the season; the shaded region indicates the convex hull of log scores achieved by each model; and the actual log scores achieved in each week are indicated with points.




Figure 2: Log scores achieved by each component model vs{.} model uncertainty as measured by the number of bins required to cover 90% of the predictive distribution. The plot summarizes results across all seasons in the training phase when all three component models produced predictions. The thick line is a smoothed estimate of mean log score at each week in the season; the shaded region indicates the convex hull of log scores achieved by each model; and the actual log scores achieved in each week are indicated with points.




Figure 3: Log scores achieved by each component model vs. wLI in the week of the season when predictions were made. The plot summarizes results across all seasons in the training phase when all three component models produced predictions. The thick line is a smoothed estimate of mean log score at each week in the season; the shaded region indicates the convex hull of log scores achieved by each model; and the actual log scores achieved in each week are indicated with points.




Figure 4: Point estimates and confidence intervals for mean log score for each model in weeks before the target (onset or peak) occurred. Estimates are obtained from a mixed effects model with a separate fixed effect mean for the interaction of model and prediction target; random effects for each combination of region, season, model, and prediction target; and lag 1 autocorrelation nested within each combination of region, season, model, and prediction target. The wider confidence interval bounds are simultaneous confidence intervals with an approximate familywise 95% coverage rate for all intervals. The inner confidence intervals are calculated separately, with approximate coverage rates of 95%. Log scores of  $-\infty$  were truncated at -10 before fitting this model.




Figure 5: Point estimates and confidence intervals for the difference in mean log score between the best model and each other model for predictions of season onset timing made before the season onset occurred. Estimates are obtained from a mixed effects model with a separate fixed effect mean for the interaction of model and prediction target; random effects for each combination of region, season, model, and prediction target; and lag 1 autocorrelation nested within each combination of region, season, model, and prediction target. The wider confidence interval bounds are simultaneous confidence intervals with an approximate familywise 95% coverage rate for all individual model means and all pairwise model contrasts for all prediction targets. The inner confidence intervals are calculated separately, with approximate coverage rates of 95%. Log scores of -Infinity were truncated at -10 before fitting this model.




Figure 6: Point estimates and confidence intervals for the difference in mean log score between the best model and each other model for predictions of peak timing made before the season onset occurred. Estimates are obtained from a mixed effects model with a separate fixed effect mean for the interaction of model and prediction target; random effects for each combination of region, season, model, and prediction target; and lag 1 autocorrelation nested within each combination of region, season, model, and prediction target. The wider confidence interval bounds are simultaneous confidence intervals with an approximate familywise 95% coverage rate for all individual model means and all pairwise model contrasts for all prediction targets. The inner confidence intervals are calculated separately, with approximate coverage rates of 95%. Log scores of -Infinity were truncated at -10 before fitting this model.




Figure 7: Point estimates and confidence intervals for the difference in mean log score between the best model and each other model for predictions of peak incidence made before the season peak occurred. The wider confidence interval bounds are simultaneous confidence intervals with an approximate familywise 95% coverage rate for all individual model means and all pairwise model contrasts for all prediction targets. The inner confidence intervals are calculated separately, with approximate coverage rates of 95%. Log scores of -Infinity were truncated at -10 before fitting this model.