

Supporting Information for

**Slow particle remineralization, rather than suppressed
disaggregation, drives efficient flux transfer through the
Eastern Tropical North Pacific Oxygen Deficient Zone**

Jacob A. Cram¹, Clara A. Fuchsman¹, Megan E. Duffy², Jessica L. Pretty³, Rachel M.
Lekanoff³, Jacquelyn A Neibauer², Shirley W. Leung², Klaus B. Huebert¹, Thomas S.
Weber⁴, Daniele Bianchi⁵, Natalya Evans⁶, Allan H. Devol², Richard G. Keil², Andrew
M.P. McDonnell³

¹Horn Point Laboratory, University of Maryland Center for Environmental Science,
Cambridge, MD, USA.

²School of Oceanography, University of Washington Seattle, Seattle, WA, USA.

³College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Fairbanks,
AK, USA.

⁴School of Arts and Sciences, University of Rochester, Rochester, NY, USA.

⁵Department of Atmospheric and Oceanic Sciences, University of California Los
Angeles, Los Angeles, CA, USA.

⁶Department of Biological Sciences, University of Southern California, Los Angeles, CA,
USA.

Contents of this file


Figures S1 to S12

Caption for Text S1


Additional Supporting Information (Files uploaded separately)

Text S1


30 Introduction
31 This file contains supplemental figures referenced in the manuscript. It also contains a
32 caption for a .PDF file containing mathematical equations underpinning the particle
33 remineralization model used.




36 Figure S1. Physical and chemical data from P16 Station 100. Located at 16.5°N
 37 152.0°W. (A) Map of the nearby tropical pacific station P16 Station 100. Colors indicate
 38 chlorophyll concentrations at the surface, averaged over all MODIS images. The red
 39 circle indicates the location of P16 Station 100. (B-D) Oceanographic parameters. The
 40 thin horizontal green line shows the location of the base of the photic zone (200 m). (B)
 41 Oxygen temperature and salinity. (C) Oxygen, and fluorescence. Because the fluorometer
 42 was broken on this cruise, fluorescence data were pulled from world ocean atlas (Garcia
 43 et al. 2014). (D) Beam attenuation and density, calculated from the salinity temperature
 44 and pressure data.




46 Figure S2. Water mass analysis at ETNP Station P2 indicates the relative contributions
 47 proportions of the three primary water masses at this site, **13°C** water (13CW), North
 48 Equatorial Pacific Intermediate Water (NEPIW) and Antarctic Intermediate Water
 49 (AAIW). Values indicate relative contributions of each water mass and are scaled so as to
 50 sum to one. The horizontal green line indicates the base of the photic zone (160m). Data
 51 are taken directly from Evans et al. (2020).




53 Figure S3. Acoustic data, measured by EK60, measured over the course of the
 54 experiment, at ETNP Station P2. Shown are data from all frequency bands. Values are in
 55 return signal intensity and have not been normalized to observed biomass. Horizontal
 56 blue lines indicate the surface and bottom of the ODZ, while the horizontal green line
 57 indicates the base of the photic zone.




59 Figure S4. A profile of particle abundances at different sizes and depths, collected on
 60 January 13 beginning at 06:13 at ETNP Station P2. Horizontal blue lines indicate the
 61 surface and bottom of the ODZ, while the horizontal green line indicates the base of the
 62 photic zone. (A) Numbers of observed particles. As the x axis is log transformed, zeros
 63 are indicated as points along the Y axis. Vertical black lines indicate 1 and 5 observed
 64 particles, respectively. (B) Particle numbers normalized to volume sampled and particle
 65 size bin width. (C) Smoothed and extrapolated particle abundances, based on a negative
 66 binomial GAM that predicts particle abundance form size and depth.







68 Figure S5. An example of observed particle size distribution spectra. These are depth
69 binned data from between 150 and 175 m deep in the water column from the cast that
70 occurred at *2017-01-13 17:51* local time at ETNP Station P2. This depth bin contains
71 total numbers of particles that were seen across 206.8 L of merged UVP image volume.
72 Points indicate (A) total numbers of observed particles and (B) particle numbers
73 normalized to volume sampled and particle size bin width. Half-dots along the x axis
74 correspond to particle size bins in which zero particles were observed. The line indicates
75 the predicted best fit line of the data. The line was fit on the binsize and volume
76 normalized data by a negative-binomial general linear model. The line in panel A
77 indicates predictions from this same model, re-scaled into absolute particle space.




79 Figure S6. GAM predicted effects of A Depth, B Day of the month in January 2017, and
80 C hour of the day on the fifth-root transformed, depth normalized, rate of change of flux
81 at ETNP Station P2. Y axis indicates the value of the component smooth functions effect
82 on Flux. Positive values associate with times and regions of the water column where flux
83 is increasing, holding other factors constant, and negative ones where it is decreasing.
84 Horizontal gray line indicates $y = 0$, corresponding to that parameter having zero effect,
85 positive or on the outcome. Only Depth has a statistically significant relationship to rate
86 of change of flux (see section 6.5).












Figure S11. An example of differences between modeled and observed particle slope. The particle size distribution at a shallow and a deeper depth, from ETNP Station P2, are shown. The model generates a prediction of the deep depth profile from the shallow depth profile and the flux attenuation between the two profiles. The model predicts more attenuation of the smallest particles than is actually observed. In practice the model compares depths that are closer together than the two shown here. In particular, the depth bin above 225m in our analysis has a midpoint of 187.5m, but we choose in this example to compare the 225m particle size profile to the profile at 112.5 m. Two depths that are far apart are shown so that the flux attenuation is large enough to be seen by eye and to provide a conceptual example of the models' function.




Figure S12. GAM predicted effects of A Depth, B Day of the month in January 2017, at ETNP Station P2. Y axis indicates the value of the component smooth functions effect on the difference between observed and modeled flux. Thus higher values correspond with greater flux of small particles than predicted by the model. Horizontal gray line indicates $y = 0$, corresponding to that parameter having zero effect, positive or on the outcome. Only Depth and Day have a statistically significant relationship to rate of change of flux (see Section 6.8).

Text S1. Full mathematical justification for the Eulerian version of the particle remineralization and sinking model (PRiSM) model. Ful document uploaded separately.