## Vaso Constriction - Logistic Regression

## January 25, 2024

First the dataset vaso is loaded.

```
library(catdata)
data(vaso)
attach(vaso)

## Das folgende Objekt ist maskiert durch .GlobalEnv:
##
## vaso
```

For the fitting of a logit model, the response is 0-1 coded. (data set contains 1 2). Moreover, the covariates vol and rate are log-transformed.

```
y <- vaso$vaso
y[vaso$vaso==2] <- 0
```

Fit of a logit-model with log-transformed covariates.

```
vaso1 <- glm(y ~ vol + rate, family=binomial)</pre>
summary(vaso1)
##
## Call:
## glm(formula = y ~ vol + rate, family = binomial)
## Deviance Residuals:
## Min 1Q Median
                                       Max
## -1.4527 -0.6110 0.1001
                          0.6181
                                    2.2775
##
## Coefficients:
   Estimate Std. Error z value Pr(>|z|)
##
## (Intercept) -2.875 1.321 -2.177 0.02946 *
## vol 5.179
                        1.865 2.778 0.00547 **
## rate
               4.562
                         1.838 2.482 0.01306 *
## ---
## Signif. codes:
## 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
```

```
##
## Null deviance: 54.040 on 38 degrees of freedom
## Residual deviance: 29.227 on 36 degrees of freedom
## AIC: 35.227
##
## Number of Fisher Scoring iterations: 6
```

Next, a logit-model with original covariates is fitted.

```
vaso2 <- glm(y ~ I(exp(vol)) + I(exp(rate)), family=binomial)</pre>
summary(vaso2)
##
## Call:
## glm(formula = y ~ I(exp(vol)) + I(exp(rate)), family = binomial)
## Deviance Residuals:
## Min 1Q Median 3Q
                                              Max
## -1.50657 -0.73464 0.03997 0.48854
                                          2.32935
## Coefficients:
## Estimate Std. Error z value Pr(>|z|)
## (Intercept) -9.5296 3.2332 -2.947 0.00320 **
## I(exp(vol)) 3.8822 1.4286 2.717 0.00658 **
## I(exp(rate)) 2.6491
                           0.9142 2.898 0.00376 **
## ---
## Signif. codes:
## 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## (Dispersion parameter for binomial family taken to be 1)
##
      Null deviance: 54.040 on 38 degrees of freedom
## Residual deviance: 29.772 on 36 degrees of freedom
## AIC: 35.772
##
## Number of Fisher Scoring iterations: 6
```