Calculating comorbidities from medical codes

Jack O. Wasey 4/9/2018

AHRQ comorbidity classification

The AHRQ keeps an updated version of the Elixhauser classification of ICD-9-CM and ICD-10-CM codes into comorbidities. They provide the data in the form of SAS code. The names of the comorbidities derived from ICD-9 and ICD-10 codes are the same. Maps contain the ICD code to comorbidity mappings; the functions that apply those mappings are called things like icd10_comorbid_ahrq.

```
#icd9_map_ahrq <- icd:::sas_parse_ahrq() # user doesn't need to do this
names(icd9_map_ahrq)
#>
   [1] "CHF"
                        "Valvular"
                                        "PHTN"
                                                        "PVD"
    [5] "HTN"
                        "HTNcx"
                                        "Paralysis"
                                                        "NeuroOther"
                        "DM"
   [9] "Pulmonary"
                                        "DMcx"
                                                       "Hypothyroid"
#> [13] "Renal"
                        "Liver"
                                        "PUD"
                                                       "HIV"
#> [17] "Lymphoma"
                        "Mets"
                                        "Tumor"
                                                       "Rheumatic"
#> [21] "Coagulopathy" "Obesity"
                                        "WeightLoss"
                                                       "FluidsLytes"
#> [25] "BloodLoss"
                        "Anemia"
                                        "Alcohol"
                                                       "Drugs"
#> [29] "Psychoses"
                        "Depression"
icd9_map_ahrq$CHF[1:5]
#> [1] "39891" "40201" "40211" "40291" "40401"
#> attr(, "icd_short_diag")
#> [1] TRUE
#> attr(,"class")
#> [1] "icd9"
                    "character"
icd10_map_ahrq$CHF[1:5]
#> [1] "I0981" "I501" "I5020" "I5021" "I5022"
#> attr(, "icd_short_diag")
#> [1] TRUE
#> attr(, "class")
#> [1] "icd10"
                    "character"
```

Quan

Quan's paper looked at indices using both ICD-10 and ICD-9-CM. Quan generated updated ICD-9-CM codes for all 30 of Elixhauser and all 17 of Charlson/Deyo's comorbidities. Thus there are two 'Quan' comorbidity mappings.

```
names(icd10_map_quan_deyo)
                                      "PVD"
#> [1] "MI"
                                                     "Stroke"
                                                                    "Dementia"
   [6] "Pulmonary"
                       "Rheumatic"
                                      " מנזק"
                                                     "LiverMild"
                                                                    "DM"
#> [11] "DMcx"
                       "Paralysis"
                                      "Renal"
                                                     "Cancer"
                                                                    "LiverSevere"
#> [16] "Mets"
                       "HIV"
names(icd10_map_quan_elix)
   [1] "CHF"
                                                         "PHTN"
                        "Arrhythmia"
                                        "Valvular"
   [5] "PVD"
                        "HTN"
                                        "HTNcx"
                                                         "Paralysis"
   [9] "NeuroOther"
                        "Pulmonary"
                                         "DM"
                                                         "DMcx"
#> [13] "Hypothyroid"
                        "Renal"
                                        "Liver"
                                                         "PUD"
#> [17] "HIV"
                                                         "Tumor"
                        "Lymphoma"
                                        "Mets"
```

```
#> [21] "Rheumatic" "Coagulopathy" "Obesity" "WeightLoss"
#> [25] "FluidsLytes" "BloodLoss" "Anemia" "Alcohol"
#> [29] "Drugs" "Psychoses" "Depression"
```