

Égalité Fraternité



Modelling plant resistance deployment: the R package landsepi

The R package *landespi* provides a general modelling framework to help compare plant resistance deployment strategies and understand the impact of epidemiological, evolutionary and


genetic factors for a wide range of pathosystems.


Strategies to improve plant resistance management rely on careful selection of resistance sources and their combination at various spatio-temporal scales.

In landsepi, the landscape is a dynamic mosaic of fields cultivated with croptypes. Each croptype is composed of either a pure cultivar or a mixture; and each cultivar may carry one or several resistance genes. Each resistance gene targets one or several pathogenicity traits, with complete or partial efficiency, and may be expressed from the beginning of the season or later (e.g. APR gene). The pathogen may adapt to these genes (restoring its pathogenicity), possibly associated with a fitness cost.

The model is based on a **spatial geometry** for describing the heterogeneous landscape and allocating different cultivars, a dispersal kernel for the dissemination of the pathogen, and a stochastic SEIR ('Susceptible-Exposed-Infectious-Removed') structure with a discrete time step for the description of the host-pathogen interaction. Cropping seasons are split by host harvests which impose potential bottlenecks to the pathogen. The model accounts for pathogen evolution (via mutation, sexual reproduction, selection and drift) and provides epidemiological, evolutionary and economic outputs to assess the performance of the simulated strategies.

The package also includes a shiny interface for pedagogical purposes.





Spatial unit: The spatial unit is a polygon. An agricultural field (i.e. a piece of land cultivated by the same croptype) may be composed of a single or several polygons

Individual: A host 'individual' is an infection unit and may correspond to a given amount of plant tissue (where a local infection may develop, e.g. fungal lesion) or a whole plant (e.g. systemic viral infection). In the first case, plant growth increases the amount of available plant tissue (hence the number of individuals) during the cropping season.

Loup Rimbaud¹, Julien Papaïx², Jean-François Rey² Jean-Loup Gaussen², Marta Zaffaroni^{3,} Frédéric Fabre³

CONTACT DETAILS

- ¹ INRAE Pathologie Végétale, 84140, Montfavet, France.
- ² INRAE BioSP, 84914, Avignon, France.
- 3 INRAE SAVE, 33882 Villenave d'Ornon, France. loup.rimbaud@inrae.fr

PUBLICATIONS

Computation of specific

outputs from raw results

Rimbaud L., Fabre F., Papaīx J., Moury B., Lannou C., Barrett L.G. and Thrall P. H. (2021). Models of plant resistance deployment. Annu. Rev. Phytopathol. 59:125-152.

<--- Outputs -->

- Rimbaud L., Papaïx J., Barrett L.G., Burdon J.J. and Thrall P.H. (2018). Mosaics, mixtures, rotations or pyramiding: What is the optimal strategy to deploy major gene resistance? Evol. Appl. 11:1791-1810.

LINKS

Evolutionary: Resistance durability

Epidemiological: Disease control

ability to reduce disease impact

Economic: Cost efficiency

 $\underline{\textbf{Homepage}}; \textbf{https://csiro-inra.pages.biosp.inrae.fr/landsepi/}$ <u>app</u>:https://shiny.biosp.inrae.fr/app_direct/landsepi/ R package: https://cran.r-project.org/web/packages/landsepi/index.html

ability to limit pathogen evolution and delay resistance breakdown

ability to overcompensate epidemic losses and management costs

ACKNOWLEDGEMENTS

landsepi

This work benefited from ANR project "Ardiv" (2019-2022, grant n°ANR-18-CE32-0004-01), AFB Ecophyto II-Levies Territodaux Project "Médée" (2002-0222), GROC grant CSP00192 and the CSIRO/INRAlinkage program