



simmr output plot Z and G U.lactuca Enteromorpha 3 2 density 1 0 1.00 0.00 0.00 0.50 0.75 1.00 0.00 0.50 0.50 0.25 0.25 0.75 0.25 0.75 1.00 Proportion

simmr output plot Z and G U.lactuca Enteromorpha 3 density 0 -1.00 0.00 0.00 0.50 0.75 1.00 0.00 0.50 0.75 0.50 0.25 0.25 0.25 0.75 1.00

Proportion





Comparison of dietary proportions for sources Z and G and Enteromorpha 1.00 -0.75 -Proportion ... 0.25 -0.00 -Z and G Enteromorpha Source

Comparison of dietary proportions for sources Z and G and Enteromorpha 1.00 -0.75 -Proportion ... 0.25 -0.00 -Z and G Enteromorpha Source

simmr output plot Z and G U.lactuca Enteromorpha 3 density 5 1 0

0.75

0.00

0.25

0.50

0.75

1.00 0.00

0.25

0.50

Proportion

1.00 0.00

0.25

0.50

0.75

1.00

simmr output plot Z and G U.lactuca Enteromorpha 5 -4 3 · density 2. 0 1.00 0.00 1.00 0.00 0.00 0.50 0.50 0.75 0.50 0.25 0.75 0.25 0.75 0.25 1.00

Proportion

Comparison of dietary proportions between sources 0.75 -Proportion -0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 0.75 -Proportion -0.25 -0.00 -Grass Enteromorpha Zostera U.lactuca Source

Comparison of dietary proportions between sources 8.0 0.6 Proportion 9.9 0.2 -0.0 Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 0.75 -Proportion 0.50 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 0.75 -Proportion 0.500 J 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion ... 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source


Comparison of dietary proportions between sources 1.00 -0.75 -Proportion ... 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion ... 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion -0.25 -0.00 -Grass U.lactuca Zostera Enteromorpha Source


Comparison of dietary proportions for source Zostera 0.75 -Proportion ... 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group



Comparison of dietary proportions for source Zostera 0.75 -Proportion 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group


Comparison of dietary proportions for source Zostera 0.75 -0.50 -Proportion 0.25 -0.00 -Period 5 Period 8 Period 2 Period 3 Period 4 Period 6 Period 7 Group


Comparison of dietary proportions for source Zostera 1.00 -0.75 -Proportion 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group





Comparison of dietary proportions for source Zostera 1.00 -0.75 -Proportion 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group


Comparison of dietary proportions for source Zostera 0.75 -0.50 -Proportion 0.25 -0.00 -Period 5 Period 8 Period 2 Period 3 Period 4 Period 6 Period 7 Group















Tracers plot 15.0 -Mixtures Period 1 Mixtures Period 2 12.5 -Mixtures Period 3 Mixtures Period 4 Mixtures Period 5 d15N\_PI Mixtures Period 6 X Mixtures Period 7 Mixtures Period 8 10.0 -0 Zostera × 0 Grass U.lactuca Enteromorpha 7.5 -+ -30 -25 -20 -15 -10 d13C\_PI










Prior and posterior distributions Zostera Grass 3 3 2 -2. 1 Туре Posterior Prior 0 -Density 0.00 0.25 0.50 0.75 1.00 0.25 0.50 0.75 0.00 1.00 Source U.lactuca Enteromorpha Zostera 3 3 Grass U.lactuca Enteromorpha 2 -2. 0 0.75 0.50 0.75 0.50 0.25 1.00 0.00 0.25 1.00 0.00 Proportion

Prior and posterior distributions: Period 1 Zostera Grass 3 3 2 2 -1 Туре Posterior Prior 0 -Density 0.00 0.25 0.50 0.75 1.00 0.25 0.50 0.75 1.00 0.00 Source U.lactuca Enteromorpha Zostera 3 3 -Grass U.lactuca Enteromorpha 2 2 -1 0 0.50 0.75 0.25 0.75 0.00 0.25 0.50 0.00 1.00 Proportion

Prior and posterior distributions: Period 2 Zostera Grass 3 3 2 2 -1 Туре Posterior Prior 0 -Density 0.00 0.25 0.50 0.75 0.25 0.50 0.75 1.00 0.00 1.00 Source U.lactuca Enteromorpha Zostera 3. 3 -Grass U.lactuca Enteromorpha 2 2 -1 0 0.50 0.75 0.25 1.00 0.00 0.25 0.50 0.75 0.00 1.00 Proportion



Prior and posterior distributions Zostera Grass 3 3 2 2 -1 Туре Posterior Prior 0 -Density 0.00 0.25 0.50 0.75 0.25 0.50 0.75 1.00 1.00 0.00 Source U.lactuca Enteromorpha Zostera Grass 3 U.lactuca Enteromorpha 4 2 -2 0 0.50 0.75 0.25 0.75 0.00 0.25 0.50 0.00 1.00 Proportion

Prior and posterior distributions: Period 1 Zostera Grass 3 -9 -2 6 -1 3 -Туре Posterior Prior 0 -Density 0.00 0.25 0.50 0.75 0.25 0.50 0.75 1.00 1.00 0.00 Source U.lactuca Enteromorpha Zostera 3 -Grass 6 -U.lactuca Enteromorpha 2 4 2 -0 0.75 0.25 0.50 1.00 0.00 0.25 0.50 0.00 0.75 1.00 Proportion

Prior and posterior distributions: Period 2 Zostera Grass 3 -3 -2 -2 -1 Туре Posterior Prior 0 -Density 0.00 0.25 0.50 0.75 0.25 0.50 0.75 1.00 1.00 0.00 Source U.lactuca Enteromorpha Zostera 3 -Grass U.lactuca 4 Enteromorpha 2 -3 2 0 0.75 0.75 0.25 0.50 1.00 0.00 0.25 0.50 0.00 Proportion

