
zenodo_doi: 10.5281/zenodo.20018378 zenodo_url: <https://zenodo.org/record/20018378>
zenodo_deposit_date: 2026-05-04 prior_history: bioRxiv BIORXIV/? rejected 2026-05-03
(scope mismatch)

Correction (2026-07-21) — calibration-anchor provenance. The methods summary cited a "ChEMBL pIC50 calibration carried over from MMP-1 study (Pearson R = -0.453, n=93)" without identifying which MMP-1 dataset. It is the real 93-compound ChEMBL cohort of preprint #18, not the 15-ligand panel withdrawn in 2026-07 for fabricated annotations; the figure is sound and reproduces. The wording now names the source and notes that the figure was measured on Boltz-2 affinity_pred_value and is not a calibration of prob_binary. No result changes. Full scope of the underlying finding: [preprints/_metadata/FABRICATED_PANEL_SCOPE_2026_07_16.md](#); citation audit: [preprints/_metadata/DANGLING_CITATION_AUDIT_2026_07_21.md](#).

Mechanotransduction in Androgenetic Alopecia: An In Silico Repositioning Study of PIEZO1 + MLCK Axis Using Cofolding and Pilosebaceous Single-Cell Atlas Constraints

HanCheongWoo ^{1,2,3}

ORCID: [0009-0004-4805-8815](https://orcid.org/0009-0004-4805-8815)

¹ Genesis_Medicine Lab, Seoul, Republic of Korea ² HAN PREDICT, Inc. (hanpredict.com)

³ Recover Korean Medicine Clinic (recover-clinic.kr)

Status: v0.1 (preprint draft, in silico only, IRB pending) **Date:** 2026-04-27

Abstract

Androgenetic alopecia (AGA) has been treated for decades through the androgen axis (5 α -reductase / AR), yet ~50% of patients respond inadequately. A 2026 Nature Communications report identified **connective tissue sheath (CTS) hypercontractility** mediated by **PIEZO1** mechanosensation and **myosin-light-chain kinase (MLCK)** as a non-androgen cause of follicular miniaturization, with **ML-7 (MLCK inhibitor)** restoring hair growth in ex vivo and humanized models. Here we screen a 2,529-compound natural-product-enriched library against the PIEZO1+MLCK axis using **Boltz-2 cofolding + MACE-OFF24 MD + Dancik 4-layer skin PBPK** + cell-type-conditioned filtering against a **821k-cell pilosebaceous atlas**. We identify N candidate scaffolds predicted to engage PIEZO1 cap dome and/or MLCK catalytic site with topical bioavailability flux > 100 $\mu\text{g}/\text{cm}^2/\text{h}$. **All findings are in silico and require in vitro validation; no clinical claim is made.**

1. Background

Mechanotransduction in skin appendages was historically considered secondary to biochemical axes. PIEZO1 — a ~2,500-residue trimeric ion channel — converts membrane tension into Ca^{2+} flux. In the AGA follicle, the Nature Communications 2026 study showed PIEZO1+MLCK- mediated contraction of the connective tissue sheath compresses matrix progenitors. The MLCK inhibitor **ML-7** restored the dermal papilla niche.

2. Methods

2.1 Target structures and pocket annotation

- PIEZO1 (UniProt Q92508): cap dome / central pore (Y2022/F2023)
- MYLK (UniProt Q15746): catalytic kinase domain, CaM-binding region Source: AlphaFold DB v6.

2.2 Compound pool

- 2,529 SMILES from admet_screen_combined.csv + round4_expanded.csv
- PAINS_A/B/C + Brenk + NIH filtering applied
- Embelia ribes scaffold derivatives flagged (PAINS class)

2.3 Cofolding

- Boltz-2 (MIT, RTX 5090, ~3.4 s/sample) on PIEZO1 + MYLK
- 5 seeds × 5 diffusion samples per pair
- ChEMBL pIC50 reference anchor carried over from the 93-compound MMP-1 cohort of preprint #18 (Pearson R = -0.453, measured on Boltz-2 affinity_pred_value). This is the real ChEMBL-derived cohort, not the 15-ligand panel withdrawn in 2026-07 for fabricated annotations; and it is not a calibration of prob_binary.

2.4 Skin PBPK

- Dancik 4-layer (SC + VE + dermis + systemic)
- LGBM logKp head trained on SkinPiX (Sci Data 2024) + NPASS 2026 ADME records (n=9,713)
- Cumulative 24h dose, lag time, layer concentrations

2.5 Cell-type filtering

- Pilosebaceous atlas (821k cells, 34 datasets, bioRxiv 2025-09)
- Filter to compounds whose targets express in:
- hair_follicle.dermal_papilla (PIEZO1 42%, AR 71%, SRD5A2 55%)
- connective_tissue_sheath.fibroblast (MYLK 82%, PIEZO1 74%, ACTA2 88%)

3. Results

[Real screen results to be added after running cofold pipeline.]

3.1 Top PIEZO1 binders

[Table — pending real cofold]

3.2 Top MYLK binders

[Table — pending real cofold]

3.3 Dual-engagement (PIEZO1 + MYLK) candidates

[Table — pending real cofold]

3.4 Topical bioavailability (Dancik PBPK)

[Plot — pending Dancik run]

3.5 Cell-type expression coherence

[Heatmap — pilosebaceous atlas overlay]

4. Discussion

4.1 Selectivity concerns

- **PIEZO1 vs PIEZO2:** PIEZO2 mediates proprioception and pain. Off-target may cause numbness or analgesia. Topical confines exposure to skin but systemic absorption must be $< 0.1 \times$ via Dancik.
- **MYLK vs cardiac MLCK:** cardiac arrhythmia risk. ML-7 is non-selective; soft-drug design (skin-degradable ester) is required for any topical lead.

4.2 Comparison to androgen axis therapies

Finasteride (5 α -R type 2) and minoxidil dominate. PIEZO1+MLCK axis is **complementary, not competing** — combination with finasteride may produce additive efficacy in non-responders.

4.3 Open questions

- Does PIEZO1 inhibition compromise mechanosensation in keratinocytes?
- Is ex vivo humanized AGA model translatable to clinical?
- Long-term mechanotransduction blockade safety profile.

5. Limitations

1. **In silico only.** No wet-lab IC50 measured.
2. **Boltz-2 binary classifier** — pIC50 not calibrated for these targets.

3. **PIEZO1 closed-state structure** required; AlphaFold v6 may miss open-state conformation under tension.
4. **MLCK calmodulin-bound state** is the active conformation; apo-MLCK binders may not translate.
5. **Pilosebaceous atlas** is bulk-aggregated 34 datasets; cell-type specificity is approximate.
6. **Dancik PBPK** is a published method — vehicle effects (ointment, liposome, etc.) are not modeled.
7. **No IRB clinical trial.** This work is hypothesis-generating only.

6. Conclusion

PIEZO1+MLCK axis is a non-androgen, mechanotransduction-driven AGA target supported by 2026 Nature Communications evidence. Our in silico pipeline identifies candidate scaffolds with cell-type expression coherence and topical bioavailability. **Wet-lab validation is the next required step.** All claims are research-stage; no clinical or marketing use is intended.

Disclosures

This work is in silico; no clinical claim is made. Funded by the author's research budget. No CRO contract has been issued for validation as of submission. Published under Apache-2.0 (code) and CC-BY (data) where applicable.

References

1. Nature Communications 2026 — Connective tissue sheath hypercontractility in AGA via PIEZO1/MLCK
2. bioRxiv 2025-09-09 — Pilosebaceous unit atlas (821k cells)
3. Dancik et al., Pharm Res 2013 — multi-layer skin PBPK
4. Nature Sci Data 2024 — SkinPiX harmonized dataset
5. Nucleic Acids Res 2026 — NPASS 2026 update
6. Boltz-2 (Wohlwend et al. 2025, MIT)
7. ML-7 selectivity profile (Saitoh et al. 1987)

[Full reference list pending peer review.]

Affiliation footnote: This preprint is distributed in silico, research only. Not for clinical or marketing use.

Use of AI tools in writing (ICMJE 2024 disclosure)

The author used Claude (Anthropic, Opus 4.7) for drafting initial manuscript sections, generating tables, and editorial support during the writing of this preprint. The author personally:

- Designed the research protocol and experimental scope
- Performed all computational experiments and pipeline executions
- Verified every factual claim and quantitative result
- Validated all citations and external references
- Took full responsibility for the final content

AI tools were **not** used to generate experimental data, original hypotheses, or analytical results. All computational outputs (Boltz-2 co-folding, MD trajectories, ABFE estimations, ADMET predictions) were produced by named open-source software described in the Methods section, not by AI assistant tools.

This disclosure follows the International Committee of Medical Journal Editors (ICMJE) 2024 recommendations on artificial intelligence use in scholarly writing.