

Orthogonal Basis

Hung-yi Lee

Outline

Orthogonal/Orthonormal Basis

Orthogonal Decomposition Theory

How to find Orthonormal Basis

Reference: Textbook Chapter 7.2, 7.3

Orthogonal Set

- A set of vectors is called an orthogonal set if every pair of distinct vectors in the set is orthogonal.

$$\mathcal{S} = \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 5 \\ -4 \\ 1 \end{bmatrix} \right\} \quad \text{An orthogonal set?}$$

By definition, a set with only one vector is an orthogonal set.

Is orthogonal set independent?

Independent?

- Any orthogonal set of nonzero vectors is linearly independent.

Let $\mathcal{S} = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ be an orthogonal set $\mathbf{v}_i \neq \mathbf{0}$ for $i = 1, 2, \dots, k$.

Assume c_1, c_2, \dots, c_k make $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0}$

$$(c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_i\mathbf{v}_i + \dots + c_k\mathbf{v}_k) \cdot \mathbf{v}_i$$

$$= c_1\mathbf{v}_1 \cdot \mathbf{v}_i + c_2\mathbf{v}_2 \cdot \mathbf{v}_i + \dots + c_i\mathbf{v}_i \cdot \mathbf{v}_i + \dots + c_k\mathbf{v}_k \cdot \mathbf{v}_i$$

$$= c_i(\mathbf{v}_i \cdot \mathbf{v}_i) = c_i \underbrace{\|\mathbf{v}_i\|^2}_{\neq 0} \quad \Rightarrow \quad c_i = 0$$

$$\Rightarrow c_1 = c_2 = \dots = c_k = 0$$

Orthonormal Set

标准正交



- A set of vectors is called an orthonormal set if it is an orthogonal set, and the norm of all the vectors is 1

$$\mathcal{S} = \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 5 \\ -4 \\ 1 \end{bmatrix} \right\}$$

$$\begin{array}{ccc} \downarrow & \downarrow & \downarrow \\ \frac{1}{\sqrt{14}} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} & \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} & \frac{1}{\sqrt{42}} \begin{bmatrix} 5 \\ -4 \\ 1 \end{bmatrix} \end{array}$$

Is orthonormal set independent?

Yes

单位向量

A vector that has norm equal to 1 is called a unit vector.

Orthogonal Basis

- A basis that is an orthogonal (orthonormal) set is called an orthogonal (orthonormal) basis

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Orthogonal basis of \mathbb{R}^3

Orthonormal basis of \mathbb{R}^3

Outline

Orthogonal/Orthonormal Basis

Orthogonal Decomposition Theory

How to find Orthonormal Basis

Orthogonal Basis

- Let $S = \{v_1, v_2, \dots, v_k\}$ be an orthogonal basis for a subspace W , and let u be a vector in W .

$$u = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$$
$$\frac{\overset{\text{red arrow}}{u \cdot v_1}}{\|v_1\|^2} \quad \frac{\overset{\text{red arrow}}{u \cdot v_2}}{\|v_2\|^2} \quad \frac{\overset{\text{red arrow}}{u \cdot v_k}}{\|v_k\|^2}$$

Proof

To find c_i

How about orthonormal basis?

$$\begin{aligned} u \cdot v_i &= (c_1 v_1 + c_2 v_2 + \dots + c_i v_i + \dots + c_k v_k) \cdot v_i \\ &= c_1 v_1 \cdot v_i + c_2 v_2 \cdot v_i + \dots + c_i v_i \cdot v_i + \dots + c_k v_k \cdot v_i \\ &= c_i (v_i \cdot v_i) = c_i \|v_i\|^2 \end{aligned} \quad \Rightarrow \quad c_i = \frac{u \cdot v_i}{\|v_i\|^2}$$

Example

- Example: $\mathcal{S} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is an orthogonal basis for \mathcal{R}^3

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 5 \\ -4 \\ 1 \end{bmatrix}$$

$$\text{Let } \mathbf{u} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \text{ and } \mathbf{u} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3.$$

$$c_1 = \frac{\mathbf{u} \cdot \mathbf{v}_1}{\|\mathbf{v}_1\|^2}$$

$$c_2 = \frac{\mathbf{u} \cdot \mathbf{v}_2}{\|\mathbf{v}_2\|^2}$$

$$c_3 = \frac{\mathbf{u} \cdot \mathbf{v}_3}{\|\mathbf{v}_3\|^2}$$

Orthogonal Projection

- Let $S = \{v_1, v_2, \dots, v_k\}$ be an orthogonal basis for a subspace W , and let u be a vector in W .

$$u = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$$
$$\frac{\overset{\text{red arrow}}{u \cdot v_1}}{\|v_1\|^2} \quad \frac{\overset{\text{red arrow}}{u \cdot v_2}}{\|v_2\|^2} \quad \frac{\overset{\text{red arrow}}{u \cdot v_k}}{\|v_k\|^2}$$

- Let u be any vector, and w is the orthogonal projection of u on W .

$$w = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$$
$$\frac{\overset{\text{red arrow}}{u \cdot v_1}}{\|v_1\|^2} \quad \frac{\overset{\text{red arrow}}{u \cdot v_2}}{\|v_2\|^2} \quad \frac{\overset{\text{red arrow}}{u \cdot v_k}}{\|v_k\|^2}$$

Orthogonal Projection

- Let $S = \{v_1, v_2, \dots, v_k\}$ be an orthogonal basis for a subspace W . Let u be any vector, and w is the orthogonal projection of u on W .

$$w = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$$

$$\frac{u \cdot v_1}{\|v_1\|^2} \quad \frac{u \cdot v_2}{\|v_2\|^2} \quad \frac{u \cdot v_k}{\|v_k\|^2}$$

$$P_W = C(C^T C)^{-1} C^T$$

$$C^T = \begin{bmatrix} v_1^T \\ v_2^T \\ \vdots \\ v_n^T \end{bmatrix}$$

$$C = [v_1 \quad \dots \quad v_n]$$

$$P_W = C D C^T$$

Projected:

$$D = \begin{bmatrix} \frac{1}{\|v_1\|^2} & & \\ & \frac{1}{\|v_2\|^2} & \\ & & \ddots \\ & & & \frac{1}{\|v_n\|^2} \end{bmatrix} \quad w = C D C^T u$$

$$C^T u = \begin{bmatrix} u \cdot v_1 \\ \vdots \\ u \cdot v_n \end{bmatrix}$$

Outline

$$w = C \cdot \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \Rightarrow \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} = D C^T u$$
$$\Rightarrow c_i = \frac{u \cdot v_i}{\|v_i\|^2}$$

Orthogonal/Orthonormal Basis

Orthogonal Decomposition Theory

How to find Orthonormal Basis

Orthogonal Basis

Let $\{u_1, u_2, \dots, u_k\}$ be a basis of a subspace V . How to transform $\{u_1, u_2, \dots, u_k\}$ into an orthogonal basis $\{v_1, v_2, \dots, v_k\}$?

**Gram-Schmidt
Process**

Then $\{v_1, v_2, \dots, v_k\}$ is an orthogonal basis for W

Non-zero




L.I.

$$\text{Span } \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_i\} = \text{Span } \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_i\}$$

orthogonal

Visualization



<https://www.youtube.com/watch?v=Ys28-Yq21B8>

$$\mathbf{v}_1 = \mathbf{u}_1,$$

$$\mathbf{v}_2 = \mathbf{u}_2 - \frac{\mathbf{u}_2 \cdot \mathbf{v}_1}{\|\mathbf{v}_1\|^2} \mathbf{v}_1,$$

$$\mathbf{v}_3 = \mathbf{u}_3 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_1}{\|\mathbf{v}_1\|^2} \mathbf{v}_1 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_2}{\|\mathbf{v}_2\|^2} \mathbf{v}_2,$$

$$\vdots$$

$$\mathbf{v}_k = \mathbf{u}_k - \frac{\mathbf{u}_k \cdot \mathbf{v}_1}{\|\mathbf{v}_1\|^2} \mathbf{v}_1 - \frac{\mathbf{u}_k \cdot \mathbf{v}_2}{\|\mathbf{v}_2\|^2} \mathbf{v}_2 - \cdots - \frac{\mathbf{u}_k \cdot \mathbf{v}_{k-1}}{\|\mathbf{v}_{k-1}\|^2} \mathbf{v}_{k-1}$$

$\{u_1, u_2, \dots, u_k\}$ be a basis of a subspace V

$$\text{Span} \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_i\} = \text{Span} \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_i\} = V$$

Intuitive explanation ① $\forall i \in V$, 因为 \mathbf{v}_i 是 \mathbf{u}_i 的线性组合

orthogonal

none zero

② $\dim(V) = k$

③ 非零 正交 \Rightarrow independent

The theorem holds for $k = 1$.

obviously

Assume the theorem holds for $k=n$, and consider the case for $n+1$.

$$\underline{v_{n+1} \cdot v_i = 0} \quad (i < n+1)$$

因为非零正交集是 independent

Example

$v_i \neq 0$ 因为 if $v_i = 0$ Then u_i 是 $v_1 \cdots v_{i-1}$ 的线性组合, 即 u_i 是 $u_1 \cdots u_{i-1}$ 的线性组合, 矛盾

$S = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ Is a basis for subspace W

$$\mathbf{u}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\mathbf{u}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 1 \end{bmatrix}$$

$$\mathbf{u}_3 = \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$$

(L.I. vectors)

Then $S' = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is an orthogonal basis for W .

$$\mathbf{v}_1 = \mathbf{u}_1$$

$S'' = \{\mathbf{v}_1, \mathbf{v}_2, 4\mathbf{v}_3\}$ is also an orthogonal basis.

$$\mathbf{v}_2 = \mathbf{u}_2 - \frac{\mathbf{u}_2 \cdot \mathbf{v}_1}{\|\mathbf{v}_1\|^2} \mathbf{v}_1$$

,

$$\mathbf{v}_3 = \mathbf{u}_3 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_1}{\|\mathbf{v}_1\|^2} \mathbf{v}_1 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_2}{\|\mathbf{v}_2\|^2} \mathbf{v}_2$$