

What can we know
from RREF?

Hung-yi Lee

Reference

- Textbook: Chapter 1.6, 1.7

Outline

- RREF v.s. Linear Combination
- RREF v.s. Independent
- RREF v.s. Rank
- RREF v.s. Span

RREF v.s. Linear Combination

Column Correspondence Theorem

$$A = [\mathbf{a}_1 \quad \cdots \quad \mathbf{a}_n] \xrightarrow{\text{RREF}} R = [\mathbf{r}_1 \quad \cdots \quad \mathbf{r}_n]$$

If \mathbf{a}_j is a linear combination of other columns of A

$$\mathbf{a}_5 = -\mathbf{a}_1 + \mathbf{a}_4$$

\mathbf{r}_j is a linear combination of the corresponding columns of R with the same coefficients

$$\mathbf{r}_5 = -\mathbf{r}_1 + \mathbf{r}_4$$

\mathbf{a}_j is a linear combination of the corresponding columns of A with the same coefficients

$$\mathbf{a}_3 = 3\mathbf{a}_1 - 2\mathbf{a}_2$$

If \mathbf{r}_j is a linear combination of other columns of R

$$\mathbf{r}_3 = 3\mathbf{r}_1 - 2\mathbf{r}_2$$

Column Correspondence Theorem - Example

$$A = \begin{array}{c} \mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4 \quad \mathbf{a}_5 \quad \mathbf{a}_6 \\ \left[\begin{array}{cccccc} 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{array} \right] \end{array} \quad R = \begin{array}{c} \mathbf{r}_1 \quad \mathbf{r}_2 \quad \mathbf{r}_3 \quad \mathbf{r}_4 \quad \mathbf{r}_5 \quad \mathbf{r}_6 \\ \left[\begin{array}{cccccc} 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right] \end{array}$$

$$\mathbf{a}_2 = 2\mathbf{a}_1$$



$$\mathbf{r}_2 = 2\mathbf{r}_1$$

$$\mathbf{a}_5 = -\mathbf{a}_1 + \mathbf{a}_4$$



$$\mathbf{r}_5 = -\mathbf{r}_1 + \mathbf{r}_4$$

Column Correspondence Theorem – Intuitive Idea

$$\begin{array}{ccc}
 & a_1 + a_2 = a_3 & \\
 & A = \begin{bmatrix} 6 & 9 & 15 \\ 8 & 0 & 8 \\ 9 & 2 & 11 \end{bmatrix} & \\
 & \downarrow & \\
 & c_1 + c_2 = c_3 & \\
 & C = \begin{bmatrix} 12 & 18 & 30 \\ 8 & 0 & 8 \\ 9 & 2 & 11 \end{bmatrix} & \text{X2}
 \end{array}$$

$$\begin{array}{ccc}
 & & d_1 + d_2 = d_3 \\
 & & D = \begin{bmatrix} 6 & 9 & 15 \\ 8 & 0 & 8 \\ 3 & -7 & -4 \end{bmatrix} \text{X-1}
 \end{array}$$

$$\begin{array}{ccc}
 b_1 + b_2 = b_3 & \leftarrow & \\
 B = \begin{bmatrix} 9 & 2 & 11 \\ 8 & 0 & 8 \\ 6 & 9 & 15 \end{bmatrix} & &
 \end{array}$$

交換

Column Correspondence Theorem (Column 間的承諾) :
 就算 row elementary operation 讓 column 變的不同，
 他們之間的關係永遠不變。

Column Correspondence Theorem – Reason

- Before we start:

Coefficient Matrix: $A \xrightarrow{\text{RREF}} R$

Augmented Matrix: $[A \ b] \xrightarrow{\text{RREF}} [R \ b']$

$$\begin{array}{c} \left[\begin{array}{ccccc|c} 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{array} \right] \quad \longrightarrow \quad \left[\begin{array}{ccccc|c} 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right] \\ [A \ b] \qquad \qquad \qquad [R \ b'] \end{array}$$

Column Correspondence Theorem – Reason

- The RREF of matrix A is R

$Ax = b$ and $Rx = b$ have
the same solution set?



- The RREF of augmented matrix $[A \quad b]$ is $[R \quad b']$

$Ax = b$ and $Rx = b'$ have
the same solution set



- The RREF of matrix A is R

$Ax = 0$ and $Rx = 0$ have
the same solution set



If $b = 0$,
then $b' = 0$.

Column Correspondence

Theorem – Reason

- The RREF of matrix A is R , $Ax = 0$ and $Rx = 0$ have the same solution set

$$A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \mathbf{a}_3 & \mathbf{a}_4 & \mathbf{a}_5 & \mathbf{a}_6 \\ 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix} \quad R = \begin{bmatrix} \mathbf{r}_1 & \mathbf{r}_2 & \mathbf{r}_3 & \mathbf{r}_4 & \mathbf{r}_5 & \mathbf{r}_6 \\ 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{c}
 \mathbf{a}_2 = 2\mathbf{a}_1 \\
 \updownarrow \\
 -2\mathbf{a}_1 + \mathbf{a}_2 = \mathbf{0}
 \end{array}
 \longleftrightarrow
 \begin{array}{c}
 \boxed{Ax = 0} \\
 x = \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}
 \end{array}
 \longleftrightarrow
 \begin{array}{c}
 \boxed{Rx = 0} \\
 x = \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}
 \end{array}
 \longleftrightarrow
 \begin{array}{c}
 \mathbf{r}_2 = 2\mathbf{r}_1 \\
 \updownarrow \\
 -2\mathbf{r}_1 + \mathbf{r}_2 = \mathbf{0}
 \end{array}$$

Column Correspondence

Theorem – Reason

- The RREF of matrix A is R , $Ax = 0$ and $Rx = 0$ have the same solution set

$$A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \mathbf{a}_3 & \mathbf{a}_4 & \mathbf{a}_5 & \mathbf{a}_6 \\ 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix} \quad R = \begin{bmatrix} \mathbf{r}_1 & \mathbf{r}_2 & \mathbf{r}_3 & \mathbf{r}_4 & \mathbf{r}_5 & \mathbf{r}_6 \\ 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{c}
 \mathbf{a}_5 = -\mathbf{a}_1 + \mathbf{a}_4 \\
 \Updownarrow \\
 \mathbf{a}_1 - \mathbf{a}_4 + \mathbf{a}_5 = 0
 \end{array}
 \longleftrightarrow
 \begin{array}{c}
 \boxed{Ax = 0} \\
 x = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix}
 \end{array}
 \longleftrightarrow
 \begin{array}{c}
 \boxed{Rx = 0} \\
 x = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix}
 \end{array}
 \longleftrightarrow
 \begin{array}{c}
 \mathbf{r}_5 = -\mathbf{r}_1 + \mathbf{r}_4 \\
 \Updownarrow \\
 \mathbf{r}_1 - \mathbf{r}_4 + \mathbf{r}_5 = 0
 \end{array}$$

How about Rows?

- Are there row correspondence theorem? **NO**

$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix}$$

$$A = \begin{bmatrix} \text{---} a_1^T \text{---} \\ \text{---} a_2^T \text{---} \\ \text{---} a_3^T \text{---} \\ \text{---} a_4^T \text{---} \end{bmatrix}$$

$$\text{Span}\{a_1, a_2, a_3, a_4\}$$

$$R = \begin{bmatrix} 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R = \begin{bmatrix} \text{---} r_1^T \text{---} \\ \text{---} r_2^T \text{---} \\ \text{---} r_3^T \text{---} \\ \text{---} r_4^T \text{---} \end{bmatrix}$$

$$\text{Span}\{r_1, r_2, r_3, r_4\}$$

=

Are they the same?

Span of Columns

$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix} \quad R = \begin{bmatrix} 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$A = [\mathbf{a}_1 \quad \cdots \quad \mathbf{a}_6]$$

$$R = [\mathbf{r}_1 \quad \cdots \quad \mathbf{r}_6]$$

$$\text{Span} \{ \mathbf{a}_1, \quad \cdots \quad , \mathbf{a}_6 \}$$

$$\text{Span} \{ \mathbf{r}_1, \quad \cdots \quad , \mathbf{r}_6 \}$$

Are they the same?

The elementary row operations change the span of columns.


NOTE

- Original Matrix v.s. RREF
 - Columns:
 - The relations between the columns are the same.
 - The span of the columns are different.
 - Rows:
 - The relations between the rows are changed.
 - The span of the rows are the same.

RREF v.s. Independent

Column Correspondence Theorem


pivot columns



$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix}$$

linear
independent

Leading entries



$$R = \begin{bmatrix} 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

linear
independent

The pivot columns are linear independent.

Column Correspondence Theorem

pivot columns

$$A = \begin{bmatrix} 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix}$$

$$\mathbf{a}_2 = 2\mathbf{a}_1$$

$$\mathbf{a}_5 = -\mathbf{a}_1 + \mathbf{a}_4$$

$$\mathbf{a}_6 = -5\mathbf{a}_1 - 3\mathbf{a}_3 + 2\mathbf{a}_4$$

Leading entries

$$R = \begin{bmatrix} 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\mathbf{r}_2 = 2\mathbf{r}_1$$

$$\mathbf{r}_5 = -\mathbf{r}_1 + \mathbf{r}_4$$

$$\mathbf{r}_6 = -5\mathbf{r}_1 - 3\mathbf{r}_3 + 2\mathbf{r}_4$$

The non-pivot columns are the linear combination of the previous pivot columns.

Independent

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

All columns are
independent



Every column is a
pivot column



Every column in
RREF(A) is standard
vector.

3X3

$$\begin{bmatrix} * & * & * \\ * & * & * \\ * & * & * \end{bmatrix}$$

Columns are linear
independent

RREF



$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Identity
matrix

Independent

All columns are
independent



Every column is a
pivot column



Every column in
RREF(A) is standard
vector.

4X3

$$\begin{bmatrix} * & * & * \\ * & * & * \\ * & * & * \\ * & * & * \end{bmatrix}$$

Columns are linear
independent

RREF



$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} I \\ 0 \end{bmatrix}$$

Independent

All columns are independent



Every column is a pivot column



Every column in $\text{RREF}(A)$ is standard vector.

3X4

$$\begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{bmatrix}$$

Columns are linearly independent



RREF



$$\begin{bmatrix} 1 & 0 & 0 & * \\ 0 & 1 & 0 & * \\ 0 & 0 & 1 & * \end{bmatrix}$$

Cannot be a pivot column



Independent

$$\begin{bmatrix} 1 & 2 \\ 1 & 2 \\ 1 & 2 \end{bmatrix}$$

因為太胖了，自己走不動

$$\begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{bmatrix}$$

(矮胖型)



The columns are dependent



$$\mathcal{S} = \left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \right\}$$

Dependent or Independent?

More than 3 vectors in \mathbb{R}^3 must be dependent.

More than m vectors in \mathbb{R}^m must be dependent.

Independent – Intuition



RREF v.s. Rank

Rank

Maximum number of
Independent Columns

||

Number of Pivot
Column

||

Number of Non-zero
rows

$R(A) =$

$$\begin{bmatrix} 1 & 2 & -1 & 2 & 1 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix}$$

Rank = ? **3**

$R(R) =$

$$\begin{bmatrix} \boxed{1} & 2 & \boxed{0} & \boxed{0} & -1 & -5 \\ 0 & 0 & \boxed{1} & 0 & 0 & -3 \\ 0 & 0 & 0 & \boxed{1} & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Rank = ? **3**

Properties of Rank from RREF


Maximum number of
Independent Columns

||


Number of Pivot
Column

||


Number of Non-zero
rows



$\text{Rank } A \leq \text{Number of columns}$



$\text{Rank } A \leq \text{Min}(\text{Number of columns, Number of rows})$



$\text{Rank } A \leq \text{Number of rows}$

Properties of Rank from RREF

- Given a $m \times n$ matrix A:

Matrix A is full rank
if $\text{Rank } A = \min(m, n)$

- $\text{Rank } A \leq \min(m, n)$
- Because “the columns of A are independent” is equivalent to “ $\text{rank } A = n$ ”
 - If $m < n$, the columns of A is dependent.

$$\begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{bmatrix}$$

3 X 4

$\text{Rank } A \leq 3$

$$\left\{ \begin{bmatrix} * \\ * \\ * \end{bmatrix}, \begin{bmatrix} * \\ * \\ * \end{bmatrix}, \begin{bmatrix} * \\ * \\ * \end{bmatrix}, \begin{bmatrix} * \\ * \\ * \end{bmatrix} \right\}$$

A matrix set has 4 vectors
belonging to \mathbb{R}^3 is dependent

- In \mathbb{R}^m , you cannot find more than m vectors that are independent.

满秩

4
3

rank A
 ≤ 3

Basic, Free Variables v.s. Rank

$$Ax = b$$

$$\begin{aligned} x_1 + 2x_2 - x_3 + 2x_4 + 5x_5 &= 2 \\ -x_1 - 2x_2 + x_3 + 2x_4 + 3x_5 &= 6 \\ 2x_1 + 4x_2 - 3x_3 + 2x_4 &= 3 \\ -3x_1 - 6x_2 + 2x_3 + 3x_5 &= 9 \end{aligned}$$

$$\begin{aligned} x_1 + 2x_2 & & -x_5 &= -5 \\ & x_3 & &= -3 \\ & & x_4 + x_5 &= 2 \\ & & \cancel{0} &= \cancel{0} \end{aligned}$$

3 useful equations

$$\begin{bmatrix} 1 & 2 & -1 & 2 & 5 & 2 \\ -1 & -2 & 1 & 2 & 3 & 6 \\ 2 & 4 & -3 & 2 & 0 & 3 \\ -3 & -6 & 2 & 0 & 3 & 9 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 2 & 0 & 0 & -1 & -5 \\ 0 & 0 & 1 & 0 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

A $\text{RREF}(A)$ b'

rank = non-zero row = 3 basic variables

nullity = No. column - non-zero row = 2 free variables

$$x_1 = -5 - 2x_2 + x_5$$

$$x_3 = -3$$

$$x_4 = 2 - x_5$$

Rank

Rank

Maximum number of
Independent Columns

Number of Pivot
Column



Number of Non-zero
rows of RREF


Number of Basic
Variables

Nullity = no. column - rank



Number of zero rows
of RREF

Number of Free
Equations



RREF v.s. Span

Consistent or not

- Given $\underline{A}x = \underline{b}$, if the reduced row echelon form of $\begin{bmatrix} A \\ b \end{bmatrix}$ is

$$\begin{bmatrix} 1 & 0 & 3 & 1 \\ 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Consistent

b is in the span of the columns of A

- Given $Ax = b$, if the reduced row echelon form of $\begin{bmatrix} A \\ b \end{bmatrix}$ is

$$\begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

inconsistent

b is NOT in the span of the columns of A

$$0 \cdot x_1 + 0 \cdot x_2 + 0 \cdot x_3 = 1$$

Consistent or not

$Ax = b$ is inconsistent (no solution)



The RREF of $[A \ b]$ is

Only the last
column is non-zero

*	*	*	*	*	*
*	*	*	*	*	*
*	*	*	*	*	*
0	0	0	0	0	d
0	0	0	0	0	0
0	0	0	0	0	0


$d \neq 0$

$\text{Rank } A \neq \text{rank } [A \ b]$


Need to know b

Consistent or not

$Ax = b$ is consistent for *every* b



RREF of $[A \ b]$ cannot have a row whose only non-zero entry is at the last column



RREF of A cannot have zero row



Rank A = no. of rows


Consistent or not

e.g. $A = \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{bmatrix}$

3 independent columns

$m \times n$ n m $m=3$ $n=4$

$Ax = b$ is consistent for **every** b



Rank $A = \text{no. of rows}$

Every b is in the span of the columns of

$$A = [a_1 \ \cdots \ a_n]$$

Every b belongs to $\text{Span}\{a_1, \dots, a_n\}$

$$\text{Span}\{a_1, \dots, a_n\} = R^m$$

m independent vectors can span R^m

这个角度也可论证有解




More than m vectors in R^m must be dependent.

m independent vectors
can span R^m



More than m vectors in
 R^m must be dependent.

- Consider R^2



Does $\mathcal{S} = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix} \right\}$ generate R^3 ? yes

independent

Full Rank: Rank = n & Rank = m

- The size of A is m x n

$$\text{Rank } A = n$$

A is square or 高瘦

1	0	*	*	*	*
0	1	*	*	*	*
⋮	⋮	⋮	⋮	⋮	⋮
⋮	⋮	⋮	⋮	⋮	⋮
0	0	*	*	*	*
0	0	*	*	*	*

$A\mathbf{x} = \mathbf{b}$ has at most one solution

The columns of A are linearly independent.

RREF of A: $\begin{bmatrix} I \\ 0 \end{bmatrix}$

All columns are pivot columns.

Full Rank: Rank = n & Rank = m

- The size of A is $m \times n$

$$\text{Rank } A = m$$

A is square or 矮胖

1	*	*	*	*	*
0	1	*	*	*	*
⋮	⋮	⋮	⋮	⋮	⋮
0	0	0	1	*	*
0	0	0	0	1	*

Every row of R contains a pivot position (leading entry).

$A\mathbf{x} = \mathbf{b}$ always have solution (at least one solution) for every \mathbf{b} in \mathcal{R}^m .

The columns of A generate \mathcal{R}^m .