

Matrix Multiplication

Hung-yi Lee

Reference

- Textbook: Chapter 2.1

Matrix Multiplication

- Given two matrices A and B, the (i, j) -entry of AB is the inner product of **row i of A** and **column j of B**

row i of A \longrightarrow

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ \vdots & \vdots & & \vdots \\ a_{i1} & a_{i2} & \cdots & a_{in} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

A

column j of B

\downarrow

$$\begin{bmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1p} \\ b_{21} & \cdots & b_{2j} & \cdots & b_{2p} \\ \vdots & & \vdots & & \vdots \\ b_{n1} & \cdots & b_{nj} & \cdots & b_{np} \end{bmatrix}$$

B

$$C = AB$$

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}$$

Matrix Multiplication

- Given two matrices A and B, the (i, j) -entry of AB is the inner product of **row i of A** and **column j of B**


$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \quad B = \begin{bmatrix} -1 & 1 \\ 3 & 2 \end{bmatrix}$$

$$\mathbf{C = AB} = \begin{bmatrix} (-1) \times 1 + 3 \times 2 & 1 \times 1 + 2 \times 2 \\ (-1) \times 3 + 3 \times 4 & 1 \times 3 + 2 \times 4 \\ (-1) \times 5 + 3 \times 6 & 1 \times 5 + 2 \times 6 \end{bmatrix}$$

Matrix Multiplication – 4 ways

- Way 1: inner product


Given two matrices A and B,
the (i, j) -entry of AB is the
inner product of **row i of A**
and **column j of B**



Matrix Multiplication – 4 ways


- Way 1: inner product

Given two matrices A and B,
the (i, j) -entry of AB is the
inner product of **row i of A**
and **column j of B**



Matrix Multiplication – 4 ways

- Way 2: Linear combination of columns



$$= \left[\begin{array}{c} b_{11}a_1 + b_{12}a_2 + \dots + b_{1n}a_n \\ b_{21}a_1 + b_{22}a_2 + \dots + b_{2n}a_n \\ \dots \end{array} \right]$$

The first column The second column

Matrix Multiplication – 4 ways

- Way 2: Linear combination of columns

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 3 & 2 \end{bmatrix} = \left[\begin{array}{c} -1 \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} + 3 \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} \quad 1 \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} \end{array} \right]$$

The first column The second column

Matrix Multiplication – 4 ways

- Way 3: Linear combination of rows

$$\begin{array}{c} a_1^T \\ a_2^T \\ \vdots \\ a_m^T \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \vdots \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \vdots \\ \text{---} \end{array} \begin{array}{c} b_1^T \\ b_2^T \\ \vdots \\ b_n^T \end{array} = \left[\begin{array}{c} a_{11}b_1^T + a_{12}b_2^T \cdots + a_{1n}b_n^T \\ a_{21}b_1^T + a_{22}b_2^T \cdots + a_{2n}b_n^T \\ \vdots \\ a_{m1}b_1^T + a_{m2}b_2^T \cdots + a_{mn}b_n^T \end{array} \right]$$

Matrix Multiplication – 4 ways

- Way 3: Linear combination of rows

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} 1[-1 \quad 1] + 2[3 \quad 2] \\ 3[-1 \quad 1] + 4[3 \quad 2] \\ 5[-1 \quad 1] + 6[3 \quad 2] \end{bmatrix}$$


The first row

The second row

The third row

Matrix Multiplication – 4 ways

- Way 4: summation of matrices



$$= a_1 b_1^T + a_2 b_2^T + \dots + a_n b_n^T$$

matrices

Matrix Multiplication – 4 ways

- Way 4: summation of matrices

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 3 & 2 \end{bmatrix} = \left[\begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} \begin{bmatrix} -1 & 1 \end{bmatrix} + \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} \begin{bmatrix} 3 & 2 \end{bmatrix} \right]$$

Diagram annotations:
- The first matrix is labeled "1 x 2".
- The second matrix is labeled "2 x 1".
- A blue arrow points from the "2 x 1" label to the "1 x 1" label, indicating the resulting dimensions of the products.

$$= \left[\begin{bmatrix} -1 & 1 \\ -3 & 3 \\ -5 & 5 \end{bmatrix} + \begin{bmatrix} 6 & 4 \\ 12 & 8 \\ 18 & 12 \end{bmatrix} \right]$$

Rank = ? (under the first matrix) Rank = ? (under the second matrix)

Augmentation and Partition

- Augment: the augment of A and B is $[A \ B]$
- Partition:

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 4 & 3 & 2 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 4 & 3 & 2 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 4 & 3 & 2 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 4 & 3 & 2 & 1 \end{bmatrix}$$

Block Multiplication

$$A = \begin{bmatrix} \boxed{1} & \boxed{3} & \boxed{4} & \boxed{2} \\ \boxed{0} & \boxed{5} & \boxed{-1} & \boxed{6} \\ \boxed{1} & \boxed{0} & \boxed{3} & \boxed{-1} \end{bmatrix}$$

$$B = \begin{bmatrix} \boxed{1} & \boxed{0} & \boxed{3} \\ \boxed{1} & \boxed{2} & \boxed{0} \\ \boxed{2} & \boxed{-1} & \boxed{2} \\ \boxed{0} & \boxed{3} & \boxed{1} \end{bmatrix}$$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad B = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$AB = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$= \begin{bmatrix} \underbrace{\phantom{A_{11}B_{11} + A_{12}B_{21}}} & \phantom{A_{11}B_{12} + A_{12}B_{22}} \\ \phantom{A_{21}B_{11} + A_{22}B_{21}} & \phantom{A_{21}B_{12} + A_{22}B_{22}} \end{bmatrix}$$

Multiply as the small
matrices are scalar

Don't switch the order

Block Multiplication

$$A = \begin{bmatrix} \boxed{1} & \boxed{3} & \boxed{4} & \boxed{2} \\ \boxed{0} & \boxed{5} & \boxed{-1} & \boxed{6} \\ \boxed{1} & \boxed{0} & \boxed{3} & \boxed{-1} \end{bmatrix}$$

"2 x 2"

$$B = \begin{bmatrix} \boxed{1} & \boxed{0} & \boxed{3} \\ \boxed{1} & \boxed{2} & \boxed{0} \\ \boxed{2} & \boxed{-1} & \boxed{2} \\ \boxed{0} & \boxed{3} & \boxed{1} \end{bmatrix}$$

"2 x 2"

不是所有


partition

相乘都有

意义

$AB =$

"2 x 2"



Block Multiplication



$$A = \left[\begin{array}{cc|cc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \hline 6 & 8 & 5 & 0 \\ -7 & 9 & 0 & 5 \end{array} \right] \quad A = \begin{bmatrix} I_2 & O \\ B & 5I_2 \end{bmatrix} \quad B = \begin{bmatrix} 6 & 8 \\ -7 & 9 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} I_2 & O \\ B & 5I_2 \end{bmatrix} \begin{bmatrix} I_2 & O \\ B & 5I_2 \end{bmatrix} = \begin{bmatrix} & \\ & \end{bmatrix}$$

$$A^3 = AA^2 = \begin{bmatrix} I_2 & O \\ B & 5I_2 \end{bmatrix} \begin{bmatrix} I_2 & O \\ 6B & 25I_2 \end{bmatrix} = \begin{bmatrix} & \\ & \end{bmatrix}$$

Matrix Multiplication - Meaning

- Multiple Input $C = AB$




$$\begin{aligned} AB &= A[b_1 \quad b_2 \quad \dots \quad b_p] \\ &= [Ab_1 \quad Ab_2 \quad \dots \quad Ab_p] \end{aligned}$$

Matrix Multiplication - Meaning

- **Composition**


- Given two function f and g , the function $g(f(.))$ is the composition $g \circ f$.




Matrix multiplication is the composition of two linear functions.

Matrix Multiplication - Meaning


- Composition




Matrix Multiplication - Meaning



Matrix Multiplication - Meaning






The composition of A and B is

$$C = [Ab_1 \quad Ab_2 \quad \cdots \quad Ab_p]$$

Matrix Multiplication

Example

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} & \\ & \end{bmatrix}$$



Not Commutative


- $AB \neq BA$

$$A = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$AB = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 3 & -1 \\ 2 & 0 \end{bmatrix}$$

$$\neq BA = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix}$$

Not Communicative



If A and B are matrices, then both AB and BA are defined if and only if A and B are square matrices?

~~X~~

$m \times n$ $n \times m$
A B

Properties

- Let A and B be $k \times m$ matrices, C be an $m \times n$ matrix, and P and Q be $n \times p$ matrices
 - For any scalar s , $s(AC) = (sA)C = A(sC)$
 - $(A + B)C = AC + BC$
 - $C(P+Q)=CP+CQ$
 - $I_k A = A = A I_m$
 - The product of any matrix and a zero matrix is a zero matrix
- Power of square matrices: $A \in \mathcal{M}_{n \times n}$, $A^k = A A \cdots A$ (k times), and by convention, $A^1 = A$, $A^0 = I_n$.

Properties

$$AC: k \times n \quad (AC)^T: n \times k$$

- Let A be $k \times m$ matrices, C be an $m \times n$ matrix,
 - $(AC)^T = ? \quad C^T A^T$

$$\begin{array}{ccc} & A^T C^T & \\ \swarrow & & \searrow \\ m \times k & & n \times m \end{array}$$



$$\begin{array}{ccc} & C^T A^T & \\ \swarrow & & \searrow \\ n \times m & & m \times k \\ \searrow & & \swarrow \\ & n \times k & \end{array}$$

Special Matrix

- Diagonal Matrix

对角上对应元素相乘即可

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix} \quad B = \begin{bmatrix} 3 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \quad \underline{AB} = \begin{bmatrix} 3 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 6 \end{bmatrix}$$

- Symmetric Matrix $A^T = A$


$$A = \begin{bmatrix} 1 & 2 & 4 \\ 2 & 3 & -1 \\ 4 & -1 & 5 \end{bmatrix} = A^T \quad B = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \neq B^T$$

AA^T and $A^T A$ are square and symmetric

$$(AA^T)^T = A^{TT}A^T = AA^T \quad (A^T A)^T = A^T A^{TT} = A^T A$$

Practical Issue

- Let A and B be $k \times m$ matrices, C be an $m \times n$ matrix, and P and Q be $n \times p$ matrices
 - $A(CP) = (AC)P$



Multiplication count: $m \times n \times p$

Practical Issue





k=1 m=1000

n=1 p=1000


- Let A and B be k x m matrices, C be an m x n matrix, and P and Q be n x p matrices

- $A(CP) = (AC)P$

经典动态规划



Practical Issue - GPU



Multiplying two 10000 X 10000 matrices

CPU It cost 21.249996 sec

GPU It cost 0.843893 sec

(GTX 980 Ti)

More than 20 times faster