

## Math 415. Final Exam. December 16, 2017

Full Name: \_\_\_\_\_

Net ID: \_\_\_\_\_

Discussion Section: \_\_\_\_\_

- There are 31 problems worth 5 points each.
  - You must not communicate with other students.
  - No books, notes, calculators, or electronic devices allowed.
  - This is a 180 minutes exam.
  - Do not turn this page until instructed to.
  - Fill in the answers on the scantron form provided. Also circle your answers on the exam itself.
  - Hand in both the exam and the scantron.
  - On the scantron make sure you bubble in **your name, your UIN and your NetID**.
  - There are several different versions of this exam.
  - Please double check that you correctly bubbled in your answer on the scantron. It is the answer on the scantron that counts!
  - Good luck!
- 

**Fill in the following information on the scantron form:**

On the first page of the scantron bubble in **your name, your UIN and your NetID**!  
On the back of the scantron bubble in the following:

95. D

96. C

1. (5 points) Let  $A$  be a  $3 \times 5$ -matrix such that

$$\text{Col}(A) = \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 3 \end{bmatrix} \right\}.$$

Then the dimension of  $\text{Col}(A^T)$  is

- (A) 1
- (B) 3
- (C) 4
- (D) 0
- (E) 2

2. (5 points) Let  $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 4 & 4 \\ 1 & 4 & 8 \end{bmatrix}$ . Let  $L$  be a lower triangular  $3 \times 3$ -matrix and  $U$  be an upper triangular matrix such that  $A = LU$ . Which of the following matrices can be  $L$ ?

(A)  $\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$

(B) None of the other answers

(C)  $\begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix}$

(D)  $\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$

(E)  $\begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$

3. (5 points) Which of the following is a basis of  $\text{Span} \left( \begin{bmatrix} 2 \\ 3 \\ -4 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ -3 \\ 4 \end{bmatrix} \right)$ ?


(A)  $\begin{bmatrix} 2 \\ 3 \\ -4 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$

(B)  $\begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 3 \end{bmatrix}$

(C)  $\begin{bmatrix} 2 \\ 3 \\ -4 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ -3 \\ 4 \end{bmatrix}$

(D)  $\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 3 \end{bmatrix}$

4. (5 points) Which matrix is the PageRank matrix for the following system of web-pages?



(A) 
$$\begin{bmatrix} 0 & 0 & 0 & 1 & 1 & 0 \\ \frac{1}{2} & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{3} & 0 & 0 & 1 \\ 0 & 0 & \frac{1}{3} & 0 & 0 & 0 \end{bmatrix}$$

(B) 
$$\begin{bmatrix} 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{3} & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & 0 & 0 & \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

(C) 
$$\begin{bmatrix} -1 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 \\ 0 & -1 & \frac{1}{3} & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & 0 & -1 & \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \\ 1 & 0 & 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

(D) 
$$\begin{bmatrix} -1 & 0 & 0 & 1 & 1 & 0 \\ \frac{1}{2} & -1 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & -1 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & \frac{1}{3} & -1 & 0 & 0 \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{3} & -1 & 0 & 1 \\ 0 & 0 & \frac{1}{3} & 0 & 0 & -1 \end{bmatrix}$$

(E) None of the other answers.

5. (5 points) Let  $A$  be an  $\ell \times m$  matrix such that for every  $\mathbf{b}$  in  $\mathbb{R}^\ell$ , the equation  $A\mathbf{x} = \mathbf{b}$  has a solution. What does this statement imply about the relative size of  $\ell$  and  $m$ ?

- (A)  $\ell = m$
- (B) nothing ( $\ell$  and  $m$  can be any positive integers)
- (C)  $\ell \geq m$
- (D)  $\ell \leq m$

6. (5 points) Consider the following two statements:

(T1) Every linearly independent set of vectors in a vector space  $V$  is a basis of  $V$  or can be extended to a basis of  $V$ .

(T2) If  $\mathbf{v}_1, \dots, \mathbf{v}_n$  are linearly independent vectors in a vector space  $V$ , then  $\dim(V) \geq n$ .

Then:

- (A) Only Statement T2 is correct.
- (B) Neither Statement T1 nor Statement T2 is correct.
- (C) Only Statement T1 is correct.
- (D) Statement T1 and Statement T2 are correct.

7. (5 points) Which of the following subsets of  $\mathbb{R}^2$  is a subspace?

$$U = \left\{ \begin{bmatrix} a \\ b \end{bmatrix} : ab \geq 0 \right\}$$

$$V = \left\{ \begin{bmatrix} 3a \\ -b \end{bmatrix} : a - b = 0 \right\}$$

$$W = \left\{ \begin{bmatrix} a \\ 2b + 1 \end{bmatrix} : a + b = 0 \right\}$$

- (A)  $V$  and  $W$ , but not  $U$
- (B)  $V$  only
- (C)  $U$ ,  $V$  and  $W$
- (D)  $W$  only
- (E) None of them

8. (5 points) Let  $A$  be an  $n \times n$  matrix, let  $D$  be a diagonal matrix and let  $P$  be an invertible matrix such that  $A = PDP^{-1}$ . Which of the following statements is always true?

- (A)  $\text{Nul}(A) = \text{Nul}(D)$
- (B)  $\dim \text{Nul}(A) \neq \dim \text{Nul}(D)$
- (C) None of the other answers
- (D)  $\dim \text{Nul}(A) = \dim \text{Nul}(D)$

9. (5 points) Consider the following two statements:

- (T1) If a vector space  $V$  has dimension  $d$ , then every set of  $d$  linearly independent vectors  $\mathbf{v}_1, \dots, \mathbf{v}_d$  in  $V$  forms a basis of  $V$ .
- (T2) Every linearly independent set of vectors in a vector space  $V$  is a basis of  $V$  or can be extended to a basis of  $V$ .

Then:

- (A) Only Statement T2 is correct.
- (B) Statement T1 and Statement T2 are correct.
- (C) Neither Statement T1 nor Statement T2 is correct.
- (D) Only Statement T1 is correct.

10. (5 points) Let  $V$  be a subspace of  $\mathbb{R}^n$ ,  $n > 0$  and let  $P$  be the projection matrix of the projection onto  $V$ . Which of the following statements is not always true?

- (A)  $\text{Nul}(P) = V^\perp$ .
- (B)  $\text{rank}(P) = \dim V$ .
- (C)  $\text{Col}(P) = V$ .
- (D)  $\text{Col}(P^T) = V^\perp$ .



11. (5 points) Let  $A$  be a  $5 \times 4$  matrix. Let  $a_1, a_2, a_3, a_4$  be the columns of  $A$  and suppose that  $a_1, a_2, a_3$  are linearly independent and  $a_4 = a_1 - a_2 + a_3$ . Which of the following is a basis of  $\text{Nul}(A)$ ?

(A)  $\begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$

(B) None of the other answers

(C)  $a_1, a_2, a_3$

(D)  $a_4$

(E)  $\begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix}$

12. (5 points) Consider the linear transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  for which

$$T\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} \frac{x+y}{2} \\ \frac{x+y}{2} \\ z \end{bmatrix}.$$

What is the matrix  $T_{\mathcal{B}\mathcal{B}}$  that represents  $T$  with respect to the basis

$$\mathcal{B} := \left( \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)$$

(A)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(B)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(C)  $\begin{bmatrix} \frac{1}{2} & -\frac{1}{2} & 0 \\ -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(D) None of the other answers.

(E)  $\begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}$

13. (5 points) Let  $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4$  be 4 row vectors of length 4 (so that the transpose  $\mathbf{a}_i^T$  belongs to  $\mathbb{R}^4$ ). Let

$$A = \begin{bmatrix} - & \mathbf{a}_1 & - \\ - & \mathbf{a}_2 & - \\ - & \mathbf{a}_3 & - \\ - & \mathbf{a}_4 & - \end{bmatrix}$$

be the  $4 \times 4$  matrix with these vectors as rows. Assume  $\det(A) = 3$ . What is the determinant of the matrix

$$B = \begin{bmatrix} -\mathbf{a}_1 \\ 3\mathbf{a}_2 - 2\mathbf{a}_3 \\ 3\mathbf{a}_3 + \mathbf{a}_4 \\ \mathbf{a}_3 \end{bmatrix}$$

(A) None of the other answers is correct.

(B)  $\det(B) = 9$ .

(C)  $\det(B) = 27$ .

(D)  $\det(B) = -9$ .

(E)  $\det(B) = -27$ .

14. (5 points) Let  $A$  be an  $n \times n$  matrix, and let  $\mathbf{v}_1, \mathbf{v}_2$  in  $\mathbb{R}^n$  be such that  $\mathbf{v}_1$  is an eigenvector of  $A$  to eigenvalue  $\lambda_1$  and  $\mathbf{v}_2$  is an eigenvector of  $A$  to eigenvalue  $\lambda_2$ . Consider the following two statements:

(T1) If  $\lambda_1 \neq \lambda_2$ , then  $\mathbf{v}_1, \mathbf{v}_2$  are linearly independent.

(T2) If  $\lambda_1 \neq \lambda_2$ , then  $\mathbf{v}_1, \mathbf{v}_2$  are orthogonal to each other.

Are these statements always correct?

(A) Neither Statement T1 nor Statement T2 is correct.

(B) Only Statement T2 is correct.

(C) Statement T1 and Statement T2 are correct.

(D) Only Statement T1 is correct.

15. (5 points) Consider the following bases:

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ of } \mathbb{R}^2, \text{ and } \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \text{ of } \mathbb{R}^3.$$

With respect to these bases the linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  is represented by the matrix

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 0 & 1 \end{bmatrix}.$$

What is  $T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right)$ ?

(A)  $\begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$

(B)  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

(C) none of the other answers

(D)  $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

(E)  $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$

16. (5 points) Let  $\mathbb{P}_2$  be the vector space of polynomials of degree at most 2. Consider the following sets of polynomials in  $\mathbb{P}_2$ .

$$\mathcal{A} = \{1, t, t + 1\}$$

$$\mathcal{B} = \{1, t, t^2\}$$

$$\mathcal{C} = \{1, 2t, t^2, t^2 - t\}.$$

Which of these sets are linearly independent?

(A)  $\mathcal{A}$  only.

(B)  $\mathcal{B}$  only.

(C)  $\mathcal{A}, \mathcal{B}$  and  $\mathcal{C}$ .

(D)  $\mathcal{B}$  and  $\mathcal{C}$  only.

(E) None of them.

17. (5 points) Let  $A$  be  $2 \times 2$ -matrix with eigenvalues  $\frac{1}{2}$  and  $\frac{1}{3}$ . What can you say about  $\lim_{k \rightarrow \infty} A^k$ ?

(A)  $\lim_{k \rightarrow \infty} A^k$  does not exist.

(B) Not enough information to say anything

(C)  $\lim_{k \rightarrow \infty} A^k = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(D)  $\lim_{k \rightarrow \infty} A^k = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

(E)  $\lim_{k \rightarrow \infty} A^k = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{3} \end{bmatrix}$

18. (5 points) Let  $A$  be a  $3 \times 5$ -matrix such that

$$\text{Col}(A) = \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 3 \end{bmatrix} \right\}.$$

Then the dimension of  $\text{Nul}(A)$  is

- (A) 4
- (B) 3
- (C) 2
- (D) 1
- (E) 0

19. (5 points) Consider the following two subsets of the vector space of  $2 \times 2$  matrices.

$$V_0 = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a = 0 \right\}, \quad V_1 = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : 0 \text{ is an eigenvalue of } \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right\}$$

Then:

- (A)  $V_0$  and  $V_1$  are subspaces.
- (B) Only  $V_0$  is a subspace.
- (C) Neither  $V_0$  nor  $V_1$  is a subspace.
- (D) Only  $V_1$  is a subspace.



20. (5 points) Consider the following basis  $\mathcal{B}$  of  $\mathbb{R}^3$ :

$$\left( \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right).$$

Let  $x = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ . What is the coordinate vector  $x_{\mathcal{B}}$  of  $x$  with respect to the basis  $\mathcal{B}$ ?

(A)  $\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

(B) None of the other answers

(C)  $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

(D)  $\begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

(E)  $\begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$

21. (5 points) Let  $\mathbb{P}^2$  be the vector space of all polynomials of degree at most 2. Let  $L$  be the linear transformation from  $\mathbb{P}^2$  to  $\mathbb{P}^2$  given by

$$L(p(t)) = 3p'(t) + 2p(t).$$

Then the matrix that represents  $L$  with respect to the bases  $\{1, t, t^2\}$  and  $\{1, t, t^2\}$  is

(A)  $\begin{bmatrix} 3 & 0 & 0 \\ 2 & 3 & 0 \\ 0 & 4 & 3 \end{bmatrix}$

(B)  $\begin{bmatrix} 2 & 0 & 0 \\ 3 & 2 & 0 \\ 0 & 6 & 2 \end{bmatrix}.$

(C)  $\begin{bmatrix} 3 & 2 & 0 \\ 0 & 3 & 4 \\ 0 & 0 & 3 \end{bmatrix}$

(D)  $\begin{bmatrix} 2 & 3 & 0 \\ 0 & 2 & 6 \\ 0 & 0 & 2 \end{bmatrix}$

22. (5 points) For which value of  $t$  is the vector  $\begin{bmatrix} 3 \\ 1+t \end{bmatrix}$  in the span of the vectors  $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$  and  $\begin{bmatrix} -2 \\ 2 \end{bmatrix}$ ?

- (A) Only for  $t = -3$ .
- (B) For no  $t$ .
- (C) Only for  $t = 2$ .
- (D) For all  $t$ .
- (E) Only for  $t = -4$ .

23. (5 points) Let  $V$  be the subset of all  $2 \times 2$ -matrices  $A$  such that  $A^T = -A$ . This is a subspace of the vector space of all  $2 \times 2$ -matrices. What is the dimension of  $V$ ?

- (A) 1
- (B) 0
- (C) 2
- (D) 4
- (E) 3

24. (5 points) A financial company has assets in countries  $A, B$  and  $C$ . Each year half of the money invested in country  $A$  stays in country  $A$ , and a quarter of the money invested in country  $A$  goes to country  $B$  and  $C$  each. For country  $B$  and  $C$ , one half of the money stays in each country and the other half is invested in country  $A$ . In the steady state, what is the percentage of the assets of the company that are invested in country  $A$ ?

- (A) None of the other answers
- (B) 25%
- (C) 50%
- (D) 100%

25. (5 points) Let  $A$  be a  $4 \times 3$ -matrix and  $\mathbf{b}$  in  $\mathbb{R}^4$ . Consider the following two statements:

- (S1) The equation  $A\mathbf{x} = \mathbf{b}$  is consistent.
- (S2) If  $A\mathbf{x} = \mathbf{b}$  is consistent, the solution to  $A\mathbf{x} = \mathbf{b}$  is unique.

Which of the two statements is correct for all such  $A$  and  $\mathbf{b}$ ?

- (A) Neither Statement S1 nor Statement S2.
- (B) Only Statement S2.
- (C) Both Statement S1 and Statement S2.
- (D) Only Statement S1.

26. (5 points) Consider the following two statements:

- (T1) If  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  are three orthonormal vectors, then the projection of  $\mathbf{v}_3$  onto the span of  $\mathbf{v}_1, \mathbf{v}_2$  is  $\mathbf{v}_3$ .
- (T2) The Gram–Schmidt process produces from a linearly independent set  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  an orthonormal set  $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$  with the property that for each  $k \leq n$  the vectors  $\{\mathbf{q}_1, \dots, \mathbf{q}_k\}$  span the same subspace as  $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ .

Then:

- (A) Neither Statement T1 nor Statement T2 is correct.
- (B) Only Statement T1 is correct.
- (C) Only Statement T2 is correct.
- (D) Statement T1 and Statement T2 are correct.

27. (5 points) Let  $W$  be the  $\text{Span}\left\{\begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}\right\}$ , and let  $y = \begin{bmatrix} 2 \\ 4 \\ 2 \\ 1 \end{bmatrix}$ . Let  $y_W$  be in  $W$  and  $y_{W^\perp}$  in  $\mathbb{R}^4$  be orthogonal to  $W$  such that  $y = y_W + y_{W^\perp}$ . Then

(A)  $y_W = \begin{bmatrix} 0 \\ 2 \\ 0 \\ 1 \end{bmatrix}$  and  $y_{W^\perp} = \begin{bmatrix} 2 \\ 2 \\ 2 \\ 0 \end{bmatrix}$

(B)  $y_W = \begin{bmatrix} 0 \\ 2.5 \\ 2 \\ 2.5 \end{bmatrix}$  and  $y_{W^\perp} = \begin{bmatrix} 2 \\ 1.5 \\ 0 \\ -1.5 \end{bmatrix}$

(C) None of the other answers

(D)  $y_W = \begin{bmatrix} 0 \\ 2 \\ 2 \\ 2 \end{bmatrix}$  and  $y_{W^\perp} = \begin{bmatrix} 2 \\ 2 \\ 0 \\ -2 \end{bmatrix}$

28. (5 points) Let  $A = \begin{bmatrix} 0 & 2 & -1 \\ -2 & 8 & -5 \\ -3 & 10 & -7 \end{bmatrix}$ ,  $\mathbf{v} = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}$  and  $\mathbf{w} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$ . Then:

- (A)  $\mathbf{w}$  is an eigenvector of  $A$  for the eigenvalue  $-2$ ,
- (B)  $\mathbf{v}$  is an eigenvector of  $A$  for the eigenvalue  $2$ ,
- (C)  $\mathbf{v}$  is an eigenvector of  $A$  for the eigenvalue  $-2$ ,
- (D) None of the other answers.
- (E)  $\mathbf{w}$  is an eigenvector of  $A$  for the eigenvalue  $2$ ,

29. (5 points) Let  $A$  be an  $m \times n$ -matrix with echelon form  $U$ . Which of the following statement is true for all such  $A$ ?

- (T1)  $\text{Nul}(A) = \text{Nul}(U)$ .
- (T2)  $\text{Col}(A^T) = \text{Col}(U^T)$ .
- (A) Only (T2).
- (B) Both (T1) and (T2)
- (C) Neither (T1) nor (T2).
- (D) Only (T1).

30. (5 points) Consider the following two statements:

(S1) There exists a subspace  $V$  of  $\mathbb{R}^7$  such that  $\dim V = \dim V^\perp$ .

(S2) If  $V$  is a subspace of  $\mathbb{R}^7$ , then the zero vector is the only vector which is in  $V$  as well as in  $V^\perp$ .

Then:

(A) Only Statement S2 is correct.

(B) Statement S1 and Statement S2 are correct.

(C) Neither Statement S1 nor Statement S2 is correct.

(D) Only Statement S1 is correct.



31. (5 points) Let  $B = \begin{bmatrix} 0 & 1 \\ 3 & 1 \\ 0 & 1 \end{bmatrix}$ . Let  $Q$  be a  $3 \times 2$ -matrix with orthonormal columns and  $R$  be an upper triangular matrix such that  $B = QR$ . Which of the following matrices can be  $R$ ?

(A)  $\begin{bmatrix} 3 & 1 \\ 0 & \sqrt{2} \end{bmatrix}$

(B)  $\begin{bmatrix} 1 & \sqrt{3} \\ 0 & 2 \end{bmatrix}$

(C)  $\begin{bmatrix} 3 & 2 \\ 0 & \sqrt{2} \end{bmatrix}$

(D)  $\begin{bmatrix} \sqrt{3} & 2 \\ 0 & 2 \end{bmatrix}$

(E) None of the other answers