Solve $x^3 = 27$.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 52 • BACKWARD 2 • Q1/5

Evaluate $\sqrt[3]{125}$.

Value: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 52 • BACKWARD 2 • Q2/5

What is $\sqrt[3]{64}$?

A. 21.3

B. 4

C. 8

D. 16

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 52 • BACKWARD 2 • Q3/5

True or false?

$\sqrt[3]{(-27)} = -3$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 52 • BACKWARD 2 • Q4/5

True or false?

$\sqrt{16} = 4$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 52 • BACKWARD 2 • Q5/5

SET 52 • HSF-BF.B.4.a

FORWARD 1

Verify inverses by composition • High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.b

Show that $f(x) = 2x^3$ and $g(x) = \sqrt[3]{(x \div 2)}$ are inverses.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q1

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.b • Set 52 • FORWARD 1 • Q1/5

$f(x) = (x + 1) \div (x - 1)$.

Find $f(f(3))$.

$f(f(3)) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q2

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.b • Set 52 • FORWARD 1 • Q2/5

Which function is its own inverse?

A. $f(x) = 1 \div x$

B. $f(x) = 2x$

C. $f(x) = x + 1$

D. $f(x) = x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S52-F1-Q3**

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.b • Set 52 • FORWARD 1 • Q3/5

True or false?

If f(g(x)) = x on the domain of g and g(f(x)) = x on the domain of f, then g is the inverse of f.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q4

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.b • Set 52 • FORWARD 1 • Q4/5

True or false?

$f(x) = x + 3$ and $g(x) = x - 3$ are not inverses, because $f(g(1)) = 7$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q5

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.b • Set 52 • FORWARD 1 • Q5/5

SET 52 • HSF-BF.B.4.a

FORWARD 2

Restrict a domain to produce an inverse • High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.d

Restrict the domain of $f(x) = x^2 + 2$ so that f has an inverse, and give the inverse.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q1

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.d • Set 52 • FORWARD 2 • Q1/5

Explain why f(x) = |x| has no inverse on all real numbers.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q2

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.d • Set 52 • FORWARD 2 • Q2/5

$f(x) = (x + 4)^2$.

What is the largest domain that contains $x = 0$ on which f is one-to-one?

A. All real numbers

B. $-8 \leq x \leq 0$

C. $x \geq 0$

D. $x \geq -4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S52-F2-Q3**

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.d • Set 52 • FORWARD 2 • Q3/5

True or false?

$f(x) = x^2$ restricted to $x \leq 0$ has the inverse $f^{-1}(x) = -\sqrt{x}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q4

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.d • Set 52 • FORWARD 2 • Q4/5

True or false?

$f(x) = x^2 + 1$ is one-to-one on all real numbers.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q5

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.4.d • Set 52 • FORWARD 2 • Q5/5

Prove that linear functions grow by equal differences and exponential functions grow by equal factors over equal intervals

MAIN	Algebra I main question	**HSF-LE.A.1.a**
BACKWARD 1	Tell linear from nonlinear functions	**8.F.A.3**
BACKWARD 2	Compare average rates of change over intervals	**HSF-IF.B.6**
FORWARD 1	Derive the formula for the sum of a finite geometric series	**HSA-SSE.B.4**
FORWARD 2	Use logarithms to turn equal factors into equal differences	**HSF-BF.B.5**

SET 53 • HSF-LE.A.1.a

MAIN

Algebra I main question • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a

$f(x) = 3x + 5$. Show that f grows by equal differences over equal intervals: find $f(x + h) - f(x)$ in simplest form, and explain why it does not depend on x.

$f(x + h) - f(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 53 • MAIN • Q1/5

$g(x) = 4 \cdot 2^x$. Show that g grows by equal factors over equal intervals: find $g(x + h) \div g(x)$ in simplest form, and explain why it does not depend on x.

$g(x + h) \div g(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 53 • MAIN • Q2/5

A table shows x = 0, 2, 4, 6 and y = 5, 11, 17, 23.

Is the relationship linear or exponential? Explain using equal intervals.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 53 • MAIN • Q3/5

A table shows x = 0, 1, 2, 3 and y = 80, 40, 20, 10.

Which description is correct?

A. Linear: y decreases by 40 each time

B. Exponential: y is multiplied by $\frac{1}{2}$ each time x increases by 1

C. Linear: y decreases by 20 each time

D. Neither linear nor exponential

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 53 • MAIN • Q4/5

True or false?

A table with y-values 2, 5, 10, 17 for x = 0, 1, 2, 3 is linear.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 53 • MAIN • Q5/5

SET 53 • HSF-LE.A.1.a

BACKWARD 1

Tell linear from nonlinear functions • Grade 8 • Functions • 8.F.A.3

Is $y = x^2 + 1$ linear? Use the points (0, 1), (1, 2), and (2, 5) to explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q1

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 1 • Q1/5

Is $y = 4 - 2x$ linear? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q2

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 1 • Q2/5

Which function is linear?

A. $y = 2^x$

B. $y = x^2$

C. $y = 0.5x - 3$

D. $y = 1 \div x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q3

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 1 • Q3/5

True or false?

The graph of a linear function is a straight line.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q4

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 1 • Q4/5

True or false?

The area of a square, $A = s^2$, is a linear function of s.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q5

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 1 • Q5/5

SET 53 • HSF-LE.A.1.a

BACKWARD 2

Compare average rates of change over intervals • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6

$f(x) = 3x + 5$.

Find the average rate of change from $x = 0$ to $x = 10$.

Rate: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 53 • BACKWARD 2 • Q1/5

$g(x) = 4 \cdot 2^x$.

Find the average rate of change from $x = 0$ to $x = 1$, and from $x = 1$ to $x = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 53 • BACKWARD 2 • Q2/5

What is the average rate of change of $5 \cdot 3^x$ from $x = 0$ to $x = 2$?

A. 15

B. 20

C. 40

D. 45

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 53 • BACKWARD 2 • Q3/5

True or false?

An exponential growth function has a larger average rate of change on later intervals of the same length.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 53 • BACKWARD 2 • Q4/5

True or false?

The average rate of change of 2^x from $x = 0$ to $x = 3$ is 2.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 53 • BACKWARD 2 • Q5/5

SET 53 • HSF-LE.A.1.a

FORWARD 1

Derive the formula for the sum of a finite geometric series • High School — Algebra II (course placement in this bank) • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

Let $S = 1 + 2 + 4 + \ldots + 2^9$. Show that $2S - S = 2^{10} - 1$, and find S.

$S =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q1

High School — Algebra II (course placement in this bank) • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 53 • FORWARD 1 • Q1/5

Use $S = a(1 - r^n) \div (1 - r)$ to find the sum of the first 6 terms of 1, 3, 9, ...

Sum: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q2

High School — Algebra II (course placement in this bank) • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 53 • FORWARD 1 • Q2/5

When deriving the sum formula, why do we multiply S by r?

A. To double the sum

B. So that $S - rS$ cancels all but two terms

C. To make every term equal

D. To make r equal 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S53-F1-Q3**

High School — Algebra II (course placement in this bank) • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 53 • FORWARD 1 • Q3/5

True or false?

For $r \neq 1$, $a + ar + \ldots + ar^{n-1} = a(1 - r^n) \div (1 - r)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q4

High School — Algebra II (course placement in this bank) • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 53 • FORWARD 1 • Q4/5

True or false?

$1 + 3 + 9 + 27 = 3^4$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q5

High School — Algebra II (course placement in this bank) • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 53 • FORWARD 1 • Q5/5

SET 53 • HSF-LE.A.1.a

FORWARD 2

Use logarithms to turn equal factors into equal differences • High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5

$g(x) = 4 \cdot 2^x$.

Show that $\log_2 g(x) = x + 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q1

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 53 • FORWARD 2 • Q1/5

$y = 10^x$.

What is log y?

log y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q2

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 53 • FORWARD 2 • Q2/5

The values y = 1, 10, 100, 1,000 occur at x = 0, 1, 2, 3.

What are the values of log y?

A. 0, 1, 2, 3

B. 0, 10, 20, 30

C. 1, 2, 3, 4

D. 1, 10, 100, 1,000

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q3

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 53 • FORWARD 2 • Q3/5

True or false?

Taking the logarithm of an exponential function $a \cdot b^x$ gives a linear function of x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q4

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 53 • FORWARD 2 • Q4/5

True or false?

$$\log(100 \cdot 10^{x}) = 100x$$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q5

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 53 • FORWARD 2 • Q5/5

Recognize situations in which a quantity changes at a constant rate per unit interval

MAIN	Algebra I main question	**HSF-LE.A.1.b**
BACKWARD 1	Find the constant of proportionality	**7.RP.A.2.b**
BACKWARD 2	Write a linear function from a rate and an initial value	**8.F.B.4**
FORWARD 1	Relate arc length and sector area to the central angle	**HSG-C.B.5**
FORWARD 2	Use radian measure as arc length on a circle	**HSF-TF.A.1**

SET 54 • HSF-LE.A.1.b

MAIN

Algebra I main question • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.b

A pool fills at a constant rate. At 2 p.m. it holds 1,500 gallons; at 5 p.m. it holds 3,300 gallons.

What is the rate, and how much water was in it at noon?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.b • Set 54 • MAIN • Q1/5

A plant is 6 cm tall on day 0 and grows 0.8 cm each day.

How tall is it on day 15?

Height: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.b • Set 54 • MAIN • Q2/5

A taxi charges the same amount for every mile.

Explain why the fare for the miles driven is a linear function of the miles.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.b • Set 54 • MAIN • Q3/5

Which situation is modeled by a linear function?

A. A cell population doubles each hour.

B. A tank loses 4 liters each minute.

C. A loan grows 4% each month.

D. A car loses 15% of its value each year.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.b • Set 54 • MAIN • Q4/5

True or false?

A savings account that earns the same dollar amount of interest each year grows linearly.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.b • Set 54 • MAIN • Q5/5

SET 54 • HSF-LE.A.1.b

BACKWARD 1

Find the constant of proportionality • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2.b

$y = 2.5x$ gives the cost of x pounds of apples.

What is the constant of proportionality?

Constant: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2.b • Set 54 • BACKWARD 1 • Q1/5

A proportional table has x = 2, 4, 6 and y = 7, 14, 21.

What is the unit rate?

Unit rate: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2.b • Set 54 • BACKWARD 1 • Q2/5

A cyclist rides 12 miles in 3 hours at a constant rate.

What is the constant of proportionality in miles per hour?

A. 15

B. 36

C. $\frac{1}{4}$

D. 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2.b • Set 54 • BACKWARD 1 • Q3/5

True or false?

In y = 9x, y increases by 9 whenever x increases by 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2.b • Set 54 • BACKWARD 1 • Q4/5

True or false?

A proportional relationship through (5, 20) has constant of proportionality 100.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2.b • Set 54 • BACKWARD 1 • Q5/5

SET 54 • HSF-LE.A.1.b

BACKWARD 2

Write a linear function from a rate and an initial value • Grade 8 • Functions • 8.F.B.4

A line passes through (2, 1,500) and (5, 3,300).

Find its slope and its y-intercept.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q1

Grade 8 • Functions • 8.F.B.4 • Set 54 • BACKWARD 2 • Q1/5

A phone battery is at 80% and drops 5 percentage points per hour.

Write a function B(t) for the charge after t hours.

B(t) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q2

Grade 8 • Functions • 8.F.B.4 • Set 54 • BACKWARD 2 • Q2/5

Which function has rate of change 0.8 and initial value 6?

A. $y = 6x + 0.8$

B. $y = 0.8x + 6$

C. $y = 6.8x$

D. $y = 6(0.8)^x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q3

Grade 8 • Functions • 8.F.B.4 • Set 54 • BACKWARD 2 • Q3/5

True or false?

A function with initial value 10 and rate of change -2 is $y = 10 - 2x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q4

Grade 8 • Functions • 8.F.B.4 • Set 54 • BACKWARD 2 • Q4/5

True or false?

The line through (0, 4) and (2, 10) has rate of change 5.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q5

Grade 8 • Functions • 8.F.B.4 • Set 54 • BACKWARD 2 • Q5/5

SET 54 • HSF-LE.A.1.b

FORWARD 1

Relate arc length and sector area to the central angle • High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5

In a circle of radius 5 cm, by how much does the arc length increase for each 1-radian increase in the central angle?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q1

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 54 • FORWARD 1 • Q1/5

A circle has radius 10.

Find the arc lengths for central angles of 0.5, 1, and 1.5 radians.

Arc lengths: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q2

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 54 • FORWARD 1 • Q2/5

The area of a sector is $A = \frac{1}{2}r^2\theta$, and $r = 4$.

By how much does A increase when θ increases by 1 radian?

A. 4

B. 2

C. 16

D. 8

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q3

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 54 • FORWARD 1 • Q3/5

True or false?

For a fixed radius, arc length is a linear function of the central angle.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q4

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 54 • FORWARD 1 • Q4/5

True or false?

Doubling the radius while keeping the angle the same doubles the area of a sector.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q5

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 54 • FORWARD 1 • Q5/5

SET 54 • HSF-LE.A.1.b

FORWARD 2

Use radian measure as arc length on a circle • High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.A.1

Convert 180° to radians.

Radians: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q1

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 54 • FORWARD 2 • Q1/5

A central angle cuts off an arc of length 6 on a circle of radius 2.

What is its radian measure?

Radians: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q2

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 54 • FORWARD 2 • Q2/5

Convert $\frac{\pi}{3}$ radians to degrees.

A. 30°

B. 90°

C. 120°

D. 60°

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-F2-Q3**

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 54 • FORWARD 2 • Q3/5

True or false?

One radian is the angle that cuts off an arc as long as the radius.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q4

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 54 • FORWARD 2 • Q4/5

True or false?

2π radians equals 180°.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q5

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 54 • FORWARD 2 • Q5/5

Recognize situations in which a quantity grows or decays by a constant percent rate per unit interval

MAIN	Algebra I main question	**HSF-LE.A.1.c**
BACKWARD 1	Read growth and decay rates from exponential functions	**HSF-IF.C.8.b**
BACKWARD 2	Solve percent problems	**7.RP.A.3**
FORWARD 1	Find times in percent-rate models with logarithms	**HSF-LE.A.4**
FORWARD 2	Use natural logarithms with continuous growth	**HSF-BF.B.5**

SET 55 • HSF-LE.A.1.c

MAIN

Algebra I main question • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.c

A car worth $24,000 loses 15% of its value each year.

Write a function V(t), and find the value after 3 years to the nearest dollar.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.c • Set 55 • MAIN • Q1/5

Explain why a quantity that grows by 6% each year grows by equal factors over equal intervals.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.c • Set 55 • MAIN • Q2/5

A town's population increases by 2.5% per year.

By what factor does it grow in 2 years? What is the percent increase over 2 years?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.c • Set 55 • MAIN • Q3/5

Which situation is exponential?

A. A worker earns $15 each hour.

B. A medicine dose decreases by 20% each hour.

C. A tank gains 3 liters each second.

D. A runner covers 200 meters each minute.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.c • Set 55 • MAIN • Q4/5

True or false?

A 10% increase followed by a 10% decrease returns a quantity to its original value.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.c • Set 55 • MAIN • Q5/5

SET 55 • HSF-LE.A.1.c

BACKWARD 1

Read growth and decay rates from exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b

$V = 24{,}000(0.85)^t$.

What is the yearly percent change?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 55 • BACKWARD 1 • Q1/5

$P = 500(1.025)^t$.

Is this growth or decay, and at what rate?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 55 • BACKWARD 1 • Q2/5

Which function shows 4% decay per year?

A. $y = 300(1.04)^t$

B. $y = 300(0.04)^t$

C. $y = 300 - 0.04t$

D. $y = 300(0.96)^t$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 55 • BACKWARD 1 • Q3/5

True or false?

$y = 50(1.5)^t$ grows by 50% per unit of t.

A. True

B. False

True or false?

$y = 80(0.75)^t$ decreases by 75% per unit of t.

A. True

B. False

SET 55 • HSF-LE.A.1.c

BACKWARD 2

Solve percent problems • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

A $24,000 car loses 15% of its value.

What is its new value?

Value: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 55 • BACKWARD 2 • Q1/5

What is 8% sales tax on $45?

Tax: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 55 • BACKWARD 2 • Q2/5

What is the price of a $60 item after a 25% discount?

A. $15

B. $45

C. $75

D. $35

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 55 • BACKWARD 2 • Q3/5

True or false?

20% of 150 is 30.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 55 • BACKWARD 2 • Q4/5

True or false?

A 50% increase of 40 is 60, and a 50% decrease of 60 is 40.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 55 • BACKWARD 2 • Q5/5

SET 55 • HSF-LE.A.1.c

FORWARD 1

Find times in percent-rate models with logarithms • High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

$V(t) = 24{,}000(0.85)^t$.

When is V = 12,000? Round to the nearest tenth of a year.

t ≈ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q1

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 55 • FORWARD 1 • Q1/5

Solve $1.025^t = 2$. Round to the nearest tenth.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q2

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 55 • FORWARD 1 • Q2/5

Which expression gives the half-life of $y = (0.9)^t$?

A. $0.9 \div 2$

B. $0.5 \div 0.9$

C. $\log 0.9 \div \log 0.5$

D. $\log 0.5 \div \log 0.9$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S55-F1-Q3**

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 55 • FORWARD 1 • Q3/5

True or false?

$\log(0.85^t) = t \log 0.85$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q4

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 55 • FORWARD 1 • Q4/5

True or false?

If $2^t = 5$, then $t = 2.5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q5

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 55 • FORWARD 1 • Q5/5

SET 55 • HSF-LE.A.1.c

FORWARD 2

Use natural logarithms with continuous growth • High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5

Write the inverse of $y = e^x$.

Inverse: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q1

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 55 • FORWARD 2 • Q1/5

Solve $e^{0.05t} = 3$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q2

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 55 • FORWARD 2 • Q2/5

What is $\ln(e^4)$?

A. 4e

B. 4

C. 1

D. e

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S55-F2-Q3**

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 55 • FORWARD 2 • Q3/5

True or false?

e raised to the power ln 7 equals 7.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q4

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 55 • FORWARD 2 • Q4/5

True or false?

$\ln 1 = 1$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q5

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 55 • FORWARD 2 • Q5/5

Construct linear and exponential functions from a graph, a description, two points, or a table

MAIN	Algebra I main question	**HSF-LE.A.2**
BACKWARD 1	Write a linear function from two points or a table	**8.F.B.4**
BACKWARD 2	Write formulas for arithmetic and geometric sequences	**HSF-BF.A.2**
FORWARD 1	Build a sinusoidal model from a description	**HSF-TF.B.5**
FORWARD 2	Find a quadratic through three points with a system	**HSA-REI.C.8**

SET 56 • HSF-LE.A.2

MAIN

Algebra I main question • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

Write the linear function whose graph passes through (2, 11) and (6, 23).

f(x) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 56 • MAIN • Q1/5

Write the exponential function $y = ab^x$ $(b > 0)$ whose graph passes through $(1, 6)$ and $(3, 54)$.

$y =$ _____


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 56 • MAIN • Q2/5

The graph of an exponential function $y = ab^x$ passes through the marked points.

Write the function.



$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 56 • MAIN • Q3/5

A table shows x = 0, 1, 2, 3 and y = 1.5, 3, 6, 12.

Which function fits?

A. $y = 2(1.5)^x$

B. $y = 1.5x + 1.5$

C. $y = 3x$

D. $y = 1.5(2)^x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 56 • MAIN • Q4/5

True or false?

The linear function with $f(0) = 10$ that decreases by 2 for each unit increase in x is $f(x) = 10(0.8)^x$.

A. True

B. False

SET 56 • HSF-LE.A.2

BACKWARD 1

Write a linear function from two points or a table • Grade 8 • Functions • 8.F.B.4

Write the linear function through $(0, -3)$ and $(4, 5)$.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q1

Grade 8 • Functions • 8.F.B.4 • Set 56 • BACKWARD 1 • Q1/5

A table shows x = 1, 2, 3 and y = 10, 7, 4.

Write the linear function.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q2

Grade 8 • Functions • 8.F.B.4 • Set 56 • BACKWARD 1 • Q2/5

Which line passes through (−1, 2) and (1, 6)?

A. $y = -2x + 4$

B. $y = 2x + 4$

C. $y = 2x + 2$

D. $y = 4x + 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q3

Grade 8 • Functions • 8.F.B.4 • Set 56 • BACKWARD 1 • Q3/5

True or false?

The line through (0, 0) and (3, 6) is $y = 2x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q4

Grade 8 • Functions • 8.F.B.4 • Set 56 • BACKWARD 1 • Q4/5

True or false?

The line through (1, 1) and (2, 3) is $y = 2x + 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q5

Grade 8 • Functions • 8.F.B.4 • Set 56 • BACKWARD 1 • Q5/5

SET 56 • HSF-LE.A.2

BACKWARD 2

Write formulas for arithmetic and geometric sequences • High School — Algebra I • Functions: Building Functions • HSF-BF.A.2

A geometric sequence has a(1) = 2 and ratio 3.

Write an explicit formula.

a(n) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.A.2 • Set 56 • BACKWARD 2 • Q1/5

Write an explicit formula for 10, 7, 4, ...

a(n) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.A.2 • Set 56 • BACKWARD 2 • Q2/5

Which is a recursive formula for 6, 3, 1.5, ...?

A. $a(n) = 6 - 3n$

B. $a(1) = 6, a(n) = 0.5a(n - 1)$

C. $a(1) = 6, a(n) = 2a(n - 1)$

D. $a(1) = 6, a(n) = a(n - 1) - 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.A.2 • Set 56 • BACKWARD 2 • Q3/5

True or false?

An arithmetic sequence is a linear function on the positive integers.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.A.2 • Set 56 • BACKWARD 2 • Q4/5

True or false?

The geometric sequence 5, 10, 20, ... has ratio 5.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.A.2 • Set 56 • BACKWARD 2 • Q5/5

SET 56 • HSF-LE.A.2

FORWARD 1

Build a sinusoidal model from a description • High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5

A tide has high water 10 ft and low water 2 ft, with period 12 hours and high tide at t = 0.

Write a cosine model h(t).

h(t) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q1

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • FORWARD 1 • Q1/5

A sinusoid has maximum 10 and minimum 2.

What are its amplitude and midline?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q2

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • FORWARD 1 • Q2/5

A model $y = \cos(Bt)$ has period 12.

What is B?

A. 12

B. $\pi \div 12$

C. $\pi \div 6$

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q3

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • FORWARD 1 • Q3/5

True or false?

A sinusoidal model with maximum 8 and minimum 0 has amplitude 4.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q4

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • FORWARD 1 • Q4/5

True or false?

$y = 3\cos(2t)$ has period 2.

A. True

B. False

SET 56 • HSF-LE.A.2

FORWARD 2

Find a quadratic through three points with a system • High School — Precalculus (course placement in this bank) • Algebra: Reasoning with Equations & Inequalities • HSA-REI.C.8

$y = ax^2 + bx + c$ passes through (0, 1), (1, 2), and (2, 7).

Write the system of equations for a, b, and c.

System: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q1

High School — Precalculus (course placement in this bank) • Algebra: Reasoning with Equations & Inequalities • HSA-REI.C.8 • Set 56 • FORWARD 2 • Q1/5

Solve that system.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q2

High School — Precalculus (course placement in this bank) • Algebra: Reasoning with Equations & Inequalities • HSA-REI.C.8 • Set 56 • FORWARD 2 • Q2/5

Which row of the augmented matrix comes from the point (2, 7)?

A. [4 2 1 | 7]

B. [2 4 1 | 7]

C. [2 2 1 | 7]

D. [4 2 7 | 1]

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S56-F2-Q3**

High School — Precalculus (course placement in this bank) • Algebra: Reasoning with Equations & Inequalities • HSA-REI.C.8 • Set 56 • FORWARD 2 • Q3/5

True or false?

Three points with different x-values that are not on one line determine exactly one parabola $y = ax^2 + bx + c$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q4

High School — Precalculus (course placement in this bank) • Algebra: Reasoning with Equations & Inequalities • HSA-REI.C.8 • Set 56 • FORWARD 2 • Q4/5

True or false?

The coefficient matrix of that system has 2 rows.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q5

High School — Precalculus (course placement in this bank) • Algebra: Reasoning with Equations & Inequalities • HSA-REI.C.8 • Set 56 • FORWARD 2 • Q5/5

Observe that exponential growth eventually exceeds linear, quadratic, and polynomial growth

MAIN	Algebra I main question	**HSF-LE.A.3**
BACKWARD 1	Tell equal differences from equal factors	**HSF-LE.A.1.a**
BACKWARD 2	Compare two functions given in different forms	**8.F.A.2**
FORWARD 1	Compare the growth of polynomial functions	**HSF-IF.C.7.c**
FORWARD 2	Compare logarithmic growth with other growth	**HSF-BF.B.5**

SET 57 • HSF-LE.A.3

MAIN

Algebra I main question • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.3

$f(x) = 100x$ and $g(x) = 2^x$.

What is the smallest whole number $x \geq 1$ for which $g(x) > f(x)$?

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.3 • Set 57 • MAIN • Q1/5

$f(x) = x^2$ and $g(x) = 1.5^x$.

Which is larger at $x = 10$? At $x = 20$? What happens for larger x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.3 • Set 57 • MAIN • Q2/5

Plan A pays $1,000 per day. Plan B pays 1 cent on day 1, and its daily pay doubles each day.

On which day does Plan B's daily pay first exceed Plan A's?

Day: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.3 • Set 57 • MAIN • Q3/5

Which function eventually has the greatest values as x increases?

A. $y = 50x^2$

B. $y = 1.01^x$

C. $y = x^3 + 500$

D. $y = 1{,}000x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.3 • Set 57 • MAIN • Q4/5

True or false?

Because $3x^2 > 2^x$ at $x = 5$, $3x^2$ stays greater than 2^x for all $x > 5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.3 • Set 57 • MAIN • Q5/5

SET 57 • HSF-LE.A.3

BACKWARD 1

Tell equal differences from equal factors • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a

A table shows x = 0, 1, 2, 3 and y = 3, 6, 12, 24.

Is it linear or exponential? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 57 • BACKWARD 1 • Q1/5

A table shows x = 0, 1, 2, 3 and y = 3, 6, 9, 12.

Is it linear or exponential? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 57 • BACKWARD 1 • Q2/5

Which function grows by equal factors over equal intervals?

A. $y = 7 + 1.1x$

B. $y = 7(1.1)^x$

C. $y = 7x$

D. $y = 1.1x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 57 • BACKWARD 1 • Q3/5

True or false?

Linear functions grow by equal differences over equal intervals.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 57 • BACKWARD 1 • Q4/5

True or false?

$y = 5x^2$ grows by equal factors over equal intervals.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.1.a • Set 57 • BACKWARD 1 • Q5/5

SET 57 • HSF-LE.A.3

BACKWARD 2

Compare two functions given in different forms • Grade 8 • Functions • 8.F.A.2

Function A is $y = 4x + 1$. Function B has $x = 0, 1, 2$ and $y = 3, 5, 7$.

Which has the greater rate of change?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q1

Grade 8 • Functions • 8.F.A.2 • Set 57 • BACKWARD 2 • Q1/5

Function A is $y = -x + 6$. Function B is the line through $(0, 2)$ and $(2, 8)$.

Which has the greater y-intercept?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q2

Grade 8 • Functions • 8.F.A.2 • Set 57 • BACKWARD 2 • Q2/5

$f(x) = 3x - 2$. Function g has $g(0) = 1$ and $g(1) = 5$, and it is linear.

Which statement is true?

A. f has the greater rate of change.

B. f has the greater initial value.

C. g has the greater rate of change and the greater initial value.

D. f and g have the same rate of change.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q3

Grade 8 • Functions • 8.F.A.2 • Set 57 • BACKWARD 2 • Q3/5

True or false?

A linear function with slope 5 increases faster than one with slope 2.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q4

Grade 8 • Functions • 8.F.A.2 • Set 57 • BACKWARD 2 • Q4/5

True or false?

$y = 2x + 10$ has greater values than $y = 3x$ for every x.

A. True

B. False

SET 57 • HSF-LE.A.3

FORWARD 1

Compare the growth of polynomial functions • High School — Algebra II (course placement in this bank) • Functions: Interpreting Functions • HSF-IF.C.7.c

For which $x > 0$ is $x^3 > 100x^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q1

High School — Algebra II (course placement in this bank) • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 57 • FORWARD 1 • Q1/5

Which grows faster for large x, x^4 or $1{,}000x^3$? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q2

High School — Algebra II (course placement in this bank) • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 57 • FORWARD 1 • Q2/5

As $x \to \infty$, what does $f(x) = x^4 - 50x^3$ do?

A. $f(x) \to -\infty$

B. $f(x) \to -50$

C. $f(x) \to \infty$

D. $f(x) \to 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S57-F1-Q3**

High School — Algebra II (course placement in this bank) • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 57 • FORWARD 1 • Q3/5

True or false?

A polynomial of higher degree with a positive leading coefficient eventually exceeds any polynomial of lower degree.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q4

High School — Algebra II (course placement in this bank) • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 57 • FORWARD 1 • Q4/5

True or false?

For every large x, $0.001x^3$ is less than $1{,}000x^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q5

High School — Algebra II (course placement in this bank) • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 57 • FORWARD 1 • Q5/5

SET 57 • HSF-LE.A.3

FORWARD 2

Compare logarithmic growth with other growth • High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5

Find $\log_2 1{,}024$.

Value: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q1

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 57 • FORWARD 2 • Q1/5

By what factor must x be multiplied for log x to increase by 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q2

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 57 • FORWARD 2 • Q2/5

Which is greatest when $x = 1{,}000{,}000$?

A. $\sqrt{x}$

B. $\log_2 x$

C. $\ln x$

D. $\log x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q3

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 57 • FORWARD 2 • Q3/5

True or false?

Eventually, log x grows more slowly than any linear function with positive slope.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q4

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 57 • FORWARD 2 • Q4/5

True or false?

$\log_2 x = 100$ when $x = 200$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q5

High School — Precalculus (course placement in this bank) • Functions: Building Functions • HSF-BF.B.5 • Set 57 • FORWARD 2 • Q5/5

Interpret the parameters of a linear or exponential function in context

MAIN	Algebra I main question	**HSF-LE.B.5**
BACKWARD 1	Interpret the parts of an expression in context	**HSA-SSE.A.1.a**
BACKWARD 2	Interpret the rate and initial value of a linear function	**8.F.B.4**
FORWARD 1	Interpret the parameters of a sinusoidal model	**HSF-TF.B.5**
FORWARD 2	Interpret the mean and standard deviation of a normal model	**HSS-ID.A.4**

SET 58 • HSF-LE.B.5

MAIN

Algebra I main question • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5

$C(m) = 0.35m + 40$ is the cost in dollars of renting a car for one day and driving m miles.

What do 0.35 and 40 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 58 • MAIN • Q1/5

$P(t) = 1{,}200(1.04)^t$ is a town's population t years after 2020.

What do 1,200 and 1.04 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 58 • MAIN • Q2/5

$A(t) = 80(0.5)^{t/6}$ is the milligrams of a medicine left in the body after t hours.

What do 80, 0.5, and 6 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 58 • MAIN • Q3/5

$V(t) = 18{,}000 - 1{,}500t$ is a machine's value in dollars after t years.

What does $-1{,}500$ mean?

A. The machine loses $1,500 of value each year.

B. The machine loses 1,500% of its value each year.

C. The machine was bought for $1,500.

D. The machine is worth $18,000 after 1,500 years.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 58 • MAIN • Q4/5

True or false?

In $y = 250(0.92)^t$, the value decreases by 92% each year.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 58 • MAIN • Q5/5

SET 58 • HSF-LE.B.5

BACKWARD 1

Interpret the parts of an expression in context • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

A plumber charges 75 + 60h dollars for h hours of work.

What does 75 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 58 • BACKWARD 1 • Q1/5

In $3{,}000(1.05)^t$, what does the factor 1.05 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 58 • BACKWARD 1 • Q2/5

A box with a square base of side s and height h has surface area $2s^2 + 4sh$.

What does 4sh represent?

A. The volume

B. The area of the four side faces

C. The perimeter of the base

D. The area of the top and bottom

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 58 • BACKWARD 1 • Q3/5

True or false?

In 500 − 20w, the term −20w shows a decrease of 20 for each unit of w.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 58 • BACKWARD 1 • Q4/5

True or false?

In 1.5x + 7, the coefficient of x is 7.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 58 • BACKWARD 1 • Q5/5

SET 58 • HSF-LE.B.5

BACKWARD 2

Interpret the rate and initial value of a linear function • Grade 8 • Functions • 8.F.B.4

Cost $y = 12x + 30$ for x hours of bike rental.

What is the initial value, and what does it mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q1

Grade 8 • Functions • 8.F.B.4 • Set 58 • BACKWARD 2 • Q1/5

The water in a tank is $y = 500 - 25t$ gallons after t minutes.

What does -25 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q2

Grade 8 • Functions • 8.F.B.4 • Set 58 • BACKWARD 2 • Q2/5

A plant's height is a linear function of time. It is 8 cm on day 0 and 20 cm on day 4.

What is the rate of change?

A. 5 cm per day

B. 20 cm per day

C. 8 cm per day

D. 3 cm per day

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q3

Grade 8 • Functions • 8.F.B.4 • Set 58 • BACKWARD 2 • Q3/5

True or false?

In y = 40x + 100, 100 is the value of y when x = 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q4

Grade 8 • Functions • 8.F.B.4 • Set 58 • BACKWARD 2 • Q4/5

True or false?

In d = 65t, 65 is the starting distance.

A. True

B. False

SET 58 • HSF-LE.B.5

FORWARD 1

Interpret the parameters of a sinusoidal model • High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5

$T(t) = 15 + 10\sin(\pi t \div 12)$ is the temperature in °C t hours after 8 a.m.

What do 15 and 10 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q1

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 58 • FORWARD 1 • Q1/5

What is the period of T, and what does it mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q2

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 58 • FORWARD 1 • Q2/5

$h(t) = 1.5\cos(\pi t \div 6) + 3$ is the depth of water in meters.

What is the greatest depth?

A. 4.5 m

B. 3 m

C. 6 m

D. 1.5 m

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S58-F1-Q3**

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 58 • FORWARD 1 • Q3/5

True or false?

In $y = A \sin(Bx) + D$, D gives the midline.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q4

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 58 • FORWARD 1 • Q4/5

True or false?

The amplitude of $y = 1.5\cos(\pi t \div 6) + 3$ is 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q5

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 58 • FORWARD 1 • Q5/5

SET 58 • HSF-LE.B.5

FORWARD 2

Interpret the mean and standard deviation of a normal model • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

Battery life is normal with mean 400 hours and standard deviation 20 hours.

What does the 20 tell you?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 58 • FORWARD 2 • Q1/5

With the same model, about what percent of batteries last between 360 and 440 hours?

Percent: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data
HSS-ID.A.4 • Set 58 • FORWARD 2 • Q2/5

Which change makes a normal curve wider?

A. Increasing the standard deviation

B. Increasing the mean

C. Decreasing the standard deviation

D. Decreasing the mean

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S58-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data
HSS-ID.A.4 • Set 58 • FORWARD 2 • Q3/5

True or false?

Changing the mean of a normal model shifts the curve left or right.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data
HSS-ID.A.4 • Set 58 • FORWARD 2 • Q4/5

True or false?

In a normal model with mean 50 and standard deviation 5, the value 60 is 10 standard deviations above the mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 58 • FORWARD 2 • Q5/5

SET 59 • HSS-ID.A.1 • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Represent data with dot plots, histograms, and box plots

MAIN	Algebra I main question	**HSS-ID.A.1**
BACKWARD 1	Display numerical data in plots	**6.SP.B.4**
BACKWARD 2	Compute measures of center and variation	**6.SP.B.5.c**
FORWARD 1	Decide whether a normal model fits a distribution	**HSS-ID.A.4**
FORWARD 2	Decide whether results are consistent with a model by simulation	**HSS-IC.A.2**

SET 59 • HSS-ID.A.1

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1

Make a dot plot of the numbers of books read by 10 students:

2, 3, 3, 4, 4, 4, 5, 5, 6, 9



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 59 • MAIN • Q1/5

Find the five-number summary of 2, 3, 3, 4, 4, 4, 5, 5, 6, 9.

Summary: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 59 • MAIN • Q2/5

For the data 2, 3, 3, 4, 4, 4, 5, 5, 6, 9, use the 1.5 × IQR rule to decide whether 9 is an outlier.

Answer: _____


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 59 • MAIN • Q3/5

The histogram shows the heights of 30 plants. Each bar includes its lower edge but not its upper edge.

How many plants are at least 20 cm tall?



A. 21 **B.** 17 **C.** 5 **D.** 12

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-M-Q4**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 59 • MAIN • Q4/5

True or false?

A box plot shows the mean of a data set.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 59 • MAIN • Q5/5

SET 59 • HSS-ID.A.1

BACKWARD 1

Display numerical data in plots • Grade 6 • Statistics & Probability • 6.SP.B.4

Which display shows every data value, a dot plot or a box plot?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q1

Grade 6 • Statistics & Probability • 6.SP.B.4 • Set 59 • BACKWARD 1 • Q1/5

Make a dot plot of 1, 2, 2, 3, 3, 3, 5.



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q2

Grade 6 • Statistics & Probability • 6.SP.B.4 • Set 59 • BACKWARD 1 • Q2/5

Which display groups data values into intervals?

A. Box plot

B. Dot plot

C. Histogram

D. Ordered list

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q3

Grade 6 • Statistics & Probability • 6.SP.B.4 • Set 59 • BACKWARD 1 • Q3/5

True or false?

A box plot is made from the five-number summary.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q4

Grade 6 • Statistics & Probability • 6.SP.B.4 • Set 59 • BACKWARD 1 • Q4/5

True or false?

A histogram shows each individual data value.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q5

Grade 6 • Statistics & Probability • 6.SP.B.4 • Set 59 • BACKWARD 1 • Q5/5

SET 59 • HSS-ID.A.1

BACKWARD 2

Compute measures of center and variation • Grade 6 • Statistics & Probability • 6.SP.B.5.c

Find the median of 3, 8, 5, 10, 4.

Median: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q1

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 59 • BACKWARD 2 • Q1/5

Find the mean of 3, 8, 5, 10, 4.

Mean: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q2

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 59 • BACKWARD 2 • Q2/5

Q1 = 12 and Q3 = 20.

What is the interquartile range?

A. 32

B. 8

C. 4

D. 16

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q3

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 59 • BACKWARD 2 • Q3/5

True or false?

The range of 2, 7, 9, 15 is 13.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q4

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 59 • BACKWARD 2 • Q4/5

True or false?

The median of 1, 2, 3, 100 is 26.5.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q5

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 59 • BACKWARD 2 • Q5/5

SET 59 • HSS-ID.A.1

FORWARD 1

Decide whether a normal model fits a distribution • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

A data set is strongly skewed to the right.

Is a normal model appropriate? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F1-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data
HSS-ID.A.4 • Set 59 • FORWARD 1 • Q1/5

Scores are normal with mean 75 and standard deviation 8.

What score is 1 standard deviation above the mean?

Score: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F1-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 59 • FORWARD 1 • Q2/5

Values are normal with mean 20 and standard deviation 4.

About what percent are between 12 and 28?

A. 99.7%

B. 50%

C. 68%

D. 95%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-F1-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data
HSS-ID.A.4 • Set 59 • FORWARD 1 • Q3/5

True or false?

In a normal distribution, the mean and the median are equal.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F1-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 59 • FORWARD 1 • Q4/5

True or false?

About 50% of a normal distribution lies within 1 standard deviation of the mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F1-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 59 • FORWARD 1 • Q5/5

SET 59 • HSS-ID.A.1

FORWARD 2

Decide whether results are consistent with a model by simulation • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2

In 100 simulated sets of 10 fair coin flips, 8 or more heads happened in 5 sets.

Estimate the probability of 8 or more heads in 10 fair flips.

Estimate: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.2 • Set 59 • FORWARD 2 • Q1/5

A spinner model says red comes up 25% of the time. In 200 spins, red came up 52 times.

What proportion is that? Is it consistent with the model?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.2 • Set 59 • FORWARD 2 • Q2/5

A model says a coin lands heads 50% of the time.

Which result most strongly questions the model?

A. 52 heads in 100 flips

B. 6 heads in 10 flips

C. 95 heads in 100 flips

D. 48 heads in 100 flips

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.A.2 • Set 59 • FORWARD 2 • Q3/5

True or false?

A result that rarely happens in simulations of a model is evidence against the model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.2 • Set 59 • FORWARD 2 • Q4/5

True or false?

If a coin lands heads 6 times in 10 flips, the coin must be unfair.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.2 • Set 59 • FORWARD 2 • Q5/5

SET 60 • HSS-ID.A.2 • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Compare the center and spread of two or more data sets

MAIN	Algebra I main question	**HSS-ID.A.2**
BACKWARD 1	Tell measures of center from measures of variation	**6.SP.A.3**
BACKWARD 2	Compare two distributions using center and variability	**7.SP.B.3**
FORWARD 1	Use simulation to decide whether a difference between treatment…	**HSS-IC.B.5**
FORWARD 2	Evaluate reports based on data	**HSS-IC.B.6**

SET 60 • HSS-ID.A.2

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2

Class A scores: 70, 75, 80, 85, 90. Class B scores: 60, 70, 80, 90, 100.

Compare the centers and the spreads.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 60 • MAIN • Q1/5

Find the population standard deviation (divide by n) of 2, 4, 4, 4, 5, 5, 7, 9.

Standard deviation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 60 • MAIN • Q2/5

Store X's daily sales have median 50 and IQR 10. Store Y's have median 45 and IQR 25.

Which store typically sells more? Which is more consistent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 60 • MAIN • Q3/5

Which measure of spread is best for skewed data with outliers?

A. Range

B. Standard deviation

C. Mean

D. Interquartile range

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 60 • MAIN • Q4/5

True or false?

Two data sets with the same mean must have the same standard deviation.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 60 • MAIN • Q5/5

SET 60 • HSS-ID.A.2

BACKWARD 1

Tell measures of center from measures of variation • Grade 6 • Statistics & Probability • 6.SP.A.3

Which describes a typical value of a data set, the median or the IQR?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q1

Grade 6 • Statistics & Probability • 6.SP.A.3 • Set 60 • BACKWARD 1 • Q1/5

What is the range of the data set 4, 4, 4, 4?

Range: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q2

Grade 6 • Statistics & Probability • 6.SP.A.3 • Set 60 • BACKWARD 1 • Q2/5

Which is a measure of variation?

A. Median

B. Mode

C. Mean

D. Interquartile range

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q3

Grade 6 • Statistics & Probability • 6.SP.A.3 • Set 60 • BACKWARD 1 • Q3/5

True or false?

A measure of center summarizes all of the values with a single number.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q4

Grade 6 • Statistics & Probability • 6.SP.A.3 • Set 60 • BACKWARD 1 • Q4/5

True or false?

The range is a measure of center.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q5

Grade 6 • Statistics & Probability • 6.SP.A.3 • Set 60 • BACKWARD 1 • Q5/5

SET 60 • HSS-ID.A.2

BACKWARD 2

Compare two distributions using center and variability • Grade 7 • Statistics & Probability • 7.SP.B.3

Two classes have mean scores 78 and 84, and each has MAD 3.

How many MADs apart are the means?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.B.3 • Set 60 • BACKWARD 2 • Q1/5

Team A's mean height is 170 cm and Team B's is 180 cm; each has MAD 5 cm.

Express the difference of the means as a multiple of the MAD.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.B.3 • Set 60 • BACKWARD 2 • Q2/5

Two means differ by 12, and each data set has MAD 4.

How many MADs is the difference?

A. 48

B. 12

C. 3

D. 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.B.3 • Set 60 • BACKWARD 2 • Q3/5

True or false?

When two distributions have little variability, a small difference in their centers is easier to notice.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.B.3 • Set 60 • BACKWARD 2 • Q4/5

True or false?

If two data sets overlap, their means must be equal.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.B.3 • Set 60 • BACKWARD 2 • Q5/5

SET 60 • HSS-ID.A.2

FORWARD 1

Use simulation to decide whether a difference between treatments is significant • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

Plants given a fertilizer averaged 24 cm; the control group averaged 20 cm. In 1,000 random regroupings of the same plants, a difference of 4 cm or more happened 12 times.

Is the difference significant? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions

HSS-IC.B.5 • Set 60 • FORWARD 1 • Q1/5

Why are subjects assigned to treatments at random in an experiment?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 60 • FORWARD 1 • Q2/5

In a regrouping simulation, a difference as large as the observed one happened 400 times out of 1,000.

What does that suggest?

A. The treatment definitely works.

B. The difference could easily be due to chance.

C. The treatment definitely does not work.

D. The experiment had no control group.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S60-F1-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 60 • FORWARD 1 • Q3/5

True or false?

A small simulated probability for the observed difference is evidence that the treatment had an effect.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions

HSS-IC.B.5 • Set 60 • FORWARD 1 • Q4/5

True or false?

In an experiment, the control group receives the treatment.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 60 • FORWARD 1 • Q5/5

SET 60 • HSS-ID.A.2

FORWARD 2

Evaluate reports based on data • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

An ad says "9 out of 10 dentists recommend Brand X," based on 10 dentists chosen by Brand X.

What is the flaw?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 60 • FORWARD 2 • Q1/5

An online poll of a website's visitors finds 80% support a policy.

Can the result be applied to all adults? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 60 • FORWARD 2 • Q2/5

Which sample is most likely to represent a school's students?

A. The first 50 students to arrive

B. The 50 students in the chess club

C. A random sample of 50 students from the school roster

D. 50 volunteers

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S60-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 60 • FORWARD 2 • Q3/5

True or false?

A larger random sample generally gives a smaller margin of error.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 60 • FORWARD 2 • Q4/5

True or false?

A survey with many volunteers is free of bias.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 60 • FORWARD 2 • Q5/5

SET 61 • HSS-ID.A.3 • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Interpret differences in shape, center, and spread, accounting for outliers

MAIN	Algebra I main question	**HSS-ID.A.3**
BACKWARD 1	Describe a distribution by its center, spread, and shape	**6.SP.A.2**
BACKWARD 2	Choose measures of center and spread that fit the data	**6.SP.B.5.d**
FORWARD 1	Use z-scores to judge how unusual a value is	**HSS-ID.A.4**
FORWARD 2	Distinguish a population parameter from a sample statistic	**HSS-IC.A.1**

SET 61 • HSS-ID.A.3

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3

Salaries at a small company, in thousands of dollars: 40, 42, 45, 45, 48, 50, 250.

Find the mean and the median. Which better describes a typical salary?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 61 • MAIN • Q1/5

If the salary of 250 is removed, what happens to the mean and to the median?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 61 • MAIN • Q2/5

A histogram of wait times has a long tail on the right.

Describe its shape and compare its mean and median.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 61 • MAIN • Q3/5

Which data set has the largest standard deviation?

A. 28, 29, 30, 31, 32

B. 25, 30, 30, 30, 35

C. 10, 20, 30, 40, 50

D. 30, 30, 30, 30, 30

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 61 • MAIN • Q4/5

True or false?

An outlier affects the median more than the mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 61 • MAIN • Q5/5

SET 61 • HSS-ID.A.3

BACKWARD 1

Describe a distribution by its center, spread, and shape • Grade 6 • Statistics & Probability • 6.SP.A.2

Data: 1, 2, 2, 3, 3, 3, 4, 4, 5.

Describe the shape of the distribution.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q1

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 61 • BACKWARD 1 • Q1/5

Which value in 2, 3, 3, 3, 4, 15 is unusual?

Value: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q2

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 61 • BACKWARD 1 • Q2/5

A dot plot has most values on the left and a few values far to the right.

What is its shape?

A. Skewed right

B. Symmetric

C. Uniform

D. Skewed left

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q3

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 61 • BACKWARD 1 • Q3/5

True or false?

A distribution can be described by its center, its spread, and its shape.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q4

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 61 • BACKWARD 1 • Q4/5

True or false?

A symmetric distribution must have an outlier.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q5

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 61 • BACKWARD 1 • Q5/5

SET 61 • HSS-ID.A.3

BACKWARD 2

Choose measures of center and spread that fit the data • Grade 6 • Statistics & Probability • 6.SP.B.5.d

For data with an extreme outlier, should you report the mean or the median as the center? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q1

Grade 6 • Statistics & Probability • 6.SP.B.5.d • Set 61 • BACKWARD 2 • Q1/5

For symmetric data with no outliers, which measures of center and spread are usually reported?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q2

Grade 6 • Statistics & Probability • 6.SP.B.5.d • Set 61 • BACKWARD 2 • Q2/5

Home prices in a town include one mansion.

Which measure best describes a typical price?

A. The mean

B. The median

C. The maximum

D. The range

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q3

Grade 6 • Statistics & Probability • 6.SP.B.5.d • Set 61 • BACKWARD 2 • Q3/5

True or false?

The median is not changed much by a single extreme value.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q4

Grade 6 • Statistics & Probability • 6.SP.B.5.d • Set 61 • BACKWARD 2 • Q4/5

True or false?

The IQR changes greatly when the largest value is made even larger.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q5

Grade 6 • Statistics & Probability • 6.SP.B.5.d • Set 61 • BACKWARD 2 • Q5/5

SET 61 • HSS-ID.A.3

FORWARD 1

Use z-scores to judge how unusual a value is • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

A distribution has mean 50 and standard deviation 4.

What is the z-score of 58?

$z =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 61 • FORWARD 1 • Q1/5

A score of 92 on a test with mean 80 and standard deviation 6, or a score of 85 on a test with mean 70 and standard deviation 10: which is better relative to its test?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 61 • FORWARD 1 • Q2/5

In a normal distribution, which z-score is the most unusual?

A. -3.1

B. -0.5

C. 1.2

D. 2.0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-F1-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data
HSS-ID.A.4 • Set 61 • FORWARD 1 • Q3/5

True or false?

A z-score of 0 means the value equals the mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data

HSS-ID.A.4 • Set 61 • FORWARD 1 • Q4/5

True or false?

A z-score of -2 is above the mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Interpreting Categorical & Quantitative Data
HSS-ID.A.4 • Set 61 • FORWARD 1 • Q5/5

SET 61 • HSS-ID.A.3

FORWARD 2

Distinguish a population parameter from a sample statistic • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

What is the difference between a population and a sample?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.1 • Set 61 • FORWARD 2 • Q1/5

A random sample of 200 students has a mean of 7.1 hours of sleep.

What does 7.1 estimate?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.1 • Set 61 • FORWARD 2 • Q2/5

A parameter describes

A. a graph

B. one individual

C. a population

D. a sample

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.1 • Set 61 • FORWARD 2 • Q3/5

True or false?

Different random samples from the same population can give different sample means.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.1 • Set 61 • FORWARD 2 • Q4/5

True or false?

A sample statistic always equals the population parameter.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.1 • Set 61 • FORWARD 2 • Q5/5

SET 62 • HSS-ID.B.5 • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Summarize categorical data in two-way frequency tables and interpret relative frequencies

MAIN	Algebra I main question	**HSS-ID.B.5**
BACKWARD 1	Read two-way tables for association	**8.SP.A.4**
BACKWARD 2	Write a part of a whole as a percent	**7.RP.A.3**
FORWARD 1	Use a two-way table to find conditional probabilities and test inde…	**HSS-CP.A.4**
FORWARD 2	Explain conditional probability and independence in everyday lang..	**HSS-CP.A.5**

SET 62 • HSS-ID.B.5

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.5

The table shows a survey of 200 students.

What percent of all the students play a sport?

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

Percent: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.5 • Set 62 • MAIN • Q1/5

What percent of Grade 10 students play a sport? What percent of Grade 9 students do?

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.5 • Set 62 • MAIN • Q2/5

Is there an association between grade and playing a sport? Explain.

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.5 • Set 62 • MAIN • Q3/5

What is the joint relative frequency of "Grade 9 and does not play a sport"?

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

A. 50%

B. 57.9%

C. 55%

D. 27.5%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-M-Q4**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.5 • Set 62 • MAIN • Q4/5

True or false?

Of the students who play a sport, more than half are in Grade 10.

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.5 • Set 62 • MAIN • Q5/5

SET 62 • HSS-ID.B.5

BACKWARD 1

Read two-way tables for association • Grade 8 • Statistics & Probability • 8.SP.A.4

12 of 20 boys and 9 of 15 girls in a class like math.

How many students in all like math?

Students: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q1

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 62 • BACKWARD 1 • Q1/5

Using the same class, is there an association between gender and liking math? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q2

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 62 • BACKWARD 1 • Q2/5

What are the row and column totals of a two-way table called?

A. Joint frequencies

B. Conditional frequencies

C. Marginal frequencies

D. Outliers

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q3

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 62 • BACKWARD 1 • Q3/5

True or false?

A two-way table shows data on two categorical variables.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q4

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 62 • BACKWARD 1 • Q4/5

True or false?

In a two-way table, every row must have the same total.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q5

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 62 • BACKWARD 1 • Q5/5

SET 62 • HSS-ID.B.5

BACKWARD 2

Write a part of a whole as a percent • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

Write 45 out of 100 as a percent.

Percent: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 62 • BACKWARD 2 • Q1/5

Write 60 out of 105 as a percent, rounded to the nearest tenth.

Percent: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 62 • BACKWARD 2 • Q2/5

27 is what percent of 60?

A. 45%

B. 60%

C. 33%

D. 27%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 62 • BACKWARD 2 • Q3/5

True or false?

55 out of 200 is 27.5%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 62 • BACKWARD 2 • Q4/5

True or false?

40 out of 95 is 40%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 62 • BACKWARD 2 • Q5/5

SET 62 • HSS-ID.B.5

FORWARD 1

Use a two-way table to find conditional probabilities and test independence • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability • HSS-CP.A.4

A student from the table is chosen at random.

Find P(plays a sport | Grade 10).

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

Probability: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.4 • Set 62 • FORWARD 1 • Q1/5

Are "Grade 10" and "plays a sport" independent? Explain with probabilities.

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.4 • Set 62 • FORWARD 1 • Q2/5

What is P(Grade 9 | plays a sport)?

	Plays a sport	Does not	Total
Grade 9	45	55	100
Grade 10	60	40	100
Total	105	95	200

A. About 0.43

B. 0.225

C. 0.525

D. 0.45

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F1-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability
HSS-CP.A.4 • Set 62 • FORWARD 1 • Q3/5

True or false?

Events A and B are independent if $P(A \mid B) = P(A)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.4 • Set 62 • FORWARD 1 • Q4/5

True or false?

P(A | B) always equals P(B | A).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.4 • Set 62 • FORWARD 1 • Q5/5

SET 62 • HSS-ID.B.5

FORWARD 2

Explain conditional probability and independence in everyday language • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability • HSS-CP.A.5

Explain in everyday language what P(late | rain) > P(late) means.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.5 • Set 62 • FORWARD 2 • Q1/5

A and B are independent, P(A) = 0.3, and P(B) = 0.5.

Find P(A and B).

Probability: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.5 • Set 62 • FORWARD 2 • Q2/5

Which pair of events is most likely independent?

A. Height and shoe size

B. Studying and passing a test

C. Flipping heads and rolling a 6

D. Rain and carrying an umbrella

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability
HSS-CP.A.5 • Set 62 • FORWARD 2 • Q3/5

True or false?

If A and B are independent, then P(A and B) = P(A) · P(B).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.5 • Set 62 • FORWARD 2 • Q4/5

True or false?

If P(A | B) = 0.8 and P(A) = 0.4, then A and B are independent.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.A.5 • Set 62 • FORWARD 2 • Q5/5

SET 63 • HSS-ID.B.6.a • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Fit a linear, quadratic, or exponential function to data and use it to solve problems

MAIN	Algebra I main question	**HSS-ID.B.6.a**
BACKWARD 1	Describe patterns in scatter plots	**8.SP.A.1**
BACKWARD 2	Write linear and exponential functions through given points	**HSF-LE.A.2**
FORWARD 1	Use logarithms with an exponential model fitted to data	**HSF-LE.A.4**
FORWARD 2	Fit a sinusoidal model to periodic data	**HSF-TF.B.5**


SET 63 • HSS-ID.B.6.a

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a

The scatter plot shows a population over 5 years.

Which type of function fits best? Explain.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 63 • MAIN • Q1/5

An exponential model for the data is $P(t) = 100(1.3)^t$.

Use it to predict the population at t = 8, to the nearest whole number.

$P(8) \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 63 • MAIN • Q2/5

Data: x = 0, 1, 2, 3, 4 and y = 2, 5, 10, 17, 26.

Which type of function fits exactly? Write it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 63 • MAIN • Q3/5

A scatter plot rises and then falls in a symmetric curve.

Which model fits best?

A. Quadratic

B. Exponential growth

C. Exponential decay

D. Linear

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 63 • MAIN • Q4/5

True or false?

Using a model to predict far outside the range of the data is always reliable.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 63 • MAIN • Q5/5

SET 63 • HSS-ID.B.6.a

BACKWARD 1

Describe patterns in scatter plots • Grade 8 • Statistics & Probability • 8.SP.A.1

Points rise from left to right, close to a line.

Describe the association.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q1

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 63 • BACKWARD 1 • Q1/5

What is an outlier in a scatter plot?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q2

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 63 • BACKWARD 1 • Q2/5

Points fall from left to right in a curved pattern.

Which describes the association?

A. Positive and nonlinear

B. Positive and linear

C. Negative and nonlinear

D. No association

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q3

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 63 • BACKWARD 1 • Q3/5

True or false?

A scatter plot shows the relationship between two quantitative variables.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q4

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 63 • BACKWARD 1 • Q4/5

True or false?

If the points show no pattern, there is a strong association.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q5

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 63 • BACKWARD 1 • Q5/5

SET 63 • HSS-ID.B.6.a

BACKWARD 2

Write linear and exponential functions through given points • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

Write the exponential function through (0, 100) and (1, 130).

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 63 • BACKWARD 2 • Q1/5

Write the linear function through (0, 2) and (4, 26).

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 63 • BACKWARD 2 • Q2/5

Which exponential function $y = ab^x$ $(b > 0)$ passes through $(0, 50)$ and $(2, 12.5)$?

A. $y = 50(0.25)^x$

B. $y = 12.5(2)^x$

C. $y = 50(0.5)^x$

D. $y = 50 - 18.75x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 63 • BACKWARD 2 • Q3/5

True or false?

$y = 3(2)^x$ passes through (2, 12).

A. True

B. False

True or false?

$y = 5(1.2)^x$ passes through (0, 6).

A. True

B. False

SET 63 • HSS-ID.B.6.a

FORWARD 1

Use logarithms with an exponential model fitted to data • High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

$P(t) = 100(1.3)^t$.

When does P reach 1,000? Round to the nearest tenth.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S63-F1-Q1**

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 63 • FORWARD 1 • Q1/5

Solve $2(1.5)^x = 30$. Round to the nearest hundredth.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F1-Q2

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 63 • FORWARD 1 • Q2/5

Which is the exact solution of $100(1.3)^t = 500$?

A. $t = \log 1.3 \div \log 5$

B. $t = 500 \div 130$

C. $t = \log 5 \div \log 1.3$

D. $t = 5 \div 1.3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S63-F1-Q3**

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 63 • FORWARD 1 • Q3/5

True or false?

If $1.3^t = 1.69$, then $t = 2$.

A. True

B. False

True or false?

If $10^t = 50$, then $t = 5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F1-Q5

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 63 • FORWARD 1 • Q5/5

SET 63 • HSS-ID.B.6.a

FORWARD 2

Fit a sinusoidal model to periodic data • High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5

Monthly average temperatures peak at 30 °C and bottom out at 10 °C.

What amplitude and midline fit a sinusoidal model?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q1

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 63 • FORWARD 2 • Q1/5

What period fits monthly temperatures that repeat every year?

Period: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q2

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 63 • FORWARD 2 • Q2/5

Daylight hours vary from 9 to 15 over a year.

Which model fits?

A. A linear model

B. A sinusoidal model with amplitude 3 and period 12 months

C. A quadratic model with vertex at month 12

D. An exponential model

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S63-F2-Q3**

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 63 • FORWARD 2 • Q3/5

True or false?

A repeating pattern in data suggests a sinusoidal model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q4

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 63 • FORWARD 2 • Q4/5

True or false?

Data that vary between 9 and 15 have amplitude 6.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q5

High School — Algebra II (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 63 • FORWARD 2 • Q5/5

Assess the fit of a function by plotting and analyzing residuals

MAIN	Algebra I main question	**HSS-ID.B.6.b**
BACKWARD 1	Use a fitted function to make predictions	**HSS-ID.B.6.a**
BACKWARD 2	Subtract rational numbers	**7.NS.A.1.c**
FORWARD 1	Use logarithms to straighten exponential data	**HSF-LE.A.4**
FORWARD 2	Evaluate reports that use fitted models	**HSS-IC.B.6**

SET 64 • HSS-ID.B.6.b

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.b

The model $\hat{y} = 2.2x + 0.8$ is fitted to the data (1, 3), (2, 5), (3, 8), (4, 9), (5, 12).

Find the residual for each point.

Residuals: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.b • Set 64 • MAIN • Q1/5

For that model, what does the residual -0.6 at $x = 4$ mean?


Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.b • Set 64 • MAIN • Q2/5

Make a residual plot for the residuals 0, −0.2, 0.6, −0.6, 0.2 at x = 1, 2, 3, 4, 5.



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.b • Set 64 • MAIN • Q3/5

A residual plot shows a clear U-shaped pattern.

What does that suggest?

A. The correlation is exactly 1.

B. A linear model is not a good fit; a curved model may fit better.

C. The linear model fits well.

D. There are no residuals.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.b • Set 64 • MAIN • Q4/5

True or false?

A good linear fit has residuals scattered randomly around 0, with no pattern.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.b • Set 64 • MAIN • Q5/5

SET 64 • HSS-ID.B.6.b

BACKWARD 1

Use a fitted function to make predictions • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a

The model is $\hat{y} = 2.2x + 0.8$.

Predict y when $x = 6$.

$\hat{y} =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 64 • BACKWARD 1 • Q1/5

The model is $y = 3x^2 - 2$.

Predict y when x = 4.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 64 • BACKWARD 1 • Q2/5

The model is $y = 50(0.9)^x$.

Predict y when x = 2.

A. 50

B. 40.5

C. 45

D. 90

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S64-B1-Q3**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 64 • BACKWARD 1 • Q3/5

True or false?

A predicted value comes from the model, not from the data table.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 64 • BACKWARD 1 • Q4/5

True or false?

If a model predicts 14 and the actual value is 14, the residual is 14.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.a • Set 64 • BACKWARD 1 • Q5/5

SET 64 • HSS-ID.B.6.b

BACKWARD 2

Subtract rational numbers • Grade 7 • The Number System • 7.NS.A.1.c

Find $8 - 7.4$.

Value: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q1

Grade 7 • The Number System • 7.NS.A.1.c • Set 64 • BACKWARD 2 • Q1/5

Find $9 - 9.6$.

Value: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q2

Grade 7 • The Number System • 7.NS.A.1.c • Set 64 • BACKWARD 2 • Q2/5

Find $5 - 5.2$.

A. -10.2

B. 10.2

C. 0.2

D. -0.2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q3

Grade 7 • The Number System • 7.NS.A.1.c • Set 64 • BACKWARD 2 • Q3/5

True or false?

$3 - (-2) = 5$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q4

Grade 7 • The Number System • 7.NS.A.1.c • Set 64 • BACKWARD 2 • Q4/5

True or false?

$-4 - 6 = -2$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q5

Grade 7 • The Number System • 7.NS.A.1.c • Set 64 • BACKWARD 2 • Q5/5

SET 64 • HSS-ID.B.6.b

FORWARD 1

Use logarithms to straighten exponential data • High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Data: $y = 1, 3, 9, 27$ at $x = 0, 1, 2, 3$.

Find $\log_3 y$ for each point.

Values: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q1

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 64 • FORWARD 1 • Q1/5

A fitted line is log y = 0.3x + 1.

Write y as an exponential function of x.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q2

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 64 • FORWARD 1 • Q2/5

Why take the logarithm of y for data that look exponential?

A. It removes all of the residuals.

B. It makes every y-value negative.

C. It doubles the data.

D. It makes the pattern linear, so a line can be fit.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S64-F1-Q3**

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 64 • FORWARD 1 • Q3/5

True or false?

If $y = 5 \cdot 2^x$, then log y is a linear function of x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q4

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 64 • FORWARD 1 • Q4/5

True or false?

$$\log(5 \cdot 2^x) = 5 + 2x$$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q5

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 64 • FORWARD 1 • Q5/5

SET 64 • HSS-ID.B.6.b

FORWARD 2

Evaluate reports that use fitted models • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

A report fits a line to the heights of children aged 5 to 15 and uses it to predict the height of a 40-year-old.

What is the problem?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 64 • FORWARD 2 • Q1/5

A report finds a strong correlation between ice cream sales and drownings and says ice cream causes drowning.

What is wrong?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions

HSS-IC.B.6 • Set 64 • FORWARD 2 • Q2/5

A residual plot shows a strong curve, but a report says the linear model is excellent.

What should you conclude?

A. The report is correct.

B. The data must be wrong.

C. The residuals are all 0.

D. The linear model may not be appropriate.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S64-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 64 • FORWARD 2 • Q3/5

True or false?

A report should state how its data were collected.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 64 • FORWARD 2 • Q4/5

True or false?

A model with a high correlation coefficient is always the right type of model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 64 • FORWARD 2 • Q5/5

SET 65 • HSS-ID.B.6.c • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Fit a linear function to a scatter plot that suggests a linear association

MAIN	Algebra I main question	**HSS-ID.B.6.c**
BACKWARD 1	Judge a line of fit informally	**8.SP.A.2**
BACKWARD 2	Write a line through two points	**HSF-LE.A.2**
FORWARD 1	Evaluate conclusions drawn from a line of fit	**HSS-IC.B.6**
FORWARD 2	Use simulation to judge whether an association could be due to ch…	**HSS-IC.B.5**

SET 65 • HSS-ID.B.6.c


MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c

The scatter plot shows hours studied and test scores.

Draw a line of fit and write its equation.

Equation: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 65 • MAIN • Q1/5

The least-squares line for the data is $\hat{y} = 4x + 58.14$.

Predict the score for 4.5 hours, to the nearest whole number.

Score: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 65 • MAIN • Q2/5

Write the equation of the line through the data points (1, 62) and (7, 86), to use as a line of fit.

Equation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 65 • MAIN • Q3/5

Which line fits the points (0, 1), (1, 3), (2, 5), (3, 7) exactly?

A. $y = 3x + 1$

B. $y = 2x + 1$

C. $y = 2x$

D. $y = x + 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 65 • MAIN • Q4/5

True or false?

A line of fit should pass through every data point.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 65 • MAIN • Q5/5

SET 65 • HSS-ID.B.6.c

BACKWARD 1

Judge a line of fit informally • Grade 8 • Statistics & Probability • 8.SP.A.2

Is a straight line a good model for points that curve strongly upward? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q1

Grade 8 • Statistics & Probability • 8.SP.A.2 • Set 65 • BACKWARD 1 • Q1/5

How can you judge whether a line of fit is good?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q2

Grade 8 • Statistics & Probability • 8.SP.A.2 • Set 65 • BACKWARD 1 • Q2/5

A line of fit should have

A. all of the points below it

B. all of the points above it

C. about as many points above it as below it

D. exactly two points on it

True or false?

Lines of fit are used for data with a linear association.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q4

Grade 8 • Statistics & Probability • 8.SP.A.2 • Set 65 • BACKWARD 1 • Q4/5

True or false?

A line of fit must pass through the origin.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q5

Grade 8 • Statistics & Probability • 8.SP.A.2 • Set 65 • BACKWARD 1 • Q5/5

SET 65 • HSS-ID.B.6.c

BACKWARD 2

Write a line through two points • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

Write the equation of the line through (2, 66) and (6, 82).

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 65 • BACKWARD 2 • Q1/5

Write the equation of the line through (0, 7) and (3, 13).

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 65 • BACKWARD 2 • Q2/5

Which line passes through (0, 10) and (5, 0)?

A. $y = 2x + 10$

B. $y = 10x - 2$

C. $y = -2x + 10$

D. $y = -0.5x + 10$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 65 • BACKWARD 2 • Q3/5

True or false?

The line through (0, 3) with slope 4 is $y = 4x + 3$.

A. True

B. False

True or false?

The line through (2, 2) and (4, 8) has slope 2.

A. True

B. False

SET 65 • HSS-ID.B.6.c

FORWARD 1

Evaluate conclusions drawn from a line of fit • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

In a study of 7 students, each extra hour of study went with about 4 more points.

Can a school claim this holds for all of its students? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 65 • FORWARD 1 • Q1/5

A report uses $\hat{y} = 4x + 58$ to predict a score of 138 for 20 hours of study.

Why is that unreasonable?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 65 • FORWARD 1 • Q2/5

Which detail makes a regression conclusion more trustworthy?

A. The prediction is far outside the data.

B. The line was drawn by hand.

C. The data came from a large random sample.

D. The sample was 5 volunteers.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S65-F1-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 65 • FORWARD 1 • Q3/5

True or false?

Observational data alone cannot prove that one variable causes another.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 65 • FORWARD 1 • Q4/5

True or false?

A line of fit from 7 students proves the effect for every student.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 65 • FORWARD 1 • Q5/5

SET 65 • HSS-ID.B.6.c

FORWARD 2

Use simulation to judge whether an association could be due to chance • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

The test scores were randomly re-paired with the study hours 1,000 times, and the slope was never as large as 4.

What does that suggest?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 65 • FORWARD 2 • Q1/5

In an experiment, a tutoring group averaged 6 points more than a control group. Random regroupings gave differences of 6 or more 30% of the time.

What is the conclusion?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions

HSS-IC.B.5 • Set 65 • FORWARD 2 • Q2/5

What is the purpose of random assignment in an experiment?

A. To make the groups alike except for the treatment

B. To make the sample smaller

C. To guarantee a significant result

D. To remove the need for a control group

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S65-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 65 • FORWARD 2 • Q3/5

True or false?

A result is statistically significant when it would rarely happen by chance alone.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 65 • FORWARD 2 • Q4/5

True or false?

A difference that happens in 40% of random simulations is statistically significant.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 65 • FORWARD 2 • Q5/5

SET 66 • HSS-ID.C.7 • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Interpret the slope and the intercept of a linear model in the context of the data

MAIN	Algebra I main question	**HSS-ID.C.7**
BACKWARD 1	Interpret the parameters of linear and exponential functions	**HSF-LE.B.5**
BACKWARD 2	Use a linear model to solve problems	**8.SP.A.3**
FORWARD 1	Interpret the parameters of continuous exponential models	**HSF-LE.A.4**
FORWARD 2	Understand a fitted slope as an estimate of a population value	**HSS-IC.A.1**

SET 66 • HSS-ID.C.7

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.7

$\hat{y} = 4x + 58$ predicts a test score from x hours of study.

Interpret the slope and the intercept.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.7 • Set 66 • MAIN • Q1/5

A model for used cars is $\hat{y} = 22{,}000 - 1{,}800x$, where x is the age in years and $\hat{y}$ is the price in dollars.

Interpret $-1{,}800$ and $22{,}000$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.7 • Set 66 • MAIN • Q2/5

$\hat{y} = 0.9x - 20$ predicts an adult's weight in pounds from height in inches.

Does the intercept make sense in context? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.7 • Set 66 • MAIN • Q3/5

$\hat{y} = 2.5x + 10$ predicts the cost in dollars of a pizza with x toppings.

What does 2.5 mean?

A. Each additional topping adds about \$2.50 to the predicted cost.

B. A pizza has 2.5 toppings.

C. A pizza costs \$2.50.

D. A plain pizza costs \$2.50.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S66-M-Q4**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.7 • Set 66 • MAIN • Q4/5

True or false?

The slope of a fitted line gives the exact change in y for every individual.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.7 • Set 66 • MAIN • Q5/5

SET 66 • HSS-ID.C.7

BACKWARD 1

Interpret the parameters of linear and exponential functions • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5

A technician charges $C = 50 + 20h$ dollars for h hours.

What does 20 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 66 • BACKWARD 1 • Q1/5

$P = 3{,}000(0.9)^t$ is a population after t years.

What does 0.9 mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 66 • BACKWARD 1 • Q2/5

$d = 300 - 60t$ is a driver's distance from home, in miles, after t hours.

What does 300 mean?

A. The starting distance from home

B. The speed

C. The distance driven each hour

D. The time to arrive home

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S66-B1-Q3**

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 66 • BACKWARD 1 • Q3/5

True or false?

In y = 1.5x + 4, increasing x by 2 increases y by 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 66 • BACKWARD 1 • Q4/5

True or false?

In $y = 8(1.25)^x$, y increases by 1.25 for each unit of x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.B.5 • Set 66 • BACKWARD 1 • Q5/5

SET 66 • HSS-ID.C.7

BACKWARD 2

Use a linear model to solve problems • Grade 8 • Statistics & Probability • 8.SP.A.3

$y = 2x + 5$ models a plant's height in cm after x weeks.

Predict the height at 6 weeks.

Height: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q1

Grade 8 • Statistics & Probability • 8.SP.A.3 • Set 66 • BACKWARD 2 • Q1/5

Using $y = 2x + 5$, after how many weeks is the plant 21 cm tall?

Weeks: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q2

Grade 8 • Statistics & Probability • 8.SP.A.3 • Set 66 • BACKWARD 2 • Q2/5

In $y = 2x + 5$, what does 5 mean?

A. The height at week 0

B. The weeks needed to grow 1 cm

C. The final height

D. The growth per week

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q3

Grade 8 • Statistics & Probability • 8.SP.A.3 • Set 66 • BACKWARD 2 • Q3/5

True or false?

The slope of a linear model is a rate of change.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q4

Grade 8 • Statistics & Probability • 8.SP.A.3 • Set 66 • BACKWARD 2 • Q4/5

True or false?

In $y = 2x + 5$, the plant grows 5 cm each week.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q5

Grade 8 • Statistics & Probability • 8.SP.A.3 • Set 66 • BACKWARD 2 • Q5/5

SET 66 • HSS-ID.C.7

FORWARD 1

Interpret the parameters of continuous exponential models • High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

$A(t) = 500e^{0.04t}$.

What does 0.04 mean? When does A double, to the nearest tenth?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q1

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 66 • FORWARD 1 • Q1/5

$N(t) = 1{,}000e^{-0.1t}$.

What does -0.1 mean? When is $N = 500$, to the nearest tenth?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q2

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 66 • FORWARD 1 • Q2/5

Solve $e^{2x} = 9$ exactly.

A. $x = \ln 3$

B. $x = 4.5$

C. $x = e^{4.5}$

D. $x = \ln 9 \div 9$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S66-F1-Q3**

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 66 • FORWARD 1 • Q3/5

True or false?

$\ln(e^{0.04t}) = 0.04t$

A. True

B. False

True or false?

If $e^k = 1$, then $k = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q5

High School — Algebra II (course placement in this bank) • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 66 • FORWARD 1 • Q5/5

SET 66 • HSS-ID.C.7

FORWARD 2

Understand a fitted slope as an estimate of a population value • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

A random sample of 50 used cars of one model gives $\hat{y} = 22{,}000 - 1{,}800x$.

What population value does $-1{,}800$ estimate?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions

HSS-IC.A.1 • Set 66 • FORWARD 2 • Q1/5

Would another random sample of 50 cars give exactly the same slope? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.1 • Set 66 • FORWARD 2 • Q2/5

Which of these is a parameter?

A. The mean price of the 50 sampled cars

B. One car's price

C. The slope from the sample

D. The mean price of all used cars of this model

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q3

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.A.1 • Set 66 • FORWARD 2 • Q3/5

True or false?

Larger random samples tend to give statistics closer to the parameter.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.A.1 • Set 66 • FORWARD 2 • Q4/5

True or false?

A statistic is a number that describes a whole population.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.A.1 • Set 66 • FORWARD 2 • Q5/5

Compute (using technology) and interpret the correlation coefficient of a linear fit

MAIN	Algebra I main question	**HSS-ID.C.8**
BACKWARD 1	Describe the direction of an association	**8.SP.A.1**
BACKWARD 2	Fit and use a line for linear data	**HSS-ID.B.6.c**
FORWARD 1	Use simulation to judge whether a correlation could be due to ch…	**HSS-IC.B.5**
FORWARD 2	Evaluate claims based on a correlation	**HSS-IC.B.6**

SET 67 • HSS-ID.C.8

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8

Use technology to find the correlation coefficient r for the data

(1, 9), (2, 8), (3, 8), (4, 6), (5, 5), (6, 5), (7, 3), (8, 2),

to the nearest hundredth. Describe the association.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 67 • MAIN • Q1/5

$r = -0.85$ for hours of TV and test score.

Interpret this value.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 67 • MAIN • Q2/5

Which value of r shows the strongest linear relationship?

A. 0.10

B. −0.92

C. −0.40

D. 0.75

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 67 • MAIN • Q3/5

Can a correlation coefficient equal 1.3? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 67 • MAIN • Q4/5

True or false?

r = 0 means there is no relationship of any kind between the variables.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 67 • MAIN • Q5/5

SET 67 • HSS-ID.C.8

BACKWARD 1

Describe the direction of an association • Grade 8 • Statistics & Probability • 8.SP.A.1

Points rise from left to right, fairly close to a line.

Describe the association.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q1

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 67 • BACKWARD 1 • Q1/5

Points for hours of sleep and number of errors fall from left to right.

Is the association positive or negative?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q2

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 67 • BACKWARD 1 • Q2/5

Which pair of variables likely has a positive association?

A. Height and arm span

B. Outdoor temperature and heating cost

C. A car's age and its value

D. Speed and travel time for a fixed trip

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q3

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 67 • BACKWARD 1 • Q3/5

True or false?

In a negative association, as one variable increases, the other tends to decrease.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q4

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 67 • BACKWARD 1 • Q4/5

True or false?

Clusters in a scatter plot mean the data are linear.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q5

Grade 8 • Statistics & Probability • 8.SP.A.1 • Set 67 • BACKWARD 1 • Q5/5

SET 67 • HSS-ID.C.8

BACKWARD 2

Fit and use a line for linear data • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c

Write a line of fit through (2, 5) and (8, 14).

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 67 • BACKWARD 2 • Q1/5

The least-squares line is $\hat{y} = 1.5x + 2$.

Predict y when x = 6.

$\hat{y} =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 67 • BACKWARD 2 • Q2/5

What does the least-squares line minimize?

A. The y-intercept

B. The sum of the squared residuals

C. The number of points

D. The slope

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 67 • BACKWARD 2 • Q3/5

True or false?

The least-squares line passes through the point $(\bar{x}, \bar{y})$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 67 • BACKWARD 2 • Q4/5

True or false?

Two people drawing lines of fit by eye always get the same line.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.B.6.c • Set 67 • BACKWARD 2 • Q5/5

SET 67 • HSS-ID.C.8

FORWARD 1

Use simulation to judge whether a correlation could be due to chance • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

In 1,000 random shufflings of the y-values, r was at least 0.9 in 2 trials. The observed r is 0.92.

Is the association likely to be due to chance?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.5 • Set 67 • FORWARD 1 • Q1/5

For a different data set, shuffled data gave $r \geq 0.3$ in 350 of 1,000 trials, and the observed r is 0.3.

What is the conclusion?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.5 • Set 67 • FORWARD 1 • Q2/5

Why shuffle the y-values in such a simulation?

A. To increase r

B. To see how large r tends to be when there is no real association

C. To remove outliers

D. To make the data linear

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S67-F1-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.5 • Set 67 • FORWARD 1 • Q3/5

True or false?

With a larger sample, a moderate correlation is less likely to be due to chance.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 67 • FORWARD 1 • Q4/5

True or false?

If shuffled data often give an r as large as the observed r, the association is significant.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.5 • Set 67 • FORWARD 1 • Q5/5

SET 67 • HSS-ID.C.8

FORWARD 2

Evaluate claims based on a correlation • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

A news story says, "r = 0.6, so 60% of the change in sales is caused by advertising."

Name two errors.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 67 • FORWARD 2 • Q1/5

A study of 8 cities finds r = 0.7 between the number of parks and happiness scores.

Why should you be cautious?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 67 • FORWARD 2 • Q2/5

Which statement about $r = -0.9$ is correct?

A. There is a strong negative linear association.

B. The slope of the line is -0.9.

C. 90% of the points are on the line.

D. An increase in x causes y to fall.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S67-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions
HSS-IC.B.6 • Set 67 • FORWARD 2 • Q3/5

True or false?

A correlation can be strong even when the relationship is not causal.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 67 • FORWARD 2 • Q4/5

True or false?

If r = 0.8, the slope of the least-squares line is 0.8.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.6 • Set 67 • FORWARD 2 • Q5/5

SET 68 • HSS-ID.C.9 • INTERPRETING CATEGORICAL & QUANTITATIVE DATA

Distinguish between correlation and causation

MAIN	Algebra I main question	**HSS-ID.C.9**
BACKWARD 1	Interpret the correlation coefficient	**HSS-ID.C.8**
BACKWARD 2	Use random samples to make valid generalizations	**7.SP.A.1**
FORWARD 1	Distinguish surveys, experiments, and observational studies	**HSS-IC.B.3**
FORWARD 2	Use random methods to make fair decisions	**HSS-MD.B.6**

SET 68 • HSS-ID.C.9

MAIN

Algebra I main question • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9

Cities with more firefighters have more fire damage.

Does hiring firefighters cause damage? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 68 • MAIN • Q1/5

Ice cream sales and sunburn rates are strongly correlated.

Name a lurking variable.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 68 • MAIN • Q2/5

How could you test whether a new study app causes higher test scores?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 68 • MAIN • Q3/5

Which study design can show that one variable causes changes in another?

A. A survey of volunteers

B. A scatter plot of existing data

C. An observational study

D. A randomized experiment

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 68 • MAIN • Q4/5

True or false?

A strong correlation between two variables proves that one causes the other.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 68 • MAIN • Q5/5

SET 68 • HSS-ID.C.9

BACKWARD 1

Interpret the correlation coefficient • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8

$r = 0.88$.

Describe the association.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 68 • BACKWARD 1 • Q1/5

Which shows the stronger linear association, $r = -0.7$ or $r = 0.5$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 68 • BACKWARD 1 • Q2/5

What is r for points that lie exactly on a line with negative slope?

A. 1

B. 0

C. -0.5

D. -1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 68 • BACKWARD 1 • Q3/5

True or false?

The correlation coefficient has no units.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 68 • BACKWARD 1 • Q4/5

True or false?

Switching the roles of x and y changes the value of r.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.8 • Set 68 • BACKWARD 1 • Q5/5

SET 68 • HSS-ID.C.9

BACKWARD 2

Use random samples to make valid generalizations • Grade 7 • Statistics & Probability • 7.SP.A.1

Why is a random sample more likely to represent a population?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 68 • BACKWARD 2 • Q1/5

A survey at a gym asks about the exercise habits of all town residents.

What is the problem?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 68 • BACKWARD 2 • Q2/5

Which is a random sample of a school's students?

A. Drawing 40 names from a list of all students

B. Asking the first 40 students in line

C. Asking your friends

D. Asking one math class

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 68 • BACKWARD 2 • Q3/5

True or false?

Generalizations about a population are valid only if the sample represents the population.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 68 • BACKWARD 2 • Q4/5

True or false?

A sample of volunteers is a random sample.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 68 • BACKWARD 2 • Q5/5

SET 68 • HSS-ID.C.9

FORWARD 1

Distinguish surveys, experiments, and observational studies • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

Researchers record students' screen time and grades without changing anything.

What kind of study is this?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.3 • Set 68 • FORWARD 1 • Q1/5

Researchers randomly assign plants to receive fertilizer or no fertilizer.

What kind of study is this?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.3 • Set 68 • FORWARD 1 • Q2/5

What is the main purpose of a survey?

A. To test whether a treatment causes an effect

B. To assign treatments to subjects

C. To control lurking variables

D. To estimate a characteristic of a population from a sample

---

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S68-F1-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.3 • Set 68 • FORWARD 1 • Q3/5

True or false?

Random selection lets you generalize to a population; random assignment lets you conclude cause and effect.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions

HSS-IC.B.3 • Set 68 • FORWARD 1 • Q4/5

True or false?

In an observational study, the researchers decide which treatment each subject gets.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Making Inferences & Justifying Conclusions HSS-IC.B.3 • Set 68 • FORWARD 1 • Q5/5

SET 68 • HSS-ID.C.9

FORWARD 2

Use random methods to make fair decisions • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6

Describe a fair way to split 20 volunteers into two groups of 10 using random numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions HSS-MD.B.6 • Set 68 • FORWARD 2 • Q1/5

A coach picks a captain by drawing one name from a hat with 12 names.

What is the probability that a given player is chosen?

Probability: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.6 • Set 68 • FORWARD 2 • Q2/5

Which method is fair for deciding who goes first among 4 players?

A. The tallest goes first

B. Each rolls a number cube; ties roll again; the highest goes first

C. The coach's favorite goes first

D. Whoever asks first goes first

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S68-F2-Q3**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.6 • Set 68 • FORWARD 2 • Q3/5

True or false?

A fair lottery gives every participant the same probability of being chosen.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q4

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.6 • Set 68 • FORWARD 2 • Q4/5

True or false?

Flipping one fair coin once is a fair way to choose one person from 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.6 • Set 68 • FORWARD 2 • Q5/5