ALGEBRA II · COMMON CORE HIGH SCHOOL MATH

Question Collection

Main • Backward Branches • Forward 1 • Forward 2 (later courses)

84 question sets • 2525 questions

CONTENTS

Set 1	**HSN-CN.A.1**	Write complex numbers in the form a + bi	7
Set 2	**HSN-CN.A.2**	Add and subtract complex numbers	44
Set 3	**HSN-CN.A.2**	Multiply complex numbers and simplify powers of i	81
Set 4	**HSN-CN.C.7**	Solve quadratic equations with complex solutions	118
Set 5	**HSN-CN.C.8**	Factor polynomials over the complex numbers	155
Set 6	**HSN-CN.C.9**	Use the Fundamental Theorem of Algebra	192
Set 7	**HSA-SSE.A.1.a**	Interpret the parts of polynomial, rational, and exponential expressions in context	229
Set 8	**HSA-SSE.A.1.b**	Interpret complicated expressions by viewing parts as single entities	266
Set 9	**HSA-SSE.A.2**	Factor higher-degree expressions by recognizing differences of squares and sums	303
Set 10	**HSA-SSE.A.2**	Rewrite expressions in quadratic form and factor	340
Set 11	**HSA-SSE.B.4**	Derive the formula for the sum of a finite geometric series	377
Set 12	**HSA-SSE.B.4**	Use the finite geometric series formula to solve problems	414
Set 13	**HSA-APR.A.1**	Add and subtract polynomials of higher degree	451
Set 14	**HSA-APR.A.1**	Multiply polynomials of higher degree	488
Set 15	**HSA-APR.B.2**	Find remainders with the Remainder Theorem	525
Set 16	**HSA-APR.B.2**	Use p(a) = 0 to find factors	562
Set 17	**HSA-APR.B.3**	Identify zeros and multiplicities from a factorization	599

CONTENTS (continued)

Set 18	**HSA-APR.B.3**	Construct a rough graph of a polynomial from its zeros	636
Set 19	**HSA-APR.C.4**	Prove polynomial identities	679
Set 20	**HSA-APR.C.4**	Use polynomial identities to describe numerical relationships	716
Set 21	**HSA-APR.C.5**	Expand with the Binomial Theorem	753
Set 22	**HSA-APR.D.6**	Rewrite rational expressions by polynomial division	790
Set 23	**HSA-APR.D.6**	Rewrite simple rational expressions by inspection	827
Set 24	**HSA-APR.D.7**	Multiply and divide rational expressions	864
Set 25	**HSA-APR.D.7**	Add and subtract rational expressions	901
Set 26	**HSA-CED.A.1**	Create and solve rational, radical, and exponential equations from contexts	938
Set 27	**HSA-CED.A.1**	Create and solve nonlinear inequalities from contexts	975
Set 28	**HSA-CED.A.2**	Create and graph nonlinear two-variable models	1012
Set 29	**HSA-CED.A.3**	Represent nonlinear constraints and judge viable options	1049
Set 30	**HSA-CED.A.4**	Rearrange rational, radical, and exponential formulas	1086
Set 31	**HSA-REI.A.2**	Solve rational equations and identify extraneous solutions	1123
Set 32	**HSA-REI.A.2**	Solve radical equations and identify extraneous solutions	1160
Set 33	**HSA-REI.D.11**	Explain why intersections of nonlinear graphs solve f(x) = g(x)	1197
Set 34	**HSA-REI.D.11**	Approximate solutions of f(x) = g(x)	1234

CONTENTS (continued)

Set 35	**HSF-TF.A.1**	Use radian measure as arc length on the unit circle	1271
Set 36	**HSF-TF.A.2**	Find exact trigonometric values beyond the first quadrant	1308
Set 37	**HSF-TF.A.2**	Explain the extension of sine and cosine to all real numbers	1345
Set 38	**HSF-TF.B.5**	Write sinusoidal models with given amplitude, period, and midline	1382
Set 39	**HSF-TF.B.5**	Interpret the parameters of sinusoidal models	1419
Set 40	**HSF-TF.C.8**	Prove the Pythagorean identity	1456
Set 41	**HSF-TF.C.8**	Use the Pythagorean identity to find trigonometric values	1493
Set 42	**HSF-IF.B.4**	Interpret key features of polynomial and rational models	1530
Set 43	**HSF-IF.B.4**	Interpret periodic models from graphs and tables	1567
Set 44	**HSF-IF.B.5**	State domains of radical, rational, logarithmic, and polynomial models	1604
Set 45	**HSF-IF.B.6**	Calculate and interpret average rates of change of nonlinear models	1641
Set 46	**HSF-IF.C.7.b**	Graph square root and cube root functions with transformations	1678
Set 47	**HSF-IF.C.7.b**	Graph piecewise-defined and absolute value functions	1715
Set 48	**HSF-IF.C.7.c**	Graph polynomial functions showing zeros and end behavior	1752
Set 49	**HSF-IF.C.7.e**	Graph exponential functions with intercepts, asymptote, and end behavior	1789
Set 50	**HSF-IF.C.7.e**	Graph logarithmic functions with intercept, asymptote, and end behavior	1826
Set 51	**HSF-IF.C.7.e**	Graph trigonometric functions showing period, midline, and amplitude	1863

CONTENTS (continued)

Set 52	**HSF-IF.C.8.b**	Rewrite exponential functions to reveal rates over other time units	1900
Set 53	**HSF-IF.C.9**	Compare properties of Algebra II functions in different representations	1937
Set 54	**HSF-BF.A.1.b**	Build models by combining Algebra II function types	1974
Set 55	**HSF-BF.B.3**	Identify vertical shifts, vertical stretches, and horizontal shifts of Algebra II functio	2011
Set 56	**HSF-BF.B.3**	Horizontal stretches, compressions, and reflections f(kx)	2048
Set 57	**HSF-BF.B.3**	Recognize even and odd functions	2085
Set 58	**HSF-BF.B.4.a**	Write inverses of radical, rational, cubic, exponential, and logarithmic functions	2122
Set 59	**HSF-LE.A.4**	Write exact logarithmic solutions of exponential equations	2159
Set 60	**HSF-LE.A.4**	Evaluate logarithmic solutions of exponential models with technology	2196
Set 61	**HSS-ID.A.4**	Estimate population percentages with a normal distribution	2233
Set 62	**HSS-ID.A.4**	Recognize data sets for which a normal model is not appropriate	2270
Set 63	**HSS-IC.A.1**	Distinguish population, sample, parameter, and statistic	2307
Set 64	**HSS-IC.A.2**	Decide whether a model is consistent with observed results using a simulation	2344
Set 65	**HSS-IC.B.3**	Identify sample surveys, experiments, and observational studies	2381
Set 66	**HSS-IC.B.3**	Explain random sampling versus random assignment	2418
Set 67	**HSS-IC.B.4**	Estimate a population proportion with a margin of error from a simulation	2455
Set 68	**HSS-IC.B.4**	Estimate a population mean with a margin of error from a simulation	2492

CONTENTS (continued)

Set 69	**HSS-IC.B.5**	Compare two treatments with a re-randomization simulation	2529
Set 70	**HSS-IC.B.6**	Evaluate claims in reports based on data	2566
Set 71	**HSS-MD.B.6**	Design and judge fair decision procedures with probabilities	2603
Set 72	**HSS-MD.B.7**	Analyze decisions and strategies with probabilities	2640
Set 73	**HSN-CN.A.3**	Use conjugates to find moduli and quotients of complex numbers	2677
Set 74	**HSF-IF.C.7.d**	Graph rational functions with zeros and asymptotes	2714
Set 75	**HSF-IF.C.7.d**	Distinguish holes from vertical asymptotes	2751
Set 76	**HSF-BF.A.1.c**	Compose functions and interpret compositions with domains	2788
Set 77	**HSF-BF.B.4.b**	Verify inverses by composition on the stated domains	2825
Set 78	**HSF-BF.B.4.c**	Read values of an inverse function from graphs and tables	2862
Set 79	**HSF-BF.B.4.d**	Restrict a domain to make a function invertible and write the inverse	2899
Set 80	**HSF-BF.B.5**	Use the inverse relationship of exponents and logarithms to solve equations	2936
Set 81	**HSF-TF.A.3**	Derive values from special triangles and relate π - x, π + x, 2π - x to x	2973
Set 82	**HSF-TF.A.4**	Explain the symmetry and periodicity of trigonometric functions with the unit circ	3010
Set 83	**HSF-TF.B.6**	Choose and justify domain restrictions that make trigonometric functions invertib	3047
Set 84	**HSF-TF.B.7**	Solve trigonometric equations in models with inverse functions	3084

Write complex numbers in the form a + bi

MAIN	Algebra II main question	**HSN-CN.A.1**
BACKWARD 1	Simplify square roots of positive numbers	**HSN-RN.A.2**
BACKWARD 2	No real number squares to a negative number	**8.EE.A.2**
BACKWARD 3	Divide each term of a sum by a number	**6.EE.A.3**
FORWARD 1	Use equality of complex numbers to find unknown real values	**HSN-CN.A.1**
FORWARD 2	Represent complex numbers as points in the complex plane	**HSN-CN.B.4**

SET 1 • HSN-CN.A.1

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Write $\sqrt{(-49)}$ in the form bi, where b is a real number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q1/5

Write $\sqrt{(-18)}$ in the form bi with b in simplest radical form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q2/5

Write $6 - \sqrt{(-20)}$ in the form a + bi. Then state its real part and its imaginary part.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q3/5

Write $(-4 + \sqrt{(-75)}) / 2$ in the form a + bi. Then state its real part and its imaginary part.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q4/5

Write (10 + √(−64)) / 4 in the form a + bi. Then state its real part and its imaginary part.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q5/5

SET 1 • HSN-CN.A.1

BACKWARD 1

Simplify square roots of positive numbers • High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2

Simplify $\sqrt{50}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q1

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q1/5

Simplify $\sqrt{72}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q2

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q2/5

Which is √12 in simplest radical form?

A. 4√3

B. 3√2

C. 6√2

D. 2√3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S1-B1-Q3**

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q3/5

True or false?

$\sqrt{(9 + 16)} = \sqrt{9} + \sqrt{16}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q4

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q4/5

Simplify $\sqrt{28}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q5

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q5/5

SET 1 • HSN-CN.A.1

BACKWARD 2

No real number squares to a negative number • Grade 8 • Expressions & Equations • 8.EE.A.2

Find all real numbers x with $x^2 = 81$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q1/5

True or false?

$(-3)^2 = -9$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q2/5

Is there a real number x with $x^2 = -4$? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q3/5

Which number has a square equal to 36?

A. 18

B. $\sqrt{-36}$

C. -6

D. -18

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q4/5

Evaluate $(-7)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q5/5

SET 1 • HSN-CN.A.1

BACKWARD 3

Divide each term of a sum by a number • Grade 6 • Expressions & Equations • 6.EE.A.3

Write $(8 + 6\sqrt{2}) / 2$ as a sum of two terms.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q1/5

Write $(-6 + 4\sqrt{5}) / 2$ as a sum of two terms.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q2/5

True or false?

$(10 + \sqrt{7}) / 5 = 2 + \sqrt{7}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q3/5

(12 − 9√3) / 3 equals

A. 4 − 3√3

B. 4 − 9√3

C. 4 − 3√1

D. 12 − 3√3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q4/5

Write $(-4 + 2\sqrt{3}) / (-2)$ as a sum of two terms.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q5/5

SET 1 • HSN-CN.A.1

FORWARD 1

Use equality of complex numbers to find unknown real values • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

x and y are real numbers and $(x - 3) + 2yi = 5 - 8i$. Find x and y.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q1/5

a and b are real numbers and $3a + (b + 1)i = -12 + 4i$. Find a and b.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q2/5

True or false?

If x and y are real and x + yi = 7, then y = 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q3/5

p and q are real and $(p + q) + (p - q)i = 10 + 2i$. What is p?

A. 6

B. 5

C. 4

D. 8

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q4/5

m is a real number. For what value of m is $(m^2 - 4) + (m + 3)i$ a real number? What real number is it then?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q5/5

SET 1 • HSN-CN.A.1

FORWARD 2

Represent complex numbers as points in the complex plane • High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4

In the complex plane the horizontal axis is real and the vertical axis is imaginary. Give the point for $-3 + 5i$.

Point: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q1

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q1/5

Which complex number is represented by the point $(4, -2)$ in the complex plane?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q2

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q2/5

Find the distance from 0 to the point 6 + 8i in the complex plane.

Distance: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q3

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q3/5

True or false?

Every real number lies on the horizontal axis of the complex plane.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q4

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q4/5

Which number lies on the vertical axis of the complex plane?

A. -7

B. $7 + i$

C. $-7i$

D. $1 - i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S1-F2-Q5**

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q5/5

Add and subtract complex numbers

MAIN	Algebra II main question	**HSN-CN.A.2**
BACKWARD 1	Combine like terms	**7.EE.A.1**
BACKWARD 2	Add and subtract integers and decimals	**7.NS.A.1**
BACKWARD 3	Write square roots of negative numbers with i	**HSN-CN.A.1**
FORWARD 1	Find real parameters that make a sum real or imaginary	**HSN-CN.A.2**
FORWARD 2	Add complex numbers as vectors in the complex plane	**HSN-CN.B.5**

SET 2 • HSN-CN.A.2

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Simplify (3 + 5i) + (−7 + 2i). Write the answer in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q1/5

Simplify $(6 - 4i) - (-2 + 9i)$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q2/5

Simplify $(-1 + \sqrt{(-16)}) + (5 - \sqrt{(-9)})$. Write the answer in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q3/5

$z_1 = 2.5 - i$, $z_2 = -1.5 + 3i$ and $z_3 = 4i$. Find $z_1 - z_2 + z_3$ in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q4/5

Find the complex number w for which $(4 - 7i) + w = -2 + 3i$.

$w =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q5/5

SET 2 • HSN-CN.A.2

BACKWARD 1

Combine like terms • Grade 7 • Expressions & Equations • 7.EE.A.1

Simplify $(3 + 5x) + (-7 + 2x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q1

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q1/5

Simplify $(6 - 4x) - (-2 + 9x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q2

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q2/5

$(2 - 3y) - (5 - y)$ equals

A. $-3 - 2y$

B. $7 - 4y$

C. $-3 + 2y$

D. $-3 - 4y$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q3

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q3/5

True or false?

$(4 + 2t) + (1 - 6t) = 5 - 4t.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q4

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q4/5

Simplify $(0.5 - x) - (1.5 + 2x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q5

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q5/5

SET 2 • HSN-CN.A.2

BACKWARD 2

Add and subtract integers and decimals • Grade 7 • The Number System • 7.NS.A.1

Evaluate $6 - (-2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q1

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q1/5

Evaluate $-4 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q2

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q2/5

$-1 - 3 + 4$ equals

A. 8

B. 0

C. -8

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q3

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q3/5

True or false?

$2.5 - (-1.5) = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q4

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q4/5

Evaluate $-2 - 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q5

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q5/5

SET 2 • HSN-CN.A.2

BACKWARD 3

Write square roots of negative numbers with i • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Write $\sqrt{(-16)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q1/5

Write $\sqrt{(-9)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q2/5

$\sqrt{(-25)}$ equals

A. -5

B. $25i$

C. $-5i$

D. $5i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S2-B3-Q3**

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q3/5

True or false?

$\sqrt{(-1)} = i$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q4/5

Write $-\sqrt{(-36)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q5/5

SET 2 • HSN-CN.A.2

FORWARD 1

Find real parameters that make a sum real or imaginary • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

k is a real number. For what value of k is $(3 + ki) + (k - 5i)$ a real number? Give that real number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q1/5

k is real. For what value of k is
$(2k - 6) + 4i - (k + 1 - i)$ a pure imaginary number (real part 0)?

$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q2/5

True or false?

The sum of a complex number and its opposite is always 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q3/5

a is real and $(a + 3i) - (2 + ai)$ is real. What is a?

A. 0

B. 2

C. 3

D. -3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q4/5

Explain why the sum of a + bi and a − bi is always a real number when a and b are real.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q5/5

SET 2 • HSN-CN.A.2

FORWARD 2

Add complex numbers as vectors in the complex plane • High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5

Plot 2 + i and 1 + 3i as points in the complex plane (real axis horizontal). What point represents their sum?

Sum point: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q1

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q1/5

The points 0, 2 + i, 1 + 3i and their sum are vertices of a quadrilateral. What kind of quadrilateral is it?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q2

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q2/5

True or false?

Subtracting w from z gives the vector from the point w to the point z.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q3

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q3/5

What is the distance between the points $5 + 2i$ and $1 - i$ in the complex plane?

A. 25

B. $\sqrt{13}$

C. 5

D. 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S2-F2-Q4**

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q4/5

Find the midpoint of the segment joining $6 - 2i$ and $-2 + 4i$, as a complex number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q5

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q5/5

Multiply complex numbers and simplify powers of i

MAIN	Algebra II main question	**HSN-CN.A.2**
BACKWARD 1	Multiply binomials	**HSA-APR.A.1**
BACKWARD 2	Use $i^2 = -1$ and the cycle of powers of i	**HSN-CN.A.1**
BACKWARD 3	Divide whole numbers to find a remainder	**4.NBT.B.6**
FORWARD 1	Find a real parameter that makes a product real	**HSN-CN.A.2**
FORWARD 2	Divide complex numbers using conjugates	**HSN-CN.A.3**

SET 3 • HSN-CN.A.2

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Multiply $(3 + 2i)(4 - 5i)$. Write the product in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q1/5

Simplify $(-2 + i)^2$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q2/5

Multiply $(5 - 3i)(5 + 3i)$. Write the product in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q3/5

Simplify $(1 + i)^3$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q4/5

Simplify $i^{23} \cdot (4 - i)$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q5/5

SET 3 • HSN-CN.A.2

BACKWARD 1

Multiply binomials • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(3 + 2x)(4 - 5x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q1/5

Expand $(-2 + x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q2/5

$(5 - 3x)(5 + 3x)$ equals

A. $25 + 9x^2$

B. $10 - 9x^2$

C. $25 - 9x^2$

D. $25 - 30x - 9x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S3-B1-Q3**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q3/5

True or false?

$(1 + x)^2 = 1 + x^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q4/5

Multiply $x(4 - x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q5/5

SET 3 • HSN-CN.A.2

BACKWARD 2

Use $i^2 = -1$ and the cycle of powers of i • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Simplify i^2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q1/5

Simplify i^3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q2/5

Simplify i^4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q3/5

True or false?

$-10i^2 = -10$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q4/5

Which equals 1?

A. i^8

B. i^9

C. i^3

D. i^6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q5/5

SET 3 • HSN-CN.A.2

BACKWARD 3

Divide whole numbers to find a remainder • Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6

Find the remainder when 23 is divided by 4.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q1

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q1/5

Find the remainder when 50 is divided by 4.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q2

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q2/5

What is the remainder when 36 is divided by 4?

A. 0

B. 4

C. 1

D. 9

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q3

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q3/5

True or false?

The remainder when 101 is divided by 4 is 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q4

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q4/5

Write 39 as 4q + r with 0 ≤ r < 4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q5

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q5/5

SET 3 • HSN-CN.A.2

FORWARD 1

Find a real parameter that makes a product real • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

a is a real number. For what value of a is $(a + 2i)(3 - i)$ a real number? Give the product.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q1/5

b is real. For what value of b is (2 + bi)(1 + i) pure imaginary (real part 0)?

b = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q2/5

True or false?

The product of two non-real complex numbers can be a real number.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q3/5

For real c, $(c + i)^2$ is real when

A. $c = 0$

B. $c = 1$

C. $c = -1$

D. never

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q4/5

Show that $(a + bi)(a - bi)$ is a real number for all real a and b.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q5/5

SET 3 • HSN-CN.A.2

FORWARD 2

Divide complex numbers using conjugates • High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3

Write (3 + 4i) / (1 + 2i) in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q1

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q1/5

Write $10 / (3 - i)$ in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q2

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q2/5

True or false?

To divide by $2 - 5i$, multiply numerator and denominator by $2 + 5i$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q3

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q3/5

(2 + i) / i equals

A. $-1 + 2i$

B. $2 + i$

C. $1 - 2i$

D. $1 + 2i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S3-F2-Q4**

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q4/5

Explain why multiplying by the conjugate of the denominator always gives a real denominator.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q5

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q5/5

Solve quadratic equations with complex solutions

MAIN	Algebra II main question	**HSN-CN.C.7**
BACKWARD 1	Use the quadratic formula and the discriminant	**HSA-REI.B.4.b**
BACKWARD 2	Write square roots of negative numbers with i	**HSN-CN.A.1**
BACKWARD 3	Solve equations of the form $(x - h)^2 = k$	**HSA-REI.B.4.b**
FORWARD 1	Find a quadratic from one complex solution	**HSN-CN.C.7**
FORWARD 2	Classify solutions for a family of quadratics	**HSN-CN.C.7**

SET 4 • HSN-CN.C.7

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^2 + 49 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q1/5

Solve $x^2 - 4x + 13 = 0$ over the complex numbers. Give exact solutions in the form a + bi.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q2/5

Solve $2x^2 + 2x + 5 = 0$ over the complex numbers. Give exact solutions in the form a + bi.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q3/5

Solve $(x + 3)^2 = -20$ over the complex numbers. Give exact solutions in the form a + bi.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q4/5

Solve $3x^2 - 6x + 7 = 0$ over the complex numbers. Give exact solutions in the form a + bi.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q5/5

SET 4 • HSN-CN.C.7

BACKWARD 1

Use the quadratic formula and the discriminant • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Find the discriminant $b^2 - 4ac$ of $x^2 - 4x + 13 = 0$.

Discriminant: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q1/5

Find the discriminant of $2x^2 + 2x + 5 = 0$.

Discriminant: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q2/5

True or false?

If the discriminant is negative, the quadratic equation has no real solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q3/5

In $3x^2 - 6x + 7 = 0$, the values of a, b and c are

A. 3, −6, −7

B. 3, 6, 7

C. −3, −6, 7

D. 3, −6, 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q4/5

Write the quadratic formula.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q5/5

SET 4 • HSN-CN.C.7

BACKWARD 2

Write square roots of negative numbers with i • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Write $\sqrt{(-36)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q1/5

Write $\sqrt{(-48)}$ in simplest form with i.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q2/5

$\sqrt{(-20)}$ equals

A. $2\sqrt{5}\,i$

B. $4\sqrt{5}\,i$

C. $-2\sqrt{5}$

D. $10i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q3/5

True or false?

$\sqrt{(-49)} = -7$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q4/5

Write $(-2 + \sqrt{(-36)}) / 4$ in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q5/5

SET 4 • HSN-CN.C.7

BACKWARD 3

Solve equations of the form $(x - h)^2 = k$ • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Solve $(x - 2)^2 = 9$ over the real numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q1/5

Solve $(x + 1)^2 = 5$ over the real numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q2/5

True or false?

Solving $x^2 = 16$ gives only $x = 4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q3/5

The solutions of $(x + 4)^2 = 25$ are

A. 5 and −5

B. −1 and 9

C. 1 and −9

D. 21 and −29

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q4/5

Solve $x^2 = 50$ over the real numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q5/5

SET 4 • HSN-CN.C.7

FORWARD 1

Find a quadratic from one complex solution • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

$3 - 2i$ is a solution of $x^2 + bx + c = 0$, where b and c are real. Find b and c.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q1/5

$-1 + 4i$ is a solution of a quadratic equation $x^2 + bx + c = 0$ with real b and c. Write the equation.

Equation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q2/5

True or false?

If 2 + i is a solution of a quadratic with real coefficients, then 2 − i is also a solution.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q3/5

Which quadratic with real coefficients has 5i as a solution?

A. $x^2 + 25 = 0$

B. $x^2 + 5 = 0$

C. $x^2 - 25 = 0$

D. $x^2 - 10x + 25 = 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q4/5

Explain why $x^2 - (2 + i)x = 0$ has the solution $2 + i$ but not the solution $2 - i$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q5/5

SET 4 • HSN-CN.C.7

FORWARD 2

Classify solutions for a family of quadratics • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

For which real values of k does $x^2 + kx + 9 = 0$ have two non-real solutions?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q1/5

For which real values of c does $x^2 - 4x + c = 0$ have non-real solutions? Give them for c = 8.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q2/5

True or false?

$x^2 + kx - 1 = 0$ has non-real solutions for some real k.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q3/5

$x^2 + 6x + m = 0$ has exactly one (repeated) real solution when $m =$

A. 3

B. 36

C. -9

D. 9

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q4/5

For $c > 4$, write the solutions of $x^2 - 4x + c = 0$ in terms of c.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q5/5

Factor polynomials over the complex numbers

MAIN	Algebra II main question	**HSN-CN.C.8**
BACKWARD 1	Factor a difference of squares	**HSA-SSE.A.2**
BACKWARD 2	Multiply complex conjugates	**HSN-CN.A.2**
BACKWARD 3	Solve quadratics with complex solutions	**HSN-CN.C.7**
FORWARD 1	Factor cubic and quartic polynomials over the complex numbers	**HSN-CN.C.8**
FORWARD 2	Write polynomials with real coefficients from complex zeros	**HSN-CN.C.9**

SET 5 • HSN-CN.C.8

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

Factor $x^2 + 25$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q1/5

Factor $x^2 + 7$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q2/5

Factor $4x^2 + 9$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q3/5

Factor $x^4 - 16$ completely into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q4/5

Factor $x^2 - 6x + 13$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q5/5

SET 5 • HSN-CN.C.8

BACKWARD 1

Factor a difference of squares • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 - 49$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q1/5

Factor $9x^2 - 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q2/5

$x^4 - 1$ equals

A. $(x - 1)^4$

B. $(x^2 - 1)^2$

C. $(x^2 - 1)(x^2 + 1)$

D. $(x^2 + 1)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S5-B1-Q3**

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q3/5

True or false?

$x^2 + 9 = (x + 3)(x - 3)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q4/5

Factor $x^2 - 5$ using radicals.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q5/5

SET 5 • HSN-CN.C.8

BACKWARD 2

Multiply complex conjugates • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Multiply $(x + 5i)(x - 5i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q1/5

Multiply $(2x + 3i)(2x - 3i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q2/5

True or false?

$(3i)^2 = -9$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q3/5

$(x - (1 + i))(x - (1 - i))$ equals

A. $x^2 + 2x + 2$

B. $x^2 - 2x + 2$

C. $x^2 + 2$

D. $x^2 - 2x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q4/5

Simplify $(i\sqrt{7})^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q5/5

SET 5 • HSN-CN.C.8

BACKWARD 3

Solve quadratics with complex solutions • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^2 + 25 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q1/5

Solve $x^2 - 6x + 13 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q2/5

True or false?

If r is a zero of p(x), then x − r is a factor of p(x).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q3/5

The zeros of $x^2 + 4$ are

A. $2i$ and $-2i$

B. $4i$ and $-4i$

C. -4

D. 2 and -2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q4/5

Solve $x^2 + 2x + 5 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q5/5

SET 5 • HSN-CN.C.8

FORWARD 1

Factor cubic and quartic polynomials over the complex numbers • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

Factor $x^3 + 8$ completely into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q1/5

Factor $x^3 + 9x$ completely into linear factors.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q2/5

Factor $x^4 + 5x^2 + 4$ completely into linear factors.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q3/5

True or false?

$x^3 - 1 = (x - 1)(x^2 + x + 1)$, and $x^2 + x + 1$ has two non-real zeros.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S5-F1-Q4**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q4/5

Which is a linear factor of $x^3 - 27$ over the complex numbers?

A. $x + 3i$

B. $x + 3$

C. $x - 3i$

D. $x - (-3/2 + (3\sqrt{3}/2)i)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S5-F1-Q5**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q5/5

SET 5 • HSN-CN.C.8

FORWARD 2

Write polynomials with real coefficients from complex zeros • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9

Write a polynomial with real coefficients, leading coefficient 1 and least degree, whose zeros include 2i and 1 − i.

p(x) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q1/5

What is the least degree of a polynomial with real coefficients that has zeros 3, 4i and 2 + i?

Degree: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q2/5

True or false?

A polynomial with real coefficients can have exactly one non-real zero.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q3/5

A cubic with real coefficients has zeros 5 and $-i$. Its third zero is

A. -5

B. $5i$

C. 1

D. i

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S5-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q4/5

Explain why a polynomial with real coefficients and degree 3 must have at least one real zero.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q5/5

SET 6 • HSN-CN.C.9 • THE COMPLEX NUMBER SYSTEM

Use the Fundamental Theorem of Algebra

MAIN	Algebra II main question	**HSN-CN.C.9**
BACKWARD 1	Find the degree of a polynomial	**HSA-SSE.A.1.a**
BACKWARD 2	Find zeros from factored form	**HSA-SSE.B.3.a**
BACKWARD 3	Solve quadratics with complex solutions	**HSN-CN.C.7**
FORWARD 1	Relate the complex zeros to the real x-intercepts of a graph	**HSA-APR.B.3**
FORWARD 2	Prove existence statements about real zeros	**HSN-CN.C.9**

SET 6 • HSN-CN.C.9

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9

How many complex zeros, counted with multiplicity, does $p(x) = 3x^5 - x^2 + 7$ have? Name the theorem you used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q1/5

List every complex zero of $p(x) = (x - 2)^3(x^2 + 9)$ with its multiplicity, and check that the multiplicities add to the degree.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q2/5

Show that the Fundamental Theorem of Algebra holds for every quadratic $p(x) = ax^2 + bx + c$ with real a, b, c and $a \neq 0$: p has exactly two complex zeros counted with multiplicity.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q3/5

Find every complex zero of $p(x) = x^4 - 81$ and check that the count agrees with the Fundamental Theorem of Algebra.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q4/5

A student says $p(x) = x^3 + x + 1$ has no zeros because it does not factor with integers. Use the Fundamental Theorem of Algebra to explain how many complex zeros it has, and explain why at least one of them is real.

Work space

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S6-M-Q5**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q5/5

SET 6 • HSN-CN.C.9

BACKWARD 1

Find the degree of a polynomial • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

What is the degree of $3x^5 - x^2 + 7$?

Degree: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q1/5

What is the degree of $(x - 2)^3(x^2 + 9)$ when multiplied out?

Degree: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q2/5

The degree of $x(x + 1)(x^2 - 4)$ is

A. 1

B. 4

C. 2

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S6-B1-Q3**

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q3/5

True or false?

The degree of $4x^3 + 2x^4$ is 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q4/5

What is the leading coefficient of $-2x^3 + 5x - 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q5/5

SET 6 • HSN-CN.C.9

BACKWARD 2

Find zeros from factored form • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a

Find the zeros of $(x - 2)(x + 5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q1/5

Find the zeros of $(x - 1)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q2/5

True or false?

A product is 0 only when one of its factors is 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q3/5

The zeros of $x(x + 4)$ are

A. 0 and 4

B. 0 and -4

C. -4 only

D. 4 only

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q4/5

Find the zeros of $(2x - 3)(x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q5/5

SET 6 • HSN-CN.C.9

BACKWARD 3

Solve quadratics with complex solutions • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^2 + 9 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q1/5

Solve $x^2 + 2x + 2 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q2/5

True or false?

$x^2 + 1 = 0$ has no complex solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q3/5

The solutions of $x^2 - 2x + 5 = 0$ are

A. $1 \pm 4i$

B. $-1 \pm 2i$

C. $1 \pm 2i$

D. $2 \pm i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q4/5

Solve $x^2 = -81$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q5/5

SET 6 • HSN-CN.C.9

FORWARD 1

Relate the complex zeros to the real x-intercepts of a graph • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

A polynomial of degree 4 with real coefficients has a graph that crosses the x-axis at exactly two points and touches it nowhere else. Each crossing is a simple zero. How many non-real zeros does it have?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q1/5

The graph of $y = x^3 - 4x^2 + 5x - 2$ touches the x-axis at $x = 1$ and crosses it at $x = 2$. List all complex zeros with multiplicity.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q2/5

True or false?

A polynomial of degree 6 whose graph never meets the x-axis has 6 non-real zeros.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q3/5

A degree-5 polynomial with real coefficients has exactly one real zero, of multiplicity 1. How many non-real zeros does it have?

A. 4

B. 5

C. 2

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S6-F1-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q4/5

Can a degree-4 polynomial with real coefficients have exactly three real zeros, each of multiplicity 1? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q5/5

SET 6 • HSN-CN.C.9

FORWARD 2

Prove existence statements about real zeros • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9

Prove that every polynomial of odd degree with real coefficients has at least one real zero.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S6-F2-Q1**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q1/5

A degree-4 polynomial with real coefficients has the zeros 1 + i and 3. What are all of its possible zero lists?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q2/5

True or false?

A degree-4 polynomial with real coefficients may have no real zeros.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q3/5

A polynomial with real coefficients has degree 7. The possible numbers of real zeros (with multiplicity) are

A. 1, 3, 5, 7

B. 7 only

C. 0, 2, 4, 6

D. 0 to 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S6-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q4/5

Explain why the Fundamental Theorem of Algebra says nothing about whether a polynomial's zeros can be found exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q5/5

Interpret the parts of polynomial, rational, and exponential expressions in context

MAIN	Algebra II main question	**HSA-SSE.A.1.a**
BACKWARD 1	Name terms, factors, and coefficients	**HSA-SSE.A.1.a**
BACKWARD 2	Translate between a percent change and a multiplying factor	**7.RP.A.3**
BACKWARD 3	Interpret a quotient as an amount per unit	**6.RP.A.2**
FORWARD 1	Build an expression whose parts have given meanings	**HSA-SSE.A.1.a**
FORWARD 2	Rewrite an exponential expression to reveal a different rate	**HSF-IF.C.8.b**

SET 7 • HSA-SSE.A.1.a

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

A town's population is modeled by $P(t) = 12{,}500(0.97)^t$, where t is the number of years after 2020. What do 12,500 and 0.97 represent, and what does $1 - 0.97$ represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q1/5

The average cost per item of making x items is $C(x) = (4{,}000 + 6x) / x$ dollars. Interpret 4,000, 6x, and the division by x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q2/5

An open box is made by cutting squares of side x cm from the corners of a 20 cm by 20 cm sheet. Its volume is $V(x) = x(20 - 2x)^2$ cm³. Interpret x, $20 - 2x$, and $(20 - 2x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q3/5

The amount of a medicine in the body t hours after a dose is $A(t) = 200(1/2)^{t/6}$ mg. Interpret 200, 1/2, and t/6.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q4/5

Maya keeps $1,500 in an account that earns 4% interest compounded once a year and also puts $200 per year in a jar (no interest). Her total after n years is $T(n) = 1500(1.04)^n + 200n$. Interpret $1500(1.04)^n$, 1.04, and 200n.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q5/5

SET 7 • HSA-SSE.A.1.a

BACKWARD 1

Name terms, factors, and coefficients • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

Name the terms of 4,000 + 6x.

Terms: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q1/5

What is the coefficient of x in $7x - 3$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q2/5

In $5x(x - 2)$, which is a factor?

A. $5x - 2$

B. $x - 10$

C. $x - 2$

D. -2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q3/5

True or false?

In 3 + 2x, the number 3 is a factor.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q4/5

Name the factors of $200(0.5)^t$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q5/5

SET 7 • HSA-SSE.A.1.a

BACKWARD 2

Translate between a percent change and a multiplying factor • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

A price is multiplied by 0.97. By what percent does it change?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q1/5

What factor multiplies an amount that increases by 4%?

Factor: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q2/5

Multiplying by 1.25 is the same as

A. increasing by 2.5%

B. increasing by 25%

C. increasing by 125%

D. decreasing by 25%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q3/5

True or false?

Multiplying by 0.8 decreases an amount by 80%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q4/5

Find $1 - 0.85$ and explain what it means for an amount multiplied by 0.85.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q5/5

SET 7 • HSA-SSE.A.1.a

BACKWARD 3

Interpret a quotient as an amount per unit • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2

A total cost of $600 is shared by 30 people. What does 600/30 represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q1/5

A car travels 150 miles in 3 hours. What does 150/3 represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q2/5

If T is a total cost for n items, then T/n is

A. the profit

B. the number of items

C. the cost per item

D. the total cost

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q3/5

True or false?

Dividing a total by the number of items gives an average per item.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q4/5

What does (4,000 + 6x)/x mean when x = 100?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q5/5

SET 7 • HSA-SSE.A.1.a

FORWARD 1

Build an expression whose parts have given meanings • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

Write an expression for the value of a $24,000 car that loses 15% of its value each year, t years after purchase.

Expression: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q1/5

A gym charges a $90 joining fee and $35 per month. Write an expression for the average cost per month over m months.

Expression: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q2/5

True or false?

An expression for a quantity that triples every 5 years, starting at 40, is $40(3)^{t/5}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q3/5

A bacteria count starts at 500 and doubles every 3 hours. Which expression gives the count after h hours?

A. $2(500)^{h/3}$

B. $500(2)^{3h}$

C. $500(3)^{h/2}$

D. $500(2)^{h/3}$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S7-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q4/5

Explain what the part 90/m of the expression 35 + 90/m means for the gym in this branch.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q5/5

SET 7 • HSA-SSE.A.1.a

FORWARD 2

Rewrite an exponential expression to reveal a different rate • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

Rewrite $12{,}500(0.97)^{t}$ (t in years) to show the monthly factor, rounded to 5 decimal places.

Expression: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q1/5

$A(t) = 200(1/2)^{t/6}$. Write A as $200 \cdot b^t$ and give b to 4 decimal places. What is the hourly percent decrease?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q2/5

True or false?

$1.04^{n} = (1.04^{1/12})^{12n}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q3/5

Which expression shows that $500(2)^{t/3}$ grows by about 26% per time unit?

A. $1.26(500)^{t}$

B. $500(1.26)^{3t}$

C. $500(1.26)^{t}$

D. $500(2.26)^{t}$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S7-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q4/5

Explain why rewriting does not change the values of the function.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q5/5

SET 8 • HSA-SSE.A.1.b • SEEING STRUCTURE IN EXPRESSIONS

Interpret complicated expressions by viewing parts as single entities

MAIN	Algebra II main question	**HSA-SSE.A.1.b**
BACKWARD 1	Squares are never negative	**8.EE.A.2**
BACKWARD 2	Powers of numbers greater than 1 and between 0 and 1	**8.EE.A.1**
BACKWARD 3	A fraction gets smaller as its denominator grows	**3.NF.A.3.d**
FORWARD 1	Find the set of all values of an expression on a domain	**HSA-SSE.A.1.b**
FORWARD 2	Use single-entity reasoning to find horizontal asymptotes	**HSF-IF.C.7.d**

SET 8 • HSA-SSE.A.1.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b

$P(r) = 5000(1 + r)^{10}$ with $r > 0$ is the value of an investment. Without calculating, explain why $P(r) > 5000$.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q1/5

The height of a ball is $h(t) = 80 - 16(t - 2)^2$ feet after t seconds. Without expanding, find the greatest height and when it occurs, and explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q2/5

$C(x) = 10 + 40/(x + 1)$ is a cost in dollars for $x \geq 0$. Explain why $C(x)$ is always more than 10 and never more than 50.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q3/5

The temperature of a drink is $T(t) = 70 - 30(0.8)^t$ °F after t minutes. Explain why T(t) is always below 70 and gets closer to 70 as t grows.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q4/5

$f(x) = 3/(x^2 + 1)$. Find the greatest value of $f(x)$ and explain why no input gives a greater value.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q5/5

SET 8 • HSA-SSE.A.1.b

BACKWARD 1

Squares are never negative • Grade 8 • Expressions & Equations • 8.EE.A.2

True or false?

$(t - 2)^2 \geq 0$ for every real t.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q1/5

What is the smallest value of $(x + 5)^2$, and for which x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q2/5

Which expression can be negative for some real x?

A. $x^2 + 1$

B. $x - 1$

C. x^2

D. $(x - 1)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q3/5

What is the smallest value of $x^2 + 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q4/5

True or false?

$-(x - 3)^2$ is never positive.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q5/5

SET 8 • HSA-SSE.A.1.b

BACKWARD 2

Powers of numbers greater than 1 and between 0 and 1 • Grade 8 • Expressions & Equations • 8.EE.A.1

Is 1.05^{10} greater than or less than 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q1/5

Is 0.8^5 greater than or less than 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q2/5

As n increases, 0.5^n

A. becomes negative

B. stays 0.5

C. gets closer to 0

D. grows without bound

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q3/5

True or false?

A positive number raised to any power is positive.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q4/5

Order from least to greatest: 0.9^2, 0.9^5, 0.9.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q5/5

SET 8 • HSA-SSE.A.1.b

BACKWARD 3

A fraction gets smaller as its denominator grows • Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d

Which is greater, 40/2 or 40/5?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q1

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q1/5

True or false?

If $x \geq 0$, then $40/(x + 1) \leq 40$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q2

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q2/5

As x grows, 6/(x + 3)

A. increases

B. becomes negative

C. decreases toward 0

D. stays 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q3

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q3/5

What is the greatest value of $12/(x + 4)$ for $x \geq 0$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q4

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q4/5

True or false?

8/10 is greater than 8/9.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q5

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q5/5

SET 8 • HSA-SSE.A.1.b

FORWARD 1

Find the set of all values of an expression on a domain • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b

Find every value that $h(t) = 80 - 16(t - 2)^2$ takes for $0 \le t \le 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q1/5

Find every value of $f(x) = 3/(x^2 + 1)$ for real x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q2/5

True or false?

For $x \geq 0$, $C(x) = 10 + 40/(x + 1)$ takes every value in $10 < C \leq 50$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q3/5

For $t \geq 0$, which set contains every value of $70 - 30(0.8)^t$?

A. $T \geq 70$

B. $40 \leq T < 70$

C. $40 < T \leq 70$

D. $0 < T < 70$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S8-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q4/5

Find every value of $g(x) = 5 - 2/(x^2 + 4)$ for real x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q5/5

SET 8 • HSA-SSE.A.1.b

FORWARD 2

Use single-entity reasoning to find horizontal asymptotes • High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d

Explain why $y = 10 + 40/(x + 1)$ has the horizontal asymptote $y = 10$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q1/5

Rewrite (2x + 7)/(x + 3) as 2 + 1/(x + 3) and state its horizontal asymptote.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q2/5

True or false?

$y = 3/(x^2 + 1)$ has the horizontal asymptote $y = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q3/5

The horizontal asymptote of $y = 5 - 2/(x - 4)$ is

A. $y = -2$

B. $y = 5$

C. $y = 4$

D. $y = 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q4/5

Does the graph of $y = 10 + 40/(x + 1)$ ever reach its horizontal asymptote? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q5/5

Factor higher-degree expressions by recognizing differences of squares and sums or differences of cubes

MAIN	Algebra II main question	**HSA-SSE.A.2**
BACKWARD 1	Take out a greatest common factor	**6.EE.A.3**
BACKWARD 2	Recognize perfect squares and perfect cubes	**8.EE.A.2**
BACKWARD 3	Multiply to check a factoring pattern	**HSA-APR.A.1**
FORWARD 1	Factor expressions whose parts are themselves expressions	**HSA-SSE.A.2**
FORWARD 2	Complete the factorization over the complex numbers	**HSN-CN.C.8**

SET 9 • HSA-SSE.A.2

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^4 - 81$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q1/5

Factor $8x^3 + 27$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q2/5

Factor $x^6 - 64$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q3/5

Factor $16x^4 - y^4$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q4/5

Factor $2x^5 - 32x$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q5/5

SET 9 • HSA-SSE.A.2

BACKWARD 1

Take out a greatest common factor • Grade 6 • Expressions & Equations • 6.EE.A.3

Factor 6x + 18.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q1/5

Factor $2x^5 - 32x$ by taking out the greatest common factor only.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q2/5

The greatest common factor of $12x^3$ and $18x$ is

A. $6x^3$

B. $6x$

C. $3x$

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q3/5

True or false?

$3x^2 + 9x = 3x(x + 3)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q4/5

Factor $5y^2 - 20y$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q5/5

SET 9 • HSA-SSE.A.2

BACKWARD 2

Recognize perfect squares and perfect cubes • Grade 8 • Expressions & Equations • 8.EE.A.2

Write $16x^4$ as a square.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q1/5

Write $8x^3$ as a cube.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q2/5

Which is a perfect cube?

A. 32

B. 16

C. 36

D. 64

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q3/5

True or false?

x^6 is both a perfect square and a perfect cube.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q4/5

Find the cube root of 27.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q5/5

SET 9 • HSA-SSE.A.2

BACKWARD 3

Multiply to check a factoring pattern • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(a + b)(a^2 - ab + b^2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q1/5

Multiply $(x - 2)(x^2 + 2x + 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q2/5

True or false?

$(x + 3)(x^2 - 6x + 9) = x^3 + 27.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q3/5

$(x^2 - 9)(x^2 + 9)$ equals

A. $x^4 - 18x^2 - 81$

B. $x^4 + 81$

C. $(x - 3)^4$

D. $x^4 - 81$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S9-B3-Q4**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q4/5

Multiply $(2x + 1)(4x^2 - 2x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q5/5

SET 9 • HSA-SSE.A.2

FORWARD 1

Factor expressions whose parts are themselves expressions • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $(x + 1)^2 - 9$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q1/5

Factor $(2x - 1)^3 + 8$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q2/5

True or false?

$a^6 - b^6$ can be factored first as a difference of squares or first as a difference of cubes.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q3/5

$(x - y)^2 - (x + y)^2$ equals

A. 0

B. $4xy$

C. $-4xy$

D. $-2y^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S9-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q4/5

Factor $x^2(x + 2) - 4(x + 2)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q5/5

SET 9 • HSA-SSE.A.2

FORWARD 2

Complete the factorization over the complex numbers • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

Factor $x^4 - 81$ completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q1/5

Factor $16x^4 - 1$ completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q2/5

True or false?

Over the complex numbers, $x^3 + 8$ has three linear factors.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q3/5

Which polynomial is fully factored over the integers but not over the complex numbers?

A. $x^2 + 9$

B. $x - 9$

C. $x^2 - 9$

D. $(x - 3)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q4

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q4/5

$2x^5 - 32x = 2x(x^2 + 4)(x + 2)(x - 2)$ over the integers. Complete the factorization over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q5/5

Rewrite expressions in quadratic form and factor

MAIN	Algebra II main question	**HSA-SSE.A.2**
BACKWARD 1	Factor quadratic trinomials	**HSA-SSE.A.2**
BACKWARD 2	Write powers as squares of other powers	**8.EE.A.1**
BACKWARD 3	Treat a repeated part as a single variable	**HSA-SSE.A.1.b**
FORWARD 1	Solve equations in quadratic form over the complex numbers	**HSN-CN.C.7**
FORWARD 2	Solve exponential equations in quadratic form	**HSF-LE.A.4**

SET 10 • HSA-SSE.A.2

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^4 - 5x^2 + 4$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q1/5

Factor $x^4 + 3x^2 - 10$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q2/5

Factor $9^x - 4 \cdot 3^x + 3$ by writing it in quadratic form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q3/5

Factor $x^6 - 9x^3 + 8$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q4/5

Factor $2x^4 - 7x^2 - 4$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q5/5

SET 10 • HSA-SSE.A.2

BACKWARD 1

Factor quadratic trinomials • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $u^2 - 5u + 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q1/5

Factor $u^2 + 3u - 10$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q2/5

$2u^2 - 7u - 4$ equals

A. $(2u + 1)(u - 4)$

B. $(u - 2)(2u + 2)$

C. $(2u + 4)(u - 1)$

D. $(2u - 1)(u + 4)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q3/5

True or false?

$u^2 - 9u + 8 = (u - 1)(u - 8)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q4/5

Factor $u^2 - 4u + 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q5/5

SET 10 • HSA-SSE.A.2

BACKWARD 2

Write powers as squares of other powers • Grade 8 • Expressions & Equations • 8.EE.A.1

Write x^4 as a power of x^2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q1/5

Write 9^x as a power of 3^x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q2/5

x^6 equals

A. $(x^4)^2$

B. $(x^3)^2$

C. $x^3 \cdot x^3 \cdot x^3$

D. $(x^2)^4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q3/5

True or false?

$4^x = (2^x)^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q4/5

Write 25^x as a power of 5^x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q5/5

SET 10 • HSA-SSE.A.2

BACKWARD 3

Treat a repeated part as a single variable • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b

In $x^4 - 5x^2 + 4$, what single variable u turns the expression into $u^2 - 5u + 4$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q1/5

Rewrite $2(x - 1)^2 + 3(x - 1) - 5$ using $u = x - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q2/5

True or false?

In $4^x - 2^x - 6$, the substitution $u = 2^x$ gives $u^2 - u - 6$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q3/5

Which substitution writes $x^6 - 9x^3 + 8$ as a quadratic?

A. $u = x^2$

B. $u = x^3$

C. $u = x^6$

D. $u = 3x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q4/5

After factoring $(u - 1)(u - 8)$ with $u = x^3$, write the result in terms of x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q5/5

SET 10 • HSA-SSE.A.2

FORWARD 1

Solve equations in quadratic form over the complex numbers • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^4 + 3x^2 - 4 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q1/5

Solve $x^4 - 5x^2 + 4 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q2/5

True or false?

$x^4 + 10x^2 + 9 = 0$ has no real solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q3/5

How many complex solutions, with multiplicity, does $x^4 - 13x^2 + 36 = 0$ have?

A. 1

B. 3

C. 4

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q4/5

Solve $x^4 + x^2 - 12 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q5/5

SET 10 • HSA-SSE.A.2

FORWARD 2

Solve exponential equations in quadratic form • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $9^x - 5 \cdot 3^x + 6 = 0$. Give exact solutions.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q1/5

Solve $4^x - 6 \cdot 2^x + 8 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q2/5

True or false?

$4^x + 2^x - 2 = 0$ has two real solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q3/5

The solutions of $25^x - 26 \cdot 5^x + 25 = 0$ are

A. 0 and 2

B. 1 and 25

C. 0 and 1

D. 1 and 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S10-F2-Q4**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q4/5

Solve $e^{2t} - 7e^{t} + 10 = 0$. Give exact solutions and decimal values to 3 decimal places.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q5/5

SET 11 • HSA-SSE.B.4 • SEEING STRUCTURE IN EXPRESSIONS

Derive the formula for the sum of a finite geometric series

MAIN	Algebra II main question	**HSA-SSE.B.4**
BACKWARD 1	Write terms of a geometric sequence	**HSF-LE.A.2**
BACKWARD 2	Distribute a factor over a sum	**6.EE.A.3**
BACKWARD 3	Use the product rule for exponents	**8.EE.A.1**
FORWARD 1	Prove the sum formula from a polynomial identity	**HSA-APR.C.4**
FORWARD 2	Describe how partial sums behave when the ratio is between -1 a...	**HSA-SSE.B.4**

SET 11 • HSA-SSE.B.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

Let $S_n = a + ar + ar^2 + \ldots + ar^{n-1}$ with $r \neq 1$. Derive $S_n = a(1 - r^n)/(1 - r)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S11-M-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q1/5

Use the multiply-and-subtract method (not the formula) to find $S = 3 + 6 + 12 + \ldots + 3 \cdot 2^9$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q2/5

Explain why the derivation of $S_n = a(1 - r^n)/(1 - r)$ does not work when $r = 1$, and find S_n in that case.

Work space

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q3/5

Derive a formula for $T = 1 + x + x^2 + \ldots + x^n$ ($n + 1$ terms, $x \neq 1$), then check it for $x = 2$ and $n = 3$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q4/5

Derive the total of 20 yearly amounts $100, 100(1.05), 100(1.05)^2, \ldots, 100(1.05)^{19}$ by the multiply-and-subtract method, and give it to the nearest cent.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q5/5

SET 11 • HSA-SSE.B.4

BACKWARD 1

Write terms of a geometric sequence • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

Write the first four terms of the geometric sequence with first term 3 and ratio 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q1/5

What is the 10th term of 3, 6, 12, ...?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q2/5

The nth term of a geometric sequence with first term a and ratio r is

A. $a + (n - 1)r$

B. $a \cdot n \cdot r$

C. ar^n

D. ar^{n-1}

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S11-B1-Q3**

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q3/5

True or false?

100, 105, 110.25, ... is geometric with ratio 1.05.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q4/5

Find the ratio of 81, 27, 9, 3, ...

Ratio: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q5/5

SET 11 • HSA-SSE.B.4

BACKWARD 2

Distribute a factor over a sum • Grade 6 • Expressions & Equations • 6.EE.A.3

Distribute: 2(3 + 6 + 12).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q1/5

Distribute: $r(a + ar + ar^2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q2/5

True or false?

$x(1 + x + x^2) = x + x^2 + x^3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q3/5

1.05(100 + 105) equals

A. 105 + 105

B. 100 + 110.25

C. 105 + 110.25

D. 205

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q4/5

Subtract: $(6 + 12 + 24 + 48) - (3 + 6 + 12 + 24)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q5/5

SET 11 • HSA-SSE.B.4

BACKWARD 3

Use the product rule for exponents • Grade 8 • Expressions & Equations • 8.EE.A.1

Simplify $r \cdot r^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q1/5

Simplify $ar^{n-1} \cdot r$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q2/5

$2 \cdot 2^9$ equals

A. 2^{10}

B. 4^{10}

C. 4^9

D. 2^9

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q3/5

True or false?

$x \cdot x^n = x^{n+1}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q4/5

Evaluate 2^{10}.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q5/5

SET 11 • HSA-SSE.B.4

FORWARD 1

Prove the sum formula from a polynomial identity • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Prove the identity

$x^n - 1 = (x - 1)(x^{n-1} + x^{n-2} + \ldots + x + 1)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q1/5

Use the identity $x^n - 1 = (x - 1)(x^{n-1} + \ldots + 1)$ to write $1 + 3 + 9 + \ldots + 3^5$ as a quotient, and evaluate it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q2/5

True or false?

The identity shows that $x - 1$ is a factor of $x^n - 1$ for every whole number $n \geq 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q3/5

Which equals $1 + 2 + 4 + \ldots + 2^9$?

A. 2^{10}

B. $2^{10} - 1$

C. $2^9 - 1$

D. $(2^9 - 1)/2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q4/5

Explain how the identity gives $a(1 - r^n)/(1 - r)$ for $a + ar + \ldots + ar^{n-1}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q5/5

SET 11 • HSA-SSE.B.4

FORWARD 2

Describe how partial sums behave when the ratio is between −1 and 1 • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

For the series $8 + 4 + 2 + 1 + \ldots$, find S_5 and S_{10} to 4 decimal places.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q1/5

Using $S_n = 16(1 - 0.5^n)$, explain why every partial sum of $8 + 4 + 2 + \ldots$ is less than 16 but can be made as close to 16 as you like.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q2/5

True or false?

For $r = 2$, the partial sums of $a + ar + ar^2 + \ldots$ $(a > 0)$ stay below a fixed number.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q3/5

For $10 - 5 + 2.5 - 1.25 + \ldots$, the partial sums get close to

A. 20/3

B. 15

C. 5

D. 20

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q4/5

A ball travels 10 m down, then rebounds and falls in a geometric pattern so the total distance after n bounces is $10 + 2 \cdot 6(1 - 0.6^n)/0.4$. What total can the distance never reach, and why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q5/5

Use the finite geometric series formula to solve problems

MAIN	Algebra II main question	**HSA-SSE.B.4**
BACKWARD 1	Identify the first term, ratio, and number of terms	**HSF-LE.A.2**
BACKWARD 2	Change a percent to a growth factor	**7.RP.A.3**
BACKWARD 3	Round results of calculations correctly	**HSN-Q.A.3**
FORWARD 1	Find the number of terms with logarithms	**HSF-LE.A.4**
FORWARD 2	Compare two payment plans with the series formula	**HSA-SSE.B.4**

SET 12 • HSA-SSE.B.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

Find the sum of the first 12 terms of 5 + 15 + 45 + ...

Sum: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q1/5

Leah deposits $250 at the end of each month into an account that earns 0.5% interest per month. The balance just after the 36th deposit is $250 + 250(1.005) + \ldots + 250(1.005)^{35}$. Find it to the nearest cent.

Balance: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q2/5

A $180,000 loan is repaid with 360 equal monthly payments P at a monthly interest rate of 0.4%. P satisfies
$180{,}000 = P/1.004 + P/1.004^2 + \ldots + P/1.004^{360}$. Use the geometric series formula to find P to the nearest cent.

P = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q3/5

A ball is dropped from 10 m and each rebound reaches 60% of the previous height. Find the total vertical distance it has traveled when it hits the ground for the 6th time, to the nearest hundredth of a meter.

Distance: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q4/5

What is the least number of terms of 2 + 6 + 18 + ... whose sum is greater than 10,000? Show the sums that decide it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q5/5

SET 12 • HSA-SSE.B.4

BACKWARD 1

Identify the first term, ratio, and number of terms • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

For 5 + 15 + 45 + ... (12 terms), give a, r, and n.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q1/5

For $250 + 250(1.005) + \ldots + 250(1.005)^{35}$, give a, r, and n.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q2/5

For 6 + 3.6 + 2.16 + 1.296 + 0.7776, r is

A. 0.4

B. 1.67

C. 0.6

D. 2.4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q3/5

True or false?

A list with exponents 0 through 35 has 35 terms.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q4/5

For $P/1.004 + P/1.004^2 + \ldots + P/1.004^{360}$, give the first term and the ratio.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q5/5

SET 12 • HSA-SSE.B.4

BACKWARD 2

Change a percent to a growth factor • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

What factor adds 0.5% interest?

Factor: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q1/5

A height is 60% of the one before. What is the ratio?

Ratio: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q2/5

0.4% per month as a factor is

A. 1.04

B. 1.004

C. 0.004

D. 1.4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q3/5

True or false?

Interest of 12% per year is a factor of 1.12.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q4/5

A loss of 15% per year is what factor?

Factor: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q5/5

SET 12 • HSA-SSE.B.4

BACKWARD 3

Round results of calculations correctly • High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3

Round 9834.0262 to the nearest cent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q1

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q1/5

Round 37.6672 to the nearest hundredth.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q2

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q2/5

True or false?

Rounding 1.005 to 1.01 before raising it to the 36th power gives the same answer to the nearest cent.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S12-B3-Q3

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q3/5

1.005^{36} is closest to

A. 1.1967

B. 1.18

C. 1.2

D. 1.036

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q4

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q4/5

Evaluate 3^{12} exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q5

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q5/5

SET 12 • HSA-SSE.B.4

FORWARD 1

Find the number of terms with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

How many end-of-month deposits of $250 at 0.5% per month are needed for the balance to reach $20,000? Use $250(1.005^n - 1)/0.005 \geq 20{,}000$.

n = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q1/5

Solve $2(3^n - 1)/2 = 3^n - 1 = 1{,}000{,}000$ for n exactly, then give the least whole number of terms whose sum is at least 1,000,000.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q2/5

True or false?

For a ratio greater than 1, more terms always give a larger sum when $a > 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q3/5

The sum $5(2^n - 1)$ first exceeds 5,000 when n =

A. 9

B. 11

C. 10

D. 1000

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q4/5

Explain why the number of terms must be rounded up, not to the nearest whole number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q5/5

SET 12 • HSA-SSE.B.4

FORWARD 2

Compare two payment plans with the series formula • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

Plan A pays $1,000 per month for 24 months. Plan B pays $500 the first month and 8% more each month after that, for 24 months. Which plan pays more in total, and by how much?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q1/5

For Plan B in this branch ($500, then 8% more each month), after how many months does its total first exceed Plan A's total of $1,000 per month for the same number of months?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q2/5

True or false?

A geometric plan with ratio greater than 1 eventually pays more in total than any plan with a fixed monthly amount.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q3/5

Plan C pays $2,000 then 5% less each month. Its total over 24 months is closest to

A. $40,000

B. $24,000

C. $48,000

D. $28,320

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q4/5

Name one assumption in comparing the plans by their totals only.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q5/5

Add and subtract polynomials of higher degree

MAIN	Algebra II main question	**HSA-APR.A.1**
BACKWARD 1	Combine like terms with the same power	**7.EE.A.1**
BACKWARD 2	Subtract a whole expression	**6.EE.A.3**
BACKWARD 3	Write polynomials in standard form	**HSA-SSE.A.1.a**
FORWARD 1	Reason about the degree of a sum or difference	**HSA-APR.A.1**
FORWARD 2	Add and subtract rational expressions	**HSA-APR.D.7**

SET 13 • HSA-APR.A.1

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Simplify $(4x^3 - 2x^2 + 7) + (-x^3 + 5x - 9)$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q1/5

Simplify $(6x^4 - x^3 + 2x) - (2x^4 + 3x^3 - x^2 + 2x - 1)$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q2/5

A company's revenue is $R(x) = -0.01x^3 + 3x^2 + 40x$ dollars and its cost is $C(x) = 0.02x^2 + 15x + 900$ dollars for x items. Write the profit $P(x) = R(x) - C(x)$ in standard form.

$P(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q3/5

Simplify $(x^5 - 3x^3 + x) - (x^5 + 2x^4 - 3x^3 - 7)$. Write the result in standard form and give its degree.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q4/5

$p(x) = 5x^3 + x^2 - 6$. Find the polynomial $q(x)$ for which $p(x) + q(x) = 2x^3 - x + 4$.

$q(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q5/5

SET 13 • HSA-APR.A.1

BACKWARD 1

Combine like terms with the same power • Grade 7 • Expressions & Equations • 7.EE.A.1

Simplify $4x^3 - x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q1

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q1/5

Simplify $-2x^2 + 5x - 3x^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q2

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q2/5

True or false?

$x^3 + x^2 = x^5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q3

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q3/5

Which pair are like terms?

A. $4x^2$ and $-x^2$

B. $4x^2$ and $4x$

C. x^3 and $3x$

D. x and 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q4

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q4/5

Simplify $7 - 9 + 2x - 2x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q5

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q5/5

SET 13 • HSA-APR.A.1

BACKWARD 2

Subtract a whole expression • Grade 6 • Expressions & Equations • 6.EE.A.3

Rewrite $-(2x^4 + 3x^3 - x^2 + 2x - 1)$ without parentheses.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q1/5

Simplify $(5 - x) - (3 - 4x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q2/5

True or false?

$a - (b - c) = a - b - c$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q3/5

$-(x^2 - 7)$ equals

A. $x^2 + 7$

B. $-x^2 - 7$

C. $-x^2 + 7$

D. $x^2 - 7$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q4/5

Simplify $10 - (x + 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q5/5

SET 13 • HSA-APR.A.1

BACKWARD 3

Write polynomials in standard form • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

Write $7 + 5x - 2x^3 + x^2$ in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q1/5

What is the leading coefficient of $3 - 4x^2 + x^4$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q2/5

The degree of $-2x^4 + x + 7$ is

A. 7

B. 1

C. 4

D. -2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q3/5

True or false?

The standard form of $2x - x^3$ is $-x^3 + 2x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q4/5

What is the constant term of $6x^3 - 2x + 9$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q5/5

SET 13 • HSA-APR.A.1

FORWARD 1

Reason about the degree of a sum or difference • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

p(x) has degree 4 and q(x) has degree 4. What degrees can p(x) + q(x) have? Give an example where the degree drops.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q1/5

For which value of k does
$(kx^3 + 2x - 1) - (5x^3 - x^2 + 3)$ have degree 2?

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q2/5

True or false?

The difference of two polynomials is always a polynomial.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q3/5

p has degree 5 and q has degree 3. The degree of $p - q$ is

A. 5

B. 2

C. 3

D. 8

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q4/5

Explain why the sum of two polynomials is a polynomial.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q5/5

SET 13 • HSA-APR.A.1

FORWARD 2

Add and subtract rational expressions • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Write 3/x + 2/(x + 1) as one rational expression, and state the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q1/5

Write $1/(x-2) - 1/(x+2)$ as one rational expression, and state the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q2/5

True or false?

Polynomials are closed under division.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q3/5

x/2 + x/3 equals

A. 5x/5

B. x/5

C. 5x/6

D. 2x/5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S13-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q4/5

Explain how adding rational expressions is like adding fractions of integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q5/5

Multiply polynomials of higher degree

MAIN	Algebra II main question	**HSA-APR.A.1**
BACKWARD 1	Distribute a monomial over a polynomial	**HSA-APR.A.1**
BACKWARD 2	Multiply powers with the same base	**8.EE.A.1**
BACKWARD 3	Multiply two binomials	**HSA-APR.A.1**
FORWARD 1	Find a coefficient of a product without full expansion	**HSA-APR.A.1**
FORWARD 2	Expand powers of binomials with the Binomial Theorem	**HSA-APR.C.5**

SET 14 • HSA-APR.A.1

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(x^2 + 3x - 2)(2x - 5)$. Write the product in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q1/5

Expand $(x - 2)^3$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q2/5

Multiply $(x^2 - x + 1)(x^2 + x + 1)$. Write the product in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q3/5

Multiply $(2x + 1)(x - 3)(x + 4)$. Write the product in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q4/5

A box has length $x + 2$, width $x - 1$, and height $2x + 3$ (in cm). Write its volume as a polynomial in standard form.

$V(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q5/5

SET 14 • HSA-APR.A.1

BACKWARD 1

Distribute a monomial over a polynomial • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $2x(x^2 + 3x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q1/5

Multiply $-5(x^2 + 3x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q2/5

True or false?

$x^2(x + 4) = x^3 + 4x^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q3/5

$-3x(x - 1)$ equals

A. $3x^2 - 3x$

B. $-3x^2 + 3x$

C. $-3x^2 - 3x$

D. $-3x - 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S14-B1-Q4**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q4/5

Multiply $x(x^2 - x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q5/5

SET 14 • HSA-APR.A.1

BACKWARD 2

Multiply powers with the same base • Grade 8 • Expressions & Equations • 8.EE.A.1

Simplify $x^2 \cdot x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q1/5

Simplify $2x \cdot 3x^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q2/5

True or false?

$x^2 \cdot x^2 = x^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q3/5

$-x \cdot x^3$ equals

A. x^4

B. $-x^3$

C. $-x^4$

D. $-x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q4/5

Simplify $(x^2)(-4x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q5/5

SET 14 • HSA-APR.A.1

BACKWARD 3

Multiply two binomials • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(x - 3)(x + 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q1/5

Multiply $(x + 2)(x - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q2/5

$(x - 2)^2$ equals

A. $x^2 + 4$

B. $x^2 - 4$

C. $x^2 - 4x + 4$

D. $x^2 - 2x + 4$

True or false?

$(2x + 1)(x - 3) = 2x^2 - 5x - 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q4/5

Multiply $(2x + 3)(x - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q5/5

SET 14 • HSA-APR.A.1

FORWARD 1

Find a coefficient of a product without full expansion • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Find the coefficient of x^2 in $(x^2 + 3x - 2)(4x^2 - x + 5)$ without expanding everything.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q1/5

For which k does $(x + k)(x^2 - 3x + 1)$ have no x^2 term?

$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q2/5

True or false?

The constant term of a product of polynomials is the product of their constant terms.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q3/5

The leading coefficient of $(2x^3 - 1)(5x^2 + x)$ is

A. 7

B. 5

C. 2

D. 10

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q4/5

Find the coefficient of x in $(x - 1)(x - 2)(x - 3)$.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q5/5

SET 14 • HSA-APR.A.1

FORWARD 2

Expand powers of binomials with the Binomial Theorem • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Expand $(x + 1)^4$ using Pascal's Triangle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q1/5

Expand $(x - 2)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q2/5

True or false?

The coefficients of $(x + y)^5$ are 1, 5, 10, 10, 5, 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q3/5

The coefficient of x^2y^2 in $(x + y)^4$ is

A. 2

B. 6

C. 4

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S14-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q4/5

Explain why the coefficients of $(x + y)^n$ are symmetric.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q5/5

Find remainders with the Remainder Theorem

MAIN	Algebra II main question	**HSA-APR.B.2**
BACKWARD 1	Evaluate a polynomial function	**HSF-IF.A.2**
BACKWARD 2	Powers of negative numbers	**7.NS.A.2**
BACKWARD 3	Dividend = divisor × quotient + remainder	**4.NBT.B.6**
FORWARD 1	Find unknown coefficients from given remainders	**HSA-APR.B.2**
FORWARD 2	Prove the Remainder Theorem	**HSA-APR.B.2**

SET 15 • HSA-APR.B.2

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Use the Remainder Theorem to find the remainder when $p(x) = x^3 - 4x^2 + 2x - 5$ is divided by $x - 3$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q1/5

Use the Remainder Theorem to find the remainder when $p(x) = 2x^4 + x^3 - 3x + 7$ is divided by $x + 2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q2/5

Use the Remainder Theorem to find the remainder when $p(x) = x^{100} + 3$ is divided by $x + 1$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q3/5

Use the Remainder Theorem to find the remainder when $p(x) = 4x^3 - 6x^2 + x - 1$ is divided by $x - 1/2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q4/5

Use the Remainder Theorem to find the remainder when $p(x) = x^4 - 5x^2 + 4x$ is divided by $x - 2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q5/5

SET 15 • HSA-APR.B.2

BACKWARD 1

Evaluate a polynomial function • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

$p(x) = x^3 - 4x^2 + 2x - 5$. Find $p(1)$.

$p(1) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q1/5

$p(x) = 2x^2 - x + 3$. Find $p(-2)$.

$p(-2) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q2/5

True or false?

If $p(x) = x^2 + 1$, then $p(-3) = -8$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q3/5

If $p(x) = x^3 - x$, then $p(2) =$

A. 4

B. 6

C. 8

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q4/5

$p(x) = 4x - 1$. Find $p(1/2)$.

$p(1/2) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q5/5

SET 15 • HSA-APR.B.2

BACKWARD 2

Powers of negative numbers • Grade 7 • The Number System • 7.NS.A.2

Evaluate $(-2)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q1/5

Evaluate $(-2)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q2/5

$(-1)^{100}$ equals

A. -100

B. -1

C. 100

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q3/5

True or false?

$-2^4 = (-2)^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q4/5

Evaluate $2(-2)^4 + (-2)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q5/5

SET 15 • HSA-APR.B.2

BACKWARD 3

Dividend = divisor × quotient + remainder • Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6

Write $47 = 5 \times q + r$ with $0 \leq r < 5$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q1

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q1/5

A division gives quotient 12 and remainder 3 with divisor 7. What was the dividend?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q2

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q2/5

True or false?

If a division leaves remainder 0, the divisor is a factor of the dividend.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q3

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q3/5

$x^3 - 1 = (x - 1)(x^2 + x + 1) + 0$. The remainder on division by $x - 1$ is

A. 0

B. 1

C. $x^2 + x + 1$

D. -1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q4

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q4/5

$p(x) = (x - 3)(x^2 - x - 1) - 8$. What is the remainder when $p(x)$ is divided by $x - 3$?

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q5

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q5/5

SET 15 • HSA-APR.B.2

FORWARD 1

Find unknown coefficients from given remainders • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

When $p(x) = x^3 + kx^2 - 4$ is divided by $x - 2$, the remainder is 12. Find k.

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q1/5

$p(x) = x^3 + ax + b$ leaves remainder 5 on division by $x - 1$ and remainder -1 on division by $x + 1$. Find a and b.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q2/5

True or false?

If p(x) leaves remainder 0 on division by $x - 4$, then $p(4) = 0$.

A. True

B. False

$p(x) = 2x^2 + kx + 1$ leaves remainder 7 on division by $x + 1$. Then $k =$

A. 4

B. -4

C. -6

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q4/5

For which k is the remainder of $x^4 - kx + 6$ divided by $x + 2$ equal to 0?

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q5/5

SET 15 • HSA-APR.B.2

FORWARD 2

Prove the Remainder Theorem • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Prove that the remainder when a polynomial p(x) is divided by $x - a$ is p(a).

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q1/5

Why must the remainder on division by $x - a$ be a constant?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q2/5

True or false?

The Remainder Theorem also works for division by $2x - 1$ with remainder $p(1/2)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q3/5

Which step uses the fact that $a - a = 0$?

A. Graphing $p(x)$

B. Dividing $p(x)$ by x

C. Factoring $p(x)$

D. Substituting $x = a$ into $p(x) = (x - a)q(x) + r$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q4/5

Explain how the proof shows that $p(a) = 0$ exactly when $x - a$ is a factor of $p(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q5/5

Use p(a) = 0 to find factors

MAIN	Algebra II main question	**HSA-APR.B.2**
BACKWARD 1	Use the Remainder Theorem	**HSA-APR.B.2**
BACKWARD 2	Factor the quadratic quotient	**HSA-SSE.A.2**
BACKWARD 3	Divide a polynomial by a linear factor	**HSA-APR.A.1**
FORWARD 1	Choose parameters so that a given binomial is a factor	**HSA-APR.B.2**
FORWARD 2	Finish factorizations over the complex numbers	**HSN-CN.C.8**

SET 16 • HSA-APR.B.2

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Show that $x - 2$ is a factor of $p(x) = x^3 - 3x^2 - 4x + 12$, then factor $p(x)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q1/5

Show that $x + 1$ is a factor of $p(x) = x^3 + 2x^2 - 5x - 6$, then factor $p(x)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q2/5

Show that $x + 3$ is a factor of $p(x) = 2x^3 + 5x^2 - 4x - 3$, then factor $p(x)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q3/5

Is $x - 4$ a factor of $p(x) = x^3 - 5x^2 + 2x + 9$? Justify with the Remainder Theorem.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q4/5

$p(x) = x^3 - 7x + 6$. Use $p(a) = 0$ to find all the zeros of p, and write p(x) in factored form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q5/5

SET 16 • HSA-APR.B.2

BACKWARD 1

Use the Remainder Theorem • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Find the remainder when $x^3 - 3x^2 - 4x + 12$ is divided by $x - 2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q1/5

Find the remainder when $x^3 + 2x^2 - 5x - 6$ is divided by $x - 1$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q2/5

True or false?

Dividing by $x + 3$ means evaluating at $x = 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q3/5

The remainder of $p(x) \div (x + 1)$ is

A. $-p(1)$

B. $p(-1)$

C. $p(0)$

D. $p(1)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q4/5

Find $p(-3)$ for $p(x) = 2x^3 + 5x^2 - 4x - 3$.

$p(-3) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q5/5

SET 16 • HSA-APR.B.2

BACKWARD 2

Factor the quadratic quotient • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 - x - 6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q1/5

Factor $2x^2 - x - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q2/5

$x^2 + x - 6$ equals

A. $(x - 3)(x + 2)$

B. $(x - 6)(x + 1)$

C. $(x + 3)(x - 2)$

D. $(x + 6)(x - 1)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q3/5

True or false?

$x^2 + 4$ factors over the integers.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q4/5

Factor $x^2 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q5/5

SET 16 • HSA-APR.B.2

BACKWARD 3

Divide a polynomial by a linear factor • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Find $q(x)$ if $x^3 - 3x^2 - 4x + 12 = (x - 2)\,q(x)$.

$q(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B3-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q1/5

Multiply $(x + 1)(x^2 + x - 6)$ to check a division.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B3-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q2/5

True or false?

If $(x - 1)(x^2 + x - 6) = x^3 - 7x + 6$, then $x^2 + x - 6$ is the quotient of $x^3 - 7x + 6 \div (x - 1)$.

A. True

B. False

$x^3 - 8 \div (x - 2)$ equals

A. $x^2 + 2x + 4$

B. $x^2 + 4$

C. $x^2 - 4$

D. $x^2 - 2x + 4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S16-B3-Q4**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q4/5

Find $q(x)$ if $2x^3 + 5x^2 - 4x - 3 = (x + 3)\, q(x)$.

$q(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B3-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q5/5

SET 16 • HSA-APR.B.2

FORWARD 1

Choose parameters so that a given binomial is a factor • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Find k so that $x - 3$ is a factor of $x^3 + kx^2 - 3x + 9$.

$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q1/5

Find a and b so that $x - 1$ and $x + 2$ are both factors of $x^3 + ax^2 + bx - 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q2/5

True or false?

If $x - 2$ and $x + 2$ are factors of $p(x)$, then $x^2 - 4$ is a factor of $p(x)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q3/5

For which k is $x + 1$ a factor of $x^4 + kx + 3$?

A. 2

B. -4

C. 4

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S16-F1-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q4/5

Write a cubic with leading coefficient 1 that has $x - 1$, $x + 1$, and $x - 4$ as factors, in standard form.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q5/5

SET 16 • HSA-APR.B.2

FORWARD 2

Finish factorizations over the complex numbers • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

$x - 1$ is a factor of $x^3 - 1$. Factor $x^3 - 1$ completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q1/5

$x + 2$ is a factor of $x^3 + 2x^2 + 9x + 18$. Factor it completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q2/5

True or false?

Every factor found with the factor theorem gives a zero of the polynomial.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q3/5

$x - 2$ is a factor of $x^3 - 2x^2 + x - 2$. The other zeros are

A. i and $-i$

B. -2

C. 1 and -1

D. $2i$ and $-2i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S16-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q4/5

$x - 3$ is a factor of $x^3 - 3x^2 + 4x - 12$. List all complex zeros.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q5/5

Identify zeros and multiplicities from a factorization

MAIN	Algebra II main question	**HSA-APR.B.3**
BACKWARD 1	Solve by the zero-product property	**HSA-REI.B.4.b**
BACKWARD 2	Factor to find hidden repeated factors	**HSA-SSE.A.2**
BACKWARD 3	An exponent counts repeated factors	**6.EE.A.1**
FORWARD 1	Build a polynomial from zeros and one more condition	**HSA-APR.B.3**
FORWARD 2	Decide what the zeros determine and what remains free	**HSA-APR.B.3**

SET 17 • HSA-APR.B.3

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

List the zeros of $p(x) = (x - 4)^2(x + 1)(2x - 3)$ and give the multiplicity of each.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q1/5

Factor $p(x) = x^3 - 6x^2 + 9x$, then list its zeros with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q2/5

List the zeros of $p(x) = (x^2 - 9)(x^2 + 2x + 1)$ with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q3/5

List the zeros of $p(x) = -2x(x + 5)^3$ with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q4/5

Factor $p(x) = x^4 - 8x^2 + 16$, then list its zeros with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q5/5

SET 17 • HSA-APR.B.3

BACKWARD 1

Solve by the zero-product property • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Solve $(x - 4)(x + 1) = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q1/5

Solve $x(2x - 3) = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q2/5

True or false?

If $(x - 2)^2 = 0$, then $x = 2$ or $x = -2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q3/5

The solutions of $(x + 5)(x - 5) = 0$ are

A. 25

B. -5 and 5

C. 0 and 5

D. 5 only

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q4/5

Solve $-2x(x + 5) = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q5/5

SET 17 • HSA-APR.B.3

BACKWARD 2

Factor to find hidden repeated factors • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 - 6x + 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q1/5

Factor $x^2 + 2x + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q2/5

$x^4 - 8x^2 + 16$ equals

A. $(x^2 + 4)^2$

B. $(x^2 - 4)^2$

C. $(x^2 - 8)^2$

D. $(x - 4)^4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q3/5

True or false?

$x^3 - 6x^2 + 9x = x(x - 3)^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q4/5

Factor $x^2 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q5/5

SET 17 • HSA-APR.B.3

BACKWARD 3

An exponent counts repeated factors • Grade 6 • Expressions & Equations • 6.EE.A.1

Write $(x - 4)^2$ as a product without an exponent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q1/5

How many factors of $(x + 5)$ are in $(x + 5)^3$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q2/5

True or false?

$(x - 2)^2(x - 2) = (x - 2)^3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q3/5

In $7(x - 1)^4$, the factor $x - 1$ appears

A. 28 times

B. 4 times

C. 1 time

D. 7 times

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q4/5

Write $(x + 1)(x + 1)(x - 3)$ with an exponent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q5/5

SET 17 • HSA-APR.B.3

FORWARD 1

Build a polynomial from zeros and one more condition • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

Write the cubic polynomial with zeros -1, 2, and 4 (each multiplicity 1) whose graph passes through $(0, 16)$.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q1/5

Write a polynomial of degree 3 with a zero of multiplicity 2 at $x = 3$, a zero at $x = -2$, and leading coefficient -1.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q2/5

True or false?

Zeros alone determine a polynomial completely.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q3/5

A quartic has zeros 1 and −1, each of multiplicity 2, and p(0) = 3. Then p(x) =

A. $-3(x-1)^2(x+1)^2$

B. $3(x^2-1)$

C. $(x-1)^2(x+1)^2$

D. $3(x-1)^2(x+1)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S17-F1-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q4/5

A polynomial has zeros 0 (multiplicity 2) and 5 (multiplicity 1), degree 3, and $p(1) = -8$. Find $p(x)$.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q5/5

SET 17 • HSA-APR.B.3

FORWARD 2

Decide what the zeros determine and what remains free • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

All polynomials with zeros 1 and 3 only (each multiplicity 1) and degree 2 have what form?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q1/5

If the degree is not fixed, give two different polynomials whose only real zeros are 1 and 3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q2/5

True or false?

Two cubic polynomials with the same three zeros (each multiplicity 1) and the same leading coefficient are equal.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q3/5

Which extra information fixes a cubic with zeros −2, 0, 2?

A. That it has degree 3

B. That 0 is a zero

C. Its end behavior only

D. Its value at x = 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q4/5

Explain why knowing only the real zeros of a degree-4 polynomial may not determine it even up to scale.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q5/5

Construct a rough graph of a polynomial from its zeros

MAIN	Algebra II main question	**HSA-APR.B.3**
BACKWARD 1	Read zeros and multiplicities	**HSA-APR.B.3**
BACKWARD 2	End behavior from the leading term	**HSF-IF.B.4**
BACKWARD 3	Find the y-intercept by evaluating at 0	**HSF-IF.A.2**
BACKWARD 4	Sign of a power of a negative or positive number	**7.NS.A.2**
FORWARD 1	Sketch graphs from standard form by factoring first	**HSF-IF.C.7.c**
FORWARD 2	Write a polynomial from its rough graph	**HSA-APR.B.3**

SET 18 • HSA-APR.B.3

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$$y = (x + 2)(x - 1)(x - 3)$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q1/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$$y = -(x + 1)^2(x - 2)$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q2/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$y = x(x - 3)^2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q3/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$y = (x+2)(x+1)(x-1)(x-3)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q4/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$$y = -(x - 1)^3(x + 2)$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 1

Read zeros and multiplicities • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

List the zeros of $(x + 2)(x - 1)(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q1/5

List the zeros of $-(x + 1)^2(x - 2)$ with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q2/5

True or false?

$x(x - 3)^2$ has a zero of multiplicity 2 at $x = 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q3/5

The zero $x = 1$ of $-(x - 1)^3(x + 2)$ has multiplicity

A. 1

B. 2

C. 4

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q4/5

List the zeros of $(x + 2)(x + 1)(x - 1)(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 2

End behavior from the leading term • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4

As x grows large and positive, what happens to x^3?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q1/5

As x grows large and negative, what happens to $-x^4$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q2/5

The leading term of $(x + 2)(x - 1)(x - 3)$ is

A. $-x^3$

B. -6

C. $3x$

D. x^3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q3/5

True or false?

A polynomial of even degree with a positive leading coefficient rises at both ends.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q4/5

What is the leading term of $-(x - 1)^3(x + 2)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 3

Find the y-intercept by evaluating at 0 • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Find y when x = 0 for $y = (x + 2)(x - 1)(x - 3)$.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q1/5

Find y when x = 0 for $y = -(x + 1)^2(x - 2)$.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q2/5

True or false?

The y-intercept of $x(x - 3)^2$ is 9.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q3/5

The y-intercept of $-(x-1)^3(x+2)$ is

A. 0

B. -2

C. 2

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q4/5

Find y when $x = 1$ for $y = x(x - 3)^2$.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 4

Sign of a power of a negative or positive number • Grade 7 • The Number System • 7.NS.A.2

Is $(-0.5)^2$ positive or negative?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q1/5

Is $(-0.5)^3$ positive or negative?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q2/5

True or false?

An even power of a negative number is positive.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q3/5

Near $x = 3$, $(x - 3)^2$ is

A. negative on both sides

B. positive on both sides

C. zero everywhere

D. positive on the left and negative on the right

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q4/5

Does $(x - 1)^3$ change sign at $x = 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q5/5

SET 18 • HSA-APR.B.3

FORWARD 1

Sketch graphs from standard form by factoring first • High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c

Factor, then sketch a rough graph showing every zero, the y-intercept, and end behavior.

$y = x^3 - x^2 - 6x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q1/5

Factor, then sketch a rough graph showing every zero, the y-intercept, and end behavior.

$$y = x^4 - 5x^2 + 4$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q2/5

True or false?

$y = x^3 + 4x$ crosses the x-axis only at 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q3/5

$y = -x^3 + 9x$ crosses the x-axis at

A. 0 and 9

B. 3 only

C. $-3, 0, 3$

D. $-9, 0, 9$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q4/5

$y = x^3 - 3x^2 + 4 = (x + 1)(x - 2)^2$. Describe its behavior at each zero.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q5/5

SET 18 • HSA-APR.B.3

FORWARD 2

Write a polynomial from its rough graph • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

A cubic's graph crosses the x-axis at −3 and touches it at 2, and passes through (0, 12). Write the polynomial.

p(x) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q1/5

A quartic rises at both ends, touches the x-axis at -1 and at 3, and passes through (0, 18). Write the polynomial.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q2/5

True or false?

A graph that touches the x-axis at x = a without crossing has a zero of even multiplicity at a.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q3/5

A cubic falls to the right and crosses the x-axis at 0, 1, and 4. Its leading coefficient is

A. zero

B. not determined by the end behavior

C. positive

D. negative

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S18-F2-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q4/5

A polynomial graph crosses at -2, touches at 1, falls to the left and rises to the right, and has y-intercept 4. Give a polynomial of least degree.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q5/5

Prove polynomial identities

MAIN	Algebra II main question	**HSA-APR.C.4**
BACKWARD 1	Expand the square of a binomial	**HSA-APR.A.1**
BACKWARD 2	Recognize a difference of squares	**HSA-SSE.A.2**
BACKWARD 3	What makes an identity	**6.EE.A.4**
FORWARD 1	Prove or disprove proposed identities	**HSA-APR.C.4**
FORWARD 2	Prove identities with the Binomial Theorem	**HSA-APR.C.5**

SET 19 • HSA-APR.C.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Prove the identity $(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q1/5

Prove the identity $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q2/5

Prove the identity $(a + b)^2 - (a - b)^2 = 4ab$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q3/5

Prove the identity $x^4 + 4 = (x^2 + 2x + 2)(x^2 - 2x + 2)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q4/5

Prove the identity $(n + 1)^3 - n^3 = 3n^2 + 3n + 1$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q5/5

SET 19 • HSA-APR.C.4

BACKWARD 1

Expand the square of a binomial • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Expand $(a + b)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q1/5

Expand $(x^2 - y^2)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q2/5

True or false?

$(a - b)^2 = a^2 - b^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q3/5

$(2xy)^2$ equals

A. $2xy^2$

B. $4xy$

C. $2x^2y^2$

D. $4x^2y^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S19-B1-Q4**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q4/5

Expand $(n + 1)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q5/5

SET 19 • HSA-APR.C.4

BACKWARD 2

Recognize a difference of squares • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $a^2 - b^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q1/5

Factor $(x^2 + 2)^2 - (2x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q2/5

True or false?

$u^2 - v^2 = (u - v)(u + v)$ for any expressions u and v.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q3/5

Which is a difference of squares?

A. $x^3 - 1$

B. $x^2 - 3x$

C. $9x^2 - 16$

D. $9x^2 + 16$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q4/5

Factor $25 - x^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q5/5

SET 19 • HSA-APR.C.4

BACKWARD 3

What makes an identity • Grade 6 • Expressions & Equations • 6.EE.A.4

Test $(a + b)^2 = a^2 + b^2$ with $a = 1$ and $b = 2$. Is it an identity?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q1/5

Explain the difference between an equation and an identity.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q2/5

True or false?

Checking an identity with three numbers proves it.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q3/5

$2(x + 3) = 2x + 6$ is

A. an identity

B. true only for $x = 3$

C. true only for $x = 0$

D. never true

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q4/5

Give values that show $a^3 + b^3 = (a + b)^3$ is not an identity.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q5/5

SET 19 • HSA-APR.C.4

FORWARD 1

Prove or disprove proposed identities • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Prove or disprove: $(a + b)^3 = a^3 + b^3 + 3ab(a + b)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • FORWARD 1 • Q1/5

Prove or disprove: $a^4 - b^4 = (a - b)^4$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • FORWARD 1 • Q2/5

True or false?

$(x - 1)(x^2 + x + 1) = x^3 - 1$ is an identity.

A. True

B. False

Which is an identity?

A. $(x + 2)^2 = x^2 + 4$

B. $(x + 1)^3 = x^3 + 1$

C. $x^2 - 4 = (x - 4)^2$

D. $(x + 2)^2 - (x - 2)^2 = 8x$

Find c so that $(x + c)^2 - (x - c)^2 = 12x$ is an identity.

c = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • FORWARD 1 • Q5/5

SET 19 • HSA-APR.C.4

FORWARD 2

Prove identities with the Binomial Theorem • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Use the Binomial Theorem with $x = y = 1$ to prove that the entries in row n of Pascal's Triangle add to 2^n.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q1/5

Use the Binomial Theorem with $x = 1$ and $y = -1$ to find $C(5,0) - C(5,1) + C(5,2) - C(5,3) + C(5,4) - C(5,5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q2/5

True or false?

$(1 + 1)^4 = 1 + 4 + 6 + 4 + 1.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q3/5

The sum of the coefficients of $(2x + 1)^3$ is

A. 9

B. 8

C. 6

D. 27

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S19-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q4/5

Use $(1 + 0.1)^3 = 1 + 3(0.1) + 3(0.01) + 0.001$ to find 1.1^3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q5/5

Use polynomial identities to describe numerical relationships

MAIN	Algebra II main question	**HSA-APR.C.4**
BACKWARD 1	Substitute values into an identity	**6.EE.A.2**
BACKWARD 2	Check a Pythagorean triple	**8.G.B.6**
BACKWARD 3	Choose a and b to fit a product	**HSA-APR.C.4**
FORWARD 1	Generate a triple with a required side	**HSA-APR.C.4**
FORWARD 2	Connect the triple identity to complex numbers	**HSN-CN.A.2**

SET 20 • HSA-APR.C.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

The identity $(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$ generates Pythagorean triples. Use $x = 3$ and $y = 2$ to find a triple.

Triple: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q1/5

Use the identity $(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$ with $x = 5$ and $y = 2$ to find a Pythagorean triple.

Triple: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q2/5

Use the identity $(a + b)(a - b) = a^2 - b^2$ to compute 47×53 mentally. Show which a and b you use.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q3/5

Use the identity $(n + 1)^2 - n^2 = 2n + 1$ to write 41 as a difference of two consecutive squares.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q4/5

Use the identity $(n + 1)^3 - n^3 = 3n^2 + 3n + 1$ to compute $13^3 - 12^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q5/5

SET 20 • HSA-APR.C.4

BACKWARD 1

Substitute values into an identity • Grade 6 • Expressions & Equations • 6.EE.A.2

Evaluate $x^2 - y^2$ for $x = 3$ and $y = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q1

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q1/5

Evaluate 2xy for x = 5 and y = 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q2

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q2/5

True or false?

For $n = 12$, $3n^2 + 3n + 1 = 469$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q3

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q3/5

For $a = 50$ and $b = 3$, $a^2 - b^2$ is

A. 2491

B. 2500

C. 2494

D. 2509

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q4

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q4/5

Evaluate 2n + 1 for n = 20.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q5

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q5/5

SET 20 • HSA-APR.C.4

BACKWARD 2

Check a Pythagorean triple • Grade 8 • Geometry • 8.G.B.6

Do 5, 12, 13 satisfy $a^2 + b^2 = c^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q1

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q1/5

Do 20, 21, 29 satisfy $a^2 + b^2 = c^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q2

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q2/5

True or false?

6, 8, 11 is a Pythagorean triple.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q3

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q3/5

Which is a Pythagorean triple?

A. 8, 15, 17

B. 6, 7, 9

C. 2, 3, 4

D. 4, 5, 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q4

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q4/5

Explain why a triangle with sides 5, 12, 13 is a right triangle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q5

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q5/5

SET 20 • HSA-APR.C.4

BACKWARD 3

Choose a and b to fit a product • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Write 47 and 53 as $50 - b$ and $50 + b$. What is b?

b = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B3-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q1/5

Write 98×102 as $(a - b)(a + b)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B3-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q2/5

True or false?

$29 \times 31 = 30^2 - 1$.

A. True

B. False

64 × 56 equals

A. $(60 - 4)^2$

B. $64^2 - 56^2$

C. $60^2 - 4^2$

D. $60^2 + 4^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S20-B3-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q4/5

Compute 98×102.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B3-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q5/5

SET 20 • HSA-APR.C.4

FORWARD 1

Generate a triple with a required side • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Use the triple identity to find x and y (whole numbers, x > y) that give a triple with hypotenuse 25 and legs 7 and 24.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q1/5

Find x and y (whole numbers, x > y) for which the identity gives a triple with a leg 2xy = 40 and hypotenuse 41.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q2/5

True or false?

Every choice of whole numbers $x > y > 0$ in the identity gives a Pythagorean triple.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q3/5

Which choice gives the triple 8, 15, 17?

A. $x = 4, y = 1$

B. $x = 5, y = 3$

C. $x = 4, y = 2$

D. $x = 3, y = 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q4/5

Explain why $x = 2y$ gives a triple with legs in the ratio $3 : 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q5/5

SET 20 • HSA-APR.C.4

FORWARD 2

Connect the triple identity to complex numbers • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Compute $(3 + 2i)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q1/5

Compute $(x + yi)^2$ in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q2/5

True or false?

The real and imaginary parts of $(x + yi)^2$ are the legs of the Pythagorean triple from the identity.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q3/5

$(5 + 2i)^2$ equals

A. $25 + 4i$

B. $21 + 20i$

C. $21 + 10i$

D. $29 + 20i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S20-F2-Q4**

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q4/5

Use $(4 + i)^2$ to find a Pythagorean triple.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q5/5

Expand with the Binomial Theorem

MAIN	Algebra II main question	**HSA-APR.C.5**
BACKWARD 1	Multiply binomials repeatedly	**HSA-APR.A.1**
BACKWARD 2	Build Pascal's Triangle by a rule	**4.OA.C.5**
BACKWARD 3	Powers of products and of negatives	**8.EE.A.1**
FORWARD 1	Find a specific term or coefficient with a condition	**HSA-APR.C.5**
FORWARD 2	Use binomial coefficients to compute probabilities	**HSS-CP.B.9**

SET 21 • HSA-APR.C.5

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Use the Binomial Theorem to expand $(x + 2)^5$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q1/5

Use the Binomial Theorem to expand $(2a - b)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q2/5

Use the Binomial Theorem to find the coefficient of x^3 in $(x + 3)^7$.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q3/5

Use the Binomial Theorem to find the term containing x^3y^3 in $(x - 2y)^6$.

Term: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q4/5

Use the first three terms of the binomial expansion of $(1 + 0.02)^5$ to estimate 1.02^5 to four decimal places.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q5/5

SET 21 • HSA-APR.C.5

BACKWARD 1

Multiply binomials repeatedly • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Expand $(x + 2)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q1/5

Expand $(x + 2)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q2/5

True or false?

$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q3/5

The number of terms in the expansion of $(x + y)^5$ is

A. 6

B. 5

C. 10

D. 32

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q4/5

Expand $(a - b)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q5/5

SET 21 • HSA-APR.C.5

BACKWARD 2

Build Pascal's Triangle by a rule • Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5

Row 4 of Pascal's Triangle is 1, 4, 6, 4, 1. Write row 5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q1

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q1/5

Write row 7 of Pascal's Triangle (row 6 is 1, 6, 15, 20, 15, 6, 1).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q2

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q2/5

True or false?

Each inner entry of Pascal's Triangle is the sum of the two entries above it.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q3

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q3/5

The fourth entry of row 7 (counting from the first 1) is

A. 20

B. 7

C. 35

D. 21

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q4

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q4/5

What is the sum of the entries in row 5?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q5

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q5/5

BACKWARD 3

Powers of products and of negatives • Grade 8 • Expressions & Equations • 8.EE.A.1

Simplify $(2a)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q1/5

Simplify $(-2y)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q2/5

True or false?

$(-b)^4 = b^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q3/5

3^4 equals

A. 27

B. 12

C. 64

D. 81

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q4/5

Simplify $(0.02)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q5/5

SET 21 • HSA-APR.C.5

FORWARD 1

Find a specific term or coefficient with a condition • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Find the constant term of $(x + 1/x)^6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q1/5

The coefficient of x^2 in $(1 + kx)^5$ is 90. Find k ($k > 0$).

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q2/5

True or false?

The middle term of $(x + y)^8$ is $70x^4y^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q3/5

The term with x^4 in $(x^2 - 1)^5$ is

A. $-5x^4$

B. $-10x^4$

C. $10x^4$

D. $5x^4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S21-F1-Q4**

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q4/5

Find the coefficient of x in $(2x - 1)^6$.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q5/5

SET 21 • HSA-APR.C.5

FORWARD 2

Use binomial coefficients to compute probabilities • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability • HSS-CP.B.9

A fair coin is tossed 5 times. Use C(5, k) to find P(exactly 3 heads).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q1/5

A spinner lands on red with probability 0.3. Find P(exactly 2 reds in 4 spins).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q2/5

True or false?

The probabilities of 0, 1, ..., n successes add to 1 because $(p + (1 - p))^n = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q3

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q3/5

For 6 fair coin tosses, P(exactly 3 heads) is

A. 6/64

B. 3/6

C. 15/64

D. 20/64

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S21-F2-Q4**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability
HSS-CP.B.9 • Set 21 • FORWARD 2 • Q4/5

Explain why C(n, k) counts the outcomes with exactly k successes in n trials.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q5/5

Rewrite rational expressions by polynomial division

MAIN	Algebra II main question	**HSA-APR.D.6**
BACKWARD 1	Write a division with a remainder as a mixed number	**4.NF.B.3**
BACKWARD 2	Multiply to check a division	**HSA-APR.A.1**
BACKWARD 3	Use the Remainder Theorem to check the remainder	**HSA-APR.B.2**
FORWARD 1	Divide with unknown coefficients and higher-degree divisors	**HSA-APR.D.6**
FORWARD 2	Use division to find slant asymptotes	**HSF-IF.C.7.d**

SET 22 • HSA-APR.D.6

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6

Rewrite $(x^3 + 2x^2 - 5x + 3)/(x - 1)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q1/5

Rewrite $(2x^3 - 3x^2 + 4x - 7)/(x^2 + 1)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q2/5

Rewrite $(x^4 - 16)/(x - 2)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q3/5

Rewrite $(6x^2 + x - 4)/(2x + 1)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q4/5

Rewrite $(x^3 - 4x + 5)/(x + 3)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q5/5

SET 22 • HSA-APR.D.6

BACKWARD 1

Write a division with a remainder as a mixed number • Grade 4 • Number & Operations–Fractions • 4.NF.B.3

Write 23/4 as a mixed number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q1

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q1/5

Write 47/5 in the form q + r/5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q2

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q2/5

True or false?

17/3 = 5 + 2/3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q3

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q3/5

31/7 equals

A. 4 + 1/7

B. 4 + 3/7

C. 3 + 4/7

D. 4 + 3/31

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q4

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q4/5

Write 100/9 in the form q + r/9.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q5

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q5/5

SET 22 • HSA-APR.D.6

BACKWARD 2

Multiply to check a division • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Compute $(x - 1)(x^2 + 3x - 2) + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B2-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 22 • BACKWARD 2 • Q1/5

Compute $(x^2 + 1)(2x - 3) + (2x - 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B2-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 22 • BACKWARD 2 • Q2/5

True or false?

$(2x + 1)(3x - 1) - 3 = 6x^2 + x - 4.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B2-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 22 • BACKWARD 2 • Q3/5

$(x + 3)(x^2 - 3x + 5)$ equals

A. $x^3 - 4x + 15$

B. $x^3 + 15$

C. $x^3 - 4x + 5$

D. $x^3 + 6x^2 - 4x + 15$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B2-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 22 • BACKWARD 2 • Q4/5

Compute $(x - 2)(x^3 + 2x^2 + 4x + 8)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B2-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 22 • BACKWARD 2 • Q5/5

SET 22 • HSA-APR.D.6

BACKWARD 3

Use the Remainder Theorem to check the remainder • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Find the remainder of $x^3 + 2x^2 - 5x + 3$ divided by $x - 1$ by evaluating.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B3-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 22 • BACKWARD 3 • Q1/5

Find the remainder of $x^3 - 4x + 5$ divided by $x + 3$ by evaluating.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B3-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 22 • BACKWARD 3 • Q2/5

True or false?

The remainder of $x^4 - 16$ divided by $x - 2$ is 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B3-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 22 • BACKWARD 3 • Q3/5

The remainder of $6x^2 + x - 4$ divided by $2x + 1$ is

A. 3

B. -4

C. 1

D. -3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B3-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 22 • BACKWARD 3 • Q4/5

Find the remainder of $2x^3 + x - 1$ divided by $x - 2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B3-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 22 • BACKWARD 3 • Q5/5

SET 22 • HSA-APR.D.6

FORWARD 1

Divide with unknown coefficients and higher-degree divisors • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6

When $x^3 + ax^2 + 2x + 4$ is divided by $x^2 - x + 1$, the remainder is a constant. Find a and the remainder.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • FORWARD 1 • Q1/5

Rewrite $(x^4 + x^2 + 1)/(x^2 + x + 1)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • FORWARD 1 • Q2/5

True or false?

Dividing a degree-5 polynomial by a degree-2 polynomial gives a quotient of degree 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • FORWARD 1 • Q3/5

The remainder of a division by $x^2 + 4$ can have degree at most

A. 0

B. 1

C. 2

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S22-F1-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • FORWARD 1 • Q4/5

Find the polynomial $a(x)$ for which $a(x)/(x - 2) = x^2 + 1 + 3/(x - 2)$.

$a(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • FORWARD 1 • Q5/5

SET 22 • HSA-APR.D.6

FORWARD 2

Use division to find slant asymptotes • High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d

Use division to find the slant asymptote of $f(x) = (x^2 + 3x + 1)/(x + 1)$.

Asymptote: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 22 • FORWARD 2 • Q1/5

Find the slant asymptote of $f(x) = (2x^2 - x - 4)/(x - 1)$.

Asymptote: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 22 • FORWARD 2 • Q2/5

True or false?

A rational function with numerator degree exactly one more than the denominator degree has a slant asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 22 • FORWARD 2 • Q3/5

The slant asymptote of $y = (x^2 + 1)/x$ is

A. $y = x$

B. $y = 1$

C. $x = 0$

D. $y = x + 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 22 • FORWARD 2 • Q4/5

Explain why the graph of $y = x + 2 - 1/(x + 1)$ gets close to $y = x + 2$ for large $|x|$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 22 • FORWARD 2 • Q5/5

Rewrite simple rational expressions by inspection

MAIN	Algebra II main question	**HSA-APR.D.6**
BACKWARD 1	Split a fraction with a sum in the numerator	**7.EE.A.1**
BACKWARD 2	Rewrite a numerator as a multiple of the denominator plus a cons…	**6.EE.A.3**
BACKWARD 3	Divide by long division when inspection fails	**HSA-APR.D.6**
FORWARD 1	Use the rewritten form to reason about values	**HSA-APR.D.6**
FORWARD 2	Graph rational functions as transformations of y = 1/x	**HSF-BF.B.3**

SET 23 • HSA-APR.D.6

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6

Rewrite $(x + 5)/(x + 2)$ by inspection in the form $q + r/(x + 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • MAIN • Q1/5

Rewrite $(2x - 1)/(x - 3)$ by inspection in the form $q + r/(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • MAIN • Q2/5

Rewrite $(x^2 + 1)/x$ by inspection in the form $q(x) + r/x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • MAIN • Q3/5

Rewrite $(3x + 7)/(x + 1)$ by inspection in the form $q + r/(x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • MAIN • Q4/5

Rewrite $x^2/(x - 1)$ by inspection in the form $q(x) + r/(x - 1)$. (Hint: $x^2 = (x^2 - 1) + 1$.)

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • MAIN • Q5/5

SET 23 • HSA-APR.D.6

BACKWARD 1

Split a fraction with a sum in the numerator • Grade 7 • Expressions & Equations • 7.EE.A.1

Write (x + 3)/x as a sum of two fractions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B1-Q1

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 23 • BACKWARD 1 • Q1/5

Simplify $(2(x - 3) + 5)/(x - 3)$ as a sum.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B1-Q2

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 23 • BACKWARD 1 • Q2/5

True or false?

$(a + b)/c = a/c + b/c$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B1-Q3

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 23 • BACKWARD 1 • Q3/5

$(x^2 + 1)/x$ equals

A. $x + 1/x$

B. $1 + 1/x$

C. $x + 1$

D. $x^2 + 1/x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B1-Q4

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 23 • BACKWARD 1 • Q4/5

True or false?

$c/(a + b) = c/a + c/b$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B1-Q5

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 23 • BACKWARD 1 • Q5/5

SET 23 • HSA-APR.D.6

BACKWARD 2

Rewrite a numerator as a multiple of the denominator plus a constant • Grade 6 • Expressions & Equations • 6.EE.A.3

Find c so that $x + 5 = (x + 2) + c$.

$c =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 23 • BACKWARD 2 • Q1/5

Find c so that $2x - 1 = 2(x - 3) + c$.

c = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 23 • BACKWARD 2 • Q2/5

True or false?

$3x + 7 = 3(x + 1) + 4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 23 • BACKWARD 2 • Q3/5

$x^2 - 1$ equals

A. $x(x - 1)$

B. $(x + 1)^2$

C. $(x - 1)^2$

D. $(x - 1)(x + 1)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 23 • BACKWARD 2 • Q4/5

Find c so that $4x + 1 = 4(x - 2) + c$.

$c =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 23 • BACKWARD 2 • Q5/5

SET 23 • HSA-APR.D.6

BACKWARD 3

Divide by long division when inspection fails • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6

Divide x^2 by $x - 1$ with long division. Give the quotient and remainder.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B3-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • BACKWARD 3 • Q1/5

Divide $3x + 7$ by $x + 1$. Give the quotient and remainder.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B3-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • BACKWARD 3 • Q2/5

True or false?

Long division and inspection give the same result.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B3-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • BACKWARD 3 • Q3/5

The remainder of $(2x - 1) \div (x - 3)$ is

A. -5

B. 2

C. -1

D. 5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B3-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • BACKWARD 3 • Q4/5

Divide $x + 5$ by $x + 2$. Give the quotient and remainder.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-B3-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • BACKWARD 3 • Q5/5

SET 23 • HSA-APR.D.6

FORWARD 1

Use the rewritten form to reason about values • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6

For which integers x is $(x + 5)/(x + 2)$ an integer? Use $1 + 3/(x + 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • FORWARD 1 • Q1/5

Explain why $(2x - 1)/(x - 3)$ can never equal 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • FORWARD 1 • Q2/5

True or false?

$(3x + 7)/(x + 1) > 3$ for every $x > -1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • FORWARD 1 • Q3/5

As x grows large, (x + 5)/(x + 2) approaches

A. 3

B. 0

C. 5/2

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • FORWARD 1 • Q4/5

For $x > 0$, what is the least value of $x + 1/x$, and where does it occur?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 23 • FORWARD 1 • Q5/5

SET 23 • HSA-APR.D.6

FORWARD 2

Graph rational functions as transformations of y = 1/x • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Describe how the graph of $y = 1 + 3/(x + 2)$ is obtained from $y = 3/x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 23 • FORWARD 2 • Q1/5

Give the asymptotes of $y = 2 + 5/(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 23 • FORWARD 2 • Q2/5

True or false?

$y = (3x + 7)/(x + 1)$ has horizontal asymptote $y = 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 23 • FORWARD 2 • Q3/5

$y = (x + 5)/(x + 2)$ is a translation of $y = 3/x$ by

A. left 2, up 1

B. right 2, up 1

C. left 2, up 5

D. left 5, up 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S23-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 23 • FORWARD 2 • Q4/5

Write $(4x - 3)/(x - 1)$ as a transformation of k/x and give its asymptotes.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S23-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 23 • FORWARD 2 • Q5/5

Multiply and divide rational expressions

MAIN	Algebra II main question	**HSA-APR.D.7**
BACKWARD 1	Multiply and divide fractions	**6.NS.A.1**
BACKWARD 2	Factor numerators and denominators	**HSA-SSE.A.2**
BACKWARD 3	Division by zero is undefined	**7.NS.A.2.b**
FORWARD 1	Simplify complex fractions	**HSA-APR.D.7**
FORWARD 2	Explain closure of rational expressions	**HSA-APR.D.7**

SET 24 • HSA-APR.D.7

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Simplify $(x^2 - 9)/(x^2 + x - 6) \cdot (x - 2)/(x - 3)$. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • MAIN • Q1/5

Simplify $(x + 2)/(3x^2) \div (x^2 - 4)/x$. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • MAIN • Q2/5

Simplify $(2x - 6)/(x + 1) \cdot (x + 1)/(x^2 - 9)$. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • MAIN • Q3/5

Simplify $(3x + 3)/(x - 4) \div (x + 1)/(x - 4)$. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • MAIN • Q4/5

Simplify $(5x^2 + 5x)/(x^2 - 4) \div (x + 1)/(x - 2)$. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • MAIN • Q5/5

SET 24 • HSA-APR.D.7

BACKWARD 1

Multiply and divide fractions • Grade 6 • The Number System • 6.NS.A.1

Compute 2/3 · 9/4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B1-Q1

Grade 6 • The Number System • 6.NS.A.1 • Set 24 • BACKWARD 1 • Q1/5

Compute 5/6 ÷ 10/9.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B1-Q2

Grade 6 • The Number System • 6.NS.A.1 • Set 24 • BACKWARD 1 • Q2/5

True or false?

$a/b \div c/d = a/b \cdot d/c$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B1-Q3

Grade 6 • The Number System • 6.NS.A.1 • Set 24 • BACKWARD 1 • Q3/5

3/8 ÷ 3/4 equals

A. 9/32

B. 2

C. 3/4

D. 1/2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B1-Q4

Grade 6 • The Number System • 6.NS.A.1 • Set 24 • BACKWARD 1 • Q4/5

Compute 14/15 · 5/7.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B1-Q5

Grade 6 • The Number System • 6.NS.A.1 • Set 24 • BACKWARD 1 • Q5/5

SET 24 • HSA-APR.D.7

BACKWARD 2

Factor numerators and denominators • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 + x - 6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 24 • BACKWARD 2 • Q1/5

Factor $5x^2 + 5x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 24 • BACKWARD 2 • Q2/5

$x^2 - 4$ equals

A. $(x - 4)(x + 1)$

B. $(x - 2)^2$

C. $(x - 2)(x + 2)$

D. $x(x - 4)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 24 • BACKWARD 2 • Q3/5

True or false?

$2x - 6 = 2(x - 3)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 24 • BACKWARD 2 • Q4/5

Factor $x^2 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 24 • BACKWARD 2 • Q5/5

SET 24 • HSA-APR.D.7

BACKWARD 3

Division by zero is undefined • Grade 7 • The Number System • 7.NS.A.2.b

For which x is 1/(x − 4) undefined?

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B3-Q1

Grade 7 • The Number System • 7.NS.A.2.b • Set 24 • BACKWARD 3 • Q1/5

For which x is $(x + 2)/(x^2 - 9)$ undefined?

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B3-Q2

Grade 7 • The Number System • 7.NS.A.2.b • Set 24 • BACKWARD 3 • Q2/5

True or false?

After canceling $(x - 3)$, $x = 3$ is still excluded from the original expression.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B3-Q3

Grade 7 • The Number System • 7.NS.A.2.b • Set 24 • BACKWARD 3 • Q3/5

Dividing by $(x + 1)/(x - 2)$ is impossible when

A. $x = -1$ only

B. $x = 0$

C. $x = -1$ or $x = 2$

D. $x = 2$ only

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B3-Q4

Grade 7 • The Number System • 7.NS.A.2.b • Set 24 • BACKWARD 3 • Q4/5

Why is $x = 0$ excluded from $(x + 2)/(3x^2)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-B3-Q5

Grade 7 • The Number System • 7.NS.A.2.b • Set 24 • BACKWARD 3 • Q5/5

SET 24 • HSA-APR.D.7

FORWARD 1

Simplify complex fractions • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Simplify (1/x + 1)/(1/x − 1). State the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 1 • Q1/5

Simplify $(1 - 4/x^2)/(1 + 2/x)$. State the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 1 • Q2/5

True or false?

$(a/b)/(c/d) = ad/(bc)$ when $b, c, d \neq 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 1 • Q3/5

$(x/2)/(x/4)$ for $x \neq 0$ equals

A. 2

B. 1/2

C. $x^2/8$

D. $8/x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F1-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 1 • Q4/5

Simplify $(x - 1/x)/(x + 1)$. State the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F1-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 1 • Q5/5

SET 24 • HSA-APR.D.7

FORWARD 2

Explain closure of rational expressions • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Explain why the product of two rational expressions is a rational expression.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 2 • Q1/5

Explain why division by a nonzero rational expression gives a rational expression.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 2 • Q2/5

True or false?

Rational expressions are closed under division by the zero expression.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 2 • Q3/5

Which system is closed under addition, subtraction, multiplication, and division by nonzero elements?

A. Whole numbers

B. Rational expressions

C. Integers

D. Polynomials

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F2-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 2 • Q4/5

Give an example showing polynomials are not closed under division.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S24-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 24 • FORWARD 2 • Q5/5

Add and subtract rational expressions

MAIN	Algebra II main question	**HSA-APR.D.7**
BACKWARD 1	Add fractions with unlike denominators	**5.NF.A.1**
BACKWARD 2	Find a least common denominator by factoring	**HSA-SSE.A.2**
BACKWARD 3	Subtract a whole numerator	**6.EE.A.3**
FORWARD 1	Decompose into partial fractions	**HSA-APR.D.7**
FORWARD 2	Solve rational equations	**HSA-REI.A.2**

SET 25 • HSA-APR.D.7

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Simplify $(x + 1)/(x^2 + x) + 1/x$. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • MAIN • Q1/5

Write $\frac{2}{(x + 1)} + \frac{3}{(x - 2)}$ as one rational expression. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • MAIN • Q2/5

Simplify $x/(x-3) - 3/(x-3)$. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • MAIN • Q3/5

Write $1/(x^2 - 1) - 1/(x - 1)$ as one rational expression. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • MAIN • Q4/5

Write $4/(x^2 - 4x) - 1/x$ as one rational expression. State all excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • MAIN • Q5/5

SET 25 • HSA-APR.D.7

BACKWARD 1

Add fractions with unlike denominators • Grade 5 • Number & Operations–Fractions • 5.NF.A.1

Compute 2/3 + 3/4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B1-Q1

Grade 5 • Number & Operations–Fractions • 5.NF.A.1 • Set 25 • BACKWARD 1 • Q1/5

Compute 5/6 − 1/4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B1-Q2

Grade 5 • Number & Operations–Fractions • 5.NF.A.1 • Set 25 • BACKWARD 1 • Q2/5

True or false?

1/2 + 1/3 = 2/5.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B1-Q3

Grade 5 • Number & Operations–Fractions • 5.NF.A.1 • Set 25 • BACKWARD 1 • Q3/5

1/4 + 1/6 equals

A. 5/12

B. 2/10

C. 2/12

D. 1/5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B1-Q4

Grade 5 • Number & Operations–Fractions • 5.NF.A.1 • Set 25 • BACKWARD 1 • Q4/5

Compute 3/10 + 2/15.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B1-Q5

Grade 5 • Number & Operations–Fractions • 5.NF.A.1 • Set 25 • BACKWARD 1 • Q5/5

SET 25 • HSA-APR.D.7

BACKWARD 2

Find a least common denominator by factoring • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 + x$ and $x^2 - 4x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 25 • BACKWARD 2 • Q1/5

What is the least common denominator of $1/(x^2 - 1)$ and $1/(x - 1)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 25 • BACKWARD 2 • Q2/5

The LCD of $1/x$ and $1/(x(x-4))$ is

A. $x - 4$

B. $x^2(x - 4)$

C. $x(x - 4)$

D. x

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 25 • BACKWARD 2 • Q3/5

True or false?

The LCD of $2/(x + 1)$ and $3/(x - 2)$ is $(x + 1)(x - 2)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 25 • BACKWARD 2 • Q4/5

What is the LCD of $1/(x^2 + x)$ and $1/x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 25 • BACKWARD 2 • Q5/5

SET 25 • HSA-APR.D.7

BACKWARD 3

Subtract a whole numerator • Grade 6 • Expressions & Equations • 6.EE.A.3

Simplify $4 - (x - 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 25 • BACKWARD 3 • Q1/5

Simplify $1 - (x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 25 • BACKWARD 3 • Q2/5

True or false?

$3 - (x - 3) = -x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 25 • BACKWARD 3 • Q3/5

$2(x - 2) - 3(x + 1)$ equals

A. $5x - 7$

B. $-x + 1$

C. $-x - 7$

D. $-x - 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 25 • BACKWARD 3 • Q4/5

Simplify $x - (3 - x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 25 • BACKWARD 3 • Q5/5

SET 25 • HSA-APR.D.7

FORWARD 1

Decompose into partial fractions • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Find A and B so that

$$\frac{5x + 1}{(x - 1)(x + 2)} = \frac{A}{x - 1} + \frac{B}{x + 2}.$$

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • FORWARD 1 • Q1/5

Find A and B so that $1/(x(x + 1)) = A/x + B/(x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • FORWARD 1 • Q2/5

True or false?

$1/(x(x + 1)) = 1/x - 1/(x + 1)$ for $x \neq 0, -1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • FORWARD 1 • Q3/5

$4/(x^2 - 4)$ equals

A. $2/(x - 2) - 2/(x + 2)$

B. $1/(x - 2) + 1/(x + 2)$

C. $4/(x - 2) - 4/(x + 2)$

D. $1/(x - 2) - 1/(x + 2)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S25-F1-Q4**

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • FORWARD 1 • Q4/5

Use $1/(n(n+1)) = 1/n - 1/(n+1)$ to find $1/(1\cdot2) + 1/(2\cdot3) + 1/(3\cdot4) + 1/(4\cdot5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F1-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 25 • FORWARD 1 • Q5/5

SET 25 • HSA-APR.D.7

FORWARD 2

Solve rational equations • High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2

Solve $2/(x + 1) + 3/(x - 2) = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F2-Q1

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 25 • FORWARD 2 • Q1/5

Solve $x/(x-3) - 3/(x-3) = 1$. Explain your answer.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F2-Q2

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 25 • FORWARD 2 • Q2/5

True or false?

A solution of a rational equation that makes a denominator 0 must be rejected.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F2-Q3

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 25 • FORWARD 2 • Q3/5

The solution of $1/x + 1/(2x) = 3$ is

A. $x = 1/2$

B. $x = 2$

C. $x = 3/2$

D. no solution

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F2-Q4

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 25 • FORWARD 2 • Q4/5

Solve $4/(x^2 - 4x) - 1/x = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S25-F2-Q5

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 25 • FORWARD 2 • Q5/5

Create and solve rational, radical, and exponential equations from contexts

MAIN	Algebra II main question	**HSA-CED.A.1**
BACKWARD 1	Express rates of work and unit rates	**6.RP.A.3.b**
BACKWARD 2	Write exponential growth models	**HSF-LE.A.2**
BACKWARD 3	Solve linear equations after clearing a denominator	**HSA-REI.B.3**
FORWARD 1	Create equations whose algebraic solutions must be checked aga.	**HSA-CED.A.1**
FORWARD 2	Solve exponential context equations with logarithms	**HSF-LE.A.4**

SET 26 • HSA-CED.A.1

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1

Ana can paint a room in 4 hours. Working together, Ana and Ben paint it in 2.4 hours. Write an equation for Ben's time b (in hours) working alone, and solve it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • MAIN • Q1/5

A car's speed s (in mph) before braking is estimated from its skid length d (in feet) by $s = \sqrt{(22.5d)}$. Write and solve an equation for the skid length of a car going 45 mph.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • MAIN • Q2/5

A town of 1,200 people doubles in population every 5 years. Write an equation for the time t (in years) when the population is 9,600, and solve it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • MAIN • Q3/5

A printer costs $500 plus $4 per shirt. Write an equation for the number of shirts x that makes the average cost per shirt $9, and solve it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • MAIN • Q4/5

A tank holds 10 L of a 20% salt solution. Write an equation for the liters x of pure water to add to make an 8% solution, and solve it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • MAIN • Q5/5

SET 26 • HSA-CED.A.1

BACKWARD 1

Express rates of work and unit rates • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.b

Ana paints a room in 4 hours. What fraction of the room does she paint per hour?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B1-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.b • Set 26 • BACKWARD 1 • Q1/5

A pump fills a pool in 6 hours. What fraction does it fill in 2 hours?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B1-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.b • Set 26 • BACKWARD 1 • Q2/5

True or false?

If two people work together, their hourly rates add.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B1-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.b • Set 26 • BACKWARD 1 • Q3/5

A machine makes 300 parts in 5 hours. Its rate is

A. 1500 parts per hour

B. 300 parts per hour

C. 60 parts per hour

D. 5/300 parts per hour

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B1-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.b • Set 26 • BACKWARD 1 • Q4/5

A job takes 2.4 hours together. What fraction is done per hour?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B1-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.b • Set 26 • BACKWARD 1 • Q5/5

SET 26 • HSA-CED.A.1

BACKWARD 2

Write exponential growth models • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

A population of 1,200 doubles every 5 years. Write the population after t years.

P(t) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B2-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 26 • BACKWARD 2 • Q1/5

A $500 deposit grows 3% per year. Write its value after t years.

V(t) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B2-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 26 • BACKWARD 2 • Q2/5

True or false?

A quantity that doubles every 5 years is multiplied by 2^4 after 20 years.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B2-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 26 • BACKWARD 2 • Q3/5

A quantity triples every year starting at 7. After t years it is

A. $7 + 3t$

B. $21t$

C. $3 \cdot 7^t$

D. $7 \cdot 3^t$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B2-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 26 • BACKWARD 2 • Q4/5

How many doublings happen in 15 years if the doubling time is 5 years?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B2-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 26 • BACKWARD 2 • Q5/5

SET 26 • HSA-CED.A.1

BACKWARD 3

Solve linear equations after clearing a denominator • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3

Solve 500 + 4x = 9x.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 26 • BACKWARD 3 • Q1/5

Solve $2 = 0.08(10 + x)$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 26 • BACKWARD 3 • Q2/5

True or false?

To solve $(500 + 4x)/x = 9$ we may multiply both sides by x when $x \neq 0$.

A. True

B. False

2025 = 22.5d gives d =

A. 2002.5

B. 45

C. 9

D. 90

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 26 • BACKWARD 3 • Q4/5

Solve $5x/12 = 1/2.4$ for x.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 26 • BACKWARD 3 • Q5/5

SET 26 • HSA-CED.A.1

FORWARD 1

Create equations whose algebraic solutions must be checked against the context • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1

A boat travels 24 km downstream and 24 km back in 5 hours. The current is 2 km/h. Write an equation for the boat's still-water speed b and solve it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • FORWARD 1 • Q1/5

The length of a rectangle is 3 more than its width w, and its diagonal is 15. Write an equation and find w.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • FORWARD 1 • Q2/5

True or false?

A negative solution of an equation for a time must be rejected.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • FORWARD 1 • Q3/5

A model gives $t = -3$ or $t = 4$ for the time (in seconds) of an event. The answer is

A. neither

B. both

C. 4 seconds

D. -3 seconds

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S26-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • FORWARD 1 • Q4/5

A pipe fills a tank in x hours and a second pipe takes 2 hours longer. Together they take 2.4 hours. Write an equation and find x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 26 • FORWARD 1 • Q5/5

SET 26 • HSA-CED.A.1

FORWARD 2

Solve exponential context equations with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

The town of 1,200 doubles every 5 years. Write and solve an equation for when the population reaches 5,000. Give t to the nearest tenth of a year. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 26 • FORWARD 2 • Q1/5

Solve $500(1.04)^t = 800$ for t. Give an exact answer and a decimal to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 26 • FORWARD 2 • Q2/5

True or false?

$2^{t/5} = 8$ can be solved without logarithms, but

$2^{t/5} = 5$ cannot be solved exactly without them.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 26 • FORWARD 2 • Q3/5

The exact solution of $3^x = 20$ is

A. 20/3

B. log(20/3)

C. log 20/log 3

D. 3 log 20

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F2-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 26 • FORWARD 2 • Q4/5

Explain why log(1.6)/log(1.04) is the time for an amount growing 4% per year to grow by 60%.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S26-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 26 • FORWARD 2 • Q5/5

SET 27 • HSA-CED.A.1 • CREATING EQUATIONS AND REASONING WITH EQUATIONS

Create and solve nonlinear inequalities from contexts

MAIN	Algebra II main question	**HSA-CED.A.1**
BACKWARD 1	Solve quadratic equations by factoring	**HSA-REI.B.4.b**
BACKWARD 2	Sign of a product on intervals	**7.NS.A.2**
BACKWARD 3	Reverse the inequality when multiplying by a negative	**HSA-REI.B.3**
FORWARD 1	Solve rational inequalities with sign charts and excluded values	**HSA-CED.A.1**
FORWARD 2	Write systems of constraints and judge viability	**HSA-CED.A.3**

SET 27 • HSA-CED.A.1

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1

A ball's height is $h = -16t^2 + 64t$ feet after t seconds. Write and solve an inequality for the times when the ball is at least 48 feet high.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • MAIN • Q1/5

A company sells x items at a price of 100 − 2x dollars each. Write and solve an inequality for the values of x that give revenue of at least $1,200.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • MAIN • Q2/5

The average cost of making x posters is (300 + 2x)/x dollars. Write and solve an inequality for the numbers of posters (x > 0) with an average cost of at most $5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • MAIN • Q3/5

A drug concentration is $C(t) = 20t/(t^2 + 4)$ mg/L after t hours. Write and solve an inequality for the times when C is at least 4 mg/L.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • MAIN • Q4/5

$500 is invested at 4% interest compounded yearly. Write an inequality for the number of whole years t until the value is at least $800, and solve it. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • MAIN • Q5/5

SET 27 • HSA-CED.A.1

BACKWARD 1

Solve quadratic equations by factoring • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Solve $t^2 - 4t + 3 = 0$.

$t =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 27 • BACKWARD 1 • Q1/5

Solve $x^2 - 50x + 600 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 27 • BACKWARD 1 • Q2/5

True or false?

The solutions of $t^2 - 5t + 4 = 0$ are 1 and 4.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 27 • BACKWARD 1 • Q3/5

The solutions of $x^2 - x - 12 = 0$ are

A. 4 and −3

B. 6 and −2

C. 12 and −1

D. −4 and 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 27 • BACKWARD 1 • Q4/5

Solve $2x^2 - 100x + 1200 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 27 • BACKWARD 1 • Q5/5

SET 27 • HSA-CED.A.1

BACKWARD 2

Sign of a product on intervals • Grade 7 • The Number System • 7.NS.A.2

Is $(t - 1)(t - 3)$ positive or negative when $t = 2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B2-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 27 • BACKWARD 2 • Q1/5

Is $(t - 1)(t - 3)$ positive or negative when $t = 5$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B2-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 27 • BACKWARD 2 • Q2/5

True or false?

A product of two negative numbers is negative.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B2-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 27 • BACKWARD 2 • Q3/5

$(x - 20)(x - 30) \leq 0$ when

A. $x \leq 20$ or $x \geq 30$

B. $x \leq 20$

C. $20 \leq x \leq 30$

D. $x \geq 30$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B2-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 27 • BACKWARD 2 • Q4/5

For which t is $(t - 1)(t - 4) < 0$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B2-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 27 • BACKWARD 2 • Q5/5

SET 27 • HSA-CED.A.1

BACKWARD 3

Reverse the inequality when multiplying by a negative • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3

Solve $-16t^2 + 64t - 48 \geq 0$ after dividing both sides by -16.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 27 • BACKWARD 3 • Q1/5

Solve $-3x \geq 12$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 27 • BACKWARD 3 • Q2/5

True or false?

Multiplying both sides of an inequality by x is always allowed.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B3-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 27 • BACKWARD 3 • Q3/5

If $x > 0$, then $(300 + 2x)/x \leq 5$ is equivalent to

A. $300 + 2x \leq 5x$

B. $300 + 2x \geq 5x$

C. $2x \leq 5$

D. $300 \leq 2x$

Solve $2 - x < 7$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 27 • BACKWARD 3 • Q5/5

SET 27 • HSA-CED.A.1

FORWARD 1

Solve rational inequalities with sign charts and excluded values • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1

Solve $(x - 3)/(x + 1) \leq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • FORWARD 1 • Q1/5

Solve $(x + 2)/(x - 4) > 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • FORWARD 1 • Q2/5

True or false?

Multiplying $(x - 3)/(x + 1) \leq 0$ by $x + 1$ gives $x - 3 \leq 0$ for all x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • FORWARD 1 • Q3/5

The solution set of $1/x > 2$ is

A. $x > 1/2$

B. $x < 1/2$

C. $x > 0$

D. $0 < x < 1/2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • FORWARD 1 • Q4/5

A model's average cost is (400 + 3x)/x dollars. For how many items x > 0 is the average cost under $7?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.1 • Set 27 • FORWARD 1 • Q5/5

SET 27 • HSA-CED.A.1

FORWARD 2

Write systems of constraints and judge viability • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3

The ball's height must be at least 48 ft and the time at most 2.5 s. Write the constraints and find the viable times.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 27 • FORWARD 2 • Q1/5

Revenue $x(100 - 2x)$ must be at least \$1,200 and at most 24 items can be made. Which x are viable?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 27 • FORWARD 2 • Q2/5

True or false?

A value that solves the inequality may still not be viable in the context.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 27 • FORWARD 2 • Q3/5

Revenue $x(100 - 2x) \geq 1200$ with x a whole number is satisfied by

A. $x = 19$

B. $x = 25$

C. $x = 31$

D. $x = 50$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 27 • FORWARD 2 • Q4/5

Explain why 12 years, not 11.98 years, answers the compound-interest question.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S27-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 27 • FORWARD 2 • Q5/5

Create and graph nonlinear two-variable models

MAIN	Algebra II main question	**HSA-CED.A.2**
BACKWARD 1	Make a table of values from an equation	**6.EE.C.9**
BACKWARD 2	Choose scales and label axes	**HSN-Q.A.1**
BACKWARD 3	Write a growth or decay model from a percent	**HSF-LE.A.2**
FORWARD 1	Create equations in more than two variables and graph a member	**HSA-CED.A.2**
FORWARD 2	Interpret key features of the created models	**HSF-IF.B.4**

SET 28 • HSA-CED.A.2

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2

A rabbit population starts at 5 and doubles every year. Write an equation for the number N after t years, then graph it for $0 \le t \le 4$ on the grid.



N = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • MAIN • Q1/5

A rectangle has area 24 square units. Write an equation for its width w in terms of its length L, then graph it for $1 \le L \le 12$.

$w =$ _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • MAIN • Q2/5

A stone is dropped from 64 feet. Its height after t seconds is 64 minus 16 times the square of t. Write the equation and graph it for $0 \leq t \leq 2$.

$h =$ _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • MAIN • Q3/5

A $20,000 car loses 15% of its value each year. Write an equation for its value V, in thousands of dollars, after t years, then graph it for $0 \leq t \leq 5$.

$V =$ _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • MAIN • Q4/5

A car's speed s (mph) is the square root of 24 times its skid length d (ft). Write the equation and graph it for $0 \le d \le 150$.

$s =$ _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • MAIN • Q5/5

SET 28 • HSA-CED.A.2

BACKWARD 1

Make a table of values from an equation • Grade 6 • Expressions & Equations • 6.EE.C.9

For $N = 200 \cdot 2^t$, find N when t = 0, 1, 2, 3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B1-Q1

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 28 • BACKWARD 1 • Q1/5

For w = 24/L, find w when L = 2, 3, 4, 6.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B1-Q2

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 28 • BACKWARD 1 • Q2/5

True or false?

For $h = 64 - 16t^2$, $h = 48$ when $t = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B1-Q3

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 28 • BACKWARD 1 • Q3/5

For $s = \sqrt{(24d)}$, s when $d = 96$ is

A. 96

B. 48

C. 2304

D. 24

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B1-Q4

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 28 • BACKWARD 1 • Q4/5

For $V = 20000(0.85)^t$, find V when t = 1 and t = 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B1-Q5

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 28 • BACKWARD 1 • Q5/5

SET 28 • HSA-CED.A.2

BACKWARD 2

Choose scales and label axes • High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.1

Values of N go from 200 to 3,200. Give a sensible step for the vertical axis on a grid with about 7 lines.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B2-Q1

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.1 • Set 28 • BACKWARD 2 • Q1/5

Which quantity goes on the horizontal axis when V depends on t?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B2-Q2

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.1 • Set 28 • BACKWARD 2 • Q2/5

True or false?

Axis labels should include units.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B2-Q3

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.1 • Set 28 • BACKWARD 2 • Q3/5

The best vertical scale for values from 0 to 64 is

A. steps of 0.5

B. steps of 1 from 0 to 10

C. steps of 100 from 0 to 1000

D. steps of 10 from 0 to 70

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B2-Q4

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.1 • Set 28 • BACKWARD 2 • Q4/5

Give the horizontal range for $0 \leq d \leq 150$ with a step of 10. How many steps is that?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B2-Q5

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.1 • Set 28 • BACKWARD 2 • Q5/5

SET 28 • HSA-CED.A.2

BACKWARD 3

Write a growth or decay model from a percent • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

Write the value of $20,000 after losing 15% per year for t years.

$V =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B3-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 28 • BACKWARD 3 • Q1/5

What is the growth factor for 4% growth?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B3-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 28 • BACKWARD 3 • Q2/5

True or false?

Losing 15% each year means multiplying by 0.15 each year.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B3-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 28 • BACKWARD 3 • Q3/5

A quantity doubles every hour starting at 200. After t hours it is

A. $2 \cdot 200^{t}$

B. $200 \cdot 2^{t}$

C. $200 + 2t$

D. $400t$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B3-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 28 • BACKWARD 3 • Q4/5

What is the decay factor for 30% decay?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-B3-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 28 • BACKWARD 3 • Q5/5

SET 28 • HSA-CED.A.2

FORWARD 1

Create equations in more than two variables and graph a member of the family • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2

The current I (amps) through a resistor is the voltage V divided by the resistance R. Write the equation, and write I as a function of R when V = 12.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • FORWARD 1 • Q1/5

The volume of a cylinder is $V = \pi r^2 h$. Write h in terms of r when $V = 100\pi$, and describe its graph for $r > 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • FORWARD 1 • Q2/5

True or false?

Fixing one variable of a three-variable equation gives a two-variable equation that can be graphed.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • FORWARD 1 • Q3/5

For $A = P(1.05)^t$ with $P = 1000$, the graph of A against t is

A. a line

B. an increasing exponential curve

C. a parabola

D. a decreasing curve

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • FORWARD 1 • Q4/5

For the area formula $A = \pi r^2$, which graph (A against r) is correct for $r \geq 0$: a line, a half parabola, or an exponential?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.2 • Set 28 • FORWARD 1 • Q5/5

SET 28 • HSA-CED.A.2

FORWARD 2

Interpret key features of the created models • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

For $N = 200 \cdot 2^t$, what does the N-intercept mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 28 • FORWARD 2 • Q1/5

For $h = 64 - 16t^2$, when does the stone hit the ground, and what is the t-intercept's meaning?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 28 • FORWARD 2 • Q2/5

True or false?

The graph of $w = 24/L$ never reaches the L-axis because a rectangle of area 24 cannot have width 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 28 • FORWARD 2 • Q3/5

For $V = 20000(0.85)^t$, the graph

A. decreases and approaches 0

B. increases

C. is a line

D. decreases to 0 at t = 6.7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 28 • FORWARD 2 • Q4/5

Describe the rate of change of $s = \sqrt{(24d)}$ as d grows.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S28-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 28 • FORWARD 2 • Q5/5

Represent nonlinear constraints and judge viable options

MAIN	Algebra II main question	**HSA-CED.A.3**
BACKWARD 1	Represent constraints with a linear system	**HSA-CED.A.3**
BACKWARD 2	Test whether a point satisfies an inequality	**6.EE.B.5**
BACKWARD 3	Solve quadratic inequalities	**HSA-CED.A.1**
FORWARD 1	Optimize within nonlinear constraints	**HSA-CED.A.3**
FORWARD 2	Locate constraint boundaries as graph intersections	**HSA-REI.D.11**

SET 29 • HSA-CED.A.3

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3

A garden has length L and width W (feet). The fencing for L + W is 30 ft and the area must be at least 200 ft². Write the constraints, find the viable widths, and decide whether W = 8 is viable.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • MAIN • Q1/5

A box has a square base of side x and a volume of 32 in^3; both x and the height h must be at most 4 in. Write the constraints, find the viable x, and decide whether x = 3 is viable.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • MAIN • Q2/5

A plan invests $1,000 at 5% compounded yearly and needs at least $1,500 within 10 years. Write the constraint and decide whether the plan is viable. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • MAIN • Q3/5

Attendance at a show is $500 - 10p$ people when tickets cost p dollars. Revenue must be at least $6,000. Write the constraints, find the viable prices, and decide whether $35 is viable.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • MAIN • Q4/5

A rectangular pen with sides x and y (m) needs area at least 50 m² and perimeter at most 40 m. Write the system and decide whether (5, 10) and (4, 12) are viable.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • MAIN • Q5/5

SET 29 • HSA-CED.A.3

BACKWARD 1

Represent constraints with a linear system • High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.3

Write a constraint for "the total of L and W is 30".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B1-Q1

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • BACKWARD 1 • Q1/5

Write a constraint for "at most 40 m of fence for a rectangle with sides x and y".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B1-Q2

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • BACKWARD 1 • Q2/5

True or false?

"At least 200" is written ≥ 200.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B1-Q3

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • BACKWARD 1 • Q3/5

"The height h must be at most 4" is

A. $h \leq 4$

B. $h \geq 4$

C. $h < 4$

D. $h = 4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B1-Q4

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • BACKWARD 1 • Q4/5

Is (12, 20) a solution of $x + y \leq 30$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B1-Q5

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • BACKWARD 1 • Q5/5

SET 29 • HSA-CED.A.3

BACKWARD 2

Test whether a point satisfies an inequality • Grade 6 • Expressions & Equations • 6.EE.B.5

Does $W = 8$ satisfy $W(30 - W) \geq 200$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 29 • BACKWARD 2 • Q1/5

Does $p = 25$ satisfy $p(500 - 10p) \geq 6000$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 29 • BACKWARD 2 • Q2/5

True or false?

(4, 12) satisfies $xy \geq 50$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 29 • BACKWARD 2 • Q3/5

Which x satisfies $x^2 \geq 8$?

A. 3

B. 2

C. 1

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 29 • BACKWARD 2 • Q4/5

Does $t = 10$ satisfy $1000(1.05)^t \geq 1500$? A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 29 • BACKWARD 2 • Q5/5

SET 29 • HSA-CED.A.3

BACKWARD 3

Solve quadratic inequalities • High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.1

Solve $W^2 - 30W + 200 \leq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B3-Q1

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.1 • Set 29 • BACKWARD 3 • Q1/5

Solve $p^2 - 50p + 600 \leq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B3-Q2

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.1 • Set 29 • BACKWARD 3 • Q2/5

True or false?

$x^2 \geq 8$ with $x > 0$ means $x \geq 2\sqrt{2}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B3-Q3

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.1 • Set 29 • BACKWARD 3 • Q3/5

The solution of $(x - 2)(x - 6) \leq 0$ is

A. $x = 2$ or $x = 6$

B. $2 \leq x \leq 6$

C. $x \leq 2$ or $x \geq 6$

D. $x \geq 6$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B3-Q4

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.1 • Set 29 • BACKWARD 3 • Q4/5

Solve $32/x^2 \leq 4$ for $x > 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-B3-Q5

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.1 • Set 29 • BACKWARD 3 • Q5/5

SET 29 • HSA-CED.A.3

FORWARD 1

Optimize within nonlinear constraints • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3

With L + W = 30, which width gives the greatest garden area, and what is that area?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • FORWARD 1 • Q1/5

Which ticket price p gives the greatest revenue $p(500 - 10p)$, and what is that revenue?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • FORWARD 1 • Q2/5

True or false?

A maximum found from the model must still satisfy every other constraint to be a viable choice.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • FORWARD 1 • Q3/5

For the pen with $xy \geq 50$ and $2x + 2y \leq 40$, which is viable?

A. (10, 10)

B. (15, 10)

C. (2, 20)

D. (4, 12)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S29-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • FORWARD 1 • Q4/5

Among whole-number widths, list all viable garden widths when the area must be at least 216 ft².

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.3 • Set 29 • FORWARD 1 • Q5/5

SET 29 • HSA-CED.A.3

FORWARD 2

Locate constraint boundaries as graph intersections • High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

Explain how the viable widths $10 \leq W \leq 20$ can be read from the graphs of $y = W(30 - W)$ and $y = 200$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 29 • FORWARD 2 • Q1/5

Where do $y = p(500 - 10p)$ and $y = 6000$ intersect?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 29 • FORWARD 2 • Q2/5

True or false?

The x-coordinates where $y = 32/x^2$ meets $y = 4$ solve $32/x^2 = 4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 29 • FORWARD 2 • Q3/5

The graphs of $y = 1000(1.05)^t$ and $y = 1500$ meet at $t \approx$

A. 15

B. 8.3

C. 10

D. 5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S29-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 29 • FORWARD 2 • Q4/5

Explain why the region above y = 200 and below the parabola shows the viable widths.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S29-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 29 • FORWARD 2 • Q5/5

Rearrange rational, radical, and exponential formulas

MAIN	Algebra II main question	**HSA-CED.A.4**
BACKWARD 1	Rearrange linear formulas	**HSA-CED.A.4**
BACKWARD 2	Undo squares and square roots	**8.EE.A.2**
BACKWARD 3	Clear fractions in an equation	**HSA-REI.B.3**
FORWARD 1	Rearrange formulas where conditions decide the result	**HSA-CED.A.4**
FORWARD 2	Rearrange exponential formulas with logarithms	**HSF-LE.A.4**

SET 30 • HSA-CED.A.4

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4

The lens formula is $1/f = 1/a + 1/b$. Solve for b and state the condition needed.

$b =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • MAIN • Q1/5

A pendulum's period is $T = 2\pi\sqrt{(L/g)}$. Solve for L.

$L =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • MAIN • Q2/5

An investment grows by $A = P(1 + r)^t$. Solve for r.

r = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • MAIN • Q3/5

The speed of a falling object is $v = \sqrt{(2gh)}$. Solve for h and state the condition on v.

$h =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • MAIN • Q4/5

In a circuit, $I = V/(R + r)$. Solve for R and state the condition needed.

$R =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • MAIN • Q5/5

SET 30 • HSA-CED.A.4

BACKWARD 1

Rearrange linear formulas • High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4

Solve P = 2L + 2W for W.

W = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B1-Q1

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • BACKWARD 1 • Q1/5

Solve d = rt for t.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B1-Q2

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • BACKWARD 1 • Q2/5

True or false?

Solving V = IR for R gives R = V/I.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B1-Q3

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • BACKWARD 1 • Q3/5

Solving $y = mx + b$ for x gives

A. $(y - b)/m$

B. $y/m - b$

C. $(y + b)/m$

D. $m(y - b)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B1-Q4

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • BACKWARD 1 • Q4/5

Solve A = (1/2)bh for h.

h = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B1-Q5

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • BACKWARD 1 • Q5/5

SET 30 • HSA-CED.A.4

BACKWARD 2

Undo squares and square roots • Grade 8 • Expressions & Equations • 8.EE.A.2

Solve $x^2 = 49$ for $x > 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 30 • BACKWARD 2 • Q1/5

Solve $\sqrt{y} = 6$.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 30 • BACKWARD 2 • Q2/5

True or false?

If $\sqrt{(L/g)} = k$, then $L/g = k^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 30 • BACKWARD 2 • Q3/5

$(\sqrt{(2gh)})^2$ equals

A. $2g\sqrt{h}$

B. $2gh$

C. $\sqrt{(2gh)}$

D. $4g^2h^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 30 • BACKWARD 2 • Q4/5

Solve $T/(2\pi) = \sqrt{m}$ for m.

m = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 30 • BACKWARD 2 • Q5/5

SET 30 • HSA-CED.A.4

BACKWARD 3

Clear fractions in an equation • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3

Solve 1/2 + 1/x = 1 for x.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 30 • BACKWARD 3 • Q1/5

Combine $1/a - 1/f$ into one fraction.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 30 • BACKWARD 3 • Q2/5

True or false?

If $1/b = (a - f)/(af)$, then $b = af/(a - f)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B3-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 30 • BACKWARD 3 • Q3/5

Solving 3 = 12/(R + 2) for R gives

A. 36

B. 4

C. 6

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 30 • BACKWARD 3 • Q4/5

Solve $I = V/(R + r)$ for $R + r$.

$R + r =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 30 • BACKWARD 3 • Q5/5

SET 30 • HSA-CED.A.4

FORWARD 1

Rearrange formulas where conditions decide the result • High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4

Solve $A = \pi r^2$ for r, given r is a length.

r = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • FORWARD 1 • Q1/5

Solve $y = (x + 1)/(x - 2)$ for x and state the condition on y.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • FORWARD 1 • Q2/5

True or false?

Solving $s = \sqrt{(24d)}$ for d gives $d = s^2/24$ for every real s.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • FORWARD 1 • Q3/5

Solving $E = (1/2)mv^2$ for v ($v \geq 0$) gives

A. $\sqrt{(2E/m)}$

B. $(2E/m)^2$

C. $\sqrt{(E/2m)}$

D. $2E/m$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • FORWARD 1 • Q4/5

Solve $P = k/V^2$ for V ($V > 0$).

$V =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Creating Equations • HSA-CED.A.4 • Set 30 • FORWARD 1 • Q5/5

SET 30 • HSA-CED.A.4

FORWARD 2

Rearrange exponential formulas with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $A = Pe^{rt}$ for t.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 30 • FORWARD 2 • Q1/5

Solve $A = P(1 + r)^t$ for t.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 30 • FORWARD 2 • Q2/5

True or false?

Solving $N = N_0 \cdot 2^{t/d}$ for t gives $t = d \cdot \log_2(N/N_0)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 30 • FORWARD 2 • Q3/5

Solving $y = 5 \cdot 10^x$ for x gives

A. y/50

B. log(y/5)

C. $10^{y/5}$

D. log y − 5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F2-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 30 • FORWARD 2 • Q4/5

Use $t = \log(A/P)/\log(1 + r)$ to find t for $A = 2000$, $P = 1000$, $r = 0.05$. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S30-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 30 • FORWARD 2 • Q5/5

Solve rational equations and identify extraneous solutions

MAIN	Algebra II main question	**HSA-REI.A.2**
BACKWARD 1	Solve proportions	**7.RP.A.2**
BACKWARD 2	Find a common denominator of rational expressions	**HSA-APR.D.7**
BACKWARD 3	Values that make a denominator zero	**7.NS.A.2.b**
FORWARD 1	Explain and construct extraneous solutions	**HSA-REI.A.2**
FORWARD 2	See rational equations as graph intersections	**HSA-REI.D.11**

SET 31 • HSA-REI.A.2

MAIN

Algebra II main question • High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2

Solve $3/(x - 1) = 6/(x + 2)$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-M-Q1

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • MAIN • Q1/5

Solve $x/(x - 2) + 1 = 2/(x - 2)$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-M-Q2

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • MAIN • Q2/5

Solve $1/x + 1/(x + 2) = 4/3$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-M-Q3

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • MAIN • Q3/5

Solve $x^2/(x - 3) = 9/(x - 3)$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-M-Q4

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • MAIN • Q4/5

Solve $2/(x + 1) - 1/(x - 1) = 2/(x^2 - 1)$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-M-Q5

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • MAIN • Q5/5

SET 31 • HSA-REI.A.2

BACKWARD 1

Solve proportions • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2

Solve $3/x = 6/10$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B1-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2 • Set 31 • BACKWARD 1 • Q1/5

Solve 4/7 = x/21.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B1-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2 • Set 31 • BACKWARD 1 • Q2/5

True or false?

$a/b = c/d$ implies $ad = bc$ when $b, d \neq 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B1-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2 • Set 31 • BACKWARD 1 • Q3/5

2/5 = 6/x gives x =

A. 12

B. 2.4

C. 3

D. 15

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B1-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2 • Set 31 • BACKWARD 1 • Q4/5

Solve $3/(x - 1) = 1$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B1-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.2 • Set 31 • BACKWARD 1 • Q5/5

SET 31 • HSA-REI.A.2

BACKWARD 2

Find a common denominator of rational expressions • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

What is the LCD of 1/x and 1/(x + 2)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 31 • BACKWARD 2 • Q1/5

What is the LCD of $\frac{2}{x+1}$, $\frac{1}{x-1}$, and $\frac{2}{x^2-1}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 31 • BACKWARD 2 • Q2/5

True or false?

The LCD of $1/(x-2)$ and $2/(x-2)$ is $(x-2)^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 31 • BACKWARD 2 • Q3/5

Multiplying $1/x + 1/(x + 2) = 4/3$ by $3x(x + 2)$ gives

A. $3 + 3 = 4$

B. $3(x + 2) + 3x = 4$

C. $3(x + 2) + 3x = 4x(x + 2)$

D. $x + 2 + x = 4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B2-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 31 • BACKWARD 2 • Q4/5

Write 1/x + 1/(x + 2) as one fraction.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 31 • BACKWARD 2 • Q5/5

SET 31 • HSA-REI.A.2

BACKWARD 3

Values that make a denominator zero • Grade 7 • The Number System • 7.NS.A.2.b

Which x-values make $x/(x - 2)$ undefined?

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B3-Q1

Grade 7 • The Number System • 7.NS.A.2.b • Set 31 • BACKWARD 3 • Q1/5

Which x-values make $2/(x^2 - 1)$ undefined?

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B3-Q2

Grade 7 • The Number System • 7.NS.A.2.b • Set 31 • BACKWARD 3 • Q2/5

True or false?

A rational equation cannot have a solution that makes any denominator zero.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B3-Q3

Grade 7 • The Number System • 7.NS.A.2.b • Set 31 • BACKWARD 3 • Q3/5

Which value is excluded from $1/x + 1/(x + 2) = 4/3$?

A. 2

B. 1

C. 4/3

D. -2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B3-Q4

Grade 7 • The Number System • 7.NS.A.2.b • Set 31 • BACKWARD 3 • Q4/5

List the excluded values of $x^2/(x - 3) = 9/(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-B3-Q5

Grade 7 • The Number System • 7.NS.A.2.b • Set 31 • BACKWARD 3 • Q5/5

SET 31 • HSA-REI.A.2

FORWARD 1

Explain and construct extraneous solutions • High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2

Explain how multiplying both sides of $x/(x - 2) + 1 = 2/(x - 2)$ by $x - 2$ creates an extraneous solution.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F1-Q1

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • FORWARD 1 • Q1/5

Write a rational equation that has the extraneous solution x = 4 and no actual solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F1-Q2

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • FORWARD 1 • Q2/5

True or false?

Every rational equation has at least one extraneous solution.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F1-Q3

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • FORWARD 1 • Q3/5

For which a does $a/(x - 1) = 2/(x - 1)$ have infinitely many solutions?

A. a = 2

B. no value

C. a = 0

D. a = 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F1-Q4

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • FORWARD 1 • Q4/5

Solve $(x + 1)/(x - 1) = 2/(x - 1) + 1$ and explain the result.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F1-Q5

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 31 • FORWARD 1 • Q5/5

SET 31 • HSA-REI.A.2

FORWARD 2

See rational equations as graph intersections • High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

Explain why $x/(x - 2) + 1 = 2/(x - 2)$ has no solution using the graphs of $y = x/(x - 2) + 1$ and $y = 2/(x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 31 • FORWARD 2 • Q1/5

Where do $y = 3/(x - 1)$ and $y = 6/(x + 2)$ intersect?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 31 • FORWARD 2 • Q2/5

True or false?

An extraneous solution corresponds to an intersection point of the two graphs.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 31 • FORWARD 2 • Q3/5

$y = x^2/(x - 3)$ and $y = 9/(x - 3)$ intersect at

A. $(3, 3/2)$

B. $(-3, 3/2)$

C. $(-3, -3/2)$

D. $(3, 0)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S31-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 31 • FORWARD 2 • Q4/5

Use a graph or table to approximate the solution of $1/x = x - 1$ with $x > 0$ to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S31-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 31 • FORWARD 2 • Q5/5

Solve radical equations and identify extraneous solutions

MAIN	Algebra II main question	**HSA-REI.A.2**
BACKWARD 1	Square and cube both sides; principal roots	**8.EE.A.2**
BACKWARD 2	Solve the resulting quadratic	**HSA-REI.B.4.b**
BACKWARD 3	Check a candidate in the original equation	**6.EE.B.5**
FORWARD 1	Solve equations with two radicals, rational exponents, and $\sqrt{(x^2)}$	**HSA-REI.A.2**
FORWARD 2	Explain extraneous solutions with graphs of radical functions	**HSF-IF.C.7.b**

SET 32 • HSA-REI.A.2

MAIN

Algebra II main question • High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2

Solve $\sqrt{(x + 7)} = x - 5$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-M-Q1

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • MAIN • Q1/5

Solve $\sqrt{(2x - 1)} = 3$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-M-Q2

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • MAIN • Q2/5

Solve $\sqrt{(x + 3)} + 3 = x$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-M-Q3

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • MAIN • Q3/5

Solve $\sqrt[3]{(x-2)} = -3$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-M-Q4

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • MAIN • Q4/5

Solve $\sqrt{(x - 1)} + 5 = 2$. Identify any extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-M-Q5

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • MAIN • Q5/5

SET 32 • HSA-REI.A.2

BACKWARD 1

Square and cube both sides; principal roots • Grade 8 • Expressions & Equations • 8.EE.A.2

Solve $\sqrt{y} = 3$.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B1-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 32 • BACKWARD 1 • Q1/5

Solve $\sqrt[3]{y} = -3$.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B1-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 32 • BACKWARD 1 • Q2/5

True or false?

$\sqrt{9} = 3$ or -3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B1-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 32 • BACKWARD 1 • Q3/5

$\sqrt{y} = -2$ has

A. $y = 2$

B. no solution

C. $y = 4$

D. $y = -4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B1-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 32 • BACKWARD 1 • Q4/5

Square both sides of $\sqrt{(x + 7)} = x - 5$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B1-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 32 • BACKWARD 1 • Q5/5

SET 32 • HSA-REI.A.2

BACKWARD 2

Solve the resulting quadratic • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Solve $x^2 - 11x + 18 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B2-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 32 • BACKWARD 2 • Q1/5

Solve $x^2 - 7x + 6 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B2-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 32 • BACKWARD 2 • Q2/5

True or false?

$(x - 5)^2 = x^2 + 25$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B2-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 32 • BACKWARD 2 • Q3/5

$x^2 - 6x + 9 = x + 3$ gives

A. $x^2 - 7x + 12 = 0$

B. $x^2 - 5x + 12 = 0$

C. $x^2 - 6x + 6 = 0$

D. $x^2 - 7x + 6 = 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B2-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 32 • BACKWARD 2 • Q4/5

Solve $x^2 - 4x = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B2-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 32 • BACKWARD 2 • Q5/5

SET 32 • HSA-REI.A.2

BACKWARD 3

Check a candidate in the original equation • Grade 6 • Expressions & Equations • 6.EE.B.5

Check $x = 2$ in $\sqrt{(x + 7)} = x - 5$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 32 • BACKWARD 3 • Q1/5

Check $x = 6$ in $\sqrt{(x + 3)} + 3 = x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 32 • BACKWARD 3 • Q2/5

True or false?

$x = 5$ satisfies $\sqrt{(2x - 1)} = 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 32 • BACKWARD 3 • Q3/5

Which value satisfies $\sqrt{(x + 3)} + 3 = x$?

A. 1

B. 0

C. 6

D. -3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 32 • BACKWARD 3 • Q4/5

Check $x = 10$ in $\sqrt{(x - 1)} + 5 = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.B.5 • Set 32 • BACKWARD 3 • Q5/5

SET 32 • HSA-REI.A.2

FORWARD 1

Solve equations with two radicals, rational exponents, and $\sqrt{(x^2)}$ • High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2

Solve $\sqrt{(x + 5)} - \sqrt{x} = 1$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F1-Q1

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • FORWARD 1 • Q1/5

Solve $x^{3/2} = 8$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F1-Q2

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • FORWARD 1 • Q2/5

Solve $\sqrt{(x^2)} = 5$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F1-Q3

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • FORWARD 1 • Q3/5

True or false?

$\sqrt{(x^2)} = x$ for every real x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F1-Q4

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • FORWARD 1 • Q4/5

The solution set of $\sqrt{(x^2)} = x$ is

A. $x \geq 0$

B. $x \leq 0$

C. $x = 0$ only

D. all real x

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F1-Q5

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 32 • FORWARD 1 • Q5/5

SET 32 • HSA-REI.A.2

FORWARD 2

Explain extraneous solutions with graphs of radical functions • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

The graphs of $y = \sqrt{(x + 7)}$ and $y = x - 5$ meet at one point. Give it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 32 • FORWARD 2 • Q1/5

The extraneous value $x = 2$ is where $y = x - 5$ meets which other graph?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 32 • FORWARD 2 • Q2/5

True or false?

The graph of $y = \sqrt{(x + 7)}$ starts at $(-7, 0)$ and increases.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 32 • FORWARD 2 • Q3/5

The graph of $y = \sqrt{(x - 1)} + 5$ lies

A. below $y = 2$

B. below the x-axis

C. at or below $y = 5$

D. at or above $y = 5$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 32 • FORWARD 2 • Q4/5

Use the graphs to explain why $\sqrt{(x - 1)} + 5 = 2$ has no solution.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S32-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 32 • FORWARD 2 • Q5/5

Explain why intersections of nonlinear graphs solve f(x) = g(x)

MAIN	Algebra II main question	**HSA-REI.D.11**
BACKWARD 1	A graph is the set of its solutions	**HSA-REI.D.10**
BACKWARD 2	Solve a linear system by graphing	**8.EE.C.8.a**
BACKWARD 3	Evaluate nonlinear functions at an input	**HSF-IF.A.2**
FORWARD 1	Relate intersections to zeros of f - g	**HSA-REI.D.11**
FORWARD 2	Find all solutions of a trigonometric equation on an interval from a…	**HSF-TF.B.7**

SET 33 • HSA-REI.D.11

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

The graphs of $y = 2^x$ and $y = x + 2$ are shown. They meet at $x = 2$ and at a negative x. Explain why $x = 2$ solves $2^x = x + 2$, and verify it.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • MAIN • Q1/5

The graphs of $y = x^3 - 3x$ and $y = x$ are shown. Use them to list every solution of $x^3 - 3x = x$, and explain why they are solutions.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • MAIN • Q2/5

The graphs of $y = 1/x$ and $y = x$ are shown. List every solution of $1/x = x$ and explain using the graphs.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • MAIN • Q3/5

The graphs of $y = \sqrt{x}$ and $y = x/2$ are shown. List every solution of $\sqrt{x} = x/2$ and verify each.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • MAIN • Q4/5

The graphs of $y = |x - 1|$ and $y = 3$ are shown. List every solution of $|x - 1| = 3$ and explain.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • MAIN • Q5/5

SET 33 • HSA-REI.D.11

BACKWARD 1

A graph is the set of its solutions • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.10

Is (2, 4) on the graph of $y = 2^x$? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.10 • Set 33 • BACKWARD 1 • Q1/5

Is (3, 6) on the graph of $y = x + 2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.10 • Set 33 • BACKWARD 1 • Q2/5

True or false?

Every point on the graph of $y = f(x)$ has the form $(x, f(x))$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.10 • Set 33 • BACKWARD 1 • Q3/5

Which point is on $y = x^3 - 3x$?

A. (0, 3)

B. (2, 6)

C. (2, 2)

D. (1, 2)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.10 • Set 33 • BACKWARD 1 • Q4/5

Is $(-1, -1)$ on the graph of $y = 1/x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.10 • Set 33 • BACKWARD 1 • Q5/5

SET 33 • HSA-REI.D.11

BACKWARD 2

Solve a linear system by graphing • Grade 8 • Expressions & Equations • 8.EE.C.8.a

The lines $y = x + 1$ and $y = -x + 5$ meet at $(2, 3)$. Explain why $x = 2$ solves $x + 1 = -x + 5$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.C.8.a • Set 33 • BACKWARD 2 • Q1/5

Where do y = 2x and y = 6 meet?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.C.8.a • Set 33 • BACKWARD 2 • Q2/5

True or false?

Two lines that do not intersect give an equation with no solution.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.C.8.a • Set 33 • BACKWARD 2 • Q3/5

$y = 3$ and $y = x - 1$ meet at $x =$

A. 4

B. 3

C. -1

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.C.8.a • Set 33 • BACKWARD 2 • Q4/5

Solve $x + 1 = -x + 5$ algebraically.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.C.8.a • Set 33 • BACKWARD 2 • Q5/5

SET 33 • HSA-REI.D.11

BACKWARD 3

Evaluate nonlinear functions at an input • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Find 2^x and $x + 2$ when $x = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 33 • BACKWARD 3 • Q1/5

Find $x^3 - 3x$ when $x = -2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 33 • BACKWARD 3 • Q2/5

True or false?

$\sqrt{4} = 4/2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 33 • BACKWARD 3 • Q3/5

$|x - 1|$ when $x = -2$ is

A. 3

B. 1

C. -1

D. -3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 33 • BACKWARD 3 • Q4/5

Find $1/x$ when $x = -1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 33 • BACKWARD 3 • Q5/5

SET 33 • HSA-REI.D.11

FORWARD 1

Relate intersections to zeros of f − g • High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

Explain why the solutions of $x^3 - 3x = x$ are the zeros of $h(x) = x^3 - 4x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • FORWARD 1 • Q1/5

Find the zeros of $h(x) = x^3 - 4x$ algebraically.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • FORWARD 1 • Q2/5

True or false?

If $y = f(x)$ and $y = g(x)$ meet at $x = a$, then the graph of $y = f(x) - g(x)$ crosses or touches the x-axis at a.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • FORWARD 1 • Q3/5

The solutions of $\sqrt{x} = x/2$ are the zeros of

A. $\sqrt{x} + x/2$

B. $\sqrt{x} - x/2$

C. $\sqrt{x} \cdot x/2$

D. $x/2$

Explain why $2^x = x + 2$ has exactly two solutions using $h(x) = 2^x - x - 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 33 • FORWARD 1 • Q5/5

SET 33 • HSA-REI.D.11

FORWARD 2

Find all solutions of a trigonometric equation on an interval from a graph • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7

On $0 \le x < 2\pi$, where does $y = \sin x$ meet $y = 1/2$? Give exact values.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 33 • FORWARD 2 • Q1/5

On $0 \le x < 2\pi$, where does $y = \cos x$ meet $y = 0$?

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 33 • FORWARD 2 • Q2/5

True or false?

sin x = 1/2 has infinitely many real solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 33 • FORWARD 2 • Q3/5

How many solutions does $\sin x = 0.3$ have on $0 \leq x < 2\pi$?

A. 1

B. 0

C. 2

D. 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S33-F2-Q4**

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 33 • FORWARD 2 • Q4/5

Explain why $\sin x = 2$ has no solution using the graph.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S33-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 33 • FORWARD 2 • Q5/5

Approximate solutions of f(x) = g(x)

MAIN	Algebra II main question	**HSA-REI.D.11**
BACKWARD 1	Read solutions of linear-quadratic systems from graphs	**HSA-REI.D.11**
BACKWARD 2	Evaluate exponential and logarithmic expressions with a calculat…	**HSF-IF.A.2**
BACKWARD 3	Round to a place value	**5.NBT.A.4**
FORWARD 1	Guarantee the accuracy of an approximation	**HSA-REI.D.11**
FORWARD 2	Compare an approximation with an exact logarithmic solution	**HSF-LE.A.4**

SET 34 • HSA-REI.D.11

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

Use a table or successive approximation to solve $2^x = 5 - x$ to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • MAIN • Q1/5

Solve $x^3 = 7 - x$ to the nearest hundredth with a table or successive approximation. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • MAIN • Q2/5

Solve $\log x = 2 - x$ (log base 10) to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • MAIN • Q3/5

The table shows $f(x) = 3(0.8)^x$ and $g(x) = x$. Between which two table values is the solution of $f(x) = g(x)$? Estimate it to the nearest tenth.

x	0	0.5	1	1.5	2	2.5
f(x)	3	2.683	2.4	2.147	1.92	1.717
g(x)	0	0.5	1	1.5	2	2.5

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • MAIN • Q4/5

Use a graph or table to find the negative solution of $2^x = x + 2$ to the nearest tenth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • MAIN • Q5/5

SET 34 • HSA-REI.D.11

BACKWARD 1

Read solutions of linear-quadratic systems from graphs • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

Where do $y = x^2$ and $y = 4$ meet?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • BACKWARD 1 • Q1/5

Read the intersection of $y = x^2$ and $y = x + 2$ from a graph or table.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • BACKWARD 1 • Q2/5

True or false?

The solutions of $f(x) = g(x)$ can be approximated from a table where $f - g$ changes sign.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • BACKWARD 1 • Q3/5

$f(1) - g(1) = -2$ and $f(2) - g(2) = 1$. A solution lies

A. below 1

B. at 1.5 exactly

C. above 2

D. between 1 and 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • BACKWARD 1 • Q4/5

$f(x) = x^2$ and $g(x) = 3$. Between which consecutive integers is the positive solution?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • BACKWARD 1 • Q5/5

SET 34 • HSA-REI.D.11

BACKWARD 2

Evaluate exponential and logarithmic expressions with a calculator • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Find $2^{1.7}$ to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 34 • BACKWARD 2 • Q1/5

Find log 1.75 (base 10) to the nearest thousandth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 34 • BACKWARD 2 • Q2/5

True or false?

$1.7^3 = 4.913$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 34 • BACKWARD 2 • Q3/5

$3(0.8)^2$ equals

A. 0.192

B. 4.8

C. 1.92

D. 2.4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 34 • BACKWARD 2 • Q4/5

Find $2^{-1.7}$ to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 34 • BACKWARD 2 • Q5/5

SET 34 • HSA-REI.D.11

BACKWARD 3

Round to a place value • Grade 5 • Number & Operations in Base Ten • 5.NBT.A.4

Round 1.7156 to the nearest hundredth.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B3-Q1

Grade 5 • Number & Operations in Base Ten • 5.NBT.A.4 • Set 34 • BACKWARD 3 • Q1/5

Round 1.944 to the nearest tenth.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B3-Q2

Grade 5 • Number & Operations in Base Ten • 5.NBT.A.4 • Set 34 • BACKWARD 3 • Q2/5

True or false?

1.755 rounded to the nearest hundredth is 1.76.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B3-Q3

Grade 5 • Number & Operations in Base Ten • 5.NBT.A.4 • Set 34 • BACKWARD 3 • Q3/5

-1.690 to the nearest tenth is

A. -1.6

B. -1.69

C. -2

D. -1.7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B3-Q4

Grade 5 • Number & Operations in Base Ten • 5.NBT.A.4 • Set 34 • BACKWARD 3 • Q4/5

Round 1.7392 to the nearest hundredth.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-B3-Q5

Grade 5 • Number & Operations in Base Ten • 5.NBT.A.4 • Set 34 • BACKWARD 3 • Q5/5

SET 34 • HSA-REI.D.11

FORWARD 1

Guarantee the accuracy of an approximation • High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

$h(x) = 2^x + x - 5$. Find $h(1.71)$ and $h(1.72)$ and explain why a solution lies between them. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • FORWARD 1 • Q1/5

Use one step of bisection on $[1, 2]$ for $x^3 + x - 7 = 0$. Which half contains the solution?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • FORWARD 1 • Q2/5

True or false?

After 4 bisection steps starting from an interval of length 1, the solution is known within 1/16.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • FORWARD 1 • Q3/5

How many bisection steps from [1, 2] guarantee an interval shorter than 0.01?

A. 5

B. 100

C. 10

D. 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S34-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • FORWARD 1 • Q4/5

Explain why checking h(1.715) and h(1.725) decides whether the solution rounds to 1.72.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 34 • FORWARD 1 • Q5/5

SET 34 • HSA-REI.D.11

FORWARD 2

Compare an approximation with an exact logarithmic solution • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $2^x = 5$ exactly, then give a decimal to the nearest hundredth. A calculator may be used.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 34 • FORWARD 2 • Q1/5

Solve $3(0.8)^x = 1$ exactly and to the nearest hundredth.
A calculator may be used.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 34 • FORWARD 2 • Q2/5

True or false?

$2^x = 5 - x$ can be solved exactly with one logarithm.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 34 • FORWARD 2 • Q3/5

Which equation can be solved exactly with logarithms?

A. $\log x = 2 - x$

B. $2^x = x + 2$

C. $5(1.2)^x = 30$

D. $3(0.8)^x = x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F2-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 34 • FORWARD 2 • Q4/5

Explain why approximation methods are needed for $2^x = x + 2$ but not for $2^x = 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S34-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 34 • FORWARD 2 • Q5/5

SET 35 • HSF-TF.A.1 • TRIGONOMETRIC FUNCTIONS

Use radian measure as arc length on the unit circle

MAIN	Algebra II main question	**HSF-TF.A.1**
BACKWARD 1	Circumference of a circle	**7.G.B.4**
BACKWARD 2	Arc length is proportional to the angle	**HSG-C.B.5**
BACKWARD 3	Degrees as fractions of a turn	**4.MD.C.5**
FORWARD 1	Measure negative angles and angles beyond one turn	**HSF-TF.A.1**
FORWARD 2	Read sine and cosine of radian angles from the unit circle	**HSF-TF.A.2**

SET 35 • HSF-TF.A.1

MAIN

Algebra II main question • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1

On the unit circle, a central angle subtends an arc of length 2.5. What is the radian measure of the angle? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-M-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • MAIN • Q1/5

Convert 150° to radians. Give an exact answer.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-M-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • MAIN • Q2/5

Convert 7π/4 radians to degrees.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-M-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • MAIN • Q3/5

A central angle of 2π/3 radians is drawn in a circle of radius 6 cm. Find the length of the arc it subtends.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-M-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • MAIN • Q4/5

Explain why one full turn measures 2π radians.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-M-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • MAIN • Q5/5

SET 35 • HSF-TF.A.1

BACKWARD 1

Circumference of a circle • Grade 7 • Geometry • 7.G.B.4

Find the circumference of a circle of radius 1.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B1-Q1

Grade 7 • Geometry • 7.G.B.4 • Set 35 • BACKWARD 1 • Q1/5

Find the circumference of a circle of radius 6 cm.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B1-Q2

Grade 7 • Geometry • 7.G.B.4 • Set 35 • BACKWARD 1 • Q2/5

True or false?

The circumference of a circle is πd.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B1-Q3

Grade 7 • Geometry • 7.G.B.4 • Set 35 • BACKWARD 1 • Q3/5

Half of the unit circle has length

A. 2π

B. 1

C. π

D. $\pi/2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B1-Q4

Grade 7 • Geometry • 7.G.B.4 • Set 35 • BACKWARD 1 • Q4/5

What length is one quarter of the unit circle?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B1-Q5

Grade 7 • Geometry • 7.G.B.4 • Set 35 • BACKWARD 1 • Q5/5

SET 35 • HSF-TF.A.1

BACKWARD 2

Arc length is proportional to the angle • High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5

A 60° angle subtends what fraction of a full circle?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B2-Q1

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 35 • BACKWARD 2 • Q1/5

Find the arc length of a 90° angle in a circle of radius 4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B2-Q2

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 35 • BACKWARD 2 • Q2/5

True or false?

Doubling the radius doubles the arc length for the same angle.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B2-Q3

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 35 • BACKWARD 2 • Q3/5

The arc length for angle θ (radians) in a circle of radius r is

A. θ/r

B. $r^2\theta$

C. $2\pi\theta$

D. $r\theta$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B2-Q4

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 35 • BACKWARD 2 • Q4/5

What fraction of the circle is an angle of π/3 radians?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B2-Q5

High School — Geometry (course placement in this bank) • Geometry: Circles • HSG-C.B.5 • Set 35 • BACKWARD 2 • Q5/5

SET 35 • HSF-TF.A.1

BACKWARD 3

Degrees as fractions of a turn • Grade 4 • Measurement & Data • 4.MD.C.5

How many degrees are in a full turn?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B3-Q1

Grade 4 • Measurement & Data • 4.MD.C.5 • Set 35 • BACKWARD 3 • Q1/5

What fraction of a turn is 150°?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B3-Q2

Grade 4 • Measurement & Data • 4.MD.C.5 • Set 35 • BACKWARD 3 • Q2/5

True or false?

A 315° angle is 7/8 of a turn.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B3-Q3

Grade 4 • Measurement & Data • 4.MD.C.5 • Set 35 • BACKWARD 3 • Q3/5

A quarter turn is

A. 90°

B. 180°

C. 45°

D. 360°

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B3-Q4

Grade 4 • Measurement & Data • 4.MD.C.5 • Set 35 • BACKWARD 3 • Q4/5

How many degrees are in 2/3 of a turn?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-B3-Q5

Grade 4 • Measurement & Data • 4.MD.C.5 • Set 35 • BACKWARD 3 • Q5/5

SET 35 • HSF-TF.A.1

FORWARD 1

Measure negative angles and angles beyond one turn • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1

Give a positive angle less than 2π that is coterminal with $-\pi/3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • FORWARD 1 • Q1/5

Give the angle between 0 and 2π that is coterminal with 13π/6.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • FORWARD 1 • Q2/5

True or false?

An arc of length 10 on the unit circle goes more than once around the circle.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • FORWARD 1 • Q3/5

An angle of $-\pi/2$ radians is traced

A. a quarter turn clockwise

B. one full turn

C. half a turn

D. a quarter turn counterclockwise

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • FORWARD 1 • Q4/5

A point travels 10 units counterclockwise around the unit circle from (1, 0). How far past one full turn is it? Give a decimal to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 35 • FORWARD 1 • Q5/5

SET 35 • HSF-TF.A.1

FORWARD 2

Read sine and cosine of radian angles from the unit circle • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

What point of the unit circle is at angle π? Give cos π and sin π.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 35 • FORWARD 2 • Q1/5

Find $\cos(3\pi/2)$ and $\sin(3\pi/2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 35 • FORWARD 2 • Q2/5

True or false?

$\cos(2\pi) = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 35 • FORWARD 2 • Q3/5

$\sin(\pi/2)$ equals

A. -1

B. 0

C. $\pi/2$

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 35 • FORWARD 2 • Q4/5

Explain why cos θ and sin θ make sense for θ = 13π/6.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S35-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 35 • FORWARD 2 • Q5/5

SET 36 • HSF-TF.A.2 • TRIGONOMETRIC FUNCTIONS

Find exact trigonometric values beyond the first quadrant

MAIN	Algebra II main question	**HSF-TF.A.2**
BACKWARD 1	Special right triangle ratios	**HSG-SRT.C.6**
BACKWARD 2	Reference angles for radian measures	**HSF-TF.A.1**
BACKWARD 3	Signs of coordinates by quadrant	**6.NS.C.6.b**
FORWARD 1	Values for angles beyond 2π and solving for angles	**HSF-TF.A.2**
FORWARD 2	Verify the Pythagorean identity at exact values	**HSF-TF.C.8**

SET 36 • HSF-TF.A.2

MAIN

Algebra II main question • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Use the unit circle to find $\sin(5\pi/6)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-M-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • MAIN • Q1/5

Use the unit circle to find cos(4π/3) exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-M-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • MAIN • Q2/5

Use the unit circle to find tan(7π/4) exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-M-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • MAIN • Q3/5

Use the unit circle to find $\sin(-\pi/2)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-M-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • MAIN • Q4/5

Use the unit circle to find cos(11π/6) exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-M-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • MAIN • Q5/5

SET 36 • HSF-TF.A.2

BACKWARD 1

Special right triangle ratios • High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6

In a 30°−60°−90° triangle with hypotenuse 1, give the side lengths.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B1-Q1

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 36 • BACKWARD 1 • Q1/5

Find sin 45° exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B1-Q2

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 36 • BACKWARD 1 • Q2/5

True or false?

$\cos 60° = 1/2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B1-Q3

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 36 • BACKWARD 1 • Q3/5

sin 60° equals

A. √2/2

B. √3/2

C. √3

D. 1/2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S36-B1-Q4**

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 36 • BACKWARD 1 • Q4/5

Find tan 30° exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B1-Q5

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 36 • BACKWARD 1 • Q5/5

SET 36 • HSF-TF.A.2

BACKWARD 2

Reference angles for radian measures • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1

Find the reference angle of 5π/6.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 36 • BACKWARD 2 • Q1/5

Find the reference angle of $4\pi/3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 36 • BACKWARD 2 • Q2/5

True or false?

The reference angle of $7\pi/4$ is $\pi/4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 36 • BACKWARD 2 • Q3/5

Which quadrant contains 11π/6?

A. I

B. IV

C. III

D. II

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 36 • BACKWARD 2 • Q4/5

Convert 4π/3 to degrees.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 36 • BACKWARD 2 • Q5/5

SET 36 • HSF-TF.A.2

BACKWARD 3

Signs of coordinates by quadrant • Grade 6 • The Number System • 6.NS.C.6.b

In which quadrant are both coordinates negative?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B3-Q1

Grade 6 • The Number System • 6.NS.C.6.b • Set 36 • BACKWARD 3 • Q1/5

What are the signs of the coordinates in quadrant II?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B3-Q2

Grade 6 • The Number System • 6.NS.C.6.b • Set 36 • BACKWARD 3 • Q2/5

True or false?

A point in quadrant IV has a positive x-coordinate.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B3-Q3

Grade 6 • The Number System • 6.NS.C.6.b • Set 36 • BACKWARD 3 • Q3/5

The point $(-\sqrt{3}/2, 1/2)$ is in quadrant

A. I

B. II

C. III

D. IV

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B3-Q4

Grade 6 • The Number System • 6.NS.C.6.b • Set 36 • BACKWARD 3 • Q4/5

Give the signs of x and y for a point in quadrant IV.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-B3-Q5

Grade 6 • The Number System • 6.NS.C.6.b • Set 36 • BACKWARD 3 • Q5/5

SET 36 • HSF-TF.A.2

FORWARD 1

Values for angles beyond 2π and solving for angles • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Find sin(13π/6) exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • FORWARD 1 • Q1/5

Find $\cos(-3\pi/4)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • FORWARD 1 • Q2/5

Find every θ with $0 \le \theta < 2\pi$ and $\cos\theta = -1/2$.

$\theta =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • FORWARD 1 • Q3/5

True or false?

$\tan(\pi/2)$ is undefined.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • FORWARD 1 • Q4/5

Which angle has $\sin\theta = -\sqrt{3}/2$ and $\cos\theta = 1/2$?

A. $5\pi/3$

B. $4\pi/3$

C. $2\pi/3$

D. $7\pi/6$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 36 • FORWARD 1 • Q5/5

SET 36 • HSF-TF.A.2

FORWARD 2

Verify the Pythagorean identity at exact values • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8

Verify $\sin^2\theta + \cos^2\theta = 1$ for $\theta = 5\pi/6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 36 • FORWARD 2 • Q1/5

Verify $\sin^2\theta + \cos^2\theta = 1$ for $\theta = 7\pi/4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 36 • FORWARD 2 • Q2/5

True or false?

$\sin^2\theta + \cos^2\theta = 1$ is true only for first-quadrant angles.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 36 • FORWARD 2 • Q3/5

If $\cos\theta = -1/2$ and θ is in quadrant III, then $\sin\theta =$

A. $\sqrt{3}/2$

B. $-\sqrt{3}/2$

C. $1/2$

D. $-1/2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 36 • FORWARD 2 • Q4/5

Explain why the values found from the unit circle always satisfy $\sin^2\theta + \cos^2\theta = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S36-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 36 • FORWARD 2 • Q5/5

SET 37 • HSF-TF.A.2 • TRIGONOMETRIC FUNCTIONS

Explain the extension of sine and cosine to all real numbers

MAIN	Algebra II main question	**HSF-TF.A.2**
BACKWARD 1	Right-triangle definitions of sine and cosine	**HSG-SRT.C.6**
BACKWARD 2	Points on the unit circle	**HSG-GPE.A.1**
BACKWARD 3	Radian measure as directed arc length	**HSF-TF.A.1**
FORWARD 1	Explain where tangent is defined and how values repeat	**HSF-TF.A.2**
FORWARD 2	Explain symmetry and periodicity	**HSF-TF.A.4**

SET 37 • HSF-TF.A.2

MAIN

Algebra II main question • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Explain how the unit circle defines cos t and sin t for every real number t.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-M-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • MAIN • Q1/5

Explain why $\sin(t + 2\pi) = \sin t$ for every real number t.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-M-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • MAIN • Q2/5

Explain why $\cos t < 0$ for $\pi/2 < t < \pi$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-M-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • MAIN • Q3/5

Explain why $-1 \le \sin t \le 1$ for every real t.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-M-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • MAIN • Q4/5

Explain how the definition gives $\sin(-\pi/6)$, and find it.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-M-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • MAIN • Q5/5

SET 37 • HSF-TF.A.2

BACKWARD 1

Right-triangle definitions of sine and cosine • High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6

Define sin A in a right triangle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B1-Q1

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 37 • BACKWARD 1 • Q1/5

In a right triangle with hypotenuse 1, what are the legs next to and opposite an acute angle θ?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B1-Q2

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 37 • BACKWARD 1 • Q2/5

True or false?

The right-triangle definition works for $\theta = 120°$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B1-Q3

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 37 • BACKWARD 1 • Q3/5

In a right triangle with hypotenuse 1, the leg opposite θ is

A. $\sin \theta$

B. $\tan \theta$

C. 1

D. $\cos \theta$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S37-B1-Q4**

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 37 • BACKWARD 1 • Q4/5

Why does the right-triangle definition fail for negative angles?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B1-Q5

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 37 • BACKWARD 1 • Q5/5

SET 37 • HSF-TF.A.2

BACKWARD 2

Points on the unit circle • High School — Geometry (course placement in this bank) • Geometry: Expressing Geometric Properties with Equations • HSG-GPE.A.1

Write the equation of the circle with center (0, 0) and radius 1.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B2-Q1

High School — Geometry (course placement in this bank) • Geometry: Expressing Geometric Properties with Equations
HSG-GPE.A.1 • Set 37 • BACKWARD 2 • Q1/5

Is $(-1/2, \sqrt{3}/2)$ on the unit circle?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B2-Q2

High School — Geometry (course placement in this bank) • Geometry: Expressing Geometric Properties with Equations HSG-GPE.A.1 • Set 37 • BACKWARD 2 • Q2/5

True or false?

Every point on the unit circle has $-1 \leq x \leq 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B2-Q3

High School — Geometry (course placement in this bank) • Geometry: Expressing Geometric Properties with Equations
HSG-GPE.A.1 • Set 37 • BACKWARD 2 • Q3/5

Which point is on the unit circle?

A. (0.9, 0.9)

B. (0.5, 0.5)

C. (1, 1)

D. (0.6, −0.8)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S37-B2-Q4**

High School — Geometry (course placement in this bank) • Geometry: Expressing Geometric Properties with Equations HSG-GPE.A.1 • Set 37 • BACKWARD 2 • Q4/5

A point on the unit circle has x = 0. Give both possible points.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B2-Q5

High School — Geometry (course placement in this bank) • Geometry: Expressing Geometric Properties with Equations
HSG-GPE.A.1 • Set 37 • BACKWARD 2 • Q5/5

SET 37 • HSF-TF.A.2

BACKWARD 3

Radian measure as directed arc length • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1

What arc length on the unit circle does t = π/6 cut off?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B3-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 37 • BACKWARD 3 • Q1/5

Which direction does $t = -\pi/6$ travel?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B3-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 37 • BACKWARD 3 • Q2/5

True or false?

$t = 2\pi$ returns to the starting point (1, 0).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B3-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 37 • BACKWARD 3 • Q3/5

An arc of length π from (1, 0) ends at

A. (0, 1)

B. (0, −1)

C. (1, 0)

D. (−1, 0)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B3-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 37 • BACKWARD 3 • Q4/5

Give a negative t that ends at $(0, -1)$.

$t =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-B3-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.1 • Set 37 • BACKWARD 3 • Q5/5

SET 37 • HSF-TF.A.2

FORWARD 1

Explain where tangent is defined and how values repeat • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Explain why tan t is undefined at $t = \pi/2 + k\pi$ (k an integer).

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • FORWARD 1 • Q1/5

Give the domain and range of $y = \sin t$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • FORWARD 1 • Q2/5

True or false?

cos t and sin t can both equal 0 for the same t.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • FORWARD 1 • Q3/5

The unit-circle definition agrees with the triangle definition for

A. $t < 0$ only

B. $\pi/2 < t < \pi$ only

C. all t

D. $0 < t < \pi/2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • FORWARD 1 • Q4/5

Explain why $\cos t = \cos(-t)$ using the unit circle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 37 • FORWARD 1 • Q5/5

SET 37 • HSF-TF.A.2

FORWARD 2

Explain symmetry and periodicity • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4

Use the unit circle to explain why $\sin(-t) = -\sin t$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 37 • FORWARD 2 • Q1/5

Is sine an even function, an odd function, or neither?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 37 • FORWARD 2 • Q2/5

True or false?

$\tan(t + \pi) = \tan t$ for every t where it is defined.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 37 • FORWARD 2 • Q3/5

The period of $y = \tan t$ is

A. π

B. 4π

C. $\pi/2$

D. 2π

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 37 • FORWARD 2 • Q4/5

Explain why cos is an even function.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S37-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 37 • FORWARD 2 • Q5/5

SET 38 • HSF-TF.B.5 • TRIGONOMETRIC FUNCTIONS

Write sinusoidal models with given amplitude, period, and midline

MAIN	Algebra II main question	**HSF-TF.B.5**
BACKWARD 1	Amplitude and midline from the maximum and minimum	**7.NS.A.3**
BACKWARD 2	Period and frequency	**6.RP.A.2**
BACKWARD 3	Vertical stretches, shifts, and reflections	**HSF-BF.B.3**
FORWARD 1	Write models from data or with a phase shift	**HSF-TF.B.5**
FORWARD 2	Solve for times in sinusoidal models	**HSF-TF.B.7**

SET 38 • HSF-TF.B.5

MAIN

Algebra II main question • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

Write a cosine function with amplitude 3, period 2π, midline $y = 1$, and a maximum at $x = 0$.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-M-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • MAIN • Q1/5

A Ferris wheel has diameter 40 m, its center is 25 m above the ground, and it turns once every 4 minutes. A rider starts at the bottom. Write the rider's height h after t minutes.

h = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-M-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • MAIN • Q2/5

Water depth at a harbor varies between 1 ft (low tide) and 9 ft (high tide) with a period of 12 hours. High tide is at t = 0. Write the depth d after t hours.

d = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-M-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • MAIN • Q3/5

A town's average temperature is lowest, 10°C, in month t = 0 and highest, 30°C, at t = 6, with a 12-month period. Write a model T(t).

T = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-M-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • MAIN • Q4/5

Write a sine function with amplitude 2, period π, and midline $y = -1$ that is at the midline and increasing at $x = 0$.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-M-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • MAIN • Q5/5

SET 38 • HSF-TF.B.5

BACKWARD 1

Amplitude and midline from the maximum and minimum • Grade 7 • The Number System • 7.NS.A.3

Find (9 + 1)/2 and (9 − 1)/2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B1-Q1

Grade 7 • The Number System • 7.NS.A.3 • Set 38 • BACKWARD 1 • Q1/5

A quantity varies between 10 and 30. Find the midline value and amplitude.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B1-Q2

Grade 7 • The Number System • 7.NS.A.3 • Set 38 • BACKWARD 1 • Q2/5

True or false?

The amplitude is the maximum minus the minimum.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B1-Q3

Grade 7 • The Number System • 7.NS.A.3 • Set 38 • BACKWARD 1 • Q3/5

For max 45 and min 5, the midline is

A. y = 20

B. y = 50

C. y = 25

D. y = 40

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B1-Q4

Grade 7 • The Number System • 7.NS.A.3 • Set 38 • BACKWARD 1 • Q4/5

A Ferris wheel has diameter 40 m with center 25 m high. Give its maximum and minimum heights.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B1-Q5

Grade 7 • The Number System • 7.NS.A.3 • Set 38 • BACKWARD 1 • Q5/5

SET 38 • HSF-TF.B.5

BACKWARD 2

Period and frequency • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2

A wheel turns once every 4 minutes. How many turns per minute?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B2-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 38 • BACKWARD 2 • Q1/5

A spring oscillates 3 times per second. What is the period?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B2-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 38 • BACKWARD 2 • Q2/5

True or false?

Frequency is the reciprocal of the period.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B2-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 38 • BACKWARD 2 • Q3/5

A period of 12 hours means a frequency of

A. 12 cycles per hour

B. 2 cycles per hour

C. 1/2 cycle per hour

D. 1/12 cycle per hour

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B2-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 38 • BACKWARD 2 • Q4/5

$y = \sin(Bx)$ has period $2\pi/B$. Find B for period 4.

B = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B2-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 38 • BACKWARD 2 • Q5/5

SET 38 • HSF-TF.B.5

BACKWARD 3

Vertical stretches, shifts, and reflections • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

Describe how $y = 3\cos x + 1$ is obtained from $y = \cos x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 38 • BACKWARD 3 • Q1/5

Describe $y = -\cos x$ compared with $y = \cos x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 38 • BACKWARD 3 • Q2/5

True or false?

$y = \cos x + 5$ has a maximum of 6.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 38 • BACKWARD 3 • Q3/5

$y = -20\cos x$ starts (at $x = 0$) at

A. its minimum

B. its maximum

C. the midline going up

D. the midline going down

Write y = sin x shifted down 1 and stretched vertically by 2.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 38 • BACKWARD 3 • Q5/5

SET 38 • HSF-TF.B.5

FORWARD 1

Write models from data or with a phase shift • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

A rider's height is 3 m at t = 0 (bottom), 27 m at the top after 15 s, and the motion repeats every 30 s. Write h(t).

h = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • FORWARD 1 • Q1/5

Sunrise times vary between 5:30 and 7:30 (as 5.5 and 7.5 hours) with period 365 days, latest at day 0. Write a model.

S = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • FORWARD 1 • Q2/5

True or false?

$y = 4\sin(x - \pi/2)$ is the same as $y = -4\cos x$.

A. True

B. False

A sine model with period 10, amplitude 3, midline 7, and a maximum at t = 2.5 is

A. $7 + 3\sin(10t)$

B. $7 + 3\sin(\pi t/5)$

C. $3 + 7\sin(\pi t/5)$

D. $7 + 3\cos(\pi t/5)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • FORWARD 1 • Q4/5

A model has a maximum of 18 at t = 3 and the next minimum of 6 at t = 7. Write a cosine model.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 38 • FORWARD 1 • Q5/5

SET 38 • HSF-TF.B.5

FORWARD 2

Solve for times in sinusoidal models • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7

For $h = 25 - 20\cos(\pi t/2)$, find the first time $t > 0$ when the rider is at 35 m. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 38 • FORWARD 2 • Q1/5

For $d = 4\cos(\pi t/6) + 5$, find all times in $0 \leq t < 12$ when the depth is 5 ft.

$t =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 38 • FORWARD 2 • Q2/5

True or false?

A sinusoidal model usually reaches a given value between its min and max twice per period.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 38 • FORWARD 2 • Q3/5

For $T = 20 - 10\cos(\pi t/6)$, $T = 25$ first when

A. $t = 2$

B. $t = 4$

C. $t = 3$

D. $t = 6$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 38 • FORWARD 2 • Q4/5

Explain why there are two solutions per period when the value is strictly between the minimum and maximum.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S38-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 38 • FORWARD 2 • Q5/5

SET 39 • HSF-TF.B.5 • TRIGONOMETRIC FUNCTIONS

Interpret the parameters of sinusoidal models

MAIN	Algebra II main question	**HSF-TF.B.5**
BACKWARD 1	Build the model from amplitude, period, and midline	**HSF-TF.B.5**
BACKWARD 2	Evaluate a sinusoidal model	**HSF-TF.A.2**
BACKWARD 3	Convert rates between units	**6.RP.A.3.d**
FORWARD 1	Predict how changing a parameter changes the situation	**HSF-TF.B.5**
FORWARD 2	Interpret periodic features from graphs and tables	**HSF-IF.B.4**

SET 39 • HSF-TF.B.5

MAIN

Algebra II main question • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

A Ferris wheel rider's height is $h(t) = 12 + 10\sin(\pi t/15)$ meters after t seconds. Give the maximum height, the minimum height, and the time for one revolution.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-M-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • MAIN • Q1/5

Hours of daylight are $D(t) = 12 + 3\sin(2\pi(t - 80)/365)$, t in days. Interpret 12, 3, and 365.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-M-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • MAIN • Q2/5

A model of blood pressure is $P(t) = 100 + 20\sin(2.5\pi t)$, t in seconds. Find the period and the number of heartbeats per minute.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-M-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • MAIN • Q3/5

A spring's position is $y = -4\cos(3\pi t)$ cm, t in seconds. Give the amplitude, period, and frequency.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-M-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • MAIN • Q4/5

Two sounds are $y = \sin(880\pi t)$ and $y = \sin(440\pi t)$, t in seconds. Give the frequency of each in cycles per second and say which is higher.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-M-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • MAIN • Q5/5

SET 39 • HSF-TF.B.5

BACKWARD 1

Build the model from amplitude, period, and midline • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

What is the amplitude of $y = 12 + 10\sin(\pi t/15)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • BACKWARD 1 • Q1/5

What is the midline of $y = 12 + 10\sin(\pi t/15)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • BACKWARD 1 • Q2/5

True or false?

The period of $y = \sin(Bt)$ is $2\pi/B$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • BACKWARD 1 • Q3/5

The period of $y = \sin(\pi t/15)$ is

A. 30

B. 15

C. $\pi/15$

D. 2π

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • BACKWARD 1 • Q4/5

Write a model with amplitude 4, period 2/3, and midline 0.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • BACKWARD 1 • Q5/5

SET 39 • HSF-TF.B.5

BACKWARD 2

Evaluate a sinusoidal model • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Evaluate $12 + 10\sin(\pi/2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 39 • BACKWARD 2 • Q1/5

Evaluate $12 + 10\sin(3\pi/2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 39 • BACKWARD 2 • Q2/5

True or false?

$-4\cos(0) = -4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 39 • BACKWARD 2 • Q3/5

100 + 20sin(π/2) equals

A. 100

B. 20

C. 120

D. 80

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 39 • BACKWARD 2 • Q4/5

Evaluate 12 + 3sin(0).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 39 • BACKWARD 2 • Q5/5

SET 39 • HSF-TF.B.5

BACKWARD 3

Convert rates between units • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.d

A cycle lasts 0.8 s. How many cycles are in 60 s?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.d • Set 39 • BACKWARD 3 • Q1/5

1.5 cycles per second is how many cycles per minute?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.d • Set 39 • BACKWARD 3 • Q2/5

True or false?

A period of 30 s means 2 revolutions per minute.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.d • Set 39 • BACKWARD 3 • Q3/5

440 cycles per second is how many cycles per millisecond?

A. 4.4

B. 44

C. 440

D. 0.44

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.d • Set 39 • BACKWARD 3 • Q4/5

A tide has period 12 hours. How many cycles in 3 days?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.d • Set 39 • BACKWARD 3 • Q5/5

SET 39 • HSF-TF.B.5

FORWARD 1

Predict how changing a parameter changes the situation • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

The Ferris wheel is replaced by one that turns twice as fast. Rewrite $h(t) = 12 + 10\sin(\pi t/15)$.

$h =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • FORWARD 1 • Q1/5

The daylight model D(t) is changed for a city farther from the equator, with daylight between 8 and 16 hours. What parameter changes, and to what?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • FORWARD 1 • Q2/5

True or false?

Changing the midline changes the period.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • FORWARD 1 • Q3/5

Multiplying B by 3 makes the period

A. 3 units longer

B. unchanged

C. three times as long

D. one third as long

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • FORWARD 1 • Q4/5

A heart rate rises from 75 to 100 beats per minute. What is the new period, in seconds?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 39 • FORWARD 1 • Q5/5

SET 39 • HSF-TF.B.5

FORWARD 2

Interpret periodic features from graphs and tables • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

A table of a tide's depth shows maxima at t = 2 and t = 14 hours. What is the period?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 39 • FORWARD 2 • Q1/5

A graph of a temperature model has a high of 30 and a low of 10. What are the midline and amplitude?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 39 • FORWARD 2 • Q2/5

True or false?

Consecutive minima of a periodic function are one period apart.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 39 • FORWARD 2 • Q3/5

On which interval does $h(t) = 12 + 10\sin(\pi t/15)$ increase?

A. $0 < t < 15$

B. $0 < t < 7.5$

C. $15 < t < 30$

D. $7.5 < t < 22.5$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S39-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 39 • FORWARD 2 • Q4/5

Explain how to read the frequency of a periodic graph.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S39-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 39 • FORWARD 2 • Q5/5

SET 40 • HSF-TF.C.8 • TRIGONOMETRIC FUNCTIONS

Prove the Pythagorean identity

MAIN	Algebra II main question	**HSF-TF.C.8**
BACKWARD 1	The Pythagorean Theorem	**8.G.B.7**
BACKWARD 2	Coordinates of the point at angle θ	**HSF-TF.A.2**
BACKWARD 3	Expand and simplify algebraic squares	**HSA-SSE.A.2**
FORWARD 1	Prove identities derived from the Pythagorean identity	**HSF-TF.C.8**
FORWARD 2	Use the angle-addition formulas	**HSF-TF.C.9**

SET 40 • HSF-TF.C.8

MAIN

Algebra II main question • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8

Prove that $\sin^2\theta + \cos^2\theta = 1$ for every angle θ, using the unit circle.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-M-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • MAIN • Q1/5

A right triangle has legs a and b, hypotenuse c, and acute angle θ opposite a. Prove $\sin^2\theta + \cos^2\theta = 1$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-M-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • MAIN • Q2/5

Prove that $1 + \tan^2\theta = 1/\cos^2\theta$ wherever $\cos\theta \neq 0$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-M-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • MAIN • Q3/5

θ is in quadrant II. Prove $\sin^2\theta + \cos^2\theta = 1$ using a reference triangle, explaining what happens to the signs.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-M-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • MAIN • Q4/5

Prove that $(\sin\theta + \cos\theta)^2 = 1 + 2\sin\theta\cos\theta$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-M-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • MAIN • Q5/5

SET 40 • HSF-TF.C.8

BACKWARD 1

The Pythagorean Theorem • Grade 8 • Geometry • 8.G.B.7

A right triangle has legs 3 and 4. Find the hypotenuse.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B1-Q1

Grade 8 • Geometry • 8.G.B.7 • Set 40 • BACKWARD 1 • Q1/5

A right triangle has hypotenuse 1 and a leg 0.6. Find the other leg.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B1-Q2

Grade 8 • Geometry • 8.G.B.7 • Set 40 • BACKWARD 1 • Q2/5

True or false?

In a right triangle, $a^2 + b^2 = c^2$ where c is the hypotenuse.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B1-Q3

Grade 8 • Geometry • 8.G.B.7 • Set 40 • BACKWARD 1 • Q3/5

Legs 5 and 12 give hypotenuse

A. 60

B. 7

C. 17

D. 13

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B1-Q4

Grade 8 • Geometry • 8.G.B.7 • Set 40 • BACKWARD 1 • Q4/5

Legs a and b and hypotenuse c: write c^2 in terms of a and b.

$c^2 =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B1-Q5

Grade 8 • Geometry • 8.G.B.7 • Set 40 • BACKWARD 1 • Q5/5

SET 40 • HSF-TF.C.8

BACKWARD 2

Coordinates of the point at angle θ • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Give the coordinates of the point on the unit circle at angle θ.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 40 • BACKWARD 2 • Q1/5

Give $\cos(2\pi/3)$ and $\sin(2\pi/3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 40 • BACKWARD 2 • Q2/5

True or false?

sin θ is the x-coordinate of the point at angle θ.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 40 • BACKWARD 2 • Q3/5

The point at angle π/4 is

A. (1/2, √3/2)

B. (1, 1)

C. (√3/2, 1/2)

D. (√2/2, √2/2)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 40 • BACKWARD 2 • Q4/5

What is the radius of the unit circle?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 40 • BACKWARD 2 • Q5/5

SET 40 • HSF-TF.C.8

BACKWARD 3

Expand and simplify algebraic squares • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Expand $(a + b)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B3-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 40 • BACKWARD 3 • Q1/5

Simplify $(a/c)^2 + (b/c)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B3-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 40 • BACKWARD 3 • Q2/5

True or false?

$(\sin \theta)^2$ is written $\sin^2\theta$.

A. True

B. False

Dividing $x^2 + y^2 = 1$ by x^2 gives

A. $1 + y^2/x^2 = 1/x^2$

B. $1 + y^2 = 1/x$

C. $y^2 = 1$

D. $x^2 + y^2 = 1/x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B3-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 40 • BACKWARD 3 • Q4/5

Expand $(s + c)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-B3-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 40 • BACKWARD 3 • Q5/5

SET 40 • HSF-TF.C.8

FORWARD 1

Prove identities derived from the Pythagorean identity • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8

Prove that $\sin^4\theta - \cos^4\theta = \sin^2\theta - \cos^2\theta$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • FORWARD 1 • Q1/5

Prove that $(1 - \cos\theta)(1 + \cos\theta) = \sin^2\theta$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • FORWARD 1 • Q2/5

True or false?

$\sin\theta + \cos\theta = 1$ for every θ.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • FORWARD 1 • Q3/5

Which equals $1 - \sin^2\theta$?

A. $\tan^2\theta$

B. $1 - \cos^2\theta$

C. $\cos\theta$

D. $\cos^2\theta$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • FORWARD 1 • Q4/5

Prove that $\cos^2\theta - \sin^2\theta = 2\cos^2\theta - 1$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 40 • FORWARD 1 • Q5/5

SET 40 • HSF-TF.C.8

FORWARD 2

Use the angle-addition formulas • High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9

Use sin(a + b) = sin a cos b + cos a sin b to find sin(75°) exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F2-Q1

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 40 • FORWARD 2 • Q1/5

Use $\cos(a + b) = \cos a \cos b - \sin a \sin b$ with $a = b = \theta$ to write $\cos 2\theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F2-Q2

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 40 • FORWARD 2 • Q2/5

True or false?

$\cos 2\theta = 1 - 2\sin^2\theta$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F2-Q3

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 40 • FORWARD 2 • Q3/5

$\sin 2\theta$ equals

A. $\sin^2\theta - \cos^2\theta$

B. $2\sin\theta$

C. $2\sin\theta\cos\theta$

D. $\sin\theta + \cos\theta$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S40-F2-Q4**

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 40 • FORWARD 2 • Q4/5

Use $(\sin\theta + \cos\theta)^2 = 1 + 2\sin\theta\cos\theta$ to write $(\sin\theta + \cos\theta)^2$ in terms of $\sin 2\theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S40-F2-Q5

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 40 • FORWARD 2 • Q5/5

SET 41 • HSF-TF.C.8 • TRIGONOMETRIC FUNCTIONS

Use the Pythagorean identity to find trigonometric values

MAIN	Algebra II main question	**HSF-TF.C.8**
BACKWARD 1	Signs of sine, cosine, and tangent by quadrant	**HSF-TF.A.2**
BACKWARD 2	Square roots of fractions	**8.EE.A.2**
BACKWARD 3	Solve $x^2 = c$	**HSA-REI.B.4.b**
FORWARD 1	Find values when the quadrant is not given, and combine values	**HSF-TF.C.8**
FORWARD 2	Find the angles with inverse trigonometric functions	**HSF-TF.B.7**

SET 41 • HSF-TF.C.8

MAIN

Algebra II main question • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8

$\sin \theta = 3/5$ and θ is in quadrant II. Find $\cos \theta$ and $\tan \theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-M-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • MAIN • Q1/5

$\cos\theta = -5/13$ and θ is in quadrant III. Find $\sin\theta$ and $\tan\theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-M-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • MAIN • Q2/5

$\tan\theta = -2$ and θ is in quadrant IV. Find $\sin\theta$ and $\cos\theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-M-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • MAIN • Q3/5

$\sin\theta = -1/3$ and θ is in quadrant IV. Find $\cos\theta$ and $\tan\theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-M-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • MAIN • Q4/5

$\cos\theta = 0.6$ and $\sin\theta < 0$. Find $\sin\theta$ and $\tan\theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-M-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • MAIN • Q5/5

SET 41 • HSF-TF.C.8

BACKWARD 1

Signs of sine, cosine, and tangent by quadrant • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

In quadrant II, which of sin, cos, tan are positive?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 41 • BACKWARD 1 • Q1/5

In quadrant III, what is the sign of tan θ?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 41 • BACKWARD 1 • Q2/5

True or false?

In quadrant IV, $\cos\theta > 0$ and $\sin\theta < 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 41 • BACKWARD 1 • Q3/5

If $\sin\theta < 0$ and $\cos\theta > 0$, θ is in quadrant

A. I

B. II

C. IV

D. III

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 41 • BACKWARD 1 • Q4/5

If $\tan\theta < 0$ and $\sin\theta > 0$, which quadrant is θ in?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 41 • BACKWARD 1 • Q5/5

SET 41 • HSF-TF.C.8

BACKWARD 2

Square roots of fractions • Grade 8 • Expressions & Equations • 8.EE.A.2

Find √(16/25).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 41 • BACKWARD 2 • Q1/5

Find √(8/9) in simplest radical form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 41 • BACKWARD 2 • Q2/5

True or false?

$\sqrt{(144/169)} = 12/13$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 41 • BACKWARD 2 • Q3/5

√(1/5) equals

A. 5

B. √5

C. 1/25

D. √5/5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 41 • BACKWARD 2 • Q4/5

Find $\sqrt{0.64}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 41 • BACKWARD 2 • Q5/5

SET 41 • HSF-TF.C.8

BACKWARD 3

Solve $x^2 = c$ • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Solve $x^2 = 16/25$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 41 • BACKWARD 3 • Q1/5

Solve $x^2 = 1 - 9/25$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 41 • BACKWARD 3 • Q2/5

True or false?

$x^2 = 0.36$ has only the solution 0.6.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B3-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 41 • BACKWARD 3 • Q3/5

$x^2 + (3/5)^2 = 1$ gives $x =$

A. $\pm 2/5$

B. $2/5$

C. $4/5$

D. $\pm 4/5$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 41 • BACKWARD 3 • Q4/5

Solve $5x^2 = 1$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 41 • BACKWARD 3 • Q5/5

SET 41 • HSF-TF.C.8

FORWARD 1

Find values when the quadrant is not given, and combine values • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8

$\sin\theta = 3/5$. List both possible values of $\cos\theta$ and the quadrants they go with.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • FORWARD 1 • Q1/5

$\cos\theta = -5/13$ and θ is in quadrant III. Find $\sin\theta \cdot \cos\theta$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • FORWARD 1 • Q2/5

True or false?

Knowing $\tan \theta = 1$ determines $\sin \theta$ uniquely.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • FORWARD 1 • Q3/5

If $\sin\theta = 0.8$ and $\tan\theta < 0$, then $\cos\theta =$

A. -0.8

B. 0.2

C. 0.6

D. -0.6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • FORWARD 1 • Q4/5

Use $1 + \tan^2\theta = 1/\cos^2\theta$ to find $\cos\theta$ if $\tan\theta = 3$ and θ is in quadrant III.

$\cos\theta =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.C.8 • Set 41 • FORWARD 1 • Q5/5

SET 41 • HSF-TF.C.8

FORWARD 2

Find the angles with inverse trigonometric functions • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7

$\sin\theta = 3/5$ with $0 \leq \theta < 2\pi$. Find both values of θ to the nearest thousandth. A calculator may be used.

$\theta \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 41 • FORWARD 2 • Q1/5

$\cos\theta = -5/13$ and θ is in quadrant III. Find θ to the nearest thousandth ($0 \leq \theta < 2\pi$). A calculator may be used.

$\theta \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 41 • FORWARD 2 • Q2/5

True or false?

arcsin gives only angles between $-\pi/2$ and $\pi/2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 41 • FORWARD 2 • Q3/5

The quadrant II angle with $\sin\theta = 3/5$ is

A. $\pi + \arcsin(3/5)$

B. $\pi - \arcsin(3/5)$

C. $\arcsin(3/5)$

D. $-\arcsin(3/5)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 41 • FORWARD 2 • Q4/5

Explain why the identity finds cos θ but the inverse function is needed to find θ itself.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S41-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 41 • FORWARD 2 • Q5/5

SET 42 • HSF-IF.B.4 • INTERPRETING FUNCTIONS

Interpret key features of polynomial and rational models

MAIN	Algebra II main question	**HSF-IF.B.4**
BACKWARD 1	Interpret key features of linear and quadratic models	**HSF-IF.B.4**
BACKWARD 2	Describe increasing and decreasing behavior	**8.F.B.5**
BACKWARD 3	Zeros and sign from factors	**HSA-APR.B.3**
FORWARD 1	Sketch a model from a verbal description of key features	**HSF-IF.B.4**
FORWARD 2	Compare key features across representations	**HSF-IF.C.9**

SET 42 • HSF-IF.B.4

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

An open box is made from a 12-in square sheet by cutting squares of side x from the corners. Its volume is $V(x) = x(12 - 2x)^2$. Use the graph to give the x that maximizes V, the maximum volume, and the interval where V increases.

Answer: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • MAIN • Q1/5

The graph of $f(x) = (x + 2)(x - 1)(x - 3)$ is shown. Give the intervals where $f(x) > 0$ and the intervals where $f(x) < 0$.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • MAIN • Q2/5

A drug concentration is $C(t) = 20t/(t^2 + 4)$ mg/L after t hours. Use the graph to give the maximum concentration, when it occurs, and the end behavior. Interpret the end behavior.

Answer: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • MAIN • Q3/5

The graph of $f(x) = x^4 - 4x^2$ is shown. Give its x-intercepts and its relative maximum and minimum points.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • MAIN • Q4/5

The average cost of making x items is A(x) = (100 + 2x)/x dollars. Use the graph to describe the end behavior and interpret it.

Answer: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • MAIN • Q5/5

SET 42 • HSF-IF.B.4

BACKWARD 1

Interpret key features of linear and quadratic models • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4

$h(t) = -16t^2 + 32t + 4$. When is the maximum height, and what is it?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • BACKWARD 1 • Q1/5

What does the y-intercept of C = 25 + 3x mean if C is the cost of x rides?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • BACKWARD 1 • Q2/5

True or false?

A parabola that opens down has a maximum.

A. True

B. False

$h(t) = -16t^2 + 32t + 4$ increases for

A. all t

B. $0 < t < 1$

C. $t > 1$

D. $1 < t < 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • BACKWARD 1 • Q4/5

Where is $y = (x - 2)(x - 6)$ negative?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • BACKWARD 1 • Q5/5

SET 42 • HSF-IF.B.4

BACKWARD 2

Describe increasing and decreasing behavior • Grade 8 • Functions • 8.F.B.5

A graph rises from x = 0 to x = 2 and then falls. Describe it in words.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B2-Q1

Grade 8 • Functions • 8.F.B.5 • Set 42 • BACKWARD 2 • Q1/5

Sketch-free: describe the graph of the concentration that rises then levels toward 0.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B2-Q2

Grade 8 • Functions • 8.F.B.5 • Set 42 • BACKWARD 2 • Q2/5

True or false?

A highest point between rising and falling parts is a relative maximum.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B2-Q3

Grade 8 • Functions • 8.F.B.5 • Set 42 • BACKWARD 2 • Q3/5

A function that falls on $1 < x < 3$ is

A. constant there

B. decreasing there

C. undefined there

D. increasing there

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B2-Q4

Grade 8 • Functions • 8.F.B.5 • Set 42 • BACKWARD 2 • Q4/5

What is a relative minimum?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B2-Q5

Grade 8 • Functions • 8.F.B.5 • Set 42 • BACKWARD 2 • Q5/5

SET 42 • HSF-IF.B.4

BACKWARD 3

Zeros and sign from factors • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

List the zeros of $(x + 2)(x - 1)(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B3-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 42 • BACKWARD 3 • Q1/5

List the zeros of $x^4 - 4x^2 = x^2(x - 2)(x + 2)$ with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B3-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 42 • BACKWARD 3 • Q2/5

True or false?

$V(x) = x(12 - 2x)^2$ has zeros 0 and 6.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B3-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 42 • BACKWARD 3 • Q3/5

$20t/(t^2 + 4) = 0$ when

A. never

B. $t = -2$

C. $t = 2$

D. $t = 0$

Is $(x + 2)(x - 1)(x - 3)$ positive or negative at $x = 2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-B3-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 42 • BACKWARD 3 • Q5/5

SET 42 • HSF-IF.B.4

FORWARD 1

Sketch a model from a verbal description of key features • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

Sketch a possible graph: a population is 50 at t = 0, rises to a maximum of 200 at t = 4, then decreases toward a level of 100 as t grows.



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • FORWARD 1 • Q1/5

Sketch a possible graph of a cubic with zeros at -1 and 3 only, positive y-intercept, falling to the right, and touching the x-axis at 3.



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • FORWARD 1 • Q2/5

True or false?

A graph can have a relative maximum that is not the absolute maximum.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • FORWARD 1 • Q3/5

A model rises to a peak and approaches 0 as t grows. Which formula fits?

A. $20t/(t^2 + 4)$

B. $20t$

C. t^2

D. $20 - t$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • FORWARD 1 • Q4/5

Write a polynomial with zeros 0 and 6 and a relative maximum in between, whose values are positive between the zeros.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 42 • FORWARD 1 • Q5/5

SET 42 • HSF-IF.B.4

FORWARD 2

Compare key features across representations • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9

$f(x) = x(12 - 2x)^2$ and g is a quadratic with vertex (3, 120). Which has the greater maximum on $0 < x < 6$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 42 • FORWARD 2 • Q1/5

$C(t) = 20t/(t^2 + 4)$ and a table of $D(t)$ gives $D(1) = 3$, $D(2) = 6$, $D(3) = 4$. Which has the greater peak among these values, and when?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 42 • FORWARD 2 • Q2/5

True or false?

Two functions with the same zeros have the same relative extrema.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 42 • FORWARD 2 • Q3/5

Which has a horizontal asymptote?

A. $(x + 2)(x - 1)(x - 3)$

B. $x^4 - 4x^2$

C. $x(12 - 2x)^2$

D. $(100 + 2x)/x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 42 • FORWARD 2 • Q4/5

Explain how to compare end behavior of a polynomial given by formula and a rational function given by a graph.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S42-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 42 • FORWARD 2 • Q5/5

SET 43 • HSF-IF.B.4 • INTERPRETING FUNCTIONS

Interpret periodic models from graphs and tables

MAIN	Algebra II main question	**HSF-IF.B.4**
BACKWARD 1	Interpret maximums and minimums on a graph	**HSF-IF.B.4**
BACKWARD 2	Read values from a table	**6.EE.C.9**
BACKWARD 3	Amplitude, period, and midline of a sinusoidal model	**HSF-TF.B.5**
FORWARD 1	Predict with periodicity	**HSF-IF.B.4**
FORWARD 2	Write a model from periodic features	**HSF-TF.B.5**

SET 43 • HSF-IF.B.4

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

The graph shows a city's temperature T (°C) t hours after midnight on two days. Give the period, the maximum temperature, and the first time it occurs.

Answer: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • MAIN • Q1/5

The table shows the water depth d (ft) at a harbor every 2 hours. Give the period, the depth at low tide, and the first time of low tide.

t (h)	0	2	4	6	8	10	12	14	16	18
d (ft)	9	7	3	1	3	7	9	7	3	1

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • MAIN • Q2/5

A piston's position p (cm) is graphed for $0 \le t \le 3$ seconds. Give the amplitude, the period, and the intervals of the first cycle where p increases.

Answer: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • MAIN • Q3/5

The table shows hours of daylight D in month m (m = 0 is January). Give the maximum, the minimum, the period, and the months in which daylight increases.

m	0	1	2	3	4	5	6	7	8	9	10	11	12
D (h)	9	9.4	10.5	12	13.5	14.6	15	14.6	13.5	12	10.5	9.4	9

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • MAIN • Q4/5

The graph shows blood pressure P (mm Hg) over 2.4 seconds. Give the maximum and minimum pressures and the number of heartbeats per minute.

Answer: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • MAIN • Q5/5

SET 43 • HSF-IF.B.4

BACKWARD 1

Interpret maximums and minimums on a graph • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4

A graph of height peaks at 45 m. What does the peak mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • BACKWARD 1 • Q1/5

A graph is lowest at (6, 1). Interpret it if it shows depth (ft) against time (h).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • BACKWARD 1 • Q2/5

True or false?

The minimum value of a graph is its lowest y-value.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • BACKWARD 1 • Q3/5

A relative maximum on a graph is

A. the last point

B. any x-intercept

C. a point higher than nearby points

D. the y-intercept

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • BACKWARD 1 • Q4/5

A graph rises on $0 < x < 2$. What does this mean for a temperature graph?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • BACKWARD 1 • Q5/5

SET 43 • HSF-IF.B.4

BACKWARD 2

Read values from a table • Grade 6 • Expressions & Equations • 6.EE.C.9

In the depth table, what is d when t = 4?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 43 • BACKWARD 2 • Q1/5

In the depth table, at which listed times is d = 9?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 43 • BACKWARD 2 • Q2/5

True or false?

A repeated pattern in a table suggests a periodic relationship.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 43 • BACKWARD 2 • Q3/5

Daylight 15 h appears in the table at

A. m = 12

B. m = 6

C. m = 0

D. m = 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 43 • BACKWARD 2 • Q4/5

By how much does d change from t = 2 to t = 4?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.C.9 • Set 43 • BACKWARD 2 • Q5/5

SET 43 • HSF-IF.B.4

BACKWARD 3

Amplitude, period, and midline of a sinusoidal model • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

Give the amplitude and midline of $T = 15 - 8\cos(\pi t/12)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B3-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • BACKWARD 3 • Q1/5

Give the period of $p = 3\sin(2\pi t)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B3-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • BACKWARD 3 • Q2/5

True or false?

A wave between 80 and 120 has amplitude 20.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B3-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • BACKWARD 3 • Q3/5

A wave with maximum 9 and minimum 1 has midline

A. y = 8

B. y = 5

C. y = 10

D. y = 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B3-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • BACKWARD 3 • Q4/5

A period of 0.8 s means how many cycles per minute?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-B3-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • BACKWARD 3 • Q5/5

SET 43 • HSF-IF.B.4

FORWARD 1

Predict with periodicity • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

Use the temperature model $T(t) = 15 - 8\cos(\pi t/12)$ to find T at t = 100 hours.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • FORWARD 1 • Q1/5

Using the depth table, predict the depth at t = 30 hours.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • FORWARD 1 • Q2/5

True or false?

If f has period 12, then f(50) = f(2).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • FORWARD 1 • Q3/5

The piston has period 1 s. At t = 7.25 s, p is

A. 0

B. 3

C. −3

D. 1.5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • FORWARD 1 • Q4/5

Using the daylight table, in which month after m = 12 does daylight next reach 15 h?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 43 • FORWARD 1 • Q5/5

SET 43 • HSF-IF.B.4

FORWARD 2

Write a model from periodic features • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

Write a cosine model for the depth table.

$d =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q1/5

Write a model for the daylight table.

D = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q2/5

True or false?

The temperature graph is modeled by $15 - 8\cos(\pi t/12)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q3/5

The blood pressure graph is modeled by

A. $100 + 20\sin(0.8t)$

B. $20 + 100\sin(2.5\pi t)$

C. $120\sin(2.5\pi t)$

D. $100 + 20\sin(2.5\pi t)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q4/5

Write a model for the piston graph.

p = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S43-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q5/5

SET 44 • HSF-IF.B.5 • INTERPRETING FUNCTIONS

State domains of radical, rational, logarithmic, and polynomial models

MAIN	Algebra II main question	**HSF-IF.B.5**
BACKWARD 1	Domains of linear, quadratic, and exponential models	**HSF-IF.B.5**
BACKWARD 2	Division by zero	**7.NS.A.2.b**
BACKWARD 3	Solve linear inequalities	**HSA-REI.B.3**
FORWARD 1	Domains defined by nonlinear conditions	**HSF-IF.B.5**
FORWARD 2	Domains and ranges of inverse functions	**HSF-BF.B.4.a**

SET 44 • HSF-IF.B.5

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5

A car's speed is $s(d) = \sqrt{(22.5d)}$ mph for a skid of d feet. Give the domain of the function and the domain that makes sense in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q1/5

The average cost of x shirts is A(x) = (500 + 4x)/x. Give the domain of the function and the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q2/5

$pH = -\log[H^+]$, where $[H^+]$ is the hydrogen ion concentration. What is the domain?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q3/5

A box from a 12-in sheet has volume $V(x) = x(12 - 2x)^2$. Give the domain of the polynomial and the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q4/5

Give the domain of $f(x) = \sqrt{(x - 2)/(x - 5)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q5/5

SET 44 • HSF-IF.B.5

BACKWARD 1

Domains of linear, quadratic, and exponential models • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

$h(t) = -16t^2 + 64t$ models height until the ball lands. Give the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q1/5

C(n) = 2n + 5 is the cost of n tickets. Give the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q2/5

True or false?

The domain of $f(x) = 2^x$ is all real numbers.

A. True

B. False

A model for the number of people uses a domain of

A. fractions only

B. all real numbers

C. whole numbers

D. negative numbers

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q4/5

Why might a model's domain be smaller than the function's domain?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q5/5

SET 44 • HSF-IF.B.5

BACKWARD 2

Division by zero • Grade 7 • The Number System • 7.NS.A.2.b

For which x is 1/(x − 5) undefined?

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B2-Q1

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q1/5

For which x is (500 + 4x)/x undefined?

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B2-Q2

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q2/5

True or false?

$3/0 = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B2-Q3

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q3/5

$(x + 1)/(x^2 - 9)$ is undefined at

A. $x = 0$

B. $x = 9$

C. $x = -1$

D. $x = \pm 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B2-Q4

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q4/5

Is 0/5 defined? Give its value.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B2-Q5

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q5/5

SET 44 • HSF-IF.B.5

BACKWARD 3

Solve linear inequalities • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3

Solve $x - 2 \geq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q1/5

Solve $12 - 2x > 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q2/5

True or false?

$22.5d \geq 0$ means $d \geq 0$.

A. True

B. False

$3 - x \geq 0$ means

A. $x \geq 3$

B. $x \leq 3$

C. $x \geq -3$

D. $x \leq -3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q4/5

Solve $2x + 8 > 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q5/5

SET 44 • HSF-IF.B.5

FORWARD 1

Domains defined by nonlinear conditions • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5

Give the domain of $f(x) = \sqrt{(9 - x^2)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q1/5

Give the domain of $g(x) = \log(x^2 - 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q2/5

Give the domain of $h(x) = 1/(x^2 - x - 6)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q3/5

True or false?

The domain of $\sqrt{(x^2)}$ is all real numbers.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q4/5

The domain of $\log(x - 1) + \sqrt{(5 - x)}$ is

A. $1 < x \leq 5$

B. $x \leq 5$

C. $1 \leq x \leq 5$

D. $x > 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q5/5

SET 44 • HSF-IF.B.5

FORWARD 2

Domains and ranges of inverse functions • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

$f(x) = \sqrt{(x - 2)}$ has domain $x \geq 2$ and range $y \geq 0$. Write f^{-1} and give its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q1/5

$f(x) = 2^x$ has range $y > 0$. What is the domain of $f^{-1}(x) = \log_2 x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q2/5

True or false?

The domain of an inverse function is the range of the original function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q3/5

$f(x) = 1/x + 3$ $(x \neq 0)$ has an inverse with domain

A. $x \neq 0$

B. $x > 3$

C. all real x

D. $x \neq 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q4/5

Explain why the inverse of $s(d) = \sqrt{(22.5d)}$ needs the restriction $s \geq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S44-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q5/5

SET 45 • HSF-IF.B.6 • INTERPRETING FUNCTIONS

Calculate and interpret average rates of change of nonlinear models

MAIN	Algebra II main question	**HSF-IF.B.6**
BACKWARD 1	Average rate of change of simpler functions	**HSF-IF.B.6**
BACKWARD 2	Slope between two points	**8.F.B.4**
BACKWARD 3	Evaluate nonlinear functions	**HSF-IF.A.2**
FORWARD 1	Compare average rates on different intervals	**HSF-IF.B.6**
FORWARD 2	Compare rates of change across representations	**HSF-IF.C.9**

SET 45 • HSF-IF.B.6

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6

A culture has $N(t) = 200 \cdot 2^t$ bacteria after t hours. Find the average rate of change from t = 1 to t = 3 and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q1/5

A stone's height is $h(t) = 64 - 16t^2$ ft. Find the average rate of change from $t = 0.5$ to $t = 1.5$ s and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q2/5

The table shows a car's value. Find the average rate of change from t = 2 to t = 4 years and interpret it.

t (years)	0	1	2	3	4	5
V ($)	20000	17000	14450	12282.50	10440.06	8874.11

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q3/5

Speed from skid length is $s(d) = \sqrt{(24d)}$ mph. Find the average rate of change from $d = 24$ to $d = 96$ ft and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q4/5

Temperature is $T(t) = 15 - 8\cos(\pi t/12)$ °C after t hours. Find the average rate of change from $t = 0$ to $t = 6$ and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q5/5

SET 45 • HSF-IF.B.6

BACKWARD 1

Average rate of change of simpler functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6

$f(x) = x^2$. Find the average rate of change from $x = 1$ to $x = 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q1/5

A table gives f(2) = 10 and f(5) = 1. Find the average rate of change.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q2/5

True or false?

The average rate of change from a to b is $(f(b) - f(a))/(b - a)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q3/5

The average rate of change of f(x) = 3x + 1 on any interval is

A. it varies

B. 1

C. 3

D. 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q4/5

What are the units of a rate if f is in dollars and x is in years?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q5/5

SET 45 • HSF-IF.B.6

BACKWARD 2

Slope between two points • Grade 8 • Functions • 8.F.B.4

Find the slope between (1, 400) and (3, 1600).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B2-Q1

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q1/5

Find the slope between (0.5, 60) and (1.5, 28).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B2-Q2

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q2/5

True or false?

A negative slope means the output decreases as the input increases.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B2-Q3

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q3/5

The slope between (24, 24) and (96, 48) is

A. 72/24

B. 1/3

C. 2

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B2-Q4

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q4/5

Find the slope between (0, 7) and (6, 15).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B2-Q5

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q5/5

SET 45 • HSF-IF.B.6

BACKWARD 3

Evaluate nonlinear functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Find $N(3)$ for $N(t) = 200 \cdot 2^t$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q1/5

Find $h(1.5)$ for $h(t) = 64 - 16t^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q2/5

True or false?

$\sqrt{(24 \cdot 96)} = 48.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q3/5

$15 - 8\cos(\pi/2)$ equals

A. 15

B. 7

C. 23

D. 8

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q4/5

Find $T(0)$ for $T(t) = 15 - 8\cos(\pi t/12)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q5/5

SET 45 • HSF-IF.B.6

FORWARD 1

Compare average rates on different intervals • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6

For $N(t) = 200 \cdot 2^t$, compare the average rates on [0, 1] and [3, 4]. What does the comparison show?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q1/5

For $s(d) = \sqrt{(24d)}$, compare the average rates on $[0, 24]$ and $[96, 150]$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q2/5

True or false?

For a function with period 24, the average rate of change over [0, 24] is 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q3/5

For $f(x) = x^3$, the average rate on $[-1, 1]$ is

A. 3

B. 0

C. 2

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S45-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q4/5

Find an interval of length 6 on which $T(t) = 15 - 8\cos(\pi t/12)$ has average rate of change $-4/3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q5/5

SET 45 • HSF-IF.B.6

FORWARD 2

Compare rates of change across representations • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9

$f(t) = 200 \cdot 2^t$, and g is given by $g(1) = 500$, $g(3) = 1500$. Which has the greater average rate on [1, 3]?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q1/5

A graph of a cooling liquid falls from 90°C at t = 0 to 50°C at t = 20 min. Compare with T(t) = 90 − 1.5t on the same interval.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q2/5

True or false?

A function with a greater average rate on one interval must be greater everywhere.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q3/5

A table shows $h(0) = 0$, $h(2) = 40$; the formula $k(t) = 5t^2$ gives

A. no rate for k

B. a greater rate for k

C. a greater rate for h

D. equal average rates on [0, 2]

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q4/5

Explain how to compare rates when one function is a graph and one is a formula.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q5/5

SET 46 • HSF-IF.C.7.b • INTERPRETING FUNCTIONS

Graph square root and cube root functions with transformations

MAIN	Algebra II main question	**HSF-IF.C.7.b**
BACKWARD 1	Graph $y = \sqrt{x}$ and $y = \sqrt[3]{x}$ from a table	**HSF-IF.C.7.b**
BACKWARD 2	Translations and reflections of graphs	**HSF-BF.B.3**
BACKWARD 3	Square and cube roots of numbers	**8.EE.A.2**
FORWARD 1	Graph radical functions with horizontal stretches and reflections	**HSF-IF.C.7.b**
FORWARD 2	Graph a radical function and its inverse	**HSF-BF.B.4.a**

SET 46 • HSF-IF.C.7.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

Graph the function on the grid. Show and label the starting point and two more points, and state the domain and range.

$$y = \sqrt{(x - 2)} + 1$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q1/5

Graph the function on the grid. Show and label the starting point and two more points, and state the domain and range.

$y = -\sqrt{(x + 3)}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q2/5

Graph the function on the grid. Show and label the starting point and two more points, and state the domain and range.

$y = 2\sqrt{x} - 3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q3/5

Graph the function on the grid. Show and label the center point and two points on each side, and state the domain and range.

$y = \sqrt[3]{(x + 1)}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q4/5

Graph the function on the grid. Show and label the center point and two more points, and state the domain and range.

$$y = -\sqrt[3]{(x - 2)} + 1$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q5/5

SET 46 • HSF-IF.C.7.b

BACKWARD 1

Graph y = √x and y = ∛x from a table • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b

Complete the table for $y = \sqrt{x}$ at $x = 0, 1, 4, 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q1/5

Complete the table for $y = \sqrt[3]{x}$ at $x = -8, -1, 0, 1, 8$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q2/5

True or false?

The graph of $y = \sqrt{x}$ starts at the origin and increases.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q3/5

The domain of $y = \sqrt[3]{x}$ is

A. $x > 0$

B. all real numbers

C. $x \leq 0$

D. $x \geq 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q4/5

Give the range of $y = \sqrt{x}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q5/5

SET 46 • HSF-IF.C.7.b

BACKWARD 2

Translations and reflections of graphs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

How is $y = \sqrt{(x - 2)} + 1$ obtained from $y = \sqrt{x}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q1/5

How is $y = -\sqrt{(x + 3)}$ obtained from $y = \sqrt{x}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q2/5

True or false?

$y = 2\sqrt{x}$ is a vertical stretch of $y = \sqrt{x}$ by 2.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q3/5

The point (0, 0) of $y = \sqrt{x}$ moves to which point for $y = \sqrt{(x - 2)} + 1$?

A. (2, 1)

B. (1, 2)

C. (−2, 1)

D. (2, −1)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q4/5

Where does (1, 1) of $y = \sqrt[3]{x}$ go under $y = \sqrt[3]{(x + 1)}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q5/5

SET 46 • HSF-IF.C.7.b

BACKWARD 3

Square and cube roots of numbers • Grade 8 • Expressions & Equations • 8.EE.A.2

Find $\sqrt{9}$ and $\sqrt[3]{(-8)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B3-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q1/5

Find $\sqrt[3]{27}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B3-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q2/5

True or false?

$\sqrt{(-4)}$ is a real number.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B3-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q3/5

$\sqrt[3]{(-1)}$ equals

A. i

B. undefined

C. -1

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B3-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q4/5

Find $\sqrt{16} - 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-B3-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q5/5

SET 46 • HSF-IF.C.7.b

FORWARD 1

Graph radical functions with horizontal stretches and reflections • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

Graph the function on the grid. Show and label the starting point and two more points, and state the domain.

$y = \sqrt{(-x)}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S46-F1-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q1/5

Show that $\sqrt{(4x)} = 2\sqrt{x}$ for $x \geq 0$ and explain what this says about the two graphs.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q2/5

True or false?

$y = \sqrt[3]{(-x)}$ is the same as $y = -\sqrt[3]{x}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q3/5

The domain of $y = \sqrt{(3 - x)}$ is

A. all reals

B. $x \geq 3$

C. $x \leq 3$

D. $x \geq -3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q4/5

Write a square root function whose graph starts at $(4, -2)$ and goes up to the left.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q5/5

SET 46 • HSF-IF.C.7.b

FORWARD 2

Graph a radical function and its inverse • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

Find the inverse of $f(x) = \sqrt{(x - 2)} + 1$ and give its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q1/5

Find the inverse of $g(x) = \sqrt[3]{(x + 1)}$.

$g^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q2/5

True or false?

The graph of an inverse is the reflection of the graph over y = x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q3/5

The point (6, 3) on $f(x) = \sqrt{(x - 2)} + 1$ gives which point on f^{-1}?

A. (6, 3)

B. (3, 6)

C. (−6, −3)

D. (3, −6)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q4/5

Explain why the inverse of $f(x) = \sqrt{(x - 2)} + 1$ needs the domain $x \geq 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S46-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q5/5

SET 47 • HSF-IF.C.7.b • INTERPRETING FUNCTIONS

Graph piecewise-defined and absolute value functions

MAIN	Algebra II main question	**HSF-IF.C.7.b**
BACKWARD 1	Evaluate and graph simple piecewise functions	**HSF-IF.C.7.b**
BACKWARD 2	Absolute value as distance	**6.NS.C.7**
BACKWARD 3	Open and closed endpoints	**6.EE.B.8**
FORWARD 1	Write piecewise functions from graphs and contexts	**HSF-IF.C.7.b**
FORWARD 2	Describe absolute value graphs as transformations, and identify…	**HSF-BF.B.3**

SET 47 • HSF-IF.C.7.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

Graph the function on the grid. Show and label each endpoint with an open or closed circle.

$f(x) = x + 3$ for $x < -1$, and

$f(x) = x^2$ for $x \geq -1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q1/5

Graph the function on the grid. Show and label the vertex and two more points.

$y = |2x - 4| - 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q2/5

Graph the function on the grid. Show and label each endpoint with an open or closed circle.

$f(x) = -2$ for $x \leq 1$, and

$f(x) = \sqrt{(x - 1)}$ for $x > 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q3/5

Graph the function on the grid. Show and label the vertex and two more points.

$y = -|x + 1| + 3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q4/5

Graph the function on the grid. Show and label each endpoint with an open or closed circle.

$f(x) = 2^x$ for $x < 1$, and

$f(x) = 4 - x$ for $x \geq 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q5/5

SET 47 • HSF-IF.C.7.b

BACKWARD 1

Evaluate and graph simple piecewise functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b

$f(x) = 2$ for $x < 0$ and $f(x) = x$ for $x \geq 0$. Find $f(-3)$ and $f(5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q1/5

For f above, find f(0).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q2/5

True or false?

A piecewise function uses different rules on different parts of its domain.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q3/5

For $f(x) = x + 3$ $(x < -1)$, x^2 $(x \geq -1)$, $f(-1)$ is

A. 3

B. 2

C. -1

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q4/5

For the same f, find f(−2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q5/5

SET 47 • HSF-IF.C.7.b

BACKWARD 2

Absolute value as distance • Grade 6 • The Number System • 6.NS.C.7

Find $|-5|$ and $|3|$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B2-Q1

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q1/5

Find $|2(1) - 4| - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B2-Q2

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q2/5

True or false?

$|x|$ is never negative.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B2-Q3

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q3/5

$|x + 1| = 0$ when

A. $x = 0$

B. $x = -1$

C. $x = 1$

D. never

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B2-Q4

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q4/5

Find $-|0 + 1| + 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B2-Q5

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q5/5

SET 47 • HSF-IF.C.7.b

BACKWARD 3

Open and closed endpoints • Grade 6 • Expressions & Equations • 6.EE.B.8

On a number line, is $x < -1$ drawn with an open or a closed circle at -1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q1/5

Is $x \geq 1$ drawn with an open or a closed circle at 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q2/5

True or false?

An open circle means the endpoint is included.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q3/5

$x \leq 1$ is shown by

A. an open circle and shading right

B. a closed circle at 1 and shading right

C. an open circle at 1 and shading left

D. a closed circle at 1 and shading left

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q4/5

Write the inequality for "all numbers greater than 1".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q5/5

SET 47 • HSF-IF.C.7.b

FORWARD 1

Write piecewise functions from graphs and contexts • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

A shipping company charges $5 for packages up to 2 lb and $5 plus $1.50 per pound over 2 lb. Write the cost C(w).

C(w) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q1/5

Write $|x - 3|$ as a piecewise function without absolute value bars.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q2/5

True or false?

A piecewise function can have a jump and still be a function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q3/5

Which piecewise function is continuous at $x = 1$?

A. 2^x for $x < 1$, $4 - x$ for $x \geq 1$

B. x^2 for $x < 1$, $2x - 1$ for $x \geq 1$

C. 2 for $x < 1$, 3 for $x \geq 1$

D. x for $x < 1$, $x + 1$ for $x \geq 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S47-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q4/5

Find k so that $f(x) = kx$ for $x < 2$ and $f(x) = x^2$ for $x \geq 2$ has no jump at 2.

$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q5/5

SET 47 • HSF-IF.C.7.b

FORWARD 2

Describe absolute value graphs as transformations, and identify symmetry • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Describe $y = |2x - 4| - 1$ as a transformation of $y = |x|$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q1/5

Write an absolute value function with vertex $(-1, 3)$ that opens down and passes through $(2, 0)$.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q2/5

True or false?

$y = |x|$ is an even function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q3/5

$y = |x - 2|$ is symmetric about

A. the origin

B. $y = 2$

C. $x = 0$

D. $x = 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S47-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q4/5

Is $y = -|x + 1| + 3$ an even function? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q5/5

SET 48 • HSF-IF.C.7.c • INTERPRETING FUNCTIONS

Graph polynomial functions showing zeros and end behavior

MAIN	Algebra II main question	**HSF-IF.C.7.c**
BACKWARD 1	Factor out common factors and special products	**HSA-SSE.A.2**
BACKWARD 2	Crossing and touching from multiplicity	**HSA-APR.B.3**
BACKWARD 3	Evaluate polynomials at test points	**HSF-IF.A.2**
FORWARD 1	Locate relative extrema of polynomial graphs	**HSF-IF.C.7.c**
FORWARD 2	Interpret polynomial graphs in context	**HSF-IF.B.4**

SET 48 • HSF-IF.C.7.c

MAIN

Algebra II main question • High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = x^3 - 4x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-M-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q1/5

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = -x^3 + 3x^2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-M-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q2/5

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = x^4 - 4x^2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-M-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q3/5

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = x^3 + x^2 - 6x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-M-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q4/5

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = -x^4 + 2x^3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-M-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q5/5

SET 48 • HSF-IF.C.7.c

BACKWARD 1

Factor out common factors and special products • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^3 - 4x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q1/5

Factor $-x^3 + 3x^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q2/5

True or false?

$$x^4 - 4x^2 = x^2(x^2 - 4) = x^2(x - 2)(x + 2).$$

A. True

B. False

$x^3 + x^2 - 6x$ equals

A. $(x + 3)(x - 2)$

B. $x(x + 3)(x - 2)$

C. $x(x - 3)(x + 2)$

D. $x(x + 6)(x - 1)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q4/5

Factor $-x^4 + 2x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q5/5

SET 48 • HSF-IF.C.7.c

BACKWARD 2

Crossing and touching from multiplicity • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

Does $y = x^2(x - 3)$ cross or touch the x-axis at 0?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q1/5

Does $y = x^3(x - 2)$ cross at 0?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q2/5

True or false?

A zero of odd multiplicity is a crossing point.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q3/5

The graph of $y = (x + 2)^2(x - 1)$ at $x = -2$

A. crosses

B. touches

C. has an asymptote

D. is undefined

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B2-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q4/5

Give the end behavior of $y = -x^4 + 2x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q5/5

SET 48 • HSF-IF.C.7.c

BACKWARD 3

Evaluate polynomials at test points • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Evaluate $x^3 - 4x$ at $x = -1$ and $x = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q1/5

Evaluate $-x^3 + 3x^2$ at $x = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q2/5

True or false?

$x^4 - 4x^2$ at $x = 1$ equals -3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q3/5

$x^3 + x^2 - 6x$ at $x = -2$ equals

A. 8

B. 0

C. -8

D. 16

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q4/5

Evaluate $-x^4 + 2x^3$ at $x = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q5/5

SET 48 • HSF-IF.C.7.c

FORWARD 1

Locate relative extrema of polynomial graphs • High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c

Use a table with steps of 0.1 (or technology) to estimate the relative maximum of $y = x^3 - 4x$ to the nearest tenth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F1-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q1/5

Use technology or a table to estimate the relative minimum of $y = x^3 + x^2 - 6x$ on $0 < x < 2$ to the nearest tenth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F1-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q2/5

True or false?

A polynomial of degree n has at most n − 1 turning points.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F1-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q3/5

How many turning points can $y = x^4 - 4x^2$ have?

A. 1

B. 4

C. 3

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F1-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q4/5

Explain why the graph of $y = x^3 - 4x$ has a turning point between 0 and 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F1-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q5/5

SET 48 • HSF-IF.C.7.c

FORWARD 2

Interpret polynomial graphs in context • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

A box's volume is $V(x) = x(12 - 2x)^2$. Which part of the polynomial graph is meaningful, and what is the maximum volume there?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q1/5

A company's profit (thousands of dollars) is $P(x) = -x^3 + 9x^2 - 15x$ for x hundred items, $0 \le x \le 8$. When is the profit positive?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q2/5

True or false?

A zero of a profit model is a break-even point.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q3/5

A population model $P(t) = -t^3 + 6t^2 + 15t + 100$ rises then falls. The relative maximum means

A. the largest population before decline

B. the starting population

C. the end behavior

D. the time the population is 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q4/5

Why does the end behavior of $V(x) = x(12 - 2x)^2$ not matter for the box?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S48-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q5/5

SET 49 • HSF-IF.C.7.e • INTERPRETING FUNCTIONS

Graph exponential functions with intercepts, asymptote, and end behavior

MAIN	Algebra II main question	**HSF-IF.C.7.e**
BACKWARD 1	Graph basic exponential functions	**HSF-IF.C.7.e**
BACKWARD 2	Zero and negative exponents	**8.EE.A.1**
BACKWARD 3	Vertical and horizontal shifts and reflections	**HSF-BF.B.3**
FORWARD 1	Compare and describe exponential graphs	**HSF-IF.C.7.e**
FORWARD 2	Graph the inverse of an exponential function	**HSF-BF.B.4.a**

SET 49 • HSF-IF.C.7.e

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Graph the function on the grid. Show and label the y-intercept, the x-intercept (as a decimal), and the asymptote (dashed).

$y = 2^x - 3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q1/5

Graph the function on the grid. Show and label the y-intercept and the asymptote (dashed), and describe the end behavior.

$y = 3(0.5)^x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q2/5

Graph the function on the grid. Show and label the y-intercept, the x-intercept, and the asymptote (dashed).

$y = -2^x + 4$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q3/5

Graph the function on the grid. Show and label the y-intercept and the asymptote (dashed).

$y = 2^{x+1}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q4/5

Graph the function on the grid. Show and label the y-intercept and the asymptote (dashed).

$y = (1/3)^x + 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q5/5

SET 49 • HSF-IF.C.7.e

BACKWARD 1

Graph basic exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e

Complete the table for $y = 2^x$ at $x = -2, -1, 0, 1, 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q1/5

What is the y-intercept of $y = 3(0.5)^x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q2/5

True or false?

$y = 2^x$ has the horizontal asymptote $y = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q3/5

$y = (0.5)^x$ is

A. linear

B. increasing

C. constant

D. decreasing

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q4/5

Does $y = 2^x$ have an x-intercept?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q5/5

SET 49 • HSF-IF.C.7.e

BACKWARD 2

Zero and negative exponents • Grade 8 • Expressions & Equations • 8.EE.A.1

Evaluate 2^{-2}.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q1/5

Evaluate $(1/3)^{-2}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q2/5

True or false?

$b^0 = 1$ for $b \neq 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q3/5

$3(0.5)^{-1}$ equals

A. 6

B. 3

C. -1.5

D. 1.5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q4/5

Evaluate $2^{-1} - 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q5/5

SET 49 • HSF-IF.C.7.e

BACKWARD 3

Vertical and horizontal shifts and reflections • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

How does $y = 2^x - 3$ relate to $y = 2^x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q1/5

How does $y = 2^{x+1}$ relate to $y = 2^{x}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q2/5

True or false?

A vertical shift moves the horizontal asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q3/5

$y = -2^x + 4$ has asymptote

A. $y = 4$

B. $y = 0$

C. $y = -4$

D. $x = 4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q4/5

Give the asymptote of $y = (1/3)^x + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q5/5

SET 49 • HSF-IF.C.7.e

FORWARD 1

Compare and describe exponential graphs • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Explain why the graphs of $y = 2^x$ and $y = (1/2)^x$ are reflections of each other over the y-axis.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q1/5

Show that $2^{x+1} = 2 \cdot 2^{x}$ and explain what this says about two transformations.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q2/5

True or false?

$y = 5 \cdot 3^x$ and $y = 3^x$ have the same asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q3/5

Which exponential graph passes through (0, 4) and (1, 2)?

A. $y = 2^x + 3$

B. $y = 4 \cdot 2^x$

C. $y = 0.5 \cdot 4^x$

D. $y = 4(0.5)^x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q4/5

Write an exponential function with asymptote $y = -1$ and y-intercept 2 that is increasing.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q5/5

SET 49 • HSF-IF.C.7.e

FORWARD 2

Graph the inverse of an exponential function • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

Write the inverse of $f(x) = 2^x$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q1/5

The point (3, 8) is on $y = 2^x$. Which point is on its inverse?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q2/5

True or false?

The inverse of $y = 2^x$ has the vertical asymptote $x = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q3/5

The inverse of $y = 2^x - 3$ is

A. $y = \log_2(x - 3)$

B. $y = \log_2 x - 3$

C. $y = 2^{x+3}$

D. $y = \log_2(x + 3)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q4/5

Give the domain and range of $y = \log_2 x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S49-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q5/5

SET 50 • HSF-IF.C.7.e • INTERPRETING FUNCTIONS

Graph logarithmic functions with intercept, asymptote, and end behavior

MAIN	Algebra II main question	**HSF-IF.C.7.e**
BACKWARD 1	Graph exponential functions	**HSF-IF.C.7.e**
BACKWARD 2	Find inverses by swapping inputs and outputs	**HSF-BF.B.4.a**
BACKWARD 3	Shift and reflect graphs	**HSF-BF.B.3**
FORWARD 1	Graph logarithms with reflections that change the domain	**HSF-IF.C.7.e**
FORWARD 2	Solve equations with logarithms	**HSF-LE.A.4**

SET 50 • HSF-IF.C.7.e

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_2 x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q1/5

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_2(x + 3)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q2/5

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_3 x - 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q3/5

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = -\log_2 x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q4/5

Graph the function on the grid. Show and label two points and the asymptote (dashed); state the domain and whether there is an x-intercept.

$$y = \log_2(x - 1) + 2$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q5/5

SET 50 • HSF-IF.C.7.e

BACKWARD 1

Graph exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e

Give three points on $y = 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q1/5

Give the asymptote of $y = 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q2/5

True or false?

$y = 3^x$ passes through (1, 3).

A. True

B. False

$y = 2^x$ passes through

A. (2, 1)

B. (−1, −2)

C. (1/2, −1)

D. (−1, 1/2)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q4/5

Give the range of $y = 3^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q5/5

SET 50 • HSF-IF.C.7.e

BACKWARD 2

Find inverses by swapping inputs and outputs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

(2, 4) is on $y = 2^x$. Which point is on its inverse?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q1/5

Write the inverse of $y = 3^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q2/5

True or false?

Swapping x and y in $y = 2^x$ gives $x = 2^y$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q3/5

Which point is on $y = \log_2 x$?

A. (3, 8)

B. (2, 4)

C. (8, 3)

D. (0, 1)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q4/5

What is $\log_2 16$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q5/5

SET 50 • HSF-IF.C.7.e

BACKWARD 3

Shift and reflect graphs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

How does $y = \log_2(x + 3)$ relate to $y = \log_2 x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q1/5

How does $y = -\log_2 x$ relate to $y = \log_2 x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q2/5

True or false?

A horizontal shift moves the vertical asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q3/5

$y = \log_2(x - 1) + 2$ has asymptote

A. $x = -1$

B. $y = 2$

C. $x = 1$

D. $x = 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q4/5

Where does (1, 0) of $y = \log_3 x$ move under $y = \log_3 x - 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q5/5

SET 50 • HSF-IF.C.7.e

FORWARD 1

Graph logarithms with reflections that change the domain • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_2(4 - x)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q1/5

Give the domain of $y = \log(x^2 - 1)$ and explain the graph has two branches.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q2/5

True or false?

$y = \log_2(2x)$ is $y = \log_2 x$ shifted up 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q3/5

The x-intercept of $y = \log_2(x + 3)$ is

A. -2

B. 1

C. -3

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S50-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q4/5

Find the x-intercept of $y = \log_2(x - 1) + 2$ exactly.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q5/5

SET 50 • HSF-IF.C.7.e

FORWARD 2

Solve equations with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $2^x = 12$ exactly and to the nearest hundredth. A calculator may be used.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q1/5

Solve $5 \cdot 10^{x} = 40$ exactly.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q2/5

True or false?

$\log_2 12$ is between 3 and 4.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q3/5

The solution of $3^x = 7$ is

A. log 7/log 3

B. log 3/log 7

C. 7/3

D. log 21

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S50-F2-Q4**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q4/5

Use the graph of $y = \log_2 x$ to estimate $\log_2 6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S50-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q5/5

SET 51 • HSF-IF.C.7.e • INTERPRETING FUNCTIONS

Graph trigonometric functions showing period, midline, and amplitude

MAIN	Algebra II main question	**HSF-IF.C.7.e**
BACKWARD 1	Unit-circle values at quarter turns	**HSF-TF.A.2**
BACKWARD 2	Read amplitude, period, and midline from a formula	**HSF-TF.B.5**
BACKWARD 3	Vertical stretches, shifts, and reflections	**HSF-BF.B.3**
FORWARD 1	Graph shifted sinusoids and the tangent function	**HSF-IF.C.7.e**
FORWARD 2	Horizontal stretches f(kx) of trigonometric graphs	**HSF-BF.B.3**

SET 51 • HSF-IF.C.7.e

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period for one cycle on $0 \leq x \leq 2\pi$.

$y = 2\sin x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q1/5

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \leq x \leq 2\pi$.

$y = \cos x + 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q2/5

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \leq x \leq 2\pi$.

$y = \sin(2x)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q3/5

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \leq x \leq 4\pi$.

$y = 3\cos(x/2)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q4/5

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \leq x \leq 2\pi$.

$y = -\sin x + 2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q5/5

SET 51 • HSF-IF.C.7.e

BACKWARD 1

Unit-circle values at quarter turns • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Give sin x at x = 0, π/2, π, 3π/2, 2π.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q1/5

Give cos x at x = 0, π/2, π, 3π/2, 2π.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q2/5

True or false?

$\sin(3\pi/2) = -1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q3/5

$\cos(\pi)$ equals

A. π

B. 1

C. -1

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q4/5

Give $\sin(\pi/4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q5/5

SET 51 • HSF-IF.C.7.e

BACKWARD 2

Read amplitude, period, and midline from a formula • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

Give the amplitude and period of $y = 3\cos(x/2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q1/5

Give the midline of $y = -\sin x + 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q2/5

True or false?

$y = \sin(2x)$ has period π.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q3/5

$y = 2\sin x$ has amplitude

A. 1

B. 2

C. π

D. 2π

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q4/5

Give the period of $y = \cos(3x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q5/5

SET 51 • HSF-IF.C.7.e

BACKWARD 3

Vertical stretches, shifts, and reflections • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

Describe $y = \cos x + 1$ compared with $y = \cos x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q1/5

Describe $y = -\sin x$ compared with $y = \sin x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q2/5

True or false?

$y = 2\sin x$ has the same zeros as $y = \sin x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q3/5

The maximum of $y = -\sin x + 2$ is

A. 1

B. 2

C. 4

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q4/5

What is the minimum of $y = \cos x + 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q5/5

SET 51 • HSF-IF.C.7.e

FORWARD 1

Graph shifted sinusoids and the tangent function • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Graph the function on the grid. Show and label the asymptotes (dashed) and three points.

$y = \tan x$ for $-\pi/2 < x < \pi/2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q1/5

Describe the graph of $y = \sin(x - \pi/2)$ compared with $y = \sin x$, and name a simpler formula for it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q2/5

True or false?

The period of $y = \tan x$ is π.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q3/5

The first maximum of $y = \cos(x - \pi/3)$ for $x \geq 0$ is at

A. $x = 2\pi/3$

B. $x = \pi/3$

C. $x = 0$

D. $x = -\pi/3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S51-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q4/5

Give the x-values of the maximum points of $y = \sin(x + \pi/4)$ on $0 \leq x < 2\pi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q5/5

SET 51 • HSF-IF.C.7.e

FORWARD 2

Horizontal stretches f(kx) of trigonometric graphs • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

How is $y = \sin(2x)$ obtained from $y = \sin x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q1/5

How is $y = \cos(x/2)$ obtained from $y = \cos x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q2/5

True or false?

Replacing x by kx (k > 1) shortens the period by the factor k.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q3/5

$y = \sin(-x)$ is the same as

A. $\cos x$

B. $\sin x$

C. $-\cos x$

D. $-\sin x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S51-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q4/5

For which k does $y = \cos(kx)$ have period 6π?

$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S51-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q5/5

SET 52 • HSF-IF.C.8.b • INTERPRETING FUNCTIONS

Rewrite exponential functions to reveal rates over other time units

MAIN	Algebra II main question	**HSF-IF.C.8.b**
BACKWARD 1	Read growth rates from exponential functions	**HSF-IF.C.8.b**
BACKWARD 2	Powers of powers and rational exponents	**HSN-RN.A.1**
BACKWARD 3	Percent change and growth factors	**7.RP.A.3**
FORWARD 1	Compare growth offers by rewriting to the same time unit	**HSF-IF.C.8.b**
FORWARD 2	Find doubling times and half-lives with logarithms	**HSF-LE.A.4**

SET 52 • HSF-IF.C.8.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

$y = 100(1.08)^t$ with t in years. Rewrite it with a monthly growth factor and give the monthly percent rate to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q1/5

$y = 500(0.97)^t$ with t in years. Rewrite it per decade and give the percent decrease per decade to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q2/5

$Q = 50(2)^{t/3}$ with t in hours. Rewrite it with an hourly growth factor and give the hourly percent rate to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q3/5

$A = 1000(1.005)^{12t}$ with t in years. Rewrite it with an annual growth factor and give the annual percent rate to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q4/5

$N = 800(0.5)^{t/10}$ with t in years. Rewrite it with a yearly factor and give the yearly percent decrease to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q5/5

SET 52 • HSF-IF.C.8.b

BACKWARD 1

Read growth rates from exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b

Is $y = 100(1.08)^t$ growth or decay, and at what percent per year?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q1/5

Is $y = 500(0.97)^t$ growth or decay, and at what percent per year?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q2/5

True or false?

$y = 2(1.25)^t$ grows 25% per unit of t.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q3/5

$y = 40(0.9)^t$ decreases by

A. 9% per unit

B. 10% per unit

C. 90% per unit

D. 0.9% per unit

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q4/5

What is the growth factor for 6.17% growth?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q5/5

SET 52 • HSF-IF.C.8.b

BACKWARD 2

Powers of powers and rational exponents • High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1

Rewrite 1.08^{t} as a power of $1.08^{1/12}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q1

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q1/5

Rewrite $2^{t/3}$ as $(2^{1/3})^{t}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q2

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q2/5

True or false?

$(b^m)^n = b^{mn}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q3

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q3/5

1.005^{12t} equals

A. $(1.005^{12})^{t}$

B. $(1.005^{t})/12$

C. $12 \cdot 1.005^{t}$

D. $1.005^{12} + t$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q4

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q4/5

Write $8^{1/3}$ as a whole number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B2-Q5

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q5/5

SET 52 • HSF-IF.C.8.b

BACKWARD 3

Percent change and growth factors • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

Write the growth factor for a 25.99% increase.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B3-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q1/5

A factor 0.7374 means what percent decrease?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B3-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q2/5

True or false?

A factor 1.006434 means about 0.64% growth.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B3-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q3/5

A 3% decrease corresponds to the factor

A. 0.03

B. 0.97

C. 0.7

D. 1.03

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B3-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q4/5

Write the percent change for a factor of 0.933.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-B3-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q5/5

SET 52 • HSF-IF.C.8.b

FORWARD 1

Compare growth offers by rewriting to the same time unit • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

Account A grows 0.5% per month. Account B grows 6.1% per year. Which grows faster? Rewrite to compare. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q1/5

A culture doubles every 3 hours; another triples every 5 hours. Which grows faster per hour? A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q2/5

True or false?

Growing 1% per month is the same as growing 12% per year.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q3/5

Which is the greatest annual growth?

A. 4% per half year

B. 0.6% per month

C. 8% per year

D. 2% per quarter

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S52-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q4/5

A half-life of 10 years and a decay of 7% per year: which decays faster? A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q5/5

SET 52 • HSF-IF.C.8.b

FORWARD 2

Find doubling times and half-lives with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Find the doubling time of $y = 100(1.08)^t$, to the nearest hundredth of a year. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q1/5

Find the half-life of $y = 500(0.97)^t$ to the nearest hundredth of a year. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q2/5

True or false?

The doubling time of $y = 50(2)^{t/3}$ is 3 hours.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q3/5

The doubling time of a quantity growing 6.17% per year is about

A. 16.2 years

B. 11.6 years

C. 6.17 years

D. 3.1 years

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q4/5

Explain why rewriting $y = 100(1.08)^t$ as $100 \cdot 2^{t/9.01}$ shows the doubling time.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S52-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q5/5

SET 53 • HSF-IF.C.9 • INTERPRETING FUNCTIONS

Compare properties of Algebra II functions in different representations

MAIN	Algebra II main question	**HSF-IF.C.9**
BACKWARD 1	Compare linear, quadratic, and exponential functions	**HSF-IF.C.9**
BACKWARD 2	Recognize a pattern in a table	**8.F.A.3**
BACKWARD 3	Domains, asymptotes, and periods of Algebra II families	**HSF-IF.B.5**
FORWARD 1	Compare rates of change and long-run behavior across represent…	**HSF-IF.C.9**
FORWARD 2	Build a difference function to compare	**HSF-BF.A.1.b**

SET 53 • HSF-IF.C.9

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9

$f(x) = 3(1.1)^x$. The table shows g. Which has the greater y-intercept, and which is greater at x = 20? A calculator may be used.

x	0	1	2	3	4
g(x)	10	15	20	25	30

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q1/5

$f(x) = \sqrt{(x - 1)}$. The graph of g is shown. Compare their domains.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q2/5

$h(t) = 12 + 10\sin(\pi t/15)$ gives one Ferris wheel's height (m, t in s). A second wheel has radius 8 m, center 10 m high, and turns once every 40 s. Compare the maximum heights and the periods.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q3/5

$f(x) = \log_2 x$. The table shows g. Which is greater at $x = 9$?

x	1	3	9	27
g(x)	0	1	2	3

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q4/5

$f(x) = (2x + 1)/(x - 3)$. The graph of g shows the horizontal asymptote $y = 1$ and the vertical asymptote $x = -2$. Compare the asymptotes.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q5/5

SET 53 • HSF-IF.C.9

BACKWARD 1

Compare linear, quadratic, and exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9

f(x) = 2x + 1 and g has table g(0) = 3, g(1) = 4. Which has the greater y-intercept?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q1/5

$f(x) = x^2$ and $g(x) = 2^x$. Which is greater at $x = 10$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q2/5

True or false?

An increasing exponential eventually exceeds any linear function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q3/5

$f(x) = -(x - 1)^2 + 5$ and g with maximum 4 in a graph. The greater maximum is

A. f: 5

B. cannot tell

C. equal

D. g: 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q4/5

A table rises by 5 each step. What kind of function is it?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q5/5

SET 53 • HSF-IF.C.9

BACKWARD 2

Recognize a pattern in a table • Grade 8 • Functions • 8.F.A.3

Is the table x: 0, 1, 2; y: 10, 15, 20 linear? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q1

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q1/5

Is the table x: 1, 3, 9, 27; y: 0, 1, 2, 3 linear? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q2

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q2/5

True or false?

A table with constant ratios of outputs is exponential.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q3

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q3/5

Which table is exponential?

A. 1, 4, 9, 16

B. 1, 2, 4, 8

C. 2, 2, 2, 2

D. 1, 3, 5, 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q4

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q4/5

What is the slope of a linear table with outputs 10, 15, 20 at x = 0, 1, 2?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q5

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q5/5

SET 53 • HSF-IF.C.9

BACKWARD 3

Domains, asymptotes, and periods of Algebra II families • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

Give the domain of $\sqrt{(x - 1)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q1/5

Give the asymptotes of $(2x + 1)/(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q2/5

True or false?

The period of $12 + 10\sin(\pi t/15)$ is 30.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q3/5

The domain of $\log_2 x$ is

A. all reals

B. $x \geq 0$

C. $x > 0$

D. $x \neq 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q4/5

A wheel of radius 8 m with center 10 m high has what maximum height?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q5/5

SET 53 • HSF-IF.C.9

FORWARD 1

Compare rates of change and long-run behavior across representations • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9

$f(x) = 3(1.1)^x$ and g has the linear table above. Estimate the first whole number x at which f exceeds g. A calculator may be used.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q1/5

Compare the average rates of change of $f(x) = \log_2 x$ and the tabled g on $1 \le x \le 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q2/5

True or false?

A function with a larger y-intercept is always larger.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q3/5

Which grows fastest for large x?

A. $\sqrt{(x - 1)}$

B. $3(1.1)^x$

C. $5x + 10$

D. $\log_2 x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q4/5

Explain why the logarithm in the table, $g = \log_3 x$, stays below $f = \log_2 x$ for $x > 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q5/5

SET 53 • HSF-IF.C.9

FORWARD 2

Build a difference function to compare • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b

Write $d(x) = f(x) - g(x)$ for $f(x) = 3(1.1)^x$ and $g(x) = 5x + 10$.

$d(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q1/5

Explain what the sign of d(x) tells you.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q2/5

True or false?

A zero of $d(x) = f(x) - g(x)$ is an x where $f(x) = g(x)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q3/5

For $h_1(t) = 12 + 10\sin(\pi t/15)$ and $h_2(t) = 10 + 8\sin(\pi t/20)$, $h_1(0) - h_2(0)$ is

A. 2

B. 22

C. -2

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q4/5

Write the difference of the two Ferris wheel heights as a function of t.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q5/5

Build models by combining Algebra II function types

MAIN	Algebra II main question	**HSF-BF.A.1.b**
BACKWARD 1	Combine linear and quadratic functions	**HSF-BF.A.1.b**
BACKWARD 2	Multiply and subtract polynomials	**HSA-APR.A.1**
BACKWARD 3	Evaluate exponential and trigonometric expressions	**HSF-IF.A.2**
FORWARD 1	Interpret the parts of a combined model	**HSF-BF.A.1.b**
FORWARD 2	Compose functions in context	**HSF-BF.A.1.c**

SET 54 • HSF-BF.A.1.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b

Revenue is $R(x) = x(50 - 0.5x)$ and cost is $C(x) = 200 + 10x$ dollars for x items. Write the profit $P(x)$ in standard form and find $P(30)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q1/5

The cost of x posters is C(x) = 1200 + 3x dollars. Write the average cost per poster A(x) and find A(100).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q2/5

A town has $P(t) = 5000(1.02)^t$ people and $F(t) = 6000 + 100t$ units of food after t years. Write the food per person and find it at t = 10 to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q3/5

A spring's amplitude is 10 cm and shrinks by 10% each second, while it oscillates once per second. Write y(t) as an exponential times a cosine and find y(1).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q4/5

Coffee starts 70°C above a 20°C room, and the difference shrinks 5% each minute. Write its temperature T(t) as a sum of functions and find T(10) to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q5/5

SET 54 • HSF-BF.A.1.b

BACKWARD 1

Combine linear and quadratic functions • High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b

$R(x) = 8x$ and $C(x) = 3x + 50$. Write $P = R - C$.

$P(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q1/5

$f(x) = x + 2$ and $g(x) = x - 2$. Write $f \cdot g$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q2/5

True or false?

Profit is revenue minus cost.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q3/5

Average cost is

A. total cost times items

B. total cost divided by the number of items

C. cost minus revenue

D. revenue divided by cost

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q4/5

$f(x) = 2x$ and $g(x) = x^2 + 1$. Find $(f + g)(3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q5/5

SET 54 • HSF-BF.A.1.b

BACKWARD 2

Multiply and subtract polynomials • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Expand $x(50 - 0.5x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q1/5

Simplify $(50x - 0.5x^2) - (200 + 10x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q2/5

True or false?

$-(200 + 10x) = -200 - 10x.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q3/5

$(3x + 1) - (x - 4)$ equals

A. $2x + 5$

B. $4x - 3$

C. $2x - 3$

D. $2x + 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q4/5

Expand $2(x^2 - 3x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B2-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q5/5

SET 54 • HSF-BF.A.1.b

BACKWARD 3

Evaluate exponential and trigonometric expressions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Evaluate $70(0.95)^{10}$ to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q1/5

Evaluate $5000(1.02)^{10}$ to the nearest whole number. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q2/5

True or false?

$\cos(2\pi) = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q3/5

$10(0.9)^2$ equals

A. 9

B. 18

C. 81

D. 8.1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q4/5

Evaluate $\cos(2\pi \cdot 0.5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q5/5

SET 54 • HSF-BF.A.1.b

FORWARD 1

Interpret the parts of a combined model • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b

In $T(t) = 20 + 70(0.95)^t$, what do 20 and $70(0.95)^t$ represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q1/5

In $y(t) = 10(0.9)^t \cos(2\pi t)$, which factor controls the size of the swings and which controls their timing?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q2/5

True or false?

In $A(x) = (1200 + 3x)/x = 1200/x + 3$, the term $1200/x$ is the fixed cost spread over x posters.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q3/5

As t grows, $T(t) = 20 + 70(0.95)^t$ approaches

A. 20

B. 0

C. 90

D. 70

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q4/5

Explain why food per person F(t)/P(t) eventually decreases.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q5/5

SET 54 • HSF-BF.A.1.b

FORWARD 2

Compose functions in context • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c

A balloon's radius is $r(t) = 2t$ cm after t seconds and its volume is $V(r) = (4/3)\pi r^3$. Write $V(r(t))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q1/5

A store takes 20% off with d(p) = 0.8p and then adds a $5 fee with f(p) = p + 5. Write f(d(p)) and find it for p = 50.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q2/5

True or false?

f(g(x)) and g(f(x)) are always equal.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q3/5

If $f(x) = x^2$ and $g(x) = x + 1$, then $f(g(2)) =$

A. 4

B. 9

C. 3

D. 5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-F2-Q4**

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q4/5

Explain how composition differs from multiplying two functions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S54-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q5/5

Identify vertical shifts, vertical stretches, and horizontal shifts of Algebra II functions

MAIN	Algebra II main question	**HSF-BF.B.3**
BACKWARD 1	Shifts and stretches of linear and quadratic graphs	**HSF-BF.B.3**
BACKWARD 2	Translate points in the plane	**8.G.A.3**
BACKWARD 3	Key points of parent functions	**HSF-IF.C.7.e**
FORWARD 1	Write transformed functions from descriptions and order of transfo..	**HSF-BF.B.3**
FORWARD 2	Read sinusoid parameters as transformations	**HSF-TF.B.5**

SET 55 • HSF-BF.B.3

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

$f(x) = \sqrt{x}$. The graph of $g(x) = f(x - h) + k$ is shown. Find h and k.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q1/5

$f(x) = 2^x$. The graph of $g(x) = a \cdot f(x)$ is shown. Find a.



a = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q2/5

$f(x) = x^3$. The graph of $g(x) = f(x - h) + k$ is shown. Find h and k.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q3/5

$f(x) = |x|$. The graph of $g(x) = a \cdot f(x - h) + k$ is shown. Find a, h, and k.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q4/5

2016

$f(x) = \log_2 x$. The graph of $g(x) = f(x) + k$ is shown. Find k.



$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q5/5

2017

SET 55 • HSF-BF.B.3

BACKWARD 1

Shifts and stretches of linear and quadratic graphs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

How is $y = (x - 3)^2 + 1$ obtained from $y = x^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q1/5

How is $y = 2x^2$ obtained from $y = x^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q2/5

True or false?

$y = f(x + 4)$ is the graph of f shifted right 4.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q3/5

The vertex of $y = (x + 2)^2 - 5$ is

A. $(2, 5)$

B. $(2, -5)$

C. $(-2, -5)$

D. $(-2, 5)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q4/5

2022

Write $y = x^2$ shifted left 1 and down 4.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q5/5

SET 55 • HSF-BF.B.3

BACKWARD 2

Translate points in the plane • Grade 8 • Geometry • 8.G.A.3

Translate (0, 0) right 2 and up 1.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q1

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q1/5

Translate (1, 1) left 2 and up 3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q2

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q2/5

True or false?

Translating (0, 0) down 2 gives (0, −2).

A. True

B. False

$(x, y) \to (x - 2, y + 3)$ moves $(0, 0)$ to

A. $(-2, 3)$

B. $(-2, -3)$

C. $(2, 3)$

D. $(3, -2)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q4

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q4/5

Which translation takes (0, 0) to (1, −2)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B2-Q5

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q5/5

SET 55 • HSF-BF.B.3

BACKWARD 3

Key points of parent functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e

Give the point of $y = \sqrt{x}$ at $x = 0$ and at $x = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q1/5

Give the y-intercept of $y = 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q2/5

True or false?

The graph of $y = \log_2 x$ passes through (1, 0).

A. True

B. False

The vertex of $y = |x|$ is

A. (1, 0)

B. (1, 1)

C. (0, 1)

D. (0, 0)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q4/5

Give the center (point of symmetry) of $y = x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q5/5

SET 55 • HSF-BF.B.3

FORWARD 1

Write transformed functions from descriptions and order of transformations • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Write g, the graph of $f(x) = 2^x$ stretched vertically by 3, then shifted down 4.

$g(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q1/5

Write g, the graph of $f(x) = \sqrt{x}$ shifted down 4, then stretched vertically by 3.

$g(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q2/5

True or false?

A vertical stretch then a shift can give a different graph than the same shift then the stretch.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q3/5

$g(x) = 2f(x) + 1$ moves the point $(3, 4)$ of f to

A. $(6, 5)$

B. $(4, 9)$

C. $(3, 9)$

D. $(3, 10)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S55-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q4/5

$g(x) = f(x + 2) - 3$ and f has asymptote $x = 0$. Where is g's vertical asymptote?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q5/5

SET 55 • HSF-BF.B.3

FORWARD 2

Read sinusoid parameters as transformations • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

Describe $y = 3\sin x + 2$ as transformations of $y = \sin x$ and give its maximum.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q1/5

Write the function for y = cos x stretched vertically by 4 and shifted down 1.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q2/5

True or false?

In $y = A \sin x + D$, D is the midline.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q3/5

$y = -2\cos x + 5$ has amplitude and midline

A. 5 and $y = 2$

B. -2 and $y = 5$

C. 2 and $y = -5$

D. 2 and $y = 5$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q4/5

A model is $y = A \sin x + D$ with maximum 9 and minimum 1. Find A ($A > 0$) and D.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S55-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q5/5

Horizontal stretches, compressions, and reflections f(kx)

MAIN	Algebra II main question	**HSF-BF.B.3**
BACKWARD 1	Reflections and dilations of points	**8.G.A.3**
BACKWARD 2	Solve kx = c for x	**6.EE.B.7**
BACKWARD 3	Period of y = sin(Bx)	**HSF-TF.B.5**
FORWARD 1	Combine horizontal stretches with shifts	**HSF-BF.B.3**
FORWARD 2	Graph exponential and trigonometric functions with horizontal cha…	**HSF-IF.C.7.e**

SET 56 • HSF-BF.B.3

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

$f(x) = \sqrt{x}$ passes through (4, 2). Describe the graph of $g(x) = \sqrt{(4x)}$ compared with f, and give the point of g with y = 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q1/5

$f(x) = \sin x$ has period 2π. Describe $g(x) = \sin(3x)$ compared with f and give its period.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q2/5

Describe the graph of $g(x) = 2^{-x}$ compared with $f(x) = 2^{x}$, and give the point of g that comes from (1, 2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q3/5

f passes through $(-2, 1)$, $(0, 3)$, and $(4, -1)$. List the corresponding points of $g(x) = f(2x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q4/5

f passes through (2, 5). Which point must be on the graph of $g(x) = f(-x/2)$? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q5/5

SET 56 • HSF-BF.B.3

BACKWARD 1

Reflections and dilations of points • Grade 8 • Geometry • 8.G.A.3

Reflect (1, 2) over the y-axis.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q1

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q1/5

Multiply the x-coordinate of $(4, -1)$ by $1/2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q2

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q2/5

True or false?

Reflecting over the y-axis changes the sign of the x-coordinate.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q3

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q3/5

(x, y) → (x/3, y) applied to (6, 2) gives

A. (18, 2)

B. (6, 2/3)

C. (2, 2)

D. (3, 2)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q4

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q4/5

Apply $(x, y) \to (-2x, y)$ to $(2, 5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B1-Q5

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q5/5

SET 56 • HSF-BF.B.3

BACKWARD 2

Solve kx = c for x • Grade 6 • Expressions & Equations • 6.EE.B.7

Solve 2x = 4.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q1/5

Solve $-x/2 = 2$.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q2/5

True or false?

If $4x = 4$, then $x = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q3/5

$3x = 2\pi$ gives $x =$

A. $2\pi/3$

B. $2\pi - 3$

C. $\pi/3$

D. 6π

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q4/5

Solve $-x = 1$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q5/5

SET 56 • HSF-BF.B.3

BACKWARD 3

Period of y = sin(Bx) • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

Give the period of $y = \sin(2x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B3-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q1/5

Give the period of $y = \cos(x/2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B3-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q2/5

True or false?

A larger B gives a shorter period.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B3-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q3/5

y = sin(4x) has period

A. 8π

B. 4π

C. π/4

D. π/2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B3-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q4/5

Which B gives period 6?

B = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-B3-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q5/5

SET 56 • HSF-BF.B.3

FORWARD 1

Combine horizontal stretches with shifts • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Rewrite $g(x) = f(2x - 6)$ in the form $f(k(x - h))$ and describe the transformation.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q1/5

f passes through (4, 1). Which point is on $g(x) = f(2x - 6)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q2/5

True or false?

$f(2x - 6)$ is $f(2x)$ shifted right 6.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q3/5

$y = \sqrt{(9x)}$ equals

A. $\sqrt{x} + 9$

B. $9\sqrt{x}$

C. $3\sqrt{x}$

D. $\sqrt{x}/3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q4/5

Write g, the graph of $f(x) = 2^x$ reflected over the y-axis and then shifted right 1.

$g(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q5/5

SET 56 • HSF-BF.B.3

FORWARD 2

Graph exponential and trigonometric functions with horizontal changes • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Give the y-intercept, asymptote, and two points of $y = 2^{-x} + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q1/5

Give the maxima of $y = \cos(2x)$ on $0 \leq x \leq 2\pi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q2/5

True or false?

$y = 3^{2x} = 9^{x}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q3/5

$y = \sin(x/2)$ completes one cycle on

A. $0 \le x \le \pi$

B. $0 \le x \le 2\pi$

C. $0 \le x \le \pi/2$

D. $0 \le x \le 4\pi$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S56-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q4/5

Explain why $y = 2^{2x}$ is a horizontal compression of $y = 2^{x}$ and also an exponential with base 4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S56-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q5/5

Recognize even and odd functions

MAIN	Algebra II main question	**HSF-BF.B.3**
BACKWARD 1	Signs of powers of negatives	**7.NS.A.2**
BACKWARD 2	Reflect points over the axes and the origin	**6.NS.C.6.b**
BACKWARD 3	Substitute -x into an expression	**HSF-IF.A.2**
FORWARD 1	Reason about sums and products of even and odd functions	**HSF-BF.B.3**
FORWARD 2	Explain trigonometric symmetry with the unit circle	**HSF-TF.A.4**

SET 57 • HSF-BF.B.3

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Is $f(x) = x^4 - 3x^2$ even, odd, or neither? Show $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q1/5

Is $f(x) = x^3 - x$ even, odd, or neither? Show $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q2/5

Is $f(x) = x^2 + x$ even, odd, or neither? Justify.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q3/5

The graph of $f(x) = x/(x^2 + 1)$ is shown. Is f even, odd, or neither? Explain from the graph.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q4/5

Is $f(x) = \cos x + x^2$ even, odd, or neither? Show $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q5/5

SET 57 • HSF-BF.B.3

BACKWARD 1

Signs of powers of negatives • Grade 7 • The Number System • 7.NS.A.2

Simplify $(-x)^4$ and $(-x)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q1/5

Simplify $(-x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q2/5

True or false?

$(-x)^5 = -x^5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q3/5

$(-2)^2 - (-2)$ equals

A. -6

B. 2

C. 6

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q4/5

Simplify $-(-x)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B1-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q5/5

SET 57 • HSF-BF.B.3

BACKWARD 2

Reflect points over the axes and the origin • Grade 6 • The Number System • 6.NS.C.6.b

Reflect (2, 3) over the y-axis.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q1

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q1/5

Rotate (2, 3) 180° about the origin.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q2

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q2/5

True or false?

A graph symmetric about the y-axis contains (−a, b) whenever it contains (a, b).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q3

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q3/5

Symmetry about the origin pairs (a, b) with

A. $(-a, b)$

B. $(-a, -b)$

C. $(a, -b)$

D. (b, a)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q4

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q4/5

A graph is symmetric about the origin and contains (1, 0.5). Name another point.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B2-Q5

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q5/5

SET 57 • HSF-BF.B.3

BACKWARD 3

Substitute −x into an expression • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

For $f(x) = x^2 + x$, find $f(-x)$.

Answer: _____

For $f(x) = x^3 - x$, find $f(-2)$ and $f(2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q2/5

True or false?

For $f(x) = 3x$, $f(-x) = -3x$.

A. True

B. False

For $f(x) = x^4 - 3x^2$, $f(-1)$ is

A. 2

B. 4

C. -2

D. -4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q4/5

For $f(x) = 1/x$, find $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q5/5

SET 57 • HSF-BF.B.3

FORWARD 1

Reason about sums and products of even and odd functions • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Prove that the sum of two odd functions is odd.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q1/5

Prove that the product of two odd functions is even.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q2/5

True or false?

The only function that is both even and odd is $f(x) = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q3/5

$f(x) = |x| + x^3$ is

A. even

B. both

C. neither

D. odd

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S57-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q4/5

Is f(x) = x sin x even, odd, or neither? Explain with the product rule above.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q5/5

SET 57 • HSF-BF.B.3

FORWARD 2

Explain trigonometric symmetry with the unit circle • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4

Use the unit circle to explain why $\cos(-x) = \cos x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q1/5

Is tan even, odd, or neither? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q2/5

True or false?

sin is an even function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q3/5

Which is an even function?

A. cos x

B. tan x

C. sin x + cos x

D. sin x

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q4/5

Use symmetry to find $\sin(-\pi/6)$ from $\sin(\pi/6) = 1/2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S57-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q5/5

Write inverses of radical, rational, cubic, exponential, and logarithmic functions

MAIN	Algebra II main question	**HSF-BF.B.4.a**
BACKWARD 1	Inverses of linear functions	**HSF-BF.B.4.a**
BACKWARD 2	Rearrange formulas with roots, powers, and fractions	**HSA-CED.A.4**
BACKWARD 3	Domain and range	**HSF-IF.B.5**
FORWARD 1	Inverses that are their own inverse, and checking by points	**HSF-BF.B.4.a**
FORWARD 2	Verify inverses by composition	**HSF-BF.B.4.b**

SET 58 • HSF-BF.B.4.a

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

Find the inverse of $f(x) = (x - 1)^3 + 2$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q1/5

Find the inverse of $f(x) = 2x/(x - 3)$, $x \neq 3$, and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q2/5

Find the inverse of $f(x) = \sqrt{(x + 4)}$, $x \geq -4$, and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q3/5

Find the inverse of $f(x) = 5 \cdot 2^x$ and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q4/5

Find the inverse of $f(x) = \log_3(x - 1)$ and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q5/5

SET 58 • HSF-BF.B.4.a

BACKWARD 1

Inverses of linear functions • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

Find the inverse of $f(x) = 3x - 6$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q1/5

Solve 2x + 5 = y for x.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q2/5

True or false?

The inverse undoes the function: $f^{-1}(f(x)) = x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q3/5

The inverse of $f(x) = x/4 + 1$ is

A. $4(x - 1)$

B. $(x - 1)/4$

C. $4x - 1$

D. $x/4 - 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q4/5

If $f(2) = 7$, what is $f^{-1}(7)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q5/5

SET 58 • HSF-BF.B.4.a

BACKWARD 2

Rearrange formulas with roots, powers, and fractions • High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4

Solve $y = \sqrt{(x + 4)}$ for x ($y \geq 0$).

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q1

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q1/5

Solve $y = 2x/(x - 3)$ for x.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q2

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q2/5

True or false?

Solving $y = (x - 1)^3 + 2$ for x gives $x = \sqrt[3]{(y - 2)} + 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q3

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q3/5

Solving $y = x^3$ for x gives

A. $y/3$

B. $\sqrt[3]{y}$

C. $\sqrt{y}$

D. y^3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q4

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q4/5

Solve $y = x/(x + 1)$ for x.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B2-Q5

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q5/5

SET 58 • HSF-BF.B.4.a

BACKWARD 3

Domain and range • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

Give the range of $f(x) = \sqrt{(x + 4)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q1/5

Give the range of $f(x) = 5 \cdot 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q2/5

True or false?

The range of $f(x) = 2x/(x - 3)$ excludes $y = 2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q3/5

The domain of $\log_3(x - 1)$ is

A. $x > 1$

B. all reals

C. $x > 0$

D. $x \geq 1$

Give the range of $f(x) = \log_3(x - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q5/5

SET 58 • HSF-BF.B.4.a

FORWARD 1

Inverses that are their own inverse, and checking by points • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

Find the inverse of $f(x) = (x + 1)/(x - 1)$. What do you notice?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q1/5

$f(x) = 5 \cdot 2^x$ passes through (1, 10). Check your inverse with this point.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q2/5

True or false?

Every function has an inverse function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q3/5

Which is its own inverse?

A. $f(x) = x^2$

B. $f(x) = x + 1$

C. $f(x) = 2x$

D. $f(x) = 1/x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q4/5

Explain how the graph shows whether $f(x) = x^3 - x$ has an inverse function.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q5/5

SET 58 • HSF-BF.B.4.a

FORWARD 2

Verify inverses by composition • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b

Verify that $f(x) = (x - 1)^3 + 2$ and $g(x) = \sqrt[3]{(x - 2)} + 1$ are inverses by computing $f(g(x))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q1/5

Compute g(f(x)) for the same pair.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q2/5

True or false?

To verify inverses, both $f(g(x)) = x$ and $g(f(x)) = x$ must hold on the domains.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q3/5

For $f(x) = \sqrt{(x + 4)}$ and $g(x) = x^2 - 4$ $(x \geq 0)$, $g(f(x)) =$

A. x for all x

B. $|x|$

C. x for $x \geq -4$

D. $x^2 + 4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q4/5

Explain why $g(x) = x^2 - 4$ without a domain restriction is not the inverse of $f(x) = \sqrt{(x + 4)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S58-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q5/5

Write exact logarithmic solutions of exponential equations

MAIN	Algebra II main question	**HSF-LE.A.4**
BACKWARD 1	Isolate the exponential expression	**HSA-REI.B.3**
BACKWARD 2	Powers of 2, 3, and 10	**8.EE.A.1**
BACKWARD 3	Logarithms as the inverse of exponentials	**HSF-BF.B.4.a**
FORWARD 1	Solve equations with different bases or variables on both sides	**HSF-LE.A.4**
FORWARD 2	Use the inverse relationship to solve logarithmic equations	**HSF-BF.B.5**

SET 59 • HSF-LE.A.4

MAIN

Algebra II main question • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $3 \cdot 2^{5t} = 48$ exactly.

$t =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q1/5

Solve $2 \cdot 10^{3t} = 9$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q2/5

Solve $5e^{2t} = 30$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q3/5

Solve $100(1.05)^t = 300$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q4/5

Solve $4 \cdot 3^{t/2} = 20$ exactly.

$t =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-M-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q5/5

SET 59 • HSF-LE.A.4

BACKWARD 1

Isolate the exponential expression • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3

Solve 3y = 48 for y.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q1/5

Solve 5y = 30 for y.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q2/5

True or false?

From $2 \cdot 10^{3t} = 9$, we get $10^{3t} = 4.5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q3/5

From $4 \cdot 3^{t/2} = 20$, $3^{t/2}$ equals

A. 80

B. 5

C. 1/5

D. 16

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q4/5

Solve 100y = 300 for y.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q5/5

SET 59 • HSF-LE.A.4

BACKWARD 2

Powers of 2, 3, and 10 • Grade 8 • Expressions & Equations • 8.EE.A.1

Write 16 as a power of 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q1/5

Write 1000 as a power of 10.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q2/5

True or false?

$3^4 = 81$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q3/5

$2^{5t} = 2^4$ gives

A. t = 4

B. 5t = 4

C. 5t = 16

D. t = 20

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q4/5

Write 1/8 as a power of 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q5/5

SET 59 • HSF-LE.A.4

BACKWARD 3

Logarithms as the inverse of exponentials • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

Rewrite $10^x = 4.5$ as a logarithm.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q1/5

Rewrite $e^{u} = 6$ as a logarithm.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q2/5

True or false?

$\log_2 8 = 3$ because $2^3 = 8$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q3/5

$3^y = 5$ means y =

A. $\log_3 5$

B. 5/3

C. $\log_5 3$

D. 3/5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q4/5

Evaluate $\log_2 32$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q5/5

SET 59 • HSF-LE.A.4

FORWARD 1

Solve equations with different bases or variables on both sides • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $2^{x} = 3^{x-1}$ exactly.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q1/5

Solve $5^{2x} = 7$ exactly.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q2/5

True or false?

$\log(ab) = \log a + \log b$ for positive a and b.

A. True

B. False

The exact solution of $e^{x+1} = 10$ is

A. 10/e

B. ln 10/1

C. ln 9

D. ln 10 − 1

Solve $3 \cdot 4^x = 2 \cdot 5^x$ exactly.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q5/5

SET 59 • HSF-LE.A.4

FORWARD 2

Use the inverse relationship to solve logarithmic equations • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5

Solve $\log_2(x + 3) = 5$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q1/5

Solve $\log x + \log(x - 3) = 1$ and check for extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q2/5

True or false?

$\log_b(b^x) = x$ for every real x.

A. True

B. False

$\ln(e^7)$ equals

A. $\ln 7$

B. 1

C. 7

D. e^7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-F2-Q4**

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q4/5

Solve ln(2x) = 3 exactly.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S59-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q5/5

Evaluate logarithmic solutions of exponential models with technology

MAIN	Algebra II main question	**HSF-LE.A.4**
BACKWARD 1	Write exact logarithmic solutions	**HSF-LE.A.4**
BACKWARD 2	Evaluate logarithms with a calculator	**HSF-IF.A.2**
BACKWARD 3	Choose an appropriate level of accuracy	**HSN-Q.A.3**
FORWARD 1	Analyze how parameters change the solution	**HSF-LE.A.4**
FORWARD 2	Rewrite models to show doubling time or half-life	**HSF-IF.C.8.b**

SET 60 • HSF-LE.A.4

MAIN

Algebra II main question • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

$2,500 earns 3.5% interest compounded yearly. Solve $2500(1.035)^t = 4000$ and interpret the answer. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q1/5

Carbon-14 has a half-life of 5730 years. Solve $80(0.5)^{t/5730} = 30$ to the nearest year and interpret it. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q2/5

Coffee cools by $T(t) = 20 + 70(0.95)^t$ °C after t minutes. When is it 50°C? Give minutes to the nearest hundredth. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q3/5

A city of 1.2 million grows by $P(t) = 1.2e^{0.015t}$ million people. When does it reach 1.5 million? Give years to the nearest hundredth. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q4/5

A dose leaves $200(0.8)^t$ mg of a drug after t hours. After which whole hour is the amount first below 50 mg? A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-M-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q5/5

SET 60 • HSF-LE.A.4

BACKWARD 1

Write exact logarithmic solutions • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $1.035^{t} = 1.6$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q1/5

Solve $e^{0.015t} = 1.25$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q2/5

True or false?

$0.8^t = 0.25$ gives $t = \log 0.25 / \log 0.8$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q3/5

$(0.5)^{t/5730} = 0.375$ gives t =

A. $0.375 \cdot 5730$

B. 5730/log 0.375

C. log 0.375/5730

D. 5730 log 0.375/log 0.5

Solve $0.95^{t} = 3/7$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q5/5

SET 60 • HSF-LE.A.4

BACKWARD 2

Evaluate logarithms with a calculator • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Evaluate log 1.6 to four decimal places. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q1/5

Evaluate ln 1.25 to four decimal places. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q2/5

True or false?

log 0.25 is negative.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q3/5

log 1.6/log 1.035 is about

A. 0.20

B. 46.0

C. 1.55

D. 13.66

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q4/5

Evaluate log 0.25/log 0.8 to two decimal places. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q5/5

SET 60 • HSF-LE.A.4

BACKWARD 3

Choose an appropriate level of accuracy • High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3

A model gives 13.66 years for yearly interest. What whole-year answer fits the context?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B3-Q1

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q1/5

Round 8108.16 years to the nearest hundred years.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B3-Q2

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q2/5

True or false?

A time found from a model should be reported with more decimal places than the data justify.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B3-Q3

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q3/5

A drug amount first drops below 50 mg at t = 6.21 h. If it is checked every hour, it is first below 50 mg at

A. 6.21 h

B. 6 h

C. 7 h

D. 5 h

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B3-Q4

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q4/5

Why might 14.88 years be reported as "about 15 years"?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-B3-Q5

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q5/5

SET 60 • HSF-LE.A.4

FORWARD 1

Analyze how parameters change the solution • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

If the interest rate changes from 3.5% to 5%, how long does it take $2,500 to reach $4,000? Give years to the nearest hundredth. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q1/5

Find the doubling time at 6% yearly interest and compare it with the Rule of 72 estimate 72/6. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q2/5

True or false?

Doubling the initial amount in $a(b)^t = d$ halves the time t needed.

A. True

B. False

In $200(0.8)^t = 50$, replacing 200 by 400 changes t by

A. doubling it

B. halving it

C. adding $\log 2/\log(1/0.8) \approx 3.11$ hours

D. nothing

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q4/5

At 3.5% yearly interest, explain why $2,500 takes as long to grow to $4,000 as $5,000 takes to grow to $8,000.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q5/5

SET 60 • HSF-LE.A.4

FORWARD 2

Rewrite models to show doubling time or half-life • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

Rewrite $2500(1.035)^t$ in the form $2500 \cdot 2^{t/D}$ and give D to the nearest hundredth. A calculator may be used.

$D \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q1/5

Rewrite $200(0.8)^t$ in the form $200(0.5)^{t/H}$ and give H to the nearest hundredth. A calculator may be used.

$H \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q2/5

True or false?

$80(0.5)^{t/5730}$ shows the half-life 5730 directly.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q3/5

$1.2e^{0.015t}$ has doubling time about

A. 46.2 years

B. 66.7 years

C. 15 years

D. 0.015 years

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S60-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q4/5

Explain why the half-life form makes it easy to answer "how much remains after 3 half-lives".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S60-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q5/5

SET 61 • HSS-ID.A.4 • STATISTICS, INFERENCE, AND DECISIONS

Estimate population percentages with a normal distribution

MAIN	Algebra II main question	**HSS-ID.A.4**
BACKWARD 1	Interpret mean and standard deviation	**HSS-ID.A.2**
BACKWARD 2	Describe the shape of a distribution	**HSS-ID.A.1**
BACKWARD 3	Percents of a whole	**6.RP.A.3.c**
FORWARD 1	Find values from percentiles of a normal model	**HSS-ID.A.4**
FORWARD 2	Use a normal model for a margin of error	**HSS-IC.B.4**

SET 61 • HSS-ID.A.4

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

Adult women's heights are approximately normal with mean 64 in and standard deviation 2.5 in. Estimate the percent between 59 and 69 inches. Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q1/5

Using the same model, estimate the percent of women taller than 66.5 inches. Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q2/5

Test scores are approximately normal with mean 500 and standard deviation 100. Estimate the percent of scores between 400 and 700. Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q3/5

Battery life is approximately normal with mean 40 h and standard deviation 4 h. Use the table to estimate the percent of batteries lasting less than 46 hours.

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q4

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q4/5

Cereal box weights are approximately normal with mean 16.0 oz and standard deviation 0.2 oz. Use the table to estimate the percent between 15.8 and 16.3 oz.

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-M-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q5/5

SET 61 • HSS-ID.A.4

BACKWARD 1

Interpret mean and standard deviation • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2

A data set has mean 64 and standard deviation 2.5. What value is one standard deviation above the mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q1/5

Which value is two standard deviations below a mean of 500 with standard deviation 100?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q2/5

True or false?

A larger standard deviation means the data are more spread out.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q3/5

46 is how many standard deviations above a mean of 40 when the standard deviation is 4?

A. 0.67

B. 4

C. 6

D. 1.5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q4/5

Compute (15.8 − 16.0)/0.2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B1-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q5/5

SET 61 • HSS-ID.A.4

BACKWARD 2

Describe the shape of a distribution • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1

Describe a histogram that is highest in the middle and falls off evenly on both sides.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q1/5

Where are the mean and median of a symmetric distribution?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q2/5

True or false?

A normal curve is symmetric about its mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q3/5

A bell-shaped histogram is called

A. skewed right

B. uniform

C. mound-shaped and symmetric

D. bimodal

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q4/5

In a symmetric distribution, what percent of the data are below the median?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B2-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q5/5

SET 61 • HSS-ID.A.4

BACKWARD 3

Percents of a whole • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c

Find 95% of 2,000.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q1/5

What is 34% + 13.5%?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q2/5

True or false?

Half of 68% is 34%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q3/5

100% − 95% split equally into two tails gives each tail

A. 5%

B. 47.5%

C. 0.5%

D. 2.5%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q4/5

Write 0.7745 as a percent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q5/5

SET 61 • HSS-ID.A.4

FORWARD 1

Find values from percentiles of a normal model • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

Above what height are the tallest 2.5% of women (mean 64 in, SD 2.5 in)? Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q1/5

Use the table: what battery life is the 84th percentile (mean 40 h, SD 4 h)?

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q2/5

True or false?

In a normal model, the median equals the mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q3/5

The 16th percentile of scores with mean 500 and SD 100 is

A. 484

B. 600

C. 340

D. 400

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-F1-Q4**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q4/5

A cereal box below 15.7 oz is rejected. Use the table to estimate the percent rejected (mean 16.0, SD 0.2).

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F1-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q5/5

SET 61 • HSS-ID.A.4

FORWARD 2

Use a normal model for a margin of error • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

Simulated sample proportions are approximately normal with mean 0.58 and SD 0.049. Within what interval do about 95% of them fall?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q1/5

What margin of error does this give for the population proportion?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q2/5

True or false?

The margin of error is about two standard deviations of the simulated statistic.

A. True

B. False

If the SD of simulated means is 0.2 hours, the margin of error is about

A. 0.1 hours

B. 0.4 hours

C. 2 hours

D. 0.2 hours

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q4/5

Explain why the normal model is useful for simulated sample proportions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S61-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q5/5

SET 62 • HSS-ID.A.4 • STATISTICS, INFERENCE, AND DECISIONS

Recognize data sets for which a normal model is not appropriate

MAIN	Algebra II main question	**HSS-ID.A.4**
BACKWARD 1	Describe skewed, symmetric, bimodal, and uniform shapes	**HSS-ID.A.3**
BACKWARD 2	Measures of center and spread for a data set	**6.SP.A.2**
BACKWARD 3	Groups combined in one data set	**HSS-ID.A.1**
FORWARD 1	Check a normal model with the 68-95-99.7 rule	**HSS-ID.A.4**
FORWARD 2	Decide whether a probability model fits observed data	**HSS-IC.A.2**

SET 62 • HSS-ID.A.4

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

The histogram shows household incomes in a town. Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q1/5

The histogram shows heights of adults in a club. Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q2/5

The histogram shows heights of all people at a school event (children and adults). Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q3/5

Waiting times at a clinic have mean 4 minutes, standard deviation 5 minutes, and minimum 0. Is a normal model appropriate? Justify using the numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q4

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q4/5

The histogram shows the results of 600 rolls of a fair die. Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-M-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q5/5

SET 62 • HSS-ID.A.4

BACKWARD 1

Describe skewed, symmetric, bimodal, and uniform shapes • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3

A histogram has a long tail to the right. Name its shape.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q1/5

A histogram has two separate peaks. Name its shape.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q2/5

True or false?

In a right-skewed distribution, the mean is usually greater than the median.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q3/5

A histogram with bars all about the same height is

A. skewed left

B. uniform

C. bimodal

D. normal

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q4/5

Which is more resistant to a long tail: the mean or the median?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B1-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q5/5

SET 62 • HSS-ID.A.4

BACKWARD 2

Measures of center and spread for a data set • Grade 6 • Statistics & Probability • 6.SP.A.2

What does the standard deviation (or another spread measure) describe?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q1

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q1/5

Data are all at least 0 with mean 4. Can many values be far below 4?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q2

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q2/5

True or false?

A distribution can be described by its center, spread, and shape.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q3

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q3/5

Which is a measure of center?

A. median

B. range

C. standard deviation

D. IQR

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q4

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q4/5

Give one value two standard deviations below a mean of 4 with SD 5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B2-Q5

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q5/5

SET 62 • HSS-ID.A.4

BACKWARD 3

Groups combined in one data set • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1

Two groups with different centers are combined. What might the histogram show?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B3-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q1/5

Name a variable whose histogram for all students in a K-12 school might have several peaks.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B3-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q2/5

True or false?

A histogram of one group can look different from a histogram of two groups combined.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B3-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q3/5

A bimodal histogram suggests

A. an error in counting

B. a small standard deviation

C. two groups in the data

D. a normal model

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B3-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q4/5

How could you get a mound-shaped distribution from the school-event heights?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-B3-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q5/5

SET 62 • HSS-ID.A.4

FORWARD 1

Check a normal model with the 68-95-99.7 rule • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

In 1,000 measurements, 70% are within 1 SD of the mean and 96% within 2 SDs. Is a normal model reasonable? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q1/5

In another data set, 85% of values are within 1 SD and 91% within 2 SDs. Is a normal model reasonable?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q2/5

True or false?

If a data set has an extreme outlier, the normal model may still fit the rest but the mean and SD are distorted.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q3/5

About what percent of normal data lies more than 3 SDs from the mean?

A. 5%

B. 3%

C. 0.3%

D. 32%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F1-Q4**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q4/5

A data set of 200 values has 25 values more than 2 SDs from the mean. Compare with a normal model.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F1-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q5/5

SET 62 • HSS-ID.A.4

FORWARD 2

Decide whether a probability model fits observed data • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2

A spinner is claimed to land on each of 4 colors equally often. In 400 spins the counts are 98, 105, 97, 100. Are the results consistent with the claim?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q1/5

A model says a coin is fair. In 50 flips there were 34 heads. In 200 simulated runs of 50 fair flips, 3 had 34 or more heads. Is the model consistent with the data?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q2/5

True or false?

A model is rejected only if the observed result would be unusual under the model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q3/5

A simulated result at least as extreme as the observed one occurs in 34 of 200 runs. The model is

A. proved true

B. rejected

C. consistent with the data

D. impossible

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q4/5

Explain the difference between judging a normal model for a data set and judging a probability model with a simulation.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S62-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q5/5

SET 63 • HSS-IC.A.1 • STATISTICS, INFERENCE, AND DECISIONS

Distinguish population, sample, parameter, and statistic

MAIN	Algebra II main question	**HSS-IC.A.1**
BACKWARD 1	Samples and populations	**7.SP.A.1**
BACKWARD 2	Draw inferences from random samples	**7.SP.A.2**
BACKWARD 3	Compute sample proportions and means	**6.SP.B.5.c**
FORWARD 1	Describe sampling variability	**HSS-IC.A.1**
FORWARD 2	Quantify variability with a margin of error	**HSS-IC.B.4**

SET 63 • HSS-IC.A.1

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

A city surveys 400 randomly chosen residents, and 62% support a new park. Identify the population, the sample, the parameter, and the statistic.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q1/5

A factory measures 50 randomly selected bolts from a day's production; their mean length is 5.02 cm. Identify the parameter and the statistic.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q2/5

A news website posts a poll; 80% of the 2,000 visitors who answered want a later school start. Can this be generalized to all students? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q3/5

Explain why a random sample allows conclusions about the whole population.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q4/5

A school uses the first 30 students arriving in the morning to estimate the average hours of sleep. Is this a random sample? Explain a possible bias.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q5/5

SET 63 • HSS-IC.A.1

BACKWARD 1

Samples and populations • Grade 7 • Statistics & Probability • 7.SP.A.1

What is a population in statistics?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q1/5

What is a sample?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q2/5

True or false?

A random sample gives every member of the population a chance to be chosen.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q3/5

Surveying every student in a school is

A. a statistic

B. an experiment

C. a sample

D. a census

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q4/5

Why are samples used instead of the whole population?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q5/5

SET 63 • HSS-IC.A.1

BACKWARD 2

Draw inferences from random samples • Grade 7 • Statistics & Probability • 7.SP.A.2

In a random sample of 50 students, 20 walk to school. Estimate how many of 600 students walk.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q1/5

Two random samples give 40% and 44%. Is that surprising?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q2/5

True or false?

Different random samples from the same population usually give slightly different results.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q3/5

From a random sample, 30% of 200 voters prefer a candidate. The best estimate for all voters is

A. 30%

B. 200

C. cannot estimate

D. 60%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q4/5

Why is a larger random sample usually better?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q5/5

SET 63 • HSS-IC.A.1

BACKWARD 3

Compute sample proportions and means • Grade 6 • Statistics & Probability • 6.SP.B.5.c

248 of 400 people say yes. What proportion is that?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B3-Q1

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q1/5

Find the mean of 5.0, 5.1, 4.9, 5.08.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B3-Q2

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q2/5

True or false?

A proportion is a number between 0 and 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B3-Q3

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q3/5

1,600 of 2,000 is

A. 16%

B. 1.25%

C. 80%

D. 20%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B3-Q4

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q4/5

Write 62% as a decimal.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B3-Q5

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q5/5

SET 63 • HSS-IC.A.1

FORWARD 1

Describe sampling variability • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

Two random samples of 400 residents give 62% and 59%. Explain why they differ and what each estimates.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q1/5

Would samples of 40 or of 400 vary more from sample to sample? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q2/5

True or false?

A parameter changes from sample to sample.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q3/5

Which is a statistic?

A. the population size

B. the mean height of 30 sampled students

C. the mean height of all students

D. the population proportion

Explain why a large voluntary-response sample can still be worse than a small random sample.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q5/5

SET 63 • HSS-IC.A.1

FORWARD 2

Quantify variability with a margin of error • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

Simulated proportions from samples of 400 with $p = 0.62$ have SD 0.024. Give a margin of error.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q1/5

Use the margin of error to give an interval for the percent of residents who support the park.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q2/5

True or false?

A margin of error describes chance variation, not bias.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q3/5

If the sample size is multiplied by 4, the margin of error is about

A. unchanged

B. halved

C. doubled

D. divided by 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q4/5

Explain what the interval says about the population percent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q5/5

SET 64 • HSS-IC.A.2 • STATISTICS, INFERENCE, AND DECISIONS

Decide whether a model is consistent with observed results using a simulation

MAIN	Algebra II main question	**HSS-IC.A.2**
BACKWARD 1	Probability as long-run relative frequency	**7.SP.C.6**
BACKWARD 2	Use simulations of compound events	**7.SP.C.8.c**
BACKWARD 3	Relative frequency of a group of outcomes	**6.RP.A.3.c**
FORWARD 1	Design a simulation to test a model	**HSS-IC.A.2**
FORWARD 2	Compare two treatments with a re-randomization simulation	**HSS-IC.B.5**

SET 64 • HSS-IC.A.2

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2

A coin is claimed to be fair. It landed heads 34 times in 50 flips. The histogram shows 200 simulated runs of 50 fair flips. Is the claim consistent with the result? Justify with the simulation.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q1/5

A spinner is claimed to land on red 25% of the time. It landed on red 13 times in 40 spins. The histogram shows 200 simulated runs. Is the claim consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q2/5

A die is claimed to be fair. It showed 16 sixes in 60 rolls. The histogram shows 200 simulated runs. Is the claim consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q3/5

A player claims to make 70% of free throws. She made 11 of 20. The histogram shows 200 simulated runs of 20 shots. Is the claim consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q4/5

A model says boys and girls are equally likely. A hospital had 58 boys in 100 births. The histogram shows 200 simulated runs. Is the model consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q5/5

SET 64 • HSS-IC.A.2

BACKWARD 1

Probability as long-run relative frequency • Grade 7 • Statistics & Probability • 7.SP.C.6

A fair coin is flipped 50 times. About how many heads do you expect?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q1/5

A spinner has P(red) = 0.25. About how many reds in 40 spins?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q2/5

True or false?

Getting exactly the expected number every time is likely.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q3/5

A fair die is rolled 60 times. Expected sixes:

A. 10

B. 16

C. 6

D. 60

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q4/5

Why is 26 heads in 50 flips not surprising?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q5/5

SET 64 • HSS-IC.A.2

BACKWARD 2

Use simulations of compound events • Grade 7 • Statistics & Probability • 7.SP.C.8.c

Describe how to simulate 50 fair coin flips with random digits.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q1/5

How could you simulate one spin of a spinner with P(red) = 0.25 using two-digit random numbers 00-99?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q2/5

True or false?

Each run of a simulation should copy the real process under the model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q3/5

A simulation has 200 runs. Each run represents

A. the final answer

B. one repetition of the whole experiment

C. one histogram bar

D. one flip

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q4/5

Why are many runs used?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q5/5

SET 64 • HSS-IC.A.2

BACKWARD 3

Relative frequency of a group of outcomes • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c

3 of 200 runs is what percent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q1/5

6 + 18 runs out of 200 is what percent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q2/5

True or false?

19 out of 200 is 9.5%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q3/5

34 of 200 is

A. 1.7%

B. 17%

C. 6.8%

D. 34%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q4/5

Is 2% less than 5%?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q5/5

SET 64 • HSS-IC.A.2

FORWARD 1

Design a simulation to test a model • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2

Describe a simulation to test whether a basketball player really makes 70% of free throws, given that she made 11 of 20.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q1/5

Why should the simulation count results "as extreme or more extreme" rather than exactly 11?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q2/5

True or false?

A result that happens in 30% of simulated runs is strong evidence against the model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q3/5

A common cutoff for "unusual" is

A. about 50% of runs

B. 100% of runs

C. about 5% of runs

D. about 25% of runs

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q4/5

Doubling the number of runs from 200 to 400 changes what about the simulation?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q5/5

SET 64 • HSS-IC.A.2

FORWARD 2

Compare two treatments with a re-randomization simulation • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

In an experiment, fertilized plants grew 2.9 cm more on average. In 200 re-randomizations, none of the differences was 2.9 cm or more. What do you conclude?



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S64-F2-Q1**

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q1/5

What model does the re-randomization assume?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q2/5

True or false?

Re-randomization is a simulation of a model, like the coin-flip simulation.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q3/5

If 60 of 200 re-randomized differences were at least the observed one, the result is

A. significant

B. proof of no effect

C. not statistically significant

D. impossible

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q4/5

Explain how this extends the coin-flip reasoning.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S64-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q5/5

SET 65 • HSS-IC.B.3 • STATISTICS, INFERENCE, AND DECISIONS

Identify sample surveys, experiments, and observational studies

MAIN	Algebra II main question	**HSS-IC.B.3**
BACKWARD 1	Statistical questions and data collection	**6.SP.A.1**
BACKWARD 2	Association is not causation	**HSS-ID.C.9**
BACKWARD 3	Random samples	**7.SP.A.1**
FORWARD 1	Identify confounding and design flaws	**HSS-IC.B.3**
FORWARD 2	Analyze the results of an experiment	**HSS-IC.B.5**

SET 65 • HSS-IC.B.3

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

Researchers randomly assign 100 volunteers to take a vitamin or a placebo and compare colds. What type of study is this, and what is its purpose?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q1/5

A pollster asks a random sample of 1,000 voters which candidate they prefer. What type of study is this?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q2/5

Researchers compare the hospital records of smokers and nonsmokers for lung disease. What type of study is this? Can it show causation?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q3/5

Which type of study best answers "Does playing music while studying improve test scores?" Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q4/5

Which type of study best answers "What percent of students at the school use the library weekly?" Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q5/5

SET 65 • HSS-IC.B.3

BACKWARD 1

Statistical questions and data collection • Grade 6 • Statistics & Probability • 6.SP.A.1

Is "How many hours do students at our school sleep?" a statistical question? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q1

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q1/5

Give a data collection method for "What do voters think?"

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q2

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q2/5

True or false?

An experiment imposes a treatment on the subjects.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q3

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q3/5

Watching and recording without imposing treatments is

A. an experiment

B. an observational study

C. a census

D. a placebo

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q4

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q4/5

What is a placebo?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B1-Q5

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q5/5

SET 65 • HSS-IC.B.3

BACKWARD 2

Association is not causation • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9

Ice cream sales and sunburns are correlated. Does ice cream cause sunburn? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q1/5

Name a lurking variable for "students with more books at home get higher scores".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q2/5

True or false?

A strong correlation proves that one variable causes the other.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q3/5

A variable that affects both variables in a study is

A. the placebo

B. the sample

C. a confounding (lurking) variable

D. the response

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q4/5

What kind of study can show cause and effect?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B2-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q5/5

SET 65 • HSS-IC.B.3

BACKWARD 3

Random samples • Grade 7 • Statistics & Probability • 7.SP.A.1

Describe a way to choose a random sample of 50 students from a school list.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B3-Q1

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 65 • BACKWARD 3 • Q1/5

Is surveying students in the cafeteria line random?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B3-Q2

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 65 • BACKWARD 3 • Q2/5

True or false?

A random sample tends to represent the population.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B3-Q3

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 65 • BACKWARD 3 • Q3/5

Which sample is random?

A. friends of the teacher

B. students who volunteer

C. names drawn from a hat with every student's name

D. the first 50 arrivals

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B3-Q4

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 65 • BACKWARD 3 • Q4/5

Why does a survey need a random sample?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-B3-Q5

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 65 • BACKWARD 3 • Q5/5

SET 65 • HSS-IC.B.3

FORWARD 1

Identify confounding and design flaws • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

Students who choose to take an SAT prep course score higher. Give a confounding variable and explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • FORWARD 1 • Q1/5

Redesign the prep-course study so it can show causation.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • FORWARD 1 • Q2/5

True or false?

An experiment without a control group can still show the treatment caused the result.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • FORWARD 1 • Q3/5

In an experiment, why use a placebo?

A. to make the study observational

B. to control for the effect of believing one is treated

C. to increase the sample size

D. to make the treatment stronger

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S65-F1-Q4**

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • FORWARD 1 • Q4/5

Why might an observational study be chosen instead of an experiment for smoking?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • FORWARD 1 • Q5/5

SET 65 • HSS-IC.B.3

FORWARD 2

Analyze the results of an experiment • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

In the vitamin experiment, the vitamin group had 1.2 fewer colds on average. A re-randomization found 40 of 200 differences at least that large. What do you conclude?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 65 • FORWARD 2 • Q1/5

If instead 2 of 200 differences were at least that large, what would you conclude?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 65 • FORWARD 2 • Q2/5

True or false?

Random assignment makes it possible to attribute a significant difference to the treatment.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 65 • FORWARD 2 • Q3/5

The re-randomization model assumes

A. the groups are different sizes

B. the treatment has no effect

C. the treatment works

D. the data are normal

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 65 • FORWARD 2 • Q4/5

Explain why an observational study cannot be analyzed in the same causal way.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S65-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 65 • FORWARD 2 • Q5/5

SET 66 • HSS-IC.B.3 • STATISTICS, INFERENCE, AND DECISIONS

Explain random sampling versus random assignment

MAIN	Algebra II main question	**HSS-IC.B.3**
BACKWARD 1	Study types and their purposes	**HSS-IC.B.3**
BACKWARD 2	Choose a random sample	**7.SP.A.1**
BACKWARD 3	Correlation and causation	**HSS-ID.C.9**
FORWARD 1	Improve an experimental design	**HSS-IC.B.3**
FORWARD 2	Use re-randomization to judge a treatment difference	**HSS-IC.B.5**

SET 66 • HSS-IC.B.3

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

A random sample of 500 adults is surveyed about exercise and sleep; more exercise is associated with better sleep. What can and cannot be concluded?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • MAIN • Q1/5

60 volunteers are randomly assigned to two diets; diet A loses more weight. What can and cannot be concluded?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • MAIN • Q2/5

A random sample of 200 students is randomly assigned to two study methods. What can be concluded?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • MAIN • Q3/5

A teacher compares test scores of students who chose to use a review app with those who did not, in her own class. What can be concluded?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • MAIN • Q4/5

Describe how to randomly assign 40 subjects to two groups of 20.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • MAIN • Q5/5

SET 66 • HSS-IC.B.3

BACKWARD 1

Study types and their purposes • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

Which type of study uses random assignment?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • BACKWARD 1 • Q1/5

Which type of study is designed to estimate a population value?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • BACKWARD 1 • Q2/5

True or false?

An observational study assigns treatments.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • BACKWARD 1 • Q3/5

Random assignment is used to

A. avoid measuring

B. make the sample larger

C. make the groups comparable

D. choose who is in the population

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • BACKWARD 1 • Q4/5

What does random sampling help you do?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • BACKWARD 1 • Q5/5

SET 66 • HSS-IC.B.3

BACKWARD 2

Choose a random sample • Grade 7 • Statistics & Probability • 7.SP.A.1

Describe how to choose 20 names at random from a list of 500.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 66 • BACKWARD 2 • Q1/5

Is a sample of volunteers random?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 66 • BACKWARD 2 • Q2/5

True or false?

A sample of volunteers may differ from the population.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 66 • BACKWARD 2 • Q3/5

Which method gives a random sample?

A. choosing the first 20

B. drawing numbered slips from a well-mixed hat

C. asking for volunteers

D. asking friends

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 66 • BACKWARD 2 • Q4/5

Why must slips in a hat be well mixed?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 66 • BACKWARD 2 • Q5/5

SET 66 • HSS-IC.B.3

BACKWARD 3

Correlation and causation • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9

An association between two variables in observational data: can it prove causation?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B3-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 66 • BACKWARD 3 • Q1/5

Give a possible lurking variable for "exercise and sleep".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B3-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 66 • BACKWARD 3 • Q2/5

True or false?

Random assignment controls lurking variables on average.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B3-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 66 • BACKWARD 3 • Q3/5

Causation can be concluded from

A. an observational study

B. a correlation alone

C. a randomized experiment

D. a survey

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B3-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 66 • BACKWARD 3 • Q4/5

Why does a correlation not show which variable causes the other?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-B3-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 66 • BACKWARD 3 • Q5/5

SET 66 • HSS-IC.B.3

FORWARD 1

Improve an experimental design • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

Explain why a control group is needed in the diet experiment.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • FORWARD 1 • Q1/5

A drug study blocks subjects by age before randomly assigning treatments. Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • FORWARD 1 • Q2/5

True or false?

In a double-blind experiment, neither subjects nor evaluators know who gets the treatment.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • FORWARD 1 • Q3/5

Which change lets the diet result be generalized to all adults?

A. measuring more often

B. using a placebo

C. adding more volunteers

D. randomly selecting the subjects from all adults

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • FORWARD 1 • Q4/5

Why are many subjects better than few in an experiment?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 66 • FORWARD 1 • Q5/5

SET 66 • HSS-IC.B.3

FORWARD 2

Use re-randomization to judge a treatment difference • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

In the diet experiment, diet A lost 2.4 kg more. In 200 re-randomizations, 5 differences were at least 2.4 kg. Is the difference significant?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 66 • FORWARD 2 • Q1/5

State the conclusion in context, including to whom it applies.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 66 • FORWARD 2 • Q2/5

True or false?

Re-randomization shuffles the group labels as if the treatment had no effect.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 66 • FORWARD 2 • Q3/5

If 70 of 200 re-randomized differences were at least 2.4 kg, conclude

A. the difference could be due to chance

B. the study was observational

C. diet B works

D. diet A works

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 66 • FORWARD 2 • Q4/5

Explain why re-randomization is valid only because the original assignment was random.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S66-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 66 • FORWARD 2 • Q5/5

SET 67 • HSS-IC.B.4 • STATISTICS, INFERENCE, AND DECISIONS

Estimate a population proportion with a margin of error from a simulation

MAIN	Algebra II main question	**HSS-IC.B.4**
BACKWARD 1	Statistics, parameters, and sampling variability	**HSS-IC.A.1**
BACKWARD 2	Proportions and percents	**6.RP.A.3.c**
BACKWARD 3	Standard deviation as typical distance from the mean	**HSS-ID.A.2**
FORWARD 1	Plan a sample size for a desired margin of error	**HSS-IC.B.4**
FORWARD 2	Evaluate reports that use margins of error	**HSS-IC.B.6**

SET 67 • HSS-IC.B.4

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

In a random sample of 100 students, 58 want a later start time. The histogram shows 200 simulated samples of 100 from a population with p = 0.58; their standard deviation is 0.049. Give a margin of error and an interval estimate.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • MAIN • Q1/5

In a random sample of 300 voters, 120 support a measure. Simulated sample proportions (n = 300, p = 0.40) have standard deviation 0.0275. Give the estimate, the margin of error, and the interval.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • MAIN • Q2/5

In a random sample of 50 shoppers, 35 use a store app. Simulated proportions (n = 50, p = 0.70) have standard deviation 0.064. Give the margin of error and the interval.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • MAIN • Q3/5

Of the three surveys (n = 50, 100, 300), which has the smallest margin of error, and why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • MAIN • Q4/5

A report says "52% of voters support the plan, with a margin of error of 3 percentage points." Interpret this statement.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • MAIN • Q5/5

SET 67 • HSS-IC.B.4

BACKWARD 1

Statistics, parameters, and sampling variability • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

In the student survey, what is the statistic?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 67 • BACKWARD 1 • Q1/5

What parameter does it estimate?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 67 • BACKWARD 1 • Q2/5

True or false?

Different random samples of 100 would give different sample proportions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 67 • BACKWARD 1 • Q3/5

The simulation histogram shows

A. the parameter exactly

B. how sample proportions vary from sample to sample

C. the population

D. one sample

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 67 • BACKWARD 1 • Q4/5

Why does a random sample matter for the margin of error to be meaningful?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 67 • BACKWARD 1 • Q5/5

SET 67 • HSS-IC.B.4

BACKWARD 2

Proportions and percents • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c

Write 120 out of 300 as a proportion.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 67 • BACKWARD 2 • Q1/5

What is 0.58 ± 0.10 as an interval?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 67 • BACKWARD 2 • Q2/5

True or false?

35 out of 50 is 70%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 67 • BACKWARD 2 • Q3/5

52% ± 3% gives

A. 52% to 55%

B. 49% to 55%

C. 50% to 54%

D. 49% to 52%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 67 • BACKWARD 2 • Q4/5

Write 0.055 as a percent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B2-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 67 • BACKWARD 2 • Q5/5

SET 67 • HSS-IC.B.4

BACKWARD 3

Standard deviation as typical distance from the mean • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2

About what percent of mound-shaped data lie within 2 standard deviations of the mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B3-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 67 • BACKWARD 3 • Q1/5

If the SD is 0.049, what is 2 SDs?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B3-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 67 • BACKWARD 3 • Q2/5

True or false?

A smaller standard deviation means less variability.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B3-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 67 • BACKWARD 3 • Q3/5

Which set of simulated proportions varies least?

A. SD 0.1

B. SD 0.0275

C. SD 0.049

D. SD 0.064

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B3-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 67 • BACKWARD 3 • Q4/5

Compute 2×0.064.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-B3-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 67 • BACKWARD 3 • Q5/5

SET 67 • HSS-IC.B.4

FORWARD 1

Plan a sample size for a desired margin of error • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

Quadrupling the sample size roughly halves the margin of error. Estimate the margin for the student survey with n = 400.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • FORWARD 1 • Q1/5

About what sample size gives a margin of error of 0.05 when $p \approx 0.5$? (Use margin $\approx 2\sqrt{(p(1-p)/n)}$.)

$n \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • FORWARD 1 • Q2/5

True or false?

Doubling the sample size halves the margin of error.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • FORWARD 1 • Q3/5

Which change most reduces the margin of error?

A. multiplying n by 9

B. doubling n

C. adding 10 people

D. repeating the same sample

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • FORWARD 1 • Q4/5

Two polls of the same population have margins 0.05 and 0.025. How do their sample sizes compare?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 67 • FORWARD 1 • Q5/5

SET 67 • HSS-IC.B.4

FORWARD 2

Evaluate reports that use margins of error • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

A poll shows candidate A at 51% and B at 49% with margin ±4%. A headline says "A leads." Evaluate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 67 • FORWARD 2 • Q1/5

A report gives 58% ± 10% from 100 students and claims "a majority of students want a later start." Evaluate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 67 • FORWARD 2 • Q2/5

True or false?

A margin of error accounts for bias from a voluntary-response sample.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 67 • FORWARD 2 • Q3/5

Which report is best supported?

A. 45% ± 2% so a majority oppose

B. 70% ± 3% support, from a random sample

C. 51% ± 5% support, so most support

D. 60% support from an online poll

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 67 • FORWARD 2 • Q4/5

What should a careful report include besides the percent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S67-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 67 • FORWARD 2 • Q5/5

SET 68 • HSS-IC.B.4 • STATISTICS, INFERENCE, AND DECISIONS

Estimate a population mean with a margin of error from a simulation

MAIN	Algebra II main question	**HSS-IC.B.4**
BACKWARD 1	Mean of a data set	**6.SP.B.5.c**
BACKWARD 2	What resampling simulates	**HSS-IC.A.1**
BACKWARD 3	Two standard deviations and the 95% range	**HSS-ID.A.2**
FORWARD 1	Use intervals to judge claims about a mean	**HSS-IC.B.4**
FORWARD 2	Compare two group means with re-randomization	**HSS-IC.B.5**

SET 68 • HSS-IC.B.4

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

A random sample of 40 students slept a mean of 6.8 hours. The histogram shows the means of 200 resamples of the data; their standard deviation is 0.20 hours. Give a margin of error and an interval estimate.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • MAIN • Q1/5

A random sample of 25 apples has mean weight 152 g. Simulated sample means have standard deviation 3.6 g. Give the margin of error and the interval.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • MAIN • Q2/5

A random sample of 60 commuters has mean commute 24.5 min. Simulated means have standard deviation 1.4 min. Give the margin of error and the interval.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • MAIN • Q3/5

Which statement interprets the sleep interval 6.4 to 7.2 hours correctly? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • MAIN • Q4/5

The apple sample is increased from 25 to 100 apples. About what margin of error do you expect?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • MAIN • Q5/5

SET 68 • HSS-IC.B.4

BACKWARD 1

Mean of a data set • Grade 6 • Statistics & Probability • 6.SP.B.5.c

Find the mean of 6, 7, 7.5, 6.5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q1

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 68 • BACKWARD 1 • Q1/5

What does a sample mean estimate?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q2

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 68 • BACKWARD 1 • Q2/5

True or false?

The mean is affected by extreme values.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q3

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 68 • BACKWARD 1 • Q3/5

The mean of 150, 152, 154 is

A. 154

B. 150

C. 152

D. 456

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q4

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 68 • BACKWARD 1 • Q4/5

If each of 40 values increases by 0.5, how does the mean change?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B1-Q5

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 68 • BACKWARD 1 • Q5/5

SET 68 • HSS-IC.B.4

BACKWARD 2

What resampling simulates • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

Why can resampling the data show how sample means vary?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 68 • BACKWARD 2 • Q1/5

In the histogram, what does one bar's height mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 68 • BACKWARD 2 • Q2/5

True or false?

Sample means vary less than individual values.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 68 • BACKWARD 2 • Q3/5

The parameter being estimated in the sleep study is

A. 6.8

B. the mean sleep of all students

C. the 40 students' mean

D. the resample SD

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 68 • BACKWARD 2 • Q4/5

Is 6.8 hours a parameter or a statistic?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 68 • BACKWARD 2 • Q5/5

SET 68 • HSS-IC.B.4

BACKWARD 3

Two standard deviations and the 95% range • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2

Compute $6.8 - 2(0.20)$ and $6.8 + 2(0.20)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B3-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 68 • BACKWARD 3 • Q1/5

Compute 2×1.4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B3-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 68 • BACKWARD 3 • Q2/5

True or false?

About 95% of a mound-shaped distribution is within 2 SDs of its mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B3-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 68 • BACKWARD 3 • Q3/5

152 ± 7.2 gives

A. 144.8 to 159.2

B. 145 to 159

C. 152 to 159.2

D. 144.8 to 152

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B3-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 68 • BACKWARD 3 • Q4/5

Is 7.5 hours inside the interval 6.4 to 7.2?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-B3-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 68 • BACKWARD 3 • Q5/5

SET 68 • HSS-IC.B.4

FORWARD 1

Use intervals to judge claims about a mean • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

A health guide says students average 8 hours of sleep. Does the interval 6.4 to 7.2 support this? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • FORWARD 1 • Q1/5

A grower claims apples average 155 g. Is the claim consistent with 144.8 to 159.2 g?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • FORWARD 1 • Q2/5

True or false?

A value inside the interval is proved to be the true mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • FORWARD 1 • Q3/5

Which change would most likely narrow the commute interval?

A. sampling 240 commuters instead of 60

B. using the median

C. rounding the data

D. sampling 30 commuters

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • FORWARD 1 • Q4/5

Explain why the sleep interval should not be applied to students at a different school.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 68 • FORWARD 1 • Q5/5

SET 68 • HSS-IC.B.4

FORWARD 2

Compare two group means with re-randomization • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

In an experiment, 20 students using a study app scored 6 points more on average than 20 without it. In 200 re-randomizations, 9 differences were at least 6. Is the difference significant at the 5% level?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 68 • FORWARD 2 • Q1/5

What model do the re-randomizations represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 68 • FORWARD 2 • Q2/5

True or false?

An interval for one mean and a re-randomization test for two means both use simulated variability.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 68 • FORWARD 2 • Q3/5

If 30 of 200 differences were at least 6, the conclusion is

A. the difference could be chance

B. the app works

C. the study failed

D. the app hurts scores

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 68 • FORWARD 2 • Q4/5

Explain how the margin of error for one mean is related to judging a difference of two means.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S68-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 68 • FORWARD 2 • Q5/5

SET 69 • HSS-IC.B.5 • STATISTICS, INFERENCE, AND DECISIONS

Compare two treatments with a re-randomization simulation

MAIN	Algebra II main question	**HSS-IC.B.5**
BACKWARD 1	Compare the means of two groups	**7.SP.B.4**
BACKWARD 2	Random assignment in experiments	**HSS-IC.B.3**
BACKWARD 3	Probability as relative frequency in simulations	**7.SP.C.6**
FORWARD 1	Two-sided comparisons and design choices	**HSS-IC.B.5**
FORWARD 2	Evaluate reports of experiments	**HSS-IC.B.6**

SET 69 • HSS-IC.B.5

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

10 plants got fertilizer and 10 did not (randomly assigned). Fertilized plants grew 2.9 cm more on average. The histogram shows 200 re-randomized differences; none was 2.9 or more. Is the difference significant? Explain.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • MAIN • Q1/5

In a caffeine experiment, the caffeine group's reaction time was 0.04 s faster on average. In 200 re-randomizations, 62 differences were at least 0.04 s. Is the difference significant?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • MAIN • Q2/5

Students randomly assigned to a study app scored 6 points higher on average. In 200 re-randomizations, 9 differences were at least 6. Is the difference significant at the 5% level?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • MAIN • Q3/5

Explain what a re-randomization simulation assumes and how it is carried out.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • MAIN • Q4/5

Using the histogram of 200 re-randomized differences, about how many were at least 2 cm? Would an observed difference of 2 cm be significant at the 5% level?



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • MAIN • Q5/5

SET 69 • HSS-IC.B.5

BACKWARD 1

Compare the means of two groups • Grade 7 • Statistics & Probability • 7.SP.B.4

Group A has mean 25.4 cm and group B has mean 22.5 cm. Find the difference A − B.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.B.4 • Set 69 • BACKWARD 1 • Q1/5

Group A: 4, 6, 8. Group B: 3, 5, 4. Find the difference in means.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.B.4 • Set 69 • BACKWARD 1 • Q2/5

True or false?

A difference in sample means could happen even if the treatments have the same effect.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.B.4 • Set 69 • BACKWARD 1 • Q3/5

Mean A = 0.52 s and mean B = 0.48 s. A − B is

A. −0.04 s

B. 0.4 s

C. 1.00 s

D. 0.04 s

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.B.4 • Set 69 • BACKWARD 1 • Q4/5

Why compare the difference with the variability in the groups?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.B.4 • Set 69 • BACKWARD 1 • Q5/5

SET 69 • HSS-IC.B.5

BACKWARD 2

Random assignment in experiments • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

Why were the plants randomly assigned to the two groups?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 69 • BACKWARD 2 • Q1/5

Describe how to randomly assign 20 plants into two groups of 10.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 69 • BACKWARD 2 • Q2/5

True or false?

Without random assignment, a re-randomization test does not justify a cause-and-effect conclusion.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 69 • BACKWARD 2 • Q3/5

Random assignment is a feature of

A. censuses

B. experiments

C. observational studies

D. surveys

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 69 • BACKWARD 2 • Q4/5

What kind of conclusion does a significant result in a randomized experiment allow?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 69 • BACKWARD 2 • Q5/5

SET 69 • HSS-IC.B.5

BACKWARD 3

Probability as relative frequency in simulations • Grade 7 • Statistics & Probability • 7.SP.C.6

62 of 200 is what percent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B3-Q1

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 69 • BACKWARD 3 • Q1/5

9 of 200 is what percent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B3-Q2

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 69 • BACKWARD 3 • Q2/5

True or false?

An event that happens in 1% of simulated runs is unusual.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B3-Q3

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 69 • BACKWARD 3 • Q3/5

0 of 200 runs suggests the observed result is

A. very unlikely under the model

B. certain

C. impossible

D. typical

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B3-Q4

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 69 • BACKWARD 3 • Q4/5

Is 4/200 below 5%?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-B3-Q5

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 69 • BACKWARD 3 • Q5/5

SET 69 • HSS-IC.B.5

FORWARD 1

Two-sided comparisons and design choices • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

In the caffeine study, the question is whether caffeine changes reaction time in either direction. How should the re-randomization results be counted?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • FORWARD 1 • Q1/5

Using the fertilizer histogram, estimate how many re-randomized differences were 2 or more in absolute value.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • FORWARD 1 • Q2/5

True or false?

Using more plants in each group usually makes the re-randomized differences less spread out.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • FORWARD 1 • Q3/5

A small observed difference can be significant when

A. the groups have very little natural variation

B. the data are rounded

C. the groups were not randomized

D. the sample is small

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • FORWARD 1 • Q4/5

Why should the decision rule (such as 5%) be chosen before seeing the data?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 69 • FORWARD 1 • Q5/5

SET 69 • HSS-IC.B.5

FORWARD 2

Evaluate reports of experiments • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

A company reports its supplement "improved memory scores by 3 points" in a study of 8 people without a control group. Evaluate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 69 • FORWARD 2 • Q1/5

A report says a new teaching method "significantly" raised scores in a randomized experiment, with 2 of 200 re-randomizations as extreme. Evaluate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 69 • FORWARD 2 • Q2/5

True or false?

"Statistically significant" means the effect is large and important.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 69 • FORWARD 2 • Q3/5

Which report is most convincing?

A. a study with no comparison group

B. testimonials from customers

C. a randomized experiment with a control group and a small simulated p-value

D. an online poll of users

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 69 • FORWARD 2 • Q4/5

What information should a report of an experiment include?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S69-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 69 • FORWARD 2 • Q5/5

SET 70 • HSS-IC.B.6 • STATISTICS, INFERENCE, AND DECISIONS

Evaluate claims in reports based on data

MAIN	Algebra II main question	**HSS-IC.B.6**
BACKWARD 1	Random samples and bias	**HSS-IC.A.1**
BACKWARD 2	Study design and causation	**HSS-IC.B.3**
BACKWARD 3	Read scales on graphs	**3.MD.B.3**
FORWARD 1	Rewrite claims so that they follow from the data	**HSS-IC.B.6**
FORWARD 2	Analyze decisions based on test results	**HSS-MD.B.7**

SET 70 • HSS-IC.B.6

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

An ad says "4 out of 5 dentists recommend our toothpaste." Only 5 dentists, chosen by the company, were asked. Evaluate the claim.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • MAIN • Q1/5

A report says "Students who eat breakfast earn higher grades, so breakfast causes higher grades." The data came from a survey. Evaluate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • MAIN • Q2/5

A bar graph shows sales of 50 and 52 thousand, with the vertical axis starting at 48 thousand, so one bar looks twice as tall. Evaluate the impression the graph gives.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • MAIN • Q3/5

A TV show asks viewers to text their vote; 90% of 10,000 texts favor a policy. The show claims most citizens support it. Evaluate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • MAIN • Q4/5

A poll shows candidate A at 51% and B at 49% with a margin of error of ±4%. A headline says "A is winning." Evaluate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • MAIN • Q5/5

SET 70 • HSS-IC.B.6

BACKWARD 1

Random samples and bias • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

Why is a voluntary-response sample biased?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 70 • BACKWARD 1 • Q1/5

What does a random sample allow?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 70 • BACKWARD 1 • Q2/5

True or false?

A very large sample cannot be biased.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 70 • BACKWARD 1 • Q3/5

Which sample is most likely biased?

A. customers who chose to write reviews

B. a random sample of 1,000

C. randomly chosen customers

D. every customer

Is "5 dentists chosen by the company" a random sample?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 70 • BACKWARD 1 • Q5/5

SET 70 • HSS-IC.B.6

BACKWARD 2

Study design and causation • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

What type of study is a survey of students' breakfast habits and grades?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 70 • BACKWARD 2 • Q1/5

What study design could test whether breakfast causes higher grades?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 70 • BACKWARD 2 • Q2/5

True or false?

Only randomized experiments can establish cause and effect.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 70 • BACKWARD 2 • Q3/5

A confounding variable is

A. the response variable

B. a variable related to both the explanatory and the response variable

C. the sample size

D. the margin of error

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 70 • BACKWARD 2 • Q4/5

Name a possible confounder for breakfast and grades.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 70 • BACKWARD 2 • Q5/5

SET 70 • HSS-IC.B.6

BACKWARD 3

Read scales on graphs • Grade 3 • Measurement & Data • 3.MD.B.3

A bar graph axis starts at 48. Bars reach 50 and 52. What are the bar values?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B3-Q1

Grade 3 • Measurement & Data • 3.MD.B.3 • Set 70 • BACKWARD 3 • Q1/5

What percent larger is 52 than 50?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B3-Q2

Grade 3 • Measurement & Data • 3.MD.B.3 • Set 70 • BACKWARD 3 • Q2/5

True or false?

An axis that does not start at 0 can make differences look larger.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B3-Q3

Grade 3 • Measurement & Data • 3.MD.B.3 • Set 70 • BACKWARD 3 • Q3/5

On a graph whose axis starts at 48, a bar of 52 looks how many times as tall as a bar of 50?

A. the same

B. 1.04 times

C. 2 times

D. 4 times

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B3-Q4

Grade 3 • Measurement & Data • 3.MD.B.3 • Set 70 • BACKWARD 3 • Q4/5

How should the graph be redrawn to be fair?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-B3-Q5

Grade 3 • Measurement & Data • 3.MD.B.3 • Set 70 • BACKWARD 3 • Q5/5

SET 70 • HSS-IC.B.6

FORWARD 1

Rewrite claims so that they follow from the data • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6

Rewrite the breakfast claim so that it is supported by a survey.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • FORWARD 1 • Q1/5

Rewrite the poll headline (51% vs 49%, ±4%) so it is accurate.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • FORWARD 1 • Q2/5

True or false?

Saying "is associated with" instead of "causes" is appropriate for an observational study.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • FORWARD 1 • Q3/5

Which conclusion is justified by a random sample of 1,000 voters showing 62% ± 3% support?

A. the sample was biased

B. most voters support the measure

C. exactly 62% support it

D. the measure will pass by 24 points

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • FORWARD 1 • Q4/5

What would the dentist claim need to be convincing?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.6 • Set 70 • FORWARD 1 • Q5/5

SET 70 • HSS-IC.B.6

FORWARD 2

Analyze decisions based on test results • High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7

A screening test is 95% accurate for people with a disease and gives false positives 5% of the time. If 1% of 10,000 people have the disease, how many positive results are true positives?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 70 • FORWARD 2 • Q1/5

A report says "a positive test means you almost surely have the disease." Evaluate using your counts.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions HSS-MD.B.7 • Set 70 • FORWARD 2 • Q2/5

True or false?

The probability of having the disease given a positive result can be low even with an accurate test.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 70 • FORWARD 2 • Q3/5

With 1% prevalence, P(disease | positive) is about

A. 0.99

B. 0.05

C. 0.95

D. 0.16

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F2-Q4

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 70 • FORWARD 2 • Q4/5

What follow-up decision makes sense after a positive screening result?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S70-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 70 • FORWARD 2 • Q5/5

SET 71 • HSS-MD.B.6 • STATISTICS, INFERENCE, AND DECISIONS

Design and judge fair decision procedures with probabilities

MAIN	Algebra II main question	**HSS-MD.B.6**
BACKWARD 1	Probabilities of equally likely outcomes	**7.SP.C.7.a**
BACKWARD 2	Sample spaces for compound events	**7.SP.C.8.b**
BACKWARD 3	Probability between 0 and 1 and redoing outcomes	**7.SP.C.5**
FORWARD 1	Design fair procedures and count expected trials	**HSS-MD.B.6**
FORWARD 2	Analyze strategies and decisions with probability	**HSS-MD.B.7**

SET 71 • HSS-MD.B.6

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6

Three friends choose a driver by flipping a coin twice: HH = Ana, HT = Ben, TH = Cy, TT = flip again. Is this fair? Justify with probabilities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6 • Set 71 • MAIN • Q1/5

Four people use a die: 1 = A, 2 = B, 3 = C, 4 = D, and 5 or 6 means roll again. Is this fair? Justify.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6 • Set 71 • MAIN • Q2/5

A hat holds 3 slips with Ana's name and 2 with Ben's. One slip is drawn to choose who goes first. Is this fair? Justify.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6 • Set 71 • MAIN • Q3/5

A random number generator gives a whole number 1-100. Numbers 1-33 choose A, 34-66 choose B, 67-99 choose C, and 100 means generate again. Is this fair? Justify.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6 • Set 71 • MAIN • Q4/5

Two dice are rolled; if the sum is even, player 1 wins, otherwise player 2 wins. Is this fair? Justify.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6 • Set 71 • MAIN • Q5/5

SET 71 • HSS-MD.B.6

BACKWARD 1

Probabilities of equally likely outcomes • Grade 7 • Statistics & Probability • 7.SP.C.7.a

A fair die is rolled. What is P(3)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.C.7.a • Set 71 • BACKWARD 1 • Q1/5

A bag has 3 red and 2 blue slips. What is P(red)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.C.7.a • Set 71 • BACKWARD 1 • Q2/5

True or false?

Each face of a fair die has the same probability.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.C.7.a • Set 71 • BACKWARD 1 • Q3/5

Two fair coins: P(HT) is

A. 1/2

B. 1/4

C. 1/8

D. 1/3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.C.7.a • Set 71 • BACKWARD 1 • Q4/5

A spinner has 4 equal sections. What is P(one section)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.C.7.a • Set 71 • BACKWARD 1 • Q5/5

SET 71 • HSS-MD.B.6

BACKWARD 2

Sample spaces for compound events • Grade 7 • Statistics & Probability • 7.SP.C.8.b

List the outcomes of flipping a coin twice.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.C.8.b • Set 71 • BACKWARD 2 • Q1/5

How many outcomes does rolling two dice have?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.C.8.b • Set 71 • BACKWARD 2 • Q2/5

True or false?

There are 3 outcomes for two coin flips: two heads, one head, no heads.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.C.8.b • Set 71 • BACKWARD 2 • Q3/5

How many outcomes for two dice give an even sum?

A. 21

B. 12

C. 18

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.C.8.b • Set 71 • BACKWARD 2 • Q4/5

List the outcomes of rolling two dice with a sum of 4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.C.8.b • Set 71 • BACKWARD 2 • Q5/5

SET 71 • HSS-MD.B.6

BACKWARD 3

Probability between 0 and 1 and redoing outcomes • Grade 7 • Statistics & Probability • 7.SP.C.5

If an outcome is ignored and the experiment is redone, how does that affect the remaining outcomes' chances?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B3-Q1

Grade 7 • Statistics & Probability • 7.SP.C.5 • Set 71 • BACKWARD 3 • Q1/5

With rerolls on 5 and 6, what is P(1) for a die?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B3-Q2

Grade 7 • Statistics & Probability • 7.SP.C.5 • Set 71 • BACKWARD 3 • Q2/5

True or false?

Probabilities of all outcomes in a sample space add to 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B3-Q3

Grade 7 • Statistics & Probability • 7.SP.C.5 • Set 71 • BACKWARD 3 • Q3/5

An event that can never happen has probability

A. 1

B. -1

C. 1/2

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B3-Q4

Grade 7 • Statistics & Probability • 7.SP.C.5 • Set 71 • BACKWARD 3 • Q4/5

Three equally likely outcomes: what is each probability?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-B3-Q5

Grade 7 • Statistics & Probability • 7.SP.C.5 • Set 71 • BACKWARD 3 • Q5/5

SET 71 • HSS-MD.B.6

FORWARD 1

Design fair procedures and count expected trials • High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.6

Design a fair way to choose one of 5 people using a fair coin.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.6 • Set 71 • FORWARD 1 • Q1/5

In the 2-flip method for 3 friends, what is the probability that a choice is made on the first try?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions HSS-MD.B.6 • Set 71 • FORWARD 1 • Q2/5

True or false?

A fair procedure must give each person an equal chance, even if it may take several tries.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.6 • Set 71 • FORWARD 1 • Q3/5

To choose among 6 people fairly with one die, you

A. roll twice and add

B. assign one face to each person

C. assign 1-3 to person A

D. reroll on 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F1-Q4

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions HSS-MD.B.6 • Set 71 • FORWARD 1 • Q4/5

A teacher picks a student from a class of 28 with a random number generator 1-30. Design a fair rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F1-Q5

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions HSS-MD.B.6 • Set 71 • FORWARD 1 • Q5/5

SET 71 • HSS-MD.B.6

FORWARD 2

Analyze strategies and decisions with probability • High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7

A game costs $1. You roll a die and win $4 if you roll a 6. Find the expected net gain per game.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 71 • FORWARD 2 • Q1/5

Is the game fair in the sense of expected value? What prize would make it fair?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 71 • FORWARD 2 • Q2/5

True or false?

A game can be fair in who is chosen but still have a negative expected value for the player.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 71 • FORWARD 2 • Q3/5

With P(win) = 1/4 and prize \$8 for a \$2 ticket, the expected net gain is

A. \$2

B. −\$2

C. \$0

D. \$6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F2-Q4

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 71 • FORWARD 2 • Q4/5

Explain the difference between a fair choice procedure and a fair game.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S71-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 71 • FORWARD 2 • Q5/5

SET 72 • HSS-MD.B.7 • STATISTICS, INFERENCE, AND DECISIONS

Analyze decisions and strategies with probabilities

MAIN	Algebra II main question	**HSS-MD.B.7**
BACKWARD 1	Probability of compound independent events	**7.SP.C.8.a**
BACKWARD 2	Two-way tables of counts	**8.SP.A.4**
BACKWARD 3	Weighted averages of outcomes	**6.RP.A.3.c**
FORWARD 1	Investigate how changing conditions changes the decision	**HSS-MD.B.7**
FORWARD 2	Find expected values of random variables	**HSS-MD.A.2**

SET 72 • HSS-MD.B.7

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7

A disease affects 1% of 10,000 people. A test detects 95% of cases and gives false positives for 5% of healthy people. Find P(disease | positive).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7 • Set 72 • MAIN • Q1/5

A shipment is accepted if 2 randomly tested items are both non-defective. If 10% of items are defective, what is the probability the shipment is accepted? (Assume independence.)

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7 • Set 72 • MAIN • Q2/5

A $30 warranty covers a repair that costs $200 and happens with probability 0.1. Compare the expected repair cost without the warranty to the warranty price, and decide.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7 • Set 72 • MAIN • Q3/5

Option A pays $10 with probability 0.5; option B pays $50 with probability 0.1. Compare the expected values and the risks.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7 • Set 72 • MAIN • Q4/5

A test gives +1 point for a correct answer and −1/4 point for a wrong one on 4-choice questions. Should a student guess randomly on a question? Justify with expected value.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7 • Set 72 • MAIN • Q5/5

SET 72 • HSS-MD.B.7

BACKWARD 1

Probability of compound independent events • Grade 7 • Statistics & Probability • 7.SP.C.8.a

P(non-defective) = 0.9 for each item. Find P(two non-defective).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.C.8.a • Set 72 • BACKWARD 1 • Q1/5

Two fair coins: P(both heads)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.C.8.a • Set 72 • BACKWARD 1 • Q2/5

True or false?

For independent events, $P(A \text{ and } B) = P(A) \cdot P(B)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.C.8.a • Set 72 • BACKWARD 1 • Q3/5

P(three non-defective) with P = 0.9 each is

A. 0.9

B. 0.27

C. 0.729

D. 2.7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.C.8.a • Set 72 • BACKWARD 1 • Q4/5

Find P(at least one defective) among two items when P(defective) = 0.1.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.C.8.a • Set 72 • BACKWARD 1 • Q5/5

SET 72 • HSS-MD.B.7

BACKWARD 2

Two-way tables of counts • Grade 8 • Statistics & Probability • 8.SP.A.4

Of 10,000 people, 1% have a disease. How many have it?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B2-Q1

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 72 • BACKWARD 2 • Q1/5

Of 9,900 healthy people, 5% test positive. How many?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B2-Q2

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 72 • BACKWARD 2 • Q2/5

True or false?

A two-way table can show test results for people with and without a disease.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B2-Q3

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 72 • BACKWARD 2 • Q3/5

If 95 sick and 495 healthy people test positive, the total positives are

A. 10,000

B. 590

C. 95

D. 495

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B2-Q4

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 72 • BACKWARD 2 • Q4/5

What fraction of positives are sick (95 of 590)? Give a decimal to the nearest thousandth.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B2-Q5

Grade 8 • Statistics & Probability • 8.SP.A.4 • Set 72 • BACKWARD 2 • Q5/5

SET 72 • HSS-MD.B.7

BACKWARD 3

Weighted averages of outcomes • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c

Find 10% of $200.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 72 • BACKWARD 3 • Q1/5

Compute 0.5 × 10 and 0.1 × 50.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 72 • BACKWARD 3 • Q2/5

True or false?

$1/4 \times 1 + 3/4 \times (-1/4) = 1/16.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 72 • BACKWARD 3 • Q3/5

0.2 × 30 + 0.8 × 0 equals

A. 30

B. 24

C. 0.2

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 72 • BACKWARD 3 • Q4/5

Compute $0.9 \times 0 + 0.1 \times 200$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 72 • BACKWARD 3 • Q5/5

SET 72 • HSS-MD.B.7

FORWARD 1

Investigate how changing conditions changes the decision • High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.B.7

In a group where 10% have the disease, find P(disease | positive) for the same test.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 72 • FORWARD 1 • Q1/5

If the warranty repair probability rises to 0.2, is the $30 warranty worth it on average?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 72 • FORWARD 1 • Q2/5

True or false?

A positive result is more convincing when the disease is more common.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 72 • FORWARD 1 • Q3/5

If students could eliminate one wrong choice before guessing (3 choices left), the expected value of guessing is

A. $1/3 - 2/3 \cdot 1/4 = 1/6$

B. 0

C. $1/16$

D. $-1/4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S72-F1-Q4**

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 72 • FORWARD 1 • Q4/5

How many items must be tested (all non-defective) for the acceptance probability to drop below 0.5 when 10% are defective?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F1-Q5

High School — Algebra II, (+) standard listed in the pathway • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.B.7 • Set 72 • FORWARD 1 • Q5/5

SET 72 • HSS-MD.B.7

FORWARD 2

Find expected values of random variables • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions • HSS-MD.A.2

A random variable X takes values 0, 1, 2 with probabilities 0.25, 0.5, 0.25. Find E(X).

E(X) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.A.2 • Set 72 • FORWARD 2 • Q1/5

A raffle sells 500 tickets at $2; one prize is $400. Find the expected net gain of one ticket.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.A.2 • Set 72 • FORWARD 2 • Q2/5

True or false?

The expected value is the long-run average of many repetitions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F2-Q3

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.A.2 • Set 72 • FORWARD 2 • Q3/5

For a fair die, E(roll) is

A. 3

B. 21

C. 3.5

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S72-F2-Q4**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.A.2 • Set 72 • FORWARD 2 • Q4/5

Explain how the warranty decision is an expected-value calculation for a random variable.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S72-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Using Probability to Make Decisions
HSS-MD.A.2 • Set 72 • FORWARD 2 • Q5/5

SET 73 • HSN-CN.A.3 • ADVANCED EXTENSION (+)

Use conjugates to find moduli and quotients of complex numbers

MAIN	Algebra II main question	**HSN-CN.A.3**
BACKWARD 1	Multiply complex numbers	**HSN-CN.A.2**
BACKWARD 2	Distance from the origin	**8.G.B.8**
BACKWARD 3	Powers of i	**HSN-CN.A.1**
FORWARD 1	Prove properties of conjugates and moduli	**HSN-CN.A.3**
FORWARD 2	Write complex numbers in polar form	**HSN-CN.B.4**

SET 73 • HSN-CN.A.3

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3

Give the conjugate of $3 - 4i$ and find $|3 - 4i|$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-M-Q1

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • MAIN • Q1/5

Write (2 + 3i)/(1 − i) in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-M-Q2

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • MAIN • Q2/5

Write (5 + i)/(2 + 3i) in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-M-Q3

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • MAIN • Q3/5

Find $|-5 + 12i|$ and compute $(-5 + 12i)(-5 - 12i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-M-Q4

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • MAIN • Q4/5

Write 1/(4 + 3i) in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-M-Q5

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • MAIN • Q5/5

SET 73 • HSN-CN.A.3

BACKWARD 1

Multiply complex numbers • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Multiply $(2 + 3i)(1 + i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 73 • BACKWARD 1 • Q1/5

Multiply $(3 - 4i)(3 + 4i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 73 • BACKWARD 1 • Q2/5

True or false?

$(a + bi)(a - bi) = a^2 + b^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 73 • BACKWARD 1 • Q3/5

$(5 + i)(2 - 3i)$ equals

A. $13 + 13i$

B. $7 - 13i$

C. $10 - 3i$

D. $13 - 13i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 73 • BACKWARD 1 • Q4/5

Multiply $(1 - i)(1 + i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 73 • BACKWARD 1 • Q5/5

SET 73 • HSN-CN.A.3

BACKWARD 2

Distance from the origin • Grade 8 • Geometry • 8.G.B.8

Find the distance from (0, 0) to (3, −4).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B2-Q1

Grade 8 • Geometry • 8.G.B.8 • Set 73 • BACKWARD 2 • Q1/5

Find the distance from (0, 0) to (−5, 12).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B2-Q2

Grade 8 • Geometry • 8.G.B.8 • Set 73 • BACKWARD 2 • Q2/5

True or false?

The distance from (0, 0) to (a, b) is $\sqrt{(a^2 + b^2)}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B2-Q3

Grade 8 • Geometry • 8.G.B.8 • Set 73 • BACKWARD 2 • Q3/5

The distance from (0, 0) to (1, 1) is

A. 0

B. 1

C. 2

D. $\sqrt{2}$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B2-Q4

Grade 8 • Geometry • 8.G.B.8 • Set 73 • BACKWARD 2 • Q4/5

Find the distance from (0, 0) to (4, 3).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B2-Q5

Grade 8 • Geometry • 8.G.B.8 • Set 73 • BACKWARD 2 • Q5/5

SET 73 • HSN-CN.A.3

BACKWARD 3

Powers of i • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Simplify i^2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B3-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 73 • BACKWARD 3 • Q1/5

Simplify $-12i \cdot i$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B3-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 73 • BACKWARD 3 • Q2/5

True or false?

$-3i^2 = 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B3-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 73 • BACKWARD 3 • Q3/5

Which is real?

A. $3i$

B. $(4 + 3i)^2$

C. $(4 + 3i)(4 - 3i)$

D. $4 + 3i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B3-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 73 • BACKWARD 3 • Q4/5

Write 13/13 − 13i/13 in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-B3-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 73 • BACKWARD 3 • Q5/5

SET 73 • HSN-CN.A.3

FORWARD 1

Prove properties of conjugates and moduli • High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3

Prove that $z \cdot \bar{z} = |z|^2$ for $z = a + bi$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F1-Q1

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • FORWARD 1 • Q1/5

Verify $|zw| = |z||w|$ for $z = 3 + 4i$ and $w = 1 + i$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F1-Q2

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • FORWARD 1 • Q2/5

True or false?

The conjugate of a real number is itself.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F1-Q3

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • FORWARD 1 • Q3/5

If $z = 2 - 5i$, then $z + \bar{z}$ is

A. $4 - 10i$

B. 0

C. $-10i$

D. 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S73-F1-Q4**

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • FORWARD 1 • Q4/5

Use $\bar{z}/|z|^2$ to write $1/z$ for $z = 1 + 2i$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F1-Q5

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 73 • FORWARD 1 • Q5/5

SET 73 • HSN-CN.A.3

FORWARD 2

Write complex numbers in polar form • High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4

Write 1 + i in polar form $r(\cos\theta + i\sin\theta)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F2-Q1

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 73 • FORWARD 2 • Q1/5

Write $2(\cos(\pi/2) + i \sin(\pi/2))$ in rectangular form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F2-Q2

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 73 • FORWARD 2 • Q2/5

True or false?

The modulus of a complex number is the r in its polar form.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F2-Q3

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 73 • FORWARD 2 • Q3/5

The polar form of -3 is

A. $3(\cos \pi + i \sin \pi)$

B. $-3(\cos \pi + i \sin \pi)$

C. $3(\cos 0 + i \sin 0)$

D. $3(\cos(\pi/2) + i \sin(\pi/2))$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F2-Q4

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 73 • FORWARD 2 • Q4/5

Explain what the modulus and the angle of a complex number represent in the complex plane.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S73-F2-Q5

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 73 • FORWARD 2 • Q5/5

SET 74 • HSF-IF.C.7.d • ADVANCED EXTENSION (+)

Graph rational functions with zeros and asymptotes

MAIN	Algebra II main question	**HSF-IF.C.7.d**
BACKWARD 1	Rewrite a rational expression by inspection	**HSA-APR.D.6**
BACKWARD 2	Domain and excluded values	**HSF-IF.B.5**
BACKWARD 3	Sign of a quotient on intervals	**7.NS.A.2**
FORWARD 1	Graph rational functions with slant asymptotes	**HSF-IF.C.7.d**
FORWARD 2	Interpret a rational model's features in context	**HSF-IF.B.4**

SET 74 • HSF-IF.C.7.d

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d

Graph the rational function on the grid. Show every zero, the y-intercept, each vertical asymptote and the horizontal asymptote (dashed lines), and the behavior near each asymptote.

$y = (x + 1)/(x - 2)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • MAIN • Q1/5

Graph the rational function on the grid. Show every zero, the y-intercept, each vertical asymptote and the horizontal asymptote (dashed lines), and the behavior near each asymptote.

$y = 2x/(x^2 - 4)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • MAIN • Q2/5

Graph the rational function on the grid. Show every zero, the y-intercept, each vertical asymptote and the horizontal asymptote (dashed lines), and the behavior near each asymptote.

$y = (x^2 - 1)/(x^2 - 9)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • MAIN • Q3/5

Graph the rational function on the grid. Show every zero, the y-intercept, each vertical asymptote and the horizontal asymptote (dashed lines), and the behavior near each asymptote.

$y = 3/(x + 1)^2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • MAIN • Q4/5

Graph the rational function on the grid. Show every zero, the y-intercept, each vertical asymptote and the horizontal asymptote (dashed lines), and the behavior near each asymptote.

$y = (2x - 4)/(x + 3)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • MAIN • Q5/5

SET 74 • HSF-IF.C.7.d

BACKWARD 1

Rewrite a rational expression by inspection • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6

Rewrite $(x + 1)/(x - 2)$ as $1 + r/(x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 74 • BACKWARD 1 • Q1/5

Rewrite $(2x - 4)/(x + 3)$ as $2 + r/(x + 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 74 • BACKWARD 1 • Q2/5

True or false?

The constant q in $q + r/(x - a)$ is the horizontal asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 74 • BACKWARD 1 • Q3/5

$(x^2 - 1)/(x^2 - 9)$ equals

A. $1 + 8/(x^2 - 9)$

B. $1 - 8/(x^2 - 9)$

C. $1 + 1/(x^2 - 9)$

D. $1/9$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S74-B1-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 74 • BACKWARD 1 • Q4/5

Rewrite $2x/(x^2 - 4)$ in factored form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 74 • BACKWARD 1 • Q5/5

SET 74 • HSF-IF.C.7.d

BACKWARD 2

Domain and excluded values • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

Give the domain of $(x + 1)/(x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 74 • BACKWARD 2 • Q1/5

Give the domain of $3/(x + 1)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 74 • BACKWARD 2 • Q2/5

True or false?

A rational function is undefined where its denominator is 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 74 • BACKWARD 2 • Q3/5

$2x/(x^2 - 4)$ is undefined at

A. $x = \pm 2$

B. $x = 4$

C. $x = 2$ only

D. $x = 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 74 • BACKWARD 2 • Q4/5

List the excluded values of $(x^2 - 1)/(x^2 - 9)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 74 • BACKWARD 2 • Q5/5

SET 74 • HSF-IF.C.7.d

BACKWARD 3

Sign of a quotient on intervals • Grade 7 • The Number System • 7.NS.A.2

Is $(x + 1)/(x - 2)$ positive or negative at $x = 0$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B3-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 74 • BACKWARD 3 • Q1/5

Is $2x/(x^2 - 4)$ positive or negative at $x = 3$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B3-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 74 • BACKWARD 3 • Q2/5

True or false?

A quotient of two negative numbers is positive.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B3-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 74 • BACKWARD 3 • Q3/5

At $x = 2.1$, $(x + 1)/(x - 2)$ is

A. large and negative

B. undefined

C. zero

D. large and positive

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B3-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 74 • BACKWARD 3 • Q4/5

Is $3/(x + 1)^2$ ever negative? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-B3-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 74 • BACKWARD 3 • Q5/5

SET 74 • HSF-IF.C.7.d

FORWARD 1

Graph rational functions with slant asymptotes • High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d

Graph the rational function on the grid. Show every zero, the y-intercept, each vertical asymptote and the slant asymptote (dashed lines), and the behavior near each asymptote.

$y = (x^2 + 1)/x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • FORWARD 1 • Q1/5

Find the slant asymptote of $y = (x^2 - x - 1)/(x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • FORWARD 1 • Q2/5

True or false?

A rational function can cross its horizontal asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • FORWARD 1 • Q3/5

$y = (3x^2 + 1)/(x^2 + 4)$ has horizontal asymptote

A. none

B. $y = 3$

C. $y = 0$

D. $y = 1/4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • FORWARD 1 • Q4/5

Write a rational function with vertical asymptote $x = 1$, horizontal asymptote $y = 2$, and zero at $x = 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 74 • FORWARD 1 • Q5/5

SET 74 • HSF-IF.C.7.d

FORWARD 2

Interpret a rational model's features in context • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

The average cost per item is $A(x) = (500 + 4x)/x$. Interpret the vertical and horizontal asymptotes in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 74 • FORWARD 2 • Q1/5

A drug concentration is $C(t) = 20t/(t^2 + 4)$. Interpret its horizontal asymptote.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 74 • FORWARD 2 • Q2/5

True or false?

For a context, only the part of a rational graph in the context domain is meaningful.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 74 • FORWARD 2 • Q3/5

In $A(x) = (500 + 4x)/x$, as x grows the average cost

A. increases

B. is constant

C. approaches 0

D. decreases toward 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 74 • FORWARD 2 • Q4/5

Find the number of items for which the average cost is $5.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S74-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 74 • FORWARD 2 • Q5/5

SET 75 • HSF-IF.C.7.d • ADVANCED EXTENSION (+)

Distinguish holes from vertical asymptotes

MAIN	Algebra II main question	**HSF-IF.C.7.d**
BACKWARD 1	Factor numerators and denominators	**HSA-SSE.A.2**
BACKWARD 2	Simplify rational expressions and keep exclusions	**HSA-APR.D.7**
BACKWARD 3	Evaluate the simplified expression	**HSF-IF.A.2**
FORWARD 1	Construct and graph rational functions with holes	**HSF-IF.C.7.d**
FORWARD 2	Relate holes to extraneous solutions	**HSA-REI.A.2**

SET 75 • HSF-IF.C.7.d

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d

Find every hole and vertical asymptote of $f(x) = (x^2 - 4)/(x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • MAIN • Q1/5

Find every hole and vertical asymptote of $f(x) = (x - 3)/(x^2 - 9)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • MAIN • Q2/5

Find every hole and vertical asymptote of $f(x) = (x^2 + x - 6)/(x^2 - x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • MAIN • Q3/5

Find every hole and vertical asymptote of $f(x) = (x + 1)/(x^2 - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • MAIN • Q4/5

Find every hole and vertical asymptote of $f(x) = x(x - 1)/(x(x + 2))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • MAIN • Q5/5

SET 75 • HSF-IF.C.7.d

BACKWARD 1

Factor numerators and denominators • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 + x - 6$ and $x^2 - x - 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 75 • BACKWARD 1 • Q1/5

Factor $x^2 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 75 • BACKWARD 1 • Q2/5

True or false?

$x^2 - 1 = (x - 1)(x + 1)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 75 • BACKWARD 1 • Q3/5

$x^2 - 4$ equals

A. $(x - 4)(x + 1)$

B. $(x - 2)^2$

C. $(x - 2)(x + 2)$

D. $x(x - 4)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 75 • BACKWARD 1 • Q4/5

Factor $x^2 + 2x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 75 • BACKWARD 1 • Q5/5

SET 75 • HSF-IF.C.7.d

BACKWARD 2

Simplify rational expressions and keep exclusions • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Simplify $(x - 3)/((x - 3)(x + 3))$ and state the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 75 • BACKWARD 2 • Q1/5

Simplify $(x + 1)/((x - 1)(x + 1))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 75 • BACKWARD 2 • Q2/5

True or false?

After simplifying, the original excluded values are still excluded.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 75 • BACKWARD 2 • Q3/5

$x(x - 1)/(x(x + 2))$ simplifies to

A. $-1/2$

B. $(x - 1)/(x + 2)$ for all x

C. $x - 1$

D. $(x - 1)/(x + 2)$, $x \neq 0, -2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B2-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 75 • BACKWARD 2 • Q4/5

Simplify $(x^2 - 4)/(x - 2)$ and state the excluded value.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 75 • BACKWARD 2 • Q5/5

SET 75 • HSF-IF.C.7.d

BACKWARD 3

Evaluate the simplified expression • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Evaluate $1/(x + 3)$ at $x = 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 75 • BACKWARD 3 • Q1/5

Evaluate $(x + 3)/(x + 1)$ at $x = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 75 • BACKWARD 3 • Q2/5

True or false?

The y-coordinate of a hole is the value of the simplified expression there.

A. True

B. False

$(x - 1)/(x + 2)$ at $x = 0$ is

A. 1/2

B. $-1/2$

C. undefined

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 75 • BACKWARD 3 • Q4/5

Evaluate $1/(x - 1)$ at $x = -1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 75 • BACKWARD 3 • Q5/5

SET 75 • HSF-IF.C.7.d

FORWARD 1

Construct and graph rational functions with holes • High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d

Write a rational function with a hole at $x = 1$, a vertical asymptote $x = -3$, and horizontal asymptote $y = 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • FORWARD 1 • Q1/5

Graph $y = (x^2 - x - 2)/(x - 2)$ on the grid. Show the hole with an open circle.



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • FORWARD 1 • Q2/5

True or false?

A rational function can have both a hole and a vertical asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • FORWARD 1 • Q3/5

$f(x) = (x - 4)^2/(x - 4)$ has

A. no special features

B. a hole at (4, 4)

C. a vertical asymptote $x = 4$

D. a hole at (4, 0)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • FORWARD 1 • Q4/5

Explain why $x = -3$ is an asymptote but $x = 3$ is a hole for $(x - 3)/(x^2 - 9)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 75 • FORWARD 1 • Q5/5

SET 75 • HSF-IF.C.7.d

FORWARD 2

Relate holes to extraneous solutions • High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2

Solve $(x^2 - 4)/(x - 2) = 4$. Explain the result using the hole.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F2-Q1

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 75 • FORWARD 2 • Q1/5

Solve $(x - 3)/(x^2 - 9) = 1/6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F2-Q2

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 75 • FORWARD 2 • Q2/5

True or false?

An extraneous solution of f(x) = c can occur exactly at a hole whose height is c.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F2-Q3

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 75 • FORWARD 2 • Q3/5

$(x + 1)/(x^2 - 1) = -1/2$ has

A. no solution

B. $x = -1$

C. $x = 0$

D. $x = 1$

Solve $(x^2 + x - 6)/(x^2 - x - 2) = 2$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S75-F2-Q5

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 75 • FORWARD 2 • Q5/5

SET 76 • HSF-BF.A.1.c • ADVANCED EXTENSION (+)

Compose functions and interpret compositions with domains

MAIN	Algebra II main question	**HSF-BF.A.1.c**
BACKWARD 1	Build combined models	**HSF-BF.A.1.b**
BACKWARD 2	Evaluate a function at an expression	**HSF-IF.A.2**
BACKWARD 3	Domains of radical, rational, and log functions	**HSF-IF.B.5**
FORWARD 1	Decompose functions into compositions	**HSF-BF.A.1.c**
FORWARD 2	Verify inverses by composition	**HSF-BF.B.4.b**

SET 76 • HSF-BF.A.1.c

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c

$f(x) = \sqrt{x}$ and $g(x) = x - 4$. Write $f(g(x))$ and $g(f(x))$ and give the domain of each.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • MAIN • Q1/5

The temperature is $T(h) = 20 - 6.5h$ °C at height h km, and a balloon's height is $h(t) = 0.3t$ km after t minutes. Write $T(h(t))$ and find the temperature at $t = 10$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • MAIN • Q2/5

$f(x) = 1/(x - 1)$ and $g(x) = x^2$. Write $f(g(x))$ and give its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • MAIN • Q3/5

A dollar amount d converts to euros by e(d) = 0.9d, and a store's price with tax in euros is p(e) = 1.2e. Write p(e(d)) and find it for d = 50.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • MAIN • Q4/5

$f(x) = \log_2 x$ and $g(x) = x^2 - 9$. Write $f(g(x))$ and give its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • MAIN • Q5/5

SET 76 • HSF-BF.A.1.c

BACKWARD 1

Build combined models • High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b

$f(x) = x^2$ and $g(x) = x + 1$. Find $f(x) + g(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 76 • BACKWARD 1 • Q1/5

Find f(x) · g(x) for the same functions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 76 • BACKWARD 1 • Q2/5

True or false?

f(g(x)) is a different combination from f(x) · g(x).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 76 • BACKWARD 1 • Q3/5

For $f(x) = 2x$ and $g(x) = x + 3$, $f(x) - g(x)$ is

A. $x - 3$

B. $2x^2 + 6x$

C. $x + 3$

D. $3x + 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 76 • BACKWARD 1 • Q4/5

Name the operation in T(h(t)).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 76 • BACKWARD 1 • Q5/5

SET 76 • HSF-BF.A.1.c

BACKWARD 2

Evaluate a function at an expression • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

$f(x) = \sqrt{x}$. Find $f(x - 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 76 • BACKWARD 2 • Q1/5

$T(h) = 20 - 6.5h$. Find $T(0.3t)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 76 • BACKWARD 2 • Q2/5

True or false?

If $f(x) = 1/(x - 1)$, then $f(x^2) = 1/(x^2 - 1)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 76 • BACKWARD 2 • Q3/5

If $g(x) = x^2 - 9$, then $g(3)$ is

A. -3

B. -6

C. 9

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 76 • BACKWARD 2 • Q4/5

p(e) = 1.2e. Find p(45).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 76 • BACKWARD 2 • Q5/5

SET 76 • HSF-BF.A.1.c

BACKWARD 3

Domains of radical, rational, and log functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

Give the domain of $\sqrt{(x - 4)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 76 • BACKWARD 3 • Q1/5

Give the domain of $\log_2 u$ in terms of u.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 76 • BACKWARD 3 • Q2/5

True or false?

$1/(x^2 - 1)$ is undefined at $x = \pm 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 76 • BACKWARD 3 • Q3/5

$x^2 - 9 > 0$ when

A. $x < -3$ or $x > 3$

B. $-3 < x < 3$

C. all x

D. $x > 9$

Give the domain of $\sqrt{x} - 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 76 • BACKWARD 3 • Q5/5

SET 76 • HSF-BF.A.1.c

FORWARD 1

Decompose functions into compositions • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c

Write $h(x) = (3x + 1)^5$ as $f(g(x))$. Give f and g.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • FORWARD 1 • Q1/5

Write $h(x) = 2^{x^2}$ as $f(g(x))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • FORWARD 1 • Q2/5

True or false?

Every composition f(g(x)) can be decomposed in only one way.

A. True

B. False

If $f(x) = x + 1$, $g(x) = 2x$, $k(x) = x^2$, then $f(g(k(3)))$ is

A. 37

B. 19

C. 64

D. 13

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S76-F1-Q4**

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • FORWARD 1 • Q4/5

Write a context for a composition V(r(t)).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 76 • FORWARD 1 • Q5/5

SET 76 • HSF-BF.A.1.c

FORWARD 2

Verify inverses by composition • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b

Verify that $f(x) = 2x + 3$ and $g(x) = (x - 3)/2$ are inverses.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 76 • FORWARD 2 • Q1/5

Compute $f(g(x))$ for $f(x) = x^3$ and $g(x) = \sqrt[3]{x}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 76 • FORWARD 2 • Q2/5

True or false?

If f(g(x)) = x for all x in g's domain and g(f(x)) = x for all x in f's domain, then f and g are inverses.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 76 • FORWARD 2 • Q3/5

For $f(x) = x + 5$ and $g(x) = x - 5$, $f(g(2))$ is

A. 2

B. -3

C. 12

D. 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 76 • FORWARD 2 • Q4/5

Explain why composition is the natural test for inverses.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S76-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 76 • FORWARD 2 • Q5/5

SET 77 • HSF-BF.B.4.b • ADVANCED EXTENSION (+)

Verify inverses by composition on the stated domains

MAIN	Algebra II main question	**HSF-BF.B.4.b**
BACKWARD 1	Find inverses algebraically	**HSF-BF.B.4.a**
BACKWARD 2	Compose functions	**HSF-BF.A.1.c**
BACKWARD 3	Square roots of squares	**HSN-RN.A.2**
FORWARD 1	Show that proposed inverses fail	**HSF-BF.B.4.b**
FORWARD 2	Read inverse values from tables and graphs	**HSF-BF.B.4.c**

SET 77 • HSF-BF.B.4.b

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b

Verify by composition that $f(x) = 2x + 3$ and $g(x) = (x - 3)/2$ are inverses.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • MAIN • Q1/5

Verify by composition that $f(x) = x^3 - 1$ and $g(x) = \sqrt[3]{(x + 1)}$ are inverses.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • MAIN • Q2/5

Verify by composition that $f(x) = 2^x$ and $g(x) = \log_2 x$ are inverses, stating the domains.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • MAIN • Q3/5

Verify by composition that $f(x) = 1/(x - 2)$ $(x \neq 2)$ and $g(x) = 1/x + 2$ $(x \neq 0)$ are inverses.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • MAIN • Q4/5

Verify that $f(x) = x^2$ with domain $x \geq 0$ and $g(x) = \sqrt{x}$ are inverses. Explain why the domain restriction is needed.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • MAIN • Q5/5

SET 77 • HSF-BF.B.4.b

BACKWARD 1

Find inverses algebraically • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

Find the inverse of $f(x) = 2x + 3$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 77 • BACKWARD 1 • Q1/5

Find the inverse of $f(x) = x^3 - 1$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 77 • BACKWARD 1 • Q2/5

True or false?

The inverse of 2^x is $\log_2 x$.

A. True

B. False

The inverse of $f(x) = 1/(x - 2)$ is

A. $1/x + 2$

B. $1/(x + 2)$

C. $2 - 1/x$

D. $x - 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 77 • BACKWARD 1 • Q4/5

Find the inverse of $f(x) = x^2$ for $x \geq 0$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 77 • BACKWARD 1 • Q5/5

SET 77 • HSF-BF.B.4.b

BACKWARD 2

Compose functions • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c

$f(x) = 2x + 3$ and $g(x) = (x - 3)/2$. Compute $f(g(5))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 77 • BACKWARD 2 • Q1/5

Compute g(f(5)) for the same pair.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 77 • BACKWARD 2 • Q2/5

True or false?

f(g(x)) means apply g first, then f.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 77 • BACKWARD 2 • Q3/5

For $f(x) = x^3$ and $g(x) = x + 1$, $f(g(x))$ is

A. $(x + 1)^3$

B. $x^3 + 1$

C. $x + 1$

D. $3x + 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 77 • BACKWARD 2 • Q4/5

Compute $\log_2(2^5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 77 • BACKWARD 2 • Q5/5

SET 77 • HSF-BF.B.4.b

BACKWARD 3

Square roots of squares • High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2

Evaluate $\sqrt{((-3)^2)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B3-Q1

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 77 • BACKWARD 3 • Q1/5

Evaluate $(\sqrt{7})^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B3-Q2

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 77 • BACKWARD 3 • Q2/5

True or false?

$\sqrt{(x^2)} = x$ for every real x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B3-Q3

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 77 • BACKWARD 3 • Q3/5

$\sqrt{(x^2)}$ equals

A. $|x|$

B. x^2

C. $-x$

D. x

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B3-Q4

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 77 • BACKWARD 3 • Q4/5

For which x is $\sqrt{(x^2)} = x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-B3-Q5

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 77 • BACKWARD 3 • Q5/5

SET 77 • HSF-BF.B.4.b

FORWARD 1

Show that proposed inverses fail • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b

Show by composition that $f(x) = 3x - 1$ and $g(x) = (x - 1)/3$ are not inverses.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • FORWARD 1 • Q1/5

$f(x) = x^2$ (all real x) and $g(x) = \sqrt{x}$. Which composition fails to equal x, and for which x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • FORWARD 1 • Q2/5

True or false?

Checking f(g(x)) = x alone is always enough.

A. True

B. False

For $f(x) = e^x$ and $g(x) = \ln x$, $g(f(x)) =$

A. $|x|$

B. x only for $x > 0$

C. x for all real x

D. $e^{\ln x}$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • FORWARD 1 • Q4/5

Fix the domain of $f(x) = (x - 1)^2$ so that $g(x) = 1 + \sqrt{x}$ is its inverse.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 77 • FORWARD 1 • Q5/5

SET 77 • HSF-BF.B.4.b

FORWARD 2

Read inverse values from tables and graphs • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c

A table gives f(0) = 1, f(1) = 3, f(2) = 7, f(3) = 15. Find $f^{-1}(7)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 77 • FORWARD 2 • Q1/5

From the same table, find $f^{-1}(f^{-1}(15))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 77 • FORWARD 2 • Q2/5

True or false?

If (a, b) is on the graph of f, then (b, a) is on the graph of f^{-1}.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 77 • FORWARD 2 • Q3/5

If $f(4) = 9$, then $f^{-1}(9)$ is

A. 1/4

B. 4

C. 13

D. 9

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 77 • FORWARD 2 • Q4/5

Explain how $f(f^{-1}(7)) = 7$ shows a composition fact using the table.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S77-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 77 • FORWARD 2 • Q5/5

SET 78 • HSF-BF.B.4.c • ADVANCED EXTENSION (+)

Read values of an inverse function from graphs and tables

MAIN	Algebra II main question	**HSF-BF.B.4.c**
BACKWARD 1	Inverse functions swap inputs and outputs	**HSF-BF.B.4.a**
BACKWARD 2	Read points from graphs and tables	**8.F.A.1**
BACKWARD 3	One-to-one functions	**HSF-BF.B.4.b**
FORWARD 1	Estimate inverse values between data points and sketch inverse...	**HSF-BF.B.4.c**
FORWARD 2	Restrict the domain to make a function invertible	**HSF-BF.B.4.d**

SET 78 • HSF-BF.B.4.c

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c

The table shows f. Find $f^{-1}(15)$ and explain.

x	0	1	2	3	4
f(x)	1	3	7	15	31

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • MAIN • Q1/5

The graph of $f(x) = 2^x - 1$ is shown. Use it to find $f^{-1}(3)$.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • MAIN • Q2/5

The table shows a decreasing function g. Find $g^{-1}(10)$ and $g^{-1}(g(2))$.

x	1	2	3	4	5
g(x)	40	25	16	10	4

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • MAIN • Q3/5

The graph of $f(x) = \sqrt[3]{x} + 1$ is shown. Use it to find $f^{-1}(0)$ and $f^{-1}(3)$.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • MAIN • Q4/5

The table gives a plane's altitude A (m) t minutes after takeoff. Find $A^{-1}(1200)$ and interpret it.

t (min)	0	1	2	3	4	5
A (m)	0	250	550	880	1200	1500

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • MAIN • Q5/5

SET 78 • HSF-BF.B.4.c

BACKWARD 1

Inverse functions swap inputs and outputs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

If $f(3) = 15$, what is $f^{-1}(15)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 78 • BACKWARD 1 • Q1/5

Write the inverse of $f(x) = 2^x - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 78 • BACKWARD 1 • Q2/5

True or false?

$f^{-1}(y)$ is the input of f that gives the output y.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 78 • BACKWARD 1 • Q3/5

If (5, 4) is on the graph of g, which point is on g^{-1}?

A. (5, 4)

B. (1/5, 1/4)

C. (−5, −4)

D. (4, 5)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 78 • BACKWARD 1 • Q4/5

Use the formula to find $f^{-1}(3)$ for $f(x) = 2^x - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 78 • BACKWARD 1 • Q5/5

SET 78 • HSF-BF.B.4.c

BACKWARD 2

Read points from graphs and tables • Grade 8 • Functions • 8.F.A.1

In the table, what is f(2)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B2-Q1

Grade 8 • Functions • 8.F.A.1 • Set 78 • BACKWARD 2 • Q1/5

From the altitude table, what is A(3)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B2-Q2

Grade 8 • Functions • 8.F.A.1 • Set 78 • BACKWARD 2 • Q2/5

True or false?

A function gives exactly one output for each input.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B2-Q3

Grade 8 • Functions • 8.F.A.1 • Set 78 • BACKWARD 2 • Q3/5

Which input of g gives 16?

A. 16

B. 3

C. 4

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B2-Q4

Grade 8 • Functions • 8.F.A.1 • Set 78 • BACKWARD 2 • Q4/5

A graph passes through (8, 3). What is the output at input 8?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B2-Q5

Grade 8 • Functions • 8.F.A.1 • Set 78 • BACKWARD 2 • Q5/5

SET 78 • HSF-BF.B.4.c

BACKWARD 3

One-to-one functions • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b

Why does a strictly increasing function have an inverse?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B3-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 78 • BACKWARD 3 • Q1/5

Does a function with f(1) = 4 and f(3) = 4 have an inverse function?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B3-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 78 • BACKWARD 3 • Q2/5

True or false?

A decreasing function can have an inverse.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B3-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 78 • BACKWARD 3 • Q3/5

Which table has an inverse function?

A. x: 1, 2, 3; y: 5, 7, 5

B. x: 1, 2, 3; y: 5, 7, 9

C. x: 1, 2, 3; y: 0, 1, 0

D. x: 1, 2, 3; y: 2, 2, 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B3-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 78 • BACKWARD 3 • Q4/5

Verify $f(f^{-1}(15)) = 15$ with the table.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-B3-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 78 • BACKWARD 3 • Q5/5

SET 78 • HSF-BF.B.4.c

FORWARD 1

Estimate inverse values between data points and sketch inverse graphs • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c

Use the altitude table to estimate the time when the plane reaches 1000 m.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • FORWARD 1 • Q1/5

Sketch the graph of the inverse of $f(x) = 2^x - 1$ on the grid, using at least three points.



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • FORWARD 1 • Q2/5

True or false?

The graph of f^{-1} is the reflection of the graph of f across $y = x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • FORWARD 1 • Q3/5

Estimate $g^{-1}(20)$ from the decreasing table of g.

A. between 1 and 2

B. between 2 and 3

C. between 4 and 5

D. between 3 and 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • FORWARD 1 • Q4/5

Use the table of f to find $(f^{-1})^{-1}(2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.c • Set 78 • FORWARD 1 • Q5/5

SET 78 • HSF-BF.B.4.c

FORWARD 2

Restrict the domain to make a function invertible • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d

$f(x) = (x - 3)^2$ is not invertible. Give a domain restriction that makes it invertible, and write the inverse.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 78 • FORWARD 2 • Q1/5

A table of h shows $h(-2) = 4$, $h(0) = 0$, $h(2) = 4$. Explain why $h^{-1}(4)$ is not defined and how to fix it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 78 • FORWARD 2 • Q2/5

True or false?

Restricting the domain can turn a non-invertible function into an invertible one.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 78 • FORWARD 2 • Q3/5

$f(x) = x^2$ restricted to $x \leq 0$ has inverse

A. $\sqrt{x}$

B. x^2

C. $-\sqrt{x}$

D. $1/x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 78 • FORWARD 2 • Q4/5

Why must the restricted piece still take every output value of the original function?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S78-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 78 • FORWARD 2 • Q5/5

SET 79 • HSF-BF.B.4.d • ADVANCED EXTENSION (+)

Restrict a domain to make a function invertible and write the inverse

MAIN	Algebra II main question	**HSF-BF.B.4.d**
BACKWARD 1	Find the vertex of a quadratic	**HSF-IF.C.8.a**
BACKWARD 2	Write inverses of one-to-one functions	**HSF-BF.B.4.a**
BACKWARD 3	Domains and ranges	**HSF-IF.B.5**
FORWARD 1	Compare different restrictions	**HSF-BF.B.4.d**
FORWARD 2	Restrict trigonometric functions to define inverses	**HSF-TF.B.6**

SET 79 • HSF-BF.B.4.d

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d

Restrict $f(x) = (x - 3)^2$ to $x \geq 3$. Write f^{-1} and its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • MAIN • Q1/5

Restrict $f(x) = x^2 + 1$ to $x \le 0$. Write f^{-1} and its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • MAIN • Q2/5

Restrict $f(x) = |x + 2|$ to $x \geq -2$. Write f^{-1} and its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • MAIN • Q3/5

Restrict $f(x) = (x + 1)^2 - 4$ to $x \geq -1$. Write f^{-1} and its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • MAIN • Q4/5

Restrict $f(x) = x^4$ to $x \geq 0$. Write f^{-1} and its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • MAIN • Q5/5

SET 79 • HSF-BF.B.4.d

BACKWARD 1

Find the vertex of a quadratic • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.a

Give the vertex of $y = (x + 1)^2 - 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.a • Set 79 • BACKWARD 1 • Q1/5

Give the vertex of $y = x^2 + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.a • Set 79 • BACKWARD 1 • Q2/5

True or false?

A parabola is increasing on one side of its vertex and decreasing on the other.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.a • Set 79 • BACKWARD 1 • Q3/5

The vertex of $y = (x - 3)^2$ is

A. (0, 9)

B. (0, 3)

C. (−3, 0)

D. (3, 0)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.a • Set 79 • BACKWARD 1 • Q4/5

On which side of the vertex is $y = (x - 3)^2$ increasing?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.a • Set 79 • BACKWARD 1 • Q5/5

SET 79 • HSF-BF.B.4.d

BACKWARD 2

Write inverses of one-to-one functions • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

Solve $y = (x - 3)^2$ for x when $x \geq 3$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 79 • BACKWARD 2 • Q1/5

Solve $y = x^2 + 1$ for x when $x \leq 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 79 • BACKWARD 2 • Q2/5

True or false?

When $x \leq 0$, the solution of $x^2 = 4$ is $x = -2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 79 • BACKWARD 2 • Q3/5

Solving $y = x + 2$ $(x \geq -2)$ for x gives

A. $2 - y$

B. $y + 2$

C. $y - 2$

D. $-y - 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 79 • BACKWARD 2 • Q4/5

Solve $y = x^4$ for x when $x \geq 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 79 • BACKWARD 2 • Q5/5

SET 79 • HSF-BF.B.4.d

BACKWARD 3

Domains and ranges • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

Give the range of $f(x) = (x - 3)^2$ for $x \geq 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 79 • BACKWARD 3 • Q1/5

Give the range of $f(x) = (x + 1)^2 - 4$ for $x \geq -1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 79 • BACKWARD 3 • Q2/5

True or false?

The domain of the inverse is the range of the restricted function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 79 • BACKWARD 3 • Q3/5

The range of $x^2 + 1$ for $x \leq 0$ is

A. $y \geq 0$

B. $y \geq 1$

C. all reals

D. $y \leq 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 79 • BACKWARD 3 • Q4/5

Give the range of $|x + 2|$ for $x \geq -2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 79 • BACKWARD 3 • Q5/5

SET 79 • HSF-BF.B.4.d

FORWARD 1

Compare different restrictions • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d

Restrict $f(x) = (x - 3)^2$ to $x \leq 3$ instead. Write the inverse.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • FORWARD 1 • Q1/5

A ball's height is $h(t) = -5(t - 2)^2 + 20$ for $0 \le t \le 4$. Which restriction gives the time as a function of height on the way down? Write it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • FORWARD 1 • Q2/5

True or false?

A function can have more than one valid restriction that makes it invertible.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • FORWARD 1 • Q3/5

Which restriction does not make $f(x) = x^2$ invertible?

A. $x \geq 0$

B. $x \leq 0$

C. $-1 \leq x \leq 1$

D. $1 \leq x \leq 5$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • FORWARD 1 • Q4/5

Why is the restriction $x \geq 0$ natural for $f(x) = x^2$ in the area formula $A = s^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 79 • FORWARD 1 • Q5/5

SET 79 • HSF-BF.B.4.d

FORWARD 2

Restrict trigonometric functions to define inverses • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6

Why is sin x not invertible on all real numbers?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 79 • FORWARD 2 • Q1/5

Give the standard restricted domain of sin x and explain why it works.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 79 • FORWARD 2 • Q2/5

True or false?

cos x is restricted to $0 \leq x \leq \pi$ to define arccos.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 79 • FORWARD 2 • Q3/5

arcsin(1) equals

A. 0

B. 3π/2

C. π/2

D. π

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 79 • FORWARD 2 • Q4/5

Explain why $0 \leq x \leq \pi$ would not work for sine.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S79-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 79 • FORWARD 2 • Q5/5

SET 80 • HSF-BF.B.5 • ADVANCED EXTENSION (+)

Use the inverse relationship of exponents and logarithms to solve equations

MAIN	Algebra II main question	**HSF-BF.B.5**
BACKWARD 1	Write exact logarithmic solutions of exponential equations	**HSF-LE.A.4**
BACKWARD 2	Check for extraneous solutions	**HSA-REI.A.2**
BACKWARD 3	Domains of logarithmic expressions	**HSF-IF.B.5**
FORWARD 1	Use log properties and change of base in equations	**HSF-BF.B.5**
FORWARD 2	Check logarithmic solutions graphically	**HSA-REI.D.11**

SET 80 • HSF-BF.B.5

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5

Solve $\log_3 x = 4$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • MAIN • Q1/5

Solve $2\log_5(x - 1) = 4$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • MAIN • Q2/5

Solve $\ln(x + 2) = 1$ exactly.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • MAIN • Q3/5

Solve $\log_2 x + \log_2(x - 2) = 3$ and check for extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • MAIN • Q4/5

Solve $\log x - \log(x - 9) = 1$ (base 10) and check.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • MAIN • Q5/5

SET 80 • HSF-BF.B.5

BACKWARD 1

Write exact logarithmic solutions of exponential equations • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $3^x = 81$.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 80 • BACKWARD 1 • Q1/5

Solve $e^x = 5$ exactly.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 80 • BACKWARD 1 • Q2/5

True or false?

$\log_b y = x$ means $b^x = y$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 80 • BACKWARD 1 • Q3/5

$5^2 = 25$ written as a logarithm is

A. $\log_5 2 = 25$

B. $\log_2 25 = 5$

C. $\log_{25} 5 = 2$

D. $\log_5 25 = 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 80 • BACKWARD 1 • Q4/5

Solve $10^x = 1000$.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 80 • BACKWARD 1 • Q5/5

SET 80 • HSF-BF.B.5

BACKWARD 2

Check for extraneous solutions • High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2

Check $x = -2$ in $\log_2 x + \log_2(x - 2) = 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B2-Q1

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 80 • BACKWARD 2 • Q1/5

Check x = 4 in the same equation.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B2-Q2

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 80 • BACKWARD 2 • Q2/5

True or false?

Combining logs can introduce solutions outside the original domain.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B2-Q3

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 80 • BACKWARD 2 • Q3/5

Which value is excluded from $\log(x - 9)$?

A. $x = 20$

B. $x = 100$

C. $x = 5$

D. $x = 10$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B2-Q4

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 80 • BACKWARD 2 • Q4/5

Solve $x(x - 2) = 8$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B2-Q5

High School — Algebra II • Algebra: Reasoning with Equations & Inequalities • HSA-REI.A.2 • Set 80 • BACKWARD 2 • Q5/5

SET 80 • HSF-BF.B.5

BACKWARD 3

Domains of logarithmic expressions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

Give the domain of $\log_5(x - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 80 • BACKWARD 3 • Q1/5

Give the domain of ln(x + 2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 80 • BACKWARD 3 • Q2/5

True or false?

log 0 is undefined.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 80 • BACKWARD 3 • Q3/5

The domain of $\log_2 x + \log_2(x - 2)$ is

A. $x > 2$

B. $x \neq 2$

C. $x > 0$

D. all reals

Give the domain of $\log x - \log(x - 9)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 80 • BACKWARD 3 • Q5/5

SET 80 • HSF-BF.B.5

FORWARD 1

Use log properties and change of base in equations • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5

Solve $\log_2(x + 6) - \log_2 x = 2$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • FORWARD 1 • Q1/5

Solve $\log_4 x = \log_2 3$. (Hint: change of base.)

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • FORWARD 1 • Q2/5

True or false?

$\log_2(x + 6) - \log_2 x = \log_2((x + 6)/x)$ for $x > 0$.

A. True

B. False

$\log(x + 1) + \log(x - 1) = \log 8$ has solution

A. no solution

B. $x = 8$

C. $x = 3$

D. $x = -3$ and 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • FORWARD 1 • Q4/5

Explain why $\log_2(x - 5) = \log_2(1 - x)$ has no solution.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 80 • FORWARD 1 • Q5/5

SET 80 • HSF-BF.B.5

FORWARD 2

Check logarithmic solutions graphically • High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11

Explain how the graphs of $y = \log_2 x + \log_2(x - 2)$ and $y = 3$ show the solution $x = 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 80 • FORWARD 2 • Q1/5

Use a graph or table to approximate the solution of $\ln x = 2 - x$ to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 80 • FORWARD 2 • Q2/5

True or false?

An extraneous solution of a log equation corresponds to no intersection point.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 80 • FORWARD 2 • Q3/5

The graphs of $y = \log x$ and $y = 1$ meet at

A. $x = e$

B. $x = 10$

C. $x = 1$

D. $x = 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S80-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 80 • FORWARD 2 • Q4/5

Why can $\ln x = 2 - x$ not be solved exactly with logarithm rules?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S80-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Reasoning with Equations & Inequalities • HSA-REI.D.11 • Set 80 • FORWARD 2 • Q5/5

SET 81 • HSF-TF.A.3 • ADVANCED EXTENSION (+)

Derive values from special triangles and relate π − x, π + x, 2π − x to x

MAIN	Algebra II main question	**HSF-TF.A.3**
BACKWARD 1	Special right triangle side ratios	**HSG-SRT.C.6**
BACKWARD 2	Unit-circle values in all quadrants	**HSF-TF.A.2**
BACKWARD 3	The Pythagorean Theorem	**8.G.B.7**
FORWARD 1	Derive more special values and the reflection identities	**HSF-TF.A.3**
FORWARD 2	Explain symmetry and periodicity	**HSF-TF.A.4**

SET 81 • HSF-TF.A.3

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3

Use a 30°−60°−90° triangle with hypotenuse 1 to derive sin(π/3) and cos(π/3).

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • MAIN • Q1/5

Use a 45°−45°−90° triangle to derive tan(π/4).

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • MAIN • Q2/5

Use $\sin(\pi - x) = \sin x$ to find $\sin(5\pi/6)$ exactly, and explain the identity with the unit circle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • MAIN • Q3/5

Use $\cos(\pi + x) = -\cos x$ to find $\cos(7\pi/6)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • MAIN • Q4/5

Use $\tan(2\pi - x) = -\tan x$ to find $\tan(5\pi/3)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • MAIN • Q5/5

SET 81 • HSF-TF.A.3

BACKWARD 1

Special right triangle side ratios • High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6

Give the sides of a $30°-60°-90°$ triangle with hypotenuse 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B1-Q1

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 81 • BACKWARD 1 • Q1/5

Give the hypotenuse of a 45°−45°−90° triangle with legs 1.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B1-Q2

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 81 • BACKWARD 1 • Q2/5

True or false?

In a 30°−60°−90° triangle, the shortest side is half the hypotenuse.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B1-Q3

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 81 • BACKWARD 1 • Q3/5

cos 30° equals

A. $\sqrt{3}$

B. $\sqrt{2}/2$

C. $1/2$

D. $\sqrt{3}/2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S81-B1-Q4**

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 81 • BACKWARD 1 • Q4/5

Give tan 60°.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B1-Q5

High School — Geometry (course placement in this bank) • Geometry: Similarity, Right Triangles & Trigonometry • HSG-SRT.C.6 • Set 81 • BACKWARD 1 • Q5/5

SET 81 • HSF-TF.A.3

BACKWARD 2

Unit-circle values in all quadrants • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Give $\cos(5\pi/6)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 81 • BACKWARD 2 • Q1/5

Give $\sin(7\pi/6)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 81 • BACKWARD 2 • Q2/5

True or false?

sin is positive in quadrant II.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 81 • BACKWARD 2 • Q3/5

Which quadrant contains 5π/3?

A. II

B. I

C. IV

D. III

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 81 • BACKWARD 2 • Q4/5

Give $\tan(\pi/3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 81 • BACKWARD 2 • Q5/5

SET 81 • HSF-TF.A.3

BACKWARD 3

The Pythagorean Theorem • Grade 8 • Geometry • 8.G.B.7

A right triangle has hypotenuse 1 and leg 1/2. Find the other leg.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B3-Q1

Grade 8 • Geometry • 8.G.B.7 • Set 81 • BACKWARD 3 • Q1/5

A right isosceles triangle has hypotenuse 1. Find each leg.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B3-Q2

Grade 8 • Geometry • 8.G.B.7 • Set 81 • BACKWARD 3 • Q2/5

True or false?

$(1/2)^2 + (\sqrt{3}/2)^2 = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B3-Q3

Grade 8 • Geometry • 8.G.B.7 • Set 81 • BACKWARD 3 • Q3/5

Legs 1 and 1 give hypotenuse

A. 2

B. $\sqrt{3}$

C. $\sqrt{2}$

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B3-Q4

Grade 8 • Geometry • 8.G.B.7 • Set 81 • BACKWARD 3 • Q4/5

Find x if $x^2 + x^2 = 1$ ($x > 0$).

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-B3-Q5

Grade 8 • Geometry • 8.G.B.7 • Set 81 • BACKWARD 3 • Q5/5

SET 81 • HSF-TF.A.3

FORWARD 1

Derive more special values and the reflection identities • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3

Derive tan(π/6) exactly from a special triangle.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • FORWARD 1 • Q1/5

Explain with the unit circle why $\cos(\pi - x) = -\cos x$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • FORWARD 1 • Q2/5

True or false?

$\sin(\pi + x) = -\sin x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • FORWARD 1 • Q3/5

$\cos(2\pi - x)$ equals

A. $\cos x$

B. $-\cos x$

C. $\sin x$

D. $-\sin x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S81-F1-Q4**

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • FORWARD 1 • Q4/5

Find $\sin(11\pi/6)$ using one of the relations.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.3 • Set 81 • FORWARD 1 • Q5/5

SET 81 • HSF-TF.A.3

FORWARD 2

Explain symmetry and periodicity • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4

Explain why $\sin(-x) = -\sin x$ with the unit circle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 81 • FORWARD 2 • Q1/5

Is cos an even or odd function? Justify.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 81 • FORWARD 2 • Q2/5

True or false?

$\sin(x + 2\pi) = \sin x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 81 • FORWARD 2 • Q3/5

$\tan(-\pi/4)$ equals

A. undefined

B. -1

C. 1

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 81 • FORWARD 2 • Q4/5

Use $2\pi - x = -x + 2\pi$ to explain why $\tan(2\pi - x) = -\tan x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S81-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 81 • FORWARD 2 • Q5/5

SET 82 • HSF-TF.A.4 • ADVANCED EXTENSION (+)

Explain the symmetry and periodicity of trigonometric functions with the unit circle

MAIN	Algebra II main question	**HSF-TF.A.4**
BACKWARD 1	Explain the unit-circle definition	**HSF-TF.A.2**
BACKWARD 2	Even and odd functions	**HSF-BF.B.3**
BACKWARD 3	Reflect points over the axes	**8.G.A.3**
FORWARD 1	Use periodicity and symmetry to evaluate and simplify	**HSF-TF.A.4**
FORWARD 2	Restrict trigonometric functions so they have inverses	**HSF-TF.B.6**

SET 82 • HSF-TF.A.4

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4

Use the unit circle to explain why $\sin(-x) = -\sin x$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • MAIN • Q1/5

Use the unit circle to explain why $\cos(x + 2\pi) = \cos x$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • MAIN • Q2/5

Use the unit circle to explain why $\tan(x + \pi) = \tan x$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • MAIN • Q3/5

Use symmetry to evaluate $\cos(-2\pi/3)$, and explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • MAIN • Q4/5

Is f(x) = sin x cos x even, odd, or neither? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • MAIN • Q5/5

SET 82 • HSF-TF.A.4

BACKWARD 1

Explain the unit-circle definition • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

What point represents angle x on the unit circle?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 82 • BACKWARD 1 • Q1/5

Which direction does a negative angle go?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 82 • BACKWARD 1 • Q2/5

True or false?

$\tan x = \sin x / \cos x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 82 • BACKWARD 1 • Q3/5

The point for angle π is

A. $(0, -1)$

B. $(-1, 0)$

C. $(0, 1)$

D. $(1, 0)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 82 • BACKWARD 1 • Q4/5

Give $\cos(2\pi/3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 82 • BACKWARD 1 • Q5/5

SET 82 • HSF-TF.A.4

BACKWARD 2

Even and odd functions • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

Define an even function.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 82 • BACKWARD 2 • Q1/5

Define an odd function.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 82 • BACKWARD 2 • Q2/5

True or false?

An odd function's graph is symmetric about the origin.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 82 • BACKWARD 2 • Q3/5

x^3 is

A. odd

B. neither

C. even

D. both

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 82 • BACKWARD 2 • Q4/5

Is the product of an odd and an even function odd or even?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 82 • BACKWARD 2 • Q5/5

SET 82 • HSF-TF.A.4

BACKWARD 3

Reflect points over the axes • Grade 8 • Geometry • 8.G.A.3

Reflect (a, b) over the x-axis.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B3-Q1

Grade 8 • Geometry • 8.G.A.3 • Set 82 • BACKWARD 3 • Q1/5

Rotate (a, b) by 180° about the origin.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B3-Q2

Grade 8 • Geometry • 8.G.A.3 • Set 82 • BACKWARD 3 • Q2/5

True or false?

Reflecting over the x-axis keeps the x-coordinate.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B3-Q3

Grade 8 • Geometry • 8.G.A.3 • Set 82 • BACKWARD 3 • Q3/5

Reflecting (0.6, 0.8) over the x-axis gives

A. (0.8, 0.6)

B. (−0.6, 0.8)

C. (−0.6, −0.8)

D. (0.6, −0.8)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B3-Q4

Grade 8 • Geometry • 8.G.A.3 • Set 82 • BACKWARD 3 • Q4/5

Rotate (√3/2, 1/2) by 180°.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-B3-Q5

Grade 8 • Geometry • 8.G.A.3 • Set 82 • BACKWARD 3 • Q5/5

SET 82 • HSF-TF.A.4

FORWARD 1

Use periodicity and symmetry to evaluate and simplify • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4

Find $\sin(17\pi/6)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • FORWARD 1 • Q1/5

Find $\tan(-11\pi/4)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • FORWARD 1 • Q2/5

True or false?

$\cos(x + \pi) = -\cos x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • FORWARD 1 • Q3/5

$\sin(x + \pi) + \sin x$ equals

A. 1

B. $\sin 2x$

C. $2 \sin x$

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • FORWARD 1 • Q4/5

Simplify $\cos(-x) + \cos(x + 2\pi)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 82 • FORWARD 1 • Q5/5

SET 82 • HSF-TF.A.4

FORWARD 2

Restrict trigonometric functions so they have inverses • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6

Explain how periodicity prevents sin x from having an inverse on all real numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 82 • FORWARD 2 • Q1/5

Give the restricted domain used for tan x and explain it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 82 • FORWARD 2 • Q2/5

True or false?

arccos gives values between 0 and π.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 82 • FORWARD 2 • Q3/5

arctan(−1) equals

A. 7π/4

B. 3π/4

C. −π/4

D. π/4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 82 • FORWARD 2 • Q4/5

Use oddness to explain why $\arcsin(-1/2) = -\arcsin(1/2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S82-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 82 • FORWARD 2 • Q5/5

SET 83 • HSF-TF.B.6 • ADVANCED EXTENSION (+)

Choose and justify domain restrictions that make trigonometric functions invertible

MAIN	Algebra II main question	**HSF-TF.B.6**
BACKWARD 1	Restrict a domain to make a function invertible	**HSF-BF.B.4.d**
BACKWARD 2	Exact values on the unit circle	**HSF-TF.A.2**
BACKWARD 3	Periodicity of trigonometric functions	**HSF-TF.A.4**
FORWARD 1	Compose trigonometric functions with their inverses	**HSF-TF.B.6**
FORWARD 2	Solve trigonometric equations with inverses	**HSF-TF.B.7**

SET 83 • HSF-TF.B.6

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6

Explain why sin x has no inverse on all real numbers, and give a restricted domain on which it does.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • MAIN • Q1/5

Give the standard restricted domain of cos x and justify it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • MAIN • Q2/5

Give the standard restricted domain of tan x and justify it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • MAIN • Q3/5

Evaluate arcsin(1/2) and explain why it is not 5π/6.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • MAIN • Q4/5

Evaluate $\arccos(-1/2)$ exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • MAIN • Q5/5

SET 83 • HSF-TF.B.6

BACKWARD 1

Restrict a domain to make a function invertible • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d

Give a domain on which $f(x) = x^2$ is invertible.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 83 • BACKWARD 1 • Q1/5

Why does the restricted function need to be increasing or decreasing?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 83 • BACKWARD 1 • Q2/5

True or false?

A function that is always increasing on an interval is one-to-one there.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 83 • BACKWARD 1 • Q3/5

$f(x) = (x - 1)^2$ is one-to-one on

A. $-1 \leq x \leq 3$

B. all reals

C. $x \geq 1$

D. $x \geq 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 83 • BACKWARD 1 • Q4/5

Write the inverse of x^2 on $x \geq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.d • Set 83 • BACKWARD 1 • Q5/5

SET 83 • HSF-TF.B.6

BACKWARD 2

Exact values on the unit circle • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Give $\sin(\pi/6)$ and $\sin(5\pi/6)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 83 • BACKWARD 2 • Q1/5

Give $\cos(2\pi/3)$ and $\cos(4\pi/3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 83 • BACKWARD 2 • Q2/5

True or false?

$\tan(\pi/4) = \tan(5\pi/4)$.

A. True

B. False

Which angle in [0, π] has cos = −1/2?

A. 2π/3

B. π/3

C. 4π/3

D. 5π/6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 83 • BACKWARD 2 • Q4/5

Give $\sin(-\pi/6)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 83 • BACKWARD 2 • Q5/5

SET 83 • HSF-TF.B.6

BACKWARD 3

Periodicity of trigonometric functions • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4

What is the period of sin x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B3-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 83 • BACKWARD 3 • Q1/5

What is the period of tan x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B3-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 83 • BACKWARD 3 • Q2/5

True or false?

A periodic function cannot be one-to-one on its whole domain.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B3-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 83 • BACKWARD 3 • Q3/5

$\sin x = 0$ for

A. $x = \pi/2 + k\pi$

B. no x

C. $x = 2k\pi$ only

D. $x = k\pi$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B3-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 83 • BACKWARD 3 • Q4/5

Name two angles with cos = 1.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-B3-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 83 • BACKWARD 3 • Q5/5

SET 83 • HSF-TF.B.6

FORWARD 1

Compose trigonometric functions with their inverses • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6

Evaluate $\arcsin(\sin(5\pi/6))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • FORWARD 1 • Q1/5

Evaluate sin(arcsin(0.3)).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • FORWARD 1 • Q2/5

True or false?

arccos(cos x) = x for every real x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • FORWARD 1 • Q3/5

arctan(tan(3π/4)) equals

A. $-\pi/4$

B. $-3\pi/4$

C. $3\pi/4$

D. $\pi/4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • FORWARD 1 • Q4/5

Give the domain and range of arccos.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 83 • FORWARD 1 • Q5/5

SET 83 • HSF-TF.B.6

FORWARD 2

Solve trigonometric equations with inverses • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7

Solve $\sin x = 0.4$ on $0 \leq x < 2\pi$ to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 83 • FORWARD 2 • Q1/5

Solve $\cos x = 0.4$ on $0 \leq x < 2\pi$ to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 83 • FORWARD 2 • Q2/5

True or false?

arcsin gives only one of the solutions of sin x = c on [0, 2π).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 83 • FORWARD 2 • Q3/5

The second solution of $\sin x = c$ in $[0, 2\pi)$ (with $0 < c < 1$) is

A. $\pi + \arcsin c$

B. $-\arcsin c$

C. $\pi - \arcsin c$

D. $2\pi - \arcsin c$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 83 • FORWARD 2 • Q4/5

Explain how the restricted range of arccos is used to find all solutions of cos x = c.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S83-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 83 • FORWARD 2 • Q5/5

SET 84 • HSF-TF.B.7 • ADVANCED EXTENSION (+)

Solve trigonometric equations in models with inverse functions

MAIN	Algebra II main question	**HSF-TF.B.7**
BACKWARD 1	Evaluate inverse trigonometric functions on restricted domains	**HSF-TF.B.6**
BACKWARD 2	Parameters of sinusoidal models	**HSF-TF.B.5**
BACKWARD 3	Symmetry of unit-circle solutions	**HSF-TF.A.2**
FORWARD 1	Find all real solutions and solve equations that factor	**HSF-TF.B.7**
FORWARD 2	Solve equations with sum and double-angle identities	**HSF-TF.C.9**

SET 84 • HSF-TF.B.7

MAIN

Algebra II main question • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7

Solve $2\sin x - 1 = 0$ for $0 \leq x < 2\pi$. Give exact values.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-M-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • MAIN • Q1/5

Solve $3\cos x = -2$ for $0 \leq x < 2\pi$ to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-M-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • MAIN • Q2/5

A Ferris wheel rider's height is $h(t) = 12 + 10\sin(\pi t/15)$ m. Find every time in $0 \le t < 30$ s when $h = 17$ m.

$t =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-M-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • MAIN • Q3/5

Harbor depth is $d(t) = 5 + 4\cos(\pi t/6)$ ft. Find every time in $0 \le t < 12$ h when $d = 7$ ft.

$t =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-M-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • MAIN • Q4/5

Solve $\tan x = 3$ for $0 \leq x < 2\pi$ to the nearest hundredth. A calculator may be used.

$x \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-M-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • MAIN • Q5/5

SET 84 • HSF-TF.B.7

BACKWARD 1

Evaluate inverse trigonometric functions on restricted domains • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6

Evaluate arcsin(1/2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 84 • BACKWARD 1 • Q1/5

Evaluate arccos(1/2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 84 • BACKWARD 1 • Q2/5

True or false?

arctan 3 is between 0 and π/2.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 84 • BACKWARD 1 • Q3/5

arccos(−2/3) is in quadrant

A. II

B. III

C. I

D. IV

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B1-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 84 • BACKWARD 1 • Q4/5

Evaluate arctan 3 to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.6 • Set 84 • BACKWARD 1 • Q5/5

SET 84 • HSF-TF.B.7

BACKWARD 2

Parameters of sinusoidal models • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

In $h(t) = 12 + 10\sin(\pi t/15)$, what is the period?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 84 • BACKWARD 2 • Q1/5

Rewrite $h = 17$ as an equation for $\sin(\pi t/15)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 84 • BACKWARD 2 • Q2/5

True or false?

$d(t) = 5 + 4\cos(\pi t/6)$ has a maximum of 9.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 84 • BACKWARD 2 • Q3/5

Isolating the cosine in $5 + 4\cos(\pi t/6) = 7$ gives

A. $\cos(\pi t/6) = 1/2$

B. $\cos(\pi t/6) = 3$

C. $\cos(\pi t/6) = 2$

D. $\cos(\pi t/6) = 7/4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 84 • BACKWARD 2 • Q4/5

What is the midline of d(t)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 84 • BACKWARD 2 • Q5/5

SET 84 • HSF-TF.B.7

BACKWARD 3

Symmetry of unit-circle solutions • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

Give both angles in [0, 2π) with sin = 1/2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B3-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 84 • BACKWARD 3 • Q1/5

Give both angles in [0, 2π) with cos = 1/2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B3-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 84 • BACKWARD 3 • Q2/5

True or false?

$\tan x = c$ has two solutions in $[0, 2\pi)$ that differ by π.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B3-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 84 • BACKWARD 3 • Q3/5

If $\cos x = c$ has solution a in $[0, \pi]$, the other in $[0, 2\pi)$ is

A. $2\pi - a$

B. $\pi + a$

C. $-a$

D. $\pi - a$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B3-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 84 • BACKWARD 3 • Q4/5

Give both angles in $[0, 2\pi)$ with $\tan = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-B3-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 84 • BACKWARD 3 • Q5/5

SET 84 • HSF-TF.B.7

FORWARD 1

Find all real solutions and solve equations that factor • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7

Write all real solutions of $2\sin x - 1 = 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F1-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • FORWARD 1 • Q1/5

Solve $2\sin^2 x - \sin x = 0$ for $0 \leq x < 2\pi$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F1-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • FORWARD 1 • Q2/5

True or false?

$\cos x = 2$ has no real solution.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F1-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • FORWARD 1 • Q3/5

How many solutions does $\sin(2x) = 1/2$ have in $0 \leq x < 2\pi$?

A. 2

B. 4

C. 8

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S84-F1-Q4**

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • FORWARD 1 • Q4/5

Solve $\cos^2 x = 1/4$ for $0 \leq x < 2\pi$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F1-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.B.7 • Set 84 • FORWARD 1 • Q5/5

SET 84 • HSF-TF.B.7

FORWARD 2

Solve equations with sum and double-angle identities • High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9

Use $\sin 2x = 2\sin x \cos x$ to solve $\sin 2x = \cos x$ for $0 \leq x < 2\pi$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F2-Q1

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 84 • FORWARD 2 • Q1/5

Use $\cos 2x = 1 - 2\sin^2 x$ to solve $\cos 2x = \sin x$ for $0 \leq x < 2\pi$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F2-Q2

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 84 • FORWARD 2 • Q2/5

True or false?

$\sin(x + \pi/2) = \cos x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F2-Q3

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 84 • FORWARD 2 • Q3/5

cos 2x equals

A. $1 + 2\sin^2 x$

B. $\cos^2 x - \sin^2 x$

C. $\cos^2 x + \sin^2 x$

D. $2\cos x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S84-F2-Q4**

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 84 • FORWARD 2 • Q4/5

Explain why identities are useful for solving equations with different trigonometric functions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S84-F2-Q5

High School — Precalculus (course placement in this bank) • Functions: Trigonometric Functions • HSF-TF.C.9 • Set 84 • FORWARD 2 • Q5/5