ALGEBRA II · COMMON CORE HIGH SCHOOL MATH

Question Collection

Main • Backward Branches • Forward 1 • Forward 2 (later courses)

84 question sets • 2525 questions

CONTENTS

Set 1	**HSN-CN.A.1**	Write complex numbers in the form a + bi	7
Set 2	**HSN-CN.A.2**	Add and subtract complex numbers	44
Set 3	**HSN-CN.A.2**	Multiply complex numbers and simplify powers of i	81
Set 4	**HSN-CN.C.7**	Solve quadratic equations with complex solutions	118
Set 5	**HSN-CN.C.8**	Factor polynomials over the complex numbers	155
Set 6	**HSN-CN.C.9**	Use the Fundamental Theorem of Algebra	192
Set 7	**HSA-SSE.A.1.a**	Interpret the parts of polynomial, rational, and exponential expressions in context	229
Set 8	**HSA-SSE.A.1.b**	Interpret complicated expressions by viewing parts as single entities	266
Set 9	**HSA-SSE.A.2**	Factor higher-degree expressions by recognizing differences of squares and sums	303
Set 10	**HSA-SSE.A.2**	Rewrite expressions in quadratic form and factor	340
Set 11	**HSA-SSE.B.4**	Derive the formula for the sum of a finite geometric series	377
Set 12	**HSA-SSE.B.4**	Use the finite geometric series formula to solve problems	414
Set 13	**HSA-APR.A.1**	Add and subtract polynomials of higher degree	451
Set 14	**HSA-APR.A.1**	Multiply polynomials of higher degree	488
Set 15	**HSA-APR.B.2**	Find remainders with the Remainder Theorem	525
Set 16	**HSA-APR.B.2**	Use p(a) = 0 to find factors	562
Set 17	**HSA-APR.B.3**	Identify zeros and multiplicities from a factorization	599

CONTENTS (continued)

Set 18	**HSA-APR.B.3**	Construct a rough graph of a polynomial from its zeros	636
Set 19	**HSA-APR.C.4**	Prove polynomial identities	679
Set 20	**HSA-APR.C.4**	Use polynomial identities to describe numerical relationships	716
Set 21	**HSA-APR.C.5**	Expand with the Binomial Theorem	753
Set 22	**HSA-APR.D.6**	Rewrite rational expressions by polynomial division	790
Set 23	**HSA-APR.D.6**	Rewrite simple rational expressions by inspection	827
Set 24	**HSA-APR.D.7**	Multiply and divide rational expressions	864
Set 25	**HSA-APR.D.7**	Add and subtract rational expressions	901
Set 26	**HSA-CED.A.1**	Create and solve rational, radical, and exponential equations from contexts	938
Set 27	**HSA-CED.A.1**	Create and solve nonlinear inequalities from contexts	975
Set 28	**HSA-CED.A.2**	Create and graph nonlinear two-variable models	1012
Set 29	**HSA-CED.A.3**	Represent nonlinear constraints and judge viable options	1049
Set 30	**HSA-CED.A.4**	Rearrange rational, radical, and exponential formulas	1086
Set 31	**HSA-REI.A.2**	Solve rational equations and identify extraneous solutions	1123
Set 32	**HSA-REI.A.2**	Solve radical equations and identify extraneous solutions	1160
Set 33	**HSA-REI.D.11**	Explain why intersections of nonlinear graphs solve f(x) = g(x)	1197
Set 34	**HSA-REI.D.11**	Approximate solutions of f(x) = g(x)	1234

CONTENTS (continued)

Set 35	**HSF-TF.A.1**	Use radian measure as arc length on the unit circle	1271
Set 36	**HSF-TF.A.2**	Find exact trigonometric values beyond the first quadrant	1308
Set 37	**HSF-TF.A.2**	Explain the extension of sine and cosine to all real numbers	1345
Set 38	**HSF-TF.B.5**	Write sinusoidal models with given amplitude, period, and midline	1382
Set 39	**HSF-TF.B.5**	Interpret the parameters of sinusoidal models	1419
Set 40	**HSF-TF.C.8**	Prove the Pythagorean identity	1456
Set 41	**HSF-TF.C.8**	Use the Pythagorean identity to find trigonometric values	1493
Set 42	**HSF-IF.B.4**	Interpret key features of polynomial and rational models	1530
Set 43	**HSF-IF.B.4**	Interpret periodic models from graphs and tables	1567
Set 44	**HSF-IF.B.5**	State domains of radical, rational, logarithmic, and polynomial models	1604
Set 45	**HSF-IF.B.6**	Calculate and interpret average rates of change of nonlinear models	1641
Set 46	**HSF-IF.C.7.b**	Graph square root and cube root functions with transformations	1678
Set 47	**HSF-IF.C.7.b**	Graph piecewise-defined and absolute value functions	1715
Set 48	**HSF-IF.C.7.c**	Graph polynomial functions showing zeros and end behavior	1752
Set 49	**HSF-IF.C.7.e**	Graph exponential functions with intercepts, asymptote, and end behavior	1789
Set 50	**HSF-IF.C.7.e**	Graph logarithmic functions with intercept, asymptote, and end behavior	1826
Set 51	**HSF-IF.C.7.e**	Graph trigonometric functions showing period, midline, and amplitude	1863

CONTENTS (continued)

Set 52	**HSF-IF.C.8.b**	Rewrite exponential functions to reveal rates over other time units	1900
Set 53	**HSF-IF.C.9**	Compare properties of Algebra II functions in different representations	1937
Set 54	**HSF-BF.A.1.b**	Build models by combining Algebra II function types	1974
Set 55	**HSF-BF.B.3**	Identify vertical shifts, vertical stretches, and horizontal shifts of Algebra II functio	2011
Set 56	**HSF-BF.B.3**	Horizontal stretches, compressions, and reflections f(kx)	2048
Set 57	**HSF-BF.B.3**	Recognize even and odd functions	2085
Set 58	**HSF-BF.B.4.a**	Write inverses of radical, rational, cubic, exponential, and logarithmic functions	2122
Set 59	**HSF-LE.A.4**	Write exact logarithmic solutions of exponential equations	2159
Set 60	**HSF-LE.A.4**	Evaluate logarithmic solutions of exponential models with technology	2196
Set 61	**HSS-ID.A.4**	Estimate population percentages with a normal distribution	2233
Set 62	**HSS-ID.A.4**	Recognize data sets for which a normal model is not appropriate	2270
Set 63	**HSS-IC.A.1**	Distinguish population, sample, parameter, and statistic	2307
Set 64	**HSS-IC.A.2**	Decide whether a model is consistent with observed results using a simulation	2344
Set 65	**HSS-IC.B.3**	Identify sample surveys, experiments, and observational studies	2381
Set 66	**HSS-IC.B.3**	Explain random sampling versus random assignment	2418
Set 67	**HSS-IC.B.4**	Estimate a population proportion with a margin of error from a simulation	2455
Set 68	**HSS-IC.B.4**	Estimate a population mean with a margin of error from a simulation	2492

CONTENTS (continued)

Set 69	**HSS-IC.B.5**	Compare two treatments with a re-randomization simulation	2529
Set 70	**HSS-IC.B.6**	Evaluate claims in reports based on data	2566
Set 71	**HSS-MD.B.6**	Design and judge fair decision procedures with probabilities	2603
Set 72	**HSS-MD.B.7**	Analyze decisions and strategies with probabilities	2640
Set 73	**HSN-CN.A.3**	Use conjugates to find moduli and quotients of complex numbers	2677
Set 74	**HSF-IF.C.7.d**	Graph rational functions with zeros and asymptotes	2714
Set 75	**HSF-IF.C.7.d**	Distinguish holes from vertical asymptotes	2751
Set 76	**HSF-BF.A.1.c**	Compose functions and interpret compositions with domains	2788
Set 77	**HSF-BF.B.4.b**	Verify inverses by composition on the stated domains	2825
Set 78	**HSF-BF.B.4.c**	Read values of an inverse function from graphs and tables	2862
Set 79	**HSF-BF.B.4.d**	Restrict a domain to make a function invertible and write the inverse	2899
Set 80	**HSF-BF.B.5**	Use the inverse relationship of exponents and logarithms to solve equations	2936
Set 81	**HSF-TF.A.3**	Derive values from special triangles and relate π - x, π + x, 2π - x to x	2973
Set 82	**HSF-TF.A.4**	Explain the symmetry and periodicity of trigonometric functions with the unit circ	3010
Set 83	**HSF-TF.B.6**	Choose and justify domain restrictions that make trigonometric functions invertib	3047
Set 84	**HSF-TF.B.7**	Solve trigonometric equations in models with inverse functions	3084

Write complex numbers in the form a + bi

MAIN	Algebra II main question	**HSN-CN.A.1**
BACKWARD 1	Simplify square roots of positive numbers	**HSN-RN.A.2**
BACKWARD 2	No real number squares to a negative number	**8.EE.A.2**
BACKWARD 3	Divide each term of a sum by a number	**6.EE.A.3**
FORWARD 1	Use equality of complex numbers to find unknown real values	**HSN-CN.A.1**
FORWARD 2	Represent complex numbers as points in the complex plane	**HSN-CN.B.4**

SET 1 • HSN-CN.A.1

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Write $\sqrt{(-49)}$ in the form bi, where b is a real number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q1/5

Write $\sqrt{(-18)}$ in the form bi with b in simplest radical form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q2/5

Write $6 - \sqrt{(-20)}$ in the form a + bi. Then state its real part and its imaginary part.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q3/5

Write $(-4 + \sqrt{(-75)}) / 2$ in the form a + bi. Then state its real part and its imaginary part.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q4/5

Write $(10 + \sqrt{(-64)}) / 4$ in the form a + bi. Then state its real part and its imaginary part.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • MAIN • Q5/5

SET 1 • HSN-CN.A.1

BACKWARD 1

Simplify square roots of positive numbers • High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2

Simplify $\sqrt{50}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q1

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q1/5

Simplify $\sqrt{72}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q2

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q2/5

Which is √12 in simplest radical form?

A. 4√3

B. 3√2

C. 6√2

D. 2√3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q3

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q3/5

True or false?

$\sqrt{(9 + 16)} = \sqrt{9} + \sqrt{16}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q4

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q4/5

Simplify $\sqrt{28}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B1-Q5

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.2 • Set 1 • BACKWARD 1 • Q5/5

SET 1 • HSN-CN.A.1

BACKWARD 2

No real number squares to a negative number • Grade 8 • Expressions & Equations • 8.EE.A.2

Find all real numbers x with $x^2 = 81$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q1/5

True or false?

$(-3)^2 = -9$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q2/5

Is there a real number x with $x^2 = -4$? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q3/5

Which number has a square equal to 36?

A. 18

B. $\sqrt{-36}$

C. -6

D. -18

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q4/5

Evaluate $(-7)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 1 • BACKWARD 2 • Q5/5

SET 1 • HSN-CN.A.1

BACKWARD 3

Divide each term of a sum by a number • Grade 6 • Expressions & Equations • 6.EE.A.3

Write $(8 + 6\sqrt{2}) / 2$ as a sum of two terms.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q1/5

Write $(-6 + 4\sqrt{5}) / 2$ as a sum of two terms.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q2/5

True or false?

$(10 + \sqrt{7}) / 5 = 2 + \sqrt{7}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q3/5

$(12 - 9\sqrt{3}) / 3$ equals

A. $4 - 3\sqrt{3}$

B. $4 - 9\sqrt{3}$

C. $4 - 3\sqrt{1}$

D. $12 - 3\sqrt{3}$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q4/5

Write $(-4 + 2\sqrt{3}) / (-2)$ as a sum of two terms.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 1 • BACKWARD 3 • Q5/5

SET 1 • HSN-CN.A.1

FORWARD 1

Use equality of complex numbers to find unknown real values • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

x and y are real numbers and $(x - 3) + 2yi = 5 - 8i$. Find x and y.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q1/5

a and b are real numbers and $3a + (b + 1)i = -12 + 4i$. Find a and b.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q2/5

True or false?

If x and y are real and x + yi = 7, then y = 7.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q3/5

p and q are real and $(p + q) + (p - q)i = 10 + 2i$. What is p?

A. 6

B. 5

C. 4

D. 8

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q4/5

m is a real number. For what value of m is $(m^2 - 4) + (m + 3)i$ a real number? What real number is it then?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 1 • FORWARD 1 • Q5/5

SET 1 • HSN-CN.A.1

FORWARD 2

Represent complex numbers as points in the complex plane • High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4

In the complex plane the horizontal axis is real and the vertical axis is imaginary. Give the point for $-3 + 5i$.

Point: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q1

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q1/5

Which complex number is represented by the point $(4, -2)$ in the complex plane?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q2

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q2/5

Find the distance from 0 to the point 6 + 8i in the complex plane.

Distance: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q3

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q3/5

True or false?

Every real number lies on the horizontal axis of the complex plane.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S1-F2-Q4

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q4/5

Which number lies on the vertical axis of the complex plane?

A. -7

B. $7 + i$

C. $-7i$

D. $1 - i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S1-F2-Q5**

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.4 • Set 1 • FORWARD 2 • Q5/5

Add and subtract complex numbers

MAIN	Algebra II main question	**HSN-CN.A.2**
BACKWARD 1	Combine like terms	**7.EE.A.1**
BACKWARD 2	Add and subtract integers and decimals	**7.NS.A.1**
BACKWARD 3	Write square roots of negative numbers with i	**HSN-CN.A.1**
FORWARD 1	Find real parameters that make a sum real or imaginary	**HSN-CN.A.2**
FORWARD 2	Add complex numbers as vectors in the complex plane	**HSN-CN.B.5**

SET 2 • HSN-CN.A.2

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Simplify (3 + 5i) + (−7 + 2i). Write the answer in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q1/5

Simplify $(6 - 4i) - (-2 + 9i)$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q2/5

Simplify $(-1 + \sqrt{(-16)}) + (5 - \sqrt{(-9)})$. Write the answer in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q3/5

$z_1 = 2.5 - i$, $z_2 = -1.5 + 3i$ and $z_3 = 4i$. Find $z_1 - z_2 + z_3$ in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q4/5

Find the complex number w for which $(4 - 7i) + w = -2 + 3i$.

$w =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • MAIN • Q5/5

SET 2 • HSN-CN.A.2

BACKWARD 1

Combine like terms • Grade 7 • Expressions & Equations • 7.EE.A.1

Simplify $(3 + 5x) + (-7 + 2x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q1

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q1/5

Simplify $(6 - 4x) - (-2 + 9x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q2

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q2/5

$(2 - 3y) - (5 - y)$ equals

A. $-3 - 2y$

B. $7 - 4y$

C. $-3 + 2y$

D. $-3 - 4y$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q3

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q3/5

True or false?

$(4 + 2t) + (1 - 6t) = 5 + 4t$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q4

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q4/5

Simplify $(0.5 - x) - (1.5 + 2x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B1-Q5

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 2 • BACKWARD 1 • Q5/5

SET 2 • HSN-CN.A.2

BACKWARD 2

Add and subtract integers and decimals • Grade 7 • The Number System • 7.NS.A.1

Evaluate $6 - (-2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q1

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q1/5

Evaluate $-4 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q2

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q2/5

$-1 - 3 + 4$ equals

A. 8

B. 0

C. -8

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q3

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q3/5

True or false?

$2.5 - (-1.5) = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q4

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q4/5

Evaluate $-2 - 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B2-Q5

Grade 7 • The Number System • 7.NS.A.1 • Set 2 • BACKWARD 2 • Q5/5

SET 2 • HSN-CN.A.2

BACKWARD 3

Write square roots of negative numbers with i • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Write $\sqrt{(-16)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q1/5

Write $\sqrt{(-9)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q2/5

$\sqrt{(-25)}$ equals

A. -5

B. $25i$

C. $-5i$

D. $5i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S2-B3-Q3**

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q3/5

True or false?

$\sqrt{(-1)} = i$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q4/5

Write $-\sqrt{(-36)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-B3-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 2 • BACKWARD 3 • Q5/5

SET 2 • HSN-CN.A.2

FORWARD 1

Find real parameters that make a sum real or imaginary • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

k is a real number. For what value of k is $(3 + ki) + (k - 5i)$ a real number? Give that real number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q1/5

k is real. For what value of k is
$(2k - 6) + 4i - (k + 1 - i)$ a pure imaginary number (real part 0)?

$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q2/5

True or false?

The sum of a complex number and its conjugate is always 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q3/5

a is real and $(a + 3i) - (2 + ai)$ is real. What is a?

A. 0

B. 2

C. 3

D. -3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q4/5

Explain why the sum of a + bi and a − bi is always a real number when a and b are real.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 2 • FORWARD 1 • Q5/5

SET 2 • HSN-CN.A.2

FORWARD 2

Add complex numbers as vectors in the complex plane • High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5

Plot 2 + i and 1 + 3i as points in the complex plane (real axis horizontal). What point represents their sum?

Sum point: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q1

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q1/5

The points 0, 2 + i, 1 + 3i and their sum are vertices of a quadrilateral. What kind of quadrilateral is it?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q2

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q2/5

True or false?

Subtracting w from z gives the vector from the point w to the point z.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q3

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q3/5

What is the distance between the points $5 + 2i$ and $1 - i$ in the complex plane?

A. 25

B. $\sqrt{13}$

C. 5

D. 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S2-F2-Q4**

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q4/5

Find the midpoint of the segment joining $6 - 2i$ and $-2 + 4i$, as a complex number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S2-F2-Q5

High School — Precalculus (course placement in this bank) • Number & Quantity: The Complex Number System • HSN-CN.B.5 • Set 2 • FORWARD 2 • Q5/5

Multiply complex numbers and simplify powers of i

MAIN	Algebra II main question	**HSN-CN.A.2**
BACKWARD 1	Multiply binomials	**HSA-APR.A.1**
BACKWARD 2	Use $i^2 = -1$ and the cycle of powers of i	**HSN-CN.A.1**
BACKWARD 3	Divide whole numbers to find a remainder	**4.NBT.B.6**
FORWARD 1	Find a real parameter that makes a product real	**HSN-CN.A.2**
FORWARD 2	Divide complex numbers using conjugates	**HSN-CN.A.3**

SET 3 • HSN-CN.A.2

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Multiply $(3 + 2i)(4 - 5i)$. Write the product in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q1/5

Simplify $(-2 + i)^2$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q2/5

Multiply $(5 - 3i)(5 + 3i)$. Write the product in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q3/5

Simplify $(1 + i)^3$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q4/5

Simplify $i^{23} \cdot (4 - i)$. Write the answer in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • MAIN • Q5/5

SET 3 • HSN-CN.A.2

BACKWARD 1

Multiply binomials • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(3 + 2x)(4 - 5x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q1/5

Expand $(-2 + x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q2/5

$(5 - 3x)(5 + 3x)$ equals

A. $25 + 9x^2$

B. $10 - 9x^2$

C. $25 - 9x^2$

D. $25 - 30x - 9x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q3/5

True or false?

$(1 + x)^2 = 1 + x^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q4/5

Multiply $x(4 - x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 3 • BACKWARD 1 • Q5/5

SET 3 • HSN-CN.A.2

BACKWARD 2

Use $i^2 = -1$ and the cycle of powers of i • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Simplify i^2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q1/5

Simplify i^3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q2/5

Simplify i^4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q3/5

True or false?

$-10i^2 = -10$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q4/5

Which equals 1?

A. i^{8}

B. i^{9}

C. i^{3}

D. i^{6}

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 3 • BACKWARD 2 • Q5/5

SET 3 • HSN-CN.A.2

BACKWARD 3

Divide whole numbers to find a remainder • Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6

Find the remainder when 23 is divided by 4.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q1

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q1/5

Find the remainder when 50 is divided by 4.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q2

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q2/5

What is the remainder when 36 is divided by 4?

A. 0

B. 4

C. 1

D. 9

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q3

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q3/5

True or false?

The remainder when 101 is divided by 4 is 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q4

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q4/5

Write 39 as 4q + r with 0 ≤ r < 4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-B3-Q5

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 3 • BACKWARD 3 • Q5/5

SET 3 • HSN-CN.A.2

FORWARD 1

Find a real parameter that makes a product real • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

a is a real number. For what value of a is $(a + 2i)(3 - i)$ a real number? Give the product.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q1/5

b is real. For what value of b is (2 + bi)(1 + i) pure imaginary (real part 0)?

b = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q2/5

True or false?

The product of two non-real complex numbers can be a real number.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q3/5

For real c, $(c + i)^2$ is real when

A. $c = 0$

B. $c = 1$

C. $c = -1$

D. never

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q4/5

Show that $(a + bi)(a - bi)$ is a real number for all real a and b.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 3 • FORWARD 1 • Q5/5

SET 3 • HSN-CN.A.2

FORWARD 2

Divide complex numbers using conjugates • High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3

Write (3 + 4i) / (1 + 2i) in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q1

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q1/5

Write $10 / (3 - i)$ in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q2

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q2/5

True or false?

To divide by 2 − 5i, multiply numerator and denominator by 2 − 5i.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q3

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q3/5

(2 + i) / i equals

A. $-1 + 2i$

B. $2 + i$

C. $1 - 2i$

D. $1 + 2i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q4

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q4/5

Explain why multiplying by the conjugate of the denominator always gives a real denominator.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S3-F2-Q5

High School — Algebra II, Advanced Extension (+) • Number & Quantity: The Complex Number System • HSN-CN.A.3 • Set 3 • FORWARD 2 • Q5/5

SET 4 • HSN-CN.C.7 • THE COMPLEX NUMBER SYSTEM

Solve quadratic equations with complex solutions

MAIN	Algebra II main question	**HSN-CN.C.7**
BACKWARD 1	Use the quadratic formula and the discriminant	**HSA-REI.B.4.b**
BACKWARD 2	Write square roots of negative numbers with i	**HSN-CN.A.1**
BACKWARD 3	Solve equations of the form $(x - h)^2 = k$	**HSA-REI.B.4.b**
FORWARD 1	Find a quadratic from one complex solution	**HSN-CN.C.7**
FORWARD 2	Classify solutions for a family of quadratics	**HSN-CN.C.7**

SET 4 • HSN-CN.C.7

MAIN

Algebra II main question • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^2 + 49 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q1/5

Solve $x^2 - 4x + 13 = 0$ over the complex numbers. Give exact solutions in the form a + bi.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q2/5

Solve $2x^2 + 2x + 5 = 0$ over the complex numbers. Give exact solutions in the form a + bi.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q3/5

Solve $(x + 3)^2 = -20$ over the complex numbers. Give exact solutions in the form a + bi.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q4/5

Solve $3x^2 - 6x + 7 = 0$ over the complex numbers. Give exact solutions in the form a + bi.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-M-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • MAIN • Q5/5

SET 4 • HSN-CN.C.7

BACKWARD 1

Use the quadratic formula and the discriminant • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Find the discriminant $b^2 - 4ac$ of $x^2 - 4x + 13 = 0$.

Discriminant: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q1/5

Find the discriminant of $2x^2 + 2x + 5 = 0$.

Discriminant: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q2/5

True or false?

If the discriminant is negative, the quadratic equation has two real solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q3/5

In $3x^2 - 6x + 7 = 0$, the values of a, b and c are

A. 3, −6, −7

B. 3, 6, 7

C. −3, −6, 7

D. 3, −6, 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q4/5

Write the quadratic formula.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 1 • Q5/5

SET 4 • HSN-CN.C.7

BACKWARD 2

Write square roots of negative numbers with i • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1

Write $\sqrt{(-36)}$ in the form bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q1/5

Write $\sqrt{(-48)}$ in simplest form with i.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q2/5

$\sqrt{(-20)}$ equals

A. $2\sqrt{5}\,i$

B. $4\sqrt{5}\,i$

C. $-2\sqrt{5}$

D. $10i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q3/5

True or false?

$\sqrt{(-49)} = -7$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q4/5

Write $(-2 + \sqrt{(-36)}) / 4$ in the form a + bi.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.1 • Set 4 • BACKWARD 2 • Q5/5

SET 4 • HSN-CN.C.7

BACKWARD 3

Solve equations of the form $(x - h)^2 = k$ • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Solve $(x - 2)^2 = 9$ over the real numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q1/5

Solve $(x + 1)^2 = 5$ over the real numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q2/5

True or false?

Solving $x^2 = 16$ gives only $x = 4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q3/5

The solutions of $(x + 4)^2 = 25$ are

A. 5 and −5

B. −1 and 9

C. 1 and −9

D. 21 and −29

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q4/5

Solve $x^2 = 50$ over the real numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 4 • BACKWARD 3 • Q5/5

SET 4 • HSN-CN.C.7

FORWARD 1

Find a quadratic from one complex solution • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

$3 - 2i$ is a solution of $x^2 + bx + c = 0$, where b and c are real. Find b and c.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q1/5

$-1 + 4i$ is a solution of a quadratic equation $x^2 + bx + c = 0$ with real b and c. Write the equation.

Equation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q2/5

True or false?

If 2 + i is a solution of a quadratic with real coefficients, then 2 − i is also a solution.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q3/5

Which quadratic with real coefficients has 5i as a solution?

A. $x^2 + 25 = 0$

B. $x^2 + 5 = 0$

C. $x^2 - 25 = 0$

D. $x^2 - 10x + 25 = 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q4/5

Explain why $x^2 - (2 + i)x = 0$ has the solution $2 + i$ but not the solution $2 - i$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 1 • Q5/5

SET 4 • HSN-CN.C.7

FORWARD 2

Classify solutions for a family of quadratics • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

For which real values of k does $x^2 + kx + 9 = 0$ have two non-real solutions?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q1/5

For which real values of c does $x^2 - 4x + c = 0$ have non-real solutions? Give them for c = 8.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q2/5

True or false?

$x^2 + kx - 1 = 0$ has non-real solutions for some real k.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q3/5

$x^2 + 6x + m = 0$ has exactly one (repeated) real solution when $m =$

A. 3

B. 36

C. -9

D. 9

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q4/5

For $c > 4$, write the solutions of $x^2 - 4x + c = 0$ in terms of c.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S4-F2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 4 • FORWARD 2 • Q5/5

Factor polynomials over the complex numbers

MAIN	Algebra II main question	**HSN-CN.C.8**
BACKWARD 1	Factor a difference of squares	**HSA-SSE.A.2**
BACKWARD 2	Multiply complex conjugates	**HSN-CN.A.2**
BACKWARD 3	Solve quadratics with complex solutions	**HSN-CN.C.7**
FORWARD 1	Factor cubic and quartic polynomials over the complex numbers	**HSN-CN.C.8**
FORWARD 2	Write polynomials with real coefficients from complex zeros	**HSN-CN.C.9**

SET 5 • HSN-CN.C.8

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

Factor $x^2 + 25$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q1/5

Factor $x^2 + 7$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q2/5

Factor $4x^2 + 9$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q3/5

Factor $x^4 - 16$ completely into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q4/5

Factor $x^2 - 6x + 13$ into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • MAIN • Q5/5

SET 5 • HSN-CN.C.8

BACKWARD 1

Factor a difference of squares • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 - 49$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q1/5

Factor $9x^2 - 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q2/5

$x^4 - 1$ equals

A. $(x - 1)^4$

B. $(x^2 - 1)^2$

C. $(x^2 - 1)(x^2 + 1)$

D. $(x^2 + 1)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q3/5

True or false?

$x^2 + 9 = (x + 3)(x - 3)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q4/5

Factor $x^2 - 5$ using radicals.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 5 • BACKWARD 1 • Q5/5

SET 5 • HSN-CN.C.8

BACKWARD 2

Multiply complex conjugates • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Multiply $(x + 5i)(x - 5i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q1/5

Multiply $(2x + 3i)(2x - 3i)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q2/5

True or false?

$(3i)^2 = 9$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q3/5

$(x - (1 + i))(x - (1 - i))$ equals

A. $x^2 + 2x + 2$

B. $x^2 - 2x + 2$

C. $x^2 + 2$

D. $x^2 - 2x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q4/5

Simplify $(i\sqrt{7})^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 5 • BACKWARD 2 • Q5/5

SET 5 • HSN-CN.C.8

BACKWARD 3

Solve quadratics with complex solutions • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^2 + 25 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q1/5

Solve $x^2 - 6x + 13 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q2/5

True or false?

If r is a zero of p(x), then x − r is a factor of p(x).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q3/5

The zeros of $x^2 + 4$ are

A. $2i$ and $-2i$

B. $4i$ and $-4i$

C. -4

D. 2 and -2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q4/5

Solve $x^2 + 2x + 5 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-B3-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 5 • BACKWARD 3 • Q5/5

SET 5 • HSN-CN.C.8

FORWARD 1

Factor cubic and quartic polynomials over the complex numbers • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

Factor $x^3 + 8$ completely into linear factors with complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q1/5

Factor $x^3 + 9x$ completely into linear factors.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q2/5

Factor $x^4 + 5x^2 + 4$ completely into linear factors.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q3/5

True or false?

$x^3 - 1 = (x - 1)(x^2 + x + 1)$, and $x^2 + x + 1$ has two non-real zeros.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F1-Q4

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q4/5

Which is a linear factor of $x^3 - 27$ over the complex numbers?

A. $x + 3i$

B. $x + 3$

C. $x - 3i$

D. $x - (-3/2 + (3\sqrt{3}/2)i)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S5-F1-Q5**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 5 • FORWARD 1 • Q5/5

SET 5 • HSN-CN.C.8

FORWARD 2

Write polynomials with real coefficients from complex zeros • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9

Write a polynomial with real coefficients, leading coefficient 1 and least degree, whose zeros include $2i$ and $1 - i$.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q1/5

What is the least degree of a polynomial with real coefficients that has zeros 3, 4i and 2 + i?

Degree: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q2/5

True or false?

A polynomial with real coefficients can have exactly one non-real zero.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q3/5

A cubic with real coefficients has zeros 5 and $-i$. Its third zero is

A. -5

B. $5i$

C. 1

D. i

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S5-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q4/5

Explain why a polynomial with real coefficients and degree 3 must have at least one real zero.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S5-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 5 • FORWARD 2 • Q5/5

SET 6 • HSN-CN.C.9 • THE COMPLEX NUMBER SYSTEM

Use the Fundamental Theorem of Algebra

MAIN	Algebra II main question	**HSN-CN.C.9**
BACKWARD 1	Find the degree of a polynomial	**HSA-SSE.A.1.a**
BACKWARD 2	Find zeros from factored form	**HSA-SSE.B.3.a**
BACKWARD 3	Solve quadratics with complex solutions	**HSN-CN.C.7**
FORWARD 1	Relate the complex zeros to the real x-intercepts of a graph	**HSA-APR.B.3**
FORWARD 2	Prove existence statements about real zeros	**HSN-CN.C.9**

SET 6 • HSN-CN.C.9

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9

How many complex zeros, counted with multiplicity, does $p(x) = 3x^5 - x^2 + 7$ have? Name the theorem you used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q1/5

List every complex zero of $p(x) = (x - 2)^3(x^2 + 9)$ with its multiplicity, and check that the multiplicities add to the degree.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q2/5

Show that the Fundamental Theorem of Algebra holds for every quadratic $p(x) = ax^2 + bx + c$ with real a, b, c and $a \neq 0$: p has exactly two complex zeros counted with multiplicity.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q3/5

Find every complex zero of $p(x) = x^4 - 81$ and check that the count agrees with the Fundamental Theorem of Algebra.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q4/5

A student says $p(x) = x^3 + x + 1$ has no zeros because it does not factor with integers. Use the Fundamental Theorem of Algebra to explain how many complex zeros it has, and explain why at least one of them is real.

Work space

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • MAIN • Q5/5

SET 6 • HSN-CN.C.9

BACKWARD 1

Find the degree of a polynomial • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

What is the degree of $3x^5 - x^2 + 7$?

Degree: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q1/5

What is the degree of $(x - 2)^3(x^2 + 9)$ when multiplied out?

Degree: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q2/5

The degree of $x(x + 1)(x^2 - 4)$ is

A. 1

B. 4

C. 2

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q3/5

True or false?

The degree of $4x^3 + 2x^4$ is 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q4/5

What is the leading coefficient of $-2x^3 + 5x - 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 6 • BACKWARD 1 • Q5/5

SET 6 • HSN-CN.C.9

BACKWARD 2

Find zeros from factored form • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a

Find the zeros of $(x - 2)(x + 5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q1/5

Find the zeros of $(x - 1)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q2/5

True or false?

A product is 0 only when one of its factors is 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q3/5

The zeros of $x(x + 4)$ are

A. 0 and 4

B. 0 and -4

C. -4 only

D. 4 only

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q4/5

Find the zeros of $(2x - 3)(x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.B.3.a • Set 6 • BACKWARD 2 • Q5/5

SET 6 • HSN-CN.C.9

BACKWARD 3

Solve quadratics with complex solutions • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^2 + 9 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q1/5

Solve $x^2 + 2x + 2 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q2/5

True or false?

$x^2 + 1 = 0$ has no complex solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q3/5

The solutions of $x^2 - 2x + 5 = 0$ are

A. $1 \pm 4i$

B. $-1 \pm 2i$

C. $1 \pm 2i$

D. $2 \pm i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q4/5

Solve $x^2 = -81$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-B3-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 6 • BACKWARD 3 • Q5/5

SET 6 • HSN-CN.C.9

FORWARD 1

Relate the complex zeros to the real x-intercepts of a graph • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

A polynomial of degree 4 with real coefficients has a graph that crosses the x-axis at exactly two points and touches it nowhere else. Each crossing is a simple zero. How many non-real zeros does it have?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q1/5

The graph of $y = x^3 - 4x^2 + 5x - 2$ touches the x-axis at $x = 1$ and crosses it at $x = 2$. List all complex zeros with multiplicity.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q2/5

True or false?

A polynomial of degree 6 whose graph never meets the x-axis has 6 non-real zeros.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q3/5

A degree-5 polynomial with real coefficients has exactly one real zero, of multiplicity 1. How many non-real zeros does it have?

A. 4

B. 5

C. 2

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q4/5

Can a degree-4 polynomial with real coefficients have exactly three real zeros, each of multiplicity 1? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 6 • FORWARD 1 • Q5/5

SET 6 • HSN-CN.C.9

FORWARD 2

Prove existence statements about real zeros • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9

Prove that every polynomial of odd degree with real coefficients has at least one real zero.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S6-F2-Q1**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q1/5

A degree-4 polynomial with real coefficients has the zeros 1 + i and 3. What are all of its possible zero lists?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q2/5

True or false?

A degree-4 polynomial with real coefficients must have at least one real zero.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q3/5

A polynomial with real coefficients has degree 7. The possible numbers of real zeros (with multiplicity) are

A. 1, 3, 5, 7

B. 7 only

C. 0, 2, 4, 6

D. 0 to 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S6-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q4/5

Explain why the Fundamental Theorem of Algebra says nothing about whether a polynomial's zeros can be found exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S6-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.9 • Set 6 • FORWARD 2 • Q5/5

SET 7 • HSA-SSE.A.1.a • SEEING STRUCTURE IN EXPRESSIONS

Interpret the parts of polynomial, rational, and exponential expressions in context

MAIN	Algebra II main question	**HSA-SSE.A.1.a**
BACKWARD 1	Name terms, factors, and coefficients	**HSA-SSE.A.1.a**
BACKWARD 2	Translate between a percent change and a multiplying factor	**7.RP.A.3**
BACKWARD 3	Interpret a quotient as an amount per unit	**6.RP.A.2**
FORWARD 1	Build an expression whose parts have given meanings	**HSA-SSE.A.1.a**
FORWARD 2	Rewrite an exponential expression to reveal a different rate	**HSF-IF.C.8.b**

SET 7 • HSA-SSE.A.1.a

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

A town's population is modeled by $P(t) = 12{,}500(0.97)^t$, where t is the number of years after 2020. What do 12,500 and 0.97 represent, and what does $1 - 0.97$ represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q1/5

The average cost per item of making x items is $C(x) = (4{,}000 + 6x) / x$ dollars. Interpret 4,000, 6x, and the division by x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q2/5

An open box is made by cutting squares of side x cm from the corners of a 20 cm by 20 cm sheet. Its volume is $V(x) = x(20 - 2x)^2$ cm³. Interpret x, $20 - 2x$, and $(20 - 2x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q3/5

The amount of a medicine in the body t hours after a dose is $A(t) = 200(1/2)^{t/6}$ mg. Interpret 200, 1/2, and t/6.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q4/5

Maya keeps $1,500 in an account that earns 4% interest compounded once a year and also puts $200 per year in a jar (no interest). Her total after n years is $T(n) = 1500(1.04)^n + 200n$. Interpret $1500(1.04)^n$, 1.04, and 200n.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • MAIN • Q5/5

SET 7 • HSA-SSE.A.1.a

BACKWARD 1

Name terms, factors, and coefficients • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

Name the terms of 4,000 + 6x.

Terms: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q1/5

What is the coefficient of x in $7x - 3$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q2/5

In $5x(x - 2)$, which is a factor?

A. $5x - 2$

B. $x - 10$

C. $x - 2$

D. -2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q3/5

True or false?

In 3 + 2x, the number 3 is a factor.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q4/5

Name the factors of $200(0.5)^t$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • BACKWARD 1 • Q5/5

SET 7 • HSA-SSE.A.1.a

BACKWARD 2

Translate between a percent change and a multiplying factor • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

A price is multiplied by 0.97. By what percent does it change?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q1/5

What factor multiplies an amount that increases by 4%?

Factor: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q2/5

Multiplying by 1.25 is the same as

A. increasing by 2.5%

B. increasing by 25%

C. increasing by 125%

D. decreasing by 25%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q3/5

True or false?

Multiplying by 0.8 decreases an amount by 80%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q4/5

Find $1 - 0.85$ and explain what it means for an amount multiplied by 0.85.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B2-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 7 • BACKWARD 2 • Q5/5

SET 7 • HSA-SSE.A.1.a

BACKWARD 3

Interpret a quotient as an amount per unit • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2

A total cost of $600 is shared by 30 people. What does 600/30 represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q1/5

A car travels 150 miles in 3 hours. What does 150/3 represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q2/5

If T is a total cost for n items, then T/n is

A. the profit

B. the number of items

C. the cost per item

D. the total cost

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q3/5

True or false?

Dividing a total by the number of items gives an average per item.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q4/5

What does (4,000 + 6x)/x mean when x = 100?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.2 • Set 7 • BACKWARD 3 • Q5/5

SET 7 • HSA-SSE.A.1.a

FORWARD 1

Build an expression whose parts have given meanings • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

Write an expression for the value of a $24,000 car that loses 15% of its value each year, t years after purchase.

Expression: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q1/5

A gym charges a $90 joining fee and $35 per month. Write an expression for the average cost per month over m months.

Expression: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q2/5

True or false?

An expression for a quantity that triples every 5 years, starting at 40, is $40(3)^{5t}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q3/5

A bacteria count starts at 500 and doubles every 3 hours. Which expression gives the count after h hours?

A. $2(500)^{h/3}$

B. $500(2)^{3h}$

C. $500(3)^{h/2}$

D. $500(2)^{h/3}$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S7-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q4/5

A gym charges a $90 joining fee and $35 per month, so the average cost per month over m months is 35 + 90/m dollars. Explain what the part 90/m means.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 7 • FORWARD 1 • Q5/5

SET 7 • HSA-SSE.A.1.a

FORWARD 2

Rewrite an exponential expression to reveal a different rate • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

Rewrite $12{,}500(0.97)^{t}$ (t in years) to show the monthly factor, rounded to 5 decimal places.

Expression: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q1/5

$A(t) = 200(1/2)^{t/6}$. Write A as $200 \cdot b^t$ and give b to 4 decimal places. What is the hourly percent decrease?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q2/5

True or false?

$1.04^{n} = (1.04^{1/12})^{12n}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q3/5

Which expression shows that $500(2)^{t/3}$ grows by about 26% per time unit?

A. $1.26(500)^{t}$

B. $500(1.26)^{3t}$

C. $500(1.26)^{t}$

D. $500(2.26)^{t}$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S7-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q4/5

Explain why rewriting does not change the values of the function.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S7-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 7 • FORWARD 2 • Q5/5

SET 8 • HSA-SSE.A.1.b • SEEING STRUCTURE IN EXPRESSIONS

Interpret complicated expressions by viewing parts as single entities

MAIN	Algebra II main question	**HSA-SSE.A.1.b**
BACKWARD 1	Squares are never negative	**8.EE.A.2**
BACKWARD 2	Powers of numbers greater than 1 and between 0 and 1	**8.EE.A.1**
BACKWARD 3	A fraction gets smaller as its denominator grows	**3.NF.A.3.d**
FORWARD 1	Find the set of all values of an expression on a domain	**HSA-SSE.A.1.b**
FORWARD 2	Use single-entity reasoning to find horizontal asymptotes	**HSF-IF.C.7.d**

SET 8 • HSA-SSE.A.1.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b

$P(r) = 5000(1 + r)^{10}$ with $r > 0$ is the value of an investment. Without calculating, explain why $P(r) > 5000$.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q1/5

The height of a ball is $h(t) = 80 - 16(t - 2)^2$ feet after t seconds. Without expanding, find the greatest height and when it occurs, and explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q2/5

$C(x) = 10 + 40/(x + 1)$ is a cost in dollars for $x \geq 0$. Explain why $C(x)$ is always more than 10 and never more than 50.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q3/5

The temperature of a drink is $T(t) = 70 - 30(0.8)^t$ °F after t minutes. Explain why T(t) is always below 70 and gets closer to 70 as t grows.

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q4/5

$f(x) = 3/(x^2 + 1)$. Find the greatest value of $f(x)$ and explain why no input gives a greater value.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • MAIN • Q5/5

SET 8 • HSA-SSE.A.1.b

BACKWARD 1

Squares are never negative • Grade 8 • Expressions & Equations • 8.EE.A.2

True or false?

$(t - 2)^2 \geq 0$ for every real t.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q1/5

What is the smallest value of $(x + 5)^2$, and for which x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q2/5

Which expression can be negative for some real x?

A. $x^2 + 1$

B. $x - 1$

C. x^2

D. $(x - 1)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q3/5

What is the smallest value of $x^2 + 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q4/5

True or false?

$-(x - 3)^2$ is never positive.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B1-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 8 • BACKWARD 1 • Q5/5

SET 8 • HSA-SSE.A.1.b

BACKWARD 2

Powers of numbers greater than 1 and between 0 and 1 • Grade 8 • Expressions & Equations • 8.EE.A.1

Is 1.05^{10} greater than or less than 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q1/5

Is 0.8^5 greater than or less than 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q2/5

As n increases, 0.5^n

A. becomes negative

B. stays 0.5

C. gets closer to 0

D. grows without bound

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q3/5

True or false?

A negative number raised to any whole-number power is negative.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q4/5

Order from least to greatest: 0.9^2, 0.9^5, 0.9.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 8 • BACKWARD 2 • Q5/5

SET 8 • HSA-SSE.A.1.b

BACKWARD 3

A fraction gets smaller as its denominator grows • Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d

Which is greater, 40/2 or 40/5?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q1

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q1/5

True or false?

If $x \geq 0$, then $40/(x + 1) \leq 40$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q2

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q2/5

As x grows, 6/(x + 3)

A. increases

B. becomes negative

C. decreases toward 0

D. stays 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q3

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q3/5

What is the greatest value of 12/(x + 4) for x ≥ 0?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q4

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q4/5

True or false?

8/10 is greater than 8/9.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-B3-Q5

Grade 3 • Number & Operations–Fractions • 3.NF.A.3.d • Set 8 • BACKWARD 3 • Q5/5

SET 8 • HSA-SSE.A.1.b

FORWARD 1

Find the set of all values of an expression on a domain • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b

Find every value that $h(t) = 80 - 16(t - 2)^2$ takes for $0 \le t \le 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q1/5

Find every value of $f(x) = 3/(x^2 + 1)$ for real x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q2/5

True or false?

For $x \geq 0$, $C(x) = 10 + 40/(x + 1)$ takes every value in $10 < C \leq 50$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q3/5

For $t \geq 0$, which set contains every value of $70 - 30(0.8)^t$?

A. $T \geq 70$

B. $40 \leq T < 70$

C. $40 < T \leq 70$

D. $0 < T < 70$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S8-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q4/5

Find every value of $g(x) = 5 - 2/(x^2 + 4)$ for real x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 8 • FORWARD 1 • Q5/5

SET 8 • HSA-SSE.A.1.b

FORWARD 2

Use single-entity reasoning to find horizontal asymptotes • High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d

Explain why $y = 10 + 40/(x + 1)$ has the horizontal asymptote $y = 10$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q1/5

Rewrite (2x + 7)/(x + 3) as 2 + 1/(x + 3) and state its horizontal asymptote.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q2/5

True or false?

$y = 3/(x^2 + 1)$ has the horizontal asymptote $y = 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q3/5

The horizontal asymptote of $y = 5 - 2/(x - 4)$ is

A. $y = -2$

B. $y = 5$

C. $y = 4$

D. $y = 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q4/5

Does the graph of $y = 10 + 40/(x + 1)$ ever reach its horizontal asymptote? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S8-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Interpreting Functions • HSF-IF.C.7.d • Set 8 • FORWARD 2 • Q5/5

Factor higher-degree expressions by recognizing differences of squares and sums or differences of cubes

MAIN	Algebra II main question	**HSA-SSE.A.2**
BACKWARD 1	Take out a greatest common factor	**6.EE.A.3**
BACKWARD 2	Recognize perfect squares and perfect cubes	**8.EE.A.2**
BACKWARD 3	Multiply to check a factoring pattern	**HSA-APR.A.1**
FORWARD 1	Factor expressions whose parts are themselves expressions	**HSA-SSE.A.2**
FORWARD 2	Complete the factorization over the complex numbers	**HSN-CN.C.8**

SET 9 • HSA-SSE.A.2

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^4 - 81$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q1/5

Factor $8x^3 + 27$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q2/5

Factor $x^6 - 64$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q3/5

Factor $16x^4 - y^4$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q4/5

Factor $2x^5 - 32x$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • MAIN • Q5/5

SET 9 • HSA-SSE.A.2

BACKWARD 1

Take out a greatest common factor • Grade 6 • Expressions & Equations • 6.EE.A.3

Factor 6x + 18.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q1/5

Factor $2x^5 - 32x$ by taking out the greatest common factor only.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q2/5

The greatest common factor of $12x^3$ and $18x$ is

A. $6x^3$

B. $6x$

C. $3x$

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q3/5

True or false?

$3x^2 + 9x = 3x(x + 9)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q4/5

Factor $5y^2 - 20y$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B1-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 9 • BACKWARD 1 • Q5/5

SET 9 • HSA-SSE.A.2

BACKWARD 2

Recognize perfect squares and perfect cubes • Grade 8 • Expressions & Equations • 8.EE.A.2

Write $16x^4$ as a square.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q1/5

Write $8x^3$ as a cube.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q2/5

Which is a perfect cube?

A. 32

B. 16

C. 36

D. 64

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q3/5

True or false?

x^6 is both a perfect square and a perfect cube.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q4/5

Find the cube root of 27.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 9 • BACKWARD 2 • Q5/5

SET 9 • HSA-SSE.A.2

BACKWARD 3

Multiply to check a factoring pattern • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(a + b)(a^2 - ab + b^2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q1/5

Multiply $(x - 2)(x^2 + 2x + 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q2/5

True or false?

$(x + 3)(x^2 - 6x + 9) = x^3 + 27.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q3/5

$(x^2 - 9)(x^2 + 9)$ equals

A. $x^4 - 18x^2 - 81$

B. $x^4 + 81$

C. $(x - 3)^4$

D. $x^4 - 81$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S9-B3-Q4**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q4/5

Multiply $(2x + 1)(4x^2 - 2x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-B3-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 9 • BACKWARD 3 • Q5/5

SET 9 • HSA-SSE.A.2

FORWARD 1

Factor expressions whose parts are themselves expressions • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $(x + 1)^2 - 9$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q1/5

Factor $(2x - 1)^3 + 8$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q2/5

True or false?

$a^6 - b^6$ can be factored first as a difference of squares or first as a difference of cubes.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q3/5

$(x - y)^2 - (x + y)^2$ equals

A. 0

B. $4xy$

C. $-4xy$

D. $-2y^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S9-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q4/5

Factor $x^2(x + 2) - 4(x + 2)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 9 • FORWARD 1 • Q5/5

SET 9 • HSA-SSE.A.2

FORWARD 2

Complete the factorization over the complex numbers • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

Factor $x^4 - 81$ completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q1/5

Factor $16x^4 - 1$ completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q2/5

True or false?

Over the complex numbers, $x^3 + 8$ has only one linear factor.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q3/5

Which polynomial is fully factored over the integers but not over the complex numbers?

A. $x^2 + 9$

B. $x - 9$

C. $x^2 - 9$

D. $(x - 3)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q4

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q4/5

$2x^5 - 32x = 2x(x^2 + 4)(x + 2)(x - 2)$ over the integers.
Complete the factorization over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S9-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 9 • FORWARD 2 • Q5/5

SET 10 • HSA-SSE.A.2 • SEEING STRUCTURE IN EXPRESSIONS

Rewrite expressions in quadratic form and factor

MAIN	Algebra II main question	**HSA-SSE.A.2**
BACKWARD 1	Factor quadratic trinomials	**HSA-SSE.A.2**
BACKWARD 2	Write powers as squares of other powers	**8.EE.A.1**
BACKWARD 3	Treat a repeated part as a single variable	**HSA-SSE.A.1.b**
FORWARD 1	Solve equations in quadratic form over the complex numbers	**HSN-CN.C.7**
FORWARD 2	Solve exponential equations in quadratic form	**HSF-LE.A.4**

SET 10 • HSA-SSE.A.2

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^4 - 5x^2 + 4$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q1/5

Factor $x^4 + 3x^2 - 10$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q2/5

Factor $9^x - 4 \cdot 3^x + 3$ by writing it in quadratic form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q3/5

Factor $x^6 - 9x^3 + 8$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q4/5

Factor $2x^4 - 7x^2 - 4$ completely over the integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • MAIN • Q5/5

SET 10 • HSA-SSE.A.2

BACKWARD 1

Factor quadratic trinomials • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $u^2 - 5u + 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q1/5

Factor $u^2 + 3u - 10$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q2/5

$2u^2 - 7u - 4$ equals

A. $(2u + 1)(u - 4)$

B. $(u - 2)(2u + 2)$

C. $(2u + 4)(u - 1)$

D. $(2u - 1)(u + 4)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q3/5

True or false?

$u^2 - 9u + 8 = (u + 1)(u - 8)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q4/5

Factor $u^2 - 4u + 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 10 • BACKWARD 1 • Q5/5

SET 10 • HSA-SSE.A.2

BACKWARD 2

Write powers as squares of other powers • Grade 8 • Expressions & Equations • 8.EE.A.1

Write x^4 as a power of x^2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q1/5

Write 9^x as a power of 3^x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q2/5

x^6 equals

A. $(x^4)^2$

B. $(x^3)^2$

C. $x^3 \cdot x^3 \cdot x^3$

D. $(x^2)^4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q3/5

True or false?

$4^x = (2^x)^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q4/5

Write 25^x as a power of 5^x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 10 • BACKWARD 2 • Q5/5

SET 10 • HSA-SSE.A.2

BACKWARD 3

Treat a repeated part as a single variable • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b

In $x^4 - 5x^2 + 4$, what single variable u turns the expression into $u^2 - 5u + 4$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q1/5

Rewrite $2(x - 1)^2 + 3(x - 1) - 5$ using $u = x - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q2/5

True or false?

In $4^x - 2^x - 6$, the substitution $u = 2^x$ gives $u^2 - u - 6$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q3/5

Which substitution writes $x^6 - 9x^3 + 8$ as a quadratic?

A. $u = x^2$

B. $u = x^3$

C. $u = x^6$

D. $u = 3x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q4/5

After factoring $(u - 1)(u - 8)$ with $u = x^3$, write the result in terms of x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-B3-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.b • Set 10 • BACKWARD 3 • Q5/5

SET 10 • HSA-SSE.A.2

FORWARD 1

Solve equations in quadratic form over the complex numbers • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7

Solve $x^4 + 3x^2 - 4 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q1/5

Solve $x^4 - 5x^2 + 4 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q2/5

True or false?

$x^4 + 10x^2 + 9 = 0$ has two real solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q3/5

How many complex solutions, with multiplicity, does $x^4 - 13x^2 + 36 = 0$ have?

A. 1

B. 3

C. 4

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q4/5

Solve $x^4 + x^2 - 12 = 0$ over the complex numbers.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F1-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.C.7 • Set 10 • FORWARD 1 • Q5/5

SET 10 • HSA-SSE.A.2

FORWARD 2

Solve exponential equations in quadratic form • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

Solve $9^x - 5 \cdot 3^x + 6 = 0$. Give exact solutions.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q1/5

Solve $4^x - 6 \cdot 2^x + 8 = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q2/5

True or false?

$4^x + 2^x - 2 = 0$ has two real solutions.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q3/5

The solutions of $25^x - 26 \cdot 5^x + 25 = 0$ are

A. 0 and 2

B. 1 and 25

C. 0 and 1

D. 1 and 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q4/5

Solve $e^{2t} - 7e^{t} + 10 = 0$. Give exact solutions and decimal values to 3 decimal places.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S10-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 10 • FORWARD 2 • Q5/5

SET 11 • HSA-SSE.B.4 • SEEING STRUCTURE IN EXPRESSIONS

Derive the formula for the sum of a finite geometric series

MAIN	Algebra II main question	**HSA-SSE.B.4**
BACKWARD 1	Write terms of a geometric sequence	**HSF-LE.A.2**
BACKWARD 2	Distribute a factor over a sum	**6.EE.A.3**
BACKWARD 3	Use the product rule for exponents	**8.EE.A.1**
FORWARD 1	Prove the sum formula from a polynomial identity	**HSA-APR.C.4**
FORWARD 2	Describe how partial sums behave when the ratio is between -1 a...	**HSA-SSE.B.4**

SET 11 • HSA-SSE.B.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

Let $S_n = a + ar + ar^2 + \ldots + ar^{n-1}$ with $r \neq 1$. Derive $S_n = a(1 - r^n)/(1 - r)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q1/5

Use the multiply-and-subtract method (not the formula) to find $S = 3 + 6 + 12 + \ldots + 3 \cdot 2^9$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q2/5

Explain why the derivation of $S_n = a(1 - r^n)/(1 - r)$ does not work when $r = 1$, and find S_n in that case.

Work space

Explanation: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q3/5

Derive a formula for $T = 1 + x + x^2 + \ldots + x^n$ ($n + 1$ terms, $x \neq 1$), then check it for $x = 2$ and $n = 3$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q4/5

Derive the total of 20 yearly amounts $100, 100(1.05), 100(1.05)^2, \ldots, 100(1.05)^{19}$ by the multiply-and-subtract method, and give it to the nearest cent.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-M-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • MAIN • Q5/5

SET 11 • HSA-SSE.B.4

BACKWARD 1

Write terms of a geometric sequence • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

Write the first four terms of the geometric sequence with first term 3 and ratio 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q1/5

What is the 10th term of 3, 6, 12, ...?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q2/5

The nth term of a geometric sequence with first term a and ratio r is

A. $a + (n - 1)r$

B. $a \cdot n \cdot r$

C. ar^n

D. ar^{n-1}

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S11-B1-Q3**

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q3/5

True or false?

100, 105, 110.25, ... is arithmetic with common difference 5.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q4/5

Find the ratio of 81, 27, 9, 3, ...

Ratio: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B1-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 11 • BACKWARD 1 • Q5/5

SET 11 • HSA-SSE.B.4

BACKWARD 2

Distribute a factor over a sum • Grade 6 • Expressions & Equations • 6.EE.A.3

Distribute: 2(3 + 6 + 12).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q1/5

Distribute: $r(a + ar + ar^2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q2/5

True or false?

$x(1 + x + x^2) = x + x^2 + x^3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q3/5

1.05(100 + 105) equals

A. 105 + 105

B. 100 + 110.25

C. 105 + 110.25

D. 205

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q4/5

Subtract: $(6 + 12 + 24 + 48) - (3 + 6 + 12 + 24)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 11 • BACKWARD 2 • Q5/5

SET 11 • HSA-SSE.B.4

BACKWARD 3

Use the product rule for exponents • Grade 8 • Expressions & Equations • 8.EE.A.1

Simplify $r \cdot r^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q1/5

Simplify $ar^{n-1} \cdot r$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q2/5

$2 \cdot 2^9$ equals

A. 2^{10}

B. 4^{10}

C. 4^9

D. 2^9

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q3/5

True or false?

$x \cdot x^n = x^{2n}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q4/5

Evaluate 2^{10}.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-B3-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 11 • BACKWARD 3 • Q5/5

SET 11 • HSA-SSE.B.4

FORWARD 1

Prove the sum formula from a polynomial identity • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Prove the identity

$x^n - 1 = (x - 1)(x^{n-1} + x^{n-2} + \ldots + x + 1)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q1/5

Use the identity $x^n - 1 = (x - 1)(x^{n-1} + \ldots + 1)$ to write $1 + 3 + 9 + \ldots + 3^5$ as a quotient, and evaluate it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q2/5

True or false?

The identity shows that $x - 1$ is a factor of $x^n - 1$ for every whole number $n \geq 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q3/5

Which equals $1 + 2 + 4 + \ldots + 2^9$?

A. 2^{10}

B. $2^{10} - 1$

C. $2^9 - 1$

D. $(2^9 - 1)/2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q4/5

Explain how the identity gives $a(1 - r^n)/(1 - r)$ for $a + ar + \ldots + ar^{n-1}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 11 • FORWARD 1 • Q5/5

SET 11 • HSA-SSE.B.4

FORWARD 2

Describe how partial sums behave when the ratio is between −1 and 1 • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

For the series $8 + 4 + 2 + 1 + \ldots$, find S_5 and S_{10} to 4 decimal places.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q1/5

Using $S_n = 16(1 - 0.5^n)$, explain why every partial sum of 8 + 4 + 2 + ... is less than 16 but can be made as close to 16 as you like.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q2/5

True or false?

For r = 2, the partial sums of $a + ar + ar^2 + \ldots$ ($a > 0$) stay below a fixed number.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q3/5

For $10 - 5 + 2.5 - 1.25 + \ldots$, the partial sums get close to

A. 20/3

B. 15

C. 5

D. 20

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q4/5

A ball travels 10 m down, then rebounds and falls in a geometric pattern so the total distance after n bounces is $10 + 2 \cdot 6(1 - 0.6^n)/0.4$. What total can the distance never reach, and why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S11-F2-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 11 • FORWARD 2 • Q5/5

Use the finite geometric series formula to solve problems

MAIN	Algebra II main question	**HSA-SSE.B.4**
BACKWARD 1	Identify the first term, ratio, and number of terms	**HSF-LE.A.2**
BACKWARD 2	Change a percent to a growth factor	**7.RP.A.3**
BACKWARD 3	Round results of calculations correctly	**HSN-Q.A.3**
FORWARD 1	Find the number of terms with logarithms	**HSF-LE.A.4**
FORWARD 2	Compare two payment plans with the series formula	**HSA-SSE.B.4**

SET 12 • HSA-SSE.B.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

Find the sum of the first 12 terms of 5 + 15 + 45 + ...

Sum: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q1/5

Leah deposits $250 at the end of each month into an account that earns 0.5% interest per month. The balance just after the 36th deposit is $250 + 250(1.005) + \ldots + 250(1.005)^{35}$. Find it to the nearest cent.

Balance: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q2/5

A $180,000 loan is repaid with 360 equal monthly payments P at a monthly interest rate of 0.4%. P satisfies
$180{,}000 = P/1.004 + P/1.004^2 + \ldots + P/1.004^{360}$. Use the geometric series formula to find P to the nearest cent.

P = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q3/5

A ball is dropped from 10 m and each rebound reaches 60% of the previous height. Find the total vertical distance it has traveled when it hits the ground for the 6th time, to the nearest hundredth of a meter.

Distance: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q4/5

What is the least number of terms of 2 + 6 + 18 + ... whose sum is greater than 10,000? Show the sums that decide it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-M-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • MAIN • Q5/5

SET 12 • HSA-SSE.B.4

BACKWARD 1

Identify the first term, ratio, and number of terms • High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2

For 5 + 15 + 45 + ... (12 terms), give a, r, and n.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q1

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q1/5

For $250 + 250(1.005) + \ldots + 250(1.005)^{35}$, give a, r, and n.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q2

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q2/5

For 6 + 3.6 + 2.16 + 1.296 + 0.7776, r is

A. 0.4

B. 1.67

C. 0.6

D. 2.4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q3

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q3/5

True or false?

A list with exponents 0 through 35 has 35 terms.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q4

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q4/5

For $P/1.004 + P/1.004^2 + \ldots + P/1.004^{360}$, give the first term and the ratio.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B1-Q5

High School — Algebra I • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.2 • Set 12 • BACKWARD 1 • Q5/5

SET 12 • HSA-SSE.B.4

BACKWARD 2

Change a percent to a growth factor • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

What factor adds 0.5% interest?

Factor: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q1/5

A height is 60% of the one before. What is the ratio?

Ratio: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q2/5

0.4% per month as a factor is

A. 1.04

B. 1.004

C. 0.004

D. 1.4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q3/5

True or false?

Interest of 12% per year is a factor of 0.12.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q4/5

A loss of 15% per year is what factor?

Factor: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B2-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 12 • BACKWARD 2 • Q5/5

SET 12 • HSA-SSE.B.4

BACKWARD 3

Round results of calculations correctly • High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3

Round 9834.0262 to the nearest cent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q1

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q1/5

Round 37.6672 to the nearest hundredth.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q2

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q2/5

True or false?

Rounding 1.005 to 1.01 before raising it to the 36th power gives the same answer to the nearest cent.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q3

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q3/5

1.005^{36} is closest to

A. 1.1967

B. 1.18

C. 1.2

D. 1.036

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q4

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q4/5

Evaluate 3^{12} exactly.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-B3-Q5

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 12 • BACKWARD 3 • Q5/5

SET 12 • HSA-SSE.B.4

FORWARD 1

Find the number of terms with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

How many end-of-month deposits of $250 at 0.5% per month are needed for the balance to reach $20,000? Use $250(1.005^n - 1)/0.005 \geq 20{,}000$.

n = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q1/5

Solve $2(3^n - 1)/2 = 3^n - 1 = 1{,}000{,}000$ for n exactly, then give the least whole number of terms whose sum is at least 1,000,000.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q2/5

True or false?

For a ratio greater than 1, more terms always give a larger sum when $a > 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q3/5

The sum $5(2^n - 1)$ first exceeds 5,000 when n =

A. 9

B. 11

C. 10

D. 1000

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q4/5

Explain why the number of terms must be rounded up, not to the nearest whole number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 12 • FORWARD 1 • Q5/5

SET 12 • HSA-SSE.B.4

FORWARD 2

Compare two payment plans with the series formula • High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4

Plan A pays $1,000 per month for 24 months. Plan B pays $500 the first month and 8% more each month after that, for 24 months. Which plan pays more in total, and by how much?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q1

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q1/5

Plan A pays $1,000 per month. Plan B pays $500 the first month and 8% more each month after that. After how many months does Plan B's total first exceed Plan A's total for the same number of months?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q2

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q2/5

True or false?

A geometric plan with ratio greater than 1 eventually pays more in total than any plan with a fixed monthly amount.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q3

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q3/5

Plan C pays $2,000 then 5% less each month. Its total over 24 months is closest to

A. $40,000

B. $24,000

C. $48,000

D. $28,320

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q4

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q4/5

Name one assumption in comparing the plans by their totals only.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S12-F2-Q5

High School — Algebra II • Algebra: Seeing Structure in Expressions • HSA-SSE.B.4 • Set 12 • FORWARD 2 • Q5/5

Add and subtract polynomials of higher degree

MAIN	Algebra II main question	**HSA-APR.A.1**
BACKWARD 1	Combine like terms with the same power	**7.EE.A.1**
BACKWARD 2	Subtract a whole expression	**6.EE.A.3**
BACKWARD 3	Write polynomials in standard form	**HSA-SSE.A.1.a**
FORWARD 1	Reason about the degree of a sum or difference	**HSA-APR.A.1**
FORWARD 2	Add and subtract rational expressions	**HSA-APR.D.7**

SET 13 • HSA-APR.A.1

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Simplify $(4x^3 - 2x^2 + 7) + (-x^3 + 5x - 9)$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q1/5

Simplify $(6x^4 - x^3 + 2x) - (2x^4 + 3x^3 - x^2 + 2x - 1)$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q2/5

A company's revenue is $R(x) = -0.01x^3 + 3x^2 + 40x$ dollars and its cost is $C(x) = 0.02x^2 + 15x + 900$ dollars for x items. Write the profit $P(x) = R(x) - C(x)$ in standard form.

$P(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q3/5

Simplify $(x^5 - 3x^3 + x) - (x^5 + 2x^4 - 3x^3 - 7)$. Write the result in standard form and give its degree.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q4/5

$p(x) = 5x^3 + x^2 - 6$. Find the polynomial $q(x)$ for which $p(x) + q(x) = 2x^3 - x + 4$.

$q(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 13 • MAIN • Q5/5

SET 13 • HSA-APR.A.1

BACKWARD 1

Combine like terms with the same power • Grade 7 • Expressions & Equations • 7.EE.A.1

Simplify $4x^3 - x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q1

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q1/5

Simplify $-2x^2 + 5x - 3x^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q2

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q2/5

True or false?

$x^3 + x^2 = x^5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q3

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q3/5

Which pair are like terms?

A. $4x^2$ and $-x^2$

B. $4x^2$ and $4x$

C. x^3 and $3x$

D. x and 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q4

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q4/5

Simplify $7 - 9 + 2x - 2x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B1-Q5

Grade 7 • Expressions & Equations • 7.EE.A.1 • Set 13 • BACKWARD 1 • Q5/5

SET 13 • HSA-APR.A.1

BACKWARD 2

Subtract a whole expression • Grade 6 • Expressions & Equations • 6.EE.A.3

Rewrite $-(2x^4 + 3x^3 - x^2 + 2x - 1)$ without parentheses.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q1/5

Simplify $(5 - x) - (3 - 4x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q2/5

True or false?

$a - (b - c) = a - b - c$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q3/5

$-(x^2 - 7)$ equals

A. $x^2 + 7$

B. $-x^2 - 7$

C. $-x^2 + 7$

D. $x^2 - 7$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q4/5

Simplify $10 - (x + 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.A.3 • Set 13 • BACKWARD 2 • Q5/5

SET 13 • HSA-APR.A.1

BACKWARD 3

Write polynomials in standard form • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a

Write $7 + 5x - 2x^3 + x^2$ in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q1/5

What is the leading coefficient of $3 - 4x^2 + x^4$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q2/5

The degree of $-2x^4 + x + 7$ is

A. 7

B. 1

C. 4

D. -2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q3/5

True or false?

The standard form of $2x - x^3$ is $2x - x^3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q4/5

What is the constant term of $6x^3 - 2x + 9$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-B3-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.1.a • Set 13 • BACKWARD 3 • Q5/5

SET 13 • HSA-APR.A.1

FORWARD 1

Reason about the degree of a sum or difference • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

p(x) has degree 4 and q(x) has degree 4. What degrees can p(x) + q(x) have? Give an example where the degree drops.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q1/5

For which value of k does
$(kx^3 + 2x - 1) - (5x^3 - x^2 + 3)$ have degree 2?

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q2/5

True or false?

The difference of two polynomials is always a polynomial.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q3/5

p has degree 5 and q has degree 3. The degree of $p - q$ is

A. 5

B. 2

C. 3

D. 8

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q4/5

Explain why the sum of two polynomials is a polynomial.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 13 • FORWARD 1 • Q5/5

SET 13 • HSA-APR.A.1

FORWARD 2

Add and subtract rational expressions • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.7

Write 3/x + 2/(x + 1) as one rational expression, and state the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q1/5

Write $\frac{1}{(x - 2)} - \frac{1}{(x + 2)}$ as one rational expression, and state the excluded values.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q2/5

True or false?

Polynomials are closed under division.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q3/5

x/2 + x/3 equals

A. 5x/5

B. x/5

C. 5x/6

D. 2x/5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S13-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q4/5

Explain how adding rational expressions is like adding fractions of integers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S13-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.D.7 • Set 13 • FORWARD 2 • Q5/5

Multiply polynomials of higher degree

MAIN	Algebra II main question	**HSA-APR.A.1**
BACKWARD 1	Distribute a monomial over a polynomial	**HSA-APR.A.1**
BACKWARD 2	Multiply powers with the same base	**8.EE.A.1**
BACKWARD 3	Multiply two binomials	**HSA-APR.A.1**
FORWARD 1	Find a coefficient of a product without full expansion	**HSA-APR.A.1**
FORWARD 2	Expand powers of binomials with the Binomial Theorem	**HSA-APR.C.5**

SET 14 • HSA-APR.A.1

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(x^2 + 3x - 2)(2x - 5)$. Write the product in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q1/5

Expand $(x - 2)^3$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q2/5

Multiply $(x^2 - x + 1)(x^2 + x + 1)$. Write the product in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q3/5

Multiply $(2x + 1)(x - 3)(x + 4)$. Write the product in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q4/5

A box has length $x + 2$, width $x - 1$, and height $2x + 3$ (in cm). Write its volume as a polynomial in standard form.

$V(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-M-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • MAIN • Q5/5

SET 14 • HSA-APR.A.1

BACKWARD 1

Distribute a monomial over a polynomial • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $2x(x^2 + 3x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q1/5

Multiply $-5(x^2 + 3x - 2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q2/5

True or false?

$x^2(x + 4) = x^3 + 4x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q3/5

$-3x(x - 1)$ equals

A. $3x^2 - 3x$

B. $-3x^2 + 3x$

C. $-3x^2 - 3x$

D. $-3x - 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S14-B1-Q4**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q4/5

Multiply $x(x^2 - x + 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 1 • Q5/5

SET 14 • HSA-APR.A.1

BACKWARD 2

Multiply powers with the same base • Grade 8 • Expressions & Equations • 8.EE.A.1

Simplify $x^2 \cdot x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q1/5

Simplify $2x \cdot 3x^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q2/5

True or false?

$x^2 \cdot x^2 = x^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q3/5

$-x \cdot x^3$ equals

A. x^4

B. $-x^3$

C. $-x^4$

D. $-x^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q4/5

Simplify $(x^2)(-4x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 14 • BACKWARD 2 • Q5/5

SET 14 • HSA-APR.A.1

BACKWARD 3

Multiply two binomials • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Multiply $(x - 3)(x + 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q1/5

Multiply $(x + 2)(x - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q2/5

$(x - 2)^2$ equals

A. $x^2 + 4$

B. $x^2 - 4$

C. $x^2 - 4x + 4$

D. $x^2 - 2x + 4$

True or false?

$(2x + 1)(x - 3) = 2x^2 - 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q4/5

Multiply $(2x + 3)(x - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-B3-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 14 • BACKWARD 3 • Q5/5

SET 14 • HSA-APR.A.1

FORWARD 1

Find a coefficient of a product without full expansion • High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Find the coefficient of x^2 in $(x^2 + 3x - 2)(4x^2 - x + 5)$ without expanding everything.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q1/5

For which k does $(x + k)(x^2 - 3x + 1)$ have no x^2 term?

$k =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q2/5

True or false?

The constant term of a product of polynomials is the product of their constant terms.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q3/5

The leading coefficient of $(2x^3 - 1)(5x^2 + x)$ is

A. 7

B. 5

C. 2

D. 10

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q4/5

Find the coefficient of x in $(x - 1)(x - 2)(x - 3)$.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.A.1 • Set 14 • FORWARD 1 • Q5/5

SET 14 • HSA-APR.A.1

FORWARD 2

Expand powers of binomials with the Binomial Theorem • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Expand $(x + 1)^4$ using Pascal's Triangle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q1/5

Expand $(x - 2)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q2/5

True or false?

The coefficients of $(x + y)^5$ are 1, 5, 10, 5, 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q3/5

The coefficient of x^2y^2 in $(x + y)^4$ is

A. 2

B. 6

C. 4

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q4/5

Explain why the coefficients of $(x + y)^n$ are symmetric.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S14-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 14 • FORWARD 2 • Q5/5

Find remainders with the Remainder Theorem

MAIN	Algebra II main question	**HSA-APR.B.2**
BACKWARD 1	Evaluate a polynomial function	**HSF-IF.A.2**
BACKWARD 2	Powers of negative numbers	**7.NS.A.2**
BACKWARD 3	Dividend = divisor × quotient + remainder	**4.NBT.B.6**
FORWARD 1	Find unknown coefficients from given remainders	**HSA-APR.B.2**
FORWARD 2	Prove the Remainder Theorem	**HSA-APR.B.2**

SET 15 • HSA-APR.B.2

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Use the Remainder Theorem to find the remainder when $p(x) = x^3 - 4x^2 + 2x - 5$ is divided by $x - 3$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q1/5

Use the Remainder Theorem to find the remainder when $p(x) = 2x^4 + x^3 - 3x + 7$ is divided by $x + 2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q2/5

Use the Remainder Theorem to find the remainder when $p(x) = x^{100} + 3$ is divided by $x + 1$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q3/5

Use the Remainder Theorem to find the remainder when $p(x) = 4x^3 - 6x^2 + x - 1$ is divided by $x - 1/2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q4/5

Use the Remainder Theorem to find the remainder when $p(x) = x^4 - 5x^2 + 4x$ is divided by $x - 2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • MAIN • Q5/5

SET 15 • HSA-APR.B.2

BACKWARD 1

Evaluate a polynomial function • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

$p(x) = x^3 - 4x^2 + 2x - 5$. Find $p(1)$.

$p(1) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q1/5

$p(x) = 2x^2 - x + 3$. Find $p(-2)$.

$p(-2) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q2/5

True or false?

If $p(x) = x^2 + 1$, then $p(-3) = -8$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q3/5

If $p(x) = x^3 - x$, then $p(2) =$

A. 4

B. 6

C. 8

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q4/5

$p(x) = 4x - 1$. Find $p(1/2)$.

$p(1/2) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 15 • BACKWARD 1 • Q5/5

SET 15 • HSA-APR.B.2

BACKWARD 2

Powers of negative numbers • Grade 7 • The Number System • 7.NS.A.2

Evaluate $(-2)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q1/5

Evaluate $(-2)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q2/5

$(-1)^{100}$ equals

A. -100

B. -1

C. 100

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q3/5

True or false?

$-2^4 = (-2)^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q4/5

Evaluate $2(-2)^4 + (-2)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B2-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 15 • BACKWARD 2 • Q5/5

SET 15 • HSA-APR.B.2

BACKWARD 3

Dividend = divisor × quotient + remainder • Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6

Write $47 = 5 \times q + r$ with $0 \leq r < 5$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q1

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q1/5

A division gives quotient 12 and remainder 3 with divisor 7. What was the dividend?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q2

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q2/5

True or false?

If a division leaves remainder 0, the divisor is a factor of the dividend.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q3

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q3/5

$x^3 - 1 = (x - 1)(x^2 + x + 1) + 0$. The remainder on division by $x - 1$ is

A. 0

B. 1

C. $x^2 + x + 1$

D. -1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q4

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q4/5

$p(x) = (x - 3)(x^2 - x - 1) - 8$. What is the remainder when $p(x)$ is divided by $x - 3$?

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-B3-Q5

Grade 4 • Number & Operations in Base Ten • 4.NBT.B.6 • Set 15 • BACKWARD 3 • Q5/5

SET 15 • HSA-APR.B.2

FORWARD 1

Find unknown coefficients from given remainders • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

When $p(x) = x^3 + kx^2 - 4$ is divided by $x - 2$, the remainder is 12. Find k.

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q1/5

$p(x) = x^3 + ax + b$ leaves remainder 5 on division by $x - 1$ and remainder -1 on division by $x + 1$. Find a and b.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q2/5

True or false?

If p(x) leaves remainder 0 on division by $x - 4$, then $p(-4) = 0$.

A. True

B. False

$p(x) = 2x^2 + kx + 1$ leaves remainder 7 on division by $x + 1$. Then $k =$

A. 4

B. -4

C. -6

D. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q4/5

For which k is the remainder of $x^4 - kx + 6$ divided by $x + 2$ equal to 0?

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 1 • Q5/5

SET 15 • HSA-APR.B.2

FORWARD 2

Prove the Remainder Theorem • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Prove that the remainder when a polynomial p(x) is divided by $x - a$ is p(a).

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q1/5

Why must the remainder on division by $x - a$ be a constant?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q2/5

True or false?

The Remainder Theorem also works for division by $2x - 1$ with remainder $p(1/2)$.

A. True

B. False

Which step uses the fact that $a - a = 0$?

A. Graphing $p(x)$

B. Dividing $p(x)$ by x

C. Factoring $p(x)$

D. Substituting $x = a$ into $p(x) = (x - a)q(x) + r$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q4/5

Explain how the proof shows that $p(a) = 0$ exactly when $x - a$ is a factor of $p(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S15-F2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 15 • FORWARD 2 • Q5/5

Use p(a) = 0 to find factors

MAIN	Algebra II main question	**HSA-APR.B.2**
BACKWARD 1	Use the Remainder Theorem	**HSA-APR.B.2**
BACKWARD 2	Factor the quadratic quotient	**HSA-SSE.A.2**
BACKWARD 3	Divide a polynomial by a linear factor	**HSA-APR.A.1**
FORWARD 1	Choose parameters so that a given binomial is a factor	**HSA-APR.B.2**
FORWARD 2	Finish factorizations over the complex numbers	**HSN-CN.C.8**

SET 16 • HSA-APR.B.2

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Show that $x - 2$ is a factor of $p(x) = x^3 - 3x^2 - 4x + 12$, then factor $p(x)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q1/5

Show that $x + 1$ is a factor of $p(x) = x^3 + 2x^2 - 5x - 6$, then factor $p(x)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q2/5

Show that $x + 3$ is a factor of $p(x) = 2x^3 + 5x^2 - 4x - 3$, then factor $p(x)$ completely.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q3/5

Is $x - 4$ a factor of $p(x) = x^3 - 5x^2 + 2x + 9$? Justify with the Remainder Theorem.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q4/5

$p(x) = x^3 - 7x + 6$. Use $p(a) = 0$ to find all the zeros of p, and write p(x) in factored form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • MAIN • Q5/5

SET 16 • HSA-APR.B.2

BACKWARD 1

Use the Remainder Theorem • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Find the remainder when $x^3 - 3x^2 - 4x + 12$ is divided by $x - 2$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q1/5

Find the remainder when $x^3 + 2x^2 - 5x - 6$ is divided by $x - 1$.

Remainder: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q2/5

True or false?

Dividing by $x + 3$ means evaluating at $x = 3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q3/5

The remainder of $p(x) \div (x + 1)$ is

A. $-p(1)$

B. $p(-1)$

C. $p(0)$

D. $p(1)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q4/5

Find $p(-3)$ for $p(x) = 2x^3 + 5x^2 - 4x - 3$.

$p(-3) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • BACKWARD 1 • Q5/5

SET 16 • HSA-APR.B.2

BACKWARD 2

Factor the quadratic quotient • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 - x - 6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q1/5

Factor $2x^2 - x - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q2/5

$x^2 + x - 6$ equals

A. $(x - 3)(x + 2)$

B. $(x - 6)(x + 1)$

C. $(x + 3)(x - 2)$

D. $(x + 6)(x - 1)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q3/5

True or false?

$x^2 + 4$ factors over the integers.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q4/5

Factor $x^2 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 16 • BACKWARD 2 • Q5/5

SET 16 • HSA-APR.B.2

BACKWARD 3

Divide a polynomial by a linear factor • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Find $q(x)$ if $x^3 - 3x^2 - 4x + 12 = (x - 2)\,q(x)$.

$q(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B3-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q1/5

Multiply $(x + 1)(x^2 + x - 6)$ to check a division.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B3-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q2/5

True or false?

If $(x - 1)(x^2 + x - 6) = x^3 - 7x + 6$, then $x^2 + x - 6$ is the quotient of $x^3 - 7x + 6 \div (x - 1)$.

A. True

B. False

$x^3 - 8 \div (x - 2)$ equals

A. $x^2 + 2x + 4$

B. $x^2 + 4$

C. $x^2 - 4$

D. $x^2 - 2x + 4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S16-B3-Q4**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q4/5

Find $q(x)$ if $2x^3 + 5x^2 - 4x - 3 = (x + 3)\, q(x)$.

$q(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-B3-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 16 • BACKWARD 3 • Q5/5

SET 16 • HSA-APR.B.2

FORWARD 1

Choose parameters so that a given binomial is a factor • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2

Find k so that $x - 3$ is a factor of $x^3 + kx^2 - 3x + 9$.

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q1/5

Find a and b so that $x - 1$ and $x + 2$ are both factors of $x^3 + ax^2 + bx - 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q2/5

True or false?

If $x - 2$ and $x + 2$ are factors of $p(x)$, then $x^2 + 4$ is a factor of $p(x)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q3/5

For which k is $x + 1$ a factor of $x^4 + kx + 3$?

A. 2

B. -4

C. 4

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q4/5

Write a cubic with leading coefficient 1 that has $x - 1$, $x + 1$, and $x - 4$ as factors, in standard form.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.2 • Set 16 • FORWARD 1 • Q5/5

SET 16 • HSA-APR.B.2

FORWARD 2

Finish factorizations over the complex numbers • High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8

$x - 1$ is a factor of $x^3 - 1$. Factor $x^3 - 1$ completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q1/5

$x + 2$ is a factor of $x^3 + 2x^2 + 9x + 18$. Factor it completely over the complex numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q2/5

True or false?

Every factor found with the factor theorem gives a zero of the polynomial.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q3/5

$x - 2$ is a factor of $x^3 - 2x^2 + x - 2$. The other zeros are

A. i and $-i$

B. -2

C. 1 and -1

D. $2i$ and $-2i$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S16-F2-Q4**

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q4/5

$x - 3$ is a factor of $x^3 - 3x^2 + 4x - 12$. List all complex zeros.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S16-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Number & Quantity: The Complex Number System • HSN-CN.C.8 • Set 16 • FORWARD 2 • Q5/5

Identify zeros and multiplicities from a factorization

MAIN	Algebra II main question	**HSA-APR.B.3**
BACKWARD 1	Solve by the zero-product property	**HSA-REI.B.4.b**
BACKWARD 2	Factor to find hidden repeated factors	**HSA-SSE.A.2**
BACKWARD 3	An exponent counts repeated factors	**6.EE.A.1**
FORWARD 1	Build a polynomial from zeros and one more condition	**HSA-APR.B.3**
FORWARD 2	Decide what the zeros determine and what remains free	**HSA-APR.B.3**

SET 17 • HSA-APR.B.3

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

List the zeros of $p(x) = (x - 4)^2(x + 1)(2x - 3)$ and give the multiplicity of each.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q1/5

Factor $p(x) = x^3 - 6x^2 + 9x$, then list its zeros with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q2/5

List the zeros of $p(x) = (x^2 - 9)(x^2 + 2x + 1)$ with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q3/5

List the zeros of $p(x) = -2x(x + 5)^3$ with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q4/5

Factor $p(x) = x^4 - 8x^2 + 16$, then list its zeros with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • MAIN • Q5/5

SET 17 • HSA-APR.B.3

BACKWARD 1

Solve by the zero-product property • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b

Solve $(x - 4)(x + 1) = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q1/5

Solve $x(2x - 3) = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q2/5

True or false?

If $(x - 2)^2 = 0$, then $x = 2$ or $x = -2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q3/5

The solutions of $(x + 5)(x - 5) = 0$ are

A. 25

B. -5 and 5

C. 0 and 5

D. 5 only

Solve $-2x(x + 5) = 0$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.4.b • Set 17 • BACKWARD 1 • Q5/5

SET 17 • HSA-APR.B.3

BACKWARD 2

Factor to find hidden repeated factors • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $x^2 - 6x + 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q1/5

Factor $x^2 + 2x + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q2/5

$x^4 - 8x^2 + 16$ equals

A. $(x^2 + 4)^2$

B. $(x^2 - 4)^2$

C. $(x^2 - 8)^2$

D. $(x - 4)^4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q3/5

True or false?

$x^3 - 6x^2 + 9x = x(x + 3)^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q4/5

Factor $x^2 - 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 17 • BACKWARD 2 • Q5/5

SET 17 • HSA-APR.B.3

BACKWARD 3

An exponent counts repeated factors • Grade 6 • Expressions & Equations • 6.EE.A.1

Write $(x - 4)^2$ as a product without an exponent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q1/5

How many factors of $(x + 5)$ are in $(x + 5)^3$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q2/5

True or false?

$(x - 2)^2(x - 2) = (x - 2)^3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q3/5

In $7(x - 1)^4$, the factor $x - 1$ appears

A. 28 times

B. 4 times

C. 1 time

D. 7 times

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q4/5

Write $(x + 1)(x + 1)(x - 3)$ with an exponent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.A.1 • Set 17 • BACKWARD 3 • Q5/5

SET 17 • HSA-APR.B.3

FORWARD 1

Build a polynomial from zeros and one more condition • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

Write the cubic polynomial with zeros -1, 2, and 4 (each multiplicity 1) whose graph passes through $(0, 16)$.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q1/5

Write a polynomial of degree 3 with a zero of multiplicity 2 at $x = 3$, a zero at $x = -2$, and leading coefficient -1.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q2/5

True or false?

Zeros alone determine a polynomial completely.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q3/5

A quartic has zeros 1 and −1, each of multiplicity 2, and p(0) = 3. Then p(x) =

A. $-3(x-1)^2(x+1)^2$

B. $3(x^2-1)$

C. $(x-1)^2(x+1)^2$

D. $3(x-1)^2(x+1)^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q4/5

A polynomial has zeros 0 (multiplicity 2) and 5 (multiplicity 1), degree 3, and $p(1) = -8$. Find $p(x)$.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 1 • Q5/5

SET 17 • HSA-APR.B.3

FORWARD 2

Decide what the zeros determine and what remains free • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

All polynomials with zeros 1 and 3 only (each multiplicity 1) and degree 2 have what form?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q1/5

If the degree is not fixed, give two different polynomials whose only real zeros are 1 and 3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q2/5

True or false?

Two cubic polynomials with the same three zeros (each multiplicity 1) must be equal.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q3/5

Which extra information fixes a cubic with zeros −2, 0, 2?

A. That it has degree 3

B. That 0 is a zero

C. Its end behavior only

D. Its value at x = 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q4/5

Explain why knowing only the real zeros of a degree-4 polynomial may not determine it even up to scale.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S17-F2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 17 • FORWARD 2 • Q5/5

Construct a rough graph of a polynomial from its zeros

MAIN	Algebra II main question	**HSA-APR.B.3**
BACKWARD 1	Read zeros and multiplicities	**HSA-APR.B.3**
BACKWARD 2	End behavior from the leading term	**HSF-IF.B.4**
BACKWARD 3	Find the y-intercept by evaluating at 0	**HSF-IF.A.2**
BACKWARD 4	Sign of a power of a negative or positive number	**7.NS.A.2**
FORWARD 1	Sketch graphs from standard form by factoring first	**HSF-IF.C.7.c**
FORWARD 2	Write a polynomial from its rough graph	**HSA-APR.B.3**


SET 18 • HSA-APR.B.3

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$$y = (x + 2)(x - 1)(x - 3)$$




REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q1/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$$y = -(x + 1)^2(x - 2)$$




REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q2/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$y = x(x - 3)^2$




REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q3/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$y = (x+2)(x+1)(x-1)(x-3)$




REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q4/5

Sketch a rough graph on the grid. Show every zero (crossing or touching the x-axis), the y-intercept, and the end behavior.

$$y = -(x - 1)^3(x + 2)$$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • MAIN • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 1

Read zeros and multiplicities • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

List the zeros of $(x + 2)(x - 1)(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q1/5

List the zeros of $-(x + 1)^2(x - 2)$ with multiplicities.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q2/5

True or false?

$x(x - 3)^2$ has a zero of multiplicity 2 at $x = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q3/5

The zero $x = 1$ of $-(x - 1)^3(x + 2)$ has multiplicity

A. 1

B. 2

C. 4

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q4/5

List the zeros of $(x + 2)(x + 1)(x - 1)(x - 3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • BACKWARD 1 • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 2

End behavior from the leading term • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4

As x grows large and positive, what happens to x^3?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q1/5

As x grows large and negative, what happens to $-x^4$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q2/5

The leading term of $(x + 2)(x - 1)(x - 3)$ is

A. $-x^3$

B. -6

C. $3x$

D. x^3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q3/5

True or false?

A polynomial of even degree with a positive leading coefficient falls at both ends.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q4/5

What is the leading term of $-(x-1)^3(x+2)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.4 • Set 18 • BACKWARD 2 • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 3

Find the y-intercept by evaluating at 0 • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

Find y when x = 0 for $y = (x + 2)(x - 1)(x - 3)$.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q1/5

Find y when x = 0 for $y = -(x + 1)^2(x - 2)$.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q2/5

True or false?

The y-intercept of $x(x - 3)^2$ is 9.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q3/5

The y-intercept of $-(x-1)^3(x+2)$ is

A. 0

B. -2

C. 2

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q4/5

Find y when $x = 1$ for $y = x(x - 3)^2$.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 18 • BACKWARD 3 • Q5/5

SET 18 • HSA-APR.B.3

BACKWARD 4

Sign of a power of a negative or positive number • Grade 7 • The Number System • 7.NS.A.2

Is $(-0.5)^2$ positive or negative?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q1/5

Is $(-0.5)^3$ positive or negative?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q2/5

True or false?

An even power of a negative number is positive.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q3/5

Near $x = 3$, $(x - 3)^2$ is

A. negative on both sides

B. positive on both sides

C. zero everywhere

D. positive on the left and negative on the right

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q4/5

Does $(x - 1)^3$ change sign at $x = 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-B4-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 18 • BACKWARD 4 • Q5/5


SET 18 • HSA-APR.B.3

FORWARD 1

Sketch graphs from standard form by factoring first • High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c

Factor, then sketch a rough graph showing every zero, the y-intercept, and end behavior.

$y = x^3 - x^2 - 6x$




REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q1/5

Factor, then sketch a rough graph showing every zero, the y-intercept, and end behavior.

$y = x^4 - 5x^2 + 4$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q2/5

True or false?

$y = x^3 + 4x$ crosses the x-axis at $x = -2$, 0, and 2.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q3/5

$y = -x^3 + 9x$ crosses the x-axis at

A. 0 and 9

B. 3 only

C. $-3, 0, 3$

D. $-9, 0, 9$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q4/5

$y = x^3 - 3x^2 + 4 = (x + 1)(x - 2)^2$. Describe its behavior at each zero.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F1-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 18 • FORWARD 1 • Q5/5

SET 18 • HSA-APR.B.3

FORWARD 2

Write a polynomial from its rough graph • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

A cubic's graph crosses the x-axis at −3 and touches it at 2, and passes through (0, 12). Write the polynomial.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q1/5

A quartic rises at both ends, touches the x-axis at -1 and at 3, and passes through (0, 18). Write the polynomial.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q2/5

True or false?

A graph that touches the x-axis at x = a without crossing has a zero of even multiplicity at a.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q3/5

A cubic falls to the right and crosses the x-axis at 0, 1, and 4. Its leading coefficient is

A. zero

B. not determined by the end behavior

C. positive

D. negative

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S18-F2-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q4/5

A polynomial graph crosses at -2, touches at 1, falls to the left and rises to the right, and has y-intercept 4. Give a polynomial of least degree.

$p(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S18-F2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 18 • FORWARD 2 • Q5/5

Prove polynomial identities

MAIN	Algebra II main question	**HSA-APR.C.4**
BACKWARD 1	Expand the square of a binomial	**HSA-APR.A.1**
BACKWARD 2	Recognize a difference of squares	**HSA-SSE.A.2**
BACKWARD 3	What makes an identity	**6.EE.A.4**
FORWARD 1	Prove or disprove proposed identities	**HSA-APR.C.4**
FORWARD 2	Prove identities with the Binomial Theorem	**HSA-APR.C.5**

SET 19 • HSA-APR.C.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Prove the identity $(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q1/5

Prove the identity $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q2/5

Prove the identity $(a + b)^2 - (a - b)^2 = 4ab$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q3/5

Prove the identity $x^4 + 4 = (x^2 + 2x + 2)(x^2 - 2x + 2)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q4/5

Prove the identity $(n + 1)^3 - n^3 = 3n^2 + 3n + 1$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • MAIN • Q5/5

SET 19 • HSA-APR.C.4

BACKWARD 1

Expand the square of a binomial • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Expand $(a + b)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q1/5

Expand $(x^2 - y^2)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q2/5

True or false?

$(a - b)^2 = a^2 - b^2$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q3/5

$(2xy)^2$ equals

A. $2xy^2$

B. $4xy$

C. $2x^2y^2$

D. $4x^2y^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q4/5

Expand $(n + 1)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 19 • BACKWARD 1 • Q5/5

SET 19 • HSA-APR.C.4

BACKWARD 2

Recognize a difference of squares • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

Factor $a^2 - b^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q1/5

Factor $(x^2 + 2)^2 - (2x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q2/5

True or false?

$u^2 - v^2 = (u - v)(u + v)$ for any expressions u and v.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q3/5

Which is a difference of squares?

A. $x^3 - 1$

B. $x^2 - 3x$

C. $9x^2 - 16$

D. $9x^2 + 16$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q4/5

Factor $25 - x^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B2-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 19 • BACKWARD 2 • Q5/5

SET 19 • HSA-APR.C.4

BACKWARD 3

What makes an identity • Grade 6 • Expressions & Equations • 6.EE.A.4

Test $(a + b)^2 = a^2 + b^2$ with $a = 1$ and $b = 2$. Is it an identity?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q1/5

Explain the difference between an equation and an identity.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q2/5

True or false?

Checking an identity with three numbers proves it.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q3/5

$2(x + 3) = 2x + 6$ is

A. an identity

B. true only for $x = 3$

C. true only for $x = 0$

D. never true

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q4/5

Give values that show $a^3 + b^3 = (a + b)^3$ is not an identity.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.A.4 • Set 19 • BACKWARD 3 • Q5/5

SET 19 • HSA-APR.C.4

FORWARD 1

Prove or disprove proposed identities • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Prove or disprove: $(a + b)^3 = a^3 + b^3 + 3ab(a + b)$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • FORWARD 1 • Q1/5

Prove or disprove: $a^4 - b^4 = (a - b)^4$.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • FORWARD 1 • Q2/5

True or false?

$(x - 1)(x^2 + x + 1) = x^3 + 1$ is an identity.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • FORWARD 1 • Q3/5

Which is an identity?

A. $(x + 2)^2 = x^2 + 4$

B. $(x + 1)^3 = x^3 + 1$

C. $x^2 - 4 = (x - 4)^2$

D. $(x + 2)^2 - (x - 2)^2 = 8x$

Find c so that $(x + c)^2 - (x - c)^2 = 12x$ is an identity.

c = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 19 • FORWARD 1 • Q5/5

SET 19 • HSA-APR.C.4

FORWARD 2

Prove identities with the Binomial Theorem • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Use the Binomial Theorem with $x = y = 1$ to prove that the entries in row n of Pascal's Triangle add to 2^n.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q1/5

Use the Binomial Theorem with $x = 1$ and $y = -1$ to find $C(5,0) - C(5,1) + C(5,2) - C(5,3) + C(5,4) - C(5,5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q2/5

True or false?

$(1 + 1)^4 = 1 + 4 + 6 + 4 + 1.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q3/5

The sum of the coefficients of $(2x + 1)^3$ is

A. 9

B. 8

C. 6

D. 27

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q4/5

Use $(1 + 0.1)^3 = 1 + 3(0.1) + 3(0.01) + 0.001$ to find 1.1^3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S19-F2-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 19 • FORWARD 2 • Q5/5

Use polynomial identities to describe numerical relationships

MAIN	Algebra II main question	**HSA-APR.C.4**
BACKWARD 1	Substitute values into an identity	**6.EE.A.2**
BACKWARD 2	Check a Pythagorean triple	**8.G.B.6**
BACKWARD 3	Choose a and b to fit a product	**HSA-APR.C.4**
FORWARD 1	Generate a triple with a required side	**HSA-APR.C.4**
FORWARD 2	Connect the triple identity to complex numbers	**HSN-CN.A.2**

SET 20 • HSA-APR.C.4

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

The identity $(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$ generates Pythagorean triples. Use $x = 3$ and $y = 2$ to find a triple.

Triple: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q1/5

Use the identity $(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$ with $x = 5$ and $y = 2$ to find a Pythagorean triple.

Triple: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q2/5

Use the identity $(a + b)(a - b) = a^2 - b^2$ to compute 47×53 mentally. Show which a and b you use.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q3/5

Use the identity $(n + 1)^2 - n^2 = 2n + 1$ to write 41 as a difference of two consecutive squares.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q4/5

Use the identity $(n + 1)^3 - n^3 = 3n^2 + 3n + 1$ to compute $13^3 - 12^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • MAIN • Q5/5

SET 20 • HSA-APR.C.4

BACKWARD 1

Substitute values into an identity • Grade 6 • Expressions & Equations • 6.EE.A.2

Evaluate $x^2 - y^2$ for $x = 3$ and $y = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q1

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q1/5

Evaluate 2xy for x = 5 and y = 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q2

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q2/5

True or false?

For $n = 12$, $3n^2 + 3n + 1 = 433$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q3

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q3/5

For $a = 50$ and $b = 3$, $a^2 - b^2$ is

A. 2491

B. 2500

C. 2494

D. 2509

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q4

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q4/5

Evaluate 2n + 1 for n = 20.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B1-Q5

Grade 6 • Expressions & Equations • 6.EE.A.2 • Set 20 • BACKWARD 1 • Q5/5

SET 20 • HSA-APR.C.4

BACKWARD 2

Check a Pythagorean triple • Grade 8 • Geometry • 8.G.B.6

Do 5, 12, 13 satisfy $a^2 + b^2 = c^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q1

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q1/5

Do 20, 21, 29 satisfy $a^2 + b^2 = c^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q2

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q2/5

True or false?

6, 8, 11 is a Pythagorean triple.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q3

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q3/5

Which is a Pythagorean triple?

A. 8, 15, 17

B. 6, 7, 9

C. 2, 3, 4

D. 4, 5, 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q4

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q4/5

Explain why a triangle with sides 5, 12, 13 is a right triangle.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B2-Q5

Grade 8 • Geometry • 8.G.B.6 • Set 20 • BACKWARD 2 • Q5/5

SET 20 • HSA-APR.C.4

BACKWARD 3

Choose a and b to fit a product • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Write 47 and 53 as $50 - b$ and $50 + b$. What is b?

b = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B3-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q1/5

Write 98×102 as $(a - b)(a + b)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B3-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q2/5

True or false?

$29 \times 31 = 30^2 + 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B3-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q3/5

64 × 56 equals

A. $(60 - 4)^2$

B. $64^2 - 56^2$

C. $60^2 - 4^2$

D. $60^2 + 4^2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S20-B3-Q4**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q4/5

Compute 98×102.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-B3-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • BACKWARD 3 • Q5/5

SET 20 • HSA-APR.C.4

FORWARD 1

Generate a triple with a required side • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4

Use the triple identity to find x and y (whole numbers, x > y) that give a triple with hypotenuse 25 and legs 7 and 24.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q1/5

Find x and y (whole numbers, x > y) for which the identity gives a triple with a leg 2xy = 40 and hypotenuse 41.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q2/5

True or false?

Every choice of whole numbers $x > y > 0$ in the identity gives a Pythagorean triple.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q3/5

Which choice gives the triple 8, 15, 17?

A. $x = 4, y = 1$

B. $x = 5, y = 3$

C. $x = 4, y = 2$

D. $x = 3, y = 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q4/5

Explain why $x = 2y$ gives a triple with legs in the ratio $3 : 4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F1-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.4 • Set 20 • FORWARD 1 • Q5/5

SET 20 • HSA-APR.C.4

FORWARD 2

Connect the triple identity to complex numbers • High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2

Compute $(3 + 2i)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q1

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q1/5

Compute $(x + yi)^2$ in the form $a + bi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q2

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q2/5

True or false?

The real and imaginary parts of $(x + yi)^2$ are the legs of the Pythagorean triple from the identity.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q3

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q3/5

$(5 + 2i)^2$ equals

A. 25 + 4i

B. 21 + 20i

C. 21 + 10i

D. 29 + 20i

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q4

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q4/5

Use $(4 + i)^2$ to find a Pythagorean triple.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S20-F2-Q5

High School — Algebra II • Number & Quantity: The Complex Number System • HSN-CN.A.2 • Set 20 • FORWARD 2 • Q5/5

Expand with the Binomial Theorem

MAIN	Algebra II main question	**HSA-APR.C.5**
BACKWARD 1	Multiply binomials repeatedly	**HSA-APR.A.1**
BACKWARD 2	Build Pascal's Triangle by a rule	**4.OA.C.5**
BACKWARD 3	Powers of products and of negatives	**8.EE.A.1**
FORWARD 1	Find a specific term or coefficient with a condition	**HSA-APR.C.5**
FORWARD 2	Use binomial coefficients to compute probabilities	**HSS-CP.B.9**

SET 21 • HSA-APR.C.5

MAIN

Algebra II main question • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Use the Binomial Theorem to expand $(x + 2)^5$. Write the result in standard form.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q1/5

Use the Binomial Theorem to expand $(2a - b)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q2/5

Use the Binomial Theorem to find the coefficient of x^3 in $(x + 3)^7$.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q3/5

Use the Binomial Theorem to find the term containing x^3y^3 in $(x - 2y)^6$.

Term: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q4/5

Use the first three terms of the binomial expansion of $(1 + 0.02)^5$ to estimate 1.02^5 to four decimal places.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-M-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • MAIN • Q5/5

SET 21 • HSA-APR.C.5

BACKWARD 1

Multiply binomials repeatedly • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

Expand $(x + 2)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q1/5

Expand $(x + 2)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q2/5

True or false?

$(x + y)^3 = x^3 + y^3$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q3/5

The number of terms in the expansion of $(x + y)^5$ is

A. 6

B. 5

C. 10

D. 32

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q4/5

Expand $(a - b)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B1-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 21 • BACKWARD 1 • Q5/5

SET 21 • HSA-APR.C.5

BACKWARD 2

Build Pascal's Triangle by a rule • Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5

Row 4 of Pascal's Triangle is 1, 4, 6, 4, 1. Write row 5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q1

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q1/5

Write row 7 of Pascal's Triangle (row 6 is 1, 6, 15, 20, 15, 6, 1).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q2

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q2/5

True or false?

Each inner entry of Pascal's Triangle is the sum of the two entries above it.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q3

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q3/5

The fourth entry of row 7 (counting from the first 1) is

A. 20

B. 7

C. 35

D. 21

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q4

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q4/5

What is the sum of the entries in row 5?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B2-Q5

Grade 4 • Operations & Algebraic Thinking • 4.OA.C.5 • Set 21 • BACKWARD 2 • Q5/5

SET 21 • HSA-APR.C.5

BACKWARD 3

Powers of products and of negatives • Grade 8 • Expressions & Equations • 8.EE.A.1

Simplify $(2a)^4$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q1/5

Simplify $(-2y)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q2/5

True or false?

$(-b)^4 = -b^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q3/5

3^4 equals

A. 27

B. 12

C. 64

D. 81

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q4/5

Simplify $(0.02)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-B3-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 21 • BACKWARD 3 • Q5/5

SET 21 • HSA-APR.C.5

FORWARD 1

Find a specific term or coefficient with a condition • High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.C.5

Find the constant term of $(x + 1/x)^6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q1

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q1/5

The coefficient of x^2 in $(1 + kx)^5$ is 90. Find k ($k > 0$).

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q2

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q2/5

True or false?

The middle term of $(x + y)^8$ is $56x^4y^4$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q3

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q3/5

The term with x^4 in $(x^2 - 1)^5$ is

A. $-5x^4$

B. $-10x^4$

C. $10x^4$

D. $5x^4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q4

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q4/5

Find the coefficient of x in $(2x - 1)^6$.

Coefficient: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F1-Q5

High School — Algebra II, (+) standard listed in the pathway • Algebra: Arithmetic with Polynomials & Rational Expressions
HSA-APR.C.5 • Set 21 • FORWARD 1 • Q5/5

SET 21 • HSA-APR.C.5

FORWARD 2

Use binomial coefficients to compute probabilities • High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability • HSS-CP.B.9

A fair coin is tossed 5 times. Use C(5, k) to find P(exactly 3 heads).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q1

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q1/5

A spinner lands on red with probability 0.3. Find P(exactly 2 reds in 4 spins).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q2

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q2/5

True or false?

The probabilities of 0, 1, ..., n successes add to 1 because $(p + (1 - p))^n = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q3

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q3/5

For 6 fair coin tosses, P(exactly 3 heads) is

A. 6/64

B. 3/6

C. 15/64

D. 20/64

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S21-F2-Q4**

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability
HSS-CP.B.9 • Set 21 • FORWARD 2 • Q4/5

Explain why C(n, k) counts the outcomes with exactly k successes in n trials.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S21-F2-Q5

High School — Probability & Statistics (course placement in this bank) • Statistics & Probability: Conditional Probability & the Rules of Probability

HSS-CP.B.9 • Set 21 • FORWARD 2 • Q5/5

Rewrite rational expressions by polynomial division

MAIN	Algebra II main question	**HSA-APR.D.6**
BACKWARD 1	Write a division with a remainder as a mixed number	**4.NF.B.3**
BACKWARD 2	Multiply to check a division	**HSA-APR.A.1**
BACKWARD 3	Use the Remainder Theorem to check the remainder	**HSA-APR.B.2**
FORWARD 1	Divide with unknown coefficients and higher-degree divisors	**HSA-APR.D.6**
FORWARD 2	Use division to find slant asymptotes	**HSF-IF.C.7.d**

SET 22 • HSA-APR.D.6

MAIN

Algebra II main question • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6

Rewrite $(x^3 + 2x^2 - 5x + 3)/(x - 1)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q1/5

Rewrite $(2x^3 - 3x^2 + 4x - 7)/(x^2 + 1)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q2/5

Rewrite $(x^4 - 16)/(x - 2)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q3/5

Rewrite $(6x^2 + x - 4)/(2x + 1)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q4/5

Rewrite $(x^3 - 4x + 5)/(x + 3)$ in the form $q(x) + r(x)/b(x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-M-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.D.6 • Set 22 • MAIN • Q5/5

SET 22 • HSA-APR.D.6

BACKWARD 1

Write a division with a remainder as a mixed number • Grade 4 • Number & Operations–Fractions • 4.NF.B.3

Write 23/4 as a mixed number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q1

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q1/5

Write 47/5 in the form q + r/5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q2

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q2/5

True or false?

17/3 = 5 + 1/3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S22-B1-Q3

Grade 4 • Number & Operations–Fractions • 4.NF.B.3 • Set 22 • BACKWARD 1 • Q3/5