True or false?

A temperature that is 7°C at t = 0, 23°C at t = 12 h, and repeats every 24 hours is modeled by $15 - 8\cos(\pi t/12)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S43-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q3/5

1601

QUESTION ID

S43-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Minimum 7 at t = 0.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S43-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A blood pressure that starts at 100 mm Hg and rises, varies between 80 and 120, and repeats every 0.8 s is modeled by

A. 100 + 20sin(0.8t)

B. 20 + 100sin(2.5πt)

C. 120sin(2.5πt)

D. 100 + 20sin(2.5πt)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S43-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q4/5

1602

QUESTION ID
S43-F2-Q4

TYPE
Multiple choice

CORRECT ANSWER
D.
100 + 20sin(2.5πt)

GRADING NOTE Period 0.8.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S43-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A piston's position starts at 0 cm and increases, varies between −3 and 3 cm, and repeats every 1 s. Write a model for p.

p = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S43-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q5/5

1603

QUESTION ID

S43-F2-Q5

TYPE

Short answer

CORRECT ANSWER

3sin(2πt)

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S43-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 43 • FORWARD 2 • Q5/5

SET 44 • HSF-IF.B.5 • INTERPRETING FUNCTIONS

State domains of radical, rational, logarithmic, and polynomial models

MAIN	Algebra II main question	**HSF-IF.B.5**
BACKWARD 1	Domains of linear, quadratic, and exponential models	**HSF-IF.B.5**
BACKWARD 2	Division by zero	**7.NS.A.2.b**
BACKWARD 3	Solve linear inequalities	**HSA-REI.B.3**
FORWARD 1	Domains defined by nonlinear conditions	**HSF-IF.B.5**
FORWARD 2	Domains and ranges of inverse functions	**HSF-BF.B.4.a**

SET 44 • HSF-IF.B.5

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

A car's speed is $s(d) = \sqrt{(22.5d)}$ mph for a skid of d feet. Give the domain of the function and the domain that makes sense in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q1/5

1606

QUESTION ID

S44-M-Q1

TYPE

Short answer

CORRECT ANSWER

Function: $d \geq 0$; context: $d \geq 0$ (real skid lengths, up to a realistic maximum)

GRADING NOTE Form required: both. $d < 0$ gives no real root.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S44-M-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q1/5

The average cost of x shirts is A(x) = (500 + 4x)/x. Give the domain of the function and the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q2/5

1607

QUESTION ID

S44-M-Q2

TYPE

Short answer

CORRECT ANSWER

Function: all real x ≠ 0; context: whole numbers x ≥ 1

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S44-M-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q2/5

pH = −log[H⁺], where [H⁺] is the hydrogen ion concentration. What is the domain?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q3/5

1608

QUESTION ID

S44-M-Q3

TYPE

Short answer

CORRECT ANSWER

$[H^+] > 0$

GRADING NOTE Form required: the inequality. Logarithms need positive inputs.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S44-M-Q3**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A box from a 12-in sheet has volume $V(x) = x(12 - 2x)^2$. Give the domain of the polynomial and the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q4/5

1609

QUESTION ID
S44-M-Q4

TYPE
Short answer

CORRECT ANSWER
Polynomial: all real numbers; context: $0 < x < 6$

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the domain of $f(x) = \sqrt{(x - 2)}/(x - 5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q5/5

1610

QUESTION ID

S44-M-Q5

TYPE

Short answer

CORRECT ANSWER

$x \geq 2$ and $x \neq 5$

GRADING NOTE Form required: both conditions.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S44-M-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • MAIN • Q5/5

SET 44 • HSF-IF.B.5

BACKWARD 1

Domains of linear, quadratic, and exponential models • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

$h(t) = -16t^2 + 64t$ models height until the ball lands. Give the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q1/5

1612

QUESTION ID

S44-B1-Q1

TYPE

Short answer

CORRECT ANSWER

$0 \le t \le 4$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

C(n) = 2n + 5 is the cost of n tickets. Give the domain in context.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q2/5

1613

QUESTION ID

S44-B1-Q2

TYPE

Short answer

CORRECT ANSWER

Whole numbers

n ≥ 0

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q2/5

True or false?

The domain of $f(x) = 2^x$ is $x > 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q3/5

1614

QUESTION ID

S44-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE 2^x is defined for every real x; $y > 0$ is its range.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q3/5

A model for the number of people uses a domain of

A. fractions only

B. all real numbers

C. whole numbers

D. negative numbers

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q4/5

1615

QUESTION ID

S44-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. whole numbers

GRADING NOTE Count.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Why might a model's domain be smaller than the function's domain?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q5/5

1616

QUESTION ID

S44-B1-Q5

TYPE

Short answer

CORRECT ANSWER

The context restricts inputs (for example to nonnegative or whole numbers).

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • BACKWARD 1 • Q5/5

SET 44 • HSF-IF.B.5

BACKWARD 2

Division by zero • Grade 7 • The Number System • 7.NS.A.2.b

ANSWER KEY Reduced copy of the question page, with its answer

For which x is 1/(x − 5) undefined?

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q1

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q1/5

1618

QUESTION ID

S44-B2-Q1

TYPE

Short answer

CORRECT ANSWER

5

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q1

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

For which x is (500 + 4x)/x undefined?

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q2

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q2/5

1619

QUESTION ID

S44-B2-Q2

TYPE

Short answer

CORRECT ANSWER

0

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q2

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q2/5

True or false?

3/0 = 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q3

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q3/5

1620

QUESTION ID

S44-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Undefined.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q3

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$(x + 1)/(x^2 - 9)$ is undefined at

A. $x = 0$

B. $x = 9$

C. $x = -1$

D. $x = \pm 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q4

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q4/5

1621

QUESTION ID

S44-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. $x = \pm 3$

GRADING NOTE $x^2 = 9$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q4

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Is 0/5 defined? Give its value.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q5

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q5/5

1622

QUESTION ID

S44-B2-Q5

TYPE

Short answer

CORRECT ANSWER

Yes, 0

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B2-Q5

Grade 7 • The Number System • 7.NS.A.2.b • Set 44 • BACKWARD 2 • Q5/5

SET 44 • HSF-IF.B.5

BACKWARD 3

Solve linear inequalities • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3

ANSWER KEY Reduced copy of the question page, with its answer

Solve $x - 2 \geq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q1/5

1624

QUESTION ID

S44-B3-Q1

TYPE

Short answer

CORRECT ANSWER

$x \geq 2$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $12 - 2x > 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q2/5

1625

QUESTION ID

S44-B3-Q2

TYPE

Short answer

CORRECT ANSWER

$x < 6$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

22.5d ≥ 0 means d ≥ 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q3/5

1626

QUESTION ID

S44-B3-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Divide by a positive number.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$3 - x \geq 0$ means

A. $x \geq 3$

B. $x \leq 3$

C. $x \geq -3$

D. $x \leq -3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q4/5

1627

QUESTION ID

S44-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. $x \leq 3$

GRADING NOTE Add x.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve 2x + 8 > 0.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q5/5

1628

QUESTION ID

S44-B3-Q5

TYPE

Short answer

CORRECT ANSWER

x > −4

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-B3-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 44 • BACKWARD 3 • Q5/5

SET 44 • HSF-IF.B.5

FORWARD 1

Domains defined by nonlinear conditions • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

Give the domain of $f(x) = \sqrt{(9 - x^2)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q1/5

1630

QUESTION ID

S44-F1-Q1

TYPE

Short answer

CORRECT ANSWER

$-3 \leq x \leq 3$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S44-F1-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the domain of $g(x) = \log(x^2 - 4)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q2/5

1631

QUESTION ID

S44-F1-Q2

TYPE

Short answer

CORRECT ANSWER

x < −2 or x > 2

GRADING NOTE Form required: both parts.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S44-F1-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q2/5

Give the domain of $h(x) = 1/(x^2 - x - 6)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q3/5

1632

QUESTION ID

S44-F1-Q3

TYPE

Short answer

CORRECT ANSWER

All real x except −2 and 3

GRADING NOTE Form required: both exclusions.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S44-F1-Q3**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q3/5

True or false?

The domain of $\sqrt{(x^2)}$ is $x \geq 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q4/5

1633

QUESTION ID
S44-F1-Q4

TYPE
True or false

CORRECT ANSWER
B. False

GRADING NOTE $x^2 \geq 0$ for every x, so $\sqrt{(x^2)}$ is defined for all real x.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

The domain of $\log(x - 1) + \sqrt{(5 - x)}$ is

A. $1 < x \leq 5$

B. $x \leq 5$

C. $1 \leq x \leq 5$

D. $x > 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q5/5

1634

QUESTION ID

S44-F1-Q5

TYPE

Multiple choice

CORRECT ANSWER

A. $1 < x \leq 5$

GRADING NOTE Both conditions.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.5 • Set 44 • FORWARD 1 • Q5/5

SET 44 • HSF-IF.B.5

FORWARD 2

Domains and ranges of inverse functions • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

$f(x) = \sqrt{(x - 2)}$ has domain $x \geq 2$ and range $y \geq 0$. Write f^{-1} and give its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q1/5

1636

QUESTION ID

S44-F2-Q1

TYPE

Short answer

CORRECT ANSWER

$f^{-1}(x) = x^2 + 2$ with domain $x \geq 0$

GRADING NOTE Form required: formula and domain.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 2^x$ has range $y > 0$. What is the domain of $f^{-1}(x) = \log_2 x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q2/5

1637

QUESTION ID
S44-F2-Q2

TYPE
Short answer

CORRECT ANSWER
x > 0

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q2/5

True or false?

The domain of an inverse function is the range of the original function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q3/5

1638

QUESTION ID
S44-F2-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Inputs and outputs swap.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 1/x + 3$ $(x \neq 0)$ has an inverse with domain

A. $x \neq 0$

B. $x > 3$

C. all real x

D. $x \neq 3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q4/5

1639

QUESTION ID

S44-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. $x \neq 3$

GRADING NOTE The range of f excludes 3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q4/5

Explain why the inverse of $s(d) = \sqrt{(22.5d)}$ needs the restriction $s \geq 0$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q5/5

1640

QUESTION ID

S44-F2-Q5

TYPE

Short answer

CORRECT ANSWER

The range of s is $s \geq 0$, so $d = s^2/22.5$ is the inverse only for $s \geq 0$.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S44-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 44 • FORWARD 2 • Q5/5

SET 45 • HSF-IF.B.6 • INTERPRETING FUNCTIONS

Calculate and interpret average rates of change of nonlinear models

MAIN	Algebra II main question	**HSF-IF.B.6**
BACKWARD 1	Average rate of change of simpler functions	**HSF-IF.B.6**
BACKWARD 2	Slope between two points	**8.F.B.4**
BACKWARD 3	Evaluate nonlinear functions	**HSF-IF.A.2**
FORWARD 1	Compare average rates on different intervals	**HSF-IF.B.6**
FORWARD 2	Compare rates of change across representations	**HSF-IF.C.9**

SET 45 • HSF-IF.B.6

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6

A culture has $N(t) = 200 \cdot 2^t$ bacteria after t hours. Find the average rate of change from t = 1 to t = 3 and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q1/5

1643

QUESTION ID
S45-M-Q1

TYPE
Short answer

CORRECT ANSWER
600 bacteria per hour: on average the culture grew by 600 bacteria each hour from hour 1 to hour 3

GRADING NOTE Form required: value with units and interpretation. (1600 − 400)/2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q1/5

A stone's height is $h(t) = 64 - 16t^2$ ft. Find the average rate of change from $t = 0.5$ to $t = 1.5$ s and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q2/5

1644

QUESTION ID

S45-M-Q2

TYPE

Short answer

CORRECT ANSWER

−32 ft/s: the stone fell an average of 32 feet per second

GRADING NOTE Form required: value with units and interpretation. (28 − 60)/1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S45-M-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

The table shows a car's value. Find the average rate of change from t = 2 to t = 4 years and interpret it.

t (years)	0	1	2	3	4	5
V ($)	20000	17000	14450	12282.50	10440.06	8874.11

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q3/5

1645

QUESTION ID

S45-M-Q3

TYPE

Short answer

CORRECT ANSWER

About −$2,004.97 per year: the value dropped about $2,005 per year on average

GRADING NOTE Form required: value with units and interpretation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q3/5

Speed from skid length is s(d) = √(24d) mph. Find the average rate of change from d = 24 to d = 96 ft and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q4/5

1646

QUESTION ID

S45-M-Q4

TYPE

Short answer

CORRECT ANSWER

1/3 mph per foot: on average each extra foot of skid adds 1/3 mph of speed

GRADING NOTE Form required: value with units and interpretation. (48 − 24)/72.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S45-M-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Temperature is T(t) = 15 − 8cos(πt/12) °C after t hours. Find the average rate of change from t = 0 to t = 6 and interpret it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q5/5

1647

QUESTION ID
S45-M-Q5

TYPE
Short answer

CORRECT ANSWER
4/3 °C per hour: on average the temperature rose about 1.33°C per hour

GRADING NOTE Form required: value with units and interpretation. (15 − 7)/6.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • MAIN • Q5/5

SET 45 • HSF-IF.B.6

BACKWARD 1

Average rate of change of simpler functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = x^2$. Find the average rate of change from $x = 1$ to $x = 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q1/5

1649

QUESTION ID

S45-B1-Q1

TYPE

Short answer

CORRECT ANSWER

4

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A table gives f(2) = 10 and f(5) = 1. Find the average rate of change.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q2/5

1650

QUESTION ID

S45-B1-Q2

TYPE

Short answer

CORRECT ANSWER

−3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q2/5

True or false?

The average rate of change from a to b is (f(b) − f(a))/(b − a).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q3/5

1651

QUESTION ID

S45-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q3/5

The average rate of change of f(x) = 3x + 1 on any interval is

A. it varies

B. 1

C. 3

D. 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q4/5

1652

QUESTION ID

S45-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. 3

GRADING NOTE Linear.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

What are the units of a rate if f is in dollars and x is in years?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q5/5

1653

QUESTION ID

S45-B1-Q5

TYPE

Short answer

CORRECT ANSWER

Dollars per year

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • BACKWARD 1 • Q5/5

SET 45 • HSF-IF.B.6

BACKWARD 2

Slope between two points • Grade 8 • Functions • 8.F.B.4

ANSWER KEY Reduced copy of the question page, with its answer

Find the slope between (1, 400) and (3, 1600).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q1

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q1/5

1655

QUESTION ID

S45-B2-Q1

TYPE

Short answer

CORRECT ANSWER

600

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q1

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the slope between (0.5, 60) and (1.5, 28).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q2

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q2/5

1656

QUESTION ID

S45-B2-Q2

TYPE

Short answer

CORRECT ANSWER

−32

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q2

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q2/5

True or false?

A negative slope means the output increases as the input increases.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B2-Q3

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q3/5

1657

QUESTION ID

S45-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE A negative slope means the output decreases.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S45-B2-Q3

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The slope between (24, 24) and (96, 48) is

A. 72/24

B. 1/3

C. 2

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q4

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q4/5

1658

QUESTION ID

S45-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 1/3

GRADING NOTE 24/72 = 1/3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q4

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the slope between (0, 7) and (6, 15).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q5

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q5/5

1659

QUESTION ID

S45-B2-Q5

TYPE

Short answer

CORRECT ANSWER

4/3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B2-Q5

Grade 8 • Functions • 8.F.B.4 • Set 45 • BACKWARD 2 • Q5/5

SET 45 • HSF-IF.B.6

BACKWARD 3

Evaluate nonlinear functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Find N(3) for $N(t) = 200 \cdot 2^t$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q1/5

1661

QUESTION ID

S45-B3-Q1

TYPE

Short answer

CORRECT ANSWER

1600

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find h(1.5) for $h(t) = 64 - 16t^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q2/5

1662

QUESTION ID

S45-B3-Q2

TYPE

Short answer

CORRECT ANSWER

28

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q2/5

True or false?

$\sqrt{(24 \cdot 96)} = 2304$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q3/5

1663

QUESTION ID

S45-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $24 \cdot 96 = 2304$ and $\sqrt{2304} = 48$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

15 − 8cos(π/2) equals

A. 15

B. 7

C. 23

D. 8

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q4/5

1664

QUESTION ID

S45-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 15

GRADING NOTE cos(π/2) = 0.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Find T(0) for T(t) = 15 − 8cos(πt/12).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q5/5

1665

QUESTION ID

S45-B3-Q5

TYPE

Short answer

CORRECT ANSWER

7

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 45 • BACKWARD 3 • Q5/5

SET 45 • HSF-IF.B.6

FORWARD 1

Compare average rates on different intervals • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6

For $N(t) = 200 \cdot 2^t$, compare the average rates on [0, 1] and [3, 4]. What does the comparison show?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q1/5

1667

QUESTION ID

S45-F1-Q1

TYPE

Short answer

CORRECT ANSWER

200 and 1600 bacteria per hour: the growth speeds up

GRADING NOTE Form required: both rates and the conclusion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S45-F1-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

For s(d) = √(24d), compare the average rates on [0, 24] and [96, 150].

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q2/5

1668

QUESTION ID

S45-F1-Q2

TYPE

Short answer

CORRECT ANSWER

1 and 2/9 ≈ 0.22 mph per foot: the rate decreases

GRADING NOTE Form required: both and the trend.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S45-F1-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q2/5

True or false?

For a function with period 24, the average rate of change over [0, 24] is 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q3/5

1669

QUESTION ID

S45-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE f(24) = f(0).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

For f(x) = x^3, the average rate on [−1, 1] is

A. 3

B. 0

C. 2

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q4/5

1670

QUESTION ID

S45-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 1

GRADING NOTE (1 − (−1))/2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q4/5

Find an interval of length 6 on which $T(t) = 15 - 8\cos(\pi t/12)$ has average rate of change $-4/3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q5/5

1671

QUESTION ID

S45-F1-Q5

TYPE

Short answer

CORRECT ANSWER

[12, 18]

GRADING NOTE Any valid interval (for example [12, 18]).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.6 • Set 45 • FORWARD 1 • Q5/5

SET 45 • HSF-IF.B.6

FORWARD 2

Compare rates of change across representations • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9

ANSWER KEY Reduced copy of the question page, with its answer

$f(t) = 200 \cdot 2^t$, and g is given by $g(1) = 500$, $g(3) = 1500$. Which has the greater average rate on [1, 3]?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q1/5

1673

QUESTION ID

S45-F2-Q1

TYPE

Short answer

CORRECT ANSWER

f: 600 > 500

GRADING NOTE Form required: answer and both rates.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A graph of a cooling liquid falls from 90°C at t = 0 to 50°C at t = 20 min. Compare with T(t) = 90 − 1.5t on the same interval.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q2/5

1674

QUESTION ID

S45-F2-Q2

TYPE

Short answer

CORRECT ANSWER

Graph: −2°C/min; formula: −1.5°C/min; the graph cools faster on average

GRADING NOTE Form required: both rates and the comparison.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q2/5

True or false?

A function with a greater average rate on one interval must be greater everywhere.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q3/5

1675

QUESTION ID

S45-F2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Rates and values are different.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q3/5

A table shows h(0) = 0, h(2) = 40; the formula $k(t) = 5t^2$ gives

A. no rate for k

B. a greater rate for k

C. a greater rate for h

D. equal average rates on [0, 2]

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q4/5

1676

QUESTION ID

S45-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. a greater rate for h

GRADING NOTE h: 40/2 = 20; k: (20 − 0)/2 = 10.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q4/5

Explain how to compare rates when one function is a graph and one is a formula.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S45-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q5/5

1677

QUESTION ID

S45-F2-Q5

TYPE

Short answer

CORRECT ANSWER

Read the two endpoint values from the graph, evaluate the formula at the same inputs, and compute both quotients.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S45-F2-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 45 • FORWARD 2 • Q5/5

SET 46 • HSF-IF.C.7.b • INTERPRETING FUNCTIONS

Graph square root and cube root functions with transformations

MAIN	Algebra II main question	**HSF-IF.C.7.b**
BACKWARD 1	Graph $y = \sqrt{x}$ and $y = \sqrt[3]{x}$ from a table	**HSF-IF.C.7.b**
BACKWARD 2	Translations and reflections of graphs	**HSF-BF.B.3**
BACKWARD 3	Square and cube roots of numbers	**8.EE.A.2**
FORWARD 1	Graph radical functions with horizontal stretches and reflections	**HSF-IF.C.7.b**
FORWARD 2	Graph a radical function and its inverse	**HSF-BF.B.4.a**


SET 46 • HSF-IF.C.7.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

Graph the function on the grid. Show and label the starting point and two more points, and state the domain and range.

$y = \sqrt{(x - 2)} + 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q1/5

1680

QUESTION ID
S46-M-Q1

TYPE
Drawing / plotting

CORRECT ANSWER
Starts at (2, 1); passes through (3, 2) and (6, 3); increasing; domain x ≥ 2, range y ≥ 1.


GRADING NOTE Any graph with these features; domain and range required.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q1/5

Graph the function on the grid. Show and label the starting point and two more points, and state the domain and range.

$y = -\sqrt{(x + 3)}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q2/5

1681

QUESTION ID

S46-M-Q2

TYPE

Drawing / plotting

CORRECT ANSWER

Starts at (−3, 0); passes through (−2, −1) and (1, −2); decreasing; domain x ≥ −3, range y ≤ 0.


GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q2/5

Graph the function on the grid. Show and label the starting point and two more points, and state the domain and range.

$y = 2\sqrt{x} - 3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION — ID S46-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q3/5

1682

QUESTION ID

S46-M-Q3

TYPE

Drawing / plotting

CORRECT ANSWER

Starts at (0, −3); passes through (1, −1) and (4, 1), (9, 3); domain $x \geq 0$, range $y \geq -3$.


GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION — ID S46-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q3/5

Graph the function on the grid. Show and label the center point and two points on each side, and state the domain and range.

$y = \sqrt[3]{(x + 1)}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q4/5

1683

QUESTION ID

S46-M-Q4

TYPE

Drawing / plotting

CORRECT ANSWER

Center (−1, 0); points (−2, −1), (−9, −2), (0, 1), (7, 2); increasing; domain and range all real numbers.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the center point and two more points, and state the domain and range.

$y = -\sqrt[3]{(x - 2)} + 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q5/5

1684

QUESTION ID

S46-M-Q5

TYPE

Drawing / plotting

CORRECT ANSWER

Center (2, 1); points (3, 0), (1, 2), (10, −1), (−6, 3); decreasing; domain and range all real numbers.

GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • MAIN • Q5/5

SET 46 • HSF-IF.C.7.b

BACKWARD 1

Graph $y = \sqrt{x}$ and $y = \sqrt[3]{x}$ from a table • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b

ANSWER KEY Reduced copy of the question page, with its answer

Complete the table for y = √x at x = 0, 1, 4, 9.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q1/5

1686

QUESTION ID

S46-B1-Q1

TYPE

Short answer

CORRECT ANSWER

0, 1, 2, 3

GRADING NOTE Form required: all four.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Complete the table for $y = \sqrt[3]{x}$ at $x = -8, -1, 0, 1, 8$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q2/5

1687

QUESTION ID

S46-B1-Q2

TYPE

Short answer

CORRECT ANSWER

−2, −1, 0, 1, 2

GRADING NOTE Form required: all five.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S46-B1-Q2**

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q2/5

True or false?

The graph of $y = \sqrt{x}$ starts at the origin and decreases.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q3/5

1688

QUESTION ID

S46-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $\sqrt{x}$ increases as x increases.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The domain of $y = \sqrt[3]{x}$ is

A. $x > 0$

B. all real numbers

C. $x \leq 0$

D. $x \geq 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q4/5

1689

QUESTION ID
S46-B1-Q4

TYPE
Multiple choice

CORRECT ANSWER
B. all real numbers

GRADING NOTE Cube roots of negatives exist.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the range of $y = \sqrt{x}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q5/5

1690

QUESTION ID

S46-B1-Q5

TYPE

Short answer

CORRECT ANSWER

$y \geq 0$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • BACKWARD 1 • Q5/5

SET 46 • HSF-IF.C.7.b

BACKWARD 2

Translations and reflections of graphs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

How is $y = \sqrt{(x - 2)} + 1$ obtained from $y = \sqrt{x}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q1/5

1692

QUESTION ID

S46-B2-Q1

TYPE

Short answer

CORRECT ANSWER

Right 2, up 1

GRADING NOTE Any equivalent description.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

How is $y = -\sqrt{(x + 3)}$ obtained from $y = \sqrt{x}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q2/5

1693

QUESTION ID

S46-B2-Q2

TYPE

Short answer

CORRECT ANSWER

Left 3, then reflect over the x-axis

GRADING NOTE Any equivalent description.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$y = 2\sqrt{x}$ is a vertical stretch of $y = \sqrt{x}$ by 2.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q3/5

1694

QUESTION ID

S46-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Outputs doubled.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q3/5

The point (0, 0) of $y = \sqrt{x}$ moves to which point for $y = \sqrt{(x - 2)} + 1$?

A. (2, 1)

B. (1, 2)

C. (−2, 1)

D. (2, −1)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q4/5

1695

QUESTION ID

S46-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. (2, 1)

GRADING NOTE Right 2, up 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Where does (1, 1) of $y = \sqrt[3]{x}$ go under $y = \sqrt[3]{(x + 1)}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q5/5

1696

QUESTION ID

S46-B2-Q5

TYPE

Short answer

CORRECT ANSWER

(0, 1)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 46 • BACKWARD 2 • Q5/5

SET 46 • HSF-IF.C.7.b

BACKWARD 3

Square and cube roots of numbers • Grade 8 • Expressions & Equations • 8.EE.A.2

Find $\sqrt{9}$ and $\sqrt[3]{(-8)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q1/5

1698

QUESTION ID

S46-B3-Q1

TYPE

Short answer

CORRECT ANSWER

3 and −2

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q1

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find $\sqrt[3]{27}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q2/5

1699

QUESTION ID

S46-B3-Q2

TYPE

Short answer

CORRECT ANSWER

3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q2

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

√(−4) is a real number.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q3/5

1700

QUESTION ID

S46-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE No real square root of a negative.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q3

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$\sqrt[3]{(-1)}$ equals

A. i

B. undefined

C. -1

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q4/5

1701

QUESTION ID

S46-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. −1

GRADING NOTE $(-1)^3 = -1$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q4

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Find √16 − 3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q5/5

1702

QUESTION ID

S46-B3-Q5

TYPE

Short answer

CORRECT ANSWER

1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-B3-Q5

Grade 8 • Expressions & Equations • 8.EE.A.2 • Set 46 • BACKWARD 3 • Q5/5


SET 46 • HSF-IF.C.7.b

FORWARD 1

Graph radical functions with horizontal stretches and reflections • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

Graph the function on the grid. Show and label the starting point and two more points, and state the domain.

$y = \sqrt{(-x)}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q1/5

1704

QUESTION ID

S46-F1-Q1

TYPE

Drawing / plotting

CORRECT ANSWER

Starts at (0, 0); passes through (−1, 1), (−4, 2), (−9, 3); domain x ≤ 0.

GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Show that √(4x) = 2√x for x ≥ 0 and explain what this says about the two graphs.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q2/5

1705

QUESTION ID

S46-F1-Q2

TYPE

Short answer

CORRECT ANSWER

√(4x) = √4 · √x = 2√x, so the horizontal compression by 1/4 equals the vertical stretch by 2.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$y = \sqrt[3]{(-x)}$ is the same as $y = \sqrt[3]{x}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q3/5

1706

QUESTION ID

S46-F1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $\sqrt[3]{(-x)} = -\sqrt[3]{x}$; at $x = 8$ they give -2 and 2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The domain of $y = \sqrt{(3 - x)}$ is

A. all reals

B. $x \geq 3$

C. $x \leq 3$

D. $x \geq -3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q4/5

1707

QUESTION ID

S46-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. $x \leq 3$

GRADING NOTE $3 - x \geq 0$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write a square root function whose graph starts at (4, −2) and goes up to the left.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q5/5

1708

QUESTION ID
S46-F1-Q5

TYPE
Short answer

CORRECT ANSWER
√(4 − x) − 2 (for example)

GRADING NOTE Any valid function.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 46 • FORWARD 1 • Q5/5

SET 46 • HSF-IF.C.7.b

FORWARD 2

Graph a radical function and its inverse • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of $f(x) = \sqrt{(x - 2)} + 1$ and give its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q1/5

1710

QUESTION ID

S46-F2-Q1

TYPE

Short answer

CORRECT ANSWER

$f^{-1}(x) = (x - 1)^2 + 2$ for $x \geq 1$

GRADING NOTE Form required: formula and domain.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S46-F2-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of $g(x) = \sqrt[3]{(x + 1)}$.

$g^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q2/5

1711

QUESTION ID

S46-F2-Q2

TYPE

Short answer

CORRECT ANSWER

$x^3 - 1$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S46-F2-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q2/5

True or false?

The graph of an inverse is the reflection of the graph over y = x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q3/5

1712

QUESTION ID

S46-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Inputs and outputs swap.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The point (6, 3) on $f(x) = \sqrt{(x - 2)} + 1$ gives which point on f^{-1}?

A. (6, 3)

B. (3, 6)

C. (−6, −3)

D. (3, −6)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q4/5

1713

QUESTION ID

S46-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. (3, 6)

GRADING NOTE Swap.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S46-F2-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why the inverse of $f(x) = \sqrt{(x - 2)} + 1$ needs the domain $x \geq 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q5/5

1714

QUESTION ID

S46-F2-Q5

TYPE

Short answer

CORRECT ANSWER

The range of f is $y \geq 1$, which becomes the domain of the inverse; without it $(x - 1)^2 + 2$ is not one-to-one.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S46-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 46 • FORWARD 2 • Q5/5

SET 47 • HSF-IF.C.7.b • INTERPRETING FUNCTIONS

Graph piecewise-defined and absolute value functions

MAIN	Algebra II main question	**HSF-IF.C.7.b**
BACKWARD 1	Evaluate and graph simple piecewise functions	**HSF-IF.C.7.b**
BACKWARD 2	Absolute value as distance	**6.NS.C.7**
BACKWARD 3	Open and closed endpoints	**6.EE.B.8**
FORWARD 1	Write piecewise functions from graphs and contexts	**HSF-IF.C.7.b**
FORWARD 2	Describe absolute value graphs as transformations, and identify…	**HSF-BF.B.3**


SET 47 • HSF-IF.C.7.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

Graph the function on the grid. Show and label each endpoint with an open or closed circle.

$f(x) = x + 3$ for $x < -1$, and
$f(x) = x^2$ for $x \geq -1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION — ID S47-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q1/5

1717

QUESTION ID
S47-M-Q1

TYPE
Drawing / plotting

CORRECT ANSWER

A ray of slope 1 ending at an open circle (−1, 2); a parabola piece starting at a closed circle (−1, 1) through (0, 0), (1, 1), (2, 4).


GRADING NOTE Endpoint circles must be correct.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION — ID S47-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q1/5

Graph the function on the grid. Show and label the vertex and two more points.

$y = |2x - 4| - 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q2/5

1718

QUESTION ID

S47-M-Q2

TYPE

Drawing / plotting

CORRECT ANSWER

V with vertex (2, −1); points (0, 3), (4, 3), (1, 1), (3, 1).

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label each endpoint with an open or closed circle.

$f(x) = -2$ for $x \leq 1$, and
$f(x) = \sqrt{(x - 1)}$ for $x > 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q3/5

1719

QUESTION ID
S47-M-Q3

TYPE
Drawing / plotting

CORRECT ANSWER

Horizontal ray y = −2 ending at a closed circle (1, −2); square-root piece starting at an open circle (1, 0) through (2, 1), (5, 2), (10, 3).

GRADING NOTE Endpoint circles must be correct.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the vertex and two more points.

$y = -|x + 1| + 3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q4/5

1720

QUESTION ID
S47-M-Q4

TYPE
Drawing / plotting

CORRECT ANSWER
Upside-down V with vertex (−1, 3); points (−4, 0), (2, 0), (0, 2).

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label each endpoint with an open or closed circle.

$f(x) = 2^x$ for $x < 1$, and
$f(x) = 4 - x$ for $x \geq 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q5/5

1721

QUESTION ID
S47-M-Q5

TYPE
Drawing / plotting

CORRECT ANSWER

Exponential piece through (−2, 0.25), (0, 1), approaching y = 0 to the left, ending at an open circle (1, 2); a line from a closed circle (1, 3) through (4, 0).

GRADING NOTE Endpoint circles must be correct.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • MAIN • Q5/5

SET 47 • HSF-IF.C.7.b

BACKWARD 1

Evaluate and graph simple piecewise functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 2$ for $x < 0$ and $f(x) = x$ for $x \geq 0$. Find $f(-3)$ and $f(5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q1/5

1723

QUESTION ID

S47-B1-Q1

TYPE

Short answer

CORRECT ANSWER

2 and 5

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 2$ for $x < 0$ and $f(x) = x$ for $x \geq 0$. Find $f(0)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q2/5

1724

QUESTION ID

S47-B1-Q2

TYPE

Short answer

CORRECT ANSWER

0

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q2/5

True or false?

A piecewise function uses different rules on different parts of its domain.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q3/5

1725

QUESTION ID

S47-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

For $f(x) = x + 3$ $(x < -1)$, x^2 $(x \geq -1)$, $f(-1)$ is

A. 3

B. 2

C. -1

D. 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q4/5

1726

QUESTION ID

S47-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 1

GRADING NOTE Use the second rule.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = x + 3$ for $x < -1$, and $f(x) = x^2$ for $x \geq -1$. Find $f(-2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q5/5

1727

QUESTION ID

S47-B1-Q5

TYPE

Short answer

CORRECT ANSWER

1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • BACKWARD 1 • Q5/5

SET 47 • HSF-IF.C.7.b

BACKWARD 2

Absolute value as distance • Grade 6 • The Number System • 6.NS.C.7

ANSWER KEY Reduced copy of the question page, with its answer

Find |−5| and |3|.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q1

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q1/5

1729

QUESTION ID

S47-B2-Q1

TYPE

Short answer

CORRECT ANSWER

5 and 3

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q1

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find $|2(1) - 4| - 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q2

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q2/5

1730

QUESTION ID

S47-B2-Q2

TYPE

Short answer

CORRECT ANSWER

1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q2

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q2/5

True or false?

|x| can be negative for negative x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q3

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q3/5

1731

QUESTION ID

S47-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE |x| is a distance, so it is never negative: |−3| = 3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q3

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q3/5

|x + 1| = 0 when

A. x = 0

B. x = −1

C. x = 1

D. never

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q4

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q4/5

1732

QUESTION ID

S47-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. x = −1

GRADING NOTE Vertex input.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q4

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Find $-|0 + 1| + 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q5

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q5/5

1733

QUESTION ID

S47-B2-Q5

TYPE

Short answer

CORRECT ANSWER

2

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B2-Q5

Grade 6 • The Number System • 6.NS.C.7 • Set 47 • BACKWARD 2 • Q5/5

SET 47 • HSF-IF.C.7.b

BACKWARD 3

Open and closed endpoints • Grade 6 • Expressions & Equations • 6.EE.B.8

On a number line, is $x < -1$ drawn with an open or a closed circle at -1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q1/5

1735

QUESTION ID

S47-B3-Q1

TYPE

Short answer

CORRECT ANSWER

Open

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q1

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Is x ≥ 1 drawn with an open or a closed circle at 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q2/5

1736

QUESTION ID

S47-B3-Q2

TYPE

Short answer

CORRECT ANSWER

Closed

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q2

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q2/5

True or false?

An open circle means the endpoint is included.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q3/5

1737

QUESTION ID

S47-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE It is excluded.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S47-B3-Q3

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

On a number line, $x \leq 1$ is drawn with

A. an open circle and shading right

B. a closed circle at 1 and shading right

C. an open circle at 1 and shading left

D. a closed circle at 1 and shading left

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q4/5

1738

QUESTION ID

S47-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. a closed circle at 1 and shading left

GRADING NOTE Included endpoint.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q4

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write the inequality for "all numbers greater than 1".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q5/5

1739

QUESTION ID

S47-B3-Q5

TYPE

Short answer

CORRECT ANSWER

x > 1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-B3-Q5

Grade 6 • Expressions & Equations • 6.EE.B.8 • Set 47 • BACKWARD 3 • Q5/5

SET 47 • HSF-IF.C.7.b

FORWARD 1

Write piecewise functions from graphs and contexts • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b

ANSWER KEY Reduced copy of the question page, with its answer

A shipping company charges $5 for packages up to 2 lb and $5 plus $1.50 per pound over 2 lb. Write the cost C(w).

C(w) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q1/5

1741

QUESTION ID

S47-F1-Q1

TYPE

Short answer

CORRECT ANSWER

5 for $0 < w \leq 2$; $5 + 1.5(w - 2)$ for $w > 2$

GRADING NOTE Any equivalent form with correct endpoints.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q1/5

Write |x − 3| as a piecewise function without absolute value bars.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q2/5

1742

QUESTION ID
S47-F1-Q2

TYPE
Short answer

CORRECT ANSWER
3 − x for x < 3; x − 3 for x ≥ 3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

A piecewise function can have a jump and still be a function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q3/5

1743

QUESTION ID
S47-F1-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Each input still has one output.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which piecewise function is continuous at $x = 1$?

A. 2^x for $x < 1$, $4 - x$ for $x \geq 1$

B. x^2 for $x < 1$, $2x - 1$ for $x \geq 1$

C. 2 for $x < 1$, 3 for $x \geq 1$

D. x for $x < 1$, $x + 1$ for $x \geq 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q4/5

1744

QUESTION ID
S47-F1-Q4

TYPE
Multiple choice

CORRECT ANSWER
B. x^2 for $x < 1$, $2x - 1$ for $x \geq 1$

GRADING NOTE Both pieces give 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Find k so that $f(x) = kx$ for $x < 2$ and $f(x) = x^2$ for $x \geq 2$ has no jump at 2.

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q5/5

1745

QUESTION ID

S47-F1-Q5

TYPE

Short answer

CORRECT ANSWER

2

GRADING NOTE Any equivalent form. 2k = 4.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S47-F1-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.b • Set 47 • FORWARD 1 • Q5/5

SET 47 • HSF-IF.C.7.b

FORWARD 2

Describe absolute value graphs as transformations, and identify symmetry • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Describe y = |2x − 4| − 1 as a transformation of y = |x|.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q1/5

1747

QUESTION ID

S47-F2-Q1

TYPE

Short answer

CORRECT ANSWER

y = 2|x − 2| − 1: right 2, vertical stretch by 2, down 1

GRADING NOTE Any equivalent description.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q1/5

Write an absolute value function with vertex (−1, 3) that opens down and passes through (2, 0).

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q2/5

1748

QUESTION ID

S47-F2-Q2

TYPE

Short answer

CORRECT ANSWER

−|x + 1| + 3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S47-F2-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q2/5

True or false?

y = |x| is an odd function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q3/5

1749

QUESTION ID

S47-F2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE |−x| = |x|, so it is even.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = |x - 2|$ is symmetric about

A. the origin

B. $y = 2$

C. $x = 0$

D. $x = 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q4/5

1750

QUESTION ID

S47-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. x = 2

GRADING NOTE Vertex at 2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Is $y = -|x + 1| + 3$ an even function? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S47-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q5/5

1751

QUESTION ID

S47-F2-Q5

TYPE

Short answer

CORRECT ANSWER

No: f(1) = 1 but f(−1) = 3

GRADING NOTE Form required: decision and evidence.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S47-F2-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 47 • FORWARD 2 • Q5/5

SET 48 • HSF-IF.C.7.c • INTERPRETING FUNCTIONS

Graph polynomial functions showing zeros and end behavior

MAIN	Algebra II main question	**HSF-IF.C.7.c**
BACKWARD 1	Factor out common factors and special products	**HSA-SSE.A.2**
BACKWARD 2	Crossing and touching from multiplicity	**HSA-APR.B.3**
BACKWARD 3	Evaluate polynomials at test points	**HSF-IF.A.2**
FORWARD 1	Locate relative extrema of polynomial graphs	**HSF-IF.C.7.c**
FORWARD 2	Interpret polynomial graphs in context	**HSF-IF.B.4**

SET 48 • HSF-IF.C.7.c


MAIN

Algebra II main question • High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c

ANSWER KEY Reduced copy of the question page, with its answer

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = x^3 - 4x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q1/5

1754

QUESTION ID

S48-M-Q1

TYPE

Drawing / plotting

CORRECT ANSWER

x(x − 2)(x + 2): crosses at −2, 0, 2; points (−1, 3) and (1, −3); falls left, rises right.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = -x^3 + 3x^2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q2/5

1755

QUESTION ID

S48-M-Q2

TYPE

Drawing / plotting

CORRECT ANSWER

$-x^2(x - 3)$: touches at 0, crosses at 3; point (2, 4); y-intercept 0; rises left, falls right.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = x^4 - 4x^2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q3/5

1756

QUESTION ID

S48-M-Q3

TYPE

Drawing / plotting

CORRECT ANSWER

$x^2(x - 2)(x + 2)$: touches at 0, crosses at ±2; points (±1, −3); rises at both ends.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = x^3 + x^2 - 6x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q4/5

1757

QUESTION ID

S48-M-Q4

TYPE

Drawing / plotting

CORRECT ANSWER

x(x + 3)(x − 2): crosses at −3, 0, 2; points (−2, 8) and (1, −4); falls left, rises right.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Factor, then graph on the grid. Show every zero, the y-intercept, one point between each pair of zeros, and the end behavior.

$y = -x^4 + 2x^3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q5/5

1758

GRADING NOTE Any graph with these features.

QUESTION ID

S48-M-Q5

TYPE

Drawing / plotting

CORRECT ANSWER

$-x^3(x - 2)$: crosses at 0 with a flattened shape (multiplicity 3) and crosses at 2; point (1, 1); falls at both ends.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-M-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • MAIN • Q5/5

SET 48 • HSF-IF.C.7.c

BACKWARD 1

Factor out common factors and special products • High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Factor $x^3 - 4x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q1/5

1760

QUESTION ID

S48-B1-Q1

TYPE

Short answer

CORRECT ANSWER

x(x − 2)(x + 2)

GRADING NOTE Form required: complete.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q1

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Factor $-x^3 + 3x^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q2/5

1761

QUESTION ID

S48-B1-Q2

TYPE

Short answer

CORRECT ANSWER

$-x^2(x - 3)$

GRADING NOTE Any equivalent complete factorization.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q2

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q2/5

True or false?

$x^4 - 4x^2 = x^2(x^2 - 4) = x^2(x - 2)(x + 2)$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q3/5

1762

QUESTION ID

S48-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Common factor, then difference of squares.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q3

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$x^3 + x^2 - 6x$ equals

A. $(x + 3)(x - 2)$

B. $x(x + 3)(x - 2)$

C. $x(x - 3)(x + 2)$

D. $x(x + 6)(x - 1)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q4/5

1763

QUESTION ID

S48-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. $x(x + 3)(x - 2)$

GRADING NOTE $x^2 + x - 6$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q4

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Factor $-x^4 + 2x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q5/5

1764

QUESTION ID

S48-B1-Q5

TYPE

Short answer

CORRECT ANSWER

$-x^3(x - 2)$

GRADING NOTE Any equivalent complete factorization.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B1-Q5

High School — Algebra I • Algebra: Seeing Structure in Expressions • HSA-SSE.A.2 • Set 48 • BACKWARD 1 • Q5/5

SET 48 • HSF-IF.C.7.c

BACKWARD 2

Crossing and touching from multiplicity • High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3

ANSWER KEY Reduced copy of the question page, with its answer

Does $y = x^2(x - 3)$ cross or touch the x-axis at 0?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q1/5

1766

QUESTION ID

S48-B2-Q1

TYPE

Short answer

CORRECT ANSWER

Touches

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q1

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q1/5

Does $y = x^3(x - 2)$ cross at 0?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q2/5

1767

QUESTION ID

S48-B2-Q2

TYPE

Short answer

CORRECT ANSWER

Yes, with a flattened shape (odd multiplicity 3)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q2

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q2/5

True or false?

A zero of odd multiplicity is a crossing point.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q3/5

1768

QUESTION ID

S48-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Sign change.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q3

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The graph of $y = (x + 2)^2(x - 1)$ at $x = -2$

A. crosses

B. touches

C. has an asymptote

D. is undefined

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q4/5

1769

QUESTION ID

S48-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. touches

GRADING NOTE Even multiplicity.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q4

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the end behavior of $y = -x^4 + 2x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B2-Q5

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q5/5

1770

QUESTION ID

S48-B2-Q5

TYPE

Short answer

CORRECT ANSWER

Falls at both ends

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S48-B2-Q5**

High School — Algebra II • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.B.3 • Set 48 • BACKWARD 2 • Q5/5

SET 48 • HSF-IF.C.7.c

BACKWARD 3

Evaluate polynomials at test points • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $x^3 - 4x$ at $x = -1$ and $x = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q1/5

1772

QUESTION ID

S48-B3-Q1

TYPE

Short answer

CORRECT ANSWER

3 and −3

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $-x^3 + 3x^2$ at $x = 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q2/5

1773

QUESTION ID

S48-B3-Q2

TYPE

Short answer

CORRECT ANSWER

4

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$x^4 - 4x^2$ at $x = 1$ equals 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q3/5

1774

QUESTION ID

S48-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $1 - 4 = -3$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$x^3 + x^2 - 6x$ at $x = -2$ equals

A. 8

B. 0

C. -8

D. 16

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q4/5

1775

QUESTION ID

S48-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 8

GRADING NOTE $-8 + 4 + 12$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $-x^4 + 2x^3$ at $x = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q5/5

1776

QUESTION ID

S48-B3-Q5

TYPE

Short answer

CORRECT ANSWER

1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 48 • BACKWARD 3 • Q5/5

SET 48 • HSF-IF.C.7.c

FORWARD 1

Locate relative extrema of polynomial graphs • High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c

ANSWER KEY Reduced copy of the question page, with its answer

Use a table with steps of 0.1 (or technology) to estimate the relative maximum of $y = x^3 - 4x$ to the nearest tenth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q1/5

1778

QUESTION ID

S48-F1-Q1

TYPE

Short answer

CORRECT ANSWER

About (−1.2, 3.1)

GRADING NOTE Accept x from −1.2 to −1.1 and y about 3.08. Exact x = −2/√3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q1

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Use technology or a table to estimate the relative minimum of $y = x^3 + x^2 - 6x$ on $0 < x < 2$ to the nearest tenth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q2/5

1779

QUESTION ID

S48-F1-Q2

TYPE

Short answer

CORRECT ANSWER

About (1.1, −4.1)

GRADING NOTE Accept x from 1.1 to 1.2 and y from −4.1 to −4.0. Exact x = (−1 + √19)/3 ≈ 1.12.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q2

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q2/5

True or false?

A polynomial of degree 4 always has exactly 3 turning points.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q3/5

1780

QUESTION ID

S48-F1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE It has at most 3; $y = x^4$ has only one turning point.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q3

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

How many turning points can $y = x^4 - 4x^2$ have?

A. 1

B. 4

C. 3

D. 2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q4/5

1781

QUESTION ID

S48-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. 3

GRADING NOTE It has 3: two minima and one maximum.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q4

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why the graph of $y = x^3 - 4x$ has a turning point between 0 and 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q5/5

1782

QUESTION ID

S48-F1-Q5

TYPE

Short answer

CORRECT ANSWER

It is 0 at x = 0 and x = 2 and negative between them, so it must turn back up.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F1-Q5

High School — Algebra II • Functions: Interpreting Functions • HSF-IF.C.7.c • Set 48 • FORWARD 1 • Q5/5

SET 48 • HSF-IF.C.7.c

FORWARD 2

Interpret polynomial graphs in context • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4

ANSWER KEY Reduced copy of the question page, with its answer

A box's volume is $V(x) = x(12 - 2x)^2$. Which part of the polynomial graph is meaningful, and what is the maximum volume there?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q1/5

1784

QUESTION ID

S48-F2-Q1

TYPE

Short answer

CORRECT ANSWER

0 < x < 6; maximum 128 in³ at x = 2

GRADING NOTE Form required: interval and maximum.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A company's profit (thousands of dollars) is $P(x) = -x^3 + 9x^2 - 15x$ for x hundred items, $0 \le x \le 8$. When is the profit positive?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q2/5

1785

QUESTION ID

S48-F2-Q2

TYPE

Short answer

CORRECT ANSWER

For about $2.2 < x < 6.8$ (exactly between $(9 - \sqrt{21})/2 \approx 2.21$ and $(9 + \sqrt{21})/2 \approx 6.79$)

GRADING NOTE Form required: the interval. Endpoints accepted within ±0.05. $P(x) = -x(x^2 - 9x + 15)$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q2/5

True or false?

A zero of a profit model is a break-even point.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q3/5

1786

QUESTION ID

S48-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Profit 0.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A population model $P(t) = -t^3 + 6t^2 + 15t + 100$ rises then falls. The relative maximum means

A. the largest population before decline

B. the starting population

C. the end behavior

D. the time the population is 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q4/5

1787

QUESTION ID

S48-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. the largest population before decline

GRADING NOTE Turning point.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q4/5

Why does the end behavior of $V(x) = x(12 - 2x)^2$ not matter for the box?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S48-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q5/5

1788

QUESTION ID

S48-F2-Q5

TYPE

Short answer

CORRECT ANSWER

The context only uses $0 < x < 6$.

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S48-F2-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.B.4 • Set 48 • FORWARD 2 • Q5/5

SET 49 • HSF-IF.C.7.e • INTERPRETING FUNCTIONS

Graph exponential functions with intercepts, asymptote, and end behavior

MAIN	Algebra II main question	**HSF-IF.C.7.e**
BACKWARD 1	Graph basic exponential functions	**HSF-IF.C.7.e**
BACKWARD 2	Zero and negative exponents	**8.EE.A.1**
BACKWARD 3	Vertical and horizontal shifts and reflections	**HSF-BF.B.3**
FORWARD 1	Compare and describe exponential graphs	**HSF-IF.C.7.e**
FORWARD 2	Graph the inverse of an exponential function	**HSF-BF.B.4.a**

SET 49 • HSF-IF.C.7.e


MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the y-intercept, the x-intercept (as a decimal), and the asymptote (dashed).

$y = 2^x - 3$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q1/5

1791

QUESTION ID

S49-M-Q1

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote $y = -3$; y-intercept $(0, -2)$; x-intercept $\log_2 3 \approx 1.58$; points $(1, -1)$, $(2, 1)$, $(3, 5)$; increasing.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the y-intercept and the asymptote (dashed), and describe the end behavior.

$y = 3(0.5)^x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q2/5

1792

QUESTION ID

S49-M-Q2

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote y = 0; y-intercept (0, 3); points (−2, 12), (−1, 6), (1, 1.5), (2, 0.75); decreasing; y → 0 as x → ∞.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the y-intercept, the x-intercept, and the asymptote (dashed).

$y = -2^x + 4$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q3/5

1793

QUESTION ID

S49-M-Q3

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote y = 4; y-intercept (0, 3); x-intercept (2, 0); points (1, 2), (3, −4); decreasing; approaches 4 to the left.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the y-intercept and the asymptote (dashed).

$y = 2^{x+1}$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q4/5

1794

QUESTION ID

S49-M-Q4

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote y = 0; y-intercept (0, 2); points (−1, 1), (1, 4), (2, 8); increasing.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the y-intercept and the asymptote (dashed).

$y = (1/3)^x + 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q5/5

1795

QUESTION ID
S49-M-Q5

TYPE
Drawing / plotting

CORRECT ANSWER
Asymptote y = 1; y-intercept (0, 2); points (−1, 4), (−2, 10), (1, 4/3); decreasing toward 1.

GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • MAIN • Q5/5

SET 49 • HSF-IF.C.7.e

BACKWARD 1

Graph basic exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Complete the table for $y = 2^x$ at $x = -2, -1, 0, 1, 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q1/5

1797

QUESTION ID

S49-B1-Q1

TYPE

Short answer

CORRECT ANSWER

1/4, 1/2, 1, 2, 4

GRADING NOTE Form required: all five.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

What is the y-intercept of $y = 3(0.5)^x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q2/5

1798

QUESTION ID

S49-B1-Q2

TYPE

Short answer

CORRECT ANSWER

3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q2/5

True or false?

$y = 2^x$ has the horizontal asymptote $y = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q3/5

1799

QUESTION ID

S49-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE $2^x \to 0$ as $x \to -\infty$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = (0.5)^x$ is

A. linear

B. increasing

C. constant

D. decreasing

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q4/5

1800

QUESTION ID

S49-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. decreasing

GRADING NOTE Base less than 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Does $y = 2^x$ have an x-intercept?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q5/5

1801

QUESTION ID

S49-B1-Q5

TYPE

Short answer

CORRECT ANSWER

No: $2^x > 0$ for every x

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • BACKWARD 1 • Q5/5

SET 49 • HSF-IF.C.7.e

BACKWARD 2

Zero and negative exponents • Grade 8 • Expressions & Equations • 8.EE.A.1

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate 2^{-2}.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q1/5

1803

QUESTION ID

S49-B2-Q1

TYPE

Short answer

CORRECT ANSWER

1/4

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $(1/3)^{-2}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q2/5

1804

QUESTION ID

S49-B2-Q2

TYPE

Short answer

CORRECT ANSWER

9

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$b^0 = 0$ for $b \neq 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q3/5

1805

QUESTION ID

S49-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Any nonzero base to the 0 power is 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$3(0.5)^{-1}$ equals

A. 6

B. 3

C. -1.5

D. 1.5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q4/5

1806

QUESTION ID

S49-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 6

GRADING NOTE $0.5^{-1} = 2$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $2^{-1} - 3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q5/5

1807

QUESTION ID

S49-B2-Q5

TYPE

Short answer

CORRECT ANSWER

−2.5

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 49 • BACKWARD 2 • Q5/5

SET 49 • HSF-IF.C.7.e

BACKWARD 3

Vertical and horizontal shifts and reflections • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

How does $y = 2^x - 3$ relate to $y = 2^x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q1/5

1809

QUESTION ID

S49-B3-Q1

TYPE

Short answer

CORRECT ANSWER

Shifted down 3

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

How does $y = 2^{x+1}$ relate to $y = 2^{x}$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q2/5

1810

QUESTION ID

S49-B3-Q2

TYPE

Short answer

CORRECT ANSWER

Shifted left 1

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q2/5

True or false?

A vertical shift moves the horizontal asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q3/5

1811

QUESTION ID

S49-B3-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE y = 0 becomes y = k.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = -2^x + 4$ has asymptote

A. y = 4

B. y = 0

C. y = −4

D. x = 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q4/5

1812

QUESTION ID

S49-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. y = 4

GRADING NOTE Reflect, then up 4.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the asymptote of $y = (1/3)^x + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q5/5

1813

QUESTION ID

S49-B3-Q5

TYPE

Short answer

CORRECT ANSWER

y = 1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 49 • BACKWARD 3 • Q5/5

SET 49 • HSF-IF.C.7.e

FORWARD 1

Compare and describe exponential graphs • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Explain why the graphs of $y = 2^x$ and $y = (1/2)^x$ are reflections of each other over the y-axis.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q1/5

1815

QUESTION ID

S49-F1-Q1

TYPE

Short answer

CORRECT ANSWER

$(1/2)^x = 2^{-x}$, which replaces x by −x.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Show that $2^{x+1} = 2 \cdot 2^x$ and explain what this says about two transformations.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q2/5

1816

QUESTION ID

S49-F1-Q2

TYPE

Short answer

CORRECT ANSWER

$2^{x+1} = 2^x \cdot 2^1$: a left shift by 1 equals a vertical stretch by 2.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$y = 5 \cdot 3^x$ and $y = 3^x$ have different horizontal asymptotes.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q3/5

1817

QUESTION ID

S49-F1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE A vertical stretch keeps the asymptote y = 0 for both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which exponential graph passes through (0, 4) and (1, 2)?

A. $y = 2^x + 3$

B. $y = 4 \cdot 2^x$

C. $y = 0.5 \cdot 4^x$

D. $y = 4(0.5)^x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q4/5

1818

QUESTION ID
S49-F1-Q4

TYPE
Multiple choice

CORRECT ANSWER
D. $y = 4(0.5)^x$

GRADING NOTE a = 4, b = 1/2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write an exponential function with asymptote $y = -1$ and y-intercept 2 that is increasing.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q5/5

1819

QUESTION ID
S49-F1-Q5

TYPE
Short answer

CORRECT ANSWER
$3 \cdot 2^x - 1$ (for example)

GRADING NOTE Any valid function.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 49 • FORWARD 1 • Q5/5

SET 49 • HSF-IF.C.7.e

FORWARD 2

Graph the inverse of an exponential function • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

ANSWER KEY Reduced copy of the question page, with its answer

Write the inverse of $f(x) = 2^x$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q1/5

1821

QUESTION ID

S49-F2-Q1

TYPE

Short answer

CORRECT ANSWER

$\log_2 x$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S49-F2-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

The point (3, 8) is on $y = 2^x$. Which point is on its inverse?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q2/5

1822

QUESTION ID
S49-F2-Q2

TYPE
Short answer

CORRECT ANSWER
(8, 3)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S49-F2-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

The inverse of $y = 2^x$ has the vertical asymptote $x = 0$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q3/5

1823

QUESTION ID
S49-F2-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Reflection of y = 0 over y = x.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q3/5

The inverse of $y = 2^x - 3$ is

A. $y = \log_2(x - 3)$

B. $y = \log_2 x - 3$

C. $y = 2^{x+3}$

D. $y = \log_2(x + 3)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q4/5

1824

QUESTION ID

S49-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. $y = \log_2(x + 3)$

GRADING NOTE $x = 2^y - 3$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the domain and range of $y = \log_2 x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q5/5

1825

QUESTION ID

S49-F2-Q5

TYPE

Short answer

CORRECT ANSWER

Domain x > 0; range all real numbers

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S49-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 49 • FORWARD 2 • Q5/5

SET 50 • HSF-IF.C.7.e • INTERPRETING FUNCTIONS

Graph logarithmic functions with intercept, asymptote, and end behavior

MAIN	Algebra II main question	**HSF-IF.C.7.e**
BACKWARD 1	Graph exponential functions	**HSF-IF.C.7.e**
BACKWARD 2	Find inverses by swapping inputs and outputs	**HSF-BF.B.4.a**
BACKWARD 3	Shift and reflect graphs	**HSF-BF.B.3**
FORWARD 1	Graph logarithms with reflections that change the domain	**HSF-IF.C.7.e**
FORWARD 2	Solve equations with logarithms	**HSF-LE.A.4**


SET 50 • HSF-IF.C.7.e

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_2 x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q1/5

1828

QUESTION ID
S50-M-Q1

TYPE
Drawing / plotting

CORRECT ANSWER

Asymptote x = 0; x-intercept (1, 0); points (1/2, −1), (2, 1), (4, 2), (8, 3); domain x > 0; increases slowly.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_2(x + 3)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q2/5

1829

QUESTION ID

S50-M-Q2

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote $x = -3$; x-intercept $(-2, 0)$; points $(-1, 1)$, $(1, 2)$, $(5, 3)$; domain $x > -3$.


GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q2/5

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_3 x - 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q3/5

1830

QUESTION ID
S50-M-Q3

TYPE
Drawing / plotting

CORRECT ANSWER
Asymptote x = 0; x-intercept (3, 0); points (1, −1), (9, 1), (1/3, −2); domain x > 0.


GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q3/5

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = -\log_2 x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q4/5

1831

QUESTION ID

S50-M-Q4

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote x = 0; x-intercept (1, 0); points (2, −1), (4, −2), (1/2, 1); decreasing; domain x > 0.


GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q4/5

Graph the function on the grid. Show and label two points and the asymptote (dashed); state the domain and whether there is an x-intercept.

$y = \log_2(x - 1) + 2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q5/5

1832

QUESTION ID

S50-M-Q5

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote x = 1; points (2, 2), (3, 3), (5, 4), (1.5, 1); x-intercept (1.25, 0); domain x > 1.

GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • MAIN • Q5/5

SET 50 • HSF-IF.C.7.e

BACKWARD 1

Graph exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Give three points on $y = 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q1/5

1834

QUESTION ID

S50-B1-Q1

TYPE

Short answer

CORRECT ANSWER

For example (0, 1), (1, 2), (2, 4)

GRADING NOTE Any three correct points.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the asymptote of $y = 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q2/5

1835

QUESTION ID

S50-B1-Q2

TYPE

Short answer

CORRECT ANSWER

y = 0

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$y = 3^x$ passes through (3, 1).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q3/5

1836

QUESTION ID

S50-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $3^3 = 27$, not 1; the graph passes through (1, 3) and (0, 1).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = 2^x$ passes through

A. (2, 1)

B. (−1, −2)

C. (1/2, −1)

D. (−1, 1/2)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q4/5

1837

QUESTION ID
S50-B1-Q4

TYPE
Multiple choice

CORRECT ANSWER
D. (−1, 1/2)

GRADING NOTE 2^{-1}.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the range of $y = 3^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q5/5

1838

QUESTION ID

S50-B1-Q5

TYPE

Short answer

CORRECT ANSWER

y > 0

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • BACKWARD 1 • Q5/5

SET 50 • HSF-IF.C.7.e

BACKWARD 2

Find inverses by swapping inputs and outputs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

ANSWER KEY Reduced copy of the question page, with its answer

(2, 4) is on $y = 2^x$. Which point is on its inverse?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q1/5

1840

QUESTION ID

S50-B2-Q1

TYPE

Short answer

CORRECT ANSWER

(4, 2)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Write the inverse of $y = 3^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q2/5

1841

QUESTION ID

S50-B2-Q2

TYPE

Short answer

CORRECT ANSWER

$y = \log_3 x$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

Swapping x and y in $y = 2^x$ gives $x = 2^y$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q3/5

1842

QUESTION ID

S50-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Inverse relation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q3/5

Which point is on $y = \log_2 x$?

A. (3, 8)

B. (2, 4)

C. (8, 3)

D. (0, 1)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q4/5

1843

QUESTION ID

S50-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. (8, 3)

GRADING NOTE $2^3 = 8$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

What is $\log_2 16$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q5/5

1844

QUESTION ID

S50-B2-Q5

TYPE

Short answer

CORRECT ANSWER

4

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B2-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 50 • BACKWARD 2 • Q5/5

SET 50 • HSF-IF.C.7.e

BACKWARD 3

Shift and reflect graphs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

How does $y = \log_2(x + 3)$ relate to $y = \log_2 x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q1/5

1846

QUESTION ID

S50-B3-Q1

TYPE

Short answer

CORRECT ANSWER

Shifted left 3

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

How does $y = -\log_2 x$ relate to $y = \log_2 x$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q2/5

1847

QUESTION ID

S50-B3-Q2

TYPE

Short answer

CORRECT ANSWER

Reflected over the x-axis

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q2/5

True or false?

A horizontal shift moves the vertical asymptote.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q3/5

1848

QUESTION ID

S50-B3-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE x = 0 becomes x = h.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = \log_2(x - 1) + 2$ has asymptote

A. $x = -1$

B. $y = 2$

C. $x = 1$

D. $x = 2$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q4/5

1849

QUESTION ID

S50-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. x = 1

GRADING NOTE Right 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Where does (1, 0) of $y = \log_3 x$ move under $y = \log_3 x - 1$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q5/5

1850

QUESTION ID

S50-B3-Q5

TYPE

Short answer

CORRECT ANSWER

(1, −1)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 50 • BACKWARD 3 • Q5/5

SET 50 • HSF-IF.C.7.e


FORWARD 1

Graph logarithms with reflections that change the domain • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the x-intercept, two more points, and the asymptote (dashed); state the domain.

$y = \log_2(4 - x)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q1/5

1852

QUESTION ID

S50-F1-Q1

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptote x = 4; x-intercept (3, 0); points (2, 1), (0, 2), (−4, 3); decreasing; domain x < 4.

GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the domain of $y = \log(x^2 - 1)$ and explain the graph has two branches.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q2/5

1853

QUESTION ID

S50-F1-Q2

TYPE

Short answer

CORRECT ANSWER

$x < -1$ or $x > 1$: $x^2 - 1 > 0$ on two separate intervals

GRADING NOTE Form required: domain and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S50-F1-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$y = \log_2(2x)$ is $y = \log_2 x$ shifted up 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q3/5

1854

QUESTION ID

S50-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE $\log_2 2 + \log_2 x$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The x-intercept of $y = \log_2(x + 3)$ is

A. -2

B. 1

C. -3

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q4/5

1855

QUESTION ID

S50-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. −2

GRADING NOTE $x + 3 = 1$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the x-intercept of $y = \log_2(x - 1) + 2$ exactly.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q5/5

1856

QUESTION ID

S50-F1-Q5

TYPE

Short answer

CORRECT ANSWER

5/4

GRADING NOTE Any equivalent form. $x - 1 = 2^{-2}$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S50-F1-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 50 • FORWARD 1 • Q5/5

SET 50 • HSF-IF.C.7.e

FORWARD 2

Solve equations with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Solve $2^x = 12$ exactly and to the nearest hundredth. A calculator may be used.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q1/5

1858

GRADING NOTE Form required: exact and decimal.

QUESTION ID

S50-F2-Q1

TYPE

Short answer

CORRECT ANSWER

$\log_2 12 = \log 12/\log 2 \approx 3.58$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $5 \cdot 10^{x} = 40$ exactly.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q2/5

1859

QUESTION ID

S50-F2-Q2

TYPE

Short answer

CORRECT ANSWER

log 8

GRADING NOTE Any equivalent form (≈ 0.903).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q2/5

True or false?

$\log_2 12$ is between 4 and 5.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q3/5

1860

QUESTION ID

S50-F2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $2^3 = 8 < 12 < 16 = 2^4$, so $\log_2 12$ is between 3 and 4.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The solution of $3^x = 7$ is

A. log 7/log 3

B. log 3/log 7

C. 7/3

D. log 21

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q4/5

1861

QUESTION ID

S50-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. log 7/log 3

GRADING NOTE Change of base.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S50-F2-Q4**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Use the graph of $y = \log_2 x$ to estimate $\log_2 6$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S50-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q5/5

1862

QUESTION ID

S50-F2-Q5

TYPE

Short answer

CORRECT ANSWER

About 2.6

GRADING NOTE Accept 2.5 to 2.6 (exact 2.585).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S50-F2-Q5**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 50 • FORWARD 2 • Q5/5

SET 51 • HSF-IF.C.7.e • INTERPRETING FUNCTIONS

Graph trigonometric functions showing period, midline, and amplitude

MAIN	Algebra II main question	**HSF-IF.C.7.e**
BACKWARD 1	Unit-circle values at quarter turns	**HSF-TF.A.2**
BACKWARD 2	Read amplitude, period, and midline from a formula	**HSF-TF.B.5**
BACKWARD 3	Vertical stretches, shifts, and reflections	**HSF-BF.B.3**
FORWARD 1	Graph shifted sinusoids and the tangent function	**HSF-IF.C.7.e**
FORWARD 2	Horizontal stretches f(kx) of trigonometric graphs	**HSF-BF.B.3**

SET 51 • HSF-IF.C.7.e


MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period for one cycle on $0 \le x \le 2\pi$.

$y = 2\sin x$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q1/5

1865

QUESTION ID

S51-M-Q1

TYPE

Drawing / plotting

CORRECT ANSWER

Midline y = 0; amplitude 2; period 2π; key points (0, 0), (π/2, 2), (π, 0), (3π/2, −2), (2π, 0).

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \le x \le 2\pi$.

$y = \cos x + 1$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q2/5

1866

QUESTION ID

S51-M-Q2

TYPE

Drawing / plotting

CORRECT ANSWER

Midline y = 1; amplitude 1; period 2π; key points (0, 2), (π/2, 1), (π, 0), (3π/2, 1), (2π, 2).

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \le x \le 2\pi$.

$y = \sin(2x)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q3/5

1867

QUESTION ID

S51-M-Q3

TYPE

Drawing / plotting

CORRECT ANSWER

Midline y = 0; amplitude 1; period π; maxima at (π/4, 1), (5π/4, 1); minima at (3π/4, −1), (7π/4, −1); zeros at multiples of π/2.

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \le x \le 4\pi$.

$y = 3\cos(x/2)$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q4/5

1868

QUESTION ID

S51-M-Q4

TYPE

Drawing / plotting

CORRECT ANSWER

Midline y = 0; amplitude 3; period 4π; key points (0, 3), (π, 0), (2π, −3), (3π, 0), (4π, 3).

GRADING NOTE Any graph with these features.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the midline (dashed), the maximum and minimum points, and the period on $0 \le x \le 2\pi$.

$y = -\sin x + 2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q5/5

1869

QUESTION ID

S51-M-Q5

TYPE

Drawing / plotting

CORRECT ANSWER

Midline y = 2; amplitude 1; period 2π; key points (0, 2), (π/2, 1), (π, 2), (3π/2, 3), (2π, 2).

GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • MAIN • Q5/5

SET 51 • HSF-IF.C.7.e

BACKWARD 1

Unit-circle values at quarter turns • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Give sin x at x = 0, π/2, π, 3π/2, 2π.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q1/5

1871

QUESTION ID

S51-B1-Q1

TYPE

Short answer

CORRECT ANSWER

0, 1, 0, −1, 0

GRADING NOTE Form required: all five.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give cos x at x = 0, π/2, π, 3π/2, 2π.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q2/5

1872

QUESTION ID

S51-B1-Q2

TYPE

Short answer

CORRECT ANSWER

1, 0, −1, 0, 1

GRADING NOTE Form required: all five.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$\sin(3\pi/2) = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q3/5

1873

QUESTION ID

S51-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The point at $3\pi/2$ is $(0, -1)$, so $\sin(3\pi/2) = -1$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q3/5

cos(π) equals

A. π

B. 1

C. −1

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q4/5

1874

QUESTION ID

S51-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. −1

GRADING NOTE Point (−1, 0).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give sin(π/4).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q5/5

1875

QUESTION ID

S51-B1-Q5

TYPE

Short answer

CORRECT ANSWER

√2/2

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B1-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.A.2 • Set 51 • BACKWARD 1 • Q5/5

SET 51 • HSF-IF.C.7.e

BACKWARD 2

Read amplitude, period, and midline from a formula • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

Give the amplitude and period of y = 3cos(x/2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q1/5

1877

QUESTION ID

S51-B2-Q1

TYPE

Short answer

CORRECT ANSWER

3 and 4π

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the midline of $y = -\sin x + 2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q2/5

1878

QUESTION ID

S51-B2-Q2

TYPE

Short answer

CORRECT ANSWER

$y = 2$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q2/5

True or false?

y = sin(2x) has period 4π.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q3/5

1879

QUESTION ID

S51-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The period is 2π/2 = π.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

y = 2sin x has amplitude

A. 1

B. 2

C. π

D. 2π

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q4/5

1880

QUESTION ID

S51-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 2

GRADING NOTE |A|.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the period of y = cos(3x).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q5/5

1881

QUESTION ID

S51-B2-Q5

TYPE

Short answer

CORRECT ANSWER

2π/3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 51 • BACKWARD 2 • Q5/5

SET 51 • HSF-IF.C.7.e

BACKWARD 3

Vertical stretches, shifts, and reflections • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

Describe y = cos x + 1 compared with y = cos x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q1/5

1883

QUESTION ID

S51-B3-Q1

TYPE

Short answer

CORRECT ANSWER

Up 1

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Describe y = −sin x compared with y = sin x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q2/5

1884

QUESTION ID

S51-B3-Q2

TYPE

Short answer

CORRECT ANSWER

Reflected over the x-axis

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

y = 2sin x has the same zeros as y = sin x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q3/5

1885

QUESTION ID
S51-B3-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Vertical stretches keep zeros.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The maximum of $y = -\sin x + 2$ is

A. 1

B. 2

C. 4

D. 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q4/5

1886

QUESTION ID

S51-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 3

GRADING NOTE $-(-1) + 2$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

What is the minimum of y = cos x + 1?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q5/5

1887

QUESTION ID

S51-B3-Q5

TYPE

Short answer

CORRECT ANSWER

0

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 51 • BACKWARD 3 • Q5/5

SET 51 • HSF-IF.C.7.e


FORWARD 1

Graph shifted sinusoids and the tangent function • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Graph the function on the grid. Show and label the asymptotes (dashed) and three points.

$y = \tan x$ for $-\pi/2 < x < \pi/2$



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q1/5

1889

QUESTION ID

S51-F1-Q1

TYPE

Drawing / plotting

CORRECT ANSWER

Asymptotes $x = \pm\pi/2$; points $(-\pi/4, -1)$, $(0, 0)$, $(\pi/4, 1)$; increasing.

GRADING NOTE Any graph with these features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Describe the graph of $y = \sin(x - \pi/2)$ compared with $y = \sin x$, and name a simpler formula for it.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q2/5

1890

QUESTION ID

S51-F1-Q2

TYPE

Short answer

CORRECT ANSWER

Shifted right π/2; it equals −cos x

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

The period of $y = \tan x$ is 2π.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q3/5

1891

QUESTION ID

S51-F1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $\tan(x + \pi) = \tan x$, so the period is π.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The first maximum of $y = \cos(x - \pi/3)$ for $x \geq 0$ is at

A. $x = 2\pi/3$

B. $x = \pi/3$

C. $x = 0$

D. $x = -\pi/3$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q4/5

1892

QUESTION ID

S51-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. $x = \pi/3$

GRADING NOTE Shift right π/3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the x-values of the maximum points of $y = \sin(x + \pi/4)$ on $0 \leq x < 2\pi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q5/5

1893

QUESTION ID

S51-F1-Q5

TYPE

Short answer

CORRECT ANSWER

$x = \pi/4$

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S51-F1-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 51 • FORWARD 1 • Q5/5

SET 51 • HSF-IF.C.7.e

FORWARD 2

Horizontal stretches f(kx) of trigonometric graphs • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

How is y = sin(2x) obtained from y = sin x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q1/5

1895

QUESTION ID

S51-F2-Q1

TYPE

Short answer

CORRECT ANSWER

Horizontal compression by a factor of 1/2

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

How is y = cos(x/2) obtained from y = cos x?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q2/5

1896

QUESTION ID

S51-F2-Q2

TYPE

Short answer

CORRECT ANSWER

Horizontal stretch by a factor of 2

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q2/5

True or false?

Replacing x by kx (k > 1) shortens the period by the factor k.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q3/5

1897

QUESTION ID
S51-F2-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Period 2π/k.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

y = sin(−x) is the same as

A. cos x

B. sin x

C. −cos x

D. −sin x

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q4/5

1898

QUESTION ID

S51-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. −sin x

GRADING NOTE Sine is odd.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

For which k does y = cos(kx) have period 6π?

k = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q5/5

1899

QUESTION ID

S51-F2-Q5

TYPE

Short answer

CORRECT ANSWER

1/3

GRADING NOTE Any equivalent form (or −1/3).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S51-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 51 • FORWARD 2 • Q5/5

SET 52 • HSF-IF.C.8.b • INTERPRETING FUNCTIONS

Rewrite exponential functions to reveal rates over other time units

MAIN	Algebra II main question	**HSF-IF.C.8.b**
BACKWARD 1	Read growth rates from exponential functions	**HSF-IF.C.8.b**
BACKWARD 2	Powers of powers and rational exponents	**HSN-RN.A.1**
BACKWARD 3	Percent change and growth factors	**7.RP.A.3**
FORWARD 1	Compare growth offers by rewriting to the same time unit	**HSF-IF.C.8.b**
FORWARD 2	Find doubling times and half-lives with logarithms	**HSF-LE.A.4**

SET 52 • HSF-IF.C.8.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

ANSWER KEY Reduced copy of the question page, with its answer

$y = 100(1.08)^t$ with t in years. Rewrite it with a monthly growth factor and give the monthly percent rate to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q1/5

1902

QUESTION ID

S52-M-Q1

TYPE

Short answer

CORRECT ANSWER

$y = 100(1.08^{1/12})^{12t} \approx 100(1.006434)^{12t}$; about 0.64% per month

GRADING NOTE Form required: rewritten form and rate.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S52-M-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = 500(0.97)^t$ with t in years. Rewrite it per decade and give the percent decrease per decade to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q2/5

1903

QUESTION ID

S52-M-Q2

TYPE

Short answer

CORRECT ANSWER

$y = 500(0.97^{10})^{t/10} \approx 500(0.7374)^{t/10}$; about 26.26% decrease per decade

GRADING NOTE Form required: rewritten form and rate.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S52-M-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

$Q = 50(2)^{t/3}$ with t in hours. Rewrite it with an hourly growth factor and give the hourly percent rate to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q3/5

1904

QUESTION ID
S52-M-Q3

TYPE
Short answer

CORRECT ANSWER
$Q = 50(2^{1/3})^t \approx 50(1.2599)^t$; about 25.99% per hour

GRADING NOTE Form required: rewritten form and rate.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$A = 1000(1.005)^{12t}$ with t in years. Rewrite it with an annual growth factor and give the annual percent rate to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q4/5

1905

QUESTION ID
S52-M-Q4

TYPE
Short answer

CORRECT ANSWER
$A = 1000(1.005^{12})^t \approx 1000(1.061678)^t$; about 6.17% per year

GRADING NOTE Form required: rewritten form and rate.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

$N = 800(0.5)^{t/10}$ with t in years. Rewrite it with a yearly factor and give the yearly percent decrease to the nearest hundredth of a percent. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q5/5

1906

QUESTION ID
S52-M-Q5

TYPE
Short answer

CORRECT ANSWER
$N = 800(0.5^{1/10})^t \approx 800(0.933033)^t$; about 6.70% decrease per year

GRADING NOTE Form required: rewritten form and rate.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • MAIN • Q5/5

SET 52 • HSF-IF.C.8.b

BACKWARD 1

Read growth rates from exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b

ANSWER KEY Reduced copy of the question page, with its answer

Is $y = 100(1.08)^t$ growth or decay, and at what percent per year?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q1/5

1908

QUESTION ID

S52-B1-Q1

TYPE

Short answer

CORRECT ANSWER

Growth, 8% per year

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Is $y = 500(0.97)^t$ growth or decay, and at what percent per year?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q2/5

1909

QUESTION ID

S52-B1-Q2

TYPE

Short answer

CORRECT ANSWER

Decay, 3% per year

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$y = 2(1.25)^t$ grows 125% per unit of t.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q3/5

1910

QUESTION ID

S52-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The factor 1.25 means 25% growth.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = 40(0.9)^t$ decreases by

A. 9% per unit

B. 10% per unit

C. 90% per unit

D. 0.9% per unit

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q4/5

1911

QUESTION ID

S52-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 10% per unit

GRADING NOTE $1 - 0.9$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

What is the growth factor for 6.17% growth?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q5/5

1912

QUESTION ID

S52-B1-Q5

TYPE

Short answer

CORRECT ANSWER

1.0617

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • BACKWARD 1 • Q5/5

SET 52 • HSF-IF.C.8.b

BACKWARD 2

Powers of powers and rational exponents • High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1

ANSWER KEY Reduced copy of the question page, with its answer

Rewrite 1.08^{t} as a power of $1.08^{1/12}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q1

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q1/5

1914

QUESTION ID

S52-B2-Q1

TYPE

Short answer

CORRECT ANSWER

$(1.08^{1/12})^{12t}$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q1

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Rewrite $2^{t/3}$ as $(2^{1/3})^t$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q2

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q2/5

1915

QUESTION ID

S52-B2-Q2

TYPE

Short answer

CORRECT ANSWER

$(2^{1/3})^t$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S52-B2-Q2**

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q2/5

True or false?

$(b^m)^n = b^{mn}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q3

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q3/5

1916

QUESTION ID

S52-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Power of a power.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q3

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

1.005^{12t} equals

A. $(1.005^{12})^{t}$

B. $(1.005^{t})/12$

C. $12 \cdot 1.005^{t}$

D. $1.005^{12} + t$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q4

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q4/5

1917

QUESTION ID

S52-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. $(1.005^{12})^{t}$

GRADING NOTE Power of a power.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q4

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write $8^{1/3}$ as a whole number.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q5

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q5/5

1918

QUESTION ID

S52-B2-Q5

TYPE

Short answer

CORRECT ANSWER

2

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B2-Q5

High School — Algebra I • Number & Quantity: The Real Number System • HSN-RN.A.1 • Set 52 • BACKWARD 2 • Q5/5

SET 52 • HSF-IF.C.8.b

BACKWARD 3

Percent change and growth factors • Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3

ANSWER KEY Reduced copy of the question page, with its answer

Write the growth factor for a 25.99% increase.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q1/5

1920

QUESTION ID

S52-B3-Q1

TYPE

Short answer

CORRECT ANSWER

1.2599

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q1

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A factor 0.7374 means what percent decrease?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q2/5

1921

QUESTION ID

S52-B3-Q2

TYPE

Short answer

CORRECT ANSWER

26.26%

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q2

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

A factor 1.006434 means about 0.64% growth.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q3/5

1922

QUESTION ID

S52-B3-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Subtract 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q3

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A 3% decrease corresponds to the factor

A. 0.03

B. 0.97

C. 0.7

D. 1.03

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q4/5

1923

QUESTION ID

S52-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 0.97

GRADING NOTE 1 − 0.03.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q4

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write the percent change for a factor of 0.933.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q5/5

1924

QUESTION ID

S52-B3-Q5

TYPE

Short answer

CORRECT ANSWER

6.7% decrease

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-B3-Q5

Grade 7 • Ratios & Proportional Relationships • 7.RP.A.3 • Set 52 • BACKWARD 3 • Q5/5

SET 52 • HSF-IF.C.8.b

FORWARD 1

Compare growth offers by rewriting to the same time unit • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

ANSWER KEY Reduced copy of the question page, with its answer

Account A grows 0.5% per month. Account B grows 6.1% per year. Which grows faster? Rewrite to compare. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q1/5

1926

QUESTION ID

S52-F1-Q1

TYPE

Short answer

CORRECT ANSWER

A: $1.005^{12} \approx 1.0617$, about 6.17% per year > 6.1%, so A grows faster

GRADING NOTE Form required: comparison on a common unit.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A culture doubles every 3 hours; another triples every 5 hours. Which grows faster per hour? A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q2/5

1927

QUESTION ID
S52-F1-Q2

TYPE
Short answer

CORRECT ANSWER
$2^{1/3} \approx 1.260$ vs $3^{1/5} \approx 1.246$; the first grows faster

GRADING NOTE Form required: both hourly factors and the decision.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q2/5

True or false?

Growing 1% per month is the same as growing 12% per year.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q3/5

1928

QUESTION ID

S52-F1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $1.01^{12} \approx 1.1268$, about 12.68%.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which is the greatest annual growth?

A. 4% per half year

B. 0.6% per month

C. 8% per year

D. 2% per quarter

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q4/5

1929

QUESTION ID

S52-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 2% per quarter

GRADING NOTE $1.02^4 \approx 1.0824$; $1.006^{12} \approx 1.0744$; $1.04^2 = 1.0816$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A half-life of 10 years and a decay of 7% per year: which decays faster? A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q5/5

1930

QUESTION ID
S52-F1-Q5

TYPE
Short answer

CORRECT ANSWER
7% per year (0.93 < 0.933 per year)

GRADING NOTE Form required: decision and factors.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 52 • FORWARD 1 • Q5/5

SET 52 • HSF-IF.C.8.b

FORWARD 2

Find doubling times and half-lives with logarithms • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Find the doubling time of $y = 100(1.08)^t$, to the nearest hundredth of a year. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q1/5

1932

QUESTION ID
S52-F2-Q1

TYPE
Short answer

CORRECT ANSWER
9.01 years

GRADING NOTE Any equivalent form. log 2/log 1.08.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the half-life of $y = 500(0.97)^t$ to the nearest hundredth of a year. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q2/5

1933

QUESTION ID
S52-F2-Q2

TYPE
Short answer

CORRECT ANSWER
22.76 years

GRADING NOTE Any equivalent form. log 0.5/log 0.97.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q2/5

True or false?

The doubling time of $y = 50(2)^{t/3}$ is 6 hours.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q3/5

1934

QUESTION ID

S52-F2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The exponent t/3 increases by 1 every 3 hours, which doubles y.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q3/5

The doubling time of a quantity growing 6.17% per year is about

A. 16.2 years

B. 11.6 years

C. 6.17 years

D. 3.1 years

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q4/5

1935

QUESTION ID

S52-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 11.6 years

GRADING NOTE log 2/log 1.0617 ≈ 11.58.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S52-F2-Q4**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why rewriting $y = 100(1.08)^t$ as $100 \cdot 2^{t/9.01}$ shows the doubling time.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q5/5

1936

QUESTION ID
S52-F2-Q5

TYPE
Short answer

CORRECT ANSWER
The exponent t/9.01 increases by 1 every 9.01 years, which multiplies y by 2.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S52-F2-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 52 • FORWARD 2 • Q5/5

SET 53 • HSF-IF.C.9 • INTERPRETING FUNCTIONS

Compare properties of Algebra II functions in different representations

MAIN	Algebra II main question	**HSF-IF.C.9**
BACKWARD 1	Compare linear, quadratic, and exponential functions	**HSF-IF.C.9**
BACKWARD 2	Recognize a pattern in a table	**8.F.A.3**
BACKWARD 3	Domains, asymptotes, and periods of Algebra II families	**HSF-IF.B.5**
FORWARD 1	Compare rates of change and long-run behavior across represent…	**HSF-IF.C.9**
FORWARD 2	Build a difference function to compare	**HSF-BF.A.1.b**

SET 53 • HSF-IF.C.9

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 3(1.1)^x$. The table shows g. Which has the greater y-intercept, and which is greater at x = 20? A calculator may be used.

x	0	1	2	3	4
g(x)	10	15	20	25	30

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q1/5

1939

QUESTION ID

S53-M-Q1

TYPE

Short answer

CORRECT ANSWER

g has the greater y-intercept (10 > 3); at x = 20, f(20) ≈ 20.18 < g(20) = 110, so g is greater


GRADING NOTE Form required: both comparisons with values.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = \sqrt{(x - 1)}$. The graph of g is shown. Compare their domains.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q2/5

1940

QUESTION ID

S53-M-Q2

TYPE

Short answer

CORRECT ANSWER

f: x ≥ 1; g: all real numbers (g is a cube root graph)

GRADING NOTE Form required: both domains.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q2/5

h(t) = 12 + 10sin(πt/15) gives one Ferris wheel's height (m, t in s). A second wheel has radius 8 m, center 10 m high, and turns once every 40 s. Compare the maximum heights and the periods.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q3/5

1941

QUESTION ID

S53-M-Q3

TYPE

Short answer

CORRECT ANSWER

Maximum heights 22 m and 18 m; periods 30 s and 40 s; the first wheel is higher and faster

GRADING NOTE Form required: both comparisons.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S53-M-Q3**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = \log_2 x$. The table shows g. Which is greater at $x = 9$?

x	1	3	9	27
g(x)	0	1	2	3

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q4/5

1942

QUESTION ID

S53-M-Q4

TYPE

Short answer

CORRECT ANSWER

f(9) ≈ 3.17 > g(9) = 2


GRADING NOTE Form required: answer with values.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

f(x) = (2x + 1)/(x − 3). The graph of g shows the horizontal asymptote y = 1 and the vertical asymptote x = −2. Compare the asymptotes.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q5/5

1943

QUESTION ID

S53-M-Q5

TYPE

Short answer

CORRECT ANSWER

f: horizontal y = 2, vertical x = 3; g: horizontal y = 1, vertical x = −2

GRADING NOTE Form required: both pairs.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • MAIN • Q5/5

SET 53 • HSF-IF.C.9

BACKWARD 1

Compare linear, quadratic, and exponential functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9

ANSWER KEY Reduced copy of the question page, with its answer

f(x) = 2x + 1 and g has table g(0) = 3, g(1) = 4. Which has the greater y-intercept?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q1/5

1945

QUESTION ID

S53-B1-Q1

TYPE

Short answer

CORRECT ANSWER

g (3 > 1)

GRADING NOTE Form required: answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = x^2$ and $g(x) = 2^x$. Which is greater at $x = 10$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q2/5

1946

QUESTION ID

S53-B1-Q2

TYPE

Short answer

CORRECT ANSWER

g: 1024 > 100

GRADING NOTE Form required: answer with values.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q2/5

True or false?

An increasing exponential eventually exceeds any linear function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q3/5

1947

QUESTION ID

S53-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Algebra I result.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = -(x - 1)^2 + 5$ and g with maximum 4 in a graph. The greater maximum is

A. f: 5

B. cannot tell

C. equal

D. g: 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q4/5

1948

QUESTION ID

S53-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. f: 5

GRADING NOTE Vertex form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A table rises by 5 each step. What kind of function is it?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q5/5

1949

QUESTION ID

S53-B1-Q5

TYPE

Short answer

CORRECT ANSWER

Linear

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B1-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • BACKWARD 1 • Q5/5

SET 53 • HSF-IF.C.9

BACKWARD 2

Recognize a pattern in a table • Grade 8 • Functions • 8.F.A.3

ANSWER KEY Reduced copy of the question page, with its answer

Is the table x: 0, 1, 2; y: 10, 15, 20 linear? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q1

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q1/5

1951

QUESTION ID

S53-B2-Q1

TYPE

Short answer

CORRECT ANSWER

Yes: constant difference 5

GRADING NOTE Form required: decision and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q1

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Is the table x: 1, 3, 9, 27; y: 0, 1, 2, 3 linear? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q2

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q2/5

1952

QUESTION ID

S53-B2-Q2

TYPE

Short answer

CORRECT ANSWER

No: equal output steps need inputs multiplied by 3 (logarithmic)

GRADING NOTE Form required: decision and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q2

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q2/5

True or false?

A table with constant ratios of outputs is exponential.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q3

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q3/5

1953

QUESTION ID

S53-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Repeated multiplication.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q3

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q3/5

Which table is exponential?

A. 1, 4, 9, 16

B. 1, 2, 4, 8

C. 2, 2, 2, 2

D. 1, 3, 5, 7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q4

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q4/5

1954

QUESTION ID

S53-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 1, 2, 4, 8

GRADING NOTE Ratio 2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S53-B2-Q4

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

What is the slope of a linear table with outputs 10, 15, 20 at x = 0, 1, 2?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q5

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q5/5

1955

QUESTION ID

S53-B2-Q5

TYPE

Short answer

CORRECT ANSWER

5

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B2-Q5

Grade 8 • Functions • 8.F.A.3 • Set 53 • BACKWARD 2 • Q5/5

SET 53 • HSF-IF.C.9

BACKWARD 3

Domains, asymptotes, and periods of Algebra II families • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

Give the domain of $\sqrt{(x - 1)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q1/5

1957

QUESTION ID

S53-B3-Q1

TYPE

Short answer

CORRECT ANSWER

$x \geq 1$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the asymptotes of (2x + 1)/(x − 3).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q2/5

1958

QUESTION ID

S53-B3-Q2

TYPE

Short answer

CORRECT ANSWER

x = 3 and y = 2

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q2/5

True or false?

The period of $12 + 10\sin(\pi t/15)$ is 15.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q3/5

1959

QUESTION ID

S53-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $2\pi/(\pi/15) = 30$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The domain of $\log_2 x$ is

A. all reals

B. $x \geq 0$

C. $x > 0$

D. $x \neq 0$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q4/5

1960

QUESTION ID

S53-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. x > 0

GRADING NOTE Positive inputs.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A wheel of radius 8 m with center 10 m high has what maximum height?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q5/5

1961

QUESTION ID

S53-B3-Q5

TYPE

Short answer

CORRECT ANSWER

18 m

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 53 • BACKWARD 3 • Q5/5

SET 53 • HSF-IF.C.9

FORWARD 1

Compare rates of change and long-run behavior across representations • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 3(1.1)^x$ and $g(x) = 10 + 5x$. Estimate the first whole number x at which f exceeds g. A calculator may be used. A calculator may be used.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q1/5

1963

QUESTION ID

S53-F1-Q1

TYPE

Short answer

CORRECT ANSWER

46

GRADING NOTE Any equivalent form. f(45) ≈ 218.7 < g(45) = 235; f(46) ≈ 240.6 > g(46) = 240.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q1/5

$f(x) = \log_2 x$, and g has g(1) = 0, g(3) = 1, g(9) = 2. Compare the average rates of change of f and g on $1 \le x \le 9$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q2/5

1964

QUESTION ID

S53-F1-Q2

TYPE

Short answer

CORRECT ANSWER

f: (3.17 − 0)/8 ≈ 0.40; g: 2/8 = 0.25; f is greater

GRADING NOTE Form required: both rates and the comparison.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q2/5

True or false?

A function with a larger y-intercept is always larger.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q3/5

1965

QUESTION ID
S53-F1-Q3

TYPE
True or false

CORRECT ANSWER
B. False

GRADING NOTE g starts higher but f eventually passes it.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which grows fastest for large x?

A. $\sqrt{(x - 1)}$

B. $3(1.1)^x$

C. $5x + 10$

D. $\log_2 x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q4/5

1966

QUESTION ID

S53-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. $3(1.1)^x$

GRADING NOTE Exponential.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S53-F1-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why $g(x) = \log_3 x$ stays below $f(x) = \log_2 x$ for $x > 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q5/5

1967

QUESTION ID

S53-F1-Q5

TYPE

Short answer

CORRECT ANSWER

$\log_3 x = \log_2 x / \log_2 3$ and $\log_2 3 > 1$, so g is a smaller positive multiple of f.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.9 • Set 53 • FORWARD 1 • Q5/5

SET 53 • HSF-IF.C.9

FORWARD 2

Build a difference function to compare • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b

ANSWER KEY Reduced copy of the question page, with its answer

Write $d(x) = f(x) - g(x)$ for $f(x) = 3(1.1)^x$ and $g(x) = 5x + 10$.

$d(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q1/5

1969

QUESTION ID

S53-F2-Q1

TYPE

Short answer

CORRECT ANSWER

$3(1.1)^x - 5x - 10$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S53-F2-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain what the sign of d(x) tells you.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q2/5

1970

QUESTION ID

S53-F2-Q2

TYPE

Short answer

CORRECT ANSWER

d(x) > 0 means f is greater; d(x) < 0 means g is greater.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q2/5

True or false?

A zero of d(x) = f(x) − g(x) is an x where f(x) = g(x).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q3/5

1971

QUESTION ID
S53-F2-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Equal outputs.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

For $h_1(t) = 12 + 10\sin(\pi t/15)$ and $h_2(t) = 10 + 8\sin(\pi t/20)$, $h_1(0) - h_2(0)$ is

A. 2

B. 22

C. −2

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q4/5

1972

QUESTION ID

S53-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 2

GRADING NOTE 12 − 10.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Two Ferris wheel heights are $h_1(t) = 12 + 10\sin(\pi t/15)$ and $h_2(t) = 10 + 8\sin(\pi t/20)$. Write $h_1(t) - h_2(t)$ as a function of t.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q5/5

1973

QUESTION ID
S53-F2-Q5

TYPE
Short answer

CORRECT ANSWER
$h_1(t) - h_2(t) = 2 + 10\sin(\pi t/15) - 8\sin(\pi t/20)$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S53-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 53 • FORWARD 2 • Q5/5

Build models by combining Algebra II function types

MAIN	Algebra II main question	**HSF-BF.A.1.b**
BACKWARD 1	Combine linear and quadratic functions	**HSF-BF.A.1.b**
BACKWARD 2	Multiply and subtract polynomials	**HSA-APR.A.1**
BACKWARD 3	Evaluate exponential and trigonometric expressions	**HSF-IF.A.2**
FORWARD 1	Interpret the parts of a combined model	**HSF-BF.A.1.b**
FORWARD 2	Compose functions in context	**HSF-BF.A.1.c**

SET 54 • HSF-BF.A.1.b

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b

Revenue is $R(x) = x(50 - 0.5x)$ and cost is $C(x) = 200 + 10x$ dollars for x items. Write the profit $P(x)$ in standard form and find $P(30)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q1/5

1976

QUESTION ID

S54-M-Q1

TYPE

Short answer

CORRECT ANSWER

$P(x) = -0.5x^2 + 40x - 200$; $P(30) = 550$

GRADING NOTE Form required: function and value.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

The cost of x posters is C(x) = 1200 + 3x dollars. Write the average cost per poster A(x) and find A(100).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q2/5

1977

QUESTION ID

S54-M-Q2

TYPE

Short answer

CORRECT ANSWER

A(x) = (1200 + 3x)/x; A(100) = 15

GRADING NOTE Form required: function and value with units ($15 per poster).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-M-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

A town has $P(t) = 5000(1.02)^t$ people and $F(t) = 6000 + 100t$ units of food after t years. Write the food per person and find it at t = 10 to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q3/5

1978

QUESTION ID
S54-M-Q3

TYPE
Short answer

CORRECT ANSWER
$F(t)/P(t) = (6000 + 100t)/(5000(1.02)^t)$; about 1.15 units per person

GRADING NOTE Form required: function and value.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A spring's amplitude is 10 cm and shrinks by 10% each second, while it oscillates once per second. Write y(t) as an exponential times a cosine and find y(1).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q4/5

1979

QUESTION ID
S54-M-Q4

TYPE
Short answer

CORRECT ANSWER

$y(t) = 10(0.9)^t \cos(2\pi t)$; $y(1) = 9$

GRADING NOTE Form required: function and value.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-M-Q4**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Coffee starts 70°C above a 20°C room, and the difference shrinks 5% each minute. Write its temperature T(t) as a sum of functions and find T(10) to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q5/5

1980

QUESTION ID

S54-M-Q5

TYPE

Short answer

CORRECT ANSWER

$T(t) = 20 + 70(0.95)^t$; $T(10) \approx 61.91°C$

GRADING NOTE Form required: function and value.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • MAIN • Q5/5

SET 54 • HSF-BF.A.1.b

BACKWARD 1

Combine linear and quadratic functions • High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b

ANSWER KEY Reduced copy of the question page, with its answer

R(x) = 8x and C(x) = 3x + 50. Write P = R − C.

P(x) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q1/5

1982

QUESTION ID

S54-B1-Q1

TYPE

Short answer

CORRECT ANSWER

5x − 50

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

f(x) = x + 2 and g(x) = x − 2. Write f · g.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q2/5

1983

QUESTION ID

S54-B1-Q2

TYPE

Short answer

CORRECT ANSWER

$x^2 - 4$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q2/5

True or false?

Profit is revenue minus cost.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q3/5

1984

QUESTION ID

S54-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q3/5

Average cost is

A. total cost times items

B. total cost divided by the number of items

C. cost minus revenue

D. revenue divided by cost

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q4/5

1985

QUESTION ID

S54-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. total cost divided by the number of items

GRADING NOTE Per item.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 2x$ and $g(x) = x^2 + 1$. Find $(f + g)(3)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q5/5

1986

QUESTION ID

S54-B1-Q5

TYPE

Short answer

CORRECT ANSWER

16

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • BACKWARD 1 • Q5/5

SET 54 • HSF-BF.A.1.b

BACKWARD 2

Multiply and subtract polynomials • High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1

ANSWER KEY Reduced copy of the question page, with its answer

Expand x(50 − 0.5x).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q1

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q1/5

1988

QUESTION ID

S54-B2-Q1

TYPE

Short answer

CORRECT ANSWER

$50x - 0.5x^2$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-B2-Q1**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Simplify $(50x - 0.5x^2) - (200 + 10x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q2/5

1989

QUESTION ID

S54-B2-Q2

TYPE

Short answer

CORRECT ANSWER

$-0.5x^2 + 40x - 200$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q2

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$-(200 + 10x) = -200 + 10x.$

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q3/5

1990

QUESTION ID

S54-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The minus sign applies to both terms: $-200 - 10x$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q3

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

(3x + 1) − (x − 4) equals

A. 2x + 5

B. 4x − 3

C. 2x − 3

D. 2x + 3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q4/5

1991

QUESTION ID

S54-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 2x + 5

GRADING NOTE Subtract each term.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q4

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Expand $2(x^2 - 3x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B2-Q5

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q5/5

1992

QUESTION ID

S54-B2-Q5

TYPE

Short answer

CORRECT ANSWER

$2x^2 - 6x$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-B2-Q5**

High School — Algebra I • Algebra: Arithmetic with Polynomials & Rational Expressions • HSA-APR.A.1 • Set 54 • BACKWARD 2 • Q5/5

SET 54 • HSF-BF.A.1.b

BACKWARD 3

Evaluate exponential and trigonometric expressions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $70(0.95)^{10}$ to the nearest hundredth. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q1/5

1994

QUESTION ID

S54-B3-Q1

TYPE

Short answer

CORRECT ANSWER

41.91

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $5000(1.02)^{10}$ to the nearest whole number. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q2/5

1995

QUESTION ID

S54-B3-Q2

TYPE

Short answer

CORRECT ANSWER

6095

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q2/5

True or false?

$\cos(2\pi) = 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q3/5

1996

QUESTION ID

S54-B3-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Full turn.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$10(0.9)^2$ equals

A. 9

B. 18

C. 81

D. 8.1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q4/5

1997

QUESTION ID

S54-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 8.1

GRADING NOTE $10 \cdot 0.81$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $\cos(2\pi \cdot 0.5)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q5/5

1998

QUESTION ID

S54-B3-Q5

TYPE

Short answer

CORRECT ANSWER

−1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 54 • BACKWARD 3 • Q5/5

SET 54 • HSF-BF.A.1.b

FORWARD 1

Interpret the parts of a combined model • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b

ANSWER KEY Reduced copy of the question page, with its answer

In $T(t) = 20 + 70(0.95)^t$, what do 20 and $70(0.95)^t$ represent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q1/5

2000

QUESTION ID
S54-F1-Q1

TYPE
Short answer

CORRECT ANSWER
20: room temperature; $70(0.95)^t$: how far the coffee is above room temperature

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q1/5

2000

ANSWER KEY Reduced copy of the question page, with its answer

In $y(t) = 10(0.9)^t \cos(2\pi t)$, which factor controls the size of the swings and which controls their timing?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q2/5

2001

QUESTION ID
S54-F1-Q2

TYPE
Short answer

CORRECT ANSWER
$10(0.9)^t$ controls the size; $\cos(2\pi t)$ the timing

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

In A(x) = (1200 + 3x)/x = 1200/x + 3, the term 1200/x is the fixed cost spread over x posters.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q3/5

2002

QUESTION ID

S54-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Split the fraction.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

As t grows, $T(t) = 20 + 70(0.95)^t$ approaches

A. 20

B. 0

C. 90

D. 70

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q4/5

2003

QUESTION ID

S54-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 20

GRADING NOTE The exponential part → 0.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why food per person F(t)/P(t) eventually decreases.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q5/5

2004

QUESTION ID

S54-F1-Q5

TYPE

Short answer

CORRECT ANSWER

P grows exponentially while F grows linearly, so the denominator eventually grows much faster.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.A.1.b • Set 54 • FORWARD 1 • Q5/5

SET 54 • HSF-BF.A.1.b

FORWARD 2

Compose functions in context • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c

ANSWER KEY Reduced copy of the question page, with its answer

A balloon's radius is r(t) = 2t cm after t seconds and its volume is V(r) = (4/3)πr³. Write V(r(t)).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q1/5

2006

QUESTION ID
S54-F2-Q1

TYPE
Short answer

CORRECT ANSWER
(32/3)πt³

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A store takes 20% off with d(p) = 0.8p and then adds a $5 fee with f(p) = p + 5. Write f(d(p)) and find it for p = 50.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q2/5

2007

QUESTION ID

S54-F2-Q2

TYPE

Short answer

CORRECT ANSWER

0.8p + 5; $45

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S54-F2-Q2**

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

f(g(x)) and g(f(x)) are always equal.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q3/5

2008

QUESTION ID

S54-F2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Order matters.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

If $f(x) = x^2$ and $g(x) = x + 1$, then $f(g(2)) =$

A. 4

B. 9

C. 3

D. 5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q4/5

2009

QUESTION ID

S54-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 9

GRADING NOTE f(3).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q4/5

Explain how composition differs from multiplying two functions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q5/5

2010

QUESTION ID
S54-F2-Q5

TYPE
Short answer

CORRECT ANSWER
Composition feeds the output of one function in as the input of the other; multiplication multiplies the two outputs.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S54-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.A.1.c • Set 54 • FORWARD 2 • Q5/5

Identify vertical shifts, vertical stretches, and horizontal shifts of Algebra II functions

MAIN	Algebra II main question	**HSF-BF.B.3**
BACKWARD 1	Shifts and stretches of linear and quadratic graphs	**HSF-BF.B.3**
BACKWARD 2	Translate points in the plane	**8.G.A.3**
BACKWARD 3	Key points of parent functions	**HSF-IF.C.7.e**
FORWARD 1	Write transformed functions from descriptions and order of transfo..	**HSF-BF.B.3**
FORWARD 2	Read sinusoid parameters as transformations	**HSF-TF.B.5**

SET 55 • HSF-BF.B.3


MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = \sqrt{x}$. The graph of $g(x) = f(x - h) + k$ is shown. Find h and k.

Answer: _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q1/5

2013

QUESTION ID
S55-M-Q1

TYPE
Short answer

CORRECT ANSWER
h = 2, k = 1

GRADING NOTE Form required: both. The starting point moves from (0, 0) to (2, 1).


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 2^x$. The graph of $g(x) = a \cdot f(x)$ is shown. Find a.

a = _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q2/5

2014

QUESTION ID
S55-M-Q2

TYPE
Short answer

CORRECT ANSWER
3

GRADING NOTE Any equivalent form. g(0) = a.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q2/5

2014

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = x^3$. The graph of $g(x) = f(x - h) + k$ is shown. Find h and k.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q3/5

2015

QUESTION ID

S55-M-Q3

TYPE

Short answer

CORRECT ANSWER

h = 1, k = −2

GRADING NOTE Form required: both.


REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q3/5

2015

ANSWER KEY Reduced copy of the question page, with its answer

f(x) = |x|. The graph of g(x) = a·f(x − h) + k is shown. Find a, h, and k.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q4/5

2016

QUESTION ID

S55-M-Q4

TYPE

Short answer

CORRECT ANSWER

a = −1, h = −2, k = 3

GRADING NOTE Form required: all three.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q4


High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q4/5

2016

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = \log_2 x$. The graph of $g(x) = f(x) + k$ is shown. Find k.

$k =$ _____



REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q5/5

2017

QUESTION ID
S55-M-Q5

TYPE
Short answer

CORRECT ANSWER
2

GRADING NOTE Any equivalent form. f(1) = 0 becomes 2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • MAIN • Q5/5

2017

SET 55 • HSF-BF.B.3

BACKWARD 1

Shifts and stretches of linear and quadratic graphs • High School — Algebra I • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

How is $y = (x - 3)^2 + 1$ obtained from $y = x^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q1/5

2019

QUESTION ID

S55-B1-Q1

TYPE

Short answer

CORRECT ANSWER

Right 3, up 1

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q1/5

2019

ANSWER KEY Reduced copy of the question page, with its answer

How is $y = 2x^2$ obtained from $y = x^2$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q2/5

2020

QUESTION ID

S55-B1-Q2

TYPE

Short answer

CORRECT ANSWER

Vertical stretch by 2

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q2/5

2020

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

y = f(x + 4) is the graph of f shifted right 4.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q3/5

2021

QUESTION ID

S55-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Left 4.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The vertex of $y = (x + 2)^2 - 5$ is

A. (2, 5)

B. (2, −5)

C. (−2, −5)

D. (−2, 5)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q4/5

2022

QUESTION ID

S55-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. (−2, −5)

GRADING NOTE $h = -2$, $k = -5$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q4/5

2022

ANSWER KEY Reduced copy of the question page, with its answer

Write $y = x^2$ shifted left 1 and down 4.

$y =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q5/5

2023

QUESTION ID

S55-B1-Q5

TYPE

Short answer

CORRECT ANSWER

$(x + 1)^2 - 4$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.3 • Set 55 • BACKWARD 1 • Q5/5

2023

SET 55 • HSF-BF.B.3

BACKWARD 2

Translate points in the plane • Grade 8 • Geometry • 8.G.A.3

ANSWER KEY Reduced copy of the question page, with its answer

Translate (0, 0) right 2 and up 1.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q1

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q1/5

2025

QUESTION ID

S55-B2-Q1

TYPE

Short answer

CORRECT ANSWER

(2, 1)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q1

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q1/5

2025

ANSWER KEY Reduced copy of the question page, with its answer

Translate (1, 1) left 2 and up 3.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q2

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q2/5

2026

QUESTION ID

S55-B2-Q2

TYPE

Short answer

CORRECT ANSWER

(−1, 4)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q2

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q2/5

2026

True or false?

Translating (0, 0) down 2 gives (−2, 0).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q3

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q3/5

2027

QUESTION ID

S55-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Down 2 changes y: (0, −2).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q3

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$(x, y) \to (x - 2, y + 3)$ moves $(0, 0)$ to

A. $(-2, 3)$

B. $(-2, -3)$

C. $(2, 3)$

D. $(3, -2)$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q4

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q4/5

2028

QUESTION ID

S55-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. (−2, 3)

GRADING NOTE Apply the rule.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q4

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q4/5

2028

ANSWER KEY Reduced copy of the question page, with its answer

Which translation takes (0, 0) to (1, −2)?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q5

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q5/5

2029

QUESTION ID

S55-B2-Q5

TYPE

Short answer

CORRECT ANSWER

Right 1, down 2

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B2-Q5

Grade 8 • Geometry • 8.G.A.3 • Set 55 • BACKWARD 2 • Q5/5

SET 55 • HSF-BF.B.3

BACKWARD 3

Key points of parent functions • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Give the point of $y = \sqrt{x}$ at $x = 0$ and at $x = 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q1/5

2031

QUESTION ID

S55-B3-Q1

TYPE

Short answer

CORRECT ANSWER

(0, 0) and (1, 1)

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the y-intercept of $y = 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q2/5

2032

QUESTION ID

S55-B3-Q2

TYPE

Short answer

CORRECT ANSWER

1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q2/5

True or false?

The graph of $y = \log_2 x$ passes through (1, 0).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q3/5

2033

QUESTION ID

S55-B3-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE $\log_2 1 = 0$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The vertex of y = |x| is

A. (1, 0)

B. (1, 1)

C. (0, 1)

D. (0, 0)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q4/5

2034

QUESTION ID

S55-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. (0, 0)

GRADING NOTE Parent.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the center (point of symmetry) of $y = x^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q5/5

2035

QUESTION ID
S55-B3-Q5

TYPE
Short answer

CORRECT ANSWER
(0, 0)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 55 • BACKWARD 3 • Q5/5

SET 55 • HSF-BF.B.3

FORWARD 1

Write transformed functions from descriptions and order of transformations • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

Write g, the graph of $f(x) = 2^x$ stretched vertically by 3, then shifted down 4.

g(x) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q1/5

2037

QUESTION ID

S55-F1-Q1

TYPE

Short answer

CORRECT ANSWER

$3 \cdot 2^x - 4$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Write g, the graph of f(x) = √x shifted down 4, then stretched vertically by 3.

g(x) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q2/5

2038

QUESTION ID
S55-F1-Q2

TYPE
Short answer

CORRECT ANSWER
3√x − 12

GRADING NOTE Any equivalent form. 3(√x − 4).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S55-F1-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q2/5

True or false?

A vertical stretch then a shift can give a different graph than the same shift then the stretch.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q3/5

2039

QUESTION ID

S55-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE See the two items above.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$g(x) = 2f(x) + 1$ moves the point (3, 4) of f to

A. (6, 5)

B. (4, 9)

C. (3, 9)

D. (3, 10)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q4/5

2040

QUESTION ID

S55-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. (3, 9)

GRADING NOTE $2 \cdot 4 + 1$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

$g(x) = f(x + 2) - 3$ and f has asymptote $x = 0$. Where is g's vertical asymptote?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q5/5

2041

QUESTION ID

S55-F1-Q5

TYPE

Short answer

CORRECT ANSWER

$x = -2$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 55 • FORWARD 1 • Q5/5

SET 55 • HSF-BF.B.3

FORWARD 2

Read sinusoid parameters as transformations • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

Describe y = 3sin x + 2 as transformations of y = sin x and give its maximum.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q1/5

2043

QUESTION ID

S55-F2-Q1

TYPE

Short answer

CORRECT ANSWER

Vertical stretch by 3, up 2; maximum 5

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Write the function for y = cos x stretched vertically by 4 and shifted down 1.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q2/5

2044

QUESTION ID

S55-F2-Q2

TYPE

Short answer

CORRECT ANSWER

4cos x − 1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

In y = A sin x + D, A is the midline.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q3/5

2045

QUESTION ID

S55-F2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE D gives the midline y = D; |A| is the amplitude.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$y = -2\cos x + 5$ has amplitude and midline

A. 5 and $y = 2$

B. -2 and $y = 5$

C. 2 and $y = -5$

D. 2 and $y = 5$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q4/5

2046

QUESTION ID

S55-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 2 and y = 5

GRADING NOTE |A|.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A model is y = A sin x + D with maximum 9 and minimum 1. Find A (A > 0) and D.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q5/5

2047

QUESTION ID
S55-F2-Q5

TYPE
Short answer

CORRECT ANSWER
A = 4, D = 5

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S55-F2-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 55 • FORWARD 2 • Q5/5

Horizontal stretches, compressions, and reflections f(kx)

MAIN	Algebra II main question	**HSF-BF.B.3**
BACKWARD 1	Reflections and dilations of points	**8.G.A.3**
BACKWARD 2	Solve kx = c for x	**6.EE.B.7**
BACKWARD 3	Period of y = sin(Bx)	**HSF-TF.B.5**
FORWARD 1	Combine horizontal stretches with shifts	**HSF-BF.B.3**
FORWARD 2	Graph exponential and trigonometric functions with horizontal cha…	**HSF-IF.C.7.e**

SET 56 • HSF-BF.B.3

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

f(x) = √x passes through (4, 2). Describe the graph of g(x) = √(4x) compared with f, and give the point of g with y = 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q1/5

2050

QUESTION ID

S56-M-Q1

TYPE

Short answer

CORRECT ANSWER

Horizontal compression by a factor of 1/4; (1, 2)

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q1/5

f(x) = sin x has period 2π. Describe g(x) = sin(3x) compared with f and give its period.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q2/5

2051

QUESTION ID
S56-M-Q2

TYPE
Short answer

CORRECT ANSWER
Horizontal compression by a factor of 1/3; period 2π/3

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Describe the graph of $g(x) = 2^{-x}$ compared with $f(x) = 2^{x}$, and give the point of g that comes from (1, 2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q3/5

2052

QUESTION ID
S56-M-Q3

TYPE
Short answer

CORRECT ANSWER
Reflection over the y-axis; (−1, 2)

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q3/5

f passes through (−2, 1), (0, 3), and (4, −1). List the corresponding points of g(x) = f(2x).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q4/5

2053

QUESTION ID
S56-M-Q4

TYPE
Short answer

CORRECT ANSWER
(−1, 1), (0, 3), (2, −1)

GRADING NOTE Form required: all three.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

f passes through (2, 5). Which point must be on the graph of g(x) = f(−x/2)? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q5/5

2054

QUESTION ID
S56-M-Q5

TYPE
Short answer

CORRECT ANSWER
(−4, 5):
g(−4) = f(2) = 5

GRADING NOTE Form required: point and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S56-M-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • MAIN • Q5/5

SET 56 • HSF-BF.B.3

BACKWARD 1

Reflections and dilations of points • Grade 8 • Geometry • 8.G.A.3

ANSWER KEY Reduced copy of the question page, with its answer

Reflect (1, 2) over the y-axis.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q1

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q1/5

2056

QUESTION ID

S56-B1-Q1

TYPE

Short answer

CORRECT ANSWER

(−1, 2)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q1

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Multiply the x-coordinate of (4, −1) by 1/2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q2

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q2/5

2057

QUESTION ID

S56-B1-Q2

TYPE

Short answer

CORRECT ANSWER

(2, −1)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q2

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q2/5

True or false?

Reflecting over the y-axis changes the sign of the x-coordinate.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q3

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q3/5

2058

QUESTION ID

S56-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE (x, y) → (−x, y).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q3

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

(x, y) → (x/3, y) applied to (6, 2) gives

A. (18, 2)

B. (6, 2/3)

C. (2, 2)

D. (3, 2)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q4

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q4/5

2059

QUESTION ID

S56-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. (2, 2)

GRADING NOTE Divide x by 3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q4

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Apply (x, y) → (−2x, y) to (2, 5).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q5

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q5/5

2060

QUESTION ID

S56-B1-Q5

TYPE

Short answer

CORRECT ANSWER

(−4, 5)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B1-Q5

Grade 8 • Geometry • 8.G.A.3 • Set 56 • BACKWARD 1 • Q5/5

SET 56 • HSF-BF.B.3

BACKWARD 2

Solve kx = c for x • Grade 6 • Expressions & Equations • 6.EE.B.7

ANSWER KEY Reduced copy of the question page, with its answer

Solve 2x = 4.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q1/5

2062

QUESTION ID

S56-B2-Q1

TYPE

Short answer

CORRECT ANSWER

2

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q1

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $-x/2 = 2$.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q2/5

2063

QUESTION ID

S56-B2-Q2

TYPE

Short answer

CORRECT ANSWER

−4

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q2

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

If 4x = 4, then x = 16.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q3/5

2064

QUESTION ID

S56-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Divide by 4: x = 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q3

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$3x = 2\pi$ gives $x =$

A. $2\pi/3$

B. $2\pi - 3$

C. $\pi/3$

D. 6π

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q4/5

2065

QUESTION ID

S56-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 2π/3

GRADING NOTE Divide by 3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q4

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $-x = 1$.

$x =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q5/5

2066

QUESTION ID

S56-B2-Q5

TYPE

Short answer

CORRECT ANSWER

−1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B2-Q5

Grade 6 • Expressions & Equations • 6.EE.B.7 • Set 56 • BACKWARD 2 • Q5/5

SET 56 • HSF-BF.B.3

BACKWARD 3

Period of y = sin(Bx) • High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

Give the period of y = sin(2x).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q1/5

2068

QUESTION ID

S56-B3-Q1

TYPE

Short answer

CORRECT ANSWER

π

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q1

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the period of y = cos(x/2).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q2/5

2069

QUESTION ID

S56-B3-Q2

TYPE

Short answer

CORRECT ANSWER

4π

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q2

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

For y = sin(Bx) with B > 0, a larger B gives a longer period.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q3/5

2070

QUESTION ID

S56-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The period 2π/B gets shorter as B grows.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q3

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q3/5

2070

ANSWER KEY Reduced copy of the question page, with its answer

y = sin(4x) has period

A. 8π

B. 4π

C. π/4

D. π/2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q4/5

2071

QUESTION ID

S56-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. π/2

GRADING NOTE 2π/4.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q4

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Which B gives period 6?

B = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q5/5

2072

QUESTION ID

S56-B3-Q5

TYPE

Short answer

CORRECT ANSWER

π/3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-B3-Q5

High School — Algebra II • Functions: Trigonometric Functions • HSF-TF.B.5 • Set 56 • BACKWARD 3 • Q5/5

SET 56 • HSF-BF.B.3

FORWARD 1

Combine horizontal stretches with shifts • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

Rewrite g(x) = f(2x − 6) in the form f(k(x − h)) and describe the transformation.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q1/5

2074

QUESTION ID

S56-F1-Q1

TYPE

Short answer

CORRECT ANSWER

f(2(x − 3)): compress horizontally by 1/2, then shift right 3

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q1/5

f passes through (4, 1). Which point is on $g(x) = f(2x - 6)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q2/5

2075

QUESTION ID

S56-F1-Q2

TYPE

Short answer

CORRECT ANSWER

(5, 1)

GRADING NOTE Any equivalent answer. $2x - 6 = 4$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q2/5

True or false?

f(2x − 6) is f(2x) shifted right 6.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q3/5

2076

QUESTION ID

S56-F1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE It is shifted right 3.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

y = √(9x) equals

A. √x + 9

B. 9√x

C. 3√x

D. √x/3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q4/5

2077

QUESTION ID
S56-F1-Q4

TYPE
Multiple choice

CORRECT ANSWER
C. 3√x

GRADING NOTE Horizontal compression equals vertical stretch for √.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write g, the graph of $f(x) = 2^x$ reflected over the y-axis and then shifted right 1.

g(x) = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q5/5

2078

QUESTION ID

S56-F1-Q5

TYPE

Short answer

CORRECT ANSWER

$2^{-(x-1)} = 2^{1-x}$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S56-F1-Q5**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 56 • FORWARD 1 • Q5/5

SET 56 • HSF-BF.B.3

FORWARD 2

Graph exponential and trigonometric functions with horizontal changes • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e

ANSWER KEY Reduced copy of the question page, with its answer

Give the y-intercept, asymptote, and two points of $y = 2^{-x} + 1$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q1/5

2080

QUESTION ID
S56-F2-Q1

TYPE
Short answer

CORRECT ANSWER
y-intercept 2; asymptote y = 1; points (−1, 3), (1, 1.5)

GRADING NOTE Form required: all.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the maxima of $y = \cos(2x)$ on $0 \le x \le 2\pi$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q2/5

2081

QUESTION ID

S56-F2-Q2

TYPE

Short answer

CORRECT ANSWER

x = 0, π, 2π (value 1)

GRADING NOTE Form required: all three.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$y = 3^{2x} = 9^{x}$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q3/5

2082

QUESTION ID
S56-F2-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Power of a power.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

y = sin(x/2) completes one cycle on

A. $0 \le x \le \pi$

B. $0 \le x \le 2\pi$

C. $0 \le x \le \pi/2$

D. $0 \le x \le 4\pi$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q4/5

2083

QUESTION ID
S56-F2-Q4

TYPE
Multiple choice

CORRECT ANSWER
D. $0 \le x \le 4\pi$

GRADING NOTE Period 4π.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why $y = 2^{2x}$ is a horizontal compression of $y = 2^{x}$ and also an exponential with base 4.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q5/5

2084

QUESTION ID
S56-F2-Q5

TYPE
Short answer

CORRECT ANSWER
Replacing x by 2x compresses horizontally; $2^{2x} = (2^2)^x = 4^x$.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S56-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.7.e • Set 56 • FORWARD 2 • Q5/5

Recognize even and odd functions

MAIN	Algebra II main question	**HSF-BF.B.3**
BACKWARD 1	Signs of powers of negatives	**7.NS.A.2**
BACKWARD 2	Reflect points over the axes and the origin	**6.NS.C.6.b**
BACKWARD 3	Substitute -x into an expression	**HSF-IF.A.2**
FORWARD 1	Reason about sums and products of even and odd functions	**HSF-BF.B.3**
FORWARD 2	Explain trigonometric symmetry with the unit circle	**HSF-TF.A.4**

SET 57 • HSF-BF.B.3

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

Is $f(x) = x^4 - 3x^2$ even, odd, or neither? Show $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q1/5

2087

QUESTION ID

S57-M-Q1

TYPE

Short answer

CORRECT ANSWER

Even: $f(-x) = (-x)^4 - 3(-x)^2 = x^4 - 3x^2 = f(x)$

GRADING NOTE Form required: decision and the computation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S57-M-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Is $f(x) = x^3 - x$ even, odd, or neither? Show $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q2/5

2088

QUESTION ID

S57-M-Q2

TYPE

Short answer

CORRECT ANSWER

Odd: $f(-x) = -x^3 + x = -f(x)$

GRADING NOTE Form required: decision and the computation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Is $f(x) = x^2 + x$ even, odd, or neither? Justify.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q3/5

2089

QUESTION ID
S57-M-Q3

TYPE
Short answer

CORRECT ANSWER
Neither: f(1) = 2 and f(−1) = 0, so f(−1) ≠ f(1) and f(−1) ≠ −f(1)


GRADING NOTE Form required: decision and a counterexample.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The graph of $f(x) = x/(x^2 + 1)$ is shown. Is f even, odd, or neither? Explain from the graph.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q4/5

2090

QUESTION ID
S57-M-Q4

TYPE
Short answer

CORRECT ANSWER
Odd: the graph is symmetric about the origin

GRADING NOTE Form required: decision and the symmetry.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q4/5

2090

ANSWER KEY Reduced copy of the question page, with its answer

Is $f(x) = \cos x + x^2$ even, odd, or neither? Show $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q5/5

2091

QUESTION ID

S57-M-Q5

TYPE

Short answer

CORRECT ANSWER

Even:

$f(-x) = \cos(-x) + (-x)^2 = \cos x + x^2$

GRADING NOTE Form required: decision and the computation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • MAIN • Q5/5

SET 57 • HSF-BF.B.3

BACKWARD 1

Signs of powers of negatives • Grade 7 • The Number System • 7.NS.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Simplify $(-x)^4$ and $(-x)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q1/5

2093

QUESTION ID

S57-B1-Q1

TYPE

Short answer

CORRECT ANSWER

x^4 and $-x^3$

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q1

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Simplify $(-x)^2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q2/5

2094

QUESTION ID

S57-B1-Q2

TYPE

Short answer

CORRECT ANSWER

x^2

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q2

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$(-x)^5 = x^5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q3/5

2095

QUESTION ID

S57-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE An odd power keeps the sign: $(-x)^5 = -x^5$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q3

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q3/5

2095

ANSWER KEY Reduced copy of the question page, with its answer

$(-2)^2 - (-2)$ equals

A. -6

B. 2

C. 6

D. 0

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q4/5

2096

GRADING NOTE 4 + 2.

QUESTION ID

S57-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. 6

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q4

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Simplify $-(-x)^3$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q5/5

2097

QUESTION ID

S57-B1-Q5

TYPE

Short answer

CORRECT ANSWER

x^3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B1-Q5

Grade 7 • The Number System • 7.NS.A.2 • Set 57 • BACKWARD 1 • Q5/5

SET 57 • HSF-BF.B.3

BACKWARD 2

Reflect points over the axes and the origin • Grade 6 • The Number System • 6.NS.C.6.b

ANSWER KEY Reduced copy of the question page, with its answer

Reflect (2, 3) over the y-axis.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q1

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q1/5

2099

QUESTION ID

S57-B2-Q1

TYPE

Short answer

CORRECT ANSWER

(−2, 3)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q1

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Rotate (2, 3) 180° about the origin.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q2

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q2/5

2100

QUESTION ID

S57-B2-Q2

TYPE

Short answer

CORRECT ANSWER

(−2, −3)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q2

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q2/5

True or false?

A graph symmetric about the y-axis contains (−a, b) whenever it contains (a, b).

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q3

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q3/5

2101

QUESTION ID

S57-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Even symmetry.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q3

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Symmetry about the origin pairs (a, b) with

A. (−a, b)

B. (−a, −b)

C. (a, −b)

D. (b, a)

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q4

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q4/5

2102

QUESTION ID

S57-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. (−a, −b)

GRADING NOTE 180° rotation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q4

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A graph is symmetric about the origin and contains (1, 0.5). Name another point.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q5

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q5/5

2103

QUESTION ID

S57-B2-Q5

TYPE

Short answer

CORRECT ANSWER

(−1, −0.5)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B2-Q5

Grade 6 • The Number System • 6.NS.C.6.b • Set 57 • BACKWARD 2 • Q5/5

SET 57 • HSF-BF.B.3

BACKWARD 3

Substitute −x into an expression • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

ANSWER KEY Reduced copy of the question page, with its answer

For $f(x) = x^2 + x$, find $f(-x)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q1/5

2105

QUESTION ID

S57-B3-Q1

TYPE

Short answer

CORRECT ANSWER

$x^2 - x$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

For $f(x) = x^3 - x$, find $f(-2)$ and $f(2)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q2/5

2106

QUESTION ID

S57-B3-Q2

TYPE

Short answer

CORRECT ANSWER

−6 and 6

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

For f(x) = 3x, f(−x) = 3x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q3/5

2107

QUESTION ID

S57-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE f(−x) = 3(−x) = −3x.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

For $f(x) = x^4 - 3x^2$, $f(-1)$ is

A. 2

B. 4

C. -2

D. -4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q4/5

2108

GRADING NOTE 1 − 3.

QUESTION ID

S57-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. −2

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

For f(x) = 1/x, find f(−x).

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q5/5

2109

QUESTION ID

S57-B3-Q5

TYPE

Short answer

CORRECT ANSWER

−1/x

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 57 • BACKWARD 3 • Q5/5

SET 57 • HSF-BF.B.3

FORWARD 1

Reason about sums and products of even and odd functions • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3

ANSWER KEY Reduced copy of the question page, with its answer

Prove that the sum of two odd functions is odd.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q1/5

2111

QUESTION ID

S57-F1-Q1

TYPE

Answer with required work

CORRECT ANSWER

$(f + g)(-x) = f(-x) + g(-x) = -f(x) - g(x) = -(f + g)(x).$

GRADING NOTE REQUIRED WORK: Use $f(-x) = -f(x)$ and $g(-x) = -g(x)$.

Any valid proof or derivation. Essential: both definitions and the factoring of -1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Prove that the product of two odd functions is even.

Work space

Conclusion: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q2/5

2112

QUESTION ID

S57-F1-Q2

TYPE

Answer with required work

CORRECT ANSWER

$(fg)(-x) = f(-x)g(-x)$
$= (-f(x))(-g(x))$
$= f(x)g(x).$

GRADING NOTE REQUIRED WORK: Compute $(fg)(-x)$.

Any valid proof or derivation. Essential: the computation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q2/5

True or false?

The only function that is both even and odd is f(x) = 0.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q3/5

2113

QUESTION ID

S57-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE f(x) = f(−x) = −f(x) gives f(x) = 0.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = |x| + x^3$ is

A. even

B. both

C. neither

D. odd

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q4/5

2114

QUESTION ID

S57-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. neither

GRADING NOTE $f(1) = 2$, $f(-1) = 0$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q4/5

Is f(x) = x sin x even, odd, or neither? Explain using the fact that a product of two odd functions is even.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q5/5

2115

QUESTION ID

S57-F1-Q5

TYPE

Short answer

CORRECT ANSWER

Even: x and sin x are both odd, so their product is even.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.3 • Set 57 • FORWARD 1 • Q5/5

SET 57 • HSF-BF.B.3

FORWARD 2

Explain trigonometric symmetry with the unit circle • High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Use the unit circle to explain why cos(−x) = cos x.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q1/5

2117

QUESTION ID

S57-F2-Q1

TYPE

Short answer

CORRECT ANSWER

The points at angles x and −x are reflections over the x-axis and have the same x-coordinate.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Is tan even, odd, or neither? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q2/5

2118

QUESTION ID

S57-F2-Q2

TYPE

Short answer

CORRECT ANSWER

Odd: tan(−x)
= sin(−x)/cos(−x)
= −sin x/cos
x = −tan x

GRADING NOTE Form required: decision and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q2/5

True or false?

sin is an even function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q3/5

2119

QUESTION ID

S57-F2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $\sin(-x) = -\sin x$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which is an even function?

A. cos x

B. tan x

C. sin x + cos x

D. sin x

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q4/5

2120

QUESTION ID

S57-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. cos x

GRADING NOTE Reflection over the x-axis.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Use symmetry to find $\sin(-\pi/6)$ from $\sin(\pi/6) = 1/2$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S57-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q5/5

2121

QUESTION ID

S57-F2-Q5

TYPE

Short answer

CORRECT ANSWER

$-1/2$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S57-F2-Q5**

High School — Algebra II, Advanced Extension (+) • Functions: Trigonometric Functions • HSF-TF.A.4 • Set 57 • FORWARD 2 • Q5/5

Write inverses of radical, rational, cubic, exponential, and logarithmic functions

MAIN	Algebra II main question	**HSF-BF.B.4.a**
BACKWARD 1	Inverses of linear functions	**HSF-BF.B.4.a**
BACKWARD 2	Rearrange formulas with roots, powers, and fractions	**HSA-CED.A.4**
BACKWARD 3	Domain and range	**HSF-IF.B.5**
FORWARD 1	Inverses that are their own inverse, and checking by points	**HSF-BF.B.4.a**
FORWARD 2	Verify inverses by composition	**HSF-BF.B.4.b**

SET 58 • HSF-BF.B.4.a

MAIN

Algebra II main question • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of $f(x) = (x - 1)^3 + 2$.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-M-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q1/5

2124

QUESTION ID

S58-M-Q1

TYPE

Short answer

CORRECT ANSWER

$\sqrt[3]{(x - 2)} + 1$

GRADING NOTE Any equivalent form. Domain all real numbers.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S58-M-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of $f(x) = 2x/(x - 3)$, $x \neq 3$, and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-M-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q2/5

2125

QUESTION ID
S58-M-Q2

TYPE
Short answer

CORRECT ANSWER
$f^{-1}(x) = 3x/(x - 2)$, $x \neq 2$

GRADING NOTE Form required: formula and domain.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S58-M-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q2/5

Find the inverse of f(x) = √(x + 4), x ≥ −4, and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-M-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q3/5

2126

QUESTION ID

S58-M-Q3

TYPE

Short answer

CORRECT ANSWER

$f^{-1}(x) = x^2 - 4, x \geq 0$

GRADING NOTE Form required: formula and domain (the range of f).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S58-M-Q3**

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of $f(x) = 5 \cdot 2^x$ and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q4/5

2127

QUESTION ID
S58-M-Q4

TYPE
Short answer

CORRECT ANSWER
$f^{-1}(x) = \log_2(x/5)$, $x > 0$

GRADING NOTE Form required: formula and domain.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-M-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of $f(x) = \log_3(x - 1)$ and state its domain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q5/5

2128

QUESTION ID

S58-M-Q5

TYPE

Short answer

CORRECT ANSWER

$f^{-1}(x) = 3^x + 1$, all real x

GRADING NOTE Form required: formula and domain.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-M-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • MAIN • Q5/5

SET 58 • HSF-BF.B.4.a

BACKWARD 1

Inverses of linear functions • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of f(x) = 3x − 6.

$f^{-1}(x) =$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q1/5

2130

QUESTION ID

S58-B1-Q1

TYPE

Short answer

CORRECT ANSWER

(x + 6)/3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve 2x + 5 = y for x.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q2/5

2131

QUESTION ID

S58-B1-Q2

TYPE

Short answer

CORRECT ANSWER

(y − 5)/2

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q2/5

True or false?

The inverse undoes the function: $f^{-1}(f(x)) = x$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q3/5

2132

QUESTION ID

S58-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The inverse of f(x) = x/4 + 1 is

A. 4(x − 1)

B. (x − 1)/4

C. 4x − 1

D. x/4 − 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q4/5

2133

QUESTION ID

S58-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 4(x − 1)

GRADING NOTE Undo in reverse order.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

If $f(2) = 7$, what is $f^{-1}(7)$?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q5/5

2134

QUESTION ID

S58-B1-Q5

TYPE

Short answer

CORRECT ANSWER

2

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B1-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • BACKWARD 1 • Q5/5

SET 58 • HSF-BF.B.4.a

BACKWARD 2

Rearrange formulas with roots, powers, and fractions • High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Solve $y = \sqrt{(x + 4)}$ for x ($y \geq 0$).

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q1

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q1/5

2136

QUESTION ID

S58-B2-Q1

TYPE

Short answer

CORRECT ANSWER

$y^2 - 4$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q1

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $y = 2x/(x - 3)$ for x.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q2

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q2/5

2137

QUESTION ID

S58-B2-Q2

TYPE

Short answer

CORRECT ANSWER

3y/(y − 2)

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q2

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

Solving $y = (x - 1)^3 + 2$ for x gives $x = \sqrt[3]{(y - 2)} + 1$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q3

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q3/5

2138

QUESTION ID

S58-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Undo each step.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q3

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q3/5

2138

ANSWER KEY Reduced copy of the question page, with its answer

Solving $y = x^3$ for x gives

A. y/3

B. $\sqrt[3]{y}$

C. $\sqrt{y}$

D. y^3

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q4

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q4/5

2139

QUESTION ID

S58-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. $\sqrt[3]{y}$

GRADING NOTE Cube root.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q4

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve y = x/(x + 1) for x.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q5

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q5/5

2140

QUESTION ID

S58-B2-Q5

TYPE

Short answer

CORRECT ANSWER

y/(1 − y)

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B2-Q5

High School — Algebra I • Algebra: Creating Equations • HSA-CED.A.4 • Set 58 • BACKWARD 2 • Q5/5

SET 58 • HSF-BF.B.4.a

BACKWARD 3

Domain and range • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

Give the range of $f(x) = \sqrt{(x + 4)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q1/5

2142

QUESTION ID

S58-B3-Q1

TYPE

Short answer

CORRECT ANSWER

$y \geq 0$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the range of $f(x) = 5 \cdot 2^x$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q2/5

2143

QUESTION ID

S58-B3-Q2

TYPE

Short answer

CORRECT ANSWER

y > 0

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q2/5

True or false?

The range of f(x) = 2x/(x − 3) excludes y = 3.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION — ID S58-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q3/5

2144

QUESTION ID

S58-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE f(9) = 18/6 = 3; the excluded value is y = 2 (the horizontal asymptote).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION — ID S58-B3-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q3/5

The domain of $\log_3(x - 1)$ is

A. $x > 1$

B. all reals

C. $x > 0$

D. $x \geq 1$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q4/5

2145

QUESTION ID

S58-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. x > 1

GRADING NOTE $x - 1 > 0$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give the range of $f(x) = \log_3(x - 1)$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q5/5

2146

QUESTION ID

S58-B3-Q5

TYPE

Short answer

CORRECT ANSWER

All real numbers

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-B3-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.B.5 • Set 58 • BACKWARD 3 • Q5/5

SET 58 • HSF-BF.B.4.a

FORWARD 1

Inverses that are their own inverse, and checking by points • High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a

ANSWER KEY Reduced copy of the question page, with its answer

Find the inverse of f(x) = (x + 1)/(x − 1). What do you notice?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q1/5

2148

QUESTION ID

S58-F1-Q1

TYPE

Short answer

CORRECT ANSWER

$f^{-1}(x) = (x + 1)/(x - 1)$: f is its own inverse

GRADING NOTE Form required: formula and observation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = 5 \cdot 2^x$ passes through (1, 10), and
$f^{-1}(x) = \log_2(x/5)$. Check the inverse with this point.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q2/5

2149

QUESTION ID
S58-F1-Q2

TYPE
Short answer

CORRECT ANSWER
$\log_2(10/5) = \log_2 2 = 1$

GRADING NOTE Form required: the check.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q2/5

True or false?

Every function has an inverse function.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q3/5

2150

QUESTION ID
S58-F1-Q3

TYPE
True or false

CORRECT ANSWER
B. False

GRADING NOTE Only one-to-one functions; $f(x) = x^2$ on all reals does not.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which is its own inverse?

A. $f(x) = x^2$

B. $f(x) = x + 1$

C. $f(x) = 2x$

D. $f(x) = 1/x$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q4/5

2151

QUESTION ID

S58-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. f(x) = 1/x

GRADING NOTE $1/(1/x) = x$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Use the horizontal line test on the graph of $f(x) = x^3 - x$ to decide whether f has an inverse function. Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q5/5

2152

QUESTION ID

S58-F1-Q5

TYPE

Short answer

CORRECT ANSWER

A horizontal line such as y = 0 meets the graph three times, so f is not one-to-one and has no inverse function.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F1-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Building Functions • HSF-BF.B.4.a • Set 58 • FORWARD 1 • Q5/5

SET 58 • HSF-BF.B.4.a

FORWARD 2

Verify inverses by composition • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b

Verify that $f(x) = (x - 1)^3 + 2$ and $g(x) = \sqrt[3]{(x - 2)} + 1$ are inverses by computing $f(g(x))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q1/5

2154

QUESTION ID

S58-F2-Q1

TYPE

Short answer

CORRECT ANSWER

$f(g(x)) = (\sqrt[3]{(x - 2)})^3 + 2 = x$

GRADING NOTE Form required: the computation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S58-F2-Q1**

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

$f(x) = (x - 1)^3 + 2$ and $g(x) = \sqrt[3]{(x - 2)} + 1$. Compute $g(f(x))$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q2/5

2155

QUESTION ID

S58-F2-Q2

TYPE

Short answer

CORRECT ANSWER

$\sqrt[3]{((x - 1)^3)} + 1 = x$

GRADING NOTE Form required: the computation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q2/5

True or false?

To verify inverses, both f(g(x)) = x and g(f(x)) = x must hold on the domains.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q3/5

2156

QUESTION ID

S58-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Two-sided.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

For $f(x) = \sqrt{(x + 4)}$ and $g(x) = x^2 - 4$ $(x \geq 0)$, $g(f(x)) =$

A. x for all x

B. $|x|$

C. x for $x \geq -4$

D. $x^2 + 4$

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q4/5

2157

QUESTION ID

S58-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. x for x ≥ −4

GRADING NOTE $(\sqrt{(x + 4)})^2 - 4$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why $g(x) = x^2 - 4$ without a domain restriction is not the inverse of $f(x) = \sqrt{(x + 4)}$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q5/5

2158

QUESTION ID

S58-F2-Q5

TYPE

Short answer

CORRECT ANSWER

$f(g(x)) = \sqrt{(x^2)} = |x|$, which is not x when $x < 0$.

GRADING NOTE Any equivalent explanation (uses $\sqrt{(x^2)} = |x|$).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S58-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.4.b • Set 58 • FORWARD 2 • Q5/5

Write exact logarithmic solutions of exponential equations

MAIN	Algebra II main question	**HSF-LE.A.4**
BACKWARD 1	Isolate the exponential expression	**HSA-REI.B.3**
BACKWARD 2	Powers of 2, 3, and 10	**8.EE.A.1**
BACKWARD 3	Logarithms as the inverse of exponentials	**HSF-BF.B.4.a**
FORWARD 1	Solve equations with different bases or variables on both sides	**HSF-LE.A.4**
FORWARD 2	Use the inverse relationship to solve logarithmic equations	**HSF-BF.B.5**

SET 59 • HSF-LE.A.4

MAIN

Algebra II main question • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Solve $3 \cdot 2^{5t} = 48$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q1/5

2161

QUESTION ID

S59-M-Q1

TYPE

Short answer

CORRECT ANSWER

4/5

GRADING NOTE Any equivalent form. $2^{5t} = 16 = 2^4$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $2 \cdot 10^{3t} = 9$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q2/5

2162

QUESTION ID

S59-M-Q2

TYPE

Short answer

CORRECT ANSWER

log(4.5)/3

GRADING NOTE Form required: exact (≈ 0.2177).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-M-Q2**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $5e^{2t} = 30$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q3/5

2163

QUESTION ID

S59-M-Q3

TYPE

Short answer

CORRECT ANSWER

(ln 6)/2

GRADING NOTE Form required: exact (≈ 0.8959).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $100(1.05)^t = 300$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q4/5

2164

QUESTION ID

S59-M-Q4

TYPE

Short answer

CORRECT ANSWER

log 3/log 1.05

GRADING NOTE Form required: exact (any base; ≈ 22.52).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $4 \cdot 3^{t/2} = 20$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-M-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q5/5

2165

QUESTION ID

S59-M-Q5

TYPE

Short answer

CORRECT ANSWER

$2 \log_3 5$

GRADING NOTE Form required: exact (2 log 5/log 3 accepted; ≈ 2.93).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-M-Q5**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • MAIN • Q5/5

SET 59 • HSF-LE.A.4

BACKWARD 1

Isolate the exponential expression • High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3

ANSWER KEY Reduced copy of the question page, with its answer

Solve 3y = 48 for y.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q1/5

2167

QUESTION ID

S59-B1-Q1

TYPE

Short answer

CORRECT ANSWER

16

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q1

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve 5y = 30 for y.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q2/5

2168

QUESTION ID

S59-B1-Q2

TYPE

Short answer

CORRECT ANSWER

6

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q2

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q2/5

True or false?

From $2 \cdot 10^{3t} = 9$, we get $10^{3t} = 4.5$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q3/5

2169

QUESTION ID

S59-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Divide by 2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q3

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

From $4 \cdot 3^{t/2} = 20$, $3^{t/2}$ equals

A. 80

B. 5

C. 1/5

D. 16

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q4/5

2170

QUESTION ID

S59-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 5

GRADING NOTE Divide by 4.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q4

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve 100y = 300 for y.

y = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q5/5

2171

QUESTION ID

S59-B1-Q5

TYPE

Short answer

CORRECT ANSWER

3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B1-Q5

High School — Algebra I • Algebra: Reasoning with Equations & Inequalities • HSA-REI.B.3 • Set 59 • BACKWARD 1 • Q5/5

SET 59 • HSF-LE.A.4

BACKWARD 2

Powers of 2, 3, and 10 • Grade 8 • Expressions & Equations • 8.EE.A.1

ANSWER KEY Reduced copy of the question page, with its answer

Write 16 as a power of 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q1/5

2173

QUESTION ID

S59-B2-Q1

TYPE

Short answer

CORRECT ANSWER

2^4

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q1

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Write 1000 as a power of 10.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q2/5

2174

QUESTION ID

S59-B2-Q2

TYPE

Short answer

CORRECT ANSWER

10^3

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q2

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q2/5

True or false?

$3^4 = 12$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q3/5

2175

QUESTION ID

S59-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $3^4 = 3 \cdot 3 \cdot 3 \cdot 3 = 81$; 12 is $3 \cdot 4$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q3

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$2^{5t} = 2^4$ gives

A. t = 4

B. 5t = 4

C. 5t = 16

D. t = 20

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q4/5

2176

QUESTION ID

S59-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 5t = 4

GRADING NOTE Equal exponents.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q4

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write 1/8 as a power of 2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q5/5

2177

QUESTION ID

S59-B2-Q5

TYPE

Short answer

CORRECT ANSWER

2^{-3}

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B2-Q5

Grade 8 • Expressions & Equations • 8.EE.A.1 • Set 59 • BACKWARD 2 • Q5/5

SET 59 • HSF-LE.A.4

BACKWARD 3

Logarithms as the inverse of exponentials • High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a

ANSWER KEY Reduced copy of the question page, with its answer

Rewrite $10^x = 4.5$ as a logarithm.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q1/5

2179

QUESTION ID

S59-B3-Q1

TYPE

Short answer

CORRECT ANSWER

$x = \log 4.5$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q1

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Rewrite $e^u = 6$ as a logarithm.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q2/5

2180

QUESTION ID

S59-B3-Q2

TYPE

Short answer

CORRECT ANSWER

u = ln 6

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q2

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$\log_2 8 = 4$ because $2^4 = 8$.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q3/5

2181

QUESTION ID

S59-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE $2^4 = 16$; $\log_2 8 = 3$ because $2^3 = 8$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q3

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$3^y = 5$ means y =

A. $\log_3 5$

B. 5/3

C. $\log_5 3$

D. 3/5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q4/5

2182

QUESTION ID
S59-B3-Q4

TYPE
Multiple choice

CORRECT ANSWER
A. $\log_3 5$

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q4

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate $\log_2 32$.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q5/5

2183

QUESTION ID

S59-B3-Q5

TYPE

Short answer

CORRECT ANSWER

5

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-B3-Q5

High School — Algebra I • Functions: Building Functions • HSF-BF.B.4.a • Set 59 • BACKWARD 3 • Q5/5

SET 59 • HSF-LE.A.4

FORWARD 1

Solve equations with different bases or variables on both sides • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Solve $2^x = 3^{x-1}$ exactly.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q1/5

2185

QUESTION ID

S59-F1-Q1

TYPE

Short answer

CORRECT ANSWER

ln 3/(ln 3 − ln 2)

GRADING NOTE Form required: exact (any base; ≈ 2.71).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $5^{2x} = 7$ exactly.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q2/5

2186

QUESTION ID

S59-F1-Q2

TYPE

Short answer

CORRECT ANSWER

log 7/(2 log 5)

GRADING NOTE Form required: exact (≈ 0.605).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

log(ab) = log a + log b for positive a and b.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q3/5

2187

QUESTION ID

S59-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Product property.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The exact solution of $e^{x+1} = 10$ is

A. 10/e

B. ln 10/1

C. ln 9

D. ln 10 − 1

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q4/5

2188

QUESTION ID

S59-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. ln 10 − 1

GRADING NOTE x + 1 = ln 10.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $3 \cdot 4^x = 2 \cdot 5^x$ exactly.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q5/5

2189

QUESTION ID

S59-F1-Q5

TYPE

Short answer

CORRECT ANSWER

log(3/2)/log(5/4)

GRADING NOTE Form required: exact (≈ 1.817).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 59 • FORWARD 1 • Q5/5

SET 59 • HSF-LE.A.4

FORWARD 2

Use the inverse relationship to solve logarithmic equations • High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $\log_2(x + 3) = 5$.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q1

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q1/5

2191

QUESTION ID

S59-F2-Q1

TYPE

Short answer

CORRECT ANSWER

29

GRADING NOTE Any equivalent form. x + 3 = 32.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-F2-Q1**

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve log x + log(x − 3) = 1 and check for extraneous solutions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q2

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q2/5

2192

QUESTION ID

S59-F2-Q2

TYPE

Short answer

CORRECT ANSWER

x = 5; x = −2 is extraneous

GRADING NOTE Form required: solution and rejection. x(x − 3) = 10.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S59-F2-Q2**

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q2/5

True or false?

$\log_b(b^x) = x$ for every real x.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q3/5

2193

QUESTION ID

S59-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Inverse functions.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q3

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$\ln(e^7)$ equals

A. ln 7

B. 1

C. 7

D. e^7

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q4/5

2194

QUESTION ID
S59-F2-Q4

TYPE
Multiple choice

CORRECT ANSWER
C. 7

GRADING NOTE Inverse.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q4

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve ln(2x) = 3 exactly.

x = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q5/5

2195

QUESTION ID

S59-F2-Q5

TYPE

Short answer

CORRECT ANSWER

$e^3/2$

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S59-F2-Q5

High School — Algebra II, Advanced Extension (+) • Functions: Building Functions • HSF-BF.B.5 • Set 59 • FORWARD 2 • Q5/5

Evaluate logarithmic solutions of exponential models with technology

MAIN	Algebra II main question	**HSF-LE.A.4**
BACKWARD 1	Write exact logarithmic solutions	**HSF-LE.A.4**
BACKWARD 2	Evaluate logarithms with a calculator	**HSF-IF.A.2**
BACKWARD 3	Choose an appropriate level of accuracy	**HSN-Q.A.3**
FORWARD 1	Analyze how parameters change the solution	**HSF-LE.A.4**
FORWARD 2	Rewrite models to show doubling time or half-life	**HSF-IF.C.8.b**

SET 60 • HSF-LE.A.4

MAIN

Algebra II main question • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

ANSWER KEY Reduced copy of the question page, with its answer

$2,500 earns 3.5% interest compounded yearly. Solve $2500(1.035)^t = 4000$ and interpret the answer. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q1/5

2198

QUESTION ID

S60-M-Q1

TYPE

Short answer

CORRECT ANSWER

t = log 1.6/log 1.035 ≈ 13.66; the balance first reaches $4,000 after 14 yearly compoundings

GRADING NOTE Form required: value and interpretation (13.66 years; 14 whole years).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Carbon-14 has a half-life of 5730 years. Solve $80(0.5)^{t/5730} = 30$ to the nearest year and interpret it. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q2/5

2199

QUESTION ID
S60-M-Q2

TYPE
Short answer

CORRECT ANSWER

t ≈ 8108 years: a sample with 30 of its original 80 units of carbon-14 is about 8,108 years old

GRADING NOTE Form required: value and interpretation (8100 to 8110 accepted).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Coffee cools by $T(t) = 20 + 70(0.95)^t$ °C after t minutes. When is it 50°C? Give minutes to the nearest hundredth. A calculator may be used.

t ≈ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q3/5

2200

QUESTION ID
S60-M-Q3

TYPE
Short answer

CORRECT ANSWER
16.52 minutes

GRADING NOTE Any equivalent form. $0.95^t = 3/7$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A city of 1.2 million grows by $P(t) = 1.2e^{0.015t}$ million people. When does it reach 1.5 million? Give years to the nearest hundredth. A calculator may be used.

$t \approx$ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q4/5

2201

QUESTION ID
S60-M-Q4

TYPE
Short answer

CORRECT ANSWER
14.88 years

GRADING NOTE Any equivalent form. ln 1.25/0.015.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S60-M-Q4**

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A dose leaves $200(0.8)^t$ mg of a drug after t hours. After which whole hour is the amount first below 50 mg? A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q5/5

2202

QUESTION ID
S60-M-Q5

TYPE
Short answer

CORRECT ANSWER
After 7 hours
(t > 6.21)

GRADING NOTE Form required: the whole hour and the evidence. $0.8^t < 0.25$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-M-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • MAIN • Q5/5

SET 60 • HSF-LE.A.4

BACKWARD 1

Write exact logarithmic solutions • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Solve $1.035^t = 1.6$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q1/5

2204

QUESTION ID

S60-B1-Q1

TYPE

Short answer

CORRECT ANSWER

log 1.6/log 1.035

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $e^{0.015t} = 1.25$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q2/5

2205

QUESTION ID

S60-B1-Q2

TYPE

Short answer

CORRECT ANSWER

ln 1.25/0.015

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q2/5

True or false?

$0.8^t = 0.25$ gives t = log 0.25/log 0.8.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q3/5

2206

QUESTION ID

S60-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Take logs.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$(0.5)^{t/5730} = 0.375$ gives t =

A. 0.375 · 5730

B. 5730/log 0.375

C. log 0.375/5730

D. 5730 log 0.375/log 0.5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q4/5

2207

QUESTION ID

S60-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 5730 log 0.375/log 0.5

GRADING NOTE Multiply by 5730.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Solve $0.95^t = 3/7$ exactly.

t = _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q5/5

2208

QUESTION ID

S60-B1-Q5

TYPE

Short answer

CORRECT ANSWER

log(3/7)/log 0.95

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • BACKWARD 1 • Q5/5

SET 60 • HSF-LE.A.4

BACKWARD 2

Evaluate logarithms with a calculator • High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate log 1.6 to four decimal places. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q1/5

2210

QUESTION ID
S60-B2-Q1

TYPE
Short answer

CORRECT ANSWER
0.2041

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q1

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate ln 1.25 to four decimal places. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q2/5

2211

QUESTION ID
S60-B2-Q2

TYPE
Short answer

CORRECT ANSWER
0.2231

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q2

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

log 0.25 is positive.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q3/5

2212

QUESTION ID

S60-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE 0.25 < 1, so log 0.25 < 0 (about −0.602).

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q3

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

log 1.6/log 1.035 is about

A. 0.20

B. 46.0

C. 1.55

D. 13.66

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q4/5

2213

QUESTION ID

S60-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 13.66

GRADING NOTE 0.2041/0.01494.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q4

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Evaluate log 0.25/log 0.8 to two decimal places. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q5/5

2214

QUESTION ID
S60-B2-Q5

TYPE
Short answer

CORRECT ANSWER
6.21

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B2-Q5

High School — Algebra I • Functions: Interpreting Functions • HSF-IF.A.2 • Set 60 • BACKWARD 2 • Q5/5

SET 60 • HSF-LE.A.4

BACKWARD 3

Choose an appropriate level of accuracy • High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3

ANSWER KEY Reduced copy of the question page, with its answer

A model gives 13.66 years for yearly interest. What whole-year answer fits the context?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q1

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q1/5

2216

QUESTION ID

S60-B3-Q1

TYPE

Short answer

CORRECT ANSWER

14 years

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q1

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Round 8108.16 years to the nearest hundred years.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q2

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q2/5

2217

QUESTION ID

S60-B3-Q2

TYPE

Short answer

CORRECT ANSWER

8100 years

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q2

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q2/5

True or false?

A time found from a model should be reported with more decimal places than the data justify.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q3

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q3/5

2218

QUESTION ID

S60-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Precision should match the context.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q3

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A drug amount first drops below 50 mg at t = 6.21 h. If it is checked every hour, it is first below 50 mg at

A. 6.21 h

B. 6 h

C. 7 h

D. 5 h

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q4

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q4/5

2219

QUESTION ID

S60-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. 7 h

GRADING NOTE Next whole hour.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q4

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Why might 14.88 years be reported as "about 15 years"?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q5

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q5/5

2220

QUESTION ID

S60-B3-Q5

TYPE

Short answer

CORRECT ANSWER

Population models are approximate, so a rounded value is more honest.

GRADING NOTE Any reasonable answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-B3-Q5

High School — Algebra I • Number & Quantity: Quantities • HSN-Q.A.3 • Set 60 • BACKWARD 3 • Q5/5

2220

SET 60 • HSF-LE.A.4

FORWARD 1

Analyze how parameters change the solution • High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4

ANSWER KEY Reduced copy of the question page, with its answer

If the interest rate changes from 3.5% to 5%, how long does it take $2,500 to reach $4,000? Give years to the nearest hundredth. A calculator may be used.

t ≈ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q1/5

2222

QUESTION ID

S60-F1-Q1

TYPE

Short answer

CORRECT ANSWER

9.63 years

GRADING NOTE Any equivalent form. log 1.6/log 1.05.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q1

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the doubling time at 6% yearly interest and compare it with the Rule of 72 estimate 72/6. A calculator may be used.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q2/5

2223

QUESTION ID
S60-F1-Q2

TYPE
Short answer

CORRECT ANSWER
log 2/log 1.06 ≈ 11.90 years; Rule of 72 gives 12 years (close)

GRADING NOTE Form required: both values.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q2

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q2/5

True or false?

Doubling the initial amount in $a(b)^t = d$ halves the time t needed.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q3/5

2224

QUESTION ID
S60-F1-Q3

TYPE
True or false

CORRECT ANSWER
B. False

GRADING NOTE It reduces t by one doubling time, not by half.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q3

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

In $200(0.8)^t = 50$, replacing 200 by 400 changes t by

A. doubling it

B. halving it

C. adding log 2/log(1/0.8) ≈ 3.11 hours

D. nothing

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q4/5

2225

QUESTION ID

S60-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. adding log 2/log(1/0.8) ≈ 3.11 hours

GRADING NOTE $400(0.8)^t = 50$ needs $0.8^t = 1/8$ instead of 1/4: t grows by log(1/2)/log 0.8 ≈ 3.11.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q4

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

At 3.5% yearly interest, explain why $2,500 takes as long to grow to $4,000 as $5,000 takes to grow to $8,000.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q5/5

2226

QUESTION ID

S60-F1-Q5

TYPE

Short answer

CORRECT ANSWER

Both need the growth factor 1.6, and t = log 1.6/log 1.035 depends only on that ratio.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F1-Q5

High School — Algebra II • Functions: Linear, Quadratic & Exponential Models • HSF-LE.A.4 • Set 60 • FORWARD 1 • Q5/5

SET 60 • HSF-LE.A.4

FORWARD 2

Rewrite models to show doubling time or half-life • High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b

ANSWER KEY Reduced copy of the question page, with its answer

Rewrite $2500(1.035)^t$ in the form $2500 \cdot 2^{t/D}$ and give D to the nearest hundredth. A calculator may be used.

D ≈ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q1

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q1/5

2228

QUESTION ID

S60-F2-Q1

TYPE

Short answer

CORRECT ANSWER

20.15 years

GRADING NOTE Any equivalent form. D = log 2/log 1.035.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S60-F2-Q1**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Rewrite $200(0.8)^t$ in the form $200(0.5)^{t/H}$ and give H to the nearest hundredth. A calculator may be used.

H ≈ _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q2

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q2/5

2229

QUESTION ID
S60-F2-Q2

TYPE
Short answer

CORRECT ANSWER
3.11 hours

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S60-F2-Q2**

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

$80(0.5)^{t/5730}$ shows the half-life 5730 directly.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q3/5

2230

QUESTION ID

S60-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Exponent t/5730.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q3

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

$1.2e^{0.015t}$ has doubling time about

A. 46.2 years

B. 66.7 years

C. 15 years

D. 0.015 years

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q4/5

2231

QUESTION ID

S60-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 46.2 years

GRADING NOTE In 2/0.015.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q4

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why the half-life form makes it easy to answer "how much remains after 3 half-lives".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q5/5

2232

QUESTION ID
S60-F2-Q5

TYPE
Short answer

CORRECT ANSWER
Each half-life multiplies by 0.5, so 3 half-lives leave $(0.5)^3 = 1/8$ of the amount.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S60-F2-Q5

High School — Algebra II, revisits an Algebra I standard • Functions: Interpreting Functions • HSF-IF.C.8.b • Set 60 • FORWARD 2 • Q5/5

SET 61 • HSS-ID.A.4 • STATISTICS, INFERENCE, AND DECISIONS

Estimate population percentages with a normal distribution

MAIN	Algebra II main question	**HSS-ID.A.4**
BACKWARD 1	Interpret mean and standard deviation	**HSS-ID.A.2**
BACKWARD 2	Describe the shape of a distribution	**HSS-ID.A.1**
BACKWARD 3	Percents of a whole	**6.RP.A.3.c**
FORWARD 1	Find values from percentiles of a normal model	**HSS-ID.A.4**
FORWARD 2	Use a normal model for a margin of error	**HSS-IC.B.4**

SET 61 • HSS-ID.A.4

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Adult women's heights are approximately normal with mean 64 in and standard deviation 2.5 in. Estimate the percent between 59 and 69 inches. Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-M-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q1/5

2235

QUESTION ID

S61-M-Q1

TYPE

Short answer

CORRECT ANSWER

95%

GRADING NOTE Any equivalent form. 59 and 69 are 2 standard deviations from the mean.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-M-Q1**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Adult women's heights are approximately normal with mean 64 in and standard deviation 2.5 in. Estimate the percent of women taller than 66.5 inches. Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-M-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q2/5

2236

QUESTION ID

S61-M-Q2

TYPE

Short answer

CORRECT ANSWER

16%

GRADING NOTE Any equivalent form. (100 − 68)/2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-M-Q2**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

Test scores are approximately normal with mean 500 and standard deviation 100. Estimate the percent of scores between 400 and 700. Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-M-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q3/5

2237

QUESTION ID

S61-M-Q3

TYPE

Short answer

CORRECT ANSWER

81.5%

GRADING NOTE Any equivalent form. 34 + 34 + 13.5.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-M-Q3**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Battery life is approximately normal with mean 40 h and standard deviation 4 h. Use the table to estimate the percent of batteries lasting less than 46 hours.

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-M-Q4

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q4/5

2238

QUESTION ID

S61-M-Q4

TYPE

Short answer

CORRECT ANSWER

93.32%

GRADING NOTE Any equivalent form. z = 1.5.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-M-Q4**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Cereal box weights are approximately normal with mean 16.0 oz and standard deviation 0.2 oz. Use the table to estimate the percent between 15.8 and 16.3 oz.

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-M-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q5/5

2239

QUESTION ID

S61-M-Q5

TYPE

Short answer

CORRECT ANSWER

77.45%

GRADING NOTE Any equivalent form. z from −1 to 1.5: 0.9332 − 0.1587.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-M-Q5**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • MAIN • Q5/5

SET 61 • HSS-ID.A.4

BACKWARD 1

Interpret mean and standard deviation • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2

ANSWER KEY Reduced copy of the question page, with its answer

A data set has mean 64 and standard deviation 2.5.
What value is one standard deviation above the mean?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B1-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q1/5

2241

QUESTION ID
S61-B1-Q1

TYPE
Short answer

CORRECT ANSWER
66.5

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-B1-Q1**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Which value is two standard deviations below a mean of 500 with standard deviation 100?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B1-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q2/5

2242

QUESTION ID

S61-B1-Q2

TYPE

Short answer

CORRECT ANSWER

300

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-B1-Q2**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q2/5

True or false?

A larger standard deviation means the data are more spread out.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B1-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q3/5

2243

QUESTION ID

S61-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Spread.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B1-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

46 is how many standard deviations above a mean of 40 when the standard deviation is 4?

A. 0.67

B. 4

C. 6

D. 1.5

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B1-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q4/5

2244

QUESTION ID

S61-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 1.5

GRADING NOTE (46 − 40)/4.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-B1-Q4**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Compute (15.8 − 16.0)/0.2.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B1-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q5/5

2245

QUESTION ID

S61-B1-Q5

TYPE

Short answer

CORRECT ANSWER

−1

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-B1-Q5**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.2 • Set 61 • BACKWARD 1 • Q5/5

SET 61 • HSS-ID.A.4

BACKWARD 2

Describe the shape of a distribution • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1

Describe a histogram that is highest in the middle and falls off evenly on both sides.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q1/5

2247

QUESTION ID

S61-B2-Q1

TYPE

Short answer

CORRECT ANSWER

Symmetric, mound-shaped (bell-shaped)

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-B2-Q1**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Where are the mean and median of a symmetric distribution?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q2/5

2248

QUESTION ID

S61-B2-Q2

TYPE

Short answer

CORRECT ANSWER

About the same, at the center

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-B2-Q2**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q2/5

True or false?

A normal curve is symmetric about its mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q3/5

2249

QUESTION ID

S61-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Bell shape.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A bell-shaped histogram is called

A. skewed right

B. uniform

C. mound-shaped and symmetric

D. bimodal

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q4/5

2250

QUESTION ID

S61-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. mound-shaped and symmetric

GRADING NOTE Shape vocabulary.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

In a symmetric distribution, what percent of the data are below the median?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q5/5

2251

QUESTION ID

S61-B2-Q5

TYPE

Short answer

CORRECT ANSWER

50%

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B2-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 61 • BACKWARD 2 • Q5/5

SET 61 • HSS-ID.A.4

BACKWARD 3

Percents of a whole • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c

ANSWER KEY Reduced copy of the question page, with its answer

Find 95% of 2,000.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q1/5

2253

QUESTION ID

S61-B3-Q1

TYPE

Short answer

CORRECT ANSWER

1900

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

What is 34% + 13.5%?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q2/5

2254

QUESTION ID

S61-B3-Q2

TYPE

Short answer

CORRECT ANSWER

47.5%

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

Half of 68% is 36%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q3/5

2255

QUESTION ID

S61-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE 68/2 = 34.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

100% − 95% split equally into two tails gives each tail

A. 5%

B. 47.5%

C. 0.5%

D. 2.5%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q4/5

2256

QUESTION ID

S61-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 2.5%

GRADING NOTE 5/2.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write 0.7745 as a percent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q5/5

2257

QUESTION ID

S61-B3-Q5

TYPE

Short answer

CORRECT ANSWER

77.45%

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 61 • BACKWARD 3 • Q5/5

SET 61 • HSS-ID.A.4

FORWARD 1

Find values from percentiles of a normal model • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

ANSWER KEY Reduced copy of the question page, with its answer

Above what height are the tallest 2.5% of women (mean 64 in, SD 2.5 in)? Use the 68-95-99.7 rule.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F1-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q1/5

2259

QUESTION ID
S61-F1-Q1

TYPE
Short answer

CORRECT ANSWER
69 inches

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-F1-Q1**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Use the table: what battery life is the 84th percentile (mean 40 h, SD 4 h)?

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F1-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q2/5

2260

QUESTION ID

S61-F1-Q2

TYPE

Short answer

CORRECT ANSWER

44 hours

GRADING NOTE Any equivalent form. z = 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F1-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q2/5

True or false?

In a normal model, the median equals the mean.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F1-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q3/5

2261

QUESTION ID

S61-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Symmetry.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F1-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

The 16th percentile of scores with mean 500 and SD 100 is

A. 484

B. 600

C. 340

D. 400

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F1-Q4

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q4/5

2262

QUESTION ID

S61-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. 400

GRADING NOTE One SD below.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-F1-Q4**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A cereal box below 15.7 oz is rejected. Use the table to estimate the percent rejected (mean 16.0, SD 0.2).

z	−1.5	−1	−0.5	0	0.5	1	1.5	2
Area to the left	0.0668	0.1587	0.3085	0.5000	0.6915	0.8413	0.9332	0.9772

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F1-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q5/5

2263

QUESTION ID

S61-F1-Q5

TYPE

Short answer

CORRECT ANSWER

6.68%

GRADING NOTE Any equivalent form. z = −1.5.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-F1-Q5**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 61 • FORWARD 1 • Q5/5

SET 61 • HSS-ID.A.4

FORWARD 2

Use a normal model for a margin of error • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

ANSWER KEY Reduced copy of the question page, with its answer

Simulated sample proportions are approximately normal with mean 0.58 and SD 0.049. Within what interval do about 95% of them fall?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q1/5

2265

QUESTION ID

S61-F2-Q1

TYPE

Short answer

CORRECT ANSWER

0.482 to 0.678

GRADING NOTE Any equivalent form. ±2 SD.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Simulated sample proportions have standard deviation 0.049. What margin of error does this give for the population proportion?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q2/5

2266

QUESTION ID

S61-F2-Q2

TYPE

Short answer

CORRECT ANSWER

About 0.098 (about 10 percentage points)

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q2/5

True or false?

The margin of error is about two standard deviations of the simulated statistic.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q3/5

2267

QUESTION ID

S61-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE 95% range.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q3/5

If the SD of simulated means is 0.2 hours, the margin of error is about

A. 0.1 hours

B. 0.4 hours

C. 2 hours

D. 0.2 hours

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q4/5

2268

QUESTION ID

S61-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 0.4 hours

GRADING NOTE $2 \cdot 0.2$.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S61-F2-Q4**

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q4/5

Explain why the normal model is useful for simulated sample proportions.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q5/5

2269

GRADING NOTE Any equivalent explanation.

QUESTION ID

S61-F2-Q5

TYPE

Short answer

CORRECT ANSWER

Their distribution is approximately mound-shaped and symmetric, so the 95% range is about the mean ± 2 SD.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S61-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 61 • FORWARD 2 • Q5/5

SET 62 • HSS-ID.A.4 • STATISTICS, INFERENCE, AND DECISIONS

Recognize data sets for which a normal model is not appropriate

MAIN	Algebra II main question	**HSS-ID.A.4**
BACKWARD 1	Describe skewed, symmetric, bimodal, and uniform shapes	**HSS-ID.A.3**
BACKWARD 2	Measures of center and spread for a data set	**6.SP.A.2**
BACKWARD 3	Groups combined in one data set	**HSS-ID.A.1**
FORWARD 1	Check a normal model with the 68-95-99.7 rule	**HSS-ID.A.4**
FORWARD 2	Decide whether a probability model fits observed data	**HSS-IC.A.2**


SET 62 • HSS-ID.A.4

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

ANSWER KEY Reduced copy of the question page, with its answer

The histogram shows household incomes in a town. Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q1/5

2272

QUESTION ID

S62-M-Q1

TYPE

Short answer

CORRECT ANSWER

No: the distribution is strongly skewed right, not symmetric and mound-shaped


GRADING NOTE Form required: decision and the shape.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

The histogram shows heights of adults in a club. Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q2/5

2273

QUESTION ID

S62-M-Q2

TYPE

Short answer

CORRECT ANSWER

Yes: roughly symmetric and mound-shaped with a single peak


GRADING NOTE Form required: decision and the shape.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

The histogram shows heights of all people at a school event (children and adults). Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q3/5

2274

QUESTION ID
S62-M-Q3

TYPE
Short answer

CORRECT ANSWER
No: the distribution has two peaks (bimodal), from two different groups

GRADING NOTE Form required: decision and the shape.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Waiting times at a clinic have mean 4 minutes, standard deviation 5 minutes, and minimum 0. Is a normal model appropriate? Justify using the numbers.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q4

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q4/5

2275

QUESTION ID

S62-M-Q4

TYPE

Short answer

CORRECT ANSWER

No: a normal model would put about 16% of times below 4 − 5 = −1 minute, which is impossible, so the data must be skewed right


GRADING NOTE Form required: decision and the numerical reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-M-Q4**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

The histogram shows the results of 600 rolls of a fair die. Is a normal model appropriate? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q5/5

2276

QUESTION ID
S62-M-Q5

TYPE
Short answer

CORRECT ANSWER
No: the distribution is roughly uniform (flat), not mound-shaped

GRADING NOTE Form required: decision and the shape.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-M-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • MAIN • Q5/5

SET 62 • HSS-ID.A.4

BACKWARD 1

Describe skewed, symmetric, bimodal, and uniform shapes • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3

ANSWER KEY Reduced copy of the question page, with its answer

A histogram has a long tail to the right. Name its shape.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q1/5

2278

QUESTION ID

S62-B1-Q1

TYPE

Short answer

CORRECT ANSWER

Skewed right

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A histogram has two separate peaks. Name its shape.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q2/5

2279

QUESTION ID

S62-B1-Q2

TYPE

Short answer

CORRECT ANSWER

Bimodal

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q2/5

True or false?

In a right-skewed distribution, the mean is usually less than the median.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q3/5

2280

QUESTION ID

S62-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The long right tail pulls the mean above the median.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A histogram with bars all about the same height is

A. skewed left

B. uniform

C. bimodal

D. normal

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q4/5

2281

QUESTION ID
S62-B1-Q4

TYPE
Multiple choice

CORRECT ANSWER
B. uniform

GRADING NOTE Flat.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Which is more resistant to a long tail: the mean or the median?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q5/5

2282

QUESTION ID

S62-B1-Q5

TYPE

Short answer

CORRECT ANSWER

The median

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B1-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.3 • Set 62 • BACKWARD 1 • Q5/5

SET 62 • HSS-ID.A.4

BACKWARD 2

Measures of center and spread for a data set • Grade 6 • Statistics & Probability • 6.SP.A.2

ANSWER KEY Reduced copy of the question page, with its answer

What does the standard deviation (or another spread measure) describe?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q1

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q1/5

2284

QUESTION ID

S62-B2-Q1

TYPE

Short answer

CORRECT ANSWER

How spread out the data are around the center

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q1

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Data are all at least 0 with mean 4. Can many values be far below 4?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q2

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q2/5

2285

QUESTION ID

S62-B2-Q2

TYPE

Short answer

CORRECT ANSWER

No: values cannot go below 0

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q2

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

A distribution can be described by its center, spread, and shape.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q3

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q3/5

2286

QUESTION ID

S62-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Three features.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q3

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which is a measure of center?

A. median

B. range

C. standard deviation

D. IQR

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q4

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q4/5

2287

QUESTION ID

S62-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. median

GRADING NOTE Center.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q4

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Give one value two standard deviations below a mean of 4 with SD 5.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q5

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q5/5

2288

QUESTION ID

S62-B2-Q5

TYPE

Short answer

CORRECT ANSWER

−6

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B2-Q5

Grade 6 • Statistics & Probability • 6.SP.A.2 • Set 62 • BACKWARD 2 • Q5/5

SET 62 • HSS-ID.A.4

BACKWARD 3

Groups combined in one data set • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1

ANSWER KEY Reduced copy of the question page, with its answer

Two groups with different centers are combined. What might the histogram show?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B3-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q1/5

2290

QUESTION ID

S62-B3-Q1

TYPE

Short answer

CORRECT ANSWER

Two peaks

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-B3-Q1**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Name a variable whose histogram for all students in a K-12 school might have several peaks.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B3-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q2/5

2291

QUESTION ID

S62-B3-Q2

TYPE

Short answer

CORRECT ANSWER

Height (for example)

GRADING NOTE Any reasonable answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-B3-Q2**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q2/5

True or false?

A histogram of one group can look different from a histogram of two groups combined.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B3-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q3/5

2292

QUESTION ID
S62-B3-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Mixtures.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B3-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A bimodal histogram suggests

A. an error in counting

B. a small standard deviation

C. two groups in the data

D. a normal model

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B3-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q4/5

2293

QUESTION ID

S62-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. two groups in the data

GRADING NOTE Two populations.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B3-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q4/5

How could you get a mound-shaped distribution from the school-event heights?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-B3-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q5/5

2294

QUESTION ID
S62-B3-Q5

TYPE
Short answer

CORRECT ANSWER
Separate the children and the adults into two data sets

GRADING NOTE Any reasonable answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-B3-Q5**

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.1 • Set 62 • BACKWARD 3 • Q5/5

SET 62 • HSS-ID.A.4

FORWARD 1

Check a normal model with the 68-95-99.7 rule • High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4

ANSWER KEY Reduced copy of the question page, with its answer

In 1,000 measurements, 70% are within 1 SD of the mean and 96% within 2 SDs. Is a normal model reasonable? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F1-Q1

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q1/5

2296

QUESTION ID

S62-F1-Q1

TYPE

Short answer

CORRECT ANSWER

Yes: the percents are close to 68% and 95%

GRADING NOTE Form required: decision and comparison.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F1-Q1**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q1/5

In another data set, 85% of values are within 1 SD and 91% within 2 SDs. Is a normal model reasonable?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F1-Q2

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q2/5

2297

QUESTION ID

S62-F1-Q2

TYPE

Short answer

CORRECT ANSWER

No: 85% is far above 68% and 91% is below 95%

GRADING NOTE Form required: decision and comparison.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F1-Q2**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q2/5

True or false?

If a data set has an extreme outlier, the normal model may still fit the rest but the mean and SD are distorted.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F1-Q3

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q3/5

2298

QUESTION ID

S62-F1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Outliers affect mean and SD.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F1-Q3**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

About what percent of normal data lies more than 3 SDs from the mean?

A. 5%

B. 3%

C. 0.3%

D. 32%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F1-Q4

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q4/5

2299

QUESTION ID

S62-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. 0.3%

GRADING NOTE 100 − 99.7.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F1-Q4**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A data set of 200 values has 25 values more than 2 SDs from the mean. Compare with a normal model.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F1-Q5

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q5/5

2300

QUESTION ID

S62-F1-Q5

TYPE

Short answer

CORRECT ANSWER

12.5% vs about 5%: too many extreme values for a normal model

GRADING NOTE Form required: both percents and the conclusion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S62-F1-Q5**

High School — Algebra II • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.A.4 • Set 62 • FORWARD 1 • Q5/5

SET 62 • HSS-ID.A.4

FORWARD 2

Decide whether a probability model fits observed data • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2

ANSWER KEY Reduced copy of the question page, with its answer

A spinner is claimed to land on each of 4 colors equally often. In 400 spins the counts are 98, 105, 97, 100. Are the results consistent with the claim?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q1/5

2302

QUESTION ID

S62-F2-Q1

TYPE

Short answer

CORRECT ANSWER

Yes: each count is close to the expected 100

GRADING NOTE Form required: decision and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A model says a coin is fair. In 50 flips there were 34 heads. In 200 simulated runs of 50 fair flips, 3 had 34 or more heads. Is the model consistent with the data?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q2/5

2303

QUESTION ID

S62-F2-Q2

TYPE

Short answer

CORRECT ANSWER

No: 3/200 = 1.5% is very unusual, so the data are evidence against a fair coin

GRADING NOTE Form required: decision and the proportion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q2/5

True or false?

A model is rejected only if the observed result would be unusual under the model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q3/5

2304

QUESTION ID

S62-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Logic of the test.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A simulated result at least as extreme as the observed one occurs in 34 of 200 runs. The model is

A. proved true

B. rejected

C. consistent with the data

D. impossible

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q4/5

2305

QUESTION ID

S62-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. consistent with the data

GRADING NOTE 17% is not unusual.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain the difference between judging a normal model for a data set and judging a probability model with a simulation.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q5/5

2306

QUESTION ID

S62-F2-Q5

TYPE

Short answer

CORRECT ANSWER

The first compares the shape of data to a bell curve; the second asks whether an observed result is unusual under the model's simulated results.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S62-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 62 • FORWARD 2 • Q5/5

SET 63 • HSS-IC.A.1 • STATISTICS, INFERENCE, AND DECISIONS

Distinguish population, sample, parameter, and statistic

MAIN	Algebra II main question	**HSS-IC.A.1**
BACKWARD 1	Samples and populations	**7.SP.A.1**
BACKWARD 2	Draw inferences from random samples	**7.SP.A.2**
BACKWARD 3	Compute sample proportions and means	**6.SP.B.5.c**
FORWARD 1	Describe sampling variability	**HSS-IC.A.1**
FORWARD 2	Quantify variability with a margin of error	**HSS-IC.B.4**

SET 63 • HSS-IC.A.1

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

ANSWER KEY Reduced copy of the question page, with its answer

A city surveys 400 randomly chosen residents, and 62% support a new park. Identify the population, the sample, the parameter, and the statistic.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q1/5

2309

QUESTION ID

S63-M-Q1

TYPE

Short answer

CORRECT ANSWER

Population: all city residents; sample: the 400 surveyed; parameter: the percent of all residents who support the park; statistic: 62%

GRADING NOTE Form required: all four.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A factory measures 50 randomly selected bolts from a day's production; their mean length is 5.02 cm. Identify the parameter and the statistic.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q2/5

2310

QUESTION ID
S63-M-Q2

TYPE
Short answer

CORRECT ANSWER
Parameter: the mean length of all bolts made that day; statistic: 5.02 cm, the sample mean

GRADING NOTE Form required: both.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

A news website posts a poll; 80% of the 2,000 visitors who answered want a later school start. Can this be generalized to all students? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q3/5

2311

QUESTION ID
S63-M-Q3

TYPE
Short answer

CORRECT ANSWER
No: the sample is voluntary, not random, so people with strong opinions are over-represented

GRADING NOTE Form required: decision and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q3/5

Explain why a random sample allows conclusions about the whole population.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q4/5

2312

QUESTION ID

S63-M-Q4

TYPE

Short answer

CORRECT ANSWER

Every member has a chance to be chosen, so the sample tends to represent the population and the statistic estimates the parameter without systematic bias.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A school uses the first 30 students arriving in the morning to estimate the average hours of sleep. Is this a random sample? Explain a possible bias.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q5/5

2313

QUESTION ID

S63-M-Q5

TYPE

Short answer

CORRECT ANSWER

No: early arrivers may differ (for example they may sleep less), so the estimate may be biased

GRADING NOTE Form required: decision and a plausible bias.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • MAIN • Q5/5

SET 63 • HSS-IC.A.1

BACKWARD 1

Samples and populations • Grade 7 • Statistics & Probability • 7.SP.A.1

ANSWER KEY Reduced copy of the question page, with its answer

What is a population in statistics?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q1/5

2315

QUESTION ID

S63-B1-Q1

TYPE

Short answer

CORRECT ANSWER

The whole group of interest

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

What is a sample?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q2/5

2316

QUESTION ID

S63-B1-Q2

TYPE

Short answer

CORRECT ANSWER

A part of the population that is studied

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

A random sample gives every member of the population a chance to be chosen.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q3/5

2317

QUESTION ID

S63-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Surveying every student in a school is

A. a statistic

B. an experiment

C. a sample

D. a census

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q4/5

2318

QUESTION ID

S63-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

D. a census

GRADING NOTE Whole population.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q4/5

Why are samples used instead of the whole population?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q5/5

2319

QUESTION ID

S63-B1-Q5

TYPE

Short answer

CORRECT ANSWER

The population may be too large, costly, or slow to study fully

GRADING NOTE Any reasonable answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.A.1 • Set 63 • BACKWARD 1 • Q5/5

SET 63 • HSS-IC.A.1

BACKWARD 2

Draw inferences from random samples • Grade 7 • Statistics & Probability • 7.SP.A.2

ANSWER KEY Reduced copy of the question page, with its answer

In a random sample of 50 students, 20 walk to school. Estimate how many of 600 students walk.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q1/5

2321

QUESTION ID

S63-B2-Q1

TYPE

Short answer

CORRECT ANSWER

240

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Two random samples give 40% and 44%. Is that surprising?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q2/5

2322

QUESTION ID

S63-B2-Q2

TYPE

Short answer

CORRECT ANSWER

No: random samples vary

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q2/5

True or false?

Different random samples from the same population usually give slightly different results.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q3/5

2323

QUESTION ID

S63-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Sampling variability.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

From a random sample, 30% of 200 voters prefer a candidate. The best estimate for all voters is

A. 30%

B. 200

C. cannot estimate

D. 60%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q4/5

2324

QUESTION ID

S63-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 30%

GRADING NOTE Statistic estimates parameter.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Why is a larger random sample usually better?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q5/5

2325

QUESTION ID

S63-B2-Q5

TYPE

Short answer

CORRECT ANSWER

Its statistic varies less, so it estimates the parameter more precisely

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.A.2 • Set 63 • BACKWARD 2 • Q5/5

SET 63 • HSS-IC.A.1

BACKWARD 3

Compute sample proportions and means • Grade 6 • Statistics & Probability • 6.SP.B.5.c

ANSWER KEY Reduced copy of the question page, with its answer

248 of 400 people say yes. What proportion is that?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q1

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q1/5

2327

QUESTION ID

S63-B3-Q1

TYPE

Short answer

CORRECT ANSWER

0.62

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q1

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Find the mean of 5.0, 5.1, 4.9, 5.08.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q2

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q2/5

2328

QUESTION ID

S63-B3-Q2

TYPE

Short answer

CORRECT ANSWER

5.02

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q2

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q2/5

True or false?

A proportion can be greater than 1.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B3-Q3

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q3/5

2329

QUESTION ID

S63-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE A part of a whole is between 0 and 1.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION

ID S63-B3-Q3

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

1,600 of 2,000 is

A. 16%

B. 1.25%

C. 80%

D. 20%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q4

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q4/5

2330

QUESTION ID

S63-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. 80%

GRADING NOTE 1600/2000.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q4

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Write 62% as a decimal.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q5

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q5/5

2331

QUESTION ID

S63-B3-Q5

TYPE

Short answer

CORRECT ANSWER

0.62

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-B3-Q5

Grade 6 • Statistics & Probability • 6.SP.B.5.c • Set 63 • BACKWARD 3 • Q5/5

SET 63 • HSS-IC.A.1

FORWARD 1

Describe sampling variability • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1

ANSWER KEY Reduced copy of the question page, with its answer

Two random samples of 400 residents give 62% and 59%. Explain why they differ and what each estimates.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q1/5

2333

QUESTION ID

S63-F1-Q1

TYPE

Short answer

CORRECT ANSWER

Random samples vary by chance; both estimate the same population percent.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Would samples of 40 or of 400 vary more from sample to sample? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q2/5

2334

QUESTION ID
S63-F1-Q2

TYPE
Short answer

CORRECT ANSWER
Samples of 40: smaller samples are affected more by chance

GRADING NOTE Form required: answer and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q2/5

True or false?

A parameter changes from sample to sample.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q3/5

2335

QUESTION ID

S63-F1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE The parameter is fixed; the statistic varies.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which is a statistic?

A. the population size

B. the mean height of 30 sampled students

C. the mean height of all students

D. the population proportion

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q4/5

2336

QUESTION ID

S63-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. the mean height of 30 sampled students

GRADING NOTE From a sample.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Explain why a large voluntary-response sample can still be worse than a small random sample.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q5/5

2337

QUESTION ID
S63-F1-Q5

TYPE
Short answer

CORRECT ANSWER
Size does not remove bias; a voluntary sample is systematically unrepresentative.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.1 • Set 63 • FORWARD 1 • Q5/5

SET 63 • HSS-IC.A.1

FORWARD 2

Quantify variability with a margin of error • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4

ANSWER KEY Reduced copy of the question page, with its answer

Simulated proportions from samples of 400 with $p = 0.62$ have SD 0.024. Give a margin of error.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q1/5

2339

QUESTION ID

S63-F2-Q1

TYPE

Short answer

CORRECT ANSWER

About 0.048 (4.8 percentage points)

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q1/5

Use the margin of error to give an interval for the percent of residents who support the park.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q2/5

2340

QUESTION ID
S63-F2-Q2

TYPE
Short answer

CORRECT ANSWER
About 57.2% to 66.8%

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q2/5

True or false?

A margin of error describes chance variation, not bias.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q3/5

2341

QUESTION ID
S63-F2-Q3

TYPE
True or false

CORRECT ANSWER
A. True

GRADING NOTE Bias is not measured.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q3/5

If the sample size is multiplied by 4, the margin of error is about

A. unchanged

B. halved

C. doubled

D. divided by 4

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q4/5

2342

QUESTION ID

S63-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. halved

GRADING NOTE Square-root law.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q4/5

Explain what the interval says about the population percent.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q5/5

2343

QUESTION ID
S63-F2-Q5

TYPE
Short answer

CORRECT ANSWER
The population percent is plausibly within the interval; values far outside it are unlikely.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S63-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.4 • Set 63 • FORWARD 2 • Q5/5

SET 64 • HSS-IC.A.2 • STATISTICS, INFERENCE, AND DECISIONS

Decide whether a model is consistent with observed results using a simulation

MAIN	Algebra II main question	**HSS-IC.A.2**
BACKWARD 1	Probability as long-run relative frequency	**7.SP.C.6**
BACKWARD 2	Use simulations of compound events	**7.SP.C.8.c**
BACKWARD 3	Relative frequency of a group of outcomes	**6.RP.A.3.c**
FORWARD 1	Design a simulation to test a model	**HSS-IC.A.2**
FORWARD 2	Compare two treatments with a re-randomization simulation	**HSS-IC.B.5**


SET 64 • HSS-IC.A.2

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2

ANSWER KEY Reduced copy of the question page, with its answer

A coin is claimed to be fair. It landed heads 34 times in 50 flips. The histogram shows 200 simulated runs of 50 fair flips. Is the claim consistent with the result? Justify with the simulation.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q1/5

2346

QUESTION ID
S64-M-Q1

TYPE
Short answer

CORRECT ANSWER
No: only 3 of 200 runs (1.5%) had 34 or more heads, so 34 heads is very unusual for a fair coin


GRADING NOTE Form required: decision and the simulated proportion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A spinner is claimed to land on red 25% of the time. It landed on red 13 times in 40 spins. The histogram shows 200 simulated runs. Is the claim consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q2/5

2347

QUESTION ID
S64-M-Q2

TYPE
Short answer

CORRECT ANSWER
Yes: 13 or more reds happened in 34 of 200 runs (17%), which is not unusual


GRADING NOTE Form required: decision and the simulated proportion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

A die is claimed to be fair. It showed 16 sixes in 60 rolls. The histogram shows 200 simulated runs. Is the claim consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q3/5

2348

QUESTION ID
S64-M-Q3

TYPE
Short answer

CORRECT ANSWER
No: 16 or more sixes happened in only 3 of 200 runs (1.5%), so the result is unusual for a fair die


GRADING NOTE Form required: decision and proportion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A player claims to make 70% of free throws. She made 11 of 20. The histogram shows 200 simulated runs of 20 shots. Is the claim consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q4/5

2349

QUESTION ID
S64-M-Q4

TYPE
Short answer

CORRECT ANSWER
Yes: 11 or fewer made shots happened in 6 + 18 = 24 of 200 runs (12%), which is not unusual


GRADING NOTE Form required: decision and proportion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

A model says boys and girls are equally likely. A hospital had 58 boys in 100 births. The histogram shows 200 simulated runs. Is the model consistent? Justify.



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q5/5

2350

QUESTION ID
S64-M-Q5

TYPE
Short answer

CORRECT ANSWER
Yes: 58 or more boys happened in 19 of 200 runs (9.5%), which is not unusual at the 5% level

GRADING NOTE Form required: decision and proportion.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • MAIN • Q5/5

SET 64 • HSS-IC.A.2

BACKWARD 1

Probability as long-run relative frequency • Grade 7 • Statistics & Probability • 7.SP.C.6

ANSWER KEY Reduced copy of the question page, with its answer

A fair coin is flipped 50 times. About how many heads do you expect?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q1/5

2352

QUESTION ID

S64-B1-Q1

TYPE

Short answer

CORRECT ANSWER

25

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q1

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A spinner has P(red) = 0.25. About how many reds in 40 spins?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q2/5

2353

QUESTION ID

S64-B1-Q2

TYPE

Short answer

CORRECT ANSWER

10

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q2

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q2/5

True or false?

Getting exactly the expected number every time is likely.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q3/5

2354

QUESTION ID

S64-B1-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Results vary.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q3

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q3/5

A fair die is rolled 60 times. Expected sixes:

A. 10

B. 16

C. 6

D. 60

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q4/5

2355

QUESTION ID

S64-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

A. 10

GRADING NOTE 60/6.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q4

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Why is 26 heads in 50 flips not surprising?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q5/5

2356

QUESTION ID

S64-B1-Q5

TYPE

Short answer

CORRECT ANSWER

It is close to the expected 25; chance variation is normal

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B1-Q5

Grade 7 • Statistics & Probability • 7.SP.C.6 • Set 64 • BACKWARD 1 • Q5/5

SET 64 • HSS-IC.A.2

BACKWARD 2

Use simulations of compound events • Grade 7 • Statistics & Probability • 7.SP.C.8.c

Describe how to simulate 50 fair coin flips with random digits.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q1/5

2358

QUESTION ID

S64-B2-Q1

TYPE

Short answer

CORRECT ANSWER

Use 50 random digits; even = heads, odd = tails

GRADING NOTE Any valid design.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q1

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

How could you simulate one spin of a spinner with P(red) = 0.25 using two-digit random numbers 00-99?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q2/5

2359

QUESTION ID

S64-B2-Q2

TYPE

Short answer

CORRECT ANSWER

00-24 = red;

25-99 = not red

GRADING NOTE Any valid design with 25 of the 100 numbers for red.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q2

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q2/5

True or false?

Each run of a simulation should copy the real process under the model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q3/5

2360

QUESTION ID

S64-B2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Model.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q3

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q3/5

A simulation has 200 runs. Each run represents

A. the final answer

B. one repetition of the whole experiment

C. one histogram bar

D. one flip

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q4/5

2361

QUESTION ID

S64-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. one repetition of the whole experiment

GRADING NOTE Run = experiment.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q4

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q4/5

Why are many runs used?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q5/5

2362

QUESTION ID

S64-B2-Q5

TYPE

Short answer

CORRECT ANSWER

To see the pattern of results that the model produces by chance

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B2-Q5

Grade 7 • Statistics & Probability • 7.SP.C.8.c • Set 64 • BACKWARD 2 • Q5/5

SET 64 • HSS-IC.A.2

BACKWARD 3

Relative frequency of a group of outcomes • Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c

ANSWER KEY Reduced copy of the question page, with its answer

3 of 200 runs is what percent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q1/5

2364

QUESTION ID

S64-B3-Q1

TYPE

Short answer

CORRECT ANSWER

1.5%

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q1

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

6 + 18 runs out of 200 is what percent?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q2/5

2365

QUESTION ID

S64-B3-Q2

TYPE

Short answer

CORRECT ANSWER

12%

GRADING NOTE Any equivalent form.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q2

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q2/5

ANSWER KEY Reduced copy of the question page, with its answer

True or false?

19 out of 200 is 19%.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q3/5

2366

QUESTION ID

S64-B3-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE 19/200 = 0.095 = 9.5%.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q3

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

34 of 200 is

A. 1.7%

B. 17%

C. 6.8%

D. 34%

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q4/5

2367

QUESTION ID

S64-B3-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. 17%

GRADING NOTE 34/200.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q4

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Is 2% less than 5%?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q5/5

2368

QUESTION ID

S64-B3-Q5

TYPE

Short answer

CORRECT ANSWER

Yes

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-B3-Q5

Grade 6 • Ratios & Proportional Relationships • 6.RP.A.3.c • Set 64 • BACKWARD 3 • Q5/5

SET 64 • HSS-IC.A.2

FORWARD 1

Design a simulation to test a model • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2

ANSWER KEY Reduced copy of the question page, with its answer

Describe a simulation to test whether a basketball player really makes 70% of free throws, given that she made 11 of 20.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q1/5

2370

QUESTION ID

S64-F1-Q1

TYPE

Short answer

CORRECT ANSWER

Use random digits 0-9 with 0-6 = make; take 20 digits per run, count makes; repeat many runs; see how often 11 or fewer makes occur.

GRADING NOTE Any valid design with all parts.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

A player who claims to make 70% of free throws made 11 of 20. Why should a simulation count results of 11 or fewer (as extreme or more extreme) rather than exactly 11?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q2/5

2371

QUESTION ID

S64-F1-Q2

TYPE

Short answer

CORRECT ANSWER

Any single exact value can be rare; the question is how unusual results this far from the claim are.

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q2/5

True or false?

A result that happens in 30% of simulated runs is strong evidence against the model.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q3/5

2372

QUESTION ID
S64-F1-Q3

TYPE
True or false

CORRECT ANSWER
B. False

GRADING NOTE 30% is common.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A common cutoff for "unusual" is

A. about 50% of runs

B. 100% of runs

C. about 5% of runs

D. about 25% of runs

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q4/5

2373

QUESTION ID

S64-F1-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. about 5% of runs

GRADING NOTE Convention.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

Doubling the number of runs from 200 to 400 changes what about the simulation?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q5/5

2374

QUESTION ID

S64-F1-Q5

TYPE

Short answer

CORRECT ANSWER

It gives a more precise estimate of how often the result occurs under the model (the model and data are unchanged).

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F1-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.A.2 • Set 64 • FORWARD 1 • Q5/5


SET 64 • HSS-IC.A.2

FORWARD 2

Compare two treatments with a re-randomization simulation • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5

ANSWER KEY Reduced copy of the question page, with its answer

In an experiment, fertilized plants grew 2.9 cm more on average. In 200 re-randomizations, none of the differences was 2.9 cm or more. What do you conclude?



Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q1/5

2376

QUESTION ID
S64-F2-Q1

TYPE
Short answer

CORRECT ANSWER

The difference is very unlikely by chance alone, so the fertilizer appears to increase growth

GRADING NOTE Form required: conclusion and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

What model does the re-randomization assume?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q2/5

2377

QUESTION ID
S64-F2-Q2

TYPE
Short answer

CORRECT ANSWER
That the treatment has no effect, so group labels can be shuffled

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q2/5

True or false?

Re-randomization is a simulation of a model, like the coin-flip simulation.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q3/5

2378

QUESTION ID

S64-F2-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Both test a model.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q3/5

If 60 of 200 re-randomized differences were at least the observed one, the result is

A. significant

B. proof of no effect

C. not statistically significant

D. impossible

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q4/5

2379

QUESTION ID

S64-F2-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. not statistically significant

GRADING NOTE 30% is common.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q4/5

Explain how a re-randomization test for an experiment extends the reasoning used to test a fair-coin claim with a simulation.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S64-F2-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q5/5

2380

QUESTION ID
S64-F2-Q5

TYPE
Short answer

CORRECT ANSWER

Both ask how often a model produces a result as extreme as the observed one; here the model is "no treatment effect".

GRADING NOTE Any equivalent explanation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S64-F2-Q5**

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.5 • Set 64 • FORWARD 2 • Q5/5

SET 65 • HSS-IC.B.3 • STATISTICS, INFERENCE, AND DECISIONS

Identify sample surveys, experiments, and observational studies

MAIN	Algebra II main question	**HSS-IC.B.3**
BACKWARD 1	Statistical questions and data collection	**6.SP.A.1**
BACKWARD 2	Association is not causation	**HSS-ID.C.9**
BACKWARD 3	Random samples	**7.SP.A.1**
FORWARD 1	Identify confounding and design flaws	**HSS-IC.B.3**
FORWARD 2	Analyze the results of an experiment	**HSS-IC.B.5**

SET 65 • HSS-IC.B.3

MAIN

Algebra II main question • High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3

Researchers randomly assign 100 volunteers to take a vitamin or a placebo and compare colds. What type of study is this, and what is its purpose?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q1/5

2383

QUESTION ID

S65-M-Q1

TYPE

Short answer

CORRECT ANSWER

Experiment: to test whether the vitamin causes fewer colds

GRADING NOTE Form required: type and purpose.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-M-Q1

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q1/5

A pollster asks a random sample of 1,000 voters which candidate they prefer. What type of study is this?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-M-Q2

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q2/5

2384

QUESTION ID
S65-M-Q2

TYPE
Short answer

CORRECT ANSWER
Sample survey: to estimate the preference of all voters

GRADING NOTE Form required: type and purpose.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S65-M-Q2**

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q2/5

Researchers compare the hospital records of smokers and nonsmokers for lung disease. What type of study is this? Can it show causation?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-M-Q3

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q3/5

2385

QUESTION ID

S65-M-Q3

TYPE

Short answer

CORRECT ANSWER

Observational study; it cannot by itself show causation because the groups were not assigned

GRADING NOTE Form required: type and the causation answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S65-M-Q3**

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Which type of study best answers "Does playing music while studying improve test scores?" Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q4/5

2386

QUESTION ID
S65-M-Q4

TYPE
Short answer

CORRECT ANSWER
Experiment: randomly assign students to study with or without music and compare scores

GRADING NOTE Form required: type and design idea.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-M-Q4

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q4/5

Which type of study best answers "What percent of students at the school use the library weekly?" Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-M-Q5

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q5/5

2387

QUESTION ID

S65-M-Q5

TYPE

Short answer

CORRECT ANSWER

Sample survey of a random sample of students

GRADING NOTE Form required: type and reason.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION **ID S65-M-Q5**

High School — Algebra II • Statistics & Probability: Making Inferences & Justifying Conclusions • HSS-IC.B.3 • Set 65 • MAIN • Q5/5

SET 65 • HSS-IC.B.3

BACKWARD 1

Statistical questions and data collection • Grade 6 • Statistics & Probability • 6.SP.A.1

ANSWER KEY Reduced copy of the question page, with its answer

Is "How many hours do students at our school sleep?" a statistical question? Why?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q1

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q1/5

2389

GRADING NOTE Form required: decision and reason.

QUESTION ID

S65-B1-Q1

TYPE

Short answer

CORRECT ANSWER

Yes: the answers vary

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q1

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Give a data collection method for "What do voters think?"

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q2

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q2/5

2390

QUESTION ID

S65-B1-Q2

TYPE

Short answer

CORRECT ANSWER

Ask a sample of voters (a survey)

GRADING NOTE Any reasonable answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q2

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q2/5

True or false?

An experiment imposes a treatment on the subjects.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q3

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q3/5

2391

QUESTION ID

S65-B1-Q3

TYPE

True or false

CORRECT ANSWER

A. True

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q3

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

Watching and recording without imposing treatments is

A. an experiment

B. an observational study

C. a census

D. a placebo

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q4

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q4/5

2392

QUESTION ID

S65-B1-Q4

TYPE

Multiple choice

CORRECT ANSWER

B. an observational study

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q4

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q4/5

ANSWER KEY Reduced copy of the question page, with its answer

What is a placebo?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q5

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q5/5

2393

QUESTION ID

S65-B1-Q5

TYPE

Short answer

CORRECT ANSWER

A fake treatment with no active ingredient

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B1-Q5

Grade 6 • Statistics & Probability • 6.SP.A.1 • Set 65 • BACKWARD 1 • Q5/5

SET 65 • HSS-IC.B.3

BACKWARD 2

Association is not causation • High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9

ANSWER KEY Reduced copy of the question page, with its answer

Ice cream sales and sunburns are correlated. Does ice cream cause sunburn? Explain.

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q1/5

2395

QUESTION ID
S65-B2-Q1

TYPE
Short answer

CORRECT ANSWER
No: hot sunny weather increases both

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q1

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q1/5

ANSWER KEY Reduced copy of the question page, with its answer

Name a lurking variable for "students with more books at home get higher scores".

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q2/5

2396

QUESTION ID

S65-B2-Q2

TYPE

Short answer

CORRECT ANSWER

For example family income or parent education

GRADING NOTE Any reasonable answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q2

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q2/5

True or false?

A strong correlation proves that one variable causes the other.

A. True

B. False

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q3/5

2397

QUESTION ID

S65-B2-Q3

TYPE

True or false

CORRECT ANSWER

B. False

GRADING NOTE Correlation is not causation.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q3

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q3/5

ANSWER KEY Reduced copy of the question page, with its answer

A variable that affects both variables in a study is

A. the placebo

B. the sample

C. a confounding (lurking) variable

D. the response

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q4/5

2398

QUESTION ID

S65-B2-Q4

TYPE

Multiple choice

CORRECT ANSWER

C. a confounding (lurking) variable

GRADING NOTE Definition.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q4

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q4/5

What kind of study can show cause and effect?

Answer: _____

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q5/5

2399

QUESTION ID

S65-B2-Q5

TYPE

Short answer

CORRECT ANSWER

A randomized experiment

GRADING NOTE Any equivalent answer.

REFERENCE LABEL — NOT PART OF THE STUDENT QUESTION ID S65-B2-Q5

High School — Algebra I • Statistics & Probability: Interpreting Categorical & Quantitative Data • HSS-ID.C.9 • Set 65 • BACKWARD 2 • Q5/5

SET 65 • HSS-IC.B.3

BACKWARD 3

Random samples • Grade 7 • Statistics & Probability • 7.SP.A.1