

Exponential Proof Complexity Lower Bounds on High-Dimensional Ramanujan Complexes

Coboundary Expansion, Linear 2-Systoles, and the $2^{\Omega(n)}$ Resolution Width Barrier

Srijan Mandal

Independent Research · ORCID: 0009-0002-6899-6185 · SSRN: 7461081
creatorofaurad@gmail.com · <https://github.com/creatorofaurad/1M>

September 16, 2026

Abstract

In this monograph, I establish an unconditional exponential lower bound of form $2^{\Omega(n)}$ for Resolution refutations of constraint systems derived from 2-dimensional simplicial Ramanujan complexes (Lubotzky–Samuels–Vishne). By moving beyond single-gate information theory and framing the refutation as an information bottleneck across topological cuts, I eliminate the classical $\mathcal{O}(\log n)$ DAG-to-tree reconvergence penalty.

The proof proceeds via two core geometric lemmas:

1. **Lemma 1 (The 2-Systole Boundary Measure):** I define a sub-clause progress measure $\mu(C)$ tracking the minimal number of 2-face axioms required to derive any clause C . By the sub-additivity of resolution inferences and the linear 2-systole $\text{Sys}_2(X) \geq \mu_0 n$, there exists a critical clause C^* at the median threshold $\frac{1}{3}\text{Sys}_2(X) \leq \mu(C^*) \leq \frac{2}{3}\text{Sys}_2(X)$ whose active boundary variables satisfy $|C^*| \geq \gamma \cdot \mu(C^*) = \Omega(n)$ via γ -coboundary expansion.
2. **Lemma 2 (The 2-Complex Resolution Lifting):** I prove that the minimal refutation width of the parity-check formula Φ_X is lower-bounded by $\text{Width}(\Phi_X \vdash \perp) \geq \alpha n$, where $\alpha = \frac{1}{3}\gamma\mu_0 > 0$.

Applying the Ben-Sasson–Wigderson Size-Width Theorem directly to the 2-complex yields:

$$\text{Size}(\Pi \vdash \perp) \geq \exp\left(\frac{(\text{Width}(\Phi_X \vdash \perp) - k)^2}{n}\right) = \exp\left(\frac{(\alpha n - 7)^2}{n}\right) = 2^{\Omega(n)}.$$

This demonstrates that 2-dimensional topological expansion creates an uncompressible memory bottleneck, forcing exponential refutation size.

Contents

1	Introduction and Theoretical Substrate	2
2	The Parity-Check Formula Φ_X	2
3	Lemma 1: The 2-Systole Boundary Measure	3

4	Lemma 2: The 2-Complex Resolution Lifting	4
5	The Exponential Size Lower Bound ($2^{\Omega(n)}$)	4
6	Conclusion and Complexity Implications	4

1 Introduction and Theoretical Substrate

A central challenge in circuit and proof complexity is the gap between formula lower bounds ($2^{\Omega(n)}$) and general DAG lower bounds ($2^{\Omega(n/\log n)}$). In standard 1-dimensional expander graphs, an adversarial DAG can evaluate intermediate linear parities, bypassing local correlation bounds.

In this work, I resolve this bottleneck by lifting the underlying structure to a ****2-Dimensional Simplicial Ramanujan Complex**** $X = (X^{(0)}, X^{(1)}, X^{(2)})$, constructed via Lubotzky–Samuels–Vishne (LSV) arithmetic lattices over local fields.

Definition 1.1 ((n, k, γ) -Ramanujan 2-Complex). A 2-dimensional simplicial complex X with n vertices is an (n, k, γ) -Ramanujan complex if:

1. Each 2-face $\sigma \in X^{(2)}$ has boundary length $|\partial\sigma| \leq k$ (with $k = 3$ or 7).
2. The 2-systole of X over $\mathbb{F}_2$ is linear in the number of vertices:

$$\text{Sys}_2(X) = \min\{|\omega|_0 \mid \omega \in Z_2(X, \mathbb{F}_2) \setminus B_2(X, \mathbb{F}_2)\} \geq \mu_0 \cdot n$$

for an absolute geometric constant $\mu_0 > 0$.

3. X exhibits γ -coboundary expansion on cochains $\eta \in C^1(X, \mathbb{F}_2)$:

$$|\delta^1 \eta| \geq \gamma \cdot \min_{\tau \in B^1(X, \mathbb{F}_2)} |\eta \oplus \tau|$$

for a spectral constant $\gamma > 0$.

2 The Parity-Check Formula Φ_X

Let X be an (n, k, γ) -Ramanujan 2-complex. We associate each 1-cell (edge) $e \in X^{(1)}$ with a Boolean variable $x_e \in \{0, 1\}$.

Definition 2.1 (Face-Parity CNF Formula Φ_X). For each 2-face $\sigma \in X^{(2)}$ with boundary 1-cycle $\partial\sigma = \{e_1, e_2, \dots, e_k\}$, we enforce the parity constraint:

$$\bigoplus_{e \in \partial\sigma} x_e = 1.$$

Expanding each parity constraint into 2^{k-1} standard CNF clauses yields the formula $\Phi_X = \bigwedge_{\sigma \in X^{(2)}} \text{CNF}(\partial\sigma)$. The initial clause width is $\text{width}(\Phi_X) = k \leq 7$.

Lemma 2.2 (Global Unsatisfiability via 2-Systole). *The formula Φ_X is unsatisfiable.*

Proof. Summing the parity constraints over all 2-faces in a closed 2-cycle $Z \in Z_2(X, \mathbb{F}_2)$ over $\mathbb{F}_2$:

$$\sum_{\sigma \in Z} \bigoplus_{e \in \partial\sigma} x_e = \bigoplus_{e \in X^{(1)}} \deg_Z(e) \cdot x_e \equiv 0 \pmod{2}$$

because every edge e in a 2-cycle appears in an even number of 2-faces ($\deg_Z(e) \equiv 0 \pmod{2}$). However, the sum of the right-hand sides is $|Z| \pmod{2} = 1 \pmod{2}$ for an odd 2-cycle. Since X contains non-trivial odd 2-cycles with $|Z| \geq \text{Sys}_2(X) \geq \mu_0 n$, Φ_X has no satisfying assignment. $\square$

3 Lemma 1: The 2-Systole Boundary Measure

Let $\Pi = (C_1, C_2, \dots, C_S)$ be a valid Resolution refutation of Φ_X terminating at the empty clause $C_S = \perp$.

Definition 3.1 (Sub-Clause Axiom Measure $\mu(C)$). For any derived clause $C \in \Pi$, define $\mu(C)$ as the minimum number of 2-face axiom sets $\text{CNF}(\partial\sigma)$ required to semantically imply C :

$$\mu(C) = \min\{|S| \mid S \subseteq X^{(2)}, \bigwedge_{\sigma \in S} \text{CNF}(\partial\sigma) \models C\}.$$

Lemma 3.2 (Properties of the Measure μ). *The measure μ satisfies:*

1. **Base Case:** For any initial axiom clause $A \in \Phi_X$, $\mu(A) = 1$.
2. **Terminal Case:** For the empty clause $\perp$, $\mu(\perp) \geq \text{Sys}_2(X) \geq \mu_0 n$.
3. **Sub-Additivity:** If clause C is derived from parents A and B via the resolution rule $\frac{A \vee x \quad B \vee \neg x}{A \vee B}$, then:

$$\mu(C) \leq \mu(A) + \mu(B).$$

Proof. If S_A implies A and S_B implies B , then $S_A \cup S_B$ semantically implies $A \vee B$. Thus $\mu(C) \leq |S_A \cup S_B| \leq \mu(A) + \mu(B)$. For $\perp$, any set S implying $\perp$ must contain a closed 2-cycle, forcing $|S| \geq \text{Sys}_2(X)$. $\square$

Theorem 3.3 (Existence of the Critical High-Width Clause). *There exists a clause $C^* \in \Pi$ such that:*

$$\frac{1}{3}\text{Sys}_2(X) \leq \mu(C^*) \leq \frac{2}{3}\text{Sys}_2(X)$$

and its active variable count satisfies:

$$|C^*| = |\text{Vars}(C^*)| \geq \gamma \cdot \mu(C^*) \geq \frac{1}{3}\gamma\mu_0 n = \Omega(n).$$

Proof. Because $\mu(\text{Axiom}) = 1$ and resolution is binary ($\mu(C) \leq \mu(A) + \mu(B)$), the measure μ cannot jump over an interval of width $> \mu(A) \geq 1$. Thus, there exists a clause C^* in Π whose measure lies in $[\frac{1}{3}\text{Sys}_2, \frac{2}{3}\text{Sys}_2]$.

Let $S \subseteq X^{(2)}$ be the minimal set of faces of size $|S| = \mu(C^*)$ implying C^* . Because $|S| \leq \frac{2}{3}\text{Sys}_2(X) < \text{Sys}_2(X)$, S cannot contain any non-trivial closed 2-cycle. Therefore, its geometric 1-boundary $\partial_2 S \in C_1(X, \mathbb{F}_2)$ is strictly non-zero.

By the γ -coboundary expansion of X :

$$|\partial_2 S| \geq \gamma \cdot |S| = \gamma \cdot \mu(C^*).$$

Every edge $e \in \partial_2 S$ appears in an odd number of faces of S . If $x_e \notin \text{Vars}(C^*)$, flipping the value of x_e would alter the satisfiability of the axioms in S while leaving C^* unaffected, contradicting the minimality of S . Therefore, every boundary edge must appear as a literal in C^* :

$$|C^*| \geq |\partial_2 S| \geq \gamma \cdot \mu(C^*) \geq \frac{1}{3}\gamma\mu_0 n = \Omega(n).$$

$\square$

4 Lemma 2: The 2-Complex Resolution Lifting

Theorem 4.1 (Resolution Refutation Width Lower Bound). *For any (n, k, γ) -Ramanujan 2-complex X , the resolution refutation width of Φ_X satisfies:*

$$\text{Width}(\Phi_X \vdash \perp) \geq \alpha \cdot n$$

where $\alpha = \frac{1}{3}\gamma\mu_0 > 0$.

Proof. The refutation width is defined as $\text{Width}(\Pi) = \max_{C \in \Pi} |C|$. By Theorem 3.3, the refutation Π contains the critical clause C^* with $|C^*| \geq \alpha n$. Since this holds for every valid resolution refutation Π of Φ_X :

$$\text{Width}(\Phi_X \vdash \perp) = \min_{\Pi \vdash \perp} \text{Width}(\Pi) \geq |C^*| \geq \alpha n = \Omega(n).$$

□

5 The Exponential Size Lower Bound ($2^{\Omega(n)}$)

We now invoke the fundamental Size-Width Theorem of Ben-Sasson and Wigderson.

Theorem 5.1 (Ben-Sasson–Wigderson Size-Width Theorem, JACM 2001). *Let Φ be an unsatisfiable CNF formula on N variables with initial clause width k . The size of any tree-like or general DAG Resolution refutation $\Pi \vdash \perp$ satisfies:*

$$\text{Size}(\Pi) \geq \exp \left(\frac{(\text{Width}(\Phi \vdash \perp) - k)^2}{2N} \right).$$

Theorem 5.2 (Master Exponential Lower Bound on Ramanujan 2-Complexes). *Let X be an (n, k, γ) -Ramanujan 2-complex with $N = |X^{(1)}| \leq c_1 n$ edges and maximum face degree $k \leq 7$. Then every Resolution refutation of the face-parity formula Φ_X has size:*

$$\text{Size}(\Pi \vdash \perp) \geq \exp \left(\frac{(\alpha n - 7)^2}{2c_1 n} \right) = 2^{\Omega(n)}.$$

Proof. Substituting $\text{Width}(\Phi_X \vdash \perp) \geq \alpha n$ (Theorem 4.1), $k \leq 7$, and $N \leq c_1 n$ into Theorem 5.1:

$$\text{Size}(\Pi \vdash \perp) \geq \exp \left(\frac{(\alpha n - 7)^2}{2c_1 n} \right) = \exp \left(\frac{\alpha^2 n^2 - 14\alpha n + 49}{2c_1 n} \right) = \exp \left(\frac{\alpha^2}{2c_1} n - \mathcal{O}(1) \right) = 2^{\Omega(n)}.$$

□

6 Conclusion and Complexity Implications

By shifting the lower-bound argument from local gate correlation to **2-dimensional coboundary expansion and linear 2-systoles**, the proof establishes:

1. ****Elimination of the Spira Loss:**** The information bottleneck is enforced by the topological boundary $|\partial_2 S| \geq \gamma |S|$, which cannot be compressed by gate sharing in a DAG.
2. ****Unconditional Proof Complexity Separation:**** For parity formulas on Ramanujan 2-complexes, Resolution refutations require size $2^{\Omega(n)}$.