YANG–MILLS MASS GAP ON THE LATTICE AND ITS REDUCTION TO ONE CUBE OF THE CONTINUUM

DEEP BHATTACHARJEE*, USHASHI BHATTACHARYA, JAMES MONTGOMERY LOCKWOOD, AND TRISTAN ROMANOV

Abstract. *We prove that four-dimensional lattice Yang–Mills theory with any compact simple gauge group has a unique infinite-volume state and an explicit transfer-matrix gap in a forest gauge, up to* $\beta_W = 0.351$ *for* SU(2) *and* $\beta_W = 0.77$ *for* SU(3). *We prove that a Dobrushin–Shlosman estimate in one cube of fixed physical size gives a mass gap in every continuum limit, and that the cube needs a side of order* $(\beta_W/\log\beta_W)^{1/4}$ *lattice spacings. We prove that the curvature of Balaban's block averages costs action independent of the scale, that a translation of the Haar measures makes it rare at every scale for the abelian theory, and that for* SU(2) *every such translation transported along a forest reaches its constant on the gauge orbit of the vacuum, which fixes the one estimate that remains open. We prove that, once the blocks are labelled by their independent links and their coarse curvature, every good window carries an exact product law in which each path link is determined by its pair, that two bad islands interact only through one kernel on the good layer between them, with covariance at most* $\frac{1}{4}\tanh(\Delta/4)$, *and that the exact response of a plaquette block at zero field,* $\frac{1}{4}(1 + I_2(\beta_W)/I_1(\beta_W))$, *bounds every Dobrushin estimate of this form by* $\beta_W = 0.3666$. *We give the explicit coupling,* $\beta_W = 5.4 \cdot 10^{21}$ *at the label radius* 10^{-8}, *past which the islands of bad blocks at one scale are rarer than their entropy. We prove that the rarity of bad labels alone cannot control their covariance and that a rough coarse background makes the constrained action nonconvex, which locates the one estimate that remains open.*

1. Introduction

Wilson's lattice gauge theory [83] assigns an element Q_e of a compact group $G \subset \mathrm{U}(N)$ to every link e of $\mathbb{Z}^4$ and weighs a configuration by $\exp(N\beta \sum_p \mathrm{Re}\,\mathrm{Tr}\,Q_p)$, where Q_p is the ordered product of the four link variables around the plaquette p.[1] At small $|\beta|$ the theory is under control: it has a unique infinite-volume state, correlations decay exponentially, and

Date: 4 November 2026.

2020 *Mathematics Subject Classification.* Primary 81T25, 82B20; Secondary 81T13, 60K35, 82B26, 22E30. PACS 11.15.Ha, 11.15.Tk, 05.50.+q, 02.50.Cw.

Key words and phrases. Lattice gauge theory, mass gap, Dobrushin uniqueness, block averages, reflection positivity.

*Corresponding author: Deep Bhattacharjee, itsdeep@live.com.

An earlier version of this paper is available at `https://hexagonmath.org/2610.00341v6`. New in this version is Section 13: the product law on good windows with each path link determined by its pair (Lemma 13.3, Proposition 13.4), the rough background with negative constrained curvature (Proposition 13.1), the nonconvex bad side and its three-preimage branch family (Propositions 13.5 and 13.6), the one-kernel interaction of two bad islands with the covariance bound $\frac{1}{4}\tanh(\Delta/4)$ (Lemma 13.9, Corollary 13.10), the floor $q^2/16$ on the covariance of rare labels (Proposition 13.12), the explicit island threshold $\beta^*(\rho, s)$ past which the islands of bad blocks at one scale are rarer than their entropy (Theorem 13.21), the coupling-cost minimality lemmas (Lemmas 13.13 and 13.14), the exact block response $\frac{1}{4}(1 + I_2(\beta_W)/I_1(\beta_W))$ and its crossing at $\beta_W = 0.3666$ (Proposition 13.15), the weighted Schur locality lemma with its perturbation ledger (Lemma 13.16, Proposition 13.17), and the positive-time spectral bounds (Lemmas 13.18 and 13.19, Proposition 13.20); the abstract, the introduction and the remark after Theorem 1.3 state them.

[1] This is the normalisation of Shen, Zhu and Zhu [74] and of our earlier paper [15]. The conventional inverse coupling is $\beta_W = N^2\beta$, so the weight is $\exp(\frac{\beta_W}{N}\sum_p \mathrm{Re}\,\mathrm{Tr}\,Q_p)$; for SU(2), identified with the unit quaternions, $\frac{1}{2}\,\mathrm{Tr}\,Q_p = \mathrm{Re}\,Q_p$ and $\beta_W = 4\beta$. Standard accounts of the lattice theory are the books of Montvay and Münster [64] and of Gattringer and Lang [39]; the Hamiltonian version is due to Kogut and Susskind [54].

the transfer matrix has a gap [67, 72].[2] The question is how far into the coupling range such statements can be proved with explicit constants, since the problem of the continuum limit [50] begins where strong coupling ends and Bałaban's weak-coupling analysis [6, 7, 8] takes over.

Two methods give explicit ranges. The Bakry–Émery method [4, 5] of Shen, Zhu and Zhu [74] bounds the Hessian of the action; in [15] we showed that the sharp bound is $4d = 16$ and that no curvature argument for the ungauged measure goes beyond $|\beta| < 1/32$. The Dobrushin method [30, 31][3] bounds the influence of each link on the conditional law of another; for SU(2) the single-link conditional law is a von Mises–Fisher law on the three-sphere, the sharp Kantorovich modulus[4] of that family is one quarter [15] (Theorem I.4 in Appendix I), and every link has eighteen partners, which gives $\beta_W < 2/9$ and nothing beyond, because the coefficients are attained.

Both limits come from the same count. A link of $\mathbb{Z}^4$ lies in six plaquettes and shares each with three other links. Gauge invariance makes part of this redundant: in a box, the variables on a forest of links can be set to the identity without changing any gauge-invariant expectation [27].[5] The forest we use is a field of straight lines with one line through every site, chosen so that every plaquette contains exactly one fixed link. Then every free link has exactly twelve free partners, two in each of its plaquettes (Lemma 2.2), and both methods improve.

Theorem 1.1 (SU(2))**.** *Let $G = \mathrm{SU}(2)$ and $0 \le \beta_W < \frac{1}{3}$. For every gauge-invariant local observable O the expectations $W_\Lambda(O)$ in boxes with free boundary converge, along every sequence of boxes that exhausts $\mathbb{Z}^4$, to a limit $\omega(O)$ that does not depend on the sequence. The state ω is invariant under translations and the hypercubic group and reflection positive;[6] truncated correlations of Lipschitz observables decay at every rate below $-\log(3\beta_W)$; and the transfer matrix has spectral gap*

$$m \ge -\log(3\beta_W) > 0.$$

For $-\frac{1}{3} < \beta_W < 0$ all statements except reflection positivity and the gap hold.

[2] Strong-coupling results of this kind go back to Osterwalder and Seiler [67], who proved exponential clustering and a gap by a convergent expansion, and to Seiler's monograph [72]; the general framework for lattice gauge fields and their continuum limits is that of Brydges, Fröhlich and Seiler [22]. More recent work treats the strong-coupling regime by probabilistic methods: the survey of Chatterjee [25], his large-N solution of SO(N) lattice gauge theory at strong coupling [26] and its planar extension by Basu and Ganguly [11], the Langevin dynamics of Shen, Zhu and Zhu [74], and the area law in an enlarged region by Cao, Nissim and Sheffield [24]. For finite gauge groups Adhikari and Cao proved correlation decay at weak coupling [1]. In two dimensions the continuum Yang–Mills measure has been constructed by Gross, King and Sengupta [42], Driver [33] and Lévy [55]; in four dimensions Bałaban's renormalisation group, as expounded by Dimock [29], gives ultraviolet stability in finite volume, and Magnen, Rivasseau and Sénéor constructed the theory with an infrared cutoff [60]. None of these results gives a mass gap in physical units.

[3] Dobrushin's theorem: if the conditional law of each variable given all others moves, in Kantorovich distance, by at most C_{xy} times the distance by which the variable at y moves, and $\sup_x \sum_y C_{xy} < 1$, then the Gibbs measure is unique and correlations decay at the rate given by the Neumann series of the matrix C. Dobrushin and Shlosman [32] replaced single variables by finite boxes, which is the form used in Section 10.

[4] The Kantorovich distance between two probability measures on a metric space is the least expected distance over their couplings, equivalently the largest difference of their integrals over 1-Lipschitz functions [80]. The modulus of a family of laws indexed by a metric space is the best Lipschitz constant of the map from the index to the law in that distance.

[5] Creutz used a maximal tree to remove the gauge freedom completely; the temporal gauge, with all links in one direction fixed, is the special case of a family of straight lines. In a maximal tree almost every link is fixed in some region and none in others, so the number of partners of a free link is not uniform, and the temporal gauge leaves fourteen partners per link. The point here is a pattern that fixes exactly one link in every plaquette.

[6] That is, $\omega(\vartheta(\bar{F})F) \ge 0$ for every F depending on the links on one side of a lattice hyperplane, ϑ being the reflection in it. It gives the Hilbert space and the transfer matrix of Osterwalder and Schrader [65, 66] and of Osterwalder and Seiler [67], reconstructed in Appendix F.

The step behind Theorem 1.1 is short once the line field is in place. The number $\frac{1}{3}$ is also the end of the road for this kind of argument.

Theorem 1.2 (Ceiling)**.** *Let F' be any set of links of $\mathbb{Z}^4$ without cycles, and give two distinct links outside F' that share a plaquette the coefficient $|\beta|$, the largest single-link Kantorovich coefficient of* $\mathrm{SU}(2)$*. Then the weighted Dobrushin condition with these coefficients fails for $\beta_W \geq \frac{1}{3}$. For the line field the coefficients are attained, so no single-link Kantorovich argument in that gauge reaches $\beta_W = \frac{1}{3}$.*

The ceiling concerns coefficients between single links. Take instead as a block the three free links of one plaquette. The free links of the line field split into such triples with period two (Lemma 5.1), the coupling inside the block plaquette is no longer paid for, and each link keeps ten outside partners. The price is the sensitivity of a law on three spheres instead of one, which we bound analytically by integrating out the middle link of the block.

Theorem 1.3 ($\mathrm{SU}(2)$ with plaquette blocks)**.** *Let $G = \mathrm{SU}(2)$ and $0 \leq \beta_W \leq 0.351$. The number $\alpha(\beta_W)$ of (5.3) is increasing, $\alpha(\frac{1}{3}) \leq 0.9428$, $\alpha(0.345) \leq 0.9805$ and $\alpha(0.351) \leq 0.99995$. The conclusions of Theorem 1.1 hold with decay at every rate below $-\frac{1}{2}\log\alpha(\beta_W)$ and a transfer-matrix gap*

$$m \geq -\tfrac{1}{2}\log\alpha(\beta_W) > 0;$$

at $\beta_W = \frac{1}{3}$ this gives $m \geq 0.0294$. For $-0.351 \leq \beta_W < 0$ all statements except reflection positivity and the gap hold.

At small coupling the rate of Theorem 1.1 is the better one, so both bounds stand side by side; Theorem 1.3 matters past $\beta_W = \frac{1}{3}$, where the gap of Theorem 1.1 has closed.

The curvature method gains as well, and for every N. Let $\operatorname{Hess} S_0$ be the Hessian of $\sum_p \operatorname{Re}\operatorname{Tr} Q_p$ on G^E and F the set of fixed links.

Theorem 1.4 (Curvature in the gauge)**.** *For every N, every field Q and every tangent vector $v = (X_e Q_e)$ with $X_e = 0$ for $e \in F$,*

$$|\operatorname{Hess} S_0(v,v)| \leq \tfrac{29}{2}|v|^2,$$

against the sharp constant 16 *without gauge fixing. Consequently, for $G = \mathrm{SU}(N)$ and $|\beta| < \frac{1}{29}$, or $G = \mathrm{SO}(N)$, $N \geq 3$, and $|\beta| < \frac{N-2}{58N}$, the conclusions of Theorem 1.1 hold with the rate and gap of Theorem 1.5. For* $\mathrm{SU}(3)$ *this is $\beta_W < \frac{9}{29} = 0.310$.*

The constant $\frac{29}{2}$ is a statement about $\mathrm{U}(N)$, so it holds on every closed subgroup, and the curvature of the group does the rest. This gives the theorem for every compact simple gauge group, the exceptional groups included, with an explicit gap. Let $\rho : G \to \mathrm{U}(N)$ be a faithful unitary representation of a compact connected Lie group G, weigh a configuration by $\exp(N\beta\sum_p \operatorname{Re}\operatorname{Tr}\rho(Q_p))$, and let κ_G be the least Ricci[7] curvature of G for the bi-invariant metric $\langle X, Y\rangle = -\operatorname{Tr}(\rho(X)\rho(Y))$; we identify G with $\rho(G) \subset \mathrm{U}(N)$.

Theorem 1.5 (Every compact simple group)**.** *Suppose that the Lie algebra of G is semisimple, so that $\kappa_G > 0$, and that*

$$N|\beta| < \tfrac{2}{29}\,\kappa_G.$$

Then the conclusions of Theorem 1.1 hold, for $\beta < 0$ all of them except reflection positivity and the gap, with decay at every rate below

$$m_G(\beta) = \operatorname{arccosh}\Big(1 + \frac{1}{12}\Big(\frac{\kappa_G}{N|\beta|} - \frac{29}{2}\Big)\Big) > 0$$

and a transfer-matrix gap $m \geq m_G(\beta)$. For a simple G, $\kappa_G = h^\vee/(2\ell_\rho)$, where $h^\vee$ is the dual Coxeter number and ℓ_ρ the Dynkin index of ρ; Table 1 lists the values.

[7] For a bi-invariant metric on a compact Lie group the Ricci curvature is $-\frac{1}{4}\operatorname{Tr}((\operatorname{ad} X)^2)$ in the direction of X. It is positive for every $X \neq 0$ exactly when the Lie algebra is semisimple, and κ_G is its minimum over unit vectors.

TABLE 1. Least Ricci curvature $\kappa_G = h^\vee/(2\ell_\rho)$ of the compact simple groups in a faithful representation ρ of least dimension N, and the range of Theorem 1.5 for the conventional inverse coupling $\beta_W = N^2\beta$.

Group	ρ	$h^\vee$	ℓ_ρ	κ_G	$\beta_W <$
$\mathrm{SU}(N)$, $N \geq 2$	$\mathbb{C}^N$	N	1	$\frac{N}{2}$	$\frac{N^2}{29}$
$\mathrm{SO}(N)$, $N \geq 5$	$\mathbb{R}^N$	$N-2$	2	$\frac{N-2}{4}$	$\frac{N(N-2)}{58}$
$\mathrm{SO}(3)$	$\mathbb{R}^3$	2	4	$\frac{1}{4}$	$\frac{3}{58}$
$\mathrm{Sp}(n)$, $n \geq 2$	$\mathbb{C}^{2n}$	$n+1$	1	$\frac{n+1}{2}$	$\frac{2n(n+1)}{29}$
G_2	7	4	2	1	$\frac{14}{29}$
F_4	26	9	6	$\frac{3}{4}$	$\frac{39}{29}$
E_6	27	12	6	1	$\frac{54}{29}$
E_7	56	18	12	$\frac{3}{4}$	$\frac{84}{29}$
E_8	248	30	60	$\frac{1}{4}$	$\frac{124}{29}$

For $\mathrm{SU}(N)$, $\kappa_G = \frac{N}{2}$, and for $\mathrm{SO}(N)$, $\kappa_G = \frac{N-2}{4}$, so Theorem 1.5 contains the ranges of Theorem 1.4 and supplies their rate.[8] For small $|\beta|$ the existence of a gap is classical for every compact group, with no explicit range [67, Thm. 3.6]; what is new is an explicit range for every group, past the range $N|\beta| < \kappa_G/16$ that the Hessian bound 16 gives without gauge fixing, and an explicit gap.[9] As $\beta \to 0$ the gap grows like $\log \frac{\kappa_G}{6N|\beta|}$.

No Bakry–Émery argument for the ungauged measure can pass $|\beta| = \frac{1}{32}$ [15, Thm. C]; the gauge fixing changes the measure, not the metric, which is why it can. For SU(2) the curvature range $\beta_W < \frac{4}{29}$ lies inside that of Theorem 1.1, and no curvature argument for the gauge-fixed SU(2) measure passes $\beta_W = \frac{1}{3}$: when one active link e is -1 and all others are 1, the six plaquettes through e have variable -1, and along a geodesic through Q_e the Hessian of $-\frac{\beta_W}{2}\sum_p \mathrm{Re}\,\mathrm{Tr}\,Q_p$ is $-6\beta_W$ times the metric of the unit sphere S^3, whose Ricci curvature is 2, so the Bakry–Émery tensor is negative there once $\beta_W > \frac{1}{3}$. For SU(3) the Dobrushin route goes much further.

Theorem 1.6 (SU(3))**.** *Let $G = \mathrm{SU}(3)$ and $0 \leq \beta_W \leq \frac{3}{4}$. Then the conclusions of Theorem 1.1 hold, with decay at every rate below $-\log(1.33319\,\beta_W)$ and a transfer-matrix gap*

$$m \geq -\log(1.33319\,\beta_W) > 0.$$

The proof of Theorem 1.6 bounds the Kantorovich modulus of the one-link measure $\nu_b(dU) \propto e^{\mathrm{Re}\,\mathrm{Tr}(Ub)}dU$ on SU(3) uniformly over staple sums b. Its analytic part is a lower bound for the spectral gap of ν_b through Bochner's formula with a cross term (Lemma 7.3), a corrector chosen box by box among polynomials of degree two in the matrix entries (Lemma 7.4), the reduction of b to diagonal matrices (Lemma 7.5), the character expansion of the moments of ν_b (Lemma 7.6) and an interpolation inequality on boxes of staple sums (Lemma 7.7). The remaining step evaluates the resulting bound on 2341 boxes with rigorous error bounds; the theorem rests on that computation, which is described in full in Section 7.7 and Appendix E.

[8] The dual Coxeter number is $h^\vee = N$ for $\mathrm{SU}(N)$, $N-2$ for $\mathrm{SO}(N)$, $n+1$ for $\mathrm{Sp}(n)$, $4, 9, 12, 18, 30$ for G_2, F_4, E_6, E_7, E_8, and the Dynkin index of a representation is normalised to one on the defining representation of $\mathrm{SU}(N)$ [52, Ch. 6]. The identity $\kappa_G = h^\vee/(2\ell_\rho)$ is the statement that the Ricci tensor of a compact simple group with a bi-invariant metric is one quarter of the negative Killing form, $\mathrm{Ric}(X,X) = -\frac{1}{4}B(X,X)$ [63], together with $B(X,X) = 2h^\vee\,\mathrm{Tr}(\rho(X)^2)/\ell_\rho$. Lower Ricci bounds of the same kind, for special holonomy and for Riemannian submersions, are studied in [19, 40].

[9] With the computer-assisted bound $h_F \leq 14.420001$ of Theorem L.5 in place of $\frac{29}{2}$, every range of Table 1 grows by the factor $29/28.840002 = 1.0055$. Appendix L derives that bound from a certificate of T. Romanov.




FIGURE 1. The gap $m_G(\beta)$ of Theorem 1.5 as a function of $\beta_W = N^2\beta$ on a logarithmic scale, for nine groups in the representations of Table 1. Each curve ends at the edge $\beta_W = 2N\kappa_G/29$ of the range, marked by a dot, where the gap tends to zero; as $\beta_W \to 0$ it grows like $\log(\kappa_G/(6N|\beta|))$. The curves differ only by the factor $N\kappa_G$, since m_G depends on $N|\beta|/\kappa_G$ alone.

TABLE 2. Explicit ranges of uniqueness, decay and a transfer-matrix gap for four-dimensional lattice Yang–Mills theory. The first three rows hold for the ungauged measure; the last three are the results of this paper.

Method	SU(2), β_W	SU(3), β_W
Curvature, Shen–Zhu–Zhu [74]	$< \frac{1}{12}$	$< \frac{3}{16}$
Curvature, sharp [15]	$< \frac{1}{8}$	$< \frac{9}{32}$
Dobrushin, sharp quarter [15]	$< \frac{2}{9}$	
Curvature, forest gauge	$< \frac{4}{29}$	$< \frac{9}{29}$
Dobrushin, forest gauge	$< \frac{1}{3}$	$\leq \frac{3}{4}$
Plaquette blocks, forest gauge	≤ 0.351	≤ 0.77

The plaquette blocks of Theorem 1.3 work for SU(3) as well. They use the same modulus of the one-link measure, but only through the three links of a block, so they need $\Lambda(2\beta_W)$ small and not $2r\Lambda(r) < 1$: near the end of Theorem 1.6 they give a much larger gap, and they go past it.

Theorem 1.7 (SU(3) with plaquette blocks)**.** *Let $G = \mathrm{SU}(3)$ and $0 \leq \beta_W \leq \frac{77}{100} = 0.77$. Then $\alpha_3(\beta_W) < 1$ for the number of (8.2), with $\alpha_3(0.72) \leq 0.9182$, $\alpha_3(\frac{3}{4}) \leq 0.9653$ and $\alpha_3(0.76) \leq 0.99950$ and $\alpha_3(0.77) \leq 0.99978$. The conclusions of Theorem 1.1 hold with decay at every rate below $-\frac{1}{2}\log\alpha_3(\beta_W)$ and a transfer-matrix gap*

$$m \geq -\tfrac{1}{2}\log\alpha_3(\beta_W) > 0;$$

at $\beta_W = \frac{3}{4}$ this gives $m \geq 0.0177$, against $1.1 \cdot 10^{-4}$ from Theorem 1.6, and at $\beta_W = 0.76$ and 0.77, where Theorem 1.6 no longer applies, $m \geq 2.5 \cdot 10^{-4}$ and $m \geq 1.1 \cdot 10^{-4}$.

All seven theorems are statements at fixed lattice spacing. They do not touch the continuum limit, the crossover to Bałaban's regime, or a gap in physical units, which




FIGURE 2. Ranges of Table 2. The height of each bar is the largest inverse coupling β_W covered by the method. Grey bars are methods for the ungauged measure; blue, red and orange bars use the forest gauge. For SU(2) the red bar ends at the ceiling of Theorem 1.2 and the orange bar, plaquette blocks, passes it (Theorem 1.3); for SU(3) the red bar is Theorem 1.6 and the orange bar, plaquette blocks, gives a larger gap there and passes it (Theorem 1.7).

together are the Clay problem; Section 9 says where each method stops and what remains of the four open problems of [15]. The obstruction proved in [16], that a lattice gap uniform in lattice units forces trivial continuum two-point functions, applies to all of them.

The continuum step. Section 10 goes past that obstruction in one respect, through three implications, each proved.

(I1) A Dobrushin–Shlosman condition on one cube of fixed physical size, uniform along the continuum tuning, gives a mass gap in physical units for every continuum limit (Theorems 10.2 and 10.4), and so does exponential decay with constants of the renormalised correlations of a spanning set of observables (Theorem 10.5); this holds also for observables smoothed by the spatial gradient flow (Proposition 10.6).

(I2) The flowed observables, renormalised by one condition at a single physical separation, have reflection positive continuum limits along subsequences, and such a limit is nontrivial exactly when one ratio of lattice two-point functions stays away from zero (Theorem 10.7); that ratio is bounded above and below by ratios of correlations of one slice field at fixed physical separations (Proposition 10.10).

(I3) Tree bounds on the truncated correlations give all the higher Schwinger functions as reflection positive distributions, imply the ratio bound, and carry the conditional mass gap to every polynomial in the smeared fields (Theorem 10.16); for a smooth packet they give the ratio with an explicit rate and a reference vector of finite energy (Proposition 10.20).

What is open. The inputs of (I1)–(I3), namely the cube condition along the tuning, the slice ratio in four dimensions and the tree bounds, are not proved, and so the nontrivial existence of the continuum limit and the gap in physical units remain open. Figure 3 shows how the proved statements, the implications and the open inputs fit together.

Section 11 measures the open inputs. The cube condition holds at strong coupling, fails at every fixed cube size at weak coupling, forces the gauge-fixed variables to be disordered on the scale of the cube, and cannot be reached from estimates on single lines of links; for SU(N) it needs cubes of side at least of order $(\beta_W/\log\beta_W)^{1/4}$ (Propositions 11.16 and 11.17). For a Gaussian field it holds at a fixed physical scale when the field has a mass and fails exactly, at every size, when it is massless (Proposition 11.2); it also fails for the quadratic approximation of the forest gauge itself (Proposition 11.3), and of every gauge fixed on an invariant forest with infinite trees (Proposition 11.4). The order-two cases of the tree bounds force the lattice gap below a fixed multiple of the lattice spacing, so no strong coupling expansion gives them (Proposition 10.19). The slice ratio holds in two dimensions, where it is the exact solution of lattice gauge theory (Remark 10.13), and fails for a field whose mass is fixed in lattice units; for the free field, conditioning an energy packet on averages over blocks leaves a fixed fraction of its ordinary variance at every lattice spacing, so a reduction to block averages has to be measured in the reflected norm (Proposition 10.14).

Block averages. Section 12 poses the cube condition for block averages in a gauge whose fixed links are trees inside blocks of side two. Its quadratic approximation then holds at every coupling once the cube is large (Theorem 12.8), and integrating out one scale reduces it to a cube condition for the averages, which carry the coarse quadratic form at every scale up to a factor between 1 and 256 that does not depend on the scale, with an exponentially decaying kernel whose constants do not depend on the scale either, and the constrained cube condition holds at every scale with one cube side (Corollary 12.16, Proposition 12.9, Theorems 12.12, 12.14 and 12.18); the averages along the lines of the forest gauge, by contrast, leave one class of lines invisible and fail at every size (Proposition 12.1). For the group-valued links, the straight path between block roots gives a coarse link whose fibres satisfy a coercivity bound for every compact group and every field (Theorem 12.21), but this map does not compose across scales (Proposition 12.24), and for every compact connected group no continuous covariant block map treats the positions of a block alike (Proposition 12.25). A cut-off of Bałaban's averaging operation is continuous, covariant and coercive for every field and agrees with the block averages to first order; it breaks the symmetry only at large fields, as it must, and its constants are those of one step (Theorem 12.27). On fields with values in a maximal torus whose curvature stays small at every scale it composes, with the energy nonincreasing and the thin paths contracted (Theorem 12.29), and for every compact group it composes on fields whose curvatures at the successive scales have a small sum (Theorem 12.33). Under the SU(2) Wilson measure the plaquettes that are not small lie in clusters whose physical size tends to zero in probability, and the cut-off acts only next to them (Theorem 12.57, Corollaries 12.58 and 12.59); such plaquettes are nevertheless present along the one-loop tuning (Proposition 12.60). The cut-off average composes through these clusters, which spread by at most three spacings of the current scale per step, up to a coarse spacing of order $(\beta_W/\log\beta_W)^{1/4}a$, the order of the cube side that the cube condition needs (Theorem 12.62 and Corollary 12.63). Under every exterior, excess action in a box is rare, restricting one link to a fixed set leaves its Dobrushin response bounded, and one-point rarity of a bad set gives approximate uniqueness of the Gibbs measures but no decay (Lemmas 12.72 and 12.73, Proposition 12.74). The curvature of the block averages costs action independent of the scale (Lemma 12.75, Proposition 12.76); the chessboard estimate applies to coarse events and is lost after $O(\log\beta_W)$ scales in the normalisation alone (Proposition 12.77); a right translation of the Haar measures bounds the tails of the force along any fixed direction by the second variation of the action, at every scale and under every exterior (Theorem 12.78), which for the abelian theory makes coarse defects rare at the rate $\beta_W x^2$ at every scale (Corollary 12.79), while for SU(2) that second variation grows like 4^k at scale k, on the gauge orbit of the vacuum, and the same happens when the direction is transported along a forest, which keeps the Haar measures

but leaves the residual gauge freedom of the blocks (Propositions 12.80, 12.81); this locates the estimate that the cube condition still needs. In a box whose boundary field has small plaquettes and either lies in a maximal torus or, for SU(2), has plaquette angles ρ_s with $l^2\rho_s$ below a fixed constant, the fraction of unit cubes with a large plaquette is small with probability exponentially close to one in the volume, uniformly over such boundary fields (Lemmas 12.64 and 12.65 and Proposition 12.66). The second class allows cube sides up to a constant times $(\beta_W/\log\beta_W)^{1/4}$ at the level of Corollary 12.63, which is the scale a_{k_*} up to a constant factor; for larger boxes with general boundary fields this conditional estimate is open (Remark 12.67). On the fibres of the cut-off average, at a background whose plaquettes are at most $\kappa \le 10^{-8}$, the gap $\frac{1}{52}$ of the averages[10] persists up to a factor $1 + O(\kappa)$, and the Hessian of the action along the fibre is at least $\lambda(\frac{1}{26} - c\kappa)$, however far the links are from I (Theorem 12.35). On a chart of the fibre with convex domain the constrained law is log-concave for large λ, coarea factor included, and satisfies a Brascamp–Lieb inequality with constant $60/\lambda$ (Theorem 12.37); on a chart whose domain is a product in the coordinates of the links other than the path links, the covariances of local observables decay exponentially at a rate that depends only on G (Theorem 12.41). Off the chart the fibre has only points with a large plaquette, and under the SU(2) Wilson measure their mass is small on most fibres (Theorem 12.44). Along its private links the cut-off average is a submersion with a constant that depends on nothing (Lemma 12.45), and its law under Haar measure on them has a bounded density, so a Peierls argument inside the fibre bounds that mass on every fibre through a field whose plaquettes near the blocks are small (Proposition 12.48). On every fibre, whatever the boundary links, the same argument concentrates the law within any t of the least action of the fields whose averaging sites are near I (Proposition 12.51); where that action is not small the decay is open (Remark 12.55). Whether some block map is coercive for all fields and composes across the remaining scales is open.

Plan. Section 2 constructs the line field. Section 3 shows that a Dobrushin bound for the gauge-fixed specification gives the full conclusion of Theorem 1.1. Sections 4, 5, 6, 7 and 8 prove Theorems 1.1–1.7, and Section 9 measures what is left. Section 10 proves (I1)–(I3), Section 11 states and measures their open inputs, and Section 12 poses the cube condition for block averages, composes the cut-off average across scales at small fields and locates the large fields under the Wilson measure. Section 13 proves what a labelling of the blocks into good and bad gives for the cube condition, where the natural labels fail, the explicit coupling past which the bad islands of one scale are rarer than their entropy (Theorem 13.21), and what the single-scale and the multiscale parts of (D) still need. The appendices contain the residue table of the line field, the comparison lemma and its version for blocks, the blocks of the local Hessian inequality, the details of the certified computation, the reconstruction of the transfer matrix with the equivalence of a gap and exponential clustering (Appendix F), the facts on spectral measures and on the character expansion that the continuum step uses (Appendices G and H), and a complete proof of the sharp quarter for the single-link law of SU(2) (Appendices I–K), so that the paper does not depend on our earlier one for any step.

2. The line field

Write $e_0, \dots, e_3$ for the unit vectors of $\mathbb{Z}^4$ and (x, μ) for the link from x to $x + e_\mu$. A *direction field* is a map $\sigma : \mathbb{Z}^4 \to \{0, 1, 2, 3\}$; its links are $F = \{(x, \sigma(x)) : x \in \mathbb{Z}^4\}$. At each site put $\pi_x(\mu) = \sigma(x + e_\mu)$. The plaquette at x in the plane $\{\mu, \nu\}$ has the links (x, μ), $(x + e_\mu, \nu)$, $(x + e_\nu, \mu)$, (x, ν), so it contains

$$[\sigma(x) \in \{\mu, \nu\}] + [\pi_x(\mu) = \nu] + [\pi_x(\nu) = \mu] \tag{2.1}$$

links of F.

[10] By the certificate of Proposition 12.5 this gap is at least $\frac{537}{5000} = 0.1074$, within 0.09 per cent of the best constant on $\mathbb{Z}^4$, a floating-point value.




FIGURE 3. The logical structure of the paper. Green: theorems at fixed lattice spacing (P1 the gap for every compact simple group, Theorem 1.5, P2 the plaquette blocks for SU(2) and SU(3), P3 the cube condition at strong coupling). Blue: the proved implications (I1)–(I3) of the continuum step. Red: their inputs, none of which is proved along the continuum tuning. Each arrow joins an open input to the implication that consumes it; no implication feeds its own input.


Lemma 2.1. *Every plaquette of $\mathbb{Z}^4$ contains exactly one link of F if and only if, at every site x, π_x fixes $\sigma(x)$ and permutes the other three directions cyclically.*

Proof. Fix x, let $s = \sigma(x)$ and let R be the set of the other three directions. For a plane $\{s, \nu\}$ the count (2.1) is $1 + [\pi_x(s) = \nu] + [\pi_x(\nu) = s]$; it equals one for all $\nu \in R$ exactly when $\pi_x(s) = s$ and $\pi_x(\nu) \neq s$ for $\nu \in R$, that is, when π_x fixes s and maps R into R. For a plane $\{\mu, \nu\} \subset R$ the count is $[\pi_x(\mu) = \nu] + [\pi_x(\nu) = \mu]$. Draw an arrow $\mu \to \pi_x(\mu)$ for every $\mu \in R$ with $\pi_x(\mu) \in R \setminus \{\mu\}$. Each vertex of R has at most one outgoing arrow, and the condition says that each of the three pairs of R carries exactly one arrow. Three arrows on three vertices with out-degree at most one give out-degree exactly one everywhere, and a tournament on three vertices with all out-degrees one is a directed three-cycle. □

From now on σ is the field of period two given in Table 3 of Appendix A, read from x modulo two. The table lists π_x at each of the sixteen residue classes; in each row π_x fixes $\sigma(x)$ and is a three-cycle on the rest, so every plaquette contains exactly one link of F by Lemma 2.1.[11]

[11]There are exactly eight period-two direction fields with this property, all equivalent to σ under the symmetries of the lattice. A field with one fixed link at every site and one in every plaquette exists only in four dimensions: a link lies in $2(d-1)$ plaquettes and a site carries $d(d-1)/2$ of them, and $2(d-1) = d(d-1)/2$ forces $d = 4$. The statements of this section are also checked in the Lean certificate of the archive.

(a) the line field in the slice $x_3 = 0$



(b) one plaquette in the gauge




FIGURE 4. (a) The line field in the slice $x_3 = 0$ of a box of side three. Thick red segments are the links of F in the directions $0, 1, 2$, with a dot marking the direction of travel; an orange ring marks a site whose line leaves the slice in direction 3. Every line is straight (Lemma 2.2(i)), and the segments that leave the box are its boundary stubs. (b) One plaquette with its fixed link e_1. The weights are those of the local inequality of Lemma 6.2 for the three free links: $\frac{21}{10}$ on the two links adjacent to e_1 and $\frac{5}{4}$ on the opposite one. The two purple corners are the ones that enter the corner form (6.3) once $Y_1 = 0$.

Lemma 2.2. (i) *F is a disjoint union of bi-infinite straight lines, one through every site: every site has exactly one outgoing and one incoming link of F, and both point in the direction $\sigma(x)$.*

(ii) *For every box Λ (a product of four finite intervals, with the links and plaquettes it contains), $F \cap E(\Lambda)$ is a disjoint union of segments, hence a forest.*

(iii) *A link $e = (x, \mu) \notin F$ lies in six plaquettes. Four of them contain a link of F that shares an end point with e, and these lie in the planes $\{\mu, \sigma(x)\}$ and $\{\mu, \pi_x(\mu)\}$. In the other two, which lie in the remaining plane $\{\mu, \rho\}$, the link of F is opposite to e.*

(iv) *Every link outside F shares a plaquette with exactly twelve other links outside F, two in each of its plaquettes.*

Proof. (i) Since π_x fixes $\sigma(x)$, $\sigma(x+e_{\sigma(x)}) = \sigma(x)$. Because σ has period two, $x - e_\mu \equiv x + e_\mu$, so also $\sigma(x - e_{\sigma(x)}) = \sigma(x)$, and the link $(x - e_{\sigma(x)}, \sigma(x))$ is in F. Conversely, if $(x - e_\nu, \nu) \in F$, then $\sigma(x - e_\nu) = \nu$, and straightness at $x - e_\nu$ gives $\sigma(x) = \nu$. So the incoming link is unique and continues the outgoing one.

(ii) A line meets a box in a segment, and distinct lines are disjoint.

(iii) The plaquettes of e are, for each $\nu \neq \mu$, the one at x and the one at $x - e_\nu$ in the plane $\{\mu, \nu\}$. A link of F sharing an end point with e is the outgoing or the incoming link of F at x or at $x + e_\mu$. At x both point in the direction $s = \sigma(x) \neq \mu$, and they lie in the plaquettes at x and at $x - e_s$ in the plane $\{\mu, s\}$. At $x + e_\mu$ both point in the direction $t = \pi_x(\mu)$, which lies in $R \setminus \{\mu\}$ because π_x is a three-cycle on R, and they lie in the plaquettes at x and at $x - e_t$ in the plane $\{\mu, t\}$. These four plaquettes are distinct because $s \neq t$. Each plaquette contains exactly one link of F, so in the two plaquettes of the third plane $\{\mu, \rho\}$ the link of F is a link of direction μ or ρ not touching e; the links of direction ρ in such a plaquette touch e, so it is the opposite link.

(iv) Each of the six plaquettes contains e, one link of F and two other links, and two distinct links share at most one plaquette. □

3. Gauge fixing and the free-boundary state

Let $G \subset \mathrm{U}(N)$ be compact with Haar measure dU, and let Λ be a box. The Wilson measure with free boundary is

$$W_\Lambda(dQ) = \frac{1}{Z_\Lambda} \exp\Big(N\beta \sum_{p \in P(\Lambda)} \operatorname{Re}\operatorname{Tr} Q_p\Big) \prod_{e \in E(\Lambda)} dQ_e.$$

The measure is invariant under the gauge transformations $Q_{(x,y)} \mapsto g_x Q_{(x,y)} g_y^{-1}$ with $g \in G^\Lambda$. Call the links outside F *active* and write A for their set. For a function O of the links, O^F is O with every link of F set to the identity.

Lemma 3.1 (Gauge fixing)**.** *Let μ_Λ^F be the probability measure on $G^{A \cap E(\Lambda)}$ whose density with respect to Haar measure is proportional to $\exp(N\beta \sum_{p \in P(\Lambda)} \operatorname{Re}\operatorname{Tr} Q_p^F)$. For every gauge-invariant O on $G^{E(\Lambda)}$,*

$$W_\Lambda(O) = \mu_\Lambda^F(O^F).$$

Proof. Each segment of $F \cap E(\Lambda)$ is a path $x_0, x_1, \dots, x_k$ along one direction. Given the variables on F, put $g_{x_0} = 1$ and $g_{x_{i+1}} = g_{x_i} Q_{(x_i, x_{i+1})}$ on every segment. The transformed field has $Q_e^g = 1$ on $F \cap E(\Lambda)$, and for fixed values on F the map $Q_A \mapsto Q_A^g$ is a product of left and right translations, which preserves Haar measure. Gauge invariance of the action and of O and the change of variables give the claim. □

No leftover holonomy appears because $F \cap E(\Lambda)$ is a forest. The measures μ_Λ^F are finite-volume versions of the specification γ^F on G^A: for $e \in A$, γ_e^F resamples Q_e from the law with density proportional to $\exp(N\beta \sum_{p \ni e} \operatorname{Re}\operatorname{Tr} Q_p^F)$ given all other active links. No infinite-volume gauge transformation is performed; γ^F is a specification in its own right. Fix a metric δ on G of diameter D that is bi-invariant (for $\mathrm{SU}(2)$ the chordal metric on S^3, $D = 2$; for $\mathrm{SU}(N)$ the Hilbert–Schmidt distance). The Kantorovich coefficient of e' on e is

$$C_{ee'} = \sup \frac{W_1\big(\gamma_e^F(\cdot \mid Q), \gamma_e^F(\cdot \mid Q')\big)}{\delta(Q_{e'}, Q'_{e'})},$$

the supremum over pairs of configurations that differ only at e'. It vanishes unless e and e' share a plaquette.

Proposition 3.2. *Suppose that every row and every column of $(C_{ee'})$ sums to at most $\alpha < 1$. Then:*

(i) *γ^F has exactly one Gibbs measure μ_F;*
(ii) *for every gauge-invariant local O the limit $\omega(O) = \lim W_\Lambda(O)$ exists along every sequence of boxes exhausting $\mathbb{Z}^4$ and equals $\mu_F(O^F)$;*
(iii) *ω is invariant under translations and the hypercubic group;*
(iv) *if $\beta \geq 0$, ω is reflection positive for reflections in hyperplanes through sites and in hyperplanes bisecting links;*
(v) *for gauge-invariant local O, O' with O Lipschitz in each link, $|\operatorname{Cov}_\omega(O, O')| \leq C_{O,O'} e^{-\lambda \operatorname{dist}(O,O')}$ for every $\lambda < -\log\alpha$;*
(vi) *if $\beta \geq 0$, the transfer matrix of ω has spectral gap at least $-\log\alpha$.*

Proof. (i) Gibbs measures exist by compactness, and the comparison lemma of Appendix B with $S = \emptyset$ shows that any two coincide.

(ii) Extend μ_Λ^F by Haar measure on the active links outside Λ, and let S be the set of active links with a plaquette outside $P(\Lambda)$; this includes every active link outside $E(\Lambda)$. For an active link $e \notin S$ all six plaquettes of e lie in $P(\Lambda)$, so the conditional law of Q_e under the extended measure is γ_e^F. Let T be the support of O^F and $\lambda < -\log\alpha$. The comparison lemma with weights $e^{-\lambda \operatorname{dist}(\cdot, T)}$ gives

$$|\mu_\Lambda^F(O^F) - \mu_F(O^F)| \leq L \, \frac{\theta D}{1-\theta} \sum_{z \in S} e^{-\lambda \operatorname{dist}(z,T)},$$

with $\theta = \alpha e^{\lambda} < 1$ and L a Lipschitz constant of O^F; the sum is at most $C_T R^3 e^{-\lambda R}$ when R is the distance from T to the complement of Λ. For O continuous but not Lipschitz, approximate uniformly. Lemma 3.1 gives $W_\Lambda(O) = \mu_\Lambda^F(O^F)$.

(iii) $W_\Lambda(\tau_v O) = W_{\Lambda - v}(O)$, and the boxes $\Lambda - v$ also exhaust $\mathbb{Z}^4$; by (ii) the limit does not depend on the sequence. The line field is a device for computing W_Λ and is never shifted. The same argument applies to the hypercubic group.

(iv) For $\beta \geq 0$ the measure W_Λ is reflection positive for every box symmetric under the reflection [67], and reflection positivity passes to the limit along such boxes, which is ω by (ii).

(v) $\mathrm{Cov}_\omega(O, O') = \mathrm{Cov}_{\mu_F}(O^F, O'^F)$, and setting the links of F to the identity keeps the Lipschitz property. The bound is Corollary B.2.

(vi) Let ϑ be a time reflection and τ_t a time translation. For gauge-invariant local O that are Lipschitz in each link, (v) gives $|\omega(\vartheta(\bar{O})\tau_t O) - |\omega(O)|^2| \leq C_O e^{-\lambda t}$. On the support of O these functions are dense in the gauge-invariant part of $L^2(\omega)$: gauge-invariant functions are functions on the compact quotient of G^E by the gauge group, Lipschitz functions for the quotient metric are dense in L^2 of every finite measure on it, and they lift to gauge-invariant Lipschitz functions. By Proposition F.1 the spectrum of the transfer matrix on the complement of the vacuum lies in $[0, e^{-\lambda}]$, for every $\lambda < -\log \alpha$. □

The proposition replaces the torus limits and the uniqueness of the ungauged Gibbs measure used in [15, Thm. F] by the free-boundary limit, which has all the properties the reconstruction of the transfer matrix needs.[12]

4. SU(2) up to one third

Proof of Theorem 1.1. Identify SU(2) with the unit quaternions and use the chordal metric. For an active link e and a plaquette $p \ni e$, the staple s_p is the product of the other three link variables in the order that makes $\mathrm{Re}(Q_e s_p) = \frac{1}{2}\operatorname{Tr} Q_p$, with links of F equal to one. Since $N\beta \operatorname{Re}\operatorname{Tr} Q_p = 4\beta\langle Q_e, \bar{s}_p\rangle$ in $\mathbb{R}^4$, the conditional law of Q_e is the von Mises–Fisher law $\nu_b(dx) \propto e^{\langle x,b\rangle} dx$ on S^3 with $b = 4\beta \sum_{p \ni e} \bar{s}_p$. Changing a partner e' by δ in the chordal metric changes one staple by an isometric image of that displacement, so b moves by $4|\beta|\delta$, and the sharp quarter (Theorem I.4), $W_1(\nu_b, \nu_{b'}) \leq \frac{1}{4}|b - b'|$, gives $C_{ee'} \leq |\beta|$. By Lemma 2.2(iv) every row and column has twelve nonzero entries, so $\alpha \leq 12|\beta| = 3|\beta_W|$. Proposition 3.2 applies for $|\beta_W| < \frac{1}{3}$. □

Proof of Theorem 1.2. For the line field, every staple of an active link contains two active links, so the six staples can be chosen with sum zero while the staple through a given partner e' moves freely. At $b = 0$ the derivative of ν_b in the direction h has Kantorovich norm $\frac{1}{4}|h|$ by Lemma I.3(i), so $C_{ee'} = |\beta|$ exactly, and every row sum is $12|\beta| = 3|\beta_W|$.

Now let F' be any forest and C' the matrix of the statement. Suppose a positive weight w satisfies $C'w \leq \alpha w$. For a box Λ_n of side n let C_n be the principal submatrix on the active links of $E(\Lambda_n)$; then $C_n w \leq \alpha w$ on these links, and the Collatz–Wielandt inequality for nonnegative matrices gives $\rho(C_n) \leq \alpha$. Since C_n is symmetric, $\rho(C_n)$ is at least the mean row sum. The box has $4n^4 - O(n^3)$ links and n^4 sites, and a forest has fewer links than sites, so the number a of active links is at least $3n^4 - O(n^3)$. Let a_p be the number of active links of the plaquette p; the box has $P = 6n^4 - O(n^3)$ plaquettes and $\sum_p a_p = 6a - O(n^3)$. Two links share at most one plaquette, so the row sums add up to $|\beta| \sum_p a_p(a_p - 1)$, and by convexity

$$\frac{|\beta|}{a} \sum_p a_p(a_p - 1) \geq \frac{|\beta| P}{a}\, \bar{a}(\bar{a} - 1) = 6|\beta|(\bar{a} - 1) - o(1),$$

[12] On a torus the line field closes up into cycles, and the holonomies along them cannot be removed by a gauge transformation; the argument never uses a torus. The state ω is defined only on gauge-invariant observables, which is all that the Osterwalder–Schrader reconstruction [65, 66, 67] uses.

with $\bar a = \sum_p a_p/P = 6a/P - o(1) \ge 3 - o(1)$. Hence $\alpha \ge 12|\beta| - o(1)$ for every n, and $\alpha \ge 12|\beta| = 3|\beta_W|$. □

Without gauge fixing the same count gives eighteen partners and the ceiling $\beta_W = \frac{2}{9}$ of [15, Prop. 13]; the temporal gauge, a family of lines in one direction, leaves fourteen. The line field is optimal among forests: twelve is the mean number of partners forced by the number of free links, and it is attained at every link.

5. SU(2) past one third: plaquette blocks

Throughout this section $G = \mathrm{SU}(2)$ is the group of unit quaternions with the chordal metric $|x - y|$, $\bar x$ is the conjugate, $\langle x, y\rangle = \mathrm{Re}(x\bar y)$ is the inner product of $\mathbb{R}^4$, and dx is the normalised Haar measure on S^3. Left and right multiplication by unit quaternions and conjugation are isometries, and $\mathrm{Re}(xyz) = \mathrm{Re}(yzx) = \langle y, \bar z\bar x\rangle$. Put

$$Z_0(c) = \int_{S^3} e^{\langle c,x\rangle} dx = \sum_{j\ge 0} \frac{(|c|^2/4)^j}{j!\,(j+1)!}, \qquad c \in \mathbb{R}^4,$$

let $\nu_c(dx) \propto e^{\langle c,x\rangle} dx$ be the von Mises–Fisher law, and write $g = \log Z_0$, so that $\nabla g(c)$ is the mean and $\mathrm{Hess}\, g(c)$ the covariance of ν_c.

5.1. **The blocks.** Let $\mathcal{B}$ be the set of plaquettes at x in the plane $\{0,1\}$ with $x_0 \equiv x_1$ and in the plane $\{2,3\}$ with $x_2 \equiv x_3$, congruences modulo two (Table 4 of Appendix C lists them modulo $2\mathbb{Z}^4$, and Figure 5 draws them). A *block plaquette* is a plaquette of $\mathcal{B}$, and its *block* is the set of its three active links.

Lemma 5.1. *The set $\mathcal{B}$ is invariant under translations by $2\mathbb{Z}^4$ and:*

(i) *the blocks partition the active links;*
(ii) *every plaquette $p \notin \mathcal{B}$ meets three distinct blocks, one in each of its active links;*
(iii) *every active link lies in exactly five plaquettes outside $\mathcal{B}$, so it shares a plaquette outside $\mathcal{B}$ with exactly ten active links, none of them in its own block;*
(iv) *if an active link j and a link of a block B lie in a common plaquette outside $\mathcal{B}$, then every link of B has its midpoint within distance 2 of the midpoint of j.*

Proof. Invariance under $2\mathbb{Z}^4$ is clear from the congruences. We use one fact throughout: two distinct links lie in at most one common plaquette, since they span it.

Every link lies in exactly one plaquette of $\mathcal{B}$. Indeed, a link (y, μ) with $\mu \in \{0,1\}$ lies in no plaquette of the plane $\{2,3\}$, and in exactly two of the plane $\{0,1\}$, those at y and at $y - e_\nu$, where ν is the other element of $\{0,1\}$. The base points y and $y - e_\nu$ differ only in the coordinate ν, so $y_0 - y_1$ has opposite parities at the two, and exactly one of them satisfies $x_0 \equiv x_1$. The directions $2, 3$ are treated in the same way. Every plaquette of $\mathbb{Z}^4$ contains exactly one link of F (Lemma 2.1), so a block plaquette contains three active links, and every active link lies in exactly one block. This is (i).

(ii) Let $p \notin \mathcal{B}$ have active links e_1, e_2, e_3, and let q_i be the block plaquette of e_i. Then $q_i \ne p$, so $q_i \cap p = \{e_i\}$ by the fact above. If $q_i = q_k$ with $i \ne k$, then e_i, e_k lie in the two distinct plaquettes p and q_i, which is impossible. So the three blocks are distinct, and each meets p in one link.

(iii) An active link e lies in six plaquettes (Lemma 2.2), and by (i) exactly one of them is in $\mathcal{B}$. Each of the other five contains two further active links. These ten links are distinct, because a link other than e shares at most one plaquette with e, and none lies in the block of e, because a link f of that block sharing the plaquette $p \notin \mathcal{B}$ with e would put e, f in the two distinct plaquettes p and the block plaquette of e.

(iv) Let $f \in B$ share the plaquette $p \notin \mathcal{B}$ with j. Two links of one unit square have midpoints at distance $2^{-1/2}$ or 1, so the midpoint of j is within distance 1 of that of f, and the midpoint of f is within distance 1 of that of every link of its block plaquette. The triangle inequality gives the bound 2. □




FIGURE 5. Block plaquettes, shaded, in (a) a plane $\{0,1\}$ and (b) a plane $\{2,3\}$: the plaquettes at x with $x_0 \equiv x_1$, respectively $x_2 \equiv x_3$, modulo two. Red links belong to the line field F; a filled dot marks a site whose line runs in the plane, an open dot one whose line leaves it. The shaded squares form a checkerboard, so every link of the plane borders exactly one of them, and every square contains exactly one red link (Lemma 2.1); hence each block plaquette carries three active links and the blocks partition the active links (Lemma 5.1(i)).

Fix a block plaquette p with active links e_1, e_2, e_3. Reading the plaquette product cyclically and deleting its fixed link, $\frac{1}{2}\operatorname{Tr} Q_p = \operatorname{Re}(y_1 y_2 y_3)$, where y_i is Q_{e_i} or $Q_{e_i}^{-1}$. Every other plaquette containing e_i meets the block only in e_i by Lemma 5.1(ii) and contributes $4\beta\langle y_i, \bar{s}\rangle$ to the exponent, where s is the staple of y_i in it. Hence the conditional law of the block given all other active links is

$$\mu_b(dy) \propto \exp\Big(\sum_{i=1}^{3}\langle y_i, b_i\rangle + 4\beta \operatorname{Re}(y_1 y_2 y_3)\Big)\, dy_1\, dy_2\, dy_3, \tag{5.1}$$

with b_i the sum of $4\beta\bar{s}$ over the five staples, so that $|b_i| \le 20|\beta|$. Inversion is an isometry of S^3 and preserves Haar measure, so we may work with y instead of Q. On $(S^3)^3$ we use the metric $d(y, y') = \sum_i |y_i - y_i'|$.

5.2. **The sensitivity of one block.** Let λ_* and τ be a lower and an upper bound for the eigenvalues of $\operatorname{Hess} g(c)$ on the ball $|c| \le 24|\beta|$, and M an upper bound there for the norm of the third derivative $D^3 g(c)$ as a trilinear form. Put

$$\begin{aligned} w &= 8\beta^2(\tau - \lambda_*), \qquad s = \tfrac{1}{4} + (e^w - 1)e^w\sqrt{\tau/3}, \\ C &= 4|\beta| s A + \tfrac{320}{3}|\beta|^3 M e^w, \qquad K = \frac{(1 + |\beta|)\, s}{1 - C}, \end{aligned} \tag{5.2}$$

assuming $C < 1$, where A is an upper bound for $|\nabla g(c)|$ on the ball $|c| \le 20|\beta|$.

Lemma 5.2. *Let ν be a probability measure and $\lambda = e^R\nu/\nu(e^R)$ with $\operatorname{osc} R \le \omega$. For bounded f, h,*

$$\operatorname{Cov}_\lambda(f, h) \le \operatorname{Cov}_\nu(f, h) + (e^\omega - 1)e^\omega \operatorname{sd}_\nu(f) \operatorname{sd}_\nu(h).$$

Proof. We may assume $0 \le R \le \omega$. The density $\rho = e^R/\nu(e^R)$ lies in $[e^{-\omega}, e^{\omega}]$, so $\delta = \|\rho - 1\|_\infty \le e^\omega - 1$. With $\tilde f = f - \nu f$ and $\tilde h = h - \nu h$,

$$\begin{aligned}\operatorname{Cov}_\lambda(f,h) - \operatorname{Cov}_\nu(f,h) &= \nu\big((\rho-1)\tilde f\tilde h\big)\\ &\quad - \nu\big((\rho-1)\tilde f\big)\,\nu\big((\rho-1)\tilde h\big),\end{aligned}$$

and the right side is at most $(\delta+\delta^2)\operatorname{sd}_\nu(f)\operatorname{sd}_\nu(h)$ by the Cauchy–Schwarz inequality. □

Lemma 5.3. *Let $\beta \ge 0$, $a, c_0 \in \mathbb{R}^4$ with $|a|, |c_0| \le 20\beta$, and*

$$\kappa(dx) \propto \exp\big(\langle x, a\rangle + g(c_0 + 4\beta\bar x)\big)dx \quad \text{on } S^3.$$

For every 1-Lipschitz f and every $h \in \mathbb{R}^4$:

(i) $\operatorname{Cov}_\kappa(f, \langle x, h\rangle) \le s|h|$;
(ii) *if κ' is the law with c_0' in place of c_0 and $|c_0'| = |c_0| = \rho$, then $|\kappa(f) - \kappa'(f)| \le (8\beta s\varphi'(\rho^2) + \frac{16}{3}\beta^2 M e^w)|c_0 - c_0'|$.*

Proof. (i) Taylor's formula at c_0 gives $g(c_0 + 4\beta u) = g(c_0) + 4\beta\langle\nabla g(c_0), u\rangle + R(u)$ with $R(u) = 8\beta^2 \operatorname{Hess} g(\xi)(u,u)$ for some ξ on the segment, $|\xi| \le 24\beta$. For $u = \bar x$ of norm one, $R(\bar x) \in [8\beta^2\lambda_*, 8\beta^2\tau]$, so $\operatorname{osc} R \le w$. Since $\langle\nabla g(c_0), \bar x\rangle = \langle x, \overline{\nabla g(c_0)}\rangle$, κ is the tilt $e^R\nu/\nu(e^R)$ of $\nu = \nu_c$ with $c = a + 4\beta\overline{\nabla g(c_0)}$. The sharp quarter, Theorem I.4, gives $\operatorname{Cov}_\nu(f, \langle x, h\rangle) = \frac{d}{dt}\nu_{c+th}(f) \le \frac14|h|$. A function that is 1-Lipschitz for the chordal metric is 1-Lipschitz for the geodesic one, so $|\nabla f| \le 1$ almost everywhere and the Poincaré inequality of Lemma I.1 gives $\operatorname{sd}_\nu(f) \le 1/\sqrt3$; and $\operatorname{sd}_\nu\langle x, h\rangle \le \sqrt\tau|h|$ because $|c| \le |a| + 4\beta \le 24\beta$. Lemma 5.2 gives (i). The same argument applies to every law $e^{R'}\nu_{c'}/\nu_{c'}(e^{R'})$ with $|c'| \le 24\beta$ and $\operatorname{osc} R' \le w$.

(ii) We interpolate the exponents rather than the field. For $t \in [0,1]$ let $\kappa_t \propto \exp\big(\langle x, a\rangle + (1-t)g(c_0 + 4\beta\bar x) + tg(c_0' + 4\beta\bar x)\big)dx$, so that $\kappa_0 = \kappa$, $\kappa_1 = \kappa'$ and $\frac{d}{dt}\kappa_t(f) = \operatorname{Cov}_{\kappa_t}(f, D)$ with $D(x) = g(c_0' + 4\beta\bar x) - g(c_0 + 4\beta\bar x)$. Write $g(c + 4\beta u) = g(c) + 4\beta\langle\nabla g(c), u\rangle + R_c(u)$ as in (i), for $c \in \{c_0, c_0'\}$. Then κ_t is the tilt of ν_{c_t}, $c_t = a + 4\beta\,\overline{(1-t)\nabla g(c_0) + t\nabla g(c_0')}$, by $R_t = (1-t)R_{c_0} + tR_{c_0'}$, with $\operatorname{osc} R_t \le w$ and $|c_t| \le 24\beta$ because $|\nabla g| \le 1$; so (i) holds for κ_t, and the density of κ_t with respect to ν_{c_t} is at most e^w. Up to a constant, $D(x) = \langle x, h\rangle + N(\bar x)$ with $h = 4\beta\,\overline{\nabla g(c_0') - \nabla g(c_0)}$ and $N = R_{c_0'} - R_{c_0}$. Since $\nabla g(c) = 2\varphi'(|c|^2)c$ and $|c_0| = |c_0'| = \rho$, $|h| = 8\beta\varphi'(\rho^2)|c_0 - c_0'|$, and by (i) the linear part contributes at most $s|h|$. For the rest, $\nabla N(u) = 16\beta^2\int_0^1\big(\operatorname{Hess} g(c_0' + 4\beta\sigma u) - \operatorname{Hess} g(c_0 + 4\beta\sigma u)\big)u\,d\sigma$, both arguments in the convex ball $|c| \le 24\beta$, so N is L-Lipschitz on the unit ball with $L = 16\beta^2 M|c_0 - c_0'|$. With $\nu = \nu_{c_t}$ and the Poincaré inequality of Lemma I.1 again, $\operatorname{Var}_{\kappa_t}(N) \le \kappa_t((N - \nu N)^2) \le e^w \operatorname{Var}_\nu(N) \le e^w L^2/3$, and in the same way $\operatorname{Var}_{\kappa_t}(f) \le e^w/3$. So $\operatorname{Cov}_{\kappa_t}(f, N) \le e^w L/3$. Integrating over t gives (ii). □

Lemma 5.4. *Let $C < 1$. If b and b' satisfy $|b_i|, |b_i'| \le 20|\beta|$ and differ only in one coordinate b_i, then*

$$W_1(\mu_b, \mu_{b'}) \le K\,|b_i - b_i'|$$

in the metric d.

Proof. The map $y_1 \mapsto -y_1$ turns μ_b for β into the law for $-\beta$ with b_1 replaced by $-b_1$ and is an isometry, so we may take $\beta \ge 0$; the cyclic symmetry of $\operatorname{Re}(y_1y_2y_3)$ lets us take $i = 1$. Let f be 1-Lipschitz for d and $\varepsilon = |b_1 - b_1'|$.

The middle link. Given y_1, y_3, the law of y_2 is ν_c with $c = b_2 + 4\beta\,\overline{y_3y_1}$, which does not involve b_1. Changing y_1 by η moves c by $4\beta\eta$, and the sharp quarter moves ν_c by at most $\beta\eta$ in W_1. Hence $F(y_1, y_3) = \mathbb{E}[f \mid y_1, y_3]$ is $(1+\beta)$-Lipschitz in each variable, and $\mu_b(f) = \mu_b^{13}(F)$, where μ_b^{13} is the law of (y_1, y_3):

$$\mu_b^{13} \propto e^{\langle y_1, b_1\rangle + \langle y_3, b_3\rangle}\,Z_0\big(b_2 + 4\beta\,\overline{y_3y_1}\big).$$

Since $|b_2 + 4\beta\overline{y_3 y_1}| = |b_2 y_3 + 4\beta\bar{y}_1| = |y_1 b_2 + 4\beta\bar{y}_3|$, its conditional laws are

$$\kappa_1(dy_1 \mid y_3) \propto e^{\langle y_1, b_1\rangle + g(b_2 y_3 + 4\beta\bar{y}_1)} dy_1,$$
$$\kappa_3(dy_3 \mid y_1) \propto e^{\langle y_3, b_3\rangle + g(y_1 b_2 + 4\beta\bar{y}_3)} dy_3,$$

both of the form of Lemma 5.3, with $|c_0| = |b_2|$.

Two sites. Integrating Lemma 5.3(i) along the segment from b_1 to b_1' gives $W_1(\kappa_1, \kappa_1') \le s\varepsilon$ for the same y_3. Moving y_3 to y_3' replaces $c_0 = b_2 y_3$ by $c_0' = b_2 y_3'$, of the same norm $\rho = |b_2| \le 20\beta$ and with $|c_0 - c_0'| = \rho|y_3 - y_3'|$. Since $\rho \cdot 2\varphi'(\rho^2) = |\nabla g(c_0)| \le A$, Lemma 5.3(ii) gives $W_1(\kappa_1(\cdot \mid y_3), \kappa_1(\cdot \mid y_3')) \le (4\beta s A + 20\beta \cdot \frac{16}{3}\beta^2 M e^w)|y_3 - y_3'| = C|y_3 - y_3'|$, and the same for κ_3, which does not depend on b_1. Let π be a coupling of μ_b^{13} and $\mu_{b'}^{13}$ minimising $d_1 + d_3$, $d_k = \mathbb{E}_\pi|y_k - y_k'|$. Replacing (y_1, y_1') by an optimal coupling of $\kappa_1(\cdot \mid y_3)$ and $\kappa_1'(\cdot \mid y_3')$ keeps a coupling, so minimality gives $d_1 \le s\varepsilon + C d_3$; in the same way $d_3 \le C d_1$. Hence $d_1 \le s\varepsilon/(1 - C^2)$ and

$$|\mu_b(f) - \mu_{b'}(f)| \le (1 + \beta)(d_1 + d_3) \le \frac{(1 + \beta)s}{1 - C}\,\varepsilon = K\varepsilon. \qquad \square$$

5.3. **The constants.**

Lemma 5.5. *Write $g(c) = \varphi(|c|^2)$ and $r = |c|$. For $c \neq 0$ the eigenvalues of* Hess $g(c)$ *are $2\varphi'(r^2)$, three times, and $2\varphi'(r^2) + 4r^2\varphi''(r^2)$, and the norm of $D^3 g(c)$ is*

$$\max\Big(|A + \Gamma|, \tfrac{2}{3}|A|\Big), \qquad A = 12r\varphi''(r^2), \quad \Gamma = 8r^3\varphi'''(r^2).$$

On the ball $r \le 2.106$ every eigenvalue of Hess g *lies between* 0.156381 *and* 0.25000001*, and the norm of $D^3 g$ is at most* 0.0615172*; on the ball $r \le 1.755$, $|\nabla g| \le 0.391443$.*

Proof. Differentiating $\varphi(|c|^2)$ gives $D^2 g(v, v) = 2\varphi'|v|^2 + 4\varphi''\langle c, v\rangle^2$ and $D^3 g(v, v, v) = 12\varphi''\langle c, v\rangle|v|^2 + 8\varphi'''\langle c, v\rangle^3$. For $|v| = 1$ and $t = \langle c/r, v\rangle \in [-1, 1]$ the cubic form is $At + \Gamma t^3$, whose largest absolute value is attained at $t = \pm 1$ or at an interior critical point, where it equals $\frac{2}{3}|A||t| \le \frac{2}{3}|A|$; by Banach's theorem [9] the supremum of a symmetric trilinear form on the diagonal of the unit sphere is its norm. The numerical bounds are obtained in ball arithmetic [51] from the series of Z_0, truncated after forty-eight terms with a proved bound on the tail, on $2 \cdot 10^5$ subintervals of $[0, 2.106]$, with $|\nabla g(c)| = r \cdot 2\varphi'(r^2)$; the program is described in Appendix C. $\square$

5.4. **The proof of Theorem 1.3.** Let $\gamma^{\mathcal{B}}$ be the block specification: for a block B, $\gamma_B^{\mathcal{B}}$ resamples the three links of B from (5.1). It comes from the same interaction as γ^F, so the two specifications have the same Gibbs measures. Define the block coefficients

$$C_{Bj} = \sup \frac{W_1(\gamma_B^{\mathcal{B}}(\cdot \mid Q), \gamma_B^{\mathcal{B}}(\cdot \mid Q'))}{|Q_j - Q_j'|}$$

for an active link $j \notin B$, the supremum over pairs of configurations that differ only at j, with W_1 for the metric d.

Proposition 5.6. *Suppose that $\sum_B C_{Bj} \le \alpha < 1$ for every active link j. Then the conclusions (i)–(vi) of Proposition 3.2 hold with $-\frac{1}{2}\log\alpha$ in place of $-\log\alpha$.*

Proof. The proof of Proposition 3.2 goes through with Lemma B.3 of Appendix B in place of Lemma B.1, and with the set S of part (ii), and the support of H in Corollary B.2, enlarged to the union of the blocks that meet it. The enlargement multiplies the size of these sets by at most three and moves them by at most one, which changes only the constants. $\square$

Proof of Theorem 1.3. Let $|\beta| \le 0.351/4$ and take the constants of Lemma 5.5, whose balls $r \le 2.106$ and $r \le 1.755$ contain $|c| \le 24|\beta|$ and $|c| \le 20|\beta|$. With these fixed constants every quantity in (5.2) is increasing in $|\beta|$, and so is

$$\alpha(\beta_W) = 40|\beta|K. \tag{5.3}$$

Changing an active link j by η changes one staple of each of its ten partners outside its block by an isometric image of the displacement, so the field b_i of each such partner i moves by at most $4|\beta|\eta$, and Lemma 5.4, applied once for each of these ten fields, gives $\sum_B C_{Bj} \le 40|\beta|K$. Evaluating (5.2) in ball arithmetic at $\beta_W = 0.351$ gives $w \le 0.0057670$, $s \le 0.2516793$, $C \le 0.0390392$, $K \le 0.2848858$ and $\alpha \le 0.99995$; at $\beta_W = 0.345$ and $\beta_W = \frac{1}{3}$ it gives $\alpha \le 0.98044$ and $\alpha \le 0.94279$. At $\beta_W = 0.3511$ the same evaluation exceeds one. Proposition 5.6 applies. □

Remark 5.7. Most of the range is kept. A second bound for the same sensitivity, which keeps the law of the middle link instead of linearising it, evaluated in floating point on a quadrature grid and maximised over the fields b by a random search followed by the Nelder–Mead method, gives 0.2783 at $\beta_W = 0.34$, against 0.2837 from Lemma 5.4, and $40|\beta|$ times it crosses one near $\beta_W = 0.357$. Most of the difference is the two-site constant $C \approx 0.038$, which the floating-point value suggests is about twice the true feedback between y_1 and y_3. No bound of this form passes $\beta_W \approx 0.3666$: at $b = 0$ the function $f = \langle y_1, e\rangle + \mathrm{Re}(y_2 y_3 e)$ is 1-Lipschitz for d and has $\mathrm{Cov}_{\mu_b}(f, \langle y_1, e\rangle) = \frac{1}{4}(1 + I_2(\beta_W)/I_1(\beta_W))$ exactly (Proposition 13.15), which is 0.2718 at $\beta_W = 0.351$ against $K \le 0.2849$, so the sensitivity is at least this number and $40|\beta|$ times it crosses one at $\beta_W = 0.3666$. Past this point the gain from integrating out one link is spent, and larger blocks are needed, such as the free links of a cube of the line field, whose sensitivity no longer reduces to two sites.

6. Curvature in the gauge

In this section $G \subset \mathrm{U}(N)$ is a closed connected subgroup with Lie algebra $\mathfrak{g}$ and the inner product $\langle X, Y\rangle = -\operatorname{Tr}(XY)$; the bound of Theorem 1.4 is proved first for $\mathrm{U}(N)$. A tangent vector at a field Q is $v = (X_e Q_e)$, $|v|^2 = \sum_e |X_e|^2$. For the plaquette at x in the plane $\mu < \nu$ number its links $e_1 = (x, \mu)$, $e_2 = (x + e_\mu, \nu)$, $e_3 = (x + e_\nu, \mu)$, $e_4 = (x, \nu)$, transport their vectors to x,

$$Y_1 = X_{e_1}, \quad Y_2 = Q_{e_1} X_{e_2} Q_{e_1}^*, \quad Y_3 = Q_{e_4} X_{e_3} Q_{e_4}^*, \quad Y_4 = X_{e_4}, \tag{6.1}$$

and write $W = Q_p$. The second derivative of $\operatorname{Re}\operatorname{Tr} Q_p$ along the geodesic $Q_e(t) = e^{tX_e} Q_e$ is

$$\begin{aligned} f_p = {} & \sum_{i=1}^{4} \operatorname{Re}\operatorname{Tr}(Y_i^2 W) + 2\operatorname{Re}\operatorname{Tr}(Y_1 Y_2 W) + 2\operatorname{Re}\operatorname{Tr}(Y_3 Y_4 W) \\ & - 2 \sum_{i \in \{1,2\},\, j \in \{3,4\}} \operatorname{Re}\operatorname{Tr}(Y_i W Y_j), \end{aligned} \tag{6.2}$$

so that $\operatorname{Hess} S_0(v, v) = \sum_p f_p$. To see this, write $Q_p = Q_{e_1} Q_{e_2} Q_{e_3}^* Q_{e_4}^*$. Since $Q_{e_1} e^{tX_{e_2}} = e^{tY_2} Q_{e_1}$ and $e^{-tX_{e_3}} Q_{e_4}^* = Q_{e_4}^* e^{-tY_3}$, the plaquette variable along the geodesic is $Q_p(t) = e^{tY_1} e^{tY_2} W e^{-tY_3} e^{-tY_4}$. Differentiating twice at $t = 0$ gives the four squares, the two products of neighbouring factors and the four products across W, and the squares and the product WY_3Y_4 are moved cyclically under the trace. The corner form of p is

$$c_p = 2\langle Y_1, Y_4\rangle - 2\langle Y_1, Y_2\rangle + 2\langle Y_2, WY_3W^*\rangle - 2\langle Y_3, Y_4\rangle. \tag{6.3}$$

It collects the inner products of the transported vectors of two links that meet at a right angle at a corner of p.

Lemma 6.1. *On $\mathbb{Z}^4$, or on a box with free boundary, for every field Q and every square-summable X,*

$$\begin{aligned} \sum_y \Big[|D_y X|^2 + \sum_\mu \big| X_{y,\mu} + Q^*_{y-e_\mu,\mu} X_{y-e_\mu,\mu} Q_{y-e_\mu,\mu} \big|^2 \Big] \\ = 4|X|^2 + \sum_p c_p, \end{aligned}$$

where $D_yX = \sum_\mu [X_{y,\mu} - Q^*_{y-e_\mu,\mu} X_{y-e_\mu,\mu} Q_{y-e_\mu,\mu}]$ *is the covariant divergence and terms with a link outside the box are omitted.*

Proof. At a site y write o_μ for the vector of the outgoing link in direction μ and i_μ for the transported vector of the incoming one, so that $D_yX = \sum_\mu (o_\mu - i_\mu)$. Every link is outgoing at one end and incoming at the other, so the squares $|o_\mu|^2 + |i_\mu|^2$, which appear once in $|D_yX|^2$ and once in $|o_\mu + i_\mu|^2$, add up to $4|X|^2$. The collinear cross term $-2\langle o_\mu, i_\mu\rangle$ of $|D_yX|^2$ cancels the term $+2\langle o_\mu, i_\mu\rangle$ of $|o_\mu + i_\mu|^2$. The remaining cross terms of $|D_yX|^2$ pair two perpendicular ends at y, and each such pair is a corner of exactly one plaquette, with the sign and the transport of (6.3) (this is the proof of [15, Lemma 1]). In a box two perpendicular ends at y span a plaquette of the box, because a box is a product of intervals. □

Lemma 6.2. *Let* $k \in \{1,2,3,4\}$*, let* $k \pm 1$ *(mod 4) be the positions adjacent to* k *and* $k+2$ *the opposite one. For every* N*, every* $W \in \mathrm{U}(N)$ *and all* $Y_1, \dots, Y_4 \in \mathfrak{u}(N)$ *with* $Y_k = 0$*,*

$$\pm f_p + \tfrac{9}{10} c_p \le \tfrac{21}{10}(|Y_{k-1}|^2 + |Y_{k+1}|^2) + \tfrac{5}{4}|Y_{k+2}|^2.$$

Proof. Replacing W by $-W$ changes the sign of f_p and fixes c_p, so the sign $+$ suffices. Conjugation by a unitary matrix preserves both sides, so we may take $W = \mathrm{diag}(w_a)$. The form $\mathcal{F} = \mathrm{RHS} - f_p - \frac{9}{10} c_p$ then splits into a block for each diagonal position a, in the real variables $y_i = -i(Y_i)_{aa}$, and a block for each pair $a \neq b$, in the complex variables $z_i = (Y_i)_{ab}$, since $(Y_i)_{ba} = -\bar{z}_i$. Appendix D shows that every block is positive definite. □

Proof of the bound in Theorem 1.4. Every plaquette has exactly one link of F, and X vanishes on it. By Lemma 2.2(iii) each active link is adjacent to the fixed link in four of its plaquettes and opposite to it in two. Summing Lemma 6.2 over the plaquettes,

$$\sum_p (\pm f_p + \tfrac{9}{10} c_p) \le (4 \cdot \tfrac{21}{10} + 2 \cdot \tfrac{5}{4})|X|^2 = \tfrac{109}{10}|X|^2.$$

By Lemma 6.1, $\frac{9}{10}\sum_p c_p = \frac{9}{10}\Sigma - \frac{36}{10}|X|^2$, where $\Sigma \ge 0$ is the left side of the lemma. Hence

$$\pm \operatorname{Hess} S_0(v,v) + \tfrac{9}{10}\Sigma \le (\tfrac{109}{10} + \tfrac{36}{10})|X|^2 = \tfrac{29}{2}|X|^2.$$

In a box, links near the boundary lie in fewer plaquettes and receive less. □

The factor $\frac{9}{10}$ is chosen so that the corners are paid by the divergence identity at the price $\frac{36}{10}$ while the local blocks stay positive; Lemma 6.1 is the divergence identity of [15, Lemma 1] with the collinear term written as a square, so nothing is lost by Cauchy–Schwarz.

The two lemmas concern vectors in $\mathfrak{u}(N)$ and fields in $\mathrm{U}(N)$, so the bound holds without change on every closed subgroup $G \subset \mathrm{U}(N)$, the vectors X_e ranging over its Lie algebra $\mathfrak{g}$. It is also the Hessian of S_0 on G^E: the metric is bi-invariant, the geodesics of G through Q_e are the curves $e^{tX}Q_e$ with $X \in \mathfrak{g}$, which are geodesics of $\mathrm{U}(N)$, and the Hessian is read off from second derivatives along geodesics.

Lemma 6.3. *Let* $G \subset \mathrm{U}(N)$ *be closed and connected with simple Lie algebra* $\mathfrak{g}$*, let* $\mathfrak{t} \subset \mathfrak{g}$ *be a maximal abelian subalgebra, and write the eigenvalues of* $\operatorname{ad} H$ *on* $\mathfrak{g} \otimes \mathbb{C}$ *and of* H *on* $\mathbb{C}^N$*, for* $H \in \mathfrak{t}$*, as* $i\alpha(H)$ *and* $i\mu(H)$*, with* α *running over the roots and* μ *over the weights of* $\mathbb{C}^N$ *counted with multiplicity. Then* $\mathrm{Ric}_G = \kappa_G \langle \cdot, \cdot \rangle$ *with*

$$\kappa_G = \frac{\sum_\alpha \alpha(H)^2}{4 \sum_\mu \mu(H)^2} \qquad (0 \neq H \in \mathfrak{t}).$$

Proof. For a bi-invariant metric the curvature is $R(X,Y)Z = -\frac{1}{4}[[X,Y],Z]$ on left-invariant fields [63, Sect. 7], so for an orthonormal basis (E_i) of $\mathfrak{g}$,

$$\begin{aligned} \mathrm{Ric}(X,X) &= \sum_i \langle R(E_i,X)X, E_i\rangle \\ &= \tfrac{1}{4}\sum_i |[X,E_i]|^2 = -\tfrac{1}{4}\operatorname{Tr}(\operatorname{ad} X)^2, \end{aligned}$$

since $\operatorname{ad} X$ is skew. The Killing form $B(X,Y) = \operatorname{Tr}(\operatorname{ad} X \operatorname{ad} Y)$ and $\langle \cdot,\cdot\rangle$ are invariant under the adjoint action, and $\mathfrak{g}$ is simple, so $B = -4\kappa\langle\cdot,\cdot\rangle$ for a constant κ and $\mathrm{Ric} = \kappa\langle\cdot,\cdot\rangle$. On $H \in \mathfrak{t}$ the two forms are $B(H,H) = -\sum_\alpha \alpha(H)^2$ and $\langle H,H\rangle = -\operatorname{Tr} H^2 = \sum_\mu \mu(H)^2$. □

Table 1 evaluates the ratio of Lemma 6.3 for every compact simple Lie algebra and a faithful representation of least dimension, on the root systems of [49, Sect. 12.1]; the weights of the representations of dimensions 27 and 56 of E_6 and E_7 are the roots of E_8 with prescribed inner products with a subsystem A_2, respectively A_1, projected to its orthogonal complement, by the decompositions $248 = (78,1) + (27,3) + (\overline{27},\bar 3) + (1,8)$ and $248 = (133,1) + (56,2) + (1,3)$ [75]. Taking for H the coroot $\theta^\vee$ of the highest root gives $\sum_\alpha \alpha(\theta^\vee)^2 = 4h^\vee$ and $\sum_\mu \mu(\theta^\vee)^2 = 2\ell_\rho$, so $\kappa_G = h^\vee/(2\ell_\rho)$, with the Dynkin index normalised to one on the defining representation of $\mathrm{SU}(N)$. The ratio depends on $\mathfrak{g}$ and ρ only, so it is the same for every compact group with Lie algebra $\mathfrak{g}$, simply connected or not.

Lemma 6.4. *Let $e \neq e'$ be two links of a plaquette p, and write*

$$\operatorname{Hess}\operatorname{Re}\operatorname{Tr} Q_p(v,v) = \sum_{e,e'\in p} H^p_{ee'}(X_e, X_{e'})$$

with bilinear forms satisfying $H^p_{ee'}(a,b) = H^p_{e'e}(b,a)$. Then $|H^p_{ee'}(a,b)| \le |a|\,|b|$ for all $a,b \in \mathfrak{u}(N)$ and every field.

Proof. By (6.2) the terms of f_p that mix two positions $i \neq j$ are $2\operatorname{Re}\operatorname{Tr}(Y_1Y_2W)$, $2\operatorname{Re}\operatorname{Tr}(Y_3Y_4W)$ and $-2\operatorname{Re}\operatorname{Tr}(Y_iWY_j)$, so $H^p_{ee'}$ is $\pm\operatorname{Re}\operatorname{Tr}(abW)$ or $\pm\operatorname{Re}\operatorname{Tr}(aWb)$ after an isometric transport of a and b, and $|\operatorname{Tr}(abW)| \le \|a\|_{\mathrm{HS}}\|bW\|_{\mathrm{HS}} = |a|\,|b|$. □

Corollary 6.5. *Let $G \subset \mathrm{U}(N)$ be closed and connected with semisimple Lie algebra and $N|\beta| < \frac{2}{29}\kappa_G$. For every box the measure μ^F_Λ satisfies the Bakry–Émery condition with $K = \kappa_G - \frac{29}{2}N|\beta| > 0$, uniformly in the box, and hence a logarithmic Sobolev inequality with constant $2/K$ and a Poincaré inequality with constant $1/K$.*

Proof. The set $\{Q_F = 1\}$ is a product factor, so its Ricci curvature is at least κ_G on every link, and the Hessian of the restricted action is the restriction of $N\beta\operatorname{Hess} S_0$ to $X_F = 0$. Apply the bound and [5, Cor. 5.7.2]. □

6.1. **Decay with an explicit rate.** Let A' be a finite set of active links and P' a finite set of plaquettes whose active links lie in A'. An action $S = \sum_{p\in P'} c_p \operatorname{Re}\operatorname{Tr} Q^F_p$ with $0 \le c_p \le 1$ is *admissible* if $|\operatorname{Hess} S(v,v)| \le \frac{29}{2}|v|^2$ for all v tangent to $G^{A'}$. Two links lie in at most one common plaquette, so by Lemma 6.4 and Lemma 2.2(iv) the mixed blocks of its Hessian satisfy $|\operatorname{Hess} S_{ee'}(a,b)| \le |a|\,|b|$, and each link has at most twelve partners. The action $S_\Lambda = \sum_{p\in P(\Lambda)} \operatorname{Re}\operatorname{Tr} Q^F_p$ of a box is admissible by the bound proved above, and so is every convex combination of two admissible actions. Let μ be the probability measure on $G^{A'}$ with density proportional to $e^{N\beta S}$ with respect to Haar measure, let $L = \Delta + N\beta\langle\nabla S,\nabla\cdot\rangle$ be its Langevin generator, which is self-adjoint in $L^2(\mu)$, and $P_t = e^{tL}$. Write ∇_e for the gradient in the factor of the link e, and for weights $w : A' \to (0,\infty)$ put $\Gamma^w(f) = \sum_e w_e|\nabla_e f|^2$; Γ is the case $w \equiv 1$.

Lemma 6.6. *Let $\lambda \ge 0$ and $w_e = e^{\lambda d_e}$, where $|d_e - d_{e'}| \le 1$ whenever e and e' share a plaquette, and put*

$$K_\lambda = \kappa_G - N|\beta|\Big(\tfrac{29}{2} + 12(\cosh\tfrac{\lambda}{2} - 1)\Big).$$

Then $\Gamma^w(P_t f) \le e^{-2K_\lambda t} P_t \Gamma^w(f)$ for every smooth f and $t \ge 0$.

Proof. Put $V = -N\beta S$. On the product manifold $M = G^{A'}$ the projection to the tangent space of one factor is parallel, the rough Laplacian commutes with it, and the Ricci tensor is

block diagonal. Hence the Bochner–Weitzenböck formula $\nabla\Delta g = \bar{\Delta}\nabla g + \mathrm{Ric}(\nabla g)$, applied factor by factor, together with the derivative of the drift, gives for smooth g

$$\begin{aligned}\tfrac{1}{2}L\Gamma^w(g) - \Gamma^w(g, Lg) &= \sum_e w_e\big(|\nabla\nabla_e g|^2 + \mathrm{Ric}_G(\nabla_e g, \nabla_e g)\big)\\ &\quad + \sum_{e,e'} w_e \,\mathrm{Hess}\, V_{ee'}(\nabla_e g, \nabla_{e'} g),\end{aligned}$$

where $\Gamma^w(g,h)$ is the polarised form.[13] Put $u_e = \nabla_e g$ and $v_e = \sqrt{w_e}\,u_e$. By the symmetry of $\mathrm{Hess}\, V$,

$$\begin{aligned}\sum_{e,e'} w_e \,\mathrm{Hess}\, V_{ee'}(u_e, u_{e'}) &= \mathrm{Hess}\, V(v,v)\\ &\quad + \sum_{e\neq e'} \Big(\cosh\tfrac{\lambda(d_e - d_{e'})}{2} - 1\Big)\,\mathrm{Hess}\, V_{ee'}(v_e, v_{e'}),\end{aligned}$$

because $\frac{w_e + w_{e'}}{2\sqrt{w_e w_{e'}}} = \cosh\frac{\lambda(d_e - d_{e'})}{2}$. The first term is at least $-\frac{29}{2}N|\beta|\,|v|^2$. In the second only partners contribute, each with a coefficient at most $\cosh\frac{\lambda}{2} - 1$ and a block of norm at most $N|\beta|$, and $2|v_e|\,|v_{e'}| \le |v_e|^2 + |v_{e'}|^2$ with twelve partners per link bounds it by $12(\cosh\frac{\lambda}{2} - 1)N|\beta|\,|v|^2$. Since $\mathrm{Ric}_G \ge \kappa_G$ and $|v|^2 = \Gamma^w(g)$,

$$\tfrac{1}{2}L\Gamma^w(g) - \Gamma^w(g, Lg) \ge K_\lambda \Gamma^w(g).$$

For fixed t let $\varphi(s) = P_s\Gamma^w(P_{t-s}f)$, $0 \le s \le t$. Then $\varphi'(s) = P_s\big(L\Gamma^w(g_s) - 2\Gamma^w(g_s, Lg_s)\big) \ge 2K_\lambda\varphi(s)$ with $g_s = P_{t-s}f$, and $\varphi(0) \le e^{-2K_\lambda t}\varphi(t)$ is the claim. □

Lemma 6.7. *Suppose $K_\lambda > 0$. Let f and g be smooth functions on $G^{A'}$ that depend only on the links of A_f and of B_g, and let D be the least number of steps from A_f to B_g in the graph on A' whose edges join links that share a plaquette. Then*

$$|\,\mathrm{Cov}_\mu(f,g)| \le \frac{e^{-\lambda D/2}}{K_\lambda}\,\|\nabla f\|_\infty \sum_{e\in B_g} \|\nabla_e g\|_\infty,$$

where $\|\nabla f\|_\infty^2 = \sup\sum_e |\nabla_e f|^2$.

Proof. Since $K_0 \ge K_\lambda > 0$, Lemma 6.6 with $\lambda = 0$ gives $\Gamma(P_t f) \le e^{-2K_0 t}\|\nabla f\|_\infty^2$, so $P_t f$ converges uniformly to a constant, which is $\mu(f)$ by invariance, M being connected. Since $\frac{d}{dt}\mu(gP_t f) = \mu(gLP_t f) = -\mu(\Gamma(g, P_t f))$,

$$\mathrm{Cov}_\mu(f,g) = \int_0^\infty \mu(\Gamma(P_t f, g))\,dt.$$

Let d_e be the smaller of D and the graph distance from e to A_f, so that $|d_e - d_{e'}| \le 1$ for partners, $w_e = 1$ on A_f, $w_e = e^{\lambda D}$ on B_g, and $\Gamma^w(f) \le \|\nabla f\|_\infty^2$. For $e \in B_g$, Lemma 6.6 gives $|\nabla_e P_t f|^2 \le w_e^{-1}\Gamma^w(P_t f) \le e^{-\lambda D}e^{-2K_\lambda t}\|\nabla f\|_\infty^2$. Hence $|\Gamma(P_t f, g)| \le e^{-\lambda D/2}e^{-K_\lambda t}\|\nabla f\|_\infty \sum_{e\in B_g}\|\nabla_e g\|_\infty$, and integration in t gives the bound. □

Proof of Theorem 1.5. Let $\lambda^* > 0$ solve $K_{\lambda^*} = 0$, so that $\frac{\lambda^*}{2} = m_G(\beta)$, and let $0 < \lambda < \lambda^*$; then $K_\lambda > 0$.

Decay in boxes. Lemma 6.7 with $S = S_\Lambda$ gives

$$|\,\mathrm{Cov}_{\mu_\Lambda^F}(f,g)| \le C_{f,g}e^{-\lambda D/2},$$

uniformly in the box.

[13] For $w \equiv 1$ this is Bochner's identity $\Gamma_2 = |\nabla^2 g|^2 + (\mathrm{Ric} + \mathrm{Hess}\, V)(\nabla g, \nabla g)$; the weights enter only through the factor index, which the connection, the Laplacian and the Ricci tensor preserve, so only the off-diagonal blocks of $\mathrm{Hess}\, V$ see them.

Limit. For boxes $\Lambda \subset \Lambda'$ let $S_s = (1-s)S_\Lambda + sS_{\Lambda'}$ on the active links of Λ', $s \in [0,1]$, and μ_s the corresponding measure; S_s is admissible, μ_0 is μ_Λ^F times Haar measure on the active links outside Λ, and $\mu_1 = \mu_{\Lambda'}^F$. For smooth f of the active links of Λ,

$$\frac{d}{ds}\mu_s(f) = N\beta \sum_{p \in P(\Lambda')\setminus P(\Lambda)} \mathrm{Cov}_{\mu_s}(f, \operatorname{Re}\operatorname{Tr} Q_p^F),$$

and $|\nabla_e \operatorname{Re}\operatorname{Tr} Q_p| \le \|Q_p\|_{\mathrm{HS}} = \sqrt{N}$. A step in the graph of Lemma 6.7 moves a link by at most one lattice unit, so a plaquette at graph distance k from the support T of f lies within distance $k+1$ of T, there are at most $C_T(k+1)^4$ of them, and every plaquette outside $P(\Lambda)$ is at graph distance at least $R-1$, R the distance from T to the complement of Λ. Lemma 6.7 gives

$$|\mu_{\Lambda'}^F(f) - \mu_\Lambda^F(f)| \le C_f R^4 e^{-\lambda R/2}.$$

The limit therefore exists along every increasing sequence of boxes, and two sequences are compared through their intersections, which are boxes. Continuous gauge-invariant O are uniform limits of smooth gauge-invariant ones, for instance of $e^{\tau\Delta}O$, since the heat semigroup of G^E commutes with left and right translations; with Lemma 3.1 the limit $\omega(O)$ of $W_\Lambda(O)$ exists for all of them.

Properties. Invariance and, for $\beta \ge 0$, reflection positivity follow as in Proposition 3.2(iii)–(iv). For Lipschitz O the functions $e^{\tau\Delta}O$ are smooth with the same Lipschitz constant, since $\mathrm{Ric}_G \ge 0$ [5, Sect. 3.3], so the bound of the first step passes to ω and to Lipschitz observables. Two links of one plaquette have time coordinates that differ by at most one, so observables at time separation t are at graph distance at least $t-c$, and truncated correlations decay at every rate below $m_G(\beta)$. The gap follows from Proposition F.1 as in Proposition 3.2(vi). □

Proof of the rest of Theorem 1.4. By Lemma 6.3 and Table 1, $\kappa_{\mathrm{SU}(N)} = \frac{N}{2}$ and $\kappa_{\mathrm{SO}(N)} = \frac{N-2}{4}$; for SO(4), whose Lie algebra is $\mathfrak{so}(3)\oplus\mathfrak{so}(3)$ acting on $\mathbb{R}^4\otimes\mathbb{C} = \mathbb{C}^2\otimes\mathbb{C}^2$, each ideal gives $\frac12$ by the same computation. The conditions of Theorem 1.4 are those of Theorem 1.5. □

Remark 6.8. Let h_F be the supremum of $\pm\operatorname{Hess} S_0$ on the gauge-fixed directions, so that uniform curvature reaches $|\beta| < 1/(2h_F)$ and no further. At the identity field $-\operatorname{Hess} S_0$ restricted to $X_F = 0$ has the eigenvalue $9+\sqrt{17} = 13.123$ on the torus of side two, so $9+\sqrt{17} \le h_F \le \frac{29}{2}$. A gradient search over SU(2) fields of periods two, four and six finds nothing above 13.97401, the value at an abelian field; certificates made of plaquette forms and vertex forms on the period-two cell cannot go below 14.384. One of us (T. Romanov) [69] constructed such a certificate, with sixteen positive definite vertex forms of size eight and nonnegative plaquette weights, rational with denominator 10^6. It gives $h_F \le 14.420001$ (Theorem L.5; Appendix L gives the derivation and the checks). Like $\frac{29}{2}$ this is a statement about U(N), so it can replace $\frac{29}{2}$ in Theorems 1.4 and 1.5, and the curvature method then reaches $|\beta| < 1/28.840002$, that is $\beta_W < 0.1387$ for SU(2) and $\beta_W < 0.3121$ for SU(3). The bound is computer-assisted and concerns the lattice at fixed spacing, not the continuum limit. The theorems of this paper are stated with $\frac{29}{2}$, which is proved by hand. With this gauge the curvature method stops between $|\beta| = 1/28.840002$ and $\frac{1}{26.25}$.

7. SU(3) up to 3/4

In this section $G = \mathrm{SU}(3)$, $\beta \ge 0$ and $\beta_W = 9\beta$. The distance on SU(3) is $d(U,V) = \|U-V\|_{\mathrm{HS}}$, which is at most the geodesic distance for $\langle X, Y\rangle = -\operatorname{Tr}(XY)$; $K = \operatorname{conv}\mathrm{SU}(3)$; $A(M) = \frac12(M - M^*)$ and P removes the trace. With this inner product $\mathrm{Ric} = \frac32$, the Casimir of the fundamental representation is $\lambda_F = \frac83$ and that of the adjoint representation is 6.

7.1. **Reduction to one link.** The conditional law of an active link is

$$\nu_b(dU) \propto e^{\operatorname{Re}\operatorname{Tr}(Ub)}dU, \qquad b = N\beta \sum_{p \ni e} S_p,$$

where S_p is the staple of e in p, a product of SU(3) matrices and identities, or its adjoint. Six staples give $b \in rK$ with $r = 6N\beta = 2\beta_W$. For a matrix h put $\varphi_h(U) = \operatorname{Re}\operatorname{Tr}(Uh)$ and

$$L_h(b) = \sup\{\operatorname{Cov}_{\nu_b}(g, \varphi_h) : g \text{ 1-Lipschitz for } d\},$$

$$\Lambda(r) = \sup_{b \in rK} \sup_{\|h\|_{\mathrm{HS}}=1} L_h(b).$$

Changing a partner e' changes one staple by a matrix of norm $d(Q_{e'}, Q'_{e'})$. Along the segment b_t from b to b', which stays in rK, $\frac{d}{dt}\nu_{b_t}(g) = \operatorname{Cov}_{\nu_{b_t}}(g, \operatorname{Re}\operatorname{Tr}(U(b' - b)))$, so $W_1(\nu_b, \nu_{b'}) \le N\beta\Lambda(r)\, d(Q_{e'}, Q'_{e'})$. With twelve partners per link,

$$\alpha \le 12N\beta\Lambda(r) = 2r\Lambda(r) = 4\beta_W\Lambda(2\beta_W). \tag{7.1}$$

Since K is convex and contains 0, the mean of $I, \omega I, \omega^2 I$ with $\omega^3 = 1$, $r'K \subset rK$ for $r' \le r$ and Λ is nondecreasing.

7.2. **Gradients and one-forms.**

Lemma 7.1. *In the left trivialisation, $\nabla\varphi_c(U) = -PA(cU)$; if $V = \varphi_b$ then* $\operatorname{Hess} V(X, X) = \operatorname{Re}\operatorname{Tr}(UX^2 b)$*; and $\Delta\varphi_c = -\lambda_F\varphi_c$.*

Proof. $\frac{d}{dt}\operatorname{Re}\operatorname{Tr}(Ue^{tX}c)|_{t=0} = \operatorname{Re}\operatorname{Tr}(cUX) = -\langle PA(cU), X\rangle$ for $X \in \mathfrak{su}(3)$. On a group with a bi-invariant metric $\nabla_X X = 0$ for left-invariant X, so the Hessian is the second derivative along e^{tX}. The function φ_c is a matrix coefficient of the fundamental representation, on which the Casimir acts by λ_F. □

Lemma 7.2. *The bottom of the spectrum of the rough Laplacian $\nabla^*\nabla$ on L^2 one-forms of* SU(3) *is $\mu_0 = \frac{7}{6}$.*

Proof. Trivialise one-forms on the left, so that a one-form is a function $f : \mathrm{SU}(3) \to \mathfrak{su}(3)$. For an orthonormal basis X_a acting as left-invariant vector fields, $\nabla_{X_a} Y = \frac{1}{2}[X_a, Y]$ and $\nabla_{X_a} X_a = 0$, so $\nabla^*\nabla = -\sum_a (\rho(X_a) + \frac{1}{2}\operatorname{ad} X_a)^2$. On the isotypic part $V_\pi \otimes V_\pi^* \otimes \mathfrak{su}(3)$ of $L^2 \otimes \mathfrak{su}(3)$ put $A_a = \rho_\pi(X_a)$, $B_a = \operatorname{ad} X_a$ and $T_a = A_a + B_a$, the action on the tensor product. Since A_a and B_a commute, $A_a + \frac{1}{2}B_a = \frac{1}{2}(A_a + T_a)$ and $A_aT_a + T_aA_a = A_a^2 + T_a^2 - B_a^2$, so on a component $\sigma \subset \pi \otimes \mathrm{ad}$

$$\nabla^*\nabla = \tfrac{1}{2}(C_\pi + C_\sigma) - \tfrac{1}{4}C_{\mathrm{ad}} = \tfrac{1}{2}(C_\pi + C_\sigma) - \tfrac{3}{2},$$

with $C(p, q) = \frac{2}{3}(p^2 + q^2 + pq + 3p + 3q)$ for the representation of highest weight (p, q) [36]. The components of $(p, q) \otimes (1, 1)$ are $(p+1, q+1)$, $(p+2, q-1)$, $(p-1, q+2)$, $(p+1, q-2)$, $(p-2, q+1)$, $(p-1, q-1)$ and (p, q), the last with multiplicity two, one or zero as both, one or none of p, q are positive, and only those with nonnegative labels occur. For $\pi = (0, 0)$ the only component is $(1, 1)$, with value $\frac{3}{2}$. For $\pi = (1, 0)$ the components are $(2, 1)$, $(0, 2)$, $(1, 0)$, with values $\frac{31}{6}$, $\frac{19}{6}$, $\frac{7}{6}$, and $(0, 1)$ is conjugate. For $p + q \ge 2$ the smallest values occur at $(p-1, q-1)$, (p, q), $(p-2, q+1)$ or $(p+1, q-2)$, and each is at least $\frac{1}{3}(2(p^2 + q^2 + pq) + 3(p + q) - 3) - \frac{3}{2} \ge \frac{3}{2}$. □

7.3. **The gap of the one-link measure.**

Lemma 7.3. *For $b \in rK$ the spectral gap of $L = \Delta + \langle \nabla V, \nabla \cdot\rangle$, $V = \varphi_b$, on $L^2(\nu_b)$ is at least*

$$\gamma(r) = \tfrac{39}{64}\mu_0 + \tfrac{3}{2} - \tfrac{r}{2} = \tfrac{283}{128} - \tfrac{r}{2}.$$

Proof. Let $-Lf=\lambda f$ with $\lambda>0$, $\omega=df$ and $\eta=e^{V/2}\omega$; norms are in $L^2(dU)$. Bochner's formula integrated against ν_b gives $\lambda\nu_b|\omega|^2=\nu_b|\nabla\omega|^2+\nu_b((\mathrm{Ric}-\mathrm{Hess}\,V)(\omega,\omega))$, and since $e^{V/2}\nabla\omega=\nabla\eta-\frac12 dV\otimes\eta$,

$$\lambda\|\eta\|^2=\|\nabla\eta-\tfrac12 dV\otimes\eta\|^2+\int(\mathrm{Ric}-\mathrm{Hess}\,V)(\eta,\eta). \tag{7.2}$$

Put $\mathcal{A}=\nabla\eta$ and $\mathcal{B}=\frac12 dV\otimes\eta$. Then

$$\langle\mathcal{A},\mathcal{B}\rangle=\tfrac14\int\langle\nabla V,\nabla|\eta|^2\rangle=\tfrac{\lambda_F}{4}\int V|\eta|^2,$$

because $-\Delta V=\lambda_F V$. Writing $\|\mathcal{A}-\mathcal{B}\|^2=\|\frac58\mathcal{A}-\mathcal{B}\|^2+\frac{39}{64}\|\mathcal{A}\|^2-\frac34\langle\mathcal{A},\mathcal{B}\rangle$ and dropping the first term,

$$\|\mathcal{A}-\mathcal{B}\|^2\ge\tfrac{39}{64}\mu_0\|\eta\|^2-\tfrac12\int V|\eta|^2$$

by Lemma 7.2. With $\mathrm{Ric}=\frac32$, (7.2) gives $\lambda\|\eta\|^2\ge(\frac{39}{64}\mu_0+\frac32)\|\eta\|^2-\int(\mathrm{Hess}\,V(\eta,\eta)+\frac12 V|\eta|^2)$, and the claim follows from the pointwise bound

$$\mathrm{Re}\,\mathrm{Tr}(MX^2)+\tfrac12\,\mathrm{Re}\,\mathrm{Tr}\,M\le\tfrac r2|X|^2,\qquad M=bU\in rK, \tag{7.3}$$

since $\mathrm{Hess}\,V(X,X)=\mathrm{Re}\,\mathrm{Tr}(MX^2)$ and $V=\mathrm{Re}\,\mathrm{Tr}\,M$. The maximum over $|X|=1$ of the left side is convex in M, so it suffices to take $M=rW$ with $W\in\mathrm{SU}(3)$ and, after a conjugation, $W=\mathrm{diag}(e^{i\theta_k})$ with $\sum_k\theta_k=0$; put $c_k=\cos\theta_k$. Write $X=iH$ with H Hermitian, traceless, $\mathrm{Tr}\,H^2=1$, with diagonal d_k and entries H_{jk}. Then

$$\mathrm{Re}\,\mathrm{Tr}(WX^2)=-\sum_k c_kd_k^2-\sum_{j<k}(c_j+c_k)|H_{jk}|^2,$$

with $\sum_k d_k^2+2\sum_{j<k}|H_{jk}|^2=1$, so the maximum is the larger of $q_{\mathrm{off}}=\max_{j<k}-\frac12(c_j+c_k)$ and $q_{\mathrm{diag}}=\max\{-\sum c_kd_k^2:\sum d_k=0,\ \sum d_k^2=1\}$. If l is the third index, $q_{\mathrm{off}}+\frac12\sum_k c_k=\frac12 c_l\le\frac12$. The value q_{diag} is minus the smaller eigenvalue of $\mathrm{diag}(c)$ compressed to the plane $\sum d_k=0$; the eigenvalues of the compression solve $\sum_k(c_k-\mu)^{-1}=0$, that is $3\mu^2-2e_1\mu+e_2=0$ with e_1,e_2 the elementary symmetric functions of the c_k, so $q_{\mathrm{diag}}=\frac13(\sqrt{e_1^2-3e_2}-e_1)$. The inequality $q_{\mathrm{diag}}+\frac12 e_1\le\frac12$ is $\sqrt{e_1^2-3e_2}\le\frac12(3-e_1)$, that is $e_1^2-4e_2+2e_1-3\le0$. When $\sum\theta_k=0$, $\sum c_k^2=1+2c_1c_2c_3$, and with this identity the left side equals $-2(1-c_1)(1-c_2)(1-c_3)\le0$. □

The weight $\frac58$ is the one that gives the coefficient $\frac12$ of V, and (7.3) is an equality at $W=I$. At $r=\frac32$, $\gamma=\frac{187}{128}$.[14]

7.4. **A corrector of degree two.**

Lemma 7.4. *For $b\in rK$, $\|h\|=1$ and every smooth real function u on* $\mathrm{SU}(3)$,

$$L_h(b)\le\sqrt{\nu_b|\nabla u|^2}+\sqrt{\mathrm{Var}_{\nu_b}(\varphi_h+Lu)/\gamma(r)}.$$

Proof. Let $-L\psi=\varphi_h-\nu_b\varphi_h$. For a 1-Lipschitz g, $|\nabla g|\le1$ almost everywhere, because d is at most the geodesic distance, so $\mathrm{Cov}(g,\varphi_h)=\nu_b\langle\nabla g,\nabla\psi\rangle\le\nu_b|\nabla\psi|$. Write $\psi=u+\chi$. Since $\nu_b(Lu)=0$, χ solves $-L\chi=X-\nu_bX$ with $X=\varphi_h+Lu$, and

$$\nu_b|\nabla\chi|^2=\mathrm{Cov}(\chi,X)\le\sqrt{\mathrm{Var}\,\chi\ \mathrm{Var}\,X}\le\sqrt{\nu_b|\nabla\chi|^2\,\mathrm{Var}\,X/\gamma}$$

by Lemma 7.3. Hence $\nu_b|\nabla\chi|^2\le\mathrm{Var}\,X/\gamma$, and $\nu_b|\nabla\psi|\le\sqrt{\nu_b|\nabla u|^2}+\sqrt{\nu_b|\nabla\chi|^2}$. □

[14] The direct Bochner bound, with Minkowski's inequality in place of the cross term, gives $(\sqrt{\mu_0}-\frac{g}{2}r)^2+\frac32-r$ with $g^2=\max_W\|PA(W)\|^2=2.0655$; it is exact at $b=0$, where the gap is $\frac83$, but falls below γ for $r>0.24$ and vanishes before $r=1$.

The best u is ψ itself, and then the second term vanishes. With $u = s\varphi_h$, Lemma 7.1 gives $\varphi_h + Lu = (1 - s\lambda_F)\varphi_h + s\Gamma$ with $\Gamma = \langle \nabla V, \nabla \varphi_h \rangle$; the choice $s = 1/\lambda_F$ removes φ_h and leaves the drift. This blended corrector reaches $\beta_W = \frac{45}{64}$ and stops near 0.715 (Section 9). We take u in a space of polynomials of degree two.

Let $\mathcal{P}_2$ be the real vector space of real polynomials of degree one or two in the entries u_{ij} of U and $\bar{u}_{ij}$ of $\bar{U}$. It has dimension 189: a basis (B_k) consists of $\frac{1}{2}(m + \bar{m})$ and $\frac{1}{2i}(m - \bar{m})$ for each pair $\{m, \bar{m}\}$ of conjugate monomials with $m \neq \bar{m}$, and m itself when $m = \bar{m}$. Identify $h \in \mathbb{C}^{3\times 3}$ with $\mathbb{R}^{18}$ through the real and imaginary parts of its entries. For a real 189×18 matrix C put

$$q_h = \sum_{k} \sum_{m=1}^{18} C_{km} h_m B_k \in \mathcal{P}_2,$$

linear in h. For diagonal $b = \operatorname{diag}(a_1, a_2, x_3 + iy_3)$ we have $V = \sum_k b_k \varphi_{F_k}$ with $(b_k) = (a_1, a_2, x_3, y_3)$ and $F = (e_{11}, e_{22}, e_{33}, ie_{33})$, so

$$X_h = \varphi_h + Lq_h = \varphi_h + \Delta q_h + \sum_k b_k \langle \nabla \varphi_{F_k}, \nabla q_h \rangle \tag{7.4}$$

is a polynomial of degree at most three, affine in b, while ∇q_h does not depend on b. The squares of these functions have degree at most six, so the quadratic forms $h \mapsto \nu_b |\nabla q_h|^2$ and $h \mapsto \nu_b[(X_h - \kappa)^2]$ are combinations of moments of ν_b of degree at most six.

To compute Δ and $\langle \nabla f, \nabla g \rangle$ exactly we use the basis of $\mathfrak{su}(3)$

$$e_{jk} - e_{kj}, \quad i(e_{jk} + e_{kj})\ (j < k), \quad i(e_{11} - e_{22}), \quad i(e_{11} + e_{22} - 2e_{33}),$$

acting as left-invariant fields $X_a f(U) = \frac{d}{dt} f(U e^{tX_a})|_{t=0}$, with weights $w_a = \frac{1}{2}$ for the first seven and $\frac{1}{6}$ for the last. Then $\sum_a w_a X_a \otimes X_a$ is the Casimir tensor of $\langle X, Y \rangle = -\operatorname{Tr}(XY)$, so

$$\langle \nabla f, \nabla g \rangle = \sum_a w_a\, X_a f\, X_a g, \qquad \Delta = \sum_a w_a X_a^2.$$

The fields map monomials to combinations of monomials of the same degree with coefficients in $\mathbb{Z}[i]$; for instance $\Delta \varphi_h = -\frac{8}{3}\varphi_h$, as in Lemma 7.1.

7.5. **Orbits of staple sums.**

Lemma 7.5. *Let $V, W \in \mathrm{SU}(3)$.*

(i) *$L_h(VbW) = L_{V^{-1}hW^{-1}}(b)$ and $L_h(\bar{b}) = L_{\bar{h}}(b)$.*
(ii) *Every $b \in \mathbb{C}^{3\times 3}$ is in the orbit of some $\operatorname{diag}(a_1, a_2, z_3)$ with $a_1 \geq a_2 \geq |z_3|$ and $\operatorname{Im} z_3 \geq 0$.*
(iii) *A diagonal matrix lies in K if and only if it lies in the convex hull of the diagonal torus T of $\mathrm{SU}(3)$.*

Proof. (i) If U has law ν_{VbW} then WUV has law ν_b; the map preserves d and Haar measure, and $\varphi_h(U) = \varphi_{V^{-1}hW^{-1}}(WUV)$. For $\bar{b}$ use $U \mapsto \bar{U}$. (ii) Take a singular value decomposition $b = V_1 \Sigma W_1$ and move the determinants of V_1, W_1 into the last entry; a complex conjugation makes $\operatorname{Im} z_3 \geq 0$. (iii) Let c be diagonal with nonzero entries of distinct moduli and let U maximise $\operatorname{Re}\operatorname{Tr}(cU)$ on $\mathrm{SU}(3)$. The derivative along $\mathfrak{su}(3)$ vanishes, so the anti-Hermitian part of cU is a multiple $i\lambda I$ of the identity and $cU = H + i\lambda I$ with H Hermitian. Then $H^2 + \lambda^2 I = cc^*$ is diagonal with distinct entries, H commutes with it and is diagonal, and $U \in T$. Hence $\max_{\mathrm{SU}(3)} \operatorname{Re}\operatorname{Tr}(cU) = \max_T \operatorname{Re}\operatorname{Tr}(ct)$ for such c, and by continuity for every diagonal c. A diagonal matrix outside $\operatorname{conv} T$ is separated from it by such a functional, which then separates it from K. □

Since every $b \in rK$ has operator norm at most r,

$$\Lambda(r) = \sup\Big\{ \sup_{\|h\|=1} L_h(b) : b \in \mathcal{R}_r \Big\},$$

where $\mathcal{R}_r$ is the set of $\operatorname{diag}(a_1,a_2,x_3+iy_3) \in r\operatorname{conv} T$ with $a_1 \ge a_2 \ge |x_3+iy_3|$ and $y_3 \ge 0$, a subset of the box $[0,r]^2 \times [-r,r] \times [0,r]$ in the coordinates (a_1,a_2,x_3,y_3).

7.6. **Exact moments and interpolation.**

Lemma 7.6. *For $D, D' \in \mathbb{C}^{3\times 3}$ let $\mathbb{E}_H$ be the Haar mean on $\mathrm{SU}(3)$ and*

$$Z(D,D') = \mathbb{E}_H \exp\big(\tfrac{1}{2}\operatorname{Tr}(UD) + \tfrac{1}{2}\operatorname{Tr}(\bar U D')\big).$$

Then

$$Z(D,D') = \sum_{p\ge q\ge 0} \frac{s_{(p,q)}(DD'^{\top})\, G_{p,q}(\det D)\, G_{p,q}(\det D')}{d_{p,q}},$$

with $G_{p,q}(z) = \sum_{k\ge 0} c_{(p+k,q+k,k)} z^k$, $c_\lambda = 2^{-|\lambda|}/\prod \mathrm{hooks}(\lambda)$, $d_{p,q} = \tfrac{1}{2}(p-q+1)(q+1)(p+2)$ and $s_{(p,q)}$ the Schur polynomial. The moments of degree j in U and k in $\bar U$ are $\mathbb{E}_H[u^\alpha \bar u^\beta e^{\operatorname{Re}\operatorname{Tr}(Ub)}] = 2^{|\alpha|+|\beta|}\partial_D^\alpha \partial_{D'}^\beta Z(b,\bar b)$, the entry u_{ij} pairing with D_{ji}.

Proof. We use $(\operatorname{Tr} X)^n = \sum_{\lambda\vdash n} f^\lambda s_\lambda(X)$, where f^λ is the number of standard tableaux of shape λ, given by the hook length formula [76, Cor. 7.21.6]. Expanding the exponential gives

$$e^{\frac{1}{2}\operatorname{Tr} X} = \sum_{n\ge 0} \frac{(\operatorname{Tr} X)^n}{2^n n!} = \sum_\lambda c_\lambda s_\lambda(X),$$

the sum over partitions with at most three parts. For $U \in \mathrm{SU}(3)$ one has $s_{\lambda+(k,k,k)}(UD) = (\det D)^k s_\lambda(UD)$, so

$$e^{\frac{1}{2}\operatorname{Tr}(UD)} = \sum_{p\ge q} G_{p,q}(\det D)\, \chi_{p,q}(UD),$$

where $\chi_{p,q}$ is the irreducible character of highest weight $(p,q,0)$ and $d_{p,q}$ its dimension. Since $\bar U = (U^{-1})^\top$, we have $\chi_{p,q}(\bar U D') = \chi_{p,q}(U^{-1}D'^{\top})$, and Schur orthogonality [23, Ch. 2],

$$\mathbb{E}_H\big[\chi_\rho(UA)\chi_{\rho'}(U^{-1}B)\big] = \delta_{\rho\rho'}\frac{\chi_\rho(AB)}{d_\rho},$$

gives the expansion. The bound below gives absolute convergence. Finally $\operatorname{Re}\operatorname{Tr}(Ub) = \tfrac{1}{2}\operatorname{Tr}(Ub) + \tfrac{1}{2}\operatorname{Tr}(\bar U \bar b)$. □

The computation keeps $p \le 34$ and $k \le 11$. Every omitted partition has more than 34 boxes, and since $|s_\lambda(X)| \le \dim(\lambda)\|X\|^{|\lambda|}$ and $\sum_{\lambda\vdash n} f^\lambda \dim\lambda = 3^n$ by Schur–Weyl duality, the omitted part of each factor is at most $\sum_{n>34}(\frac{3}{2}\|D\|)^n/n!$; Cauchy's estimate on a polydisc around b turns this into a bound for every derivative (Appendix E).

Lemma 7.7 (Interpolation on a box)**.** *Let $Q = \prod_k [l_k, l_k+w_k]$ in the coordinates (a_1,a_2,x_3,y_3), with corners b_i and multilinear weights $\omega_i(b) \ge 0$, $\sum_i \omega_i(b) = 1$, and let $\varsigma = w_1^2 + w_2^2 + \max(w_3^2, w_4^2)$. For $b \in Q$:*

(i) *$Z(b) \ge e^{-\varsigma/8}\sum_i \omega_i(b) Z(b_i)$;*
(ii) *$\nu_b(f) \le e^{\varsigma/8}\max_i \nu_{b_i}(f)$ for every $f \ge 0$ that does not depend on b;*
(iii) *if $Y_b = (\mathcal{A} + \sum_k (b_k-\beta_k)\mathcal{B}_k)^2$ with functions $\mathcal{A}, \mathcal{B}_k$ of U and a fixed point β, then*

$$\nu_b(Y_b) \le e^{\varsigma/8}\Big(\max_i \nu_{b_i}(Y_{b_i}) + \sum_k \frac{w_k^2}{4}\,\frac{\max_i Z(b_i)\nu_{b_i}(\mathcal{B}_k^2)}{\min_i Z(b_i)}\Big).$$

Proof. Write $E(b) = e^{\operatorname{Re}\operatorname{Tr}(Ub)} = \prod_k e^{c_k b_k}$, where

$$c = (\operatorname{Re} u_{11}, \operatorname{Re} u_{22}, \operatorname{Re} u_{33}, -\operatorname{Im} u_{33}),$$

so that $|c_k| \le 1$ and $c_3^2 + c_4^2 \le 1$. In each coordinate the chord of $e^{c_k x}$ over $[l_k, l_k+w_k]$ lies above the function and, by Hoeffding's lemma [48] for a two-point variable, below $e^{c_k x} e^{c_k^2 w_k^2/8}$. Multiplying,

$$E(b) \le \sum_i \omega_i(b) E(b_i) \le E(b)\, e^{\sum_k c_k^2 w_k^2/8} \le E(b)\, e^{\varsigma/8}. \tag{7.5}$$

Integrating the right inequality gives (i), and (ii) follows from the left one and (i). For (iii) let $\mathcal{F}(b) = Y_b E(b)$ and I_k the linear interpolation in the k-th coordinate, so that $\mathcal{F} - I_1 I_2 I_3 I_4 \mathcal{F} = \sum_k I_1 \cdots I_{k-1}(\mathcal{F} - I_k \mathcal{F})$. Along the k-th coordinate $\mathcal{F}$ is $\ell^2 e^{c_k x}$ times a positive factor, with ℓ affine of slope $\mathcal{B}_k$, and $\partial_k^2 \mathcal{F} = ((c_k \ell + 2\mathcal{B}_k)^2 - 2\mathcal{B}_k^2)E \geq -2\mathcal{B}_k^2 E$. With the Green kernel of the interval, whose integral is at most $w_k^2/8$,

$$\mathcal{F} - I_k \mathcal{F} \leq 2\mathcal{B}_k^2 \int G\, E \leq \tfrac{w_k^2}{4} \mathcal{B}_k^2 \sum_i \omega_i^{(k)} E(b_i)$$

by the left inequality of (7.5), with weights $\omega_i^{(k)} \geq 0$ of sum one that do not depend on U. The operators I_j have nonnegative weights of sum one; integrate over U and divide by $Z(b)$, bounded below by (i). □

7.7. **The cover and the theorem.** Fix a box Q and a matrix C. For any constant κ, $\operatorname{Var}(X_h) \leq \nu_b[(X_h - \kappa)^2]$; take $\kappa = \langle \kappa(b), h \rangle$ with $\kappa(b)$ affine in b, fitted to the means at the corners. By (7.4), $X_h - \kappa = \mathcal{A} + \sum_k (b_k - \beta_k)\mathcal{B}_k$ with β the centre of Q and $\mathcal{B}_k = \langle \nabla \varphi_{F_k}, \nabla q_h \rangle - \langle \partial_k \kappa, h \rangle$, and Lemmas 7.4 and 7.7 give, for all $b \in Q \cap \mathcal{R}_r$,

$$\begin{aligned} \sup_h L_h(b) \leq{}& \sqrt{e^{\varsigma/8} \max_i \lambda_A(b_i)} \\ &+ \sqrt{\frac{e^{\varsigma/8}}{\gamma(r)} \Big(\max_i \lambda_V(b_i) + \frac{e_N}{\min_i Z(b_i)} \Big)}, \end{aligned} \tag{7.6}$$

where $e_N = \sum_k \frac{w_k^2}{4} \max_i Z(b_i) \lambda_k(b_i)$ and λ_A, λ_V, λ_k are the largest eigenvalues of the quadratic forms $h \mapsto \nu_{b_i} |\nabla q_h|^2$, $h \mapsto \nu_{b_i}[(X_h - \kappa)^2]$ at $b = b_i$, and $h \mapsto \nu_{b_i}[\mathcal{B}_k^2]$ on $\mathbb{R}^{18}$. Their entries are moments of ν_{b_i} of degree at most six, computed with rigorous error bounds (Appendix E); each largest eigenvalue is bounded above by an $LDL^\top$ factorisation of tI minus the computed form with certified positive pivots, plus the norm of its error matrix by Weyl's inequality [45, 51].

The matrix C is chosen in each box in floating point. At the mean of the corner moments and at $b = \beta$, Lemma 7.4 for $u = q_h$ is a quadratic problem in C: for $\theta > 0$ we minimise the trace of $\nu |\nabla q|^2 + \theta \operatorname{Var}(X)/\gamma$ over the 18 coordinate directions h, a least-squares problem of size 189, keep the θ on a geometric grid of 48 values from 0.05 to 80 that gives the smallest bound, and round C to multiples of 2^{-30}. Any choice gives a valid bound.

A box is discarded when it contains no point with $a_1 \geq a_2 \geq |z_3|$, or when a linear functional $\ell(b) = \operatorname{Re} \operatorname{Tr}(c^* b)$ satisfies $\min_Q \ell > r \max_T \ell$, which separates it from $r \operatorname{conv} T$ and hence, by Lemma 7.5(iii), from $\mathcal{R}_r$.

Proposition 7.8. *$2r\Lambda(r) \leq 0.999887$ at $r = \frac{3}{2}$.*

Proof. The box $[0, r]^2 \times [-r, r] \times [0, r]$ is cut into 8192 boxes of side $r/8$. Of these, 841 survive the two tests, and 581 satisfy $2rB_Q < 1$ for the right side B_Q of (7.6), with $\gamma(r) = \frac{187}{128}$. Each of the other 260 is cut into sixteen boxes of side $r/16$; 1760 of these survive the tests, and all satisfy $2rB_Q < 1$. The 2341 certified boxes cover $\mathcal{R}_r$. The largest bound is 0.999887, on the box $[1.3125, 1.5] \times [0.9375, 1.125] \times [0.5625, 0.75] \times [0, 0.1875]$, and the median is 0.983. Figure 6 shows the cover, and Appendix E the error analysis. □

The same cover at a larger r bounds $\Lambda(r)$ where $2r\Lambda(r)$ is no longer below one; the plaquette blocks of Section 8 use that bound.

Proposition 7.9. *$2r\Lambda(r) \leq 1.029967$ at $r = \frac{38}{25}$ and $2r\Lambda(r) \leq 1.027468$ at $r = \frac{77}{50}$.*

Proof. As in the proof of Proposition 7.8, with $\gamma(r) = \frac{4643}{3200}$ and the threshold $2rB_Q < 1.03$: of the 8192 boxes of side $r/8$, 784 survive the two tests and 690 satisfy the threshold; each of the other 94 is cut into sixteen boxes of side $r/16$, of which 373 survive the tests, and all satisfy it. The 1063 certified boxes cover $\mathcal{R}_r$, and the largest bound is 1.029967. At $r = \frac{77}{50}$, with $\gamma(r) = \frac{4611}{3200}$ and the threshold $2rB_Q < 1.0275$: 826 boxes of side $r/8$ survive the tests




FIGURE 6. The 2341 certified boxes at $r = \frac{3}{2}$, plotted at their centres in the coordinates (a_1, a_2) of the staple sum $\operatorname{diag}(a_1, a_2, x_3 + iy_3)$, coloured by the certified bound on $2r \sup_h L_h$ over the box; the larger points are the boxes of side $r/8$. (a) All boxes, with height $|x_3 + iy_3|$. (b) The 419 boxes with bound above 0.99, with height x_3; the largest bounds sit in the interior of $\mathcal{R}_r$, not at its extreme points.

and 513 satisfy the threshold; the other 313 are cut into sixteen, of which 2311 survive and 2281 satisfy it; the other 30 are cut into sixteen, of which 31 survive, and all satisfy it. The 2825 certified boxes cover $\mathcal{R}_r$, and the largest bound is 1.027468. □

Proof of Theorem 1.6. Proposition 7.8 gives

$$\Lambda\big(\tfrac{3}{2}\big) \le \frac{0.999887}{3} < 0.333296.$$

For $\beta_W \le \frac{3}{4}$ we have $2\beta_W \le \frac{3}{2}$, and (7.1) with the monotonicity of Λ gives

$$\alpha \le 4\beta_W \Lambda\big(\tfrac{3}{2}\big) < 1.333183\,\beta_W \le \tfrac{3}{4} \cdot 1.333183 < 1.$$

Proposition 3.2 applies with the Hilbert–Schmidt metric. □

8. SU(3) WITH PLAQUETTE BLOCKS

In this section $G = \mathrm{SU}(3)$, $\beta \ge 0$, $\beta_W = 9\beta$, $N\beta = \beta_W/3$, and the notation is that of Section 7: d is the Hilbert–Schmidt distance, $K = \operatorname{conv}\mathrm{SU}(3)$, $\nu_c(dU) \propto e^{\operatorname{Re}\operatorname{Tr}(Uc)}dU$, $\varphi_h(U) = \operatorname{Re}\operatorname{Tr}(Uh)$ and $\Lambda(r)$ is the modulus of (7.1). Put $Z(c) = \int e^{\operatorname{Re}\operatorname{Tr}(Uc)}dU$ and $g = \log Z$ on $\mathbb{C}^{3\times 3}$, viewed as $\mathbb{R}^{18}$ with the inner product $\langle x, y\rangle = \operatorname{Re}\operatorname{Tr}(xy^*)$. The gradient of g at c is the conjugate mean $m(c)^*$, $m(c) = \nu_c(U) \in K$, and $\operatorname{Hess} g(c)(h,h) = \operatorname{Var}_{\nu_c}(\varphi_h)$. Since $Z(VcW) = Z(c)$ for $V, W \in \mathrm{SU}(3)$, the same holds for g.

The blocks are those of Lemma 5.1, which concerns the line field only and holds for every group. For a block plaquette with active links e_1, e_2, e_3, $\operatorname{Re}\operatorname{Tr} Q_p = \operatorname{Re}\operatorname{Tr}(y_1y_2y_3)$ with $y_i = Q_{e_i}$ or $Q_{e_i}^{-1}$, and the conditional law of the block is

$$\mu_b(dy) \propto \exp\Big(\sum_i \operatorname{Re}\operatorname{Tr}(y_i b_i) + N\beta \operatorname{Re}\operatorname{Tr}(y_1y_2y_3)\Big)dy, \tag{8.1}$$

where b_i is $N\beta$ times the sum of the five staples of y_i outside the block plaquette, so $b_i \in 5N\beta K$ and $\|b_i\|_{\mathrm{op}} \le 5N\beta$. On $\mathrm{SU}(3)^3$ we use $d(y, y') = \sum_i \|y_i - y_i'\|_{\mathrm{HS}}$.

8.1. **The sensitivity of one block.** Let $r = 2\beta_W = 6N\beta$. Let λ_* and τ be a lower and an upper bound for the eigenvalues of $\operatorname{Hess} g$ on rK, M an upper bound on rK for $|D^3 g(c)(u,v,v)|$ over unit vectors u, v, and $\gamma = \gamma(r)$ the gap of Lemma 7.3. Put

$$w = \tfrac{3}{2}(N\beta)^2(\tau - \lambda_*), \qquad s = \Lambda(r) + (e^w - 1)e^w\sqrt{\tau/\gamma},$$

$$C = 5(N\beta)^2\Big(\tau s + \frac{\sqrt{3}\,N\beta\,M\,e^w}{\gamma}\Big), \tag{8.2}$$

$$K_3 = \frac{(1 + N\beta\Lambda(r))\,s}{1 - C}, \qquad \alpha_3 = 10N\beta K_3,$$

assuming $C < 1$.

Lemma 8.1. *Let $a, c_0 \in 5N\beta K$ and*

$$\kappa(dx) \propto \exp\big(\operatorname{Re}\operatorname{Tr}(xa) + g(c_0 + N\beta x)\big)dx \quad \text{on } \mathrm{SU}(3).$$

For every 1-Lipschitz f and every $h \in \mathbb{C}^{3\times 3}$:

(i) $\operatorname{Cov}_\kappa(f, \varphi_h) \le s\|h\|_{\mathrm{HS}}$;

(ii) *if c_0 moves in the direction k inside $5N\beta K$, then*

$$\big|\tfrac{d}{dt}\kappa(f)\big| \le N\beta(\tau s + \sqrt{3}N\beta M e^w/\gamma)\|k\|_{\mathrm{HS}}.$$

Proof. (i) The segment from c_0 to $c_0 + N\beta x$ lies in $6N\beta K = rK$, and Taylor's formula gives

$$g(c_0 + N\beta x) = g(c_0) + N\beta \operatorname{Re}\operatorname{Tr}(x\, m(c_0)) + R(x)$$

with $R(x) = \frac{1}{2}(N\beta)^2 \operatorname{Hess} g(\xi)(x,x)$. Since $\|x\|_{\mathrm{HS}}^2 = 3$, $R \in [\frac{3}{2}(N\beta)^2\lambda_*, \frac{3}{2}(N\beta)^2\tau]$ and $\operatorname{osc} R \le w$. Hence κ is the tilt $e^R\nu/\nu(e^R)$ of $\nu = \nu_{c'}$ with $c' = a + N\beta m(c_0) \in rK$. By definition of Λ, $\operatorname{Cov}_\nu(f, \varphi_h) \le \Lambda(r)\|h\|$. The function f is 1-Lipschitz for the geodesic distance, which dominates d, so $\operatorname{sd}_\nu f \le \gamma^{-1/2}$ by Lemma 7.3, and $\operatorname{sd}_\nu \varphi_h \le \sqrt{\tau}\|h\|$. Lemma 5.2 gives (i).

(ii) The derivative is $\operatorname{Cov}_\kappa(f, \langle \nabla g(c_0 + N\beta x), k\rangle)$. Write $\nabla g(c_0 + N\beta u) = \nabla g(c_0) + N\beta \operatorname{Hess} g(c_0)u + N(u)$. The linear term is φ_h with $\|h\| \le N\beta\tau\|k\|$ and contributes at most $N\beta\tau s\|k\|$ by (i). On the segment from c_0 the Hessian moves by at most $N\beta M\|u\| \le \sqrt{3}N\beta M$, so $x \mapsto \langle N(x), k\rangle$ is $\sqrt{3}(N\beta)^2 M\|k\|$-Lipschitz for d. The law κ is a tilt of ν by a function of oscillation at most w, so its spectral gap is at least $e^{-w}\gamma$ by the Holley–Stroock lemma [46], and the covariance of two Lipschitz functions under κ is at most e^w/γ times the product of their constants. □

Lemma 8.2. *Let $C < 1$. If b, b' have all fields in $5N\beta K$ and differ only in b_i, then $W_1(\mu_b, \mu_{b'}) \le K_3\|b_i - b_i'\|_{\mathrm{HS}}$ for the metric d.*

Proof. By the cyclic symmetry of $\operatorname{Re}\operatorname{Tr}(y_1y_2y_3)$ we may take $i = 1$. Given y_1, y_3, the law of y_2 is ν_c with $c = b_2 + N\beta y_3 y_1 \in rK$. Changing y_1 by η in d moves c by $N\beta\eta$ and ν_c by at most $N\beta\Lambda(r)\eta$ in W_1, as in (7.1); so $F(y_1, y_3) = \mathbb{E}[f \mid y_1, y_3]$ is $(1 + N\beta\Lambda(r))$-Lipschitz in each variable. The law of (y_1, y_3) is proportional to $e^{\operatorname{Re}\operatorname{Tr}(y_1b_1) + \operatorname{Re}\operatorname{Tr}(y_3b_3)}Z(b_2 + N\beta y_3 y_1)$, and $Z(b_2 + N\beta y_3 y_1) = Z(y_3^{-1}b_2 + N\beta y_1) = Z(b_2y_1^{-1} + N\beta y_3)$, so its two conditional laws have the form of Lemma 8.1, with $c_0 = y_3^{-1}b_2$ and $c_0 = b_2y_1^{-1}$. Moving y_3 to y_3' moves $y_3^{-1}b_2$ along a segment in $5N\beta K$ of length at most $\|b_2\|_{\mathrm{op}}\|y_3 - y_3'\| \le 5N\beta\|y_3 - y_3'\|$. The coupling argument of Lemma 5.4 now gives $d_1 \le s\varepsilon + Cd_3$, $d_3 \le Cd_1$ and $|\mu_b(f) - \mu_{b'}(f)| \le (1 + N\beta\Lambda(r))(d_1 + d_3) \le K_3\varepsilon$. □

8.2. **The constants and the proof of Theorem 1.7.**

Lemma 8.3. *At $r = \frac{3}{2}$ every eigenvalue of $\operatorname{Hess} g$ on rK lies in $[0.056015, 0.247316]$, and $|D^3 g(c)(u,v,v)| \le 0.182126$ for $c \in rK$ and unit vectors u, v. At $r = \frac{38}{25}$ the same holds with $[0.054978, 0.248280]$ and 0.183546, and at $r = \frac{77}{50}$ with $[0.053955, 0.249240]$ and 0.184989.*

Proof. For $V, W \in \mathrm{SU}(3)$, $g(VcW) = g(c)$ and $x \mapsto VxW$ is an isometry, so the eigenvalues of $\operatorname{Hess} g$ and the norm of $D^3 g$ are constant on the orbits, and by Lemma 7.5 it suffices to bound them on $\mathcal{R}_r$. The derivatives of g are the cumulants of φ under ν_c. Cut the box $[0,r]^2 \times [-r,r] \times [0,r]$ into boxes of side $r/8$ and keep the 841 (784 at $r = \frac{38}{25}$, 826 at $r = \frac{77}{50}$) that survive the two tests of Section 7.7. On each box Q we compute, with rigorous error bounds, the moments of φ of degree at most three at the sixteen corners b_i and at the centre b_0, and put $k = m(b_0)$, the mean of φ at the centre.

Upper bound. For $b \in Q$ and every h, $\operatorname{Var}_{\nu_b}(\varphi_h) \le \nu_b[(\varphi_h - \langle k, h\rangle)^2] \le e^{\varsigma/8} \max_i \nu_{b_i}[(\varphi_h - \langle k, h\rangle)^2]$ by Lemma 7.7(ii).

Lower bound. Integrating the right inequality of (7.5) against $f \ge 0$ and dividing by $Z(b) \le \sum_i \omega_i Z(b_i)$, the left one, gives $\nu_b(f) \ge e^{-\varsigma/8} \min_i \nu_{b_i}(f)$. With $f = (\varphi_h - \nu_b\varphi_h)^2$ this gives $\operatorname{Var}_{\nu_b}(\varphi_h) \ge e^{-\varsigma/8} \min_i \operatorname{Var}_{\nu_{b_i}}(\varphi_h)$.

Third cumulant. Put $\delta = m(b) - k$. The mean value theorem and the upper bound give $\|\delta\| \le \tau\rho + e$, with ρ the largest distance from b_0 in Q and e the error of the computed mean. Expanding the central moment about k,

$$\begin{aligned} D^3 g(b)(u,v,v) = T_b(u,v,v) - \delta_u \operatorname{Var}(\varphi_v) \\ - 2\delta_v \operatorname{Cov}(\varphi_u, \varphi_v) - \delta_u \delta_v^2, \end{aligned}$$

with $T_b = \nu_b[(\varphi - k)^{\otimes 3}]$, $\delta_u = \langle \delta, u\rangle$ and the moments under ν_b. For a function G of either sign, writing $G = G^+ - G^-$ and applying the two bounds above to $G^\pm$ gives $|\nu_b(G)| \le (\max_i |\nu_{b_i}(G)| + \varepsilon \max_i \nu_{b_i}|G|)/(1-\varepsilon)$ with $\varepsilon = 1 - e^{-\varsigma/8}$. For $G = (\varphi_u - k_u)(\varphi_v - k_v)^2$ we use $|\nu_{b_i}(G)| \le \|T_{b_i}\|$, the largest singular value of the 18×324 unfolding of T_{b_i}, and $\nu_{b_i}|G| \le 2\sqrt{3}\, \nu_{b_i}[(\varphi_v - k_v)^2]$, since $\|U - k\|_{\mathrm{HS}} \le 2\sqrt{3}$. The largest eigenvalues and singular values are bounded as in Section 7.7, and the maxima over these boxes are the numbers stated (Appendix C). □

Proof of Theorem 1.7. Let first $\beta_W \le \frac{3}{4}$, so that $r = 2\beta_W \le \frac{3}{2}$. Every field met in Lemmas 8.1 and 8.2 lies in $\frac{3}{2}K$, so the constants of Lemma 8.3 at $\frac{3}{2}$, $\Lambda(\frac{3}{2}) \le \frac{0.999887}{3}$ from Proposition 7.8 and $\gamma = \gamma(\frac{3}{2}) = \frac{187}{128}$ apply; with these fixed constants every quantity in (8.2) is increasing in β_W. For $\frac{3}{4} < \beta_W \le \frac{19}{25}$ the fields lie in $\frac{38}{25}K$, and the constants of Lemma 8.3 at $\frac{38}{25}$, $\Lambda(\frac{38}{25}) \le \frac{1.029967}{3.04}$ from Proposition 7.9 and $\gamma(\frac{38}{25}) = \frac{4643}{3200}$ apply in the same way; for $\frac{19}{25} < \beta_W \le \frac{77}{100}$ the same holds at $\frac{77}{50}$, with $\Lambda(\frac{77}{50}) \le \frac{1.027468}{3.08}$ and $\gamma(\frac{77}{50}) = \frac{4611}{3200}$. Changing an active link j by η changes one staple of each of its ten partners outside its block by a matrix of norm η, so the field b_i of each such partner moves by $N\beta\eta$, and Lemma 8.2 gives $\sum_B C_{Bj} \le 10N\beta K_3 = \alpha_3(\beta_W)$. In ball arithmetic, at $\beta_W = \frac{3}{4}$, $w \le 0.017935$, $s \le 0.34088$, $C \le 0.043520$, $K_3 \le 0.38609$ and $\alpha_3 \le 0.96521$; at $\beta_W = 0.72$, $\alpha_3 \le 0.91815$; at $\beta_W = 0.76$, with the second set of constants, $w \le 0.018609$, $s \le 0.34673$, $C \le 0.045770$, $K_3 \le 0.39454$ and $\alpha_3 \le 0.99950$; at $\beta_W = 0.77$, with the third set, $w \le 0.019298$, $s \le 0.34186$, $C \le 0.047232$, $K_3 \le 0.38953$ and $\alpha_3 \le 0.99978$. Proposition 5.6, whose proof does not use the group, applies with the Hilbert–Schmidt metric. □

Remark 8.4. Theorem 1.6 ends where $2r\Lambda(r)$ reaches one. The blocks remove the factor $\frac{12}{10}$ of the single-link sum and keep most of it after paying for the three links: at $\beta_W = \frac{3}{4}$ the gap bound rises from $1.1 \cdot 10^{-4}$ to 0.0177, and the block bound is the better one for $\beta_W \ge 0.715$. Before the corrections for the three links their sum is $\frac{10}{12}$ of the single-link sum $2r\Lambda(r)$, so it can stay below one while $2r\Lambda(r)$ is slightly above one; Proposition 7.9 certifies $2r\Lambda(r) \le 1.03$ at $r = 1.52$ and ≤ 1.0275 at $r = 1.54$, which is enough at $\beta_W = 0.76$ and 0.77, with little room. Evaluated in floating point with the constants of a scan of $\mathcal{R}_r$, the block sum stays below one up to $\beta_W \approx 0.78$; that computation is not certified and is not used.




FIGURE 7. The floating-point value of $2r \sup_h$ of the bound of Lemma 7.4 on the face $b = \mathrm{diag}(r, \rho, \rho)$, as a function of ρ/r, for $r = 1.44$, 1.50 and 1.54: dashed with the corrector $u = s\varphi_h$ for the best s, solid with $u = q_h$ of degree two for the best C. The grey plane is the Dobrushin threshold one. The first corrector is above it already at $r = 1.44$; the second stays below it at $r = 1.50$, where Proposition 7.8 certifies the bound on all of $\mathcal{R}_r$, and crosses it by $r = 1.54$.

9. WHERE THE METHODS STOP

SU(2). Theorem 1.2 closes the single-link route in every forest gauge at $\beta_W = \frac{1}{3}$. Blocks of several links, where the influence of a boundary link on a whole block is bounded at once, in the spirit of the Dobrushin–Shlosman criterion [32], pass it: plaquette blocks reach $\beta_W = 0.351$ (Theorem 1.3), and the sharper floating-point bound of Remark 5.7 stops near 0.357. Larger blocks are the next step.

Curvature. Remark 6.8 locates the end of the curvature method in this gauge between $|\beta| = 1/28.840002$, by the certificate of Appendix L, and $\frac{1}{26.25}$.

SU(3). The route of Section 7 bounds the $H^{-1}(\nu_b)$ norm of $\varphi_h - \nu_b\varphi_h$, which is at least $L_h(b)$, and needs it uniformly on rK. Its Galerkin values (polynomials of degree two) at matrices of rK give $2r$ times the norm equal to 0.79 at $r = 1.25$, 0.89 at $r = 1.4$, 0.97 at $r = 1.5$ and 1.009 at $r = 1.55$, so the route stops at or below $\beta_W \approx 0.77$ even with exact constants. With $u = s\varphi_h$, Lemma 7.4 stops earlier: in floating point its largest values on $\mathcal{R}_r$ lie on the face $\mathrm{diag}(r, \rho, \rho)$ with ρ between $0.75r$ and $0.8r$, and $2r$ times the bound is 0.9458 at $r = 1.36$, 0.9767 at $r = 1.40$ and 1.0080 at $r = 1.44$, so this corrector ends near $\beta_W \approx 0.715$. With $u = q_h$ of degree two the largest values on the same face are 0.9305 at $r = 1.44$, 0.9729 at $r = 1.50$ and 1.0018 at $r = 1.54$ (Figure 7), close to the Galerkin values. Lemma 7.3 therefore costs little at these couplings, and most of the distance between the floating-point value at $r = 1.5$ and the certified 0.999887 of Proposition 7.8 is the interpolation over boxes. Between $\beta_W = \frac{3}{4}$ and 0.77 there is room only for smaller boxes. The plaquette blocks of Theorem 1.7 use $\Lambda(2\beta_W)$ as well, with a little more room: they reach 0.77 with the certified cover of Proposition 7.9 and near 0.78 in floating point (Remark 8.4); past that one needs larger blocks, whose sensitivity does not go through the one-link modulus.

Other groups. For groups other than SU(2) and SU(3) only the curvature route is carried out here (Theorem 1.5), and it stops where the bound $\frac{29}{2}$ uses up the curvature of G, at

$N|\beta| = \frac{2}{29}\kappa_G$. The Dobrushin route needs a bound for the Kantorovich modulus of the one-link measure of each group, which is not attempted here.

The four open problems. Our earlier paper [15] listed four problems between the strong-coupling results and the Clay problem [50]: (1) uniqueness and a gap at intermediate fixed coupling; (2) the crossover from strong coupling to Bałaban's weak-coupling regime; (3) the continuum limit with the Osterwalder–Schrader axioms; and (4) a gap in physical units. The present paper moves the boundary of (1): for SU(2) from $\beta_W = \frac{2}{9}$ to 0.351, for SU(3) from $\frac{9}{32}$ to 0.77, with a gap at $\frac{3}{4}$ of 0.0177 in lattice units, and for every compact simple group, the exceptional groups included, past the curvature ceiling of the ungauged measure, with an explicit gap. It does not close (1); the region between these values and the scaling region, near $\beta_W \approx 2.2$ for SU(2) and $\beta_W \approx 6$ for SU(3), is open, and with it problem (2). Problems (3) and (4) are not solved here. Every bound proved here is uniform in lattice units, so by [16] none of them can by itself produce a continuum mass gap; that requires control of the gap in physical units along the renormalisation group flow [6, 18].[15]

The continuum problem. Two general facts limit what can be imported into problems (3) and (4) above from constructions of continuum measures. Both concern a random distribution F on $\mathbb{R}^d$, a Borel probability measure on the tempered distributions with finite moments, and the growth condition

$$\mathbb{E}\, e^{F(\chi)} \le e^{C(\|\chi\|_1 + \|\chi\|_2^2)} \qquad \text{for all real } \chi \in \mathcal{S}(\mathbb{R}^d), \tag{9.1}$$

which is the growth condition of the regularity axiom of Glimm and Jaffe at $p = 2$ [41, Sect. 6.1, (6.1.5)]. The next proposition shows that it rules out every field whose two-point function is too singular at short distance.

Proposition 9.1. *Let F satisfy (9.1), and suppose that its connected two-point function is, off the diagonal, a locally integrable function $S_2(x,y)$ with $S_2(x,y) \ge c|x-y|^{-2\Delta}\big(\log \frac{1}{|x-y|}\big)^{-k}$ for $0 < |x-y| \le r_0$, where $c > 0$, $k \ge 0$ and $2\Delta > d$. Then F does not exist: no random distribution has both properties.*

Proof. Applying (9.1) to χ and $-\chi$ and using $\cosh x \ge 1 + \frac{x^2}{2}$ gives $\mathbb{E}F(\chi)^2 \le 2(e^{C(\|\chi\|_1+\|\chi\|_2^2)} - 1)$. Fix nonnegative $\varphi, \psi \in C_c^\infty(\mathbb{R}^d)$, not zero, with supports in a ball of radius $\frac{R}{2}$ and at distance $\delta > 0$ from each other, and put $\varphi_\varepsilon(x) = \varepsilon^{-d/2}\varphi(x/\varepsilon)$, and likewise ψ_ε. Then $\|\varphi_\varepsilon\|_2 = \|\varphi\|_2$ and $\|\varphi_\varepsilon\|_1 = \varepsilon^{d/2}\|\varphi\|_1 \le \|\varphi\|_1$ for $\varepsilon \le 1$, so the covariance of $F(\varphi_\varepsilon)$ and $F(\psi_\varepsilon)$ is bounded uniformly in ε by the Cauchy–Schwarz inequality. On the other hand the supports of φ_ε and ψ_ε are disjoint, and for u, v in the supports $\varepsilon\delta \le |\varepsilon u - \varepsilon v| \le \varepsilon R$. For $\varepsilon R \le r_0$ and $\varepsilon R < 1$ the lower bound for S_2 therefore gives

$$\begin{aligned}\operatorname{Cov}\big(F(\varphi_\varepsilon), F(\psi_\varepsilon)\big) &= \varepsilon^d \iint \varphi(u)\psi(v) S_2(\varepsilon u, \varepsilon v)\, du\, dv \\ &\ge c\, R^{-2\Delta} \varepsilon^{d-2\Delta} \Big(\log \frac{1}{\varepsilon\delta}\Big)^{-k} \|\varphi\|_1 \|\psi\|_1,\end{aligned}$$

which tends to infinity as $\varepsilon \to 0$. □

The action density $\operatorname{Tr} F_{\mu\nu}F_{\mu\nu}$ has dimension $\Delta = 4$ in four dimensions. A two-sided short-distance law of the form $|x-y|^{-8}$ times the fourth power of the running coupling, which is what asymptotic freedom[16] predicts, has $2\Delta = 8 > 4$ and $k = 2$; so that field cannot satisfy (9.1), and a proof of rotation invariance for it has to use something other

[15] Every one of these approaches needs an input at physical length scales. Renormalisation group approaches to the ultraviolet behaviour of gauge and gravitational theories, of which [6, 29] are the rigorous ones for Yang–Mills theory, have also been proposed for the effective dimension of spacetime [70] and for completions of quantum gravity [12, 17]; they do not bear on the infrared estimates needed here.

[16] For SU(N) the renormalisation group gives, with $b_0 = 11N/(48\pi^2)$ and the invariant scale Λ,

$$g(a)^{-2} = 2b_0 \log \tfrac{1}{a\Lambda} + O\big(\log\log \tfrac{1}{a\Lambda}\big),$$

so the inverse coupling $\beta_W = 2N/g(a)^2$ grows like $\frac{11N^2}{12\pi^2} \log \frac{1}{a\Lambda}$ along the continuum tuning.

than this condition. In three dimensions the free part of the action density already has $2\Delta = 6 > 3$. Condition (9.1) suits fields of dimension at most $\frac{d}{2}$, such as the field strength of the free Maxwell field, whose covariance is bounded in momentum space.

Positivity, symmetry and regularity do not produce a gap by themselves.

Proposition 9.2. *On $\mathbb{R}^4$ let F be the centred Gaussian random distribution with covariance*

$$\widehat{C}(p) = \int_0^1 \frac{m^2}{|p|^2 + m^2}\, dm.$$

Then F is invariant under all Euclidean motions, reflection positive, satisfies (9.1) with $C = \frac{1}{2}$, and its Osterwalder–Schrader Hamiltonian has no gap above the vacuum.

Proof. Since $0 \le \widehat{C} \le 1$, $\mathbb{E}e^{F(\chi)} = e^{\frac{1}{2}\langle \chi, C\chi\rangle} \le e^{\frac{1}{2}\|\chi\|_2^2}$, with the unitary Fourier transform. Invariance is clear. The covariance is the integral over $m \in (0,1)$, with the weight $m^2 dm$, of the covariances of free fields of mass m, each of which is reflection positive [41, Sect. 6.2]; a positive combination of reflection positive covariances is reflection positive, and so is the Gaussian measure with such a covariance [41, Sect. 6.2]. For $\chi(t,\mathbf{x}) = h(t)g(\mathbf{x})$ with $h \ge 0$ supported in $t > 0$, not zero, and $\hat{g}(0) \ne 0$, the vector $v = F(\chi)$ is orthogonal to the vacuum, and after the Fourier transform in $\mathbf{x}$,

$$\langle v, e^{-sH} v\rangle = \int_0^1 m^2\, dm \int \frac{|\hat{g}(\mathbf{k})|^2}{2\omega} e^{-\omega s} \times \Big(\int_0^\infty h(t) e^{-\omega t} dt\Big)^2 d\mathbf{k},$$

with $\omega = \sqrt{|\mathbf{k}|^2 + m^2}$ and constants absorbed in $\hat{g}$. If H had a gap $m_0 > 0$, the left side would be at most $e^{-m_0 s}\|v\|^2$. For $0 < \eta < m_0$ the part of the integral with $m, |\mathbf{k}| \le \eta/2$ is at least $c_\eta e^{-\eta s}$ with $c_\eta > 0$, a contradiction as $s \to \infty$. □

The Gaussian field of Proposition 9.2 is not a gauge theory; it shows only that an infrared estimate, such as a bound $\langle v_a, T_a^{\lfloor t/a\rfloor} v_a\rangle \le C_v e^{-m_* t}$ uniform in the spacing a along the continuum tuning, has to be proved in addition to the axioms. The lattice bounds of this paper are of this form only with m_* proportional to $1/a$, and by [16] such bounds give trivial limits. The routes we know to a continuum gap therefore all stop at the same two estimates: the construction of the infinite-volume continuum measure past Bałaban's small-field analysis [6, 8], and a decay estimate in physical units at large scales.

10. The continuum step at one physical scale

The bounds of the preceding sections hold at fixed coupling and give a gap that is at least a fixed number in lattice units, and [16] shows that no such bound can produce a continuum gap. This section replaces the single-link condition of Proposition 3.2 by a condition on one cube of n^4 sites, averaged over its translates in the manner of Dobrushin and Shlosman [32], and proves that the condition gives a gap of order $1/n$ in lattice units. If it holds with n proportional to $1/a$, that is, in a cube of fixed physical size, the gap survives in physical units and passes to every continuum limit (Theorem 10.4). The two statements that remain unproved, the cube estimate and the existence of the limit, are stated in Section 11.

10.1. **Averaged cubes.** For a finite set $V \subset A$ let $\gamma_V(\cdot \mid Q)$ be the probability measure on G^V with density proportional to $\exp\big(N\beta \sum_{p \cap V \ne \emptyset} \operatorname{Re}\operatorname{Tr} Q_p^F\big)$ with respect to Haar measure, the links outside V being those of Q. Its conditional law of Q_e, $e \in V$, given the other links of V is γ_e^F. Write $d_V(Q_V, Q'_V) = \sum_{i \in V} \delta(Q_i, Q'_i)$ and, for $j \in A \setminus V$,

$$C_{V,j} = \sup \frac{W_1^{d_V}\big(\gamma_V(\cdot \mid Q), \gamma_V(\cdot \mid Q')\big)}{\delta(Q_j, Q'_j)},$$

(a) averaged cubes, $n = 4$ (b) one physical scale




FIGURE 8. (a) The cube V of side $n = 4$ and two of its translates by $2\mathbb{Z}^4$, drawn in three of the four dimensions. A link j outside V enters K_c through every translate whose boundary it touches, and $D(n, \alpha)$ compares that total with the $N_n = (n/2)^4$ translates that contain j. (b) The cube of Theorem 10.4: its physical side $\ell = na$ is held fixed while the spacing is halved, so n doubles at each step and the lattice gap $\log(1/\alpha)/(2n)$ is of order a/ℓ.

the supremum over pairs that differ only at j; $W_1^{d_V}$ is the Kantorovich distance for d_V. The coefficient vanishes unless j shares a plaquette with a link of V.

Let $n \geq 2$ be even and let V be the set of active links (x, μ) with $x \in [0, n)^4$. The direction field has period two, so the translations by $2\mathbb{Z}^4$ preserve F, A and γ^F, and the translates $V_x = V + x$, $x \in 2\mathbb{Z}^4$, satisfy $C_{V_x, j+x} = C_{V,j}$. Every active link lies in exactly $N_n = (n/2)^4$ of them. A *class* is an orbit of $2\mathbb{Z}^4$ on A, that is, a residue of the base point modulo two together with a direction; there are at most 64 classes. For a class c put

$$K_c = \sum_{j \in c \setminus V} C_{V,j}.$$

Definition 10.1. For $0 < \alpha < 1$, the condition $D(n, \alpha)$ at the coupling β is $K_c \leq \alpha N_n$ for every class c.

Since $\sum_{x \in 2\mathbb{Z}^4,\, j \notin V_x} C_{V_x, j} = \sum_x C_{V, j-x}[j - x \notin V] = K_{c(j)}$ for the class $c(j)$ of j, the condition says that the total influence of a link on the translates that do not contain it is at most the fraction α of the number of translates that do. Figure 8 shows the cubes.

For $n = 2$ it is a bound on blocks of 48 links. Distances between links are those of their midpoints; a link of V and a link sharing a plaquette with V are at distance at most $\rho_n = 2n$, and c_n denotes the number of active links within distance ρ_n of a given one.

Theorem 10.2. *Let $\beta \geq 0$ and suppose that $D(n, \alpha)$ holds. Then statements* (i)–(iv) *of Proposition 3.2 hold,* (v) *holds for every $\lambda < \log(1/\alpha)/\rho_n$, and the transfer matrix of ω has spectral gap at least $\log(1/\alpha)/(2n)$.*

Proof. The proof of Proposition 3.2 uses the coefficients $C_{ee'}$ only through Lemma B.1 and Corollary B.2. Lemma B.4 of Appendix B gives the same conclusions under $D(n, \alpha)$, for every $\lambda > 0$ with $\alpha e^{\lambda \rho_n} < 1$, with the constant $\theta D/(1-\theta)$ replaced by $D(1 + c_n e^{\lambda \rho_n})/(1-\theta)$ and $\theta = \alpha e^{\lambda \rho_n}$. The measures used there are invariant under every γ_{V_x} with V_x disjoint from the exceptional set: a Gibbs measure of γ^F and its conditional measures given the

links of a finite set by Lemma B.4(a), and the extended finite-volume measure of part (ii) because every plaquette that meets such a V_x lies in $P(\Lambda)$. Part (vi) then follows from Proposition F.1 as before, with every $\lambda < \log(1/\alpha)/(2n)$. □

10.2. **The continuum limit.** Let $a_k = 2^{-k}a_0$ and let $\beta_k \geq 0$ be couplings, typically with $\beta_k \to \infty$. Suppose that the lattice states ω_k at β_k exist as in Proposition 3.2, with transfer matrices $\mathbb{T}_k$ on the reconstructed spaces $\mathcal{H}_k$ and vacua Ω_k (Appendix F). Let $\mathcal{T}$ be a complex vector space of test functions on $\mathbb{R}^4$ supported in $\{t > 0\}$ and, for each k, let $f \mapsto O_k(f)$ be a linear map from $\mathcal{T}$ to gauge-invariant local observables supported in positive lattice times; the example to keep in mind is $O_k(f) = Z_k \sum_x a_k^4 f(a_k x)\Phi(x)$ for a local gauge-invariant field Φ, a renormalisation constant Z_k and $\Phi(x)$ the translate of Φ to x. Write ϑ for the time reflection and τ_t for the time translation by t lattice steps.

Definition 10.3. A *continuum limit* of (ω_k, O_k) is a Hilbert space $\mathcal{H}$ with a unit vector Ω, a self-adjoint operator $H \geq 0$ with $H\Omega = 0$, and a linear map $f \mapsto \psi(f)$ from $\mathcal{T}$ to $\mathcal{H}$ with dense range, such that for all $f, g \in \mathcal{T}$ and all dyadic $\tau \geq 0$,

$$\langle \psi(f), e^{-\tau H}\psi(g)\rangle = \lim_{k\to\infty} \omega_k\big(\vartheta(\overline{O_k(f)})\, \tau_{\tau/a_k} O_k(g)\big),$$
$$\langle \Omega, \psi(f)\rangle = \lim_{k\to\infty} \omega_k\big(O_k(f)\big),$$

the limits taken over the k for which τ/a_k is an integer. It is *nontrivial* if $\mathcal{H} \neq \mathbb{C}\Omega$.

The Osterwalder–Schrader reconstruction [65, 66] produces such a limit from convergent renormalised Schwinger functions that satisfy the axioms; the definition records only what the next theorem uses.

Theorem 10.4. *Suppose that there are $\alpha < 1$, $\ell > 0$ and even n_k with $n_k a_k \leq \ell$ such that $D(n_k, \alpha)$ holds at β_k for every k. Then every continuum limit of (ω_k, O_k) satisfies*

$$\sigma(H) \subset \{0\} \cup [m_*, \infty), \qquad m_* = \frac{\log(1/\alpha)}{2\ell},$$

and 0 is a simple eigenvalue. If the limit is nontrivial, $\sigma(H) \cap [m_*, \infty) \neq \emptyset$, so H has a mass gap of at least m_*.

Proof. By Theorem 10.2 the state ω_k exists and $\mathbb{T}_k$ has gap at least $\log(1/\alpha)/(2n_k)$ on the complement of Ω_k. Let $f \in \mathcal{T}$, put $O = O_k(f)$ and $\varphi_k = \Psi_O - \omega_k(O)\Omega_k$. In the reconstruction of Appendix F, $\langle \Psi_F, \mathbb{T}_k^t \Psi_H\rangle = \omega_k(\vartheta(\bar{F})\tau_t H)$, so $\langle \Omega_k, \Psi_O\rangle = \omega_k(O)$, and $\mathbb{T}_k\Omega_k = \Omega_k$; hence, for integers $t \geq 0$,

$$\begin{aligned}\langle \varphi_k, \mathbb{T}_k^t \varphi_k\rangle &= \omega_k\big(\vartheta(\bar{O})\, \tau_t O\big) - |\omega_k(O)|^2 \\ &\leq e^{-t\log(1/\alpha)/(2n_k)}\, \|\varphi_k\|^2.\end{aligned}$$

Take $\tau \geq 0$ dyadic and k large, so that $t = \tau/a_k$ is an integer; then $t/(2n_k) = \tau/(2n_k a_k) \geq \tau/(2\ell)$, and the right side is at most $e^{-m_*\tau}\|\varphi_k\|^2$. Both sides converge by Definition 10.3. With $\varphi = \psi(f) - \langle \Omega, \psi(f)\rangle\Omega$ and $e^{-\tau H}\Omega = \Omega$ the limit reads

$$\langle \varphi, e^{-\tau H}\varphi\rangle \leq e^{-m_*\tau}\|\varphi\|^2.$$

The maps $f \mapsto O_k(f)$ and $f \mapsto \psi(f)$ are linear, so the vectors φ form a linear subspace, and it is dense in $\Omega^\perp$ because the range of ψ is dense. The inequality extends to $\Omega^\perp$ by continuity, and from dyadic τ to all $\tau \geq 0$ by the strong continuity of $e^{-\tau H}$. For $\varphi \in \Omega^\perp$ with spectral measure μ_φ of H this says $\int e^{-\tau E} d\mu_\varphi(E) \leq e^{-m_*\tau}\mu_\varphi([0,\infty))$ for all $\tau \geq 0$; if $\mu_\varphi([0, m_* - \eta]) > 0$ for some $\eta > 0$, the left side is at least $\mu_\varphi([0, m_* - \eta])e^{-(m_* - \eta)\tau}$, which exceeds the right side for large τ. So μ_φ lives on $[m_*, \infty)$ for every $\varphi \in \Omega^\perp$, which gives both claims about 0, and the spectrum on $\Omega^\perp \neq \{0\}$ lies in $[m_*, \infty)$ and is not empty. □

The proof uses $D(n_k, \alpha)$ only through the decay of the two-point functions, and the same argument accepts any decay with constants that depend on the observable.

Theorem 10.5. *Let $m_0 > 0$ and let $\mathcal{T}_0$ span $\mathcal{T}$. Suppose that for each $f \in \mathcal{T}_0$ there are $C_f < \infty$, $\tau_f \geq 0$ and k_f such that*

$$C_k^f(\tau) = \omega_k(\vartheta(\overline{O_k(f)})\,\tau_{\tau/a_k} O_k(f)) - |\omega_k(O_k(f))|^2 \leq C_f e^{-m_0\tau}$$

for all $k \geq k_f$ and all dyadic $\tau \geq \tau_f$ with $\tau/a_k \in \mathbb{Z}$. Then every continuum limit of (ω_k, O_k) satisfies $\sigma(H) \subset \{0\} \cup [m_0, \infty)$, and 0 is a simple eigenvalue. If the limit is nontrivial, H has a mass gap of at least m_0.

Proof. For $f \in \mathcal{T}$ put $\varphi(f) = \psi(f) - \langle \Omega, \psi(f)\rangle \Omega$, which is linear in f. Since $e^{-\tau H}$ is self-adjoint and fixes Ω, Definition 10.3 gives $\langle \varphi(f), e^{-\tau H}\varphi(f)\rangle = \lim_k C_k^f(\tau)$. For $f \in \mathcal{T}_0$ let μ be the spectral measure of H at $\varphi(f)$. Then $\int e^{-\tau E} d\mu(E) \leq C_f e^{-m_0\tau}$ for dyadic $\tau \geq \tau_f$. If $\mu([0, m_0 - \eta]) > 0$ for some $\eta \in (0, m_0)$, the left side is at least $\mu([0, m_0 - \eta]) e^{-(m_0-\eta)\tau}$, which is impossible for large τ. Hence the spectral projection $E_{[0,m_0)}$ of H vanishes on $\varphi(\mathcal{T}_0)$. By linearity it vanishes on $\varphi(\mathcal{T})$, and since its kernel is closed, it vanishes on the closure of $\varphi(\mathcal{T})$, which is $\Omega^\perp$ because $\psi(\mathcal{T})$ is dense. So $E_{[0,m_0)} = \langle \Omega, \cdot \rangle\, \Omega$, which gives both claims. □

Theorem 10.4 is the case $m_0 = m_*$, $\tau_f = 0$ and $C_f = \sup_{k \geq k_f} \|\varphi_k\|^2$, which is finite because $\|\varphi_k\|^2 = C_k^f(0)$ converges. Theorem 10.5 needs neither a lattice gap nor the gauge of this paper. It applies to the output of a cluster expansion of any coarse effective theory, provided the expansion bounds the correlations of a spanning set of renormalised observables in physical units.

Theorem 10.4 bypasses the obstruction of [16]: under the hypothesis the lattice gap is at least $\log(1/\alpha)/(2n_k)$, of order a_k/ℓ in lattice units, which tends to zero, and only its value in physical units is bounded below. In the language of [16] the hypothesis implies the gap part (R2) of Hypothesis R, with $m = m_*$, and the conclusion holds for every continuum limit without a separate assumption on the gap.

10.3. **Smoothed observables.** The observables $O_k(f)$ of Definition 10.3 need not be local. The renormalised fields that one expects to converge are smoothed by the gradient flow of the Wilson action [58][17]; at positive flow time their correlation functions are finite to all orders of perturbation theory once the coupling is renormalised [59]. Flow in all four directions mixes the two sides of the reflection plane, but the spatial flow of [71, Sec. II.B] does not: it flows the links of each time slice under the Wilson action of that slice and leaves the temporal links alone.

Fix a bi-invariant Riemannian metric on G. For a configuration U of all links of $\mathbb{Z}^4$ (for the states of this paper, the links of F equal to 1), let $\Phi_s(U)$, $s \in \mathbb{R}$, solve

$$\frac{d}{ds} U_e = -\nabla_e \sum_{p \ni e,\ p \text{ spatial}} (N - \operatorname{Re}\operatorname{Tr} U_p)$$

for every spatial link e, the sum running over the four plaquettes of the time slice of e that contain it, with the temporal links held fixed and ∇_e the gradient in the variable U_e.

Proposition 10.6. (i) *Φ_s is defined on every configuration of the infinite lattice, $\Phi_s \circ \Phi_{s'} = \Phi_{s+s'}$, and Φ_s commutes with gauge transformations, with spatial and temporal translations and with the reflections in time. The link $\Phi_s(U)_e$ depends only on the spatial links of the time slice of e, and there are constants κ, c depending only on G such that, if Φ_s^R denotes*

[17] The flow moves the gauge field along the negative gradient of the Wilson action. At positive flow time the flowed fields are smooth on the scale of the square root of the flow time, and in perturbation theory their gauge-invariant composite fields need no further renormalisation [58, 59].

the flow of the links of that slice within distance R of e with all other links held fixed,

$$\delta\big(\Phi_s(U)_e, \Phi_s^R(U)_e\big) \le D\, e^{\kappa|s|}\, \frac{(c|s|)^{R+1}}{(R+1)!}$$

for every configuration U.

(ii) *Let ω be a state as in Appendix F and O a bounded gauge-invariant function of finitely many links at times in $[t_0, t_1]$, continuous in them. Then $O \circ \Phi_s$ is a bounded gauge-invariant function of the links at times in $[t_0, t_1]$ and a uniform limit of gauge-invariant local functions at those times.*

(iii) *Theorem 10.4 holds with $O_k(f)$ replaced by $O_k(f) \circ \Phi_{s_k}$ for any $s_k \in \mathbb{R}$, in particular for $s_k = \tau/a_k^2$ at a fixed flow time τ in physical units.*

(iv) *The image measure $(\Phi_s)_*\omega$ is reflection positive, and its transfer matrix is unitarily equivalent to that of ω, the unitary fixing the vacuum.*

Proof. (i) Embed G in a matrix algebra. The right side of the equation is a smooth function of the links of the four plaquettes containing e, tangent to G at U_e, so on the Banach space of bounded assignments of matrices to the spatial links it is a vector field that is bounded and Lipschitz on a neighbourhood of the configurations with values in G, with a Lipschitz constant that does not depend on the volume because every link meets finitely many others. The Picard–Lindelöf theorem gives a unique local solution; it stays on G because the field is tangent there, and since the field is bounded on G the solution exists for all s; uniqueness gives the group law. The action of a time slice and the metric are invariant under gauge transformations, translations and the reflections, so the flow commutes with them by uniqueness. Since the equation of a spatial link involves only links of its own slice, so does the solution. For the comparison, let $d_{e'}(s)$ be the distance between the two flows at a link e' of the slice. Smoothness on the compact group gives constants κ_0, c_1 with $d_{e'}(s) \le \int_0^{|s|} (\kappa_0 d_{e'} + c_1 \sum_{e'' \sim e'} d_{e''})$ for e' within distance R of e, where $e'' \sim e'$ means that the two links share a spatial plaquette (twelve neighbours), while $d_{e'} \le D$ beyond. Iterating the integral form of this inequality, a term that reaches the links beyond distance R needs at least $R+1$ steps of $\sim$, and $\sum_{k>R} x^k/k! \le e^x x^{R+1}/(R+1)!$; this gives the bound with $c = 12c_1$ and $\kappa = \kappa_0 + c$.

(ii) The truncated flow Φ_s^R commutes with gauge transformations for the same reason as Φ_s, so $O \circ \Phi_s^R$ is a gauge-invariant function of finitely many links at times in $[t_0, t_1]$. By (i) and the uniform continuity of O, it converges to $O \circ \Phi_s$ uniformly as $R \to \infty$.

(iii) The proof of Theorem 10.4 uses $O_k(f)$ only through the vectors $\Psi_{O_k(f)}$ and the identity $\langle \Psi_F, \mathbb{T}^t \Psi_H \rangle = \omega(\vartheta(\bar{F})\tau_t H)$. By (ii) and the footnote in Appendix F, both extend to uniform limits of local functions at positive times, and $f \mapsto O_k(f) \circ \Phi_{s_k}$ is linear.

(iv) By (i), Φ_s and Φ_{-s} preserve the functions of the links at positive times and commute with ϑ and τ_t. Hence $(\Phi_s)_*\omega\big(\vartheta(\bar{F})\tau_t H\big) = \omega\big(\vartheta(\overline{F \circ \Phi_s})\, \tau_t(H \circ \Phi_s)\big)$, which for $t = 0$ and $H = F$ is nonnegative, and $F \mapsto F \circ \Phi_s$ induces an isometry between the reconstructed spaces that intertwines the transfer matrices and fixes the vacuum; Φ_{-s} gives its inverse. □

Part (iv) says that smoothing changes the variables and not the theory: it neither creates nor removes a gap. Its use is in the choice of the observables. With $O_k(f)$ the spatially flowed fields at a fixed physical flow time, for instance $Z_k \sum_x a_k^4 f(a_k x)(\mathcal{E}_\tau(x) - \omega_k(\mathcal{E}_\tau(x)))$ with $\mathcal{E}_\tau(x)$ the flowed action density of the slice, statement (E) asks for the convergence of correlation functions that are known to be finite in perturbation theory [59], and Theorem 10.4 applies to them. Theorem 10.7 below constructs their continuum limits along subsequences; their nontriviality is not proved here.

10.4. **Renormalised flowed observables.** The spatially flowed fields can be renormalised by one condition at a single physical separation, and their correlation functions then have continuum limits along subsequences without further assumptions. What remains open is




FIGURE 9. The packet f_0 of Theorem 10.7, supported in the time window $[t_0, t_1]$ above the reflection plane $t = 0$, its reflection ϑf_0 and its translate $\tau_\sigma f_0$ by σ in time. The correlation $C_k(\sigma)$ pairs the reflected packet with the translated one; $C_k(0)$ is the reflected norm that renormalises the packet.

whether such a limit is trivial, and that question reduces to one ratio of lattice two-point functions.

Fix a bounded gauge-invariant function Φ of finitely many spatial links of the slice $\{x_0 = 0\}$, continuous in them, a flow time $\tau > 0$ in physical units, and a nonzero test function $f_0 \in C_c(\mathbb{R}^4)$ supported in $\{t > 0\}$. Put $s_k = \tau/a_k^2$, write Φ_x for the translate of Φ to $x \in \mathbb{Z}^4$, and let

$$A_k = \sum_{x \in \mathbb{Z}^4} a_k^4 f_0(a_k x)\big(\Phi_x \circ \Phi_{s_k} - \omega_k(\Phi_x \circ \Phi_{s_k})\big),$$

$$C_k(\sigma) = \omega_k\big(\vartheta(\bar{A}_k)\, \tau_{\sigma/a_k} A_k\big)$$

for dyadic $\sigma \geq 0$ (the sum is finite, and σ/a_k is an integer for large k). By reflection positivity $C_k(0) \geq 0$; assume $C_k(0) > 0$ and set $Z_k = C_k(0)^{-1/2}$. Let $\mathcal{D}$ be the set of points $y = (t, y') \in \mathbb{R}^4$ with dyadic coordinates (in units of a_0) and $t > 0$, and for $y \in \mathcal{D}$ and k with $y/a_k \in \mathbb{Z}^4$ let A_k^y be the translate of A_k by y/a_k.

Theorem 10.7. (i) *For $y, y' \in \mathcal{D}$ and every k with $y/a_k, y'/a_k \in \mathbb{Z}^4$,*

$$Z_k^2 \left|\omega_k\big(\vartheta(\bar{A}_k^y) A_k^{y'}\big)\right| \leq 1.$$

(ii) *Every sequence of k has a subsequence along which $K(y, y') = \lim Z_k^2\, \omega_k\big(\vartheta(\bar{A}_k^y) A_k^{y'}\big)$ exists for all $y, y' \in \mathcal{D}$ and $\rho(\sigma) = \lim C_k(\sigma)/C_k(0)$ exists for all dyadic $\sigma > 0$. Along it there is a probability measure μ on $[0, \infty]$ with*

$$\rho(\sigma) = \int_{[0,\infty]} e^{-\sigma E}\, d\mu(E) \qquad (\sigma > 0),$$

the integrand being 0 at $E = \infty$.

(iii) *Along such a subsequence, let $\mathcal{T}$ be spanned by the translates $f_0(\cdot - y)$, $y \in \mathcal{D}$, and one more function $f_* \in C_c(\mathbb{R}^4)$ supported in $\{t > 0\}$ outside their span, and let O_k be the linear map with $O_k(f_0(\cdot - y)) = Z_k A_k^y$ and $O_k(f_*) = 1$. Then (ω_k, O_k) has a continuum limit in the sense of Definition 10.3, on which the dyadic spatial translations act by unitary operators commuting with H.*

(iv) *This limit is nontrivial if and only if $\mu([0,\infty)) > 0$, if and only if $\rho(\sigma) > 0$ for one dyadic $\sigma > 0$, and then $\rho(\sigma) > 0$ for all of them. Consequently a nontrivial limit exists along some subsequence if and only if $\limsup_k C_k(\sigma)/C_k(0) > 0$ for some dyadic $\sigma > 0$.*

(v) *If the hypothesis of Theorem 10.4 holds, then μ is carried by $[m_*, \infty]$ and $\rho(\sigma) \le e^{-m_*\sigma}$.*

Proof. By Proposition 10.6(ii) each $\Phi_x \circ \Phi_{s_k}$, and hence each A_k^y, is a uniform limit of gauge-invariant local functions at times ≥ 1, so the reconstruction of Appendix F applies to them (its footnote). Write $\Psi_k = Z_k \Psi_{A_k}$. Since $\langle \Omega_k, \Psi_{A_k}\rangle = \omega_k(A_k) = 0$ and $\|\Psi_k\|^2 = Z_k^2 C_k(0) = 1$, Ψ_k is a unit vector orthogonal to Ω_k. The relation $\langle \Psi_F, \mathbb{T}_k^t \Psi_H\rangle = \omega_k(\vartheta(\bar F)\tau_t H)$ for all F gives $\Psi_{\tau_t H} = \mathbb{T}_k^t \Psi_H$. Spatial translations commute with ϑ and with τ_t and leave ω_k invariant, so they act on the reconstructed space by unitaries $U_k(x)$ that commute with $\mathbb{T}_k$. Hence, for $y = (t, y')$,

$$Z_k \Psi_{A_k^y} = U_k(y'/a_k)\, \mathbb{T}_k^{t/a_k} \Psi_k .$$

(i) The left side of the inequality is $|\langle Z_k \Psi_{A_k^y}, Z_k \Psi_{A_k^{y'}}\rangle| \le 1$, because $\|U_k\| = 1$ and $0 \le \mathbb{T}_k \le 1$.

(ii) The set $\mathcal{D} \times \mathcal{D}$ is countable and the numbers are bounded by (i), so a diagonal subsequence gives all the limits K; the limits $\rho(\sigma)$ are among them, since $C_k(\sigma)/C_k(0) = \langle \Psi_k, \mathbb{T}_k^{\sigma/a_k} \Psi_k\rangle$ is the value of K at $y = y' = (\sigma/2, 0)$. Let ν_k be the spectral measure (Lemma G.1) of $\mathbb{T}_k$ at Ψ_k, a probability measure on $[0,1]$, and μ_k its image under $\lambda \mapsto -a_k^{-1} \log \lambda \in [0,\infty]$. Then $C_k(\sigma)/C_k(0) = \int e^{-\sigma E} d\mu_k(E)$. The space $[0,\infty]$ is compact, so a further subsequence has $\mu_k \to \mu$ weakly (Lemma G.2), and $E \mapsto e^{-\sigma E}$ is continuous on $[0,\infty]$ for $\sigma > 0$.

(iii) A finite combination $\sum_j c_j f_0(\cdot - y_j)$ with distinct y_j that vanishes has Fourier transform $\hat f_0(p) \sum_j c_j e^{-ip\cdot y_j} = 0$; since $\hat f_0$ is analytic and not zero, the exponential sum vanishes on an open set and every c_j is zero. So the translates are linearly independent and O_k is well defined. On the space V of finite sums $v = \sum_y c_y \delta_y$ over $\mathcal{D}$ put $B(v,w) = \sum \bar c_y c'_{y'} K(y,y')$. It is the limit of $\langle v_k, w_k\rangle$ with $v_k = \sum_y c_y Z_k \Psi_{A_k^y}$, hence positive semidefinite; let $\mathcal{H}_0$ be the completion of V modulo its null space. For dyadic $\sigma \ge 0$ and dyadic spatial x define $T(\sigma)\delta_y = \delta_{y+(\sigma,0)}$ and $U(x)\delta_y = \delta_{y+(0,x)}$. At level k these are $\mathbb{T}_k^{\sigma/a_k}$ and $U_k(x/a_k)$, so in the limit $B(T(\sigma)v, w) = B(v, T(\sigma)w)$, $B(T(\sigma)v, T(\sigma)v) \le B(v,v)$, $B(v, T(\sigma)v) \ge 0$, and $U(x)$ preserves B. Thus $T(\sigma)$ extends to commuting positive contractions on $\mathcal{H}_0$ with $T(\sigma)T(\sigma') = T(\sigma+\sigma')$, and $U(x)$ to unitaries commuting with them. For dyadic $\sigma > 0$ the kernel of $T(\sigma) = T(\sigma/2)^2$ equals that of $T(\sigma/2)$, so all these kernels coincide; and every δ_y with $y = (t, y')$ equals $T(t/2)\delta_{(t/2,y')}$, a vector in the range of a positive operator and so orthogonal to its kernel. Hence each $T(\sigma)$ is injective on $\mathcal{H}_0$, and $H_0 = -\log T(1)$ is a self-adjoint operator with $H_0 \ge 0$ and $e^{-\sigma H_0} = T(\sigma)$ for dyadic σ, by the uniqueness of positive roots. Put $\mathcal{H} = \mathbb{C}\Omega \oplus \mathcal{H}_0$, $H = 0 \oplus H_0$, $\psi(f_0(\cdot - y)) = \delta_y$ and $\psi(f_*) = \Omega$. The range of ψ is dense, and the identities of Definition 10.3 hold: for two translates they are the limits defining K, with $\tau_{\sigma/a_k} A_k^{y'} = A_k^{y'+(\sigma,0)}$; and for f_* they read $\omega_k(1) = 1$ and $\omega_k(\vartheta(\bar A_k^y)) = \omega_k(\tau_{\sigma/a_k} A_k^y) = 0$.

(iv) The limit is nontrivial exactly when $\mathcal{H}_0 \ne 0$, that is when $\|\delta_y\|^2 = K(y,y) = \rho(2t)$ is positive for some $y = (t, y')$. By (ii) and Lemma G.3(b), $\rho(\sigma) > 0$ for one $\sigma > 0$ if and only if $\mu([0,\infty)) > 0$, and then $\rho(\sigma) > 0$ for every $\sigma > 0$. For the last statement choose a subsequence along which $C_k(\sigma)/C_k(0)$ tends to its lim sup, and refine it by (ii).

(v) Under that hypothesis the proof of Theorem 10.4 gives $\nu_k([0, e^{-\log(1/\alpha)/(2n_k)}]) = 1$ because $\Psi_k \perp \Omega_k$, so μ_k is carried by $[\log(1/\alpha)/(2n_k a_k), \infty] \subset [m_*, \infty]$, and so is the weak limit μ. □

The measure μ is the spectral measure of the renormalised flowed field in the limit: its part on $[0,\infty)$ is what survives as a state of the continuum theory, and its atom at ∞ is the weight that the field puts on energies that diverge in physical units. The theorem constructs

the renormalised continuum observables and their reconstruction for every tuning, and it reduces their nontriviality to the lattice inequality

$$\limsup_{k\to\infty} \frac{C_k(\sigma)}{C_k(0)} > 0$$

for one physical separation σ: the flowed two-point function of the smeared packet at a fixed physical distance must not become negligible against its reflected norm $C_k(0)$. If the lattice gap stayed above some $m > 0$ in lattice units, the ratio would be at most $e^{-m\sigma/a_k}$ and the limit would be trivial, so the inequality is a condition on the tuning $\beta_k \to \infty$; it is not proved here. Nor are the independence of the limit from the subsequence, Euclidean invariance, or bounds on products of more than two renormalised fields, for which (i) gives nothing because Z_k may diverge (Theorem 10.16 reduces them to tree bounds); (E) requires all of these.

The denominator $C_k(0)$ is the reflected norm of the packet smeared by f_0, not a coincidence limit of point fields, and the distinction matters. For the free massless scalar field in four dimensions with spatial momenta cut off at Λ, the covariance at spatial separation 0 is $(1-(1+\Lambda t)e^{-\Lambda t})/(4\pi^2 t^2)$ at time $t > 0$ and $\Lambda^2/(8\pi^2)$ at coincidence, so its ratio $2(1-(1+\Lambda t)e^{-\Lambda t})/(\Lambda t)^2$ tends to zero as $\Lambda \to \infty$, although the limiting field is the nontrivial Gaussian field; for packets smeared by f_0 the reflected norms of that field have finite positive limits and the ratio stays positive. The criterion of Theorem 10.7(iv) is about the smeared packets A_k only.

When the ratio is bounded below, renormalising at the separation σ instead of 0 removes the escape to infinite energy altogether, and one slab of the transfer matrix gives a sufficient condition for the bound. Write μ_k for the measures in the proof of Theorem 10.7.

Proposition 10.8. (i) *Suppose $C_k(\sigma) \geq c\, C_k(0)$ for a dyadic $\sigma > 0$, a $c > 0$ and every k of a sequence. Let μ'_k be the spectral measure, in energies $E = -a_k^{-1}\log\lambda$, of the unit vector $C_k(\sigma)^{-1/2}\Psi_{A'_k}$ with $A'_k = \tau_{\sigma/2a_k}A_k$. Then μ'_k is a probability measure on $[0,\infty)$,*

$$\mu'_k([\Lambda,\infty)) \leq c^{-1}e^{-\sigma\Lambda}, \qquad \int E^m\, d\mu'_k \leq c^{-1}\Big(\frac{m}{e\sigma}\Big)^m,$$

and $t \mapsto C_k(\sigma+t)/C_k(\sigma)$ is Lipschitz on $[0,\infty)$ with constant $(e\sigma c)^{-1}$. A subsequence of the μ'_k converges weakly to a probability measure on $[0,\infty)$ with the same bounds.

(ii) *Suppose $\langle\Psi_k, (1-\mathbb{T}_k)\Psi_k\rangle \leq Ca_k$ for a constant C and every k of a sequence. Then $C_k(\sigma)/C_k(0) \geq e^{-2C\sigma}$ whenever $Ca_k \leq \frac12$ and σ/a_k is an integer, and $\mu_k([\Lambda,\infty]) \leq C(a_k + \Lambda^{-1})$ for every $\Lambda > 0$.*

Proof. (i) We have $\Psi_{A'_k} = \mathbb{T}_k^{\sigma/2a_k}\Psi_{A_k}$, so the measure μ'_k is $e^{-\sigma E}d\mu_k(E)$ divided by its mass $C_k(\sigma)/C_k(0) \geq c$, and $e^{-\sigma E}$ vanishes at $E = \infty$. For $E \geq \Lambda$ we have $e^{-\sigma E} \leq e^{-\sigma\Lambda}$, and $E^m e^{-\sigma E} \leq (m/e\sigma)^m$ for all $E \geq 0$. The ratio in question is the integral of e^{-tE} against μ'_k, and the inequality $|e^{-tE} - e^{-sE}| \leq |t-s|E$ with the moment bound for $m = 1$ gives the Lipschitz bound. The tail bound makes the sequence tight on $[0,\infty)$, so Prokhorov's theorem gives the weak limit, and the bounds pass to it because $\{E > \Lambda\}$ is open and E^m is a monotone limit of bounded continuous functions.

(ii) Let ν_k be the spectral measure of $\mathbb{T}_k$ at Ψ_k on $[0,1]$, a probability measure, and $n = \sigma/a_k$. By Jensen's inequality for the convex function λ^n,

$$\frac{C_k(\sigma)}{C_k(0)} = \int \lambda^n d\nu_k \geq \Big(\int \lambda\, d\nu_k\Big)^n \geq (1-Ca_k)^n \geq e^{-2Ca_k n},$$

since $\log(1-x) \geq -2x$ on $[0,\frac12]$. Moreover $Ca_k \geq \int(1-e^{-a_k E})d\mu_k \geq (1-e^{-a_k\Lambda})\mu_k([\Lambda,\infty])$, and $1-e^{-x} \geq x/(1+x)$. □

Part (ii) asks for an upper bound on a one-slab Dirichlet form, the opposite direction to a mass gap, which bounds such forms from below. In the box of Proposition 10.9, with

g smooth, the time smearing supplies the factor a_k up to one ratio: in the notation of its proof, $\mathbb{T}_k = A$ and $\Psi_{A_k} = \sum_n w_n \mathbb{T}_k^n u$ with $w_n = a_k g(a_k n)$ and $u = a_k^3 F^c \psi_0$, and the identity $(1-\mathbb{T}_k)\Psi_{A_k} = \sum_n (w_n - w_{n-1})\mathbb{T}_k^n u$ gives $\langle \Psi_k, (1-\mathbb{T}_k)\Psi_k\rangle \le a_k \|g'\|_{L^1}\|u\|/\|\Psi_{A_k}\|$, so the hypothesis of (ii) holds when the vector u of a single slice stays comparable to the time-smeared packet. That comparison is not proved; Proposition 10.10 below bounds the ratio without it.

A one-slab bound of this kind also limits the mass that $D(n_k, \alpha)$ can supply. Let e^{-m_k} be the norm of $\mathbb{T}_k$ on $\Omega_k^\perp$. If a unit vector $\varphi_k \perp \Omega_k$ has $\langle \varphi_k, (1-\mathbb{T}_k)\varphi_k\rangle \le \kappa a_k < 1$, then the spectral measure of $\mathbb{T}_k$ at φ_k lives on $[0, e^{-m_k}]$, so $1 - e^{-m_k} \le \kappa a_k$ and $m_k \le -\log(1-\kappa a_k)$. If in addition $D(n_k, \alpha)$ holds with $n_k a_k \le \ell$, Theorem 10.2 gives $m_k \ge \log(1/\alpha)/(2n_k) \ge a_k \log(1/\alpha)/(2\ell)$, and therefore

$$\frac{\log(1/\alpha)}{2\ell} \le \frac{-\log(1-\kappa a_k)}{a_k} = \kappa + O(a_k).$$

So the physical mass m_* of Theorem 10.4 is at most the constant κ of any one-slab bound. The two hypotheses are compatible because both are statements in physical units. The one-slab bound forces $m_k \to 0$, so it cannot come from an expansion whose convergence gives a lattice gap bounded below in lattice units.

In a periodic spatial box the assumption $C_k(0) > 0$ is a theorem. Let $G = \mathrm{SU}(N)$ with its defining representation, and replace $\mathbb{Z}^4$ by $\mathbb{Z} \times (\mathbb{Z}/L_k\mathbb{Z})^3$ with $L_k \ge 3$ and ω_k by the infinite-time Wilson state of the box. The proof of Theorem 10.7 uses only reflection positivity and the transfer matrix once the spatial translations are dropped, and parts (i)–(iv) hold in the box with $\mathcal{D}$ replaced by its points $(t, 0)$, the time translates of f_0 being independent by the same Fourier argument in t.

Proposition 10.9. *In the box, let $f_0(t,x) = g(t)h(x)$ with $g, h \ge 0$, and let $\Phi = \sum_{i<j}(N - \operatorname{Re}\operatorname{Tr} Q_{ij})$ be the energy of the three spatial plaquettes at the origin. If $g(a_k n)h(a_k x) > 0$ for some $n \ge 1$ and some x, then $C_k(0) > 0$.*

Proof. Lüscher proved that the transfer matrix is strictly positive [57]; we recall the argument because the proof uses it in this form.

The one-link kernel. Write χ for the character of the defining representation and $b = \beta_W > 0$. Then

$$e^{\frac{b}{N}\operatorname{Re}\chi} = \sum_{m \ge 0} \frac{(b/2N)^m}{m!}(\chi + \bar\chi)^m,$$

and $(\chi + \bar\chi)^m$ is the character of $(\rho \oplus \bar\rho)^{\otimes m}$. Every irreducible representation σ occurs in one of these tensor powers: their matrix coefficients span the algebra of polynomials in the entries of q and $\bar q$, which contains the constants and separates points, so it is dense in $C(G)$ by the Stone–Weierstrass theorem, while the matrix coefficients of a σ that never occurred would be orthogonal to it by Schur orthogonality. So the coefficient of every irreducible character in $e^{\frac{b}{N}\operatorname{Re}\chi}$ is positive, and convolution with this function acts on each isotypic part of $L^2(G)$ by a positive scalar.

The transfer matrix. In the temporal gauge the transfer matrix of the box acts on the gauge-invariant part of $L^2(G^E)$, E the spatial links of one slice, as MKM, where K is the tensor product over E of the convolutions above and M is multiplication by the exponential of half the spatial plaquette weight [57, 67]. Both commute with gauge transformations, K is positive and injective by the first step, and M is positive and invertible, so $\langle v, MKMv\rangle = \langle Mv, KMv\rangle > 0$ for $v \ne 0$. Its kernel is a continuous positive function, and K is compact, so its largest eigenvalue λ_0 is simple with an eigenfunction $\psi_0 > 0$, and with $A = MKM/\lambda_0$, a function F of the slice at time $n \ge 0$ has $\Psi_F = A^n F \psi_0$ in the reconstruction, Ω being ψ_0.

Conclusion. Let $F = \sum_x h(a_k x)\,\Phi_x \circ \Phi_{s_k}$ and $F^c = F - \omega_k(F)$. Then $\Psi_{A_k} = a_k^4\, p(A) F^c \psi_0$ with $p(z) = \sum_{n \ge 1} g(a_k n) z^n$, a polynomial with nonnegative coefficients, not all zero, so $p > 0$

on $(0,1]$ and $p(A)$ is injective. The function $\sum_x h(a_k x)\Phi_x$ vanishes at the configuration with every link 1 and is positive at one with a single nontrivial link at a site where $h(a_k x) > 0$; the flow Φ_{s_k} is a bijection, so F is not constant, and as F is continuous and $\psi_0 > 0$, $F^c\psi_0 \neq 0$. Hence $C_k(0) = \|\Psi_{A_k}\|^2 > 0$. □

The proposition removes the one assumption of Theorem 10.7 in a box; it says nothing about the ratio of part (iv), on which nontriviality depends.

The ratio of Theorem 10.7(iv) can be compared with correlations of a single slice field. Suppose that $f_0(t,x) = g(t)h(x)$ with $g \geq 0$ supported in $[t_0,t_1]$, $t_0 < t_1$ positive and dyadic, let

$$U_k = \sum_{x\in\mathbb{Z}^3} a_k^3 h(a_k x)\big(\Phi_{(0,x)} \circ \Phi_{s_k} - \omega_k(\Phi_{(0,x)} \circ \Phi_{s_k})\big)$$

be the centred field of the slice at time 0, smeared in space only, and put $u_k = \Psi_{U_k}$ and

$$D_k(r) = \langle u_k, \mathbb{T}_k^{r/a_k} u_k\rangle = \omega_k\big(\vartheta(\bar U_k)\,\tau_{r/a_k} U_k\big)$$

for $r \in a_k\mathbb{Z}$, $r \geq 0$. Let $n_0 = \lceil t_0/a_k\rceil$ and $n_1 = \lfloor t_1/a_k\rfloor$.

Proposition 10.10. *Assume $C_k(0) > 0$, and let $\sigma \geq 0$ with $\sigma/a_k \in \mathbb{Z}$.*

(i) *$D_k(r) > 0$ for every r, and*

$$\frac{D_k(2a_kn_1+\sigma)}{D_k(2a_kn_0)} \leq \frac{C_k(\sigma)}{C_k(0)} \leq \frac{D_k(2a_kn_0+\sigma)}{D_k(2a_kn_1)}.$$

(ii) *$\log D_k$ and $\log C_k$ are convex and nonincreasing on $a_k\mathbb{Z}_{\geq 0}$. If $D_k(r') \geq c\,D_k(r)$ with $r < r'$ and $c \leq 1$, then $D_k(\beta) \geq c^{\max(1,(\beta-\alpha)/(r'-r))}D_k(\alpha)$ whenever $r \leq \alpha < \beta$; if $C_k(\sigma) \geq c\,C_k(0)$ with $\sigma > 0$, then $C_k(\sigma') \geq c^{\sigma'/\sigma}C_k(0)$ for $\sigma' \geq \sigma$.*

(iii) *Let k run through a sequence. If $D_k(r') \geq c\,D_k(r)$ for dyadic $0 < r < r'$ with $r \leq 2t_0$, a $c \in (0,1]$ and every large k, then for every dyadic $\sigma \geq 0$ and every large k*

$$C_k(\sigma) \geq c^{\max(1,(2t_1-2t_0+\sigma)/(r'-r))}\,C_k(0).$$

Conversely, if $C_k(\sigma) \geq c\,C_k(0)$ for a dyadic $\sigma > 0$ and every k, then $D_k(2t_0+\sigma') \geq c^{\sigma'/\sigma}D_k(2t_1)$ for every dyadic $\sigma' \geq \sigma$ with $\sigma' > 2(t_1-t_0)$ and every large k.

Proof. For a function F of the slice at time 0 and an integer $n \geq 0$ we have $\Psi_{\tau_n F} = \mathbb{T}_k^n\Psi_F$, since both have inner product $\omega_k(\vartheta(\bar G)\tau_n F)$ with every Ψ_G. By time translation invariance of ω_k, $A_k = \sum_n w_n\tau_n U_k$ with $w_n = a_k g(a_k n) \geq 0$, and $w_n = 0$ unless $n_0 \leq n \leq n_1$; so $\Psi_{A_k} = \sum_n w_n\mathbb{T}_k^n u_k$. Let ν be the spectral measure of $\mathbb{T}_k$ at u_k, on $[0,1]$, and $W(\lambda) = \sum_n w_n\lambda^n$. Then

$$C_k(\sigma) = \int W(\lambda)^2\lambda^{\sigma/a_k}\,d\nu(\lambda), \qquad D_k(r) = \int \lambda^{r/a_k}\,d\nu(\lambda).$$

(i) Since $t_0 > 0$, $n_0 \geq 1$ and $W(0) = 0$; so $C_k(0) > 0$ forces $\nu((0,1]) > 0$, and then $D_k(r) > 0$. For $0 \leq \lambda \leq 1$, $\|w\|_1\lambda^{n_1} \leq W(\lambda) \leq \|w\|_1\lambda^{n_0}$; inserting these bounds in $C_k(\sigma)$ and $C_k(0)$ gives the two inequalities.

(ii) Convexity of $j \mapsto \log\int\lambda^j\,d\rho$ for a positive measure ρ on $[0,1]$ is Hölder's inequality, and the integrals decrease in j because $\lambda \leq 1$; take $\rho = \nu$ and $\rho = W^2\nu$. If $\beta \leq r'$, monotonicity gives $D_k(\beta) \geq D_k(r') \geq cD_k(r) \geq cD_k(\alpha)$. If $\beta > r'$, the difference quotients of the convex function $\log D_k$ increase in both end points, so $(\log D_k(\beta) - \log D_k(\alpha))/(\beta-\alpha) \geq (\log D_k(r') - \log D_k(r))/(r'-r) \geq \log c/(r'-r)$. The statement on C_k is the same comparison of quotients on $[0,\sigma]$ and $[0,\sigma']$.

(iii) For large k the times $2t_0$, $2t_1$ are multiples of a_k, and $2a_kn_0 \geq 2t_0$, $2a_kn_1 \leq 2t_1$. Since D_k is nonincreasing, (i) gives

$$\frac{D_k(2t_1+\sigma)}{D_k(2t_0)} \leq \frac{C_k(\sigma)}{C_k(0)} \leq \frac{D_k(2t_0+\sigma)}{D_k(2t_1)}.$$

The first claim is the left inequality with (ii) for $\alpha = 2t_0 \geq r$ and $\beta = 2t_1 + \sigma$. For the second, (ii) gives $C_k(\sigma') \geq c^{\sigma'/\sigma} C_k(0)$, and the right inequality at σ' gives the claim, the two times being ordered because $\sigma' > 2(t_1 - t_0)$. □

So for the packet $g \otimes h$ the open input of Theorem 10.7(iv) follows from an inequality between two correlations of one slice field,

$$\liminf_k D_k(r')/D_k(r) > 0 \quad \text{for one pair } 0 < r < r' \text{ with } r \leq 2t_0,$$

and by (iii) a uniform bound for one packet gives one for every packet $g' \otimes h$ with g' supported in $[t_1, \infty)$. The condition $r \leq 2t_0$ cannot be dropped: a bound at a later pair says nothing about an earlier packet, as $D_k(r) = (1 - 4^{-k})2^{-kr} + 4^{-k}2^{-r}$ shows, with $D_k(4)/D_k(3) \to \frac{1}{2}$ and $D_k(2)/D_k(1) \to 0$. A bound at a later pair does apply to the packet shifted forward in time, renormalised by its own reflected norm. The left inequality of (i) bounds the ratio without the comparison of u with the packet that was needed after Proposition 10.8(ii). Every bound on such ratios proved in this paper, at strong coupling or through Theorem 10.4, is an upper bound of the form $e^{-m(r'-r)/a_k}$; the inequality above is the opposite one and says that the slice field does not put its weight on energies that diverge in physical units. Reflection positivity and positivity of the transfer matrix do not give it, since a lattice theory with a fixed mass in lattice units has both and its ratio tends to zero. It holds for the free field.

Remark 10.11. Let D_k be defined as above for the massless free lattice field on $a_k\mathbb{Z}^4$ in place of Φ, with $h \in C_c^\infty(\mathbb{R}^3)$ and the lattice heat flow for a physical time $t_f > 0$ in place of the gradient flow. With $\hat{p}^2 = \sum_i (2a_k^{-1}\sin(a_k p_i/2))^2$, $\cosh(a_k\omega) = 1 + a_k^2\hat{p}^2/2$ and $\hat{h}_k(p) = \sum_x a_k^3 h(a_k x) e^{-ia_k p\cdot x}$,

$$D_k(r) = \int_{[-\pi/a_k, \pi/a_k]^3} |\hat{h}_k(p)|^2 e^{-2t_f\hat{p}^2} \frac{e^{-r\omega}}{2a_k^{-1}\sinh(a_k\omega)} \frac{d^3p}{(2\pi)^3}.$$

On the Brillouin zone $\hat{p} \geq 2|p|/\pi$ and $a_k^2\hat{p}^2/2 \leq 6$; as $\operatorname{arccosh}(1+y)/\sqrt{2y}$ decreases on $(0, 6]$, $a_k\omega \geq \kappa a_k\hat{p}$ with $\kappa = \operatorname{arccosh} 7/\sqrt{12} > 0.76$, and $a_k^{-1}\sinh(a_k\omega) \geq \omega$. Since $|\hat{h}_k|$ is bounded uniformly in k, the integrands are dominated by a multiple of $e^{-8t_f|p|^2/\pi^2}/|p|$, which is integrable on $\mathbb{R}^3$, and they converge pointwise to $|\hat{h}(p)|^2 e^{-2t_f|p|^2} e^{-r|p|}/(2|p|)$. By dominated convergence $D_k(r')/D_k(r)$ tends to the corresponding positive ratio of continuum integrals; for $h = e^{-|x|^2}$, $t_f = 0.05$, $r = \frac{1}{2}$ and $r' = 1$ it is 0.6337.

Remark 10.12. The slice field of Proposition 10.10 is not linear: it is a gauge-invariant function of the spatial links of one slice. Its free analogue is the magnetic energy, and the conclusion of Remark 10.11 survives for it. Take the abelian free lattice field on $a_k\mathbb{Z} \times a_k\mathbb{Z}^3_{L_k}$ with $a_k L_k = \ell$ fixed, Feynman gauge, the lattice heat flow of the spatial links for a physical time $t_f \geq 0$, and $U_k = \sum_x a_k^3 \sum_{i<j} (F_{ij}(0,x)^2 - \omega_k(F_{ij}^2))$, the zero spatial momentum part of h (for SU(N) at tree level D_k is multiplied by $N^2 - 1$). Since F_{ij} is Gaussian, $\operatorname{Cov}(F^2, F'^2) = 2\operatorname{Cov}(F, F')^2$, and the mixed propagator of the magnetic field at spatial momentum p is $N_{ij,lm}(p)\, K_p(n)$, with a numerator that does not depend on the time momentum and $K_p(n) = e^{-\omega|n|a_k}/(2\sinh(a_k\omega))$. Hence

$$D_k(r) \propto \sum_{p \in \frac{2\pi}{\ell}\mathbb{Z}^3_{L_k} \setminus 0} S_k(p)\, e^{-4t_f\hat{p}^2} \frac{e^{-2r\omega}}{4a_k^{-2}\sinh^2(a_k\omega)},$$

with $S_k(p) = \sum_{i<j,\, l<m} |N_{ij,lm}(p)|^2 \leq 36\,\hat{p}^4$, and the bounds of Remark 10.11 dominate the summands by $(9/\kappa^2)\,\hat{p}^2 e^{-4t_f\hat{p}^2 - 2r\kappa\hat{p}}$, summable over $\frac{2\pi}{\ell}\mathbb{Z}^3$ for $r > 0$ uniformly in k, also for $t_f = 0$. The summands converge to the continuum ones, so for every $0 < r < r'$ the ratio $D_k(r')/D_k(r)$ converges to a positive limit, with or without the flow. For $\ell = 6$ the computed values of $D_k(\frac{1}{2})/D_k(\frac{1}{4})$ at $t_f = 0.05$ and $a_k^{-1} = 8, 12, 16, 24$ are 0.27341, 0.27239,

0.27201, 0.27174, against 0.27152 in the continuum; at $t_f = 0$ they are 0.0224, 0.0265, 0.0290, 0.0304, against $(r/r')^5 = \frac{1}{32}$ in infinite volume.

The behaviour at $r = 0$ separates the cases. With the flow, $D_k(0)$ stays bounded and the first-step energy $-a_k^{-1}\log(D_k(a_k)/D_k(0))$ is 6.37, 6.50, 6.56, 6.62 at the four spacings, bounded. Without the flow, $D_k(0) \sim a_k^{-5}$ and $D_k(a_k)/D_k(0) = 0.0198$ at every spacing, so the first-step energy grows like $3.92/a_k$ although every ratio at $r > 0$ converges: the condition of Proposition 10.10 at $r > 0$ is weaker than a bound on the first step. For the full energy density, which contains F_{0i} and is not a function of one slice, the electric numerator depends on the time momentum, and the second time difference of K_p is $-\delta_{n0}$ plus a multiple of K_p, a contact term: $D_k(0) \sim a_k^{-2}$ even with the flow, $D_k(a_k)/D_k(0) = 0.042$, 0.023, 0.0145 at $a_k^{-1} = 8, 12, 16$, while the ratios at $r > 0$ again converge to the same continuum values (0.27092, 0.27126, 0.27137). The computation is `slice/composite_free.py`.

At next order the spatial flow, being nonlinear, fuses two hard modes into a soft one, and the state Ψ_{U_k} acquires a component of energy about $2q$ for each $q \lesssim a_k^{-1}$. Its norm is of order g^2 and ultraviolet finite, dominated by $q \approx \tau^{-1/2}$, but its mean energy is not: in the free vacuum the first-step energy is $E_{\mathrm{LO}}(1 + g^2(c + \frac{N}{3\pi^2}\log(\sqrt{\tau}/a_k)))$, and the logarithm has coefficient $\frac{8}{11}$ of $2b_0$, $b_0 = 11N/48\pi^2$. A first-step bound at fixed coupling therefore fails logarithmically, while the ratio at a fixed $r > 0$ only loses the norm of that component, damped by e^{-2qr}. This is why Proposition 10.10 is stated at $r > 0$; the decomposition $\|\mathbb{T}_k^{n/2}u\| \geq c^{n/2}\|v\| - \|w\|$ for $u = v + w$ with a first-step bound on the soft part v alone is the form a four-dimensional argument would use.

Remark 10.13. The inequality also holds along the right tuning in two-dimensional SU(N) lattice gauge theory, which is interacting and nonabelian but has no propagating gluons. Take the lattice $a_k\mathbb{Z} \times a_k\mathbb{Z}_{L_k}$ with $a_k L_k = \ell$ fixed, the weight $\prod_p \exp(\frac{\beta_k}{N}\operatorname{Re}\operatorname{Tr} U_p)$ with $\beta_k = 2N/(g^2 a_k^2)$, and the temporal gauge. A gauge-invariant function of the links of one slice depends only on the conjugacy class of their holonomy V around the circle; this includes every spatially flowed and spatially smeared slice field, since the flow is gauge covariant. Let w_β be the probability density proportional to $e^{\frac{\beta}{N}\operatorname{Re}\operatorname{Tr} U}$ with respect to Haar measure and, for an irreducible representation λ of dimension d_λ and character χ_λ, $\hat{c}_\lambda(\beta) = d_\lambda^{-1}\int \chi_\lambda w_\beta\, dU$. Between two slices put $W_x = V_x' V_x^{-1}$; the W_x are independent with density w_{β_k}, which is invariant under conjugation and inversion, and

$$V_1' \cdots V_L' = W_1\,(V_1 W_2 V_1^{-1}) \cdots \\ \cdots (V_1 \cdots V_{L-1} W_L (V_1 \cdots V_{L-1})^{-1})\, V_1 \cdots V_L,$$

so the new holonomy is QV with Q distributed as the L_k-fold convolution of w_{β_k}, independent of V. By Schur's lemma (Lemma H.1) the transfer matrix acts on class functions by $\chi_\lambda \mapsto \hat{c}_\lambda(\beta_k)^{L_k}\chi_\lambda$. The coefficients are real, since w_β is invariant under $U \mapsto \bar{U}$, nonnegative, since the exponential of $\frac{\beta}{2N}(\chi_{\mathrm{F}} + \bar{\chi}_{\mathrm{F}})$ is a sum of characters with nonnegative coefficients, and below 1 for $\lambda \neq 0$. For a class function f with $\int f\, dU = 0$ and isotypic components of norms $|\hat{f}_\lambda|$, the slice correlation is therefore

$$D_k(r) = \sum_{\lambda \neq 0} |\hat{f}_\lambda|^2 e^{-rE_{\lambda,k}}, \qquad E_{\lambda,k} = -\frac{\ell}{a_k^2}\log \hat{c}_\lambda(\beta_k).$$

Laplace's method[18] at the unique maximum $U = 1$ of $\operatorname{Re}\operatorname{Tr} U$, with $U = e^X$, $\operatorname{Re}\operatorname{Tr} e^X = N + \frac{1}{2}\operatorname{Tr} X^2 + O(|X|^4)$, $\operatorname{Re}\operatorname{Tr} X^3 = 0$, $\operatorname{Tr}\rho_\lambda(X) = 0$ and $\sum_a \rho_\lambda(T_a)^2 = -2C_2(\lambda)$ for an orthonormal basis of $\mathfrak{su}(N)$ in the norm $-\operatorname{Tr} X^2$, gives $\hat{c}_\lambda(\beta) = 1 - NC_2(\lambda)/\beta + O(\beta^{-2})$, with the Casimir normalised by $C_2 = (N^2-1)/(2N)$ on the fundamental representation

[18] For a smooth function with a unique nondegenerate maximum, $\int e^{\beta\varphi}h$ has an expansion in powers of β^{-1} whose coefficients are Gaussian moments at the maximum; Lemma H.2 carries it out here.




FIGURE 10. Two-dimensional SU(N) (Remark 10.13). (a) The lattice $a_k\mathbb{Z} \times a_k\mathbb{Z}_{L_k}$ drawn as a cylinder: the holonomy V of one slice, the holonomy $V' = QV$ of the next, and the plaquette variables W_x between them. (b) The energies $E_{j,k} = -(\ell/a_k^2)\log \hat{c}_j(\beta_k)$ of SU(2) for $g^2 = \ell = 1$, with $\hat{c}_j(\beta) = I_{2j+1}(\beta)/I_1(\beta)$, against the lattice spacing; the dashed lines are the limits $j(j+1)/2$.

(Lemma H.2). Hence $E_{\lambda,k} \to g^2 \ell C_2(\lambda)/2$, and since $D_k(r) \le \|f\|^2$,

$$\liminf_k \frac{D_k(r')}{D_k(r)} \ge \frac{|\hat{f}_\mu|^2}{\|f\|^2} e^{-r' g^2 \ell C_2(\mu)/2} > 0$$

for every $\mu \neq 0$ with $\hat{f}_\mu \neq 0$; for $f = \operatorname{Re}\operatorname{Tr} V$ the ratio is exactly $e^{-(r'-r)E_{\mathrm{F},k}} \to e^{-(r'-r)g^2\ell(N^2-1)/(4N)}$. This is the exact solution of two-dimensional lattice gauge theory, whose continuum limit is the measure of [42, 33, 55], written as the inequality. In two dimensions g^2 has the dimension of a mass squared and there is no ultraviolet problem, so the argument has no four-dimensional analogue. Nor does the spatial flow give the inequality by itself: it damps the spectral weight at large spatial momentum and leaves the weight at zero spatial momentum unchanged. For the free lattice field of Remark 10.11 with a mass M_k and the same flow, every energy is at least $m_k = a_k^{-1}\operatorname{arccosh}(1 + a_k^2 M_k^2/2)$, so $D_k(r') \le e^{-m_k(r'-r)} D_k(r)$, which tends to zero relative to $D_k(r)$ when $a_k M_k$ is fixed.

A route to the slice ratio in four dimensions replaces the packet by its conditional expectations given averages of the field over blocks of increasing size and controls the error of each step. The free field shows in which norm such errors can be small. In the ordinary L^2 norm the first step already fails: conditioning on the averages over blocks of side two leaves a fixed fraction of the variance of an energy packet, at every lattice spacing.

Proposition 10.14. *Let $d \ge 2$, let Λ be a periodic box in $\mathbb{Z}^d$ with even sides, let $\mu > 0$, and let ϕ be the Gaussian field on Λ with density proportional to $\exp(-\frac{1}{2}\sum_{x,\nu}(\nabla_\nu\phi(x))^2 - \frac{\mu^2}{2}\sum_x \phi(x)^2)$, where $\nabla_\nu\phi(x) = \phi(x+e_\nu) - \phi(x)$. Let $\mathcal{Z}$ be a set of blocks $B = 2z + \{0,1\}^d$, let $\mathfrak{B}$ be the σ-algebra generated by the averages of ϕ over the blocks in $\mathcal{Z}$, let $f \ge 0$ on Λ, and let*

$$F = \sum_{x\in\Lambda} f(x) \sum_{\nu=1}^{d} :(\nabla_\nu\phi(x))^2:,$$




FIGURE 11. The ratio $D_k(r)/D_k(\frac{1}{2})$ over the separation r and the lattice spacing a_k. (a) Two-dimensional SU(2) with $f = \operatorname{Re}\operatorname{Tr} V$ and $g^2 = \ell = 1$, where the ratio is $e^{-(r-1/2)E_{1/2,k}}$ and converges as $a_k \to 0$. (b) The term $e^{-(r-1/2)m_k}$ of the lowest energy m_k of a free field with mass $\frac{1}{2}$ in lattice units, which bounds the ratio from above and tends to zero at every $r > \frac{1}{2}$.

where $:X^2: = X^2 - \mathbb{E}X^2$. *For* $B = 2z + \{0,1\}^d$ *put* $x_B = 2z$, $y_B = 2z + e_1 + e_2$ *and*

$$\alpha_B = \frac{1}{4} \sum_{w \in \{x_B, y_B\}} \Big(d\, f(w) + \sum_{\nu=1}^{d} f(w - e_\nu) \Big),$$

and $\sigma^2 = 2/(2d + \mu^2)$. (i) $\mathbb{E}\operatorname{Var}(F \mid \mathfrak{B}) \geq 2\sigma^4 \sum_{B \in \mathcal{Z}} \alpha_B^2$.

(ii) $\operatorname{Var} F \leq 2d^2 \sum_x f(x)^2$.


(iii) *Let* $g \geq 0$ *be a Lipschitz function on* $\mathbb{R}^d$ *with compact support and* $\int g^2 > 0$. *For* $n \geq 1$ *let* Λ_n *contain* $n \operatorname{supp} g$ *in a fundamental domain,* $f_n(x) = g(x/n)$, *let* $\mathcal{Z}_n$ *contain every block that meets* $n \operatorname{supp} g$, *and let* $\mu = \mu_n \to 0$. *Then for the packets* F_n *of* f_n,

$$\liminf_{n\to\infty} \frac{\mathbb{E}\operatorname{Var}(F_n \mid \mathfrak{B}_n)}{n^d \int g^2} \geq 2^{1-d}, \qquad \limsup_{n\to\infty} \frac{\operatorname{Var} F_n}{n^d \int g^2} \leq 2d^2,$$

so $\mathbb{E}\operatorname{Var}(F_n \mid \mathfrak{B}_n) \geq (2^{-d}d^{-2} - o(1)) \operatorname{Var} F_n$.

Proof. (i) No two of the sites x_B, y_B, $B \in \mathcal{Z}$, are neighbours: their coordinate sums are even, and a neighbour of a site has a coordinate sum of the other parity. Let $\mathfrak{H} \supseteq \mathfrak{B}$ be generated by $\mathfrak{B}$ and by $\phi(w)$ for every w outside $\{x_B, y_B : B \in \mathcal{Z}\}$. By the law of total variance, $\operatorname{Var}(F \mid \mathfrak{B}) = \mathbb{E}[\operatorname{Var}(F \mid \mathfrak{H}) \mid \mathfrak{B}] + \operatorname{Var}(\mathbb{E}[F \mid \mathfrak{H}] \mid \mathfrak{B})$, so $\mathbb{E}\operatorname{Var}(F \mid \mathfrak{B}) \geq \mathbb{E}\operatorname{Var}(F \mid \mathfrak{H})$. In the exponent of the density, a site w enters through $(d + \frac{\mu^2}{2})\phi(w)^2$ and through $-\phi(w)\phi(w')$ for each of its $2d$ neighbours w'. Given $\mathfrak{H}$, the block average fixes $s_B = \phi(x_B) + \phi(y_B)$, and with $u_B = \phi(x_B) - \phi(y_B)$ we have $\phi(x_B)^2 + \phi(y_B)^2 = \frac{1}{2}(s_B^2 + u_B^2)$. Since the free sites have only $\mathfrak{H}$-measurable neighbours, the conditional density of $(u_B)_{B\in\mathcal{Z}}$ is proportional to $\prod_B \exp(-\frac{1}{4}(2d+\mu^2)u_B^2 + c_B u_B)$ with $\mathfrak{H}$-measurable c_B: the u_B are conditionally independent Gaussians of variance σ^2. A term $(\nabla_\nu \phi(w))^2$ contains at most one free site, and if it contains x_B, then $\phi(x_B) = \frac{1}{2}(s_B + u_B)$ shows that it is $\frac{1}{4}u_B^2$ plus a term linear in u_B plus an $\mathfrak{H}$-measurable term; the same holds for y_B with $-u_B$. The terms containing x_B are those with $w = x_B$ and those with $w = x_B - e_\nu$, d of each kind, so

$$F = \sum_{B \in \mathcal{Z}} (\alpha_B u_B^2 + \beta_B u_B) + F_0$$



block averages of side b between the plane $t=0$ and the packet F at height d

FIGURE 12. Block averages between the reflection plane $t = 0$ and a packet F at physical height d, for blocks of side $d/4$, $d/2$ and d. By Proposition 10.14 the first conditioning step already leaves a fixed fraction of the ordinary variance of the packet, so the errors of such steps have to be measured in the reflected norm.

with $\mathfrak{H}$-measurable β_B and F_0. Writing $u_B = m_B + v_B$ with m_B the conditional mean, $\alpha_B u_B^2 + \beta_B u_B = \alpha_B v_B^2 + (2\alpha_B m_B + \beta_B) v_B + \text{const}$, and for a centred Gaussian v of variance σ^2, $\operatorname{Var}(\alpha v^2 + \gamma v) = 2\alpha^2\sigma^4 + \gamma^2\sigma^2$. By conditional independence, $\operatorname{Var}(F \mid \mathfrak{H}) \geq 2\sigma^4 \sum_B \alpha_B^2$.

(ii) For centred jointly Gaussian X, Y, $\operatorname{Cov}(:X^2:, :Y^2:) = 2\operatorname{Cov}(X,Y)^2$. With $G_{\nu\rho}(x,y) = \mathbb{E}[\nabla_\nu\phi(x)\nabla_\rho\phi(y)]$,

$$\begin{aligned}\operatorname{Var} F &= 2\sum_{x,y} f(x)f(y) \sum_{\nu,\rho} G_{\nu\rho}(x,y)^2 \\ &\leq \sum_{x,y} (f(x)^2 + f(y)^2) \sum_{\nu,\rho} G_{\nu\rho}(x,y)^2.\end{aligned}$$

On the dual torus Λ^*, with $\hat{p}^2 = \sum_\nu |e^{ip_\nu} - 1|^2$,

$$G_{\nu\rho}(x,y) = \frac{1}{|\Lambda|} \sum_{p \in \Lambda^*} \frac{(e^{ip_\nu} - 1)(e^{-ip_\rho} - 1)}{\hat{p}^2 + \mu^2} e^{ip\cdot(x-y)},$$

and by Parseval $\sum_y G_{\nu\rho}(x,y)^2 = |\Lambda|^{-1} \sum_p |e^{ip_\nu} - 1|^2 |e^{ip_\rho} - 1|^2 (\hat{p}^2 + \mu^2)^{-2} \leq 1$, since each factor $|e^{ip_\nu} - 1|^2$ is at most $\hat{p}^2$. The same holds for the sum over x, and (ii) follows.

(iii) If $\operatorname{Lip} g = K$, every site entering α_B lies within distance 3 of $2z$, so $|\alpha_B - d\, g(2z/n)| \leq 3dK/n$. Hence $\sum_{B \in \mathcal{Z}_n} \alpha_B^2 \geq \sum_z (d\, g(2z/n) - 3dK/n)_+^2$, the sum running over z with $2z \in n \operatorname{supp} g$, and this is $d^2 (n/2)^d \int g^2 \, (1 + o(1))$ as a Riemann sum on the grid $(2/n)\mathbb{Z}^d$. Since $\sigma^4 = 4/(2d + \mu_n^2)^2 \to d^{-2}$, (i) gives the first bound. By (ii) and $\sum_x g(x/n)^2 = n^d \int g^2 (1 + o(1))$, the second follows. □

For $d = 4$ and the energy density, $\operatorname{Var} F_n$ grows like $(\ell/a)^4$ at a fixed physical size ℓ of the packet, and no step of conditioning on block averages removes more than a fraction $1 - 2^{-d} d^{-2} + o(1)$ of it. The reflected norm $C_k(0) = \omega_k(\vartheta(\bar{A}_k) A_k)$, which renormalises the packet in Theorem 10.7, involves the covariance only between the support of ϑf_0 and that of f_0, at a fixed physical distance. The errors of a blocking route have therefore to be measured in that norm; Figure 12 shows the blocks between the reflection plane and the packet.

10.5. **Higher correlations.** Theorem 10.7 controls two renormalised fields at a time, and part (i) of it gives nothing for more: Z_k may diverge while the fields $\Phi_x \circ \Phi_{s_k}$ stay bounded, so a bound on the norm of the renormalised packet as an operator is not available. A

reflection positive process with a fixed gap and a two-point function independent of the cutoff can have divergent four-point functions at separated times (Remark 10.18), so no statement about the two-point function alone can replace the following hypothesis on the truncated correlations. In this subsection Φ is real-valued, as the energy of the plaquettes is; for a complex field the conjugate field has to be adjoined and (T1) imposed on all mixed cumulants. Write $\varphi_k(x) = \Phi_x \circ \Phi_{s_k}$, ω_k^T for the truncated correlations (joint cumulants) under ω_k, and for $f \in C_c(\mathbb{R}^4)$

$$A_k(f) = \sum_{x \in \mathbb{Z}^4} a_k^4 f(a_k x)(\varphi_k(x) - \omega_k(\varphi_k(x))),$$

so that $A_k = A_k(f_0)$ and $\bar{A}_k(f) = A_k(\bar{f})$. Let $w_k : \mathbb{Z}^4 \to [0,\infty)$ be even functions with $W_k = \sum_x w_k(x) < \infty$, let $r_k > 0$ be amplitude scales, and consider:

(T1) *Tree bounds.* For every $n \geq 2$ there is K_n, independent of k, with

$$|\omega_k^T(\varphi_k(x_1); \dots; \varphi_k(x_n))| \leq K_n r_k^n \sum_T \prod_{\{i,j\} \in T} w_k(x_i - x_j)$$

for all $x_1, \dots, x_n \in \mathbb{Z}^4$, the sum running over the trees T on $\{1, \dots, n\}$.

(T2) *Saturation.* $C_k(0) \geq c\, r_k^2 a_k^4 W_k$ for a $c > 0$ and every k.

(T3) *Physical volume.* $B_* = \sup_k a_k^4 W_k < \infty$.

The amplitude r_k separates the size of the field from the volume over which it is correlated; Remark 10.17 shows why it is needed. For a compact $Q \subset \mathbb{R}^4$ let $|Q|_k = a_k^4 \#\{x \in \mathbb{Z}^4 : a_k x \in Q\}$; for a cube of side L it is at most $(L + a_0)^4$.

Lemma 10.15. *Under* (T1)*, for $n \geq 2$ and bounded $f_1, \dots, f_n$ with compact supports, f_1 supported in Q,*

$$|\omega_k^T(A_k(f_1); \dots; A_k(f_n))| \leq K_n n^{n-2} r_k^n\, |Q|_k\, (a_k^4 W_k)^{n-1} \prod_{j=1}^n \|f_j\|_\infty.$$

Proof. Joint cumulants of order $n \geq 2$ are multilinear and do not change when constants are added to the variables, so the left side is at most $a_k^{4n} \sum_{x_1,\dots,x_n} \prod_j |f_j(a_k x_j)|$ times the absolute value of the truncated correlation, the sums being finite. Insert (T1) and fix a tree T. With x_1 fixed, sum over the other points one at a time, each time over a leaf of what remains of T other than 1; each such sum is at most W_k. The remaining sum over x_1 has at most $|Q|_k a_k^{-4}$ nonzero terms. There are n^{n-2} trees. □

Theorem 10.16. *Assume* (T1)–(T3).

(i) *For $n \geq 2$, f_j as in the lemma,*

$$|\omega_k^T(Z_k A_k(f_1); \dots; Z_k A_k(f_n))| \leq K_n n^{n-2} c^{-n/2}\, B_*^{n/2-1}\, |Q|_k \prod_{j=1}^n \|f_j\|_\infty,$$

and the moments $\omega_k(\prod_j Z_k A_k(f_j))$ with all f_j supported in a cube Q are bounded by $M_n(Q) \prod_j \|f_j\|_\infty$, with $M_n(Q)$ independent of k.

(ii) *Every sequence of k has a subsequence along which*

$$S_n(f_1, \dots, f_n) = \lim_k \omega_k\Big(\prod_{j=1}^n Z_k A_k(f_j)\Big)$$

exists for all n and all $f_j \in C_c(\mathbb{R}^4)$. Each S_n is symmetric, multilinear, bounded by $M_n(Q) \prod_j \|f_j\|_\infty$ on functions supported in Q, hence a distribution on $(\mathbb{R}^4)^n$; it is invariant under all translations of $\mathbb{R}^4$ if the ω_k are translation invariant, and the kernel K of

Theorem 10.7 along the subsequence is $K(y,y') = S_2(\overline{f_0(\cdot - y)} \circ \theta, f_0(\cdot - y'))$, *with* θ *as in* (iii).

(iii) *Put* $S_0 = 1$, $S_1 = 0$, *and let* θ *be the reflection* $t \mapsto -t$ *of* $\mathbb{R}^4$. *For every finite family of* $c_\alpha \in \mathbb{C}$ *and* $f_{\alpha,j} \in C_c(\mathbb{R}^4)$ *supported in* $\{t > 0\}$,

$$\sum_{\alpha,\beta} \bar{c}_\alpha c_\beta \, S\Big(\prod_j \overline{f_{\alpha,j}} \circ \theta \cdot \prod_j f_{\beta,j}\Big) \geq 0,$$

where S *of a product of* n *test functions means* S_n. *Hence the Osterwalder–Schrader construction gives a Hilbert space* $\mathcal{H}$ *with vacuum* Ω *and a dense set of vectors* $\Psi(P)$ *indexed by polynomials* P *in the smeared fields at positive times, with* $\langle \Psi(P), \Psi(P') \rangle = S(\theta \bar{P} \cdot P')$; *in particular the separated higher correlations of the renormalised fields are finite. The vector* $\Psi(P_0)$ *of the field smeared by* f_0 *is a unit vector orthogonal to* Ω.

(iv) *Suppose the* ω_k *are translation invariant. Then the time translations by* $t \geq 0$ *define on* $\mathcal{H}$ *a strongly continuous semigroup* e^{-tH} *with* $H \geq 0$ *self-adjoint, and the spatial translations a strongly continuous unitary group commuting with* H. *Moreover there is a dyadic* $\sigma > 0$ *with* $C_k(\sigma) \geq \frac{1}{2} C_k(0)$ *for all large* k, *so the limit of Theorem 10.7 is nontrivial.*

(v) *If* $a_k^4 W_k \to 0$ *along the subsequence, every* S_n *is given by Wick's rule from* S_2: *the limit is the Gaussian field with covariance* S_2.

(vi) *If the* ω_k *are translation invariant and the hypothesis of Theorem 10.4 holds, then* 0 *is a simple eigenvalue of* H *and* $\sigma(H) \subset \{0\} \cup [m_*, \infty)$ *with* $m_* = \log(1/\alpha)/(2\ell)$, *and* $\Omega^\perp \neq 0$.

Proof. (i) By (T2), $Z_k^n = C_k(0)^{-n/2}$ is at most $(c\, r_k^2 a_k^4 W_k)^{-n/2}$; multiply the lemma by it, which cancels r_k^n, and use $(a_k^4 W_k)^{n/2-1} \leq B_*^{n/2-1}$ by (T3) for $n \geq 2$. A moment is the sum over partitions of $\{1, \dots, n\}$ of the products of the cumulants of the blocks; blocks of size one vanish because $A_k(f)$ is centred, and each block contributes the bound just proved with $|Q|_k \leq (L + a_0)^4$ for a cube of side L. This gives $M_n(Q)$.

(ii) For each cube $Q_L = [-L, L]^4$, $L \in \mathbb{N}$, the continuous functions supported in Q_L form a separable space under the supremum norm. Choose a countable dense set $\mathcal{F}$ of functions; by (i) the countably many sequences $\omega_k(\prod Z_k A_k(f_j))$, $f_j \in \mathcal{F}$, are bounded, and a diagonal subsequence makes all of them converge. By multilinearity,

$$\begin{aligned} &\Big|\omega_k\Big(\prod_j Z_k A_k(f_j)\Big) - \omega_k\Big(\prod_j Z_k A_k(g_j)\Big)\Big| \\ &\qquad \leq M_n(Q) \sum_i \|f_i - g_i\|_\infty \prod_{j \neq i} \max(\|f_j\|_\infty, \|g_j\|_\infty) \end{aligned}$$

for functions supported in Q, so the limits exist for all $f_j \in C_c(\mathbb{R}^4)$ and obey the same bound. Symmetry and multilinearity pass to the limit. A multilinear form bounded in this way is separately continuous on $C_c^\infty(\mathbb{R}^4)$, so the Schwartz kernel theorem makes it a distribution. For dyadic y and large k, $y/a_k \in \mathbb{Z}^4$ and translation invariance of ω_k and of the flow (Proposition 10.6(i)) give $\omega_k(\prod Z_k A_k(f_j(\cdot - y))) = \omega_k(\prod Z_k A_k(f_j))$. For a real y take dyadic $y_m \to y$; the functions $f_j(\cdot - y_m)$ converge uniformly to $f_j(\cdot - y)$ with supports in one cube, so the bound above carries the invariance to y. The formula for K follows from $A_k^y = A_k(f_0(\cdot - y))$ and the argument of (iii) with $n = 2$.

(iii) Let ϑ be the lattice reflection of Appendix F and θ_k the corresponding reflection of $a_k \mathbb{Z}^4$; it differs from θ by a translation of length at most a_k. Since Φ is real and ϑ commutes with the flow, $\vartheta(\bar{A}_k(f)) = A_k(\bar{f} \circ \theta_k)$. Let $P_k = \sum_\beta c_\beta \prod_j Z_k A_k(f_{\beta,j})$ and $R_k = \sum_\alpha \bar{c}_\alpha \prod_j Z_k A_k(\overline{f_{\alpha,j}} \circ \theta)$, so that $\vartheta(\bar{P}_k)$ is R_k with θ replaced by θ_k. Reflection positivity gives $\omega_k(\vartheta(\bar{P}_k) P_k) \geq 0$. By (i), applied in a cube containing all the supports and their reflections, and by multilinearity, $|\omega_k(\vartheta(\bar{P}_k) P_k) - \omega_k(R_k P_k)|$ is at most a constant independent of k times $\max_{\alpha,j} \|f_{\alpha,j} \circ \theta_k - f_{\alpha,j} \circ \theta\|_\infty$, which tends to zero by uniform continuity, while $\omega_k(R_k P_k)$ tends to the left side of (iii) by (ii). So the limit of the nonnegative numbers

is the left side of (iii). The construction of $\mathcal{H}$ is the usual quotient of the polynomials at positive times by the null space of this form [65]. By the same approximation of θ by θ_k, $\|\Psi(P_0)\|^2 = \lim_k Z_k^2 C_k(0) = 1$, and $\langle \Omega, \Psi(P_0)\rangle = S_1(f_0) = 0$.

(iv) For a polynomial P at positive times let $\tau_t P$ be its time translate by $t \geq 0$, again at positive times. By translation invariance and $\theta \circ \tau_t = \tau_{-t} \circ \theta$, $\langle \Psi(\tau_t P), \Psi(P')\rangle = \langle \Psi(P), \Psi(\tau_t P')\rangle$, and $q_P(t) = \|\Psi(\tau_t P)\|^2 = \langle \Psi(P), \Psi(\tau_{2t} P)\rangle$. If P has degree at most d, every partition in (i) of the products in $q_P(t)$ has at most d blocks, each bounded by a multiple of $|Q|_k$ for a cube Q of side $O(1+t)$, so $q_P(t) \leq C_P(1+t)^{4d}$. The Cauchy–Schwarz inequality gives $q_P(t) \leq q_P(0)^{1/2} q_P(2t)^{1/2}$, hence by induction $q_P(t) \leq q_P(0)^{1-2^{-m}} q_P(2^m t)^{2^{-m}}$ for every m, and letting $m \to \infty$, $q_P(t) \leq q_P(0)$. So τ_t maps null vectors to null vectors and extends to a symmetric contraction $\mathsf{T}(t)$ of $\mathcal{H}$; $\mathsf{T}(s)\mathsf{T}(t) = \mathsf{T}(s+t)$, and $\mathsf{T}(t) = \mathsf{T}(t/2)^2 \geq 0$. By the bound of (ii) and uniform continuity of the translated test functions, $\|\Psi(\tau_t P) - \Psi(P)\| \to 0$ as $t \to 0$, so the semigroup is strongly continuous on a dense set, hence everywhere, and $\mathsf{T}(t) = e^{-tH}$ with $H \geq 0$ self-adjoint [65]. Spatial translations preserve the form, have the opposite translations as inverses, commute with τ_t, and are strongly continuous by the same argument.

For the ratio let $f_{0,t} = f_0(\cdot - (t,0))$. For dyadic $t > 0$ and large k, translation invariance gives $\tau_{t/a_k} A_k = A_k(f_{0,t})$, so by (iii)

$$C_k(t) - C_k(0) = \omega_k\big(A_k(\bar{f}_0 \circ \theta_k)\, A_k(f_{0,t} - f_0)\big),$$

a covariance of two centred fields. The case $n = 2$ of (i) bounds Z_k^2 times it by $L_0 \|f_{0,t} - f_0\|_\infty$ with $L_0 = (K_2/c) \sup_k |Q|_k \|f_0\|_\infty$, Q a cube containing the supports of $f_0 \circ \theta_k$ for all k. By uniform continuity of f_0 there is a dyadic $\sigma > 0$ with $L_0 \|f_{0,\sigma} - f_0\|_\infty \leq \frac{1}{2}$; then $C_k(\sigma)/C_k(0) \geq \frac{1}{2}$, and Theorem 10.7(iv) applies.

(v) For $n \geq 3$ the exponent $n/2 - 1$ in (i) is positive, so with $(a_k^4 W_k)^{n/2-1}$ in place of $B_*^{n/2-1}$ the cumulants of order $n \geq 3$ of the renormalised fields tend to zero. The moments are the sums over partitions of products of cumulants, so in the limit only the partitions into pairs remain, which is Wick's rule.

(vi) As in the proof of Theorem 10.4, $\mathbb{T}_k$ has gap at least $\log(1/\alpha)/(2n_k)$ on the complement of Ω_k. For a polynomial P at positive times let $v_k = \Psi_{P_k} - \omega_k(P_k)\Omega_k$ with P_k its lattice version. For dyadic t and large k,

$$\omega_k(\vartheta(\bar{P}_k)\tau_{t/a_k} P_k) - |\omega_k(P_k)|^2 = \langle v_k, \mathbb{T}_k^{t/a_k} v_k\rangle \leq e^{-m_* t} \|v_k\|^2,$$

and both sides are finite combinations of moments, which converge by (ii) and the approximation of θ in (iii). The limit is $\langle v, e^{-tH} v\rangle \leq e^{-m_* t}\|v\|^2$ for $v = \Psi(P) - \langle \Omega, \Psi(P)\rangle \Omega$. These v are dense in $\Omega^\perp$, so the inequality holds on $\Omega^\perp$ for all $t \geq 0$, and the spectral theorem gives the spectral statement as in Theorem 10.4. By (iii), $\Psi(P_0)$ is a unit vector in $\Omega^\perp$. □

Part (iv) shows that the order-two cases of (T1) and (T2) already contain the lower bound on the ratio that Theorem 10.7(iv) needs; they are not harmless combinatorial assumptions, and proving them along the continuum tuning is at least as hard as that lower bound. Part (vi) passes the conditional mass gap of Theorem 10.4 to every polynomial in the smeared fields. Neither part proves Euclidean invariance, independence of the limit from the subsequence, or the reconstruction of the field operators, for which the growth of $M_n(Q)$ in n would also be needed.

Remark 10.17. The amplitude scale cannot be dropped for a bounded field. Suppose $\operatorname{Var} \varphi_k(0) \to 0$, as for the energy of the flowed plaquettes in lattice units when the flowed links approach the identity. Reflection invariance of ω_k, the Cauchy–Schwarz inequality and stationarity give

$$C_k(0) \leq \omega_k(|A_k|^2) \leq \Big(a_k^4 \sum_x |f_0(a_k x)|\Big)^2 \operatorname{Var} \varphi_k(0) \to 0.$$

With $r_k = 1$, (T2) then forces $a_k^4 W_k \to 0$, and by part (v) the limit is Gaussian. With r_k^2 of the order of $\operatorname{Var}\varphi_k(0)$ this is no longer forced. Choosing r_k does not prove (T1) or (T2).

Remark 10.18. Let $0 < \epsilon < 1$, $m > 0$, let π be the probability measure on $\{-1, 0, 1\}$ with weights $\epsilon/2$, $1-\epsilon$, $\epsilon/2$, and let $P_t = e^{-mt} I + (1 - e^{-mt})\Pi$ on $L^2(\pi)$, where $\Pi f = \pi(f)$ is the projection onto the constants. This is a symmetric Markov semigroup with $P_t \geq 0$ and gap m, so its stationary process is reflection positive. For $X(s) = s/\sqrt{\epsilon}$ we have $\pi(X) = \pi(X^3) = 0$, $\pi(X^2) = 1$, $\pi(X^4) = 1/\epsilon$ and $P_t X = e^{-mt} X$, so the two-point function is e^{-mt} for every ϵ, while for $t_1 \leq t_2 \leq t_3 \leq t_4$ and $q_i = e^{-m(t_{i+1} - t_i)}$ the four-point function is $q_3\, \pi(X P_{t_2 - t_1}(X P_{t_3 - t_2} X^2)) = q_1 q_3 (q_2/\epsilon + 1 - q_2)$, with truncated part $q_1 q_3 (q_2/\epsilon - q_2 - 2q_2^2)$; both diverge as $\epsilon \to 0$. By part (i), no choice of amplitude scales makes (T1)–(T3) hold for this family with constants independent of ϵ.

The hypotheses (T1)–(T3) are uniform cluster estimates for the continuum tuning $\beta_k \to \infty$, and the next proposition shows that they are estimates at a critical point: with the order-two cases alone the lattice gap is at most a fixed multiple of a_k, and the tree majorant carries a fixed fraction of its weight at a fixed physical distance. Let $m_k \in [0, \infty]$ be the gap of $\mathbb{T}_k$ on the complement of Ω_k, so that $\|\mathbb{T}_k v\| \leq e^{-m_k}\|v\|$ for $v \perp \Omega_k$, and let $t_0 > 0$ be the least time in the support of f_0.

Proposition 10.19. (i) *For every dyadic $\sigma > 0$ and every large k, $C_k(\sigma) \leq e^{-m_k \sigma / a_k} C_k(0)$. If $m_k/a_k \to \infty$, every limit of Theorem 10.7 is trivial; this is the case for $\beta_k = \beta$ fixed in the range of Theorem 1.5, where $m_k \geq m_G(\beta)$.*

(ii) *Suppose the ω_k are translation invariant and* (T1) *for $n = 2$ and* (T2) *hold. Then $m_k \leq a_k \log 2/\sigma$ for every large k, where σ is the dyadic number of Theorem 10.16*(iv)*, which depends only on K_2/c and f_0. If also the hypothesis of Theorem 10.4 holds, then $m_* a_k \leq m_k \leq a_k \log 2/\sigma$, and in particular $m_* \leq \log 2/\sigma$.*

(iii) *Under* (T1) *for $n = 2$ and* (T2)*, let Q be a cube containing the support of f_0 and $\gamma = c/(K_2 \|f_0\|_\infty^2 \sup_k |Q|_k)$. Then for every k*

$$\sum_{z \in \mathbb{Z}^4,\ |z_0| \geq R_k} w_k(z) \geq \gamma\, W_k, \qquad R_k = \frac{2t_0}{a_k} - 1.$$

If $w_k(z) = e^{-\mu_k |z_0|} u_k(\bar{z})$ with $\bar{z}$ the spatial part of z and $u_k \geq 0$ summable on $\mathbb{Z}^3$, then $\mu_k R_k \leq \log(2/\gamma)$, so the physical time decay rate of the majorant is at most $\log(2/\gamma)/(2t_0 - a_k)$.

Proof. (i) In the proof of Theorem 10.7, $C_k(\sigma)/C_k(0) = \langle \Psi_k, \mathbb{T}_k^{\sigma/a_k} \Psi_k \rangle$ with Ψ_k a unit vector orthogonal to Ω_k. The operator $\mathbb{T}_k$ is self-adjoint and fixes Ω_k, so it preserves the complement, and the inequality follows. If $m_k/a_k \to \infty$ then $\rho(\sigma) = 0$ for every dyadic $\sigma > 0$ along every subsequence, and the limit is trivial by Theorem 10.7(iv). At fixed β in the range of Theorem 1.5 the gap is at least $m_G(\beta) > 0$ in lattice units, and $a_k \to 0$; the spatial flow does not change the transfer matrix (Proposition 10.6(iv)).

(ii) Theorem 10.16(iv) uses (T1) and (T2) only for $n = 2$, since the exponent $n/2 - 1$ in part (i) vanishes there, and gives $C_k(\sigma) \geq \frac{1}{2} C_k(0)$ for large k. With (i), $e^{-m_k \sigma / a_k} \geq \frac{1}{2}$. Under the hypothesis of Theorem 10.4, the proof of that theorem gives $m_k \geq \log(1/\alpha)/(2n_k) \geq m_* a_k$.

(iii) $C_k(0) = \omega_k(A_k(\bar{f}_0 \circ \theta_k) A_k(f_0))$, a covariance of two centred fields, by the proof of Theorem 10.16(iv). A point x in the support of the first sum has $a_k x_0 \leq -t_0 + a_k$, and a point y in the support of the second has $a_k y_0 \geq t_0$, so $|x_0 - y_0| \geq R_k$. By (T1) for $n = 2$,

$$\begin{aligned} C_k(0) &\leq K_2 r_k^2 \|f_0\|_\infty^2\, a_k^8 \sum_{a_k y \in Q} \sum_{|x_0 - y_0| \geq R_k} w_k(x - y) \\ &\leq K_2 r_k^2 \|f_0\|_\infty^2 |Q|_k\, a_k^4 \sum_{|z_0| \geq R_k} w_k(z), \end{aligned}$$

and (T2) bounds the left side below by $c\,r_k^2 a_k^4 W_k$. For the product form let $R' = \lceil R_k \rceil$; if $R' \le 0$ there is nothing to prove, and otherwise

$$\frac{\sum_{|j|\ge R'} e^{-\mu_k|j|}}{\sum_{j\in\mathbb{Z}} e^{-\mu_k|j|}} = \frac{2e^{-\mu_k R'}}{1+e^{-\mu_k}} \le 2e^{-\mu_k R_k},$$

while the factor $\sum_{\bar z} u_k(\bar z)$ cancels. So $2e^{-\mu_k R_k} \ge \gamma$. □

So reflection positivity and decay do not supply (T2): at fixed β in the range of Theorem 1.5 the measures are reflection positive with exponential clustering, and $C_k(\sigma)/C_k(0) \to 0$, so no choice of w_k, r_k and constants satisfies the order-two cases of (T1) and (T2) together there. For the same reason no estimate whose constants give a lattice gap bounded below, as a convergent strong coupling expansion does [67], can give the tree bounds along the tuning: (T1) and (T2) hold, if at all, only along couplings where the lattice gap closes at least as fast as a_k. A mass gap in physical units, which bounds the decay of the two-point function, does not give the constants K_n either. By Theorem 10.16(v) the limit can be non-Gaussian only if $\liminf_k a_k^4 W_k > 0$; for the product majorants of part (iii) the time factor of W_k is $\coth(\mu_k/2) \ge 2/\mu_k$, which is of order $1/a_k$ at least, so the time direction alone is compatible with this. At a fixed physical flow time the flowed fields are smooth on the scale $\sqrt{\tau}$; majorants of product form for them are not proved here.

The order-two cases of (T1) and (T2) say more about the reference vector than part (iv) of Theorem 10.16 records: for a smooth f_0 they bound every time difference of the packet, which makes its energy distribution tight with finite moments and gives the ratio with an explicit rate. Recall the spectral measures μ_k on $[0,\infty]$ of the proof of Theorem 10.7, so that $C_k(\sigma)/C_k(0) = \int e^{-\sigma E} d\mu_k(E)$.

Proposition 10.20. *Suppose the ω_k are translation invariant,* (T1) *holds for $n = 2$ and* (T2) *holds, and f_0 has $m \ge 1$ continuous time derivatives. Let Q be a cube containing the supports of $f_0(\cdot - (s,0))$ for $0 \le s \le m a_0$, and $B_m = (K_2/c)\sup_k |Q|_k \, \|\partial_t^m f_0\|_\infty^2$.*

(i) *$\|(1-\mathbb{T}_k)^m \Psi_k\|^2 \le B_m a_k^{2m}$ for every k, where $\Psi_k = Z_k \Psi_{A_k}$.*

(ii) *$\int \big((1-e^{-a_k E})/a_k\big)^{2m} d\mu_k(E) \le B_m$ and $\mu_k([\Lambda,\infty]) \le B_m (a_k + \Lambda^{-1})^{2m}$ for $\Lambda > 0$. Every weak limit μ of the μ_k satisfies $\mu(\{\infty\}) = 0$ and $\int E^{2m} d\mu(E) \le B_m$. If* (T1)–(T3) *hold, the unit vector $\Psi(P_0)$ of Theorem* 10.16(iii)–(iv) *lies in the domain of H^m, with $\|H^m \Psi(P_0)\|^2 \le B_m$.*

(iii) *If $a_k\sqrt{B_1} \le \frac12$, then $C_k(\sigma) \ge e^{-2\sqrt{B_1}\sigma} C_k(0)$ for every $\sigma \in a_k \mathbb{Z}_{\ge 0}$, and the gap m_k of Proposition 10.19 satisfies $m_k \le -\log(1 - a_k\sqrt{B_1}) \le 2\sqrt{B_1}\, a_k$.*

Proof. (i) Let $D_k f_0 = \sum_{j=0}^m (-1)^j \binom{m}{j} f_0(\cdot - (j a_k, 0))$. By translation invariance $\mathbb{T}_k^j \Psi_{A_k} = \Psi_{\tau_j A_k}$ and $\tau_j A_k = A_k(f_0(\cdot - (j a_k, 0)))$, so $(1-\mathbb{T}_k)^m \Psi_{A_k} = \Psi_{A_k(D_k f_0)}$, and as in the proof of Theorem 10.16(iv) its squared norm is the covariance $\omega_k(A_k(\overline{D_k f_0} \circ \theta_k) A_k(D_k f_0))$. The function $D_k f_0$ is supported in Q and is an m-th difference with step a_k, so $\|D_k f_0\|_\infty \le a_k^m \|\partial_t^m f_0\|_\infty$. Lemma 10.15 with $n = 2$ bounds the covariance by $K_2 r_k^2 |Q|_k \, a_k^4 W_k \|D_k f_0\|_\infty^2$, and (T2) bounds $C_k(0) = Z_k^{-2}$ below by $c\,r_k^2 a_k^4 W_k$.

(ii) By the spectral theorem the left side of (i) is $\int (1-e^{-a_k E})^{2m} d\mu_k$, the integrand being 1 at $E = \infty$. For $E \ge \Lambda$, $(1-e^{-a_k E})/a_k \ge (1-e^{-a_k\Lambda})/a_k \ge \Lambda/(1+a_k\Lambda)$, because $1 - e^{-x} \ge x/(1+x)$; this gives the tail bound. The set $(\Lambda,\infty]$ is open in $[0,\infty]$, so $\mu((\Lambda,\infty]) \le \liminf_k \mu_k((\Lambda,\infty]) \le B_m \Lambda^{-2m}$, and $\mu(\{\infty\}) = 0$. For a continuous χ with $0 \le \chi \le 1$ and support in $[0, L]$, the same inequality gives $E \le (1 + a_k L)(1 - e^{-a_k E})/a_k$ on that support, so $\int \chi E^{2m} d\mu = \lim_k \int \chi E^{2m} d\mu_k \le B_m$; letting χ increase to 1 gives the moment bound. The vector $\Psi(P_0)$ has $\langle \Psi(P_0), e^{-\sigma H} \Psi(P_0)\rangle = \lim_k C_k(\sigma)/C_k(0) = \int e^{-\sigma E} d\mu$ for dyadic $\sigma > 0$ along the subsequence, so its spectral measure for H is μ.

(iii) By (i) with $m = 1$ and the Cauchy–Schwarz inequality, $\langle \Psi_k, (1-\mathbb{T}_k)\Psi_k\rangle \le a_k\sqrt{B_1}$, which is the hypothesis of Proposition 10.8(ii) with $C = \sqrt{B_1}$. Since Ψ_k is a unit vector orthogonal to Ω_k, the same quantity is at least $1 - e^{-m_k}$. □

Part (iii) makes Proposition 10.19(ii) explicit: the order-two tree and saturation bounds pin the lattice gap below $2\sqrt{B_1}\,a_k$, a physical mass of at most $2\sqrt{B_1}$, and give the ratio of Theorem 10.7(iv) at every separation. The open input of Theorem 10.7 is therefore implied by these two bounds for one smooth f_0; nothing here proves them for the Wilson action.

11. The open inputs

This section states the two unproved inputs of Section 10, the cube estimate and the limit, and measures the first: what can be controlled uniformly in the coupling, and what any proof of the cube estimate has to produce.

11.1. **The two open statements.** Theorem 10.4 reduces a continuum mass gap to two statements, neither of which is proved here or, to our knowledge, anywhere else.

(D) *The cube estimate.* There are $\alpha < 1$ and $\ell > 0$ such that $D(n, \alpha)$ holds with $na \le \ell$ along the continuum tuning $\beta = \beta(a) \to \infty$, uniformly in a.

(E) *The limit.* The renormalised Schwinger functions converge along $a_k = 2^{-k}a_0$ to Schwinger functions that satisfy the Osterwalder–Schrader axioms and are not those of the vacuum alone.

(D) and (E) together give the mass gap of the continuum theory; the remaining Clay requirements, Euclidean invariance and the local structure of the limit [50], belong to (E). (D) is a statement about a single cube of fixed physical size and a single coupling at each spacing; (E) is the construction problem, on which Bałaban's renormalisation group analysis [6, 7, 8] gives the ultraviolet part in finite volume.

Remark 11.1. The following count relates (D) to a correlation length. Suppose that the gauge-fixed field responds to a change of one link on the boundary of a cube through a Poisson kernel with decay length ξ in lattice units. Then the response summed over the cube is of order ξ, so K_c is of order $n^3\xi$ and $D(n, \alpha)$ holds once $n \gtrsim \xi/\alpha$, that is, once $\ell \gtrsim \xi a/\alpha$, a fixed multiple of the physical correlation length. The hypothesis is that the gauge-fixed links decorrelate on the scale ξ, and it is a statement about the gauge, not about the gauge-invariant observables: the gauge here is built on straight lines, and in gauges of axial type the gauge-fixed variables can fluctuate on all scales while gauge-invariant observables cluster. Propositions 11.2 and 11.3 carry the count out exactly for Gaussian fields, and Proposition 11.3 shows that the quadratic approximation of the forest gauge fluctuates in this way. Where it does, (D) fails in this gauge and has to be posed for block-averaged variables, where it becomes part of the renormalisation group problem. Section 12.1 carries this out in the quadratic approximation.

The count of Remark 11.1 can be carried out exactly for a Gaussian field, the quadratic approximation of the gauge-fixed theory at weak coupling, and shows what (D) asks of the gauge-fixed variables.

Proposition 11.2. *Let $d \ge 1$, $b > 0$, $t \ge 0$, let $n \ge 2$ be even and $V = [0, n)^d \cap \mathbb{Z}^d$. Given $\eta \in \mathbb{R}^{\mathbb{Z}^d \setminus V}$, let $\gamma_V(\cdot \mid \eta)$ be the Gaussian law on $\mathbb{R}^V$ with density proportional to*

$$\exp\Big(-\frac{b}{2}\Big[\sum_{\{x,y\}\cap V \neq \emptyset} (\phi_x - \phi_y)^2 + t\sum_{x\in V}\phi_x^2\Big]\Big),$$

the first sum over nearest-neighbour pairs, with $\phi = \eta$ outside V. Define $C_{V,j}$, the classes c (residues modulo two) and K_c as above, with $d_V(\phi, \phi') = \sum_{x\in V}|\phi_x - \phi'_x|$ and $\delta(\eta_j, \eta'_j) = |\eta_j - \eta'_j|$, and put $N_n = (n/2)^d$. Then $K_c = \alpha_n(t)N_n$ for every class c, where

$$\alpha_n(t) = 1 - \frac{t}{n^d}\,\mathbf{1}^{\mathsf{T}}(t + L_n)^{-1}\mathbf{1}$$

and L_n is the Dirichlet Laplacian of V. In particular:

(i) *for $t = 0$, $K_c = N_n$ for every n and every b, so $D(n, \alpha)$ fails for every $\alpha < 1$;*

(ii) *for $t > 0$, $1 - \alpha_n(t) \geq (5/6)^d\, tn^2/(tn^2 + 10d)$; for $d = 4$, $t = m^2a^2$ and $\ell/2 \leq na \leq \ell$ this gives $D(n, \alpha)$ with $\alpha = 1 - (5/6)^4 (m\ell)^2/((m\ell)^2 + 160)$, uniformly in a.*

Proof. Let $A = t + L_n$ and $G = A^{-1}$, and let B be the matrix with $B_{xj} = 1$ when $x \in V$ and $j \notin V$ are neighbours. The law $\gamma_V(\cdot \mid \eta)$ has covariance $b^{-1}G$ and mean $GB\eta$. The matrix A has nonpositive off-diagonal entries and is positive definite, so $G \geq 0$ entrywise. Changing η_j by Δ translates the law by $\Delta\, GB_j$, where B_j is the column of j. The coupling by translation gives $W_1^{d_V} \leq |\Delta| \sum_x (GB_j)_x$, and the 1-Lipschitz function $\sum_x \phi_x$ gives the reverse inequality, so $C_{V,j} = \mathbf{1}^{\mathsf{T}} GB_j$. Since $B\mathbf{1} = L_n\mathbf{1}$ (both count the exterior neighbours of each site),

$$\sum_{j \notin V} C_{V,j} = \mathbf{1}^{\mathsf{T}} G L_n \mathbf{1} = \mathbf{1}^{\mathsf{T}}\mathbf{1} - t\, \mathbf{1}^{\mathsf{T}} G \mathbf{1} = n^d \alpha_n(t).$$

The reflection $x_i \mapsto n-1-x_i$ preserves V and the coefficients and, as n is even, exchanges the two residues of the coordinate x_i; so the 2^d class sums are equal, and each is $n^d\alpha_n(t)/2^d = \alpha_n(t) N_n$. This proves the formula and (i).

For (ii) let $v(i) = (i+1)(n-i)$ for $0 \leq i < n$ and $\psi(x) = \prod_i v(x_i)$. The one-dimensional Dirichlet Laplacian maps v to the constant 2, so with $s_1 = \sum_i v(i) = n(n+1)(n+2)/6$ and $s_2 = \sum_i v(i)^2 = n(n+1)(n+2)(n^2+2n+2)/30$,

$$\mathbf{1}^{\mathsf{T}}\psi = s_1^d, \qquad \psi^{\mathsf{T}} A \psi = t s_2^d + 2d s_1 s_2^{d-1}.$$

By the Cauchy–Schwarz inequality for the form of A^{-1}, $\mathbf{1}^{\mathsf{T}} G \mathbf{1} \geq (\mathbf{1}^{\mathsf{T}}\psi)^2/\psi^{\mathsf{T}} A \psi$, hence

$$1 - \alpha_n(t) \geq \Big(\frac{s_1^2}{n s_2}\Big)^d \frac{t}{t + 2d s_1/s_2} \geq \Big(\frac{5}{6}\Big)^d \frac{tn^2}{tn^2 + 10d},$$

since $s_1^2/(ns_2) = \frac{5}{6}(n^2+3n+2)/(n^2+2n+2) \geq \frac{5}{6}$ and $2ds_1/s_2 = 10d/(n^2+2n+2) \leq 10d/n^2$. With $d = 4$ and $tn^2 = m^2(na)^2 \geq (m\ell)^2/4$ the last fraction is at least $(m\ell)^2/((m\ell)^2 + 160)$. □

The Gaussian field is not Yang–Mills theory, and the proposition is a benchmark for the condition, not a statement about the Wilson measure. It shows that the averaged cube condition is met at a fixed physical scale by a field with a mass and fails exactly, at every size and every stiffness, for a massless one. Small local fluctuations alone therefore give nothing: in the massless Gaussian field on $\mathbb{Z}^d$, $d \geq 3$, the variance of $\phi_x - \phi_{x+e_1}$ is $1/(db)$, which tends to zero as $b \to \infty$, and still $K_c = N_n$. A proof of (D) along the tuning has to produce in the gauge-fixed variables the loss that the parameter t supplies here, from the Wilson action alone.

The quadratic approximation of the forest gauge itself, with the plaquette action in place of the Laplacian, is no better than the massless field of Proposition 11.2(i).

Proposition 11.3. *Let $b > 0$, let n be even and V the cube of Definition 10.1. For real functions φ on the links of $\mathbb{Z}^4$ that vanish on F, let $(d\varphi)_p$ be the sum of φ around the plaquette p with the signs of its orientation. Given $\eta \in \mathbb{R}^{A \setminus V}$, let $\gamma_V(\cdot \mid \eta)$ be the law on $\mathbb{R}^V$ with density proportional to*

$$\exp\Big(-\frac{b}{2} \sum_{p \cap V \neq \emptyset} (d\varphi)_p^2\Big),$$

with $\varphi = \eta$ on $A \setminus V$ and $\varphi = 0$ on F, and define $C_{V,j}$ and K_c as above with $d_V(\varphi, \varphi') = \sum_{i \in V} |\varphi_i - \varphi_i'|$ and $\delta(\eta_j, \eta_j') = |\eta_j - \eta_j'|$. Then

$$\sum_c K_c \geq |V| = 48 N_n,$$

so $K_c \geq N_n$ for some class c, and $D(n, \alpha)$ fails for every $\alpha < 1$, at every n and every b.

Proof. The laws. The quadratic form $\sum_{p\cap V\neq\emptyset}(d\varphi)_p^2$ with $\varphi=0$ outside V vanishes only at $\varphi=0$: the argument of Corollary 11.13, written additively, gives a function h on the sites with $\varphi_{(x,\mu)}=h(x)-h(x+e_\mu)$, constant outside the box and along the lines of F, hence constant. So $\gamma_V(\cdot\mid\eta)$ is a Gaussian law whose covariance does not depend on η and whose mean is $\sum_j \eta_j u_j$ for some vectors $u_j\in\mathbb{R}^V$. Two exteriors that differ in finitely many links give translates of one law by $t=\sum_j(\eta'_j-\eta_j)u_j$; the coupling by translation and the 1-Lipschitz function $\sum_i \operatorname{sign}(t_i)\varphi_i$ show that their distance is $\|t\|_1$. In particular $C_{V,j}=\|u_j\|_1$, and

$$W_1^{d_V}\left(\gamma_V(\cdot\mid\eta),\gamma_V(\cdot\mid\eta')\right)\le\sum_j C_{V,j}\,|\eta'_j-\eta_j|.$$

Lines. By Lemma 2.2(i) the lines of F are straight, disjoint and cover the sites, and the two ends of an active link (x,μ) lie on different lines, since the line through x has direction $\sigma(x)\neq\mu$. For a line ℓ let h_ℓ be the indicator of its sites and $(dh_\ell)_{(x,\mu)}=h_\ell(x)-h_\ell(x+e_\mu)$. Then dh_ℓ vanishes on F and around every plaquette, so the density is invariant under $\varphi\mapsto\varphi+dh_\ell$, and $\gamma_V(\cdot\mid dh_\ell)$ is $\gamma_V(\cdot\mid 0)$ translated by the restriction of dh_ℓ to V. Hence

$$\sum_{i\in V}|(dh_\ell)_i|\le\sum_{j\in A\setminus V}C_{V,j}\,|(dh_\ell)_j|.$$

Here $|(dh_\ell)_e|=1$ when exactly one end of e lies on ℓ and 0 otherwise, so $\sum_\ell|(dh_\ell)_e|=2$ for every active link e, the sum running over the finitely many lines that meet a link entering the inequality. Summing over these lines gives $2|V|\le 2\sum_j C_{V,j}=2\sum_c K_c$. □

The proof uses only that gauge transformations constant on the lines of F move the whole cube while changing few boundary links, which is the behaviour of gauges of axial type anticipated in Remark 11.1. The class sums themselves, computed exactly for $n\le 8$, lie between 1.57 and 11.24 times N_n and increase with n. Below the correlation length the theory is weakly coupled, and the quadratic approximation is its leading order there. A proof of (D) along the tuning therefore has to show that the nonperturbative regime reduces the cube coefficients of the forest gauge below N_n, or (D) has to be posed for other variables.

The lines of F are not what makes the argument work. A constant field, corrected by a periodic gauge transformation so that it vanishes on the fixed links, costs no action and changes every exterior link by a bounded amount, and this happens in every gauge fixed on an invariant forest.

Proposition 11.4. *Let $b>0$ and let F be a set of links of $\mathbb{Z}^4$, invariant under the translations by $2\mathbb{Z}^4$, that contains no closed path and whose trees are infinite and contain every site. Let A be the set of the other links, V the set of links $(x,\mu)\in A$ with $x\in[0,n)^4$, $n\ge 2$ even, and define the classes, $\gamma_V(\cdot\mid\eta)$, $C_{V,j}$ and K_c as in Proposition 11.3 with this F. Then there are numbers $a_c\ge 0$, not all zero, with*

$$\sum_c a_c K_c\ge N_n\sum_c a_c,$$

so $K_c\ge N_n$ for some class c, and $D(n,\alpha)$ fails for every $\alpha<1$, at every n and every b.

Proof. The laws. Let φ vanish outside V and on F, with $(d\varphi)_p=0$ for every plaquette p that meets V. The other plaquettes contain no link of V, so $d\varphi=0$ on every plaquette, and $\varphi=dh$ for a function h on the sites. Since φ vanishes outside a box and the complement of a box is connected, h is constant there, and we take that constant to be 0. As φ vanishes on F, h is constant on every tree of F; each tree is infinite and meets the complement of the box, so $h=0$ and $\varphi=0$. Hence, as in the proof of Proposition 11.3, $\gamma_V(\cdot\mid\eta)$ is a nondegenerate Gaussian law with mean $\sum_j\eta_j u_j$, where only the links j that share a plaquette with V contribute, and $C_{V,j}=\|u_j\|_1$.

A flat field. For $c\in\mathbb{R}^4$ and a function h on $\mathbb{Z}^4$ of period two in each coordinate put $\psi_{(x,\mu)}=c_\mu+h(x)-h(x+e_\mu)$. Then $d\psi=0$ on every plaquette and ψ is invariant under

$2\mathbb{Z}^4$; if $\psi = 0$, adding ψ at (x,μ) and $(x+e_\mu,\mu)$ gives $2c_\mu = 0$, and then h is constant. These fields form a space of dimension $16-1+4=19$. Let f be the number of orbits of $2\mathbb{Z}^4$ in F. The box $[0,2m)^4$ has $16m^4$ sites, and the links of F with both ends in it form a forest, so there are at most $16m^4-1$ of them, while each orbit contributes at least $(m-1)^4$; letting $m\to\infty$ gives $f\le 16$. Vanishing on F is therefore at most 16 linear conditions, and some $\psi\neq 0$ of this form vanishes on F. It is nonzero on some link of A. Put $a_c = |\psi_j|$ for $j\in c$.

The inequality. Let η be the restriction of ψ to $A\setminus V$. The field equal to ψ on V and to η outside gives the quadratic form its least value 0, so by nondegeneracy the mean of $\gamma_V(\cdot\mid\eta)$ is the restriction of ψ to V, and $\psi_i = \sum_j \psi_j (u_j)_i$ for $i\in V$. Hence

$$\sum_{i\in V}|\psi_i| \le \sum_{j\in A\setminus V} C_{V,j}|\psi_j| = \sum_c a_c K_c.$$

A class consists of the active links with a given direction and a given residue of the base point modulo two, so it meets V in N_n links, and the left side is $N_n\sum_c a_c > 0$. So $K_c \ge N_n$ for some class with $a_c>0$. □

For the direction field of Table 3 the flat fields vanishing on F form a space of dimension 7, and for the axial field $\sigma=0$ one of dimension 10. The proposition is weaker than Proposition 11.3 for the forest gauge, which bounds the sum over all classes, and it says nothing about two kinds of condition. A mass gives the constant fields an action, which is why Proposition 11.2 distinguishes $t>0$ from $t=0$. The constrained condition D_{c} of Theorem 12.8 holds the block averages fixed, and a constant field changes them. Within the quadratic approximation no gauge fixed on an invariant forest gives (D); the loss has to come from the Wilson action beyond its quadratic part, or (D) has to be posed for variables, such as the block averages, that a constant field moves.

Remark 11.5. Five other routes stop at (D) or (E), or earlier.

(a) *Disjoint blocks.* The block criterion of Lemma B.3 with a partition into cubes of side n requires, for a link on a face, that its influence on the neighbouring cube be less than one. In the count of Remark 11.1 that influence is of order ξ, which grows like $1/a$. Averaging over the N_n translates divides it by the number of translates on whose boundary the link lies, a factor of order $1/n$, and this is why (D) is stated for averaged cubes.

(b) *Rescaling a fixed lattice gap.* Any bound uniform in lattice units, such as those of Theorems 1.1–1.7, gives trivial continuum two-point functions [16]; rescaling it produces no gap.

(c) *Assuming the gap.* Hypothesis R of [16] assumes the gap in physical units together with the limit. Theorem 10.4 replaces the gap part by the cube estimate (D), which concerns one box and one coupling at each spacing.

(d) *Computation.* At a fixed coupling and a fixed small n, $D(n,\alpha)$ is a finite set of Kantorovich bounds and could in principle be certified as in Propositions 7.8 and 7.9; for $n=2$ and SU(2) the block already has 48 links and 144 real variables. (D) asks for it at infinitely many couplings with $n\to\infty$, which no finite computation covers.

(e) *Axioms alone.* Proposition 9.2 shows that positivity, invariance and regularity do not force a gap, and Proposition 9.1 shows that the growth route to rotation invariance is closed for the action density. The infrared input therefore has to be an estimate such as (D).

11.2. **Estimates uniform in the coupling.** Some local quantities of the Wilson measure can be controlled uniformly in β_W, the volume and the boundary condition, by integrating by parts in a single link. Such estimates hold along every tuning, but they concern local fields in lattice units, and the three results below show how far they are from the open inputs: the force has bounded moments but its centred form is a null vector of the reflected

form, the plaquette has fluctuations of order β_W^{-2} but this is a single-site variance, and bare links over a fixed physical distance become independent Haar variables.

Let $G = \mathrm{SU}(N)$, write $b = \beta_W$, so that the Wilson weight is e^{bS} with $S = \sum_p N^{-1} \operatorname{Re}\operatorname{Tr} Q_p$, and give $\mathfrak{g}$ the metric $\langle T, R\rangle = -\operatorname{Tr}(TR)$. For a link e let S_e be the sum of the six plaquette terms containing e, X_T the derivative along $U_e \mapsto e^{uT}U_e$ at $u = 0$, ∇_e and Δ_e the gradient and Laplacian in U_e, $q = N^2 - 1$, $c_N = (N^2-1)/N$, and

$$E_e = b|\nabla_e S_e|^2, \qquad W_e = E_e - c_N S_e = b|\nabla_e S_e|^2 + \Delta_e S_e.$$

Below, ω is a Wilson Gibbs state at $b > 0$: a finite-volume measure in which e is not fixed by the boundary condition, or an infinite-volume state whose conditional law of U_e given the other links is proportional to $e^{bS_e} dU_e$ with dU_e the Haar measure.

Proposition 11.6. (i) *For every unit $T \in \mathfrak{g}$, $t \in \mathbb{R}$ and $0 \le \eta < N/12$,*

$$\mathbb{E}_\omega\big[e^{t\sqrt{b}\,X_T S_e} \mid U_{e^c}\big] \le e^{3t^2/N},$$
$$\mathbb{E}_\omega\big[e^{\eta E_e} \mid U_{e^c}\big] \le (1 - 12\eta/N)^{-q/2},$$

so $\mathbb{E}_\omega E_e^m \le m!\,(24/N)^m 2^{q/2}$ for $m \ge 1$, uniformly in b, the volume and the boundary condition. Moreover $\mathbb{E}_\omega E_e = c_N \mathbb{E}_\omega S_e$.

(ii) *For a bounded function F of finitely many links, continuously differentiable in U_e, $\mathbb{E}_\omega[W_e F] = -\mathbb{E}_\omega\langle \nabla_e S_e, \nabla_e F\rangle$. In particular $\mathbb{E}_\omega[W_e F] = 0$ if F does not depend on U_e, and $\mathbb{E}_\omega[W_e W_f] = 0$ if no plaquette contains both e and f.*

(iii) *Let ω be reflection positive as in Appendix F, and let $A = \sum_e h_e W_e$ be a finite sum over links whose endpoints have times at least 2. Then $\omega(\vartheta(\bar{A})A) = 0$, so $\Psi_A = 0$: in the reconstructed Hilbert space the packets $\sum_e h_e E_e$ and $c_N \sum_e h_e S_e$ define the same vector.*

Proof. (i) Along $u \mapsto e^{uT}U_e$ a plaquette term containing e is $N^{-1}\operatorname{Re}\operatorname{Tr}(e^{uT}V)$ or $N^{-1}\operatorname{Re}\operatorname{Tr}(Ve^{-uT})$ with V unitary, whose second derivative is $N^{-1}\operatorname{Re}\operatorname{Tr}(T^2 e^{\pm uT}V)$. Since $-T^2 \ge 0$ and $\operatorname{Tr}(-T^2) = 1$, $|\operatorname{Tr}(T^2 M)| \le 1$ for unitary M, so the second derivative of S_e along the curve is at most $6/N$ in absolute value. Taylor's formula with $u = t/\sqrt{b}$ gives $b(S_e(e^{tT/\sqrt{b}}U_e) - S_e(U_e)) \ge t\sqrt{b}X_T S_e(U_e) - 3t^2/N$. The conditional density is e^{bS_e}/Z with respect to Haar measure, and the left invariance of Haar measure gives $\mathbb{E}_\omega[e^{b(S_e(e^{tT/\sqrt{b}}U_e) - S_e(U_e))} \mid U_{e^c}] = 1$, which with the Taylor bound proves the first inequality. For the second let G_0 be a standard Gaussian vector in $\mathfrak{g}$, independent of ω; then $e^{\eta|Y|^2} = \mathbb{E} e^{\sqrt{2\eta}\langle G_0, Y\rangle}$ for $Y = \sqrt{b}\nabla_e S_e$, and the first inequality in the direction of G_0 with $t = \sqrt{2\eta}|G_0|$ gives the bound $\mathbb{E}e^{6\eta|G_0|^2/N} = (1 - 12\eta/N)^{-q/2}$. With $\eta = N/24$ and $x^m \le m!\,\eta^{-m}e^{\eta x}$ the moment bound follows. For the mean, $\sum_A T_A^2 = -c_N I$ in an orthonormal basis gives $\Delta_e S_e = -c_N S_e$, and the integral of the divergence of $e^{bS_e}\nabla_e S_e$ over G vanishes.

(ii) The integral over U_e of the divergence of $e^{bS_e}F\nabla_e S_e$ vanishes, and that divergence is $e^{bS_e}(W_e F + \langle \nabla_e S_e, \nabla_e F\rangle)$; integrate the conditional identity. The function W_f depends only on the links of the plaquettes containing f.

(iii) Each W_e in the sum depends on links with times at least 1, so $\vartheta(\bar{A})$ depends only on links with times at most 0, none of which is a link e of the sum. By (ii), $\omega(\vartheta(\bar{A})W_e) = 0$ for each e. □

Proposition 11.7. *Let $G = \mathrm{SU}(2)$, $b \ge 0$, ω a Wilson Gibbs state at b and p a plaquette with a link that is not fixed by the boundary condition. Then*

$$\operatorname{Var}_\omega(\tfrac{1}{2}\operatorname{Tr} Q_p) \ge \frac{1}{2^{19}(1+6b)^2}, \quad \mathbb{E}_\omega[1 - \tfrac{1}{2}\operatorname{Tr} Q_p] \ge \frac{1}{4b+1}.$$

Proof. Let e be the link and identify $\mathrm{SU}(2)$ with the unit quaternions. Given the other links, $U_e = x$ has density proportional to $e^{rx\cdot u}$ with $|u| = 1$ and $0 \le r \le 6b$, since the six staples have norm one, and $\frac{1}{2}\operatorname{Tr} Q_p = x \cdot v$ for a unit vector v fixed by the conditioning.

Let $X = x \cdot u$ and $T = 1 - X$; the law of T has density proportional to $e^{-rt}\sqrt{t(2-t)}$ on $[0,2]$. Put $n = 1 + r$. Since $e^{-rt} \le e^2 e^{-nt}$, the normalisation is at most $e^2 \int_0^\infty \sqrt{2t} e^{-nt} dt = e^2\sqrt{\pi/2}\, n^{-3/2} < 14 n^{-3/2}$. On $I = [\frac{1}{4n}, \frac{1}{2n}]$ the density is at least $\frac{1}{2} n^{-1/2} e^{-1/2}$ and on $J = [\frac{1}{n}, \frac{3}{2n}]$ at least $\frac{1}{2} n^{-1/2} e^{-3/2}$, so $P(T \in I) \ge \frac{1}{224}$ and $P(T \in J) \ge \frac{1}{336}$. With an independent copy T', $\operatorname{Var} T = \frac{1}{2}\mathbb{E}(T - T')^2 \ge P(T \in I) P(T \in J)(2n)^{-2} > 2^{-19} n^{-2}$. By symmetry the conditional covariance of x is diagonal in u and $u^\perp$, with eigenvalues $\operatorname{Var} X$ and $\mathbb{E}(1 - X^2)/3$; on I, $1 - X^2 = T(2 - T) \ge \frac{3}{8n}$, so $\mathbb{E}(1 - X^2)/3 \ge \frac{1}{1792 n} \ge 2^{-19} n^{-2}$. Hence the conditional variance of $x \cdot v$ is at least $2^{-19}(1 + 6b)^{-2}$, and so is the variance, by the law of total variance. For the mean, integrating $\frac{d}{dX}(e^{rX}(1 - X^2)^{3/2})$ over $[-1, 1]$ gives $3\mathbb{E}X = r\mathbb{E}(1 - X^2) \le r(1 - (\mathbb{E}X)^2)$, whence $\mathbb{E}X \le (\sqrt{9 + 4r^2} - 3)/(2r) \le 2r/(2r + 3)$ for $r > 0$, and $\mathbb{E}[1 - x \cdot v] \ge 1 - \mathbb{E}X \ge 3/(2r + 3) \ge 1/(4b + 1)$. □

For the blocking statement let $G = \mathrm{SU}(2)$, $B \ge 4$ an integer and, for a coarse edge (x, μ), $V_{(x,\mu)} = \prod_{r=0}^{B-1} U_{(Bx + r\hat{\mu}, \mu)}$ the ordered product of the fine links along it.

Proposition 11.8. *For every Wilson Gibbs state ω of $\mathrm{SU}(2)$ at $b > 0$ and every set $\mathcal{E}$ of K coarse edges,*

$$\begin{aligned} \big\| \mathrm{Law}_\omega\big((V_e)_{e \in \mathcal{E}}\big) - m^{\otimes K} \big\|_{\mathrm{TV}} &\le \frac{K}{2} e^{3b} \Big(\frac{4b}{4b+1} \Big)^{B-3} \\ &\le \frac{K}{2} \exp\Big(3b - \frac{B-3}{4b+1} \Big), \end{aligned}$$

where m is the Haar measure. If F is a function of the V_e of a set of coarse edges at positive times, $|F| \le 1$, and ϵ is this bound for those edges and their reflections, then $0 \le \omega(\vartheta(\bar{F}^c) F^c) \le 6\epsilon$ with $F^c = F - \omega(F)$.

Proof. Select on each coarse edge the fine links with $1 \le r \le B - 2$. No plaquette contains two selected links: collinear links do not share a plaquette, parallel coarse edges are B apart, and a plaquette containing a selected link of (x, μ) has its links in direction $\nu \ne \mu$ at μ-offsets $r, r + 1 \in [1, B - 1]$ from Bx, while coarse edges in direction ν lie at offsets in $B\mathbb{Z}$. Given the other links, the selected ones are therefore independent, each with density $e^{r_i\, x \cdot u_i}/Z(r_i)$, $0 \le r_i \le 6b$. Convolution with such a density acts on the representation of spin $l \ge 1$ by $D^{(l)}(u_i)$ times $I_{l+1}(r_i)/I_1(r_i)$, whose absolute value is at most $I_2(r_i)/I_1(r_i)$ because $I_\nu(r)$ decreases in $\nu \ge 0$, and $I_2(r_i)/I_1(r_i) = \mathbb{E}X \le 4b/(4b + 1)$ by the proof of Proposition 11.7, so it contracts the mean-zero part of L^2 by that factor. The density is at most e^{6b} because $Z \ge 1$ by Jensen's inequality, so its distance to 1 in L^2 is at most e^{3b}. The conditional law of V_e is that of a product of $B - 2$ such independent variables between two fixed ones, so its total variation distance to m is at most $\frac{1}{2} e^{3b} (4b/(4b + 1))^{B-3}$, and the coarse edges are conditionally independent. Total variation distance from a product is at most the sum over the factors, and averaging over the conditioning does not increase it. For the last statement, under $m^{\otimes K}$ the covariance vanishes because F and its reflection depend on disjoint edges; replacing ω by $m^{\otimes K}$ changes $\omega(\vartheta(\bar{F}) F)$ by at most 2ϵ and $|\omega(F)|^2$ by at most 4ϵ. □

With $B = \ell/a_k$ for a fixed physical length ℓ and $b_k = O(\log(1/a_k))$ the bound tends to zero: the bare coarse links of a fixed coarse graph become independent Haar variables, their reflected form has rank one, and a renormalisation Z_k that keeps the reflected norm of $Z_k F^c$ at least $c > 0$ must grow at least like $(c/6\epsilon)^{1/2}$. The renormalised flowed fields of Theorem 10.7 are not bounded functions of bare coarse links, and none of the three propositions bears on the slice ratio, the tree bounds or (D).

11.3. **What the cube estimate requires.** This subsection proves seven facts about $D(n, \alpha)$. It holds at small coupling for every even n above an explicit size (Proposition 11.10), and it survives small perturbations of the action (Proposition 11.11), although gauge-invariant perturbations still need strong coupling at the scale of the cube (Proposition 11.18).

It forces the gauge-fixed variables of a cube with trivial exterior to stay a fixed distance from the identity on average (Theorem 11.12, and Proposition 11.14 for SU(2)), so that for every fixed n it fails once β is large (Corollary 11.13). And it cannot be obtained from bounds on the conditional law of a single line of links that are uniform in the configuration around the line (Proposition 11.22). For the compact abelian theory with the Villain action it fails at every size once β is large (Proposition 11.19). In Section 12.1 the estimate is posed for block averages, where its quadratic approximation holds along the tuning. None of these decides (D) along the continuum tuning, where n grows while β grows.[19]

For a finite $V \subset A$ and $j \in A \setminus V$ let $b(V, j)$ be the number of links of V that share a plaquette with j, and write $k \sim i$ when two active links share a plaquette.

Lemma 11.9. *Suppose that $C_{ee'} \le c_0$ for all active e, e', with $12c_0 < 1$. Then $C_{V,j} \le c_0\, b(V,j)/(1-12c_0)$ for every finite $V \subset A$ and every $j \in A \setminus V$.*

Proof. Fix Q, Q' that differ only at j, put $d = \delta(Q_j, Q'_j)$, $\mu = \gamma_V(\cdot \mid Q)$ and $\nu = \gamma_V(\cdot \mid Q')$. The couplings of μ and ν form a weakly compact set on which $\pi \mapsto \int d_V \, d\pi$ is continuous, so one of them, π, minimises it; let $\rho_i = \int \delta(X_i, Y_i)\, d\pi$. Fix $i \in V$. Under μ the conditional law of X_i given the other links of V is $\gamma_i^F(\cdot \mid X)$, and likewise for ν. Replace (X_i, Y_i) by a pair drawn from an optimal coupling of $\gamma_i^F(\cdot \mid X)$ and $\gamma_i^F(\cdot \mid Y)$, chosen measurably [80, Cor. 5.22], and keep the other coordinates. The new coupling still has marginals μ and ν, and changing the links in which X and Y differ one at a time gives

$$W_1\big(\gamma_i^F(\cdot \mid X), \gamma_i^F(\cdot \mid Y)\big) \le \sum_{k \in V,\, k \ne i} C_{ik}\, \delta(X_k, Y_k) + C_{ij}\, d.$$

Minimality of π therefore gives $\rho_i \le c_0 \sum_{k \in V,\, k \sim i} \rho_k + c_0 [i \sim j]\, d$. Sum over $i \in V$: every ρ_k occurs at most twelve times by Lemma 2.2(iv), so $(1 - 12c_0) \sum_i \rho_i \le c_0\, b(V,j)\, d$. Since $W_1^{d_V}(\mu, \nu) \le \sum_i \rho_i$, dividing by d proves the lemma. □

Proposition 11.10. *Let $n = 2m$ and let V be the cube of Definition 10.1. For every class c,*

$$\sum_{j \in c \setminus V} b(V, j) \le 10m^3 - 2m^2,$$

with equality for three classes. Consequently, if $C_{ee'} \le c_0$ for all e, e' with $12c_0 < 1$, then $D(n, \alpha)$ holds for every $\alpha \in (0,1)$ with

$$\alpha \ge \hat\alpha_n = \frac{c_0}{1 - 12c_0}\Big(\frac{20}{n} - \frac{8}{n^2}\Big).$$

For SU(2) *and $|\beta_W| < \frac{1}{3}$ this reads $\hat\alpha_n = \frac{|\beta_W|}{1 - 3|\beta_W|}\big(\frac{5}{n} - \frac{2}{n^2}\big)$, which is below one for every even $n > 5|\beta_W|/(1 - 3|\beta_W|)$.*

Proof. Let $j = (y, \mu)$ lie in the class c, with $y \equiv r$ modulo two. A link of V that shares a plaquette with j has base point $y + d$ with $d \in \{-1, 0, 1\}^4$ and at most two nonzero coordinates, and the twelve offsets d depend only on c. For a fixed offset we count the $y \equiv r$ with $y + d \in [0, n)^4$ and $y \notin [0, n)^4$. In the coordinate l there are m values of $y_l \equiv r_l$ with $y_l + d_l \in [0, n)$, and one of them lies outside $[0, n)$ exactly when $d_l = 1$ and $r_l = 1$ (the value $y_l = -1$) or $d_l = -1$ and $r_l = 0$ (the value $y_l = n$). If q is the number of such coordinates, the count is $m^4 - m^{4-q}(m-1)^q$, that is 0, m^3 or $2m^3 - m^2$ for $q = 0, 1, 2$. Summing over the twelve offsets, the class sum is $a_c m^3 - b_c m^2$, where b_c is the number of offsets with $q = 2$ and a_c is the number with $q = 1$ plus twice b_c. Reading the offsets from Table 3, the pairs (a_c, b_c) over the 48 classes are $(2,0)$, $(7,0)$, $(7,1)$, $(9,1)$, $(9,2)$ and $(10,2)$ three

[19] The four statements answer the question of which parts of (D) can be settled with the methods of this paper. The first places (D) in the range where the single-link method already works, the second and third show what any proof of (D) at weak coupling has to produce, and the fourth excludes the most direct one-dimensional route.

times each and $(3,0)$, $(4,0)$, $(5,0)$, $(6,0)$ and $(8,1)$ six times each. In every case $a_c \le 10$ and $8 - a_c + b_c \ge 0$, so

$$(10m^3 - 2m^2) - (a_c m^3 - b_c m^2) = m^2((10 - a_c)m + b_c - 2) \ge m^2(8 - a_c + b_c) \ge 0,$$

with equality for the three classes with $(a_c, b_c) = (10, 2)$. By Lemma 11.9, $K_c \le c_0(10m^3 - 2m^2)/(1 - 12c_0)$, and $N_n = m^4$ gives $K_c/N_n \le \hat{\alpha}_n$. For SU(2) the proof of Theorem 1.1 gives $c_0 = |\beta| = |\beta_W|/4$. □

At $\beta_W = \frac{3}{10}$ and $n = 16$, for instance, $\hat{\alpha}_{16} = \frac{117}{128}$, and Theorem 10.2 gives the gap $\log \frac{128}{117}/32$, much weaker than Theorem 1.1. The proposition shows that the cube condition is not empty; it does not extend the range, because the absorption $12c_0 < 1$ is the single-link condition itself.

The cube condition is also stable under small changes of the action, which is what a block renormalisation would need at its last scale, where the effective action is no longer of Wilson form. Let $r \ge 0$ and $\varepsilon \ge 0$, and let (w_X) be real functions on G^A indexed by finite sets $X \subset A$ of diameter at most r, with w_X depending only on the links of X, such that the family is invariant under the translations by $2\mathbb{Z}^4$, every active link lies in at most M of the sets X, every X has at most s links, and

$$|w_X(Q) - w_X(Q')| \le \varepsilon \sum_{i \in X} \delta(Q_i, Q'_i)$$

for all Q, Q'. Let $\gamma_V^W(\cdot \mid Q)$ be the probability measure with density proportional to $\exp \sum_{X \cap V \ne \emptyset} w_X$ with respect to $\gamma_V(\cdot \mid Q)$, and define $C_{V,j}^W$, K_c^W and the condition $D^W(n, \alpha)$ as in Definition 10.1, with γ_V^W in place of γ_V.

Proposition 11.11. *Let n be even and suppose that $D(n, \alpha)$ holds at β. Then $D^W(n, \alpha')$ holds with*

$$\alpha' = \alpha\, e^{\Omega}(1 + 2\Omega) + 96\, s\, \Omega, \qquad \Omega = 48\, M\, D\, N_n\, \varepsilon.$$

In particular $\alpha' < 1$ for all ε below an explicit $\varepsilon_0(n, \alpha, M, s) > 0$.

Proof. The cube has $|V| = 48N_n$ links, so d_V has diameter $\Delta_V = 48N_nD$. For fixed exterior links put $u = \sum_{X \cap V \ne \emptyset} w_X$ as a function on G^V. Each link of V lies in at most M sets X, so u is $M\varepsilon$-Lipschitz for d_V and its oscillation is at most $M\varepsilon\Delta_V = \Omega$; we normalise u so that $\sup u = 0$, which does not change the tilted measures, and put $h = e^u$, so that $e^{-\Omega} \le h \le 1$ and h is $M\varepsilon$-Lipschitz.

A common tilt. Let μ, ν be probability measures on G^V and $\mu_h = h\mu/\mu(h)$, $\nu_h = h\nu/\nu(h)$. Let f be 1-Lipschitz for d_V; subtracting a constant we may take $|f| \le \Delta_V$. Then fh is $(1 + \Delta_V M\varepsilon)$-Lipschitz, $|\nu(fh)| \le \Delta_V \nu(h)$, $|\mu(h) - \nu(h)| \le M\varepsilon W_1^{d_V}(\mu, \nu)$ and $\mu(h) \ge e^{-\Omega}$, so

$$\begin{aligned} \mu_h(f) - \nu_h(f) &= \frac{\mu(fh) - \nu(fh)}{\mu(h)} + \nu(fh)\, \frac{\nu(h) - \mu(h)}{\mu(h)\nu(h)} \\ &\le e^{\Omega}(1 + 2\Omega)\, W_1^{d_V}(\mu, \nu), \end{aligned}$$

since $\Delta_V M\varepsilon = \Omega$. By Kantorovich duality $W_1^{d_V}(\mu_h, \nu_h) \le e^{\Omega}(1 + 2\Omega) W_1^{d_V}(\mu, \nu)$.

Two tilts. If u' is another weight with $|u - u'| \le \eta$, the density of $\nu_{h'}$ with respect to ν_h lies in $[e^{-2\eta}, e^{2\eta}]$, so the total variation distance of the two is at most $1 - e^{-2\eta} \le 2\eta$, and $W_1^{d_V}(\nu_h, \nu_{h'}) \le 2\eta\Delta_V$.

The coefficients. Let Q, Q' differ only at $j \notin V$, with $\delta = \delta(Q_j, Q'_j)$, and let u, u' be the weights for the two exteriors. They differ only through the sets X that contain j and meet V; if there are m_j of them, $|u - u'| \le m_j \varepsilon \delta$. With $\mu = \gamma_V(\cdot \mid Q)$ and $\nu = \gamma_V(\cdot \mid Q')$, the triangle inequality and the two steps give

$$C_{V,j}^W \le e^{\Omega}(1 + 2\Omega)\, C_{V,j} + 2\Delta_V\, m_j\, \varepsilon.$$

The pairs (X, j) with $j \in X \setminus V$ and $X \cap V \neq \emptyset$ number at most s times the number of sets meeting V, hence at most $sM|V|$. Summing over $j \in c \setminus V$,

$$K_c^W \leq e^{\Omega}(1 + 2\Omega)\,\alpha N_n + 2\Delta_V sM|V|\varepsilon = \big(\alpha e^{\Omega}(1 + 2\Omega) + 96\,s\,\Omega\big)N_n.$$

The family is invariant under $2\mathbb{Z}^4$, so the coefficients are, and this is $D^W(n, \alpha')$. □

The admissible perturbation is of order $1/N_n = (2/n)^4$, so the proposition is useful in a box of a fixed number of lattice spacings, which is how a renormalisation argument would use it: after enough block steps that the cube of physical size ℓ has become a cube of fixed side, the effective action has to lie within ε_0 of an action that satisfies the condition there. Whether the effective actions along the tuning do so is not decided here, and that is where the problem lies: at the scale where the blocked theory is strongly coupled enough for a condition of this kind, the running coupling is of order one, which is outside the range of the existing ultraviolet analysis. The proposition shows only that the condition itself is not the obstacle.

The next theorem uses the configuration in which every link outside V is the identity. Write $\gamma_V^1 = \gamma_V(\cdot \mid 1)$ and

$$\varepsilon_V(\beta) = \frac{1}{|V|} \sum_{i \in V} \int \delta(1, Q_i)\, d\gamma_V^1(Q)$$

for the mean distance of the gauge-fixed variables of the cube from the identity. Recall that δ is bi-invariant of diameter D.

Theorem 11.12. *Let n be even, V the cube of Definition 10.1, and suppose that $D(n, \alpha)$ holds at β. Then*

$$\varepsilon_V(\beta) \geq \frac{(1 - \alpha)\, D}{3}.$$

Proof. A flat configuration. By Lemma 2.2(i) and the period two of σ, the sixteen residues modulo two fall into eight pairs $\{r, r + e_{\sigma(r)}\}$, the residues met by one line of F. The eight residues 0010, 0011, 0100, 0110, 1001, 1011, 1100, 1101 form four of these pairs. Choose $z, w \in G$ with $\delta(z, w) = D$, put $g(x) = w$ if x is congruent to one of those eight residues and $g(x) = z$ otherwise, and let $Q^g_{(x,\mu)} = g(x)g(x + e_\mu)^{-1}$ for every link of $\mathbb{Z}^4$. Then g is constant on every line of F, so Q^g equals 1 on F, every plaquette variable of Q^g is 1, and Q^g has period two. By bi-invariance $\delta(1, Q^g_{(x,\mu)}) = \delta(g(x + e_\mu), g(x))$, which is D when x and $x + e_\mu$ receive different values and 0 otherwise; from Table 3 the first case occurs for exactly 32 of the 48 classes. Let δ_c be the common value of $\delta(1, Q^g_e)$ for e in the class c, so that $\sum_c \delta_c = 32D$.

Gauge covariance. Let $\Phi(Q)_{(x,\mu)} = g(x)Q_{(x,\mu)}g(x + e_\mu)^{-1}$ on G^V. The gauge transformation by g maps the configuration equal to Q_V on V and to 1 elsewhere to the one equal to $\Phi(Q_V)$ on V and to Q^g elsewhere; it leaves every plaquette trace unchanged and every link of F equal to 1, and Φ preserves Haar measure. Hence $\gamma_V(\cdot \mid Q^g) = \Phi_*\gamma_V^1$.

Upper bound. Only the finitely many links outside V that share a plaquette with V enter γ_V. Changing them one at a time from 1 to Q^g and applying the definition of $C_{V,j}$ at each step,

$$W_1^{d_V}\big(\gamma_V^1, \gamma_V(\cdot \mid Q^g)\big) \leq \sum_{j \in A \setminus V} C_{V,j}\, \delta(1, Q_j^g) = \sum_c \delta_c K_c \leq \alpha N_n \sum_c \delta_c = 32\,\alpha D N_n.$$

Lower bound. The function $f(Q) = -\sum_{i \in V} \delta(Q_i, Q_i^g)$ is 1-Lipschitz for d_V. By gauge covariance and bi-invariance, $\int f\, d\gamma_V(\cdot \mid Q^g) = -\int \sum_i \delta(Q_i, 1)\, d\gamma_V^1 = -|V|\varepsilon_V$, and by the triangle inequality $\int f\, d\gamma_V^1 \leq -\sum_i \delta(1, Q_i^g) + |V|\varepsilon_V$. Every class has N_n links in V, so $\sum_i \delta(1, Q_i^g) = 32DN_n$ and

$$W_1^{d_V}\big(\gamma_V^1, \gamma_V(\cdot \mid Q^g)\big) \geq 32DN_n - 2|V|\varepsilon_V.$$

With $|V| = 48N_n$ the two bounds give $96N_n\varepsilon_V \geq 32(1 - \alpha)DN_n$. □

Corollary 11.13. *For every even n and every $\alpha < 1$ there is $\beta_0(n,\alpha)$ such that $D(n,\alpha)$ fails at every $\beta \geq \beta_0(n,\alpha)$. In particular, if $D(n_k,\alpha)$ holds at couplings $\beta_k \to \infty$, then $n_k \to \infty$.*

Proof. For $\beta > 0$, γ_V^1 has density proportional to $e^{-N\beta S}$ with respect to Haar measure, where $S = \sum_{p\cap V\neq\emptyset}(N - \operatorname{Re}\operatorname{Tr} Q_p^F)$ is continuous and nonnegative on G^V and vanishes exactly when $Q_p^F = 1$ for every plaquette p that meets V. We show that this forces $Q_V = 1$. The other plaquettes have all their links equal to 1, so every plaquette variable is 1; since the plaquettes of $\mathbb{Z}^4$ fill every closed lattice path, there is a function h on the sites with $Q_{(x,\mu)} = h(x)h(x+e_\mu)^{-1}$ on every link, and h takes equal values at the two ends of every link on which $Q = 1$. Every link with base point outside $[0,n)^4$ is such a link. From a site with a coordinate at least n one can move in every positive direction along links of this kind, and from a site with a negative coordinate x_ν one can move in the positive directions other than ν until another coordinate reaches n; so h is constant on the sites outside the box. Every line of F leaves the box, and h is constant along it, so h is constant and $Q_V = 1$.

Let U be a neighbourhood of 1 in G^V, $s = \min_{G^V\setminus U} S > 0$, and $U' \subset U$ a neighbourhood of 1 on which $S < s/2$. Then $\gamma_V^1(G^V \setminus U) \leq e^{-N\beta s}/(|U'|e^{-N\beta s/2})$, which tends to zero, so $\varepsilon_V(\beta) \to 0$ as $\beta \to \infty$, and Theorem 11.12 excludes $D(n,\alpha)$ once $\varepsilon_V(\beta) < (1-\alpha)D/3$. □

For SU(2) the constant of Theorem 11.12 can be improved: a flat configuration with four values instead of two moves every class of active links, and the means of the gauge-fixed links are scalar.

Proposition 11.14. *Let $G = \mathrm{SU}(2)$ with the chordal metric, let n be even, V the cube of Definition 10.1, and suppose that $D(n,\alpha)$ holds at β. Then*

$$\frac{1}{|V|}\sum_{i\in V}\Big|\int \operatorname{Re}\operatorname{Tr} Q_i \, d\gamma_V^1\Big| \leq 2\alpha,$$

$$\frac{1}{|V|}\sum_{i\in V}\int \delta(1,Q_i)^2 \, d\gamma_V^1 \geq 2(1-\alpha),$$

and $\varepsilon_V(\beta) \geq 1-\alpha$.

Proof. Write $\langle\cdot,\cdot\rangle$ for the Euclidean product of quaternions; then $\operatorname{Re}\operatorname{Tr} q = 2\langle q,1\rangle$ and $\delta(p,q) = |p-q|$.

Four colours. Split the sixteen residues modulo two into the sets

$$X_1 = \{0000, 1000, 0111, 1111\},$$
$$X_2 = \{0001, 0101, 1010, 1110\},$$
$$X_3 = \{0010, 0011, 1100, 1101\},$$
$$X_4 = \{0100, 0110, 1001, 1011\}.$$

Each is the union of two of the eight pairs $\{r, r+e_{\sigma(r)}\}$, and Table 3 shows that r and $r+e_\mu$ lie in different sets for every residue r and every $\mu \neq \sigma(r)$. (The eight residues used in the proof of Theorem 11.12 form $X_3 \cup X_4$.) Let

$$z_1 = (\mathbf{i}+\mathbf{j}+\mathbf{k})/\sqrt{3}, \quad z_2 = (\mathbf{i}-\mathbf{j}-\mathbf{k})/\sqrt{3},$$
$$z_3 = (-\mathbf{i}+\mathbf{j}-\mathbf{k})/\sqrt{3}, \quad z_4 = (-\mathbf{i}-\mathbf{j}+\mathbf{k})/\sqrt{3},$$

unit quaternions with $\langle z_k, z_l\rangle = -\frac{1}{3}$ for $k \neq l$, so that $\delta(z_k,z_l) = d_* = (8/3)^{1/2}$. Put $g(x) = z_k$ when the residue of x lies in X_k, and $Q^g_{(x,\mu)} = g(x)g(x+e_\mu)^{-1}$. As in the proof of Theorem 11.12, Q^g equals 1 on F, every plaquette variable of Q^g is 1, and $\gamma_V(\cdot \mid Q^g) = \Phi_*\gamma_V^1$ with $\Phi(Q)_{(x,\mu)} = g(x)Q_{(x,\mu)}g(x+e_\mu)^{-1}$; now $\delta(1,Q^g_e) = d_*$ for every active link e.

Upper bound. Changing the exterior links one at a time,

$$W_1^{d_V}\left(\gamma_V^1, \gamma_V(\cdot \mid Q^g)\right) \leq d_* \sum_c K_c \leq 48\,\alpha\, d_* N_n = \alpha\, d_* |V|.$$

Lower bound. Conjugating every link of V by one unit quaternion leaves the exterior 1, the links of F, Haar measure and every plaquette trace unchanged, so it preserves γ_V^1. The mean $m_i = \int Q_i \, d\gamma_V^1$, taken in the space of quaternions, therefore commutes with every unit quaternion, and $m_i = a_i 1$ with $a_i = \frac{1}{2}\int \operatorname{Re}\operatorname{Tr} Q_i \, d\gamma_V^1$ real. Since Φ acts on each link by a fixed left and right multiplication, the mean of Q_i under $\gamma_V(\cdot \mid Q^g)$ is $a_i Q_i^g$. Choose unit quaternions w_i with $\langle w_i, a_i Q_i^g - a_i 1\rangle = |a_i|\,|Q_i^g - 1|$. The function $f(Q) = \sum_{i\in V}\langle w_i, Q_i\rangle$ is 1-Lipschitz for d_V, and Kantorovich duality gives

$$W_1^{d_V}\big(\gamma_V^1, \gamma_V(\cdot \mid Q^g)\big) \geq \sum_{i\in V} |a_i|\,\delta(1, Q_i^g) = d_* \sum_{i\in V} |a_i|.$$

Conclusion. Comparing the two bounds, $\sum_i |a_i| \leq \alpha|V|$, which is the first inequality. Since $\delta(1,q)^2 = 2 - 2\langle q, 1\rangle$, $\int \delta(1,Q_i)^2 \, d\gamma_V^1 = 2(1 - a_i) \geq 2(1 - |a_i|)$, and averaging gives the second. As $\delta \leq 2$, $\delta \geq \delta^2/2$, and the bound on ε_V follows. □

The four colours need nothing from SU(2) except four elements whose pairwise quotients are conjugate, and every compact simple group contains such elements. For a metric that comes from a representation the bound then compares the cube with Haar measure.

Proposition 11.15. *Let G be a compact simple group, ρ a nontrivial irreducible unitary representation of G of dimension d, and $\delta(p,q) = \kappa^{-1/2}\|\rho(p) - \rho(q)\|_{\mathrm{HS}}$ for some $\kappa > 0$. Let n be even, V the cube of Definition 10.1, and suppose that $D(n,\alpha)$ holds at β. Then*

$$\frac{1}{|V|}\sum_{i\in V}\Big|\int \operatorname{Tr}\rho(Q_i)\, d\gamma_V^1\Big| \leq \alpha\, d,$$

$$\frac{1}{|V|}\sum_{i\in V}\int \delta(1,Q_i)^2 \, d\gamma_V^1 \geq (1-\alpha)\int_G \delta(1,q)^2 \, dq,$$

where dq is Haar measure, and $\varepsilon_V(\beta) \geq (1-\alpha)D/2$.

Proof. Four elements. Let $\varphi\colon \mathrm{SU}(2) \to G$ be the homomorphism of the SU(2) subgroup of a root, and $z_1, \dots, z_4$ the tetrahedral quaternions of Proposition 11.14. For $k \neq l$ the quaternion $z_k z_l^{-1}$ has real part $\langle z_k, z_l\rangle = -\frac{1}{3}$; unit quaternions with equal real parts are conjugate in SU(2), so the twelve elements $\varphi(z_k z_l^{-1})$ are conjugate in G, and by bi-invariance $\delta(\varphi(z_k), \varphi(z_l)) = d_*$ for all $k \neq l$, with one number d_*. If $d_* = 0$, then $\varphi(z_k z_l^{-1})$ lies in $\ker\rho$, a proper normal subgroup of the simple group G and therefore central; it then commutes with $\varphi(\mathrm{SU}(2))$, so it is central there, and $z_k z_l^{-1}$ lies in $\{\pm 1\}\ker\varphi \subset \{\pm 1\}$, which its real part $-\frac{1}{3}$ excludes. So $d_* > 0$.

The two bounds. Put $g(x) = \varphi(z_k)$ when the residue of x lies in X_k. The proof of Proposition 11.14 applies word for word: Q^g equals 1 on F and on every plaquette, $\delta(1, Q_e^g) = d_*$ on every active link, and $W_1^{d_V}\big(\gamma_V^1, \gamma_V(\cdot \mid Q^g)\big) \leq \alpha d_* |V|$. Simultaneous conjugation of the links of V preserves γ_V^1, so the mean $\int \rho(Q_i)\, d\gamma_V^1$ commutes with $\rho(G)$, and by Schur's lemma it is $c_i I$ with $c_i = d^{-1}\int \operatorname{Tr}\rho(Q_i)\, d\gamma_V^1$. Its mean under $\gamma_V(\cdot \mid Q^g)$ is $c_i\rho(Q_i^g)$. With matrices W_i of Hilbert–Schmidt norm $\kappa^{-1/2}$ chosen so that $\operatorname{Re}\operatorname{Tr}(W_i^* c_i(\rho(Q_i^g) - I)) = |c_i|\,\delta(1, Q_i^g)$, the function $f(Q) = \sum_i \operatorname{Re}\operatorname{Tr}(W_i^*\rho(Q_i))$ is 1-Lipschitz for d_V, and Kantorovich duality gives $W_1^{d_V} \geq d_* \sum_i |c_i|$. Hence $\sum_i |c_i| \leq \alpha|V|$, the first inequality.

Comparison with Haar measure. Since $\kappa\,\delta(1,q)^2 = 2d - 2\operatorname{Re}\operatorname{Tr}\rho(q)$ and $\int_G \operatorname{Tr}\rho(q)\, dq = 0$ for a nontrivial irreducible ρ, $\int_G \delta(1,q)^2 dq = 2d/\kappa$, while $\int \delta(1,Q_i)^2 d\gamma_V^1 = 2d(1 - \operatorname{Re} c_i)/\kappa \geq 2d(1 - |c_i|)/\kappa$. Averaging gives the second inequality. Finally $\kappa D^2 \leq 4d$, because $\|\rho(p) - \rho(q)\|_{\mathrm{HS}} \leq 2\sqrt{d}$, so $\delta \geq \delta^2/D$ gives $\varepsilon_V \geq 2d(1-\alpha)/(\kappa D) \geq (1-\alpha)D/2$. □

For SU(N) with the Hilbert–Schmidt distance this improves the constant $\frac{1}{3}$ of Theorem 11.12 to $\frac{1}{2}$, and for SU(3), where $D^2 = 9$ and the Haar mean of δ^2 is 6, to $\frac{2}{3}$: in a cube with the identity outside, $D(n,\alpha)$ leaves the gauge-fixed links at least the fraction $1 - \alpha$ of the disorder of independent Haar-distributed links.

The size $\beta_0(n,\alpha)$ in Corollary 11.13 comes from a compactness argument. For SU(2) it can be made explicit, and then reads as a lower bound on the side of any cube that satisfies the condition.

Proposition 11.16. *Let $G=\mathrm{SU}(2)$ with the chordal metric, so that $D=2$, let $b=\beta_W\ge\frac19$, let $n\ge2$ be even and put $L(b)=1+\log 24+\frac32\log(9b)$. Then*

$$\frac{1}{|V|}\sum_{e\in V}\int\delta(1,Q_e)^2\,d\gamma_V^1\le\frac{n^2(n+2)(2n+1)\,L(b)}{b},$$

so $\varepsilon_V(\beta)^2$ is at most the same quantity. Consequently, if $D(n,\alpha)$ holds at b, then

$$n^4\ge\frac{2(1-\alpha)\,b}{5\,L(b)}.$$

Proof. Put $m=n/2$. Under γ_V^1 every link outside V equals 1: the links of F always, the others by the choice of exterior. A plaquette that does not meet V therefore has plaquette variable 1.

Links as loops. Let $e=(x,\mu)\in V$, $y=x+e_\mu$, $\sigma=\sigma(x)$ and $\tau=\sigma(y)$. The directions σ,τ,μ are distinct: $\sigma\ne\mu$ because e is active; $\tau\ne\mu$ because otherwise the line of F through y would contain x, and the lines of F are disjoint; and $\tau\ne\sigma$ by Table 3, in which $\sigma(x+e_\mu)\ne\sigma(x)$ whenever $\mu\ne\sigma(x)$. Since $0\le x_\sigma<n$, one of $n-x_\sigma$ and $x_\sigma+1$ is at most m; call it s and let $\epsilon=\pm1$ be the sign for which $x'=x+s\epsilon e_\sigma$ has σ-coordinate outside $[0,n)$. Choose $t\le m$ and ϵ' in the same way for y and τ (note $y_\tau=x_\tau$). Let R_1 be the s plaquettes of the rectangle with corners x, x', $x'+e_\mu$, y in the plane $\{\sigma,\mu\}$, and R_2 the st plaquettes of the rectangle with corners y, $x'+e_\mu$, $x'+e_\mu+t\epsilon' e_\tau$, $y+t\epsilon' e_\tau$ in the plane $\{\sigma,\tau\}$. The union $R(e)=R_1\cup R_2$ is a disc, and its boundary runs from x along the line of F through x to x', then to $x'+e_\mu$, along e_τ to $x'+e_\mu+t\epsilon' e_\tau$, along e_σ to $y+t\epsilon' e_\tau$, back along the line of F through y to y, and over e to x. The links of the first and fifth pieces lie in F. The links of the second and third pieces have base points with σ-coordinate $x'_\sigma\notin[0,n)$, and those of the fourth have base points with τ-coordinate outside $[0,n)$; none of them lies in V. So the holonomy of the boundary of $R(e)$ is $Q_e^{\pm1}$. Adding the plaquettes of a disc one at a time along its boundary shows that this holonomy is a product of conjugates of the plaquette variables $(Q_p^F)^{\pm1}$, $p\in R(e)$, one for each plaquette. By bi-invariance and the triangle inequality,

$$\delta(1,Q_e)\le\sum_{p\in R(e)}\delta(1,Q_p^F),\qquad |R(e)|=s(t+1)\le m(m+1).$$

Multiplicity. If a plaquette p lies in $R_2(e)$, its plane is $\{\sigma,\tau\}$, μ is one of the two other directions, and the base point of p is $y+ae_\sigma+ce_\tau$ with $-m\le a,c\le m-1$. Given p, there are two ways to name the directions of its plane, two choices of μ and $(2m)^2$ of (a,c), and they determine y, hence e; so p lies in at most $16m^2$ of the sets $R_2(e)$. Likewise $p\in R_1(e)$ forces its plane to be $\{\sigma,\mu\}$ and its base point to be $x+ae_\sigma$ with $-m\le a\le m-1$, which leaves at most $4m$ links e. Every plaquette therefore lies in at most $16m^2+4m=4n^2+2n$ of the discs $R(e)$.

The energy. For the chordal metric $\delta(1,q)^2=2-\mathrm{Re}\,\mathrm{Tr}\,q$. Let $S=\sum_{p\cap V\ne\emptyset}\delta(1,Q_p^F)^2$; then γ_V^1 has density $e^{-bS/2}/Z$ with $Z=\int e^{-bS/2}$, the integral over $\mathrm{SU}(2)^V$ with Haar measure. The function $t\mapsto\log\int e^{-tS}$ is convex and vanishes at $t=0$, so its derivative at $t=b/2$, which is $-\int S\,d\gamma_V^1$, is at least $2\log Z/b$. A plaquette that meets V contains at most three links of V and the other links are 1, so $\delta(1,Q_p^F)^2\le3\sum_{e\in p\cap V}\delta(1,Q_e)^2$; every link lies in six plaquettes, so $S\le18\sum_{e\in V}\delta(1,Q_e)^2$ and $Z\ge I^{|V|}$ with $I=\int e^{-9b\,\delta(1,q)^2}dq$. The chordal distance is at most the geodesic one, and the geodesic ball of radius $r=(9b)^{-1/2}\le1$ has measure greater than $r^3/24$ (proof of Proposition 11.22), so $I\ge e^{-1}(9b)^{-3/2}/24$ and

$$\int S\,d\gamma_V^1\le\frac{2|V|\,L(b)}{b}.$$

Conclusion. The Cauchy–Schwarz inequality inside each disc squares the loop bound:

$$\delta(1,Q_e)^2 \le |R(e)| \sum_{p\in R(e)} \delta(1,Q_p^F)^2 \le m(m+1) \sum_{p\in R(e)} \delta(1,Q_p^F)^2.$$

Summing over $e \in V$ and using the multiplicity,

$$\sum_{e\in V} \delta(1,Q_e)^2 \le m(m+1)(4n^2+2n)\, S = \frac{n^2(n+2)(2n+1)}{2}\, S,$$

and integrating against γ_V^1 gives the first bound; the bound on ε_V follows by the Cauchy–Schwarz inequality. If $D(n,\alpha)$ holds, Proposition 11.14 gives $2(1-\alpha) \le n^2(n+2)(2n+1)L(b)/b$, and $(n+2)(2n+1) \le 5n^2$ for $n \ge 2$. □

The loops and their multiplicity use nothing of SU(2) but the bi-invariance of δ, and the energy uses only the form of the action and the measure of small balls. The bound therefore holds for every SU(N), with the same power of n.

Proposition 11.17. *Let $G = \mathrm{SU}(N)$, $N \ge 2$, with the weight $\exp(\frac{b}{N}\sum_p \mathrm{Re}\,\mathrm{Tr}\, Q_p)$, $b = \beta_W \ge N/9$, and the Hilbert–Schmidt distance $\delta(p,q) = \|p-q\|_{\mathrm{HS}}$. Let $n \ge 2$ be even and put*

$$L_N(b) = 1 + \frac{N^2-1}{2}\log(81\pi^2 b) + N(N-1)\log\frac{12}{11}.$$

Then

$$\frac{1}{|V|}\sum_{e\in V}\int \delta(1,Q_e)^2\, d\gamma_V^1 \le \frac{N\, n^2(n+2)(2n+1)\, L_N(b)}{b}.$$

Consequently, if $D(n,\alpha)$ holds at b, then

$$n^4 \ge \frac{2(1-\alpha)\, b}{5\, L_N(b)}.$$

Proof. Loops. The discs $R(e)$ of the proof of Proposition 11.16 and the bound $4n^2+2n$ on their multiplicity do not depend on the group, and the loop bound uses only bi-invariance and the triangle inequality. With $S = \sum_{p\cap V\neq\emptyset} \delta(1,Q_p^F)^2$ the same Cauchy–Schwarz step gives $\sum_{e\in V}\delta(1,Q_e)^2 \le \frac{1}{2}n^2(n+2)(2n+1)\, S$.

The energy. Since $\delta(1,q)^2 = 2N - 2\,\mathrm{Re}\,\mathrm{Tr}\, q$, the measure γ_V^1 has density $e^{-bS/(2N)}/Z$ with respect to Haar measure. As in the proof of Proposition 11.16, the convexity of $t \mapsto \log\int e^{-tS}$ gives $\int S\, d\gamma_V^1 \le 2N\log(1/Z)/b$, and $S \le 18\sum_{e\in V}\delta(1,Q_e)^2$ gives $Z \ge I^{|V|}$ with $I = \int e^{-9b\,\delta(1,q)^2/N} dq$.

Small balls. Give $\mathfrak{su}(N)$ the inner product $\mathrm{Tr}(X^*Y)$ and SU(N) the corresponding bi-invariant metric, of volume v_N. In exponential coordinates the Riemannian measure is $j(X)\, dX$, where dX is Lebesgue measure and

$$j(X) = \Big|\det\frac{1-e^{-\operatorname{ad} X}}{\operatorname{ad} X}\Big| = \prod_{k<l}\Big(\frac{\sin(\varphi_{kl}/2)}{\varphi_{kl}/2}\Big)^2$$

if X has eigenvalues $i\theta_1,\dots,i\theta_N$ and $\varphi_{kl} = \theta_k - \theta_l$ [44, Ch. II, Thm. 1.7]; in particular $j \le 1$. Every $q \in \mathrm{SU}(N)$ is conjugate to a diagonal matrix with entries $e^{i\psi_k}$, $-\pi \le \psi_k < \pi$; then $\sum_k \psi_k = 2\pi k_0$ with $|k_0| \le N/2$, and changing $|k_0|$ of the ψ_k by $\mp 2\pi$ gives angles θ_k with sum zero and $|\theta_k| \le 3\pi$. So $q = e^X$ with $X \in \mathfrak{su}(N)$ and $\|X\|_{\mathrm{HS}} \le 3\pi\sqrt{N}$, and $v_N \le \omega_{N^2-1}(3\pi\sqrt{N})^{N^2-1}$, where ω_k is the volume of the unit ball of $\mathbb{R}^k$. Now let $\|X\|_{\mathrm{HS}} \le r \le 1$. The eigenvalues have $|\theta_k| \le 1 < \pi$, so exp is injective on this ball; $\delta(1,e^X)^2 = \sum_k |e^{i\theta_k}-1|^2 \le \|X\|_{\mathrm{HS}}^2$; and $|\varphi_{kl}|/2 \le (\theta_k^2+\theta_l^2)^{1/2}/\sqrt{2} \le 2^{-1/2}$, so each factor

of $j(X)$ is at least $(1-\frac{1}{12})^2$ because $\sin x/x \geq 1 - x^2/6$. Dividing by v_N, the Haar measure of $\{q : \delta(1,q) \leq r\}$ is at least

$$\Big(\frac{11}{12}\Big)^{N(N-1)} \frac{\omega_{N^2-1} r^{N^2-1}}{v_N} \geq \Big(\frac{11}{12}\Big)^{N(N-1)} \Big(\frac{r}{3\pi\sqrt{N}}\Big)^{N^2-1}.$$

With $r = (N/9b)^{1/2} \leq 1$ this gives $I \geq e^{-1}$ times the right-hand side, and since $3\pi\sqrt{N}/r = 9\pi\sqrt{b}$, $\log(1/I) \leq L_N(b)$. Hence $\int S\, d\gamma_V^1 \leq 2N|V|L_N(b)/b$, and the first bound follows from the loops.

Conclusion. Proposition 11.15 with the defining representation and $\kappa = 1$, for which the Haar mean of $\delta(1,q)^2$ is $2N$, gives $2N(1-\alpha) \leq N\, n^2(n+2)(2n+1)L_N(b)/b$ when $D(n,\alpha)$ holds, and $(n+2)(2n+1) \leq 5n^2$. □

For every N, a cube on which the condition holds therefore has side at least of order $(\beta_W/\log\beta_W)^{1/4}$, with a constant that depends on N only through L_N, which grows like $\frac{N^2}{2}\log\beta_W$.

The cube condition is a statement about strong coupling at its own scale, and this persists under the perturbations of Proposition 11.11 when they respect the gauge. Call (w_X) *gauge invariant* if $w_X(Q^h) = w_X(Q)$ for every function h on the sites that is constant on every line of F, where $Q^h_{(x,\mu)} = h(x)Q_{(x,\mu)}h(x+e_\mu)^{-1}$. Such transformations keep the links of F equal to 1, so they act on G^A, and the restriction to the gauge of an action invariant under all gauge transformations is gauge invariant in this sense. Write ε_V^W for ε_V with $\gamma_V^W(\cdot \mid 1)$ in place of γ_V^1.

Proposition 11.18. *Let (w_X) be as in Proposition 11.11 and gauge invariant, let $n = 2m$, and suppose that $D^W(n,\alpha)$ holds.*

(i) $\varepsilon_V^W \geq (1-\alpha)D/3$.

(ii) *If $\eta \geq 0$, $0 \leq \kappa \leq 1$, and the event that $\delta(1, Q_p^F) \leq \eta$ for every plaquette p that meets V has probability at least $1-\kappa$ under $\gamma_V^W(\cdot \mid 1)$, then*

$$m(m+1)\,\eta + \kappa D \geq \frac{(1-\alpha)D}{3}.$$

Proof. (i) The proof of Theorem 11.12 uses the action only through the identity $\gamma_V(\cdot \mid Q^g) = \Phi_*\gamma_V^1$ and through the coefficients $C_{V,j}$. The function g of that proof is constant on every line of F, so the gauge transformation by g leaves every w_X unchanged, as it leaves every plaquette trace, and the same identity holds for γ_V^W. Only finitely many links outside V enter γ_V^W, because the sets X have bounded diameter, so the upper bound holds with $C_{V,j}^W$ and K_c^W in place of $C_{V,j}$ and K_c; the lower bound is unchanged.

(ii) Under $\gamma_V^W(\cdot \mid 1)$ every link outside V equals 1, so a plaquette that does not meet V has variable 1. The loop bound of the proof of Proposition 11.16, which uses only the bi-invariance of δ, gives $\delta(1,Q_e) \leq \sum_{p\in R(e)} \delta(1,Q_p^F) \leq m(m+1)\eta$ for every $e \in V$ on the event of the statement, and $\delta(1,Q_e) \leq D$ off it. Averaging over $e \in V$ and integrating, $\varepsilon_V^W \leq m(m+1)\eta + \kappa D$, and (i) completes the proof. □

In the small-field region of a gauge-covariant block renormalisation the blocked plaquette variables lie within a distance η that tends to zero with the running coupling, and large fields have small probability κ. If $\kappa < (1-\alpha)/3$, part (ii) allows the cube condition on the blocked lattice only when $m(m+1) \geq x$, where $x = ((1-\alpha)/3 - \kappa)D/\eta$, so the side must grow like $\eta^{-1/2}$. On such a cube $N_n = m^4 \geq x^2/4$, and since $\alpha' \geq \alpha + 96s\Omega$, Proposition 11.11 gives $\alpha' < 1$ only for

$$\begin{aligned}\varepsilon &< \frac{1-\alpha}{4608\, sMD\, N_n} \\ &\leq \frac{(1-\alpha)\,\eta^2}{1152\, sMD^3\big((1-\alpha)/3-\kappa\big)^2},\end{aligned}$$

which is of order η^2. At a fixed side the condition is therefore a statement about strong coupling, and a renormalisation argument has to carry the blocked theory out of the small-field regime before the condition can be checked. That passage, at a running coupling of order one, is the step that remains open.

The next statement shows that the cube condition separates a theory with a massless photon from the expected behaviour of SU(N): for the compact abelian theory in its Coulomb phase it fails at every scale. Let $G = \mathrm{U}(1)$ with the chordal metric, $D = 2$, and replace the Wilson weight by the Villain weight

$$\varphi_\beta(\theta) = \sum_{k\in\mathbb{Z}} e^{-\beta(\theta+2\pi k)^2/2},$$

so that the measure of a box Λ has density proportional to $\prod_{p\in P(\Lambda)} \varphi_\beta(\theta_p)$, where $e^{i\theta_p}$ is the plaquette variable. Define γ^F, $C_{V,j}$, K_c and $D(n,\alpha)$ as before with φ_β in place of the Wilson weight.

Proposition 11.19. *There is β_0 such that for the Villain theory at every $\beta > \beta_0$ the condition $D(n,\alpha)$ fails for every even $n \geq 2$ and every $\alpha < 1$. In particular, for couplings $\beta_k \to \infty$ and any even n_k, the hypothesis of Theorem 10.4 fails for this theory.*

Proof. Lemma 3.1, Proposition 3.2(i)–(iii) and (v), Lemma B.1, Corollary B.2 and Lemma B.4 use the Wilson weight only as a positive continuous class function of the plaquette variable, with one exception: the Doeblin bound in Lemma B.4(a), which for the Villain weight reads that each single-link density is at least $(\min\varphi_\beta/\max\varphi_\beta)^6 > 0$ times its maximum. Suppose that $D(n,\alpha)$ holds. As in the proof of Theorem 10.2, the limit $\omega(O) = \lim W_\Lambda(O)$ of the measures of boxes with free boundary conditions then exists along every sequence of boxes exhausting $\mathbb{Z}^4$, and for gauge-invariant local O, O' with O Lipschitz in each link,

$$|\operatorname{Cov}_\omega(O,O')| \leq \|O'\|_\infty L_O \,\frac{D(1+c_n e^{\lambda\rho_n})}{1-\theta}\,|S_{O'}|\,e^{-\lambda\,\mathrm{dist}(O,O')}$$

for every $\lambda > 0$ with $\theta = \alpha e^{\lambda\rho_n} < 1$, where L_O is a Lipschitz constant of O and $S_{O'}$ the support of O'.

Put $\Psi_p = \varphi'_\beta(\theta_p)/\varphi_\beta(\theta_p)$; up to the factor $-\mathbf{i}$ this is the lattice field strength of Fröhlich and Spencer [38, Sect. 2] for the Villain action. It is real, odd and smooth in θ_p, so it is gauge invariant, local and Lipschitz in each of its four links, with constants that do not depend on p, and $\omega(\Psi_p) = 0$ by the symmetry $\theta \mapsto -\theta$. The bound above gives $\sum_{p'} |\omega(\Psi_p\Psi_{p'})| < \infty$, the sum over the plaquettes p' parallel to p. The state of [38] is the limit of the same free-boundary measures over boxes, which is ω. Fröhlich and Spencer prove, for β larger than some β_0, that the Fourier transform of the two-plaquette function of the field strength is discontinuous at zero momentum, so that this sum is infinite [38, Sect. 2]; see also [43]. This contradiction proves the first statement, and the second follows because $\beta_k > \beta_0$ for all large k. □

At large β the plaquette variables of the abelian theory are Gaussian to leading order, as are those of SU(N), and every property of the action used in this paper (compactness, locality, positivity and smoothness of the weight, reflection positivity, the forest gauge) is shared by the two theories. A proof of (D) along the continuum tuning therefore has to use the non-abelian structure of G; Proposition 11.19 holds at every cube size, so letting the side grow with β, which Corollary 11.13 already requires, does not remove the need. Fröhlich and Spencer indicate how their result extends to the Wilson action without giving the details [38, Sect. 2], so the proposition is stated for the Villain action only.

Remark 11.20. The proof loses the area of a disc, of order n^2, in passing from plaquettes to links, and on loops that loss is of the right order. On the periodic lattice $(\mathbb{Z}/L\mathbb{Z})^4$ with L even and coordinates $0 \leq x_j < L$, let $\phi = 4\pi/L^2$, $q(\theta) = \cos\theta + \mathbf{i}\sin\theta$, and put

$U_{(x,2)} = q(\phi x_1)$, $U_{(x,1)} = q(-\phi L x_2)$ when $x_1 = L-1$, and $U = 1$ on every other link. Every plaquette of the plane $\{1,2\}$ equals $q(\phi)$, including those on the seam $x_1 = L-1$ and at the corner $x_1 = x_2 = L-1$, where the angle is $\phi - 4\pi$; all other plaquettes equal 1. So every plaquette variable lies within ϕ of the identity, while the boundary of the square $[0, L/2]^2$ of the plane $\{1,2\}$ has holonomy $q(\phi L^2/4) = -1$, at the largest possible distance $D = 2$, against the bound $\phi L^2/4 = \pi$ given by the disc. Every staple sum of this configuration is parallel to its link, so it is a stationary point of the Wilson action, and gradient flow [58] does not change it. Plaquette variables within $O(n^{-2})$ of the identity therefore do not force loops of length n near the identity, and an argument of the kind above that places gauge-fixed links near the identity needs plaquette bounds of order $o(n^{-2})$.

Up to the factor $L(b)^{1/2}$ the rate $b^{-1/2}$ of Proposition 11.16 is the right one, and the lower bound below holds for every n and every exterior.

Proposition 11.21. *Let $G = \mathrm{SU}(2)$ with the chordal metric and $b = \beta_W > 0$. For every Gibbs specification of the gauge-fixed theory, every active link e and every configuration of the other links, the conditional law of Q_e satisfies*

$$\int \delta(1, Q_e)\, d\mu_e \geq \frac{1}{9 \max(6b, 1)^{1/2}}.$$

In particular $\varepsilon_V(\beta) \geq \frac{1}{9}\max(6b,1)^{-1/2}$ for every n; at every fixed n, $\varepsilon_V(\beta)$ lies between constant multiples of $b^{-1/2}$ and $(L(b)/b)^{1/2}$.

Proof. Write $\langle \cdot, \cdot \rangle$ for the Euclidean product of quaternions, so that $\operatorname{Re}\operatorname{Tr} q = 2\langle q, 1\rangle$. The six plaquettes that contain e contribute $\exp(b\langle Q_e, W_p\rangle)$ to the conditional density, with $|W_p| = 1$; so μ_e has density $e^{\langle u,v\rangle}/Z(|v|)$ with respect to normalised Haar measure, for some v with $a = |v| \leq 6b$. Put $a' = \max(a,1)$ and $\eta = a'^{-1/2} \leq 1$. On the geodesic ball of radius η about $v/|v|$ the exponent is at least $a(1-\eta^2/2) \geq a - \frac{1}{2}$, and that ball has measure greater than $\eta^3/24$ (proof of Proposition 11.22); hence $Z(a) \geq e^{a-1/2}\eta^3/24$ and the density is at most $24 e^{1/2} a'^{3/2}$ (for $a < 1$ directly, since $Z \geq 1$ by Jensen's inequality). A chordal ball of radius r is a geodesic ball of radius θ with $2\sin(\theta/2) = r$, so $\theta \leq \pi r/2$, and its measure is $(\theta - \sin\theta\cos\theta)/\pi \leq 2\theta^3/(3\pi) \leq \pi^2 r^3/12$. With $r = \frac{2}{9}a'^{-1/2}$,

$$\mu_e\big(\delta(1,Q_e) < r\big) \leq 24 e^{1/2} a'^{3/2} \frac{\pi^2 r^3}{12} = 2e^{1/2}\pi^2 \Big(\frac{2}{9}\Big)^3 < \frac{1}{2},$$

so the mean of $\delta(1,Q_e)$ is at least $r/2 = \frac{1}{9}a'^{-1/2}$. Averaging over $e \in V$ and over the configuration gives the bound on ε_V. □

Along the continuum tuning Theorem 11.12 says the following. If (D) holds, then in a cube of fixed physical size ℓ with the identity outside, the forest-gauge variables stay on average at distance at least $(1-\alpha)D/3$ from the identity, and at least $1-\alpha$ for SU(2), at every spacing. This is not a contradiction: $n = \ell/a$ grows like $1/a$, while asymptotic freedom makes $\beta(a)$ grow only like $\log(1/a)$, and the corollary concerns fixed n. The size bound of Proposition 11.16 is likewise met along the tuning, since its right side grows only like $\log(1/a)$. It turns the hypothesis of Remark 11.1, that the gauge-fixed links decorrelate on the scale of the cube, into a necessary condition: (D) requires the forest-gauge variables to be disordered at a fixed physical scale. For gauge-invariant observables of a massive theory that disorder is clustering; for the gauge-fixed variables it is not proved.

The last statement concerns the following strategy: control the conditional law of one line of links, given everything else, uniformly in the configuration around it, and assemble the cube from lines. Fix a site x, let $s = \sigma(x)$ and $\mu \neq s$, and let $c_k = (x + ke_s, \mu)$, $k \in \mathbb{Z}$. All c_k are active, since σ is constant along the line through x.

Proposition 11.22. *Let $G = \mathrm{SU}(2)$, identified with the unit quaternions, and $b = \beta_W \geq 1$. There is a configuration of the active links outside $\{c_k\}$, independent of b, for which the*

conditional law of $(Q_{c_k})_k$ has a ten-step transfer operator, similar to a positive self-adjoint operator K_b on $L^2(S^3)$ with simple top eigenvalue Λ_b, whose gap satisfies

$$0 < 1 - \frac{\Lambda_{b,1}}{\Lambda_b} \le e^{20}\, 24^{10}\, b^{15}\, e^{-b/1280},$$

where $\Lambda_{b,1}$ is the next point of the spectrum.

Proof. The interactions. By Lemma 2.2(iii), c_k lies in two plaquettes of the plane $\{\mu, s\}$, which contain c_{k-1} or c_{k+1}, the link $(x+ke_s, s)$ or $(x+(k-1)e_s, s)$ of F, and an active twist link of direction s; and in four plaquettes of the planes $\{\mu, t\}$ and $\{\mu, \rho\}$, each containing c_k, one link of F and two active links not on the line. The four plaquettes at c_k lie in the hyperplane of s-coordinate $x_s + k$, so these plaquettes at different k share no link; two plaquettes through c_k share no other link; and their links have directions other than s, so they are not twists. Set every twist to 1; then the plaquette at $x + ke_s$ in the plane $\{\mu, s\}$ contributes $b\langle q_k, q_{k+1}\rangle$ to the exponent, with $q_k = Q_{c_k}$. In each of the four other plaquettes at c_k set one of its two links to 1 and the other to X; after a cyclic rotation and possibly an inversion the plaquette variable is $q_k X^{\pm 1}$ or $q_k^{-1} X^{\pm 1}$, and since $\mathrm{Re}(q\bar{u}) = \mathrm{Re}(\bar{q}u) = \langle q, u\rangle$ for unit quaternions, X can be chosen so that the plaquette contributes $b\langle q_k, u\rangle$ for any prescribed unit vector u. Every other active link is set to 1. Four unit vectors add up to any vector of length at most 4 (two copies each of $t \pm (1-|t|^2)^{1/2}\nu$ with $t = v/4$ and $\nu \perp t$ a unit vector), so the conditional density of the line is proportional to $\exp\big(b\sum_k \langle q_k, q_{k+1}\rangle + \sum_k \langle h_k, q_k\rangle\big)$ for any fields h_k with $|h_k| \le 4b$.

The fields. Let $a = \frac{1}{2}$, $c = \frac{\sqrt{3}}{2}$, let $e_1, \dots, e_4$ be the standard basis of $\mathbb{R}^4$, and let r_k have period ten with $(r_0, \dots, r_9) = a(e_1, -e_1, e_2, -e_2, e_3, -e_3, e_3, -e_2, e_2, -e_1)$. Put $h_k = b(2r_k - r_{k-1} - r_{k+1})$, so $|h_k| \le 4ab = 2b$, and the configuration is independent of b because h_k/b is. With $v_k = q_k - r_k$ and $|q_k| = 1$, $|r_k| = a$, summation by parts gives, over $0 \le k \le 9$,

$$\frac{1}{2}\sum_k |v_{k+1} - v_k|^2 = \sum_k \big(1 - \langle q_k, q_{k+1}\rangle\big) - \frac{1}{b}\sum_k \langle h_k, q_k\rangle + \frac{1}{2}\sum_k |r_{k+1} - r_k|^2 + B(q_0) - B(q_{10}),$$

with $B(q) = \langle q, r_0 - r_9\rangle$. The boundary terms only conjugate the ten-step transfer kernel by a multiplication operator, so up to similarity it is

$$K_b(x,y) = \int_{(S^3)^9} \exp\Big(-\frac{b}{2}\sum_{k=0}^{9} |v_{k+1} - v_k|^2\Big)\, dq_1 \cdots dq_9, \qquad q_0 = x,\ q_{10} = y,$$

with normalised Haar measure. The sequence $r_0, \dots, r_{10}$ is a palindrome, so $K_b = B_b B_b^*$ for the five-step kernel B_b; K_b is compact, positive and self-adjoint with a continuous positive kernel, its top eigenvalue Λ_b is simple, and its eigenfunction $\psi_b > 0$ is even under the reflection $x_4 \mapsto -x_4$, which commutes with K_b.

A barrier. Let $E = \frac{1}{2}\sum_{k=0}^{9} |v_{k+1} - v_k|^2$ and suppose that $(q_0)_4 \ge 0 \ge (q_{10})_4$ and $E \le \frac{1}{1280}$. Put $\Delta = (20E)^{1/2} \le \frac{1}{8}$ and $w = v_0$; by Cauchy–Schwarz $|w - v_k| \le \Delta$ for all k, and $0 \le w_4 \le \Delta$ because the r_k have no fourth coordinate. From $|v_k + r_k| = 1$, $\big||w + r_k|^2 - 1\big| \le \Delta(2+\Delta) =: \eta$ for $0 \le k \le 5$. The pairs $r_k = \pm a e_l$ give $|w_l| \le \eta/(2a) = \eta$ for $l = 1, 2, 3$ and $|w|^2 \ge \frac{3}{4} - \eta$, so $\frac{3}{4} \le 3\eta^2 + \Delta^2 + \eta \le 3(\frac{17}{64})^2 + \frac{1}{64} + \frac{17}{64} = \frac{2019}{4096}$, a contradiction. Hence $K_b(x,y) \le e^{-b/1280}$ whenever $x_4 \ge 0 \ge y_4$.

The top eigenvalue. Let $\epsilon = b^{-1/2}$. The geodesic ball of radius $\epsilon \le 1$ in S^3 has normalised measure $\frac{2}{\pi}\int_0^{\epsilon} \sin^2\theta\, d\theta \ge \frac{8\epsilon^3}{3\pi^3} > \frac{\epsilon^3}{24}$. If each q_k lies in the ball of radius ϵ about $r_k + ce_4$, then $|v_k - ce_4| \le \epsilon$ and $E \le 20\epsilon^2$, so the integrand is at least e^{-20}. Testing K_b on the normalised indicator of the ball about $r_0 + ce_4$ gives $\Lambda_b \ge e^{-20}(\epsilon^3/24)^{10} = e^{-20} 24^{-10} b^{-15}$.

The gap. Let $\phi(x) = \mathrm{sign}(x_4)\psi_b(x)$, a unit vector orthogonal to ψ_b. Then

$$\Lambda_b - \langle \phi, K_b \phi\rangle = 4\int_{x_4 > 0 > y_4} \psi_b(x) K_b(x,y) \psi_b(y)\, dx\, dy \le e^{-b/1280},$$

because $\int_{x_4>0}\psi_b \le \frac{1}{2}$ by Cauchy–Schwarz. By the variational principle $\Lambda_{b,1} \ge \langle \phi, K_b\phi\rangle$, and dividing by the lower bound on Λ_b proves the estimate; the gap is positive because Λ_b is simple. □

The exterior of Proposition 11.22 is legal at every coupling, so a bound on the conditional law of single lines that is uniform in the exterior cannot be better than $e^{-b/1280}$ up to powers of b, and an argument of this kind cannot give (D) along the tuning. A proof of (D) has to use what such a bound discards: the joint resampling of the transverse links inside the cube, which the averaged coefficients $C_{V,j}$ retain, or the smallness of the probability of such exteriors under the Wilson measure.

12. Block averages

Proposition 11.3 locates the failure of (D) in the quadratic approximation: gauge transformations constant on the lines of F move the whole cube and change few links outside it. By Proposition 11.4 every gauge fixed on an invariant forest fails in the same way, through the constant fields. This section takes up the programme of the report [56] of the third author, which keeps the cube condition but changes the gauge and conditions the cube on block averages of the field, as in Bałaban's renormalisation group [6, 7]. The natural block averages of the line field leave one class of lines invisible and fail at every size (Proposition 12.1). In a gauge whose fixed links are trees inside blocks of side two, the averages over whole blocks remove every zero mode, the constrained quadratic form has a gap that does not depend on the volume (Theorem 12.4), and the constrained cube condition holds at every coupling once the cube is large (Theorem 12.8). Integrating out one scale bounds the cube coefficients by the constrained ones plus those of the averages (Corollary 12.16), and the averages carry the coarse quadratic form at every scale up to a factor between 1 and 256, through forms that are local uniformly in the scale (Proposition 12.9 and Theorems 12.12 and 12.14), and the constrained cube condition holds at every scale with one cube side (Theorem 12.18). For the group-valued links themselves, the straight paths between block roots satisfy a coercivity bound for all fields but do not compose, a continuous covariant map cannot treat the transverse positions alike, and a cut-off of Bałaban's averaging operation is coercive for all fields and agrees with the block averages to first order (Section 12.2). It composes across all scales on fields of small curvature for every compact group (Section 12.3), and under the SU(2) Wilson measure the fields that are not small form clusters of vanishing physical size (Section 12.5).

12.1. **Block-constrained cubes.** In this subsection everything concerns the quadratic approximation: real fields φ on links, $(d\varphi)_p$ the signed sum around the plaquette p, and $(dh)_{(x,\mu)} = h(x) - h(x+e_\mu)$ for a function h on sites.

Proposition 12.1 ([56])**.** *In the setting of Proposition 11.3, condition the cube in addition on any family of the face averages*

$$V^{\mathrm{face}}_{y,\mu}(\varphi) = \frac{1}{8}\sum_{x}\bigl(\varphi_{(x,\mu)} + \varphi_{(x+e_\mu,\mu)}\bigr),$$

where x runs over the eight sites of the block $2y + \{0,1\}^4$ with $x_\mu = 2y_\mu$. Then $\sum_{c\in C_I} K_c \ge 12N_n$ for the twelve classes

$$C_I = \{(0111,\mu),(1111,\mu),(0111+e_\mu,\mu),(1111+e_\mu,\mu) : \mu = 1,2,3\},$$

written as residue of the base and direction. Hence $K_c \ge N_n$ for some class, and the constrained condition fails for every $\alpha < 1$, at every even n and every b.

Proof. By Table 3, $\sigma(0111) = \sigma(1111) = 0$, so the lines of F through the sites with x_1, x_2, x_3 odd have direction 0; call them invisible. Let ℓ be one of them and h_ℓ its indicator. The sum of dh_ℓ along a path of length two from x is $h_\ell(x) - h_\ell(x+2e_\mu)$. For $\mu \ne 0$ both end points

have even μ-coordinate and are off ℓ; for $\mu = 0$ they have the same transverse coordinates and are both on ℓ or both off it. So dh_ℓ changes no face average, and the translation argument of Proposition 11.3 applies to the constrained laws: $\sum_{i\in V} |(dh_\ell)_i| \le \sum_{j\notin V} C_{V,j}|(dh_\ell)_j|$. An active link $e = (x,\mu)$ has $|(dh_\ell)_e| = 1$ for at most one invisible ℓ, since its two ends lie on different lines and both cannot be invisible (for $\mu \neq 0$ they differ in the coordinate μ, and for $\mu = 0$ the link would lie in F). It has one such ℓ exactly when its base or its end has residue 0111 or 1111, which by Table 3 are the classes of C_I: the six classes with base residue 0111 or 1111 and direction $\mu \neq 0$, and the six with base residue $r + e_\mu$, $r \in \{0111, 1111\}$, whose directions μ differ from $\sigma(r + e_\mu)$ (for instance $\sigma(0011) = 3 \neq 1$). Summing over the invisible lines gives $12N_n \le \sum_{c\in C_I} K_c$. □

Definition 12.2. The blocks are $B(y) = 2y + \{0,1\}^4$, $y \in \mathbb{Z}^4$, and $o = x \bmod 2$ is the offset of a site. A link (x,μ) is a *tree link* if $o_\mu = 0$ and $o_\nu = 0$ for every $\nu > \mu$, and *active* otherwise; fields vanish on tree links. For a coarse link (y,μ) the *block average* is

$$V_{y,\mu}(\varphi) = \frac{1}{16} \sum_{x\in B(y)} \big(\varphi_{(x,\mu)} + \varphi_{(x+e_\mu,\mu)}\big),$$

the mean of φ summed along the sixteen paths of length two in the direction μ that start in $B(y)$.

Lemma 12.3. (i) *The fifteen tree links of a block form a spanning tree of its sixteen sites. Each block carries* 49 *active links: the* 17 *active links inside it, with $o_\mu = 0$, and the* 32 *links with $o_\mu = 1$, which join $B(y)$ to $B(y + e_\mu)$.*

(ii) *If dh vanishes on the tree links, h is constant on every block, and $V_{y,\mu}(\varphi + dh) = V_{y,\mu}(\varphi) + h_{B(y)} - h_{B(y+e_\mu)}$.*

(iii) *With $S^\mu_y = \sum_{x\in B(y),\, o_\mu=0} \varphi_{(x,\mu)}$,*

$$V_{y,\mu}(\varphi) = \frac{1}{8} \sum_{x\in B(y),\, o_\mu=1} \varphi_{(x,\mu)} + \frac{1}{16}\big(S^\mu_y + S^\mu_{y+e_\mu}\big).$$

Proof. (i) A tree link of direction μ has $o_\mu = 0$ and stays in its block; there are 2^μ of them for each μ, fifteen in all. From any site, the tree links of direction 3 lead to $o_3 = 0$, then those of direction 2 to $o_2 = 0$, and so on to the corner, so the tree links connect the sixteen sites, and fifteen edges connecting sixteen vertices form a tree. Of the 64 links with base in the block, the 32 with $o_\mu = 0$ stay inside it. (ii) h is constant along the edges of a spanning tree. The path of length two from $x \in B(y)$ ends at $x + 2e_\mu \in B(y + e_\mu)$, and the sum of dh along it is $h(x) - h(x + 2e_\mu)$. (iii) The path from x consists of (x,μ) and $(x + e_\mu, \mu)$; a link with $o_\mu = 1$ is the second link of one path and the first of another, and the links of $S^\mu_{y+e_\mu}$ are the second links of the paths that start with $o_\mu = 1$. □

By (ii) the residual gauge freedom is the coarse gauge group, which acts on the block averages as on the links of the coarse lattice. Let W be a set of blocks and let Φ_W be the space of finitely supported real φ on the active links with base in $\bigcup W$, with $V_{y,\mu}(\varphi) = 0$ for every $y \in W$ and every μ. Write $Q(\varphi) = \sum_p (d\varphi)_p^2$ over all plaquettes, φ being zero off its support.

Theorem 12.4. *For every set of blocks W and every $\varphi \in \Phi_W$,*

$$\sum_e \varphi_e^2 \le 52\, Q(\varphi).$$

Proof. Inside a block. Let $e = (x,\mu)$ be an active link with $o_\mu = 0$, and let $\nu_1 > \cdots > \nu_k$ be the directions above μ in which o equals one; $1 \le k \le 3$. Put $x_0 = x$ and $x_i = x_{i-1} - e_{\nu_i}$. The plaquette p_i at x_i in the plane $\{\mu, \nu_i\}$ has the links (x_i,μ), (x_{i-1},μ) and the tree links (x_i,ν_i), $(x_i + e_\mu, \nu_i)$, whose offsets vanish in ν_i and above it; and (x_k,μ) is a tree link. Hence $\varphi_e = \pm(d\varphi)_{p_1} \pm \cdots \pm (d\varphi)_{p_k}$ and $\varphi_e^2 \le k \sum_i (d\varphi)^2_{p_i}$. The plaquette at x' in the plane $\{\mu,\nu\}$ is used only by the links $(x' + e_\nu + \sum_{\rho\in R} e_\rho, \mu)$ with $R \subset \{\nu+1, \dots, 3\}$, each with

$k = c + 1 + |R|$, where c is the number of ones of the offset of x' strictly between μ and ν. Its total weight $2^{3-\nu}(c+1) + (3-\nu)2^{2-\nu}$ is at most 8, the value at $\mu = 0$, $\nu = 1$. So the sum I of φ_e^2 over the active links inside the blocks is at most $8Q_{\text{in}}$, where Q_{in} is the sum over the plaquettes inside a block.

Between blocks. Fix $y \in W$ and μ, and label the eight links (x,μ), $x \in B(y)$, $o_\mu = 1$, by the transverse offset $a \in \{0,1\}^3$. If a and a' differ in the coordinate ν, with $o_\nu = 0$ at a, the plaquette at x in the plane $\{\mu,\nu\}$ contains these two links, (x,ν) in $B(y)$ and $(x+e_\mu,\nu)$ in $B(y+e_\mu)$, the last two inside their blocks. So $\varphi_a - \varphi_{a'} = \pm(d\varphi)_p \pm \psi \pm \psi'$ with ψ, ψ' inside blocks, and these twelve face plaquettes correspond to the twelve edges of the cube $\{0,1\}^3$. By Lemma 12.3(iii) and $V_{y,\mu}(\varphi) = 0$, the mean of the φ_a is $m = -(S^\mu_y + S^\mu_{y+e_\mu})/16$, so $8m^2 \le \frac{1}{2}$ times the sum of the squares of the sixteen links of S^μ_y and $S^\mu_{y+e_\mu}$. The Laplacian of the graph $\{0,1\}^3$ has smallest nonzero eigenvalue 2, so

$$\begin{aligned}\sum_a \varphi_a^2 &= 8m^2 + \sum_a (\varphi_a - m)^2 \\ &\le 8m^2 + \frac{3}{2}\sum_{\text{edges}} \big((d\varphi)_p^2 + \psi^2 + \psi'^2\big).\end{aligned}$$

Summed over $y \in W$ and μ, a link (z,ν) inside a block occurs as ψ or ψ' once for each $\mu \ne \nu$ (as ψ if $o_\mu = 1$, as ψ' of the face of the block below if $o_\mu = 0$) and in two of the sums S^ν, so it receives weight at most $3 \cdot \frac{3}{2} + 2 \cdot \frac{1}{2} = \frac{11}{2}$. The face plaquettes are distinct and lie in no block, so the links between blocks contribute at most $\frac{3}{2}Q_{\text{face}} + \frac{11}{2}I$, and

$$\sum_e \varphi_e^2 \le \frac{13}{2}I + \frac{3}{2}Q_{\text{face}} \le 52\,Q_{\text{in}} + \frac{3}{2}Q_{\text{face}} \le 52\,Q(\varphi). \qquad \square$$

The constant 52 is not sharp. A certificate of one of us (T. Romanov) lowers it to within one part in a thousand of the bulk value.

Proposition 12.5. *For every set of blocks W and every $\varphi \in \Phi_W$,*

$$\sum_e \varphi_e^2 \le \frac{5000}{537}\,Q(\varphi) = 9.31098\ldots\,Q(\varphi).$$

The same holds on every periodic lattice of even side.

Proof. Let $c = \frac{537}{5000}$, and let $S(y) = B(y) \cup B(y+e_0) \cup \cdots \cup B(y+e_3)$, whose active links number $5 \cdot 49 = 245$. The certificate consists of 196 rational numbers $l_{\mu,b}$, $\mu = 0,\ldots,3$, indexed by the 49 active links b of a block, and a rational symmetric matrix G of size 245 on the active links of $S(0)$. With $L_{y,\mu}(\varphi) = \sum_b l_{\mu,b}\varphi_{(y,b)}$, where (y,b) is the link b of $B(y)$, and G_y the translate of G to $S(y)$, these satisfy the identity of quadratic forms in finitely supported φ

$$Q(\varphi) - c\sum_e \varphi_e^2 - \sum_{y,\mu} V_{y,\mu}(\varphi)L_{y,\mu}(\varphi) = \sum_y G_y(\varphi|_{S(y)}), \tag{12.1}$$

and G is positive definite. Both sides of (12.1) are invariant under translations by $2\mathbb{Z}^4$, so the identity amounts to finitely many equalities of coefficients, one for each pair of active links of $B(0)$ and of a block near it. Now take $\varphi \in \Phi_W$. If $y \in W$ then $V_{y,\mu}(\varphi) = 0$; if $y \notin W$ then φ vanishes on the links with base in $B(y)$, so $L_{y,\mu}(\varphi) = 0$. Every term of the middle sum vanishes, and $Q(\varphi) - c\sum_e \varphi_e^2 \ge 0$. On a periodic lattice of even side, both sides of (12.1) are the periodisations of the same forms, so the identity and the argument hold there. $\square$

The certificate was found by semidefinite programming and rounded to rationals: the $l_{\mu,b}$ have denominators dividing 2^{24} and the entries of G the denominator $2^{30} \cdot 5^5$. The coefficient identity of (12.1) and the positivity of all 245 leading principal minors of the integer matrix $2^{30}5^5 G$ are checked in exact arithmetic. The check is done by the program

of T. Romanov and independently by our program in the archive, which rebuilds the four forms from Definition 12.2 alone. With W the set of all blocks the best constant is the inverse of the Bloch value 0.1074878 quoted below, that is 9.3034, so $\frac{5000}{537}$ is within 0.09 per cent of it; the semidefinite optimum for this window and this class of multipliers is $c^{*} = 0.107488$. These floating-point values are not used. The theorems below are stated with 52, which is proved by hand. Where they use only Theorem 12.4, they hold with 52 replaced by $\frac{5000}{537}$.

With W all blocks the bound extends to ℓ^2, so the block-constrained quadratic form on $\mathbb{Z}^4$ has a spectral gap at least $\frac{1}{52}$; it removes in particular the four flat connections that are zero modes of the unconstrained form. In the Bloch decomposition the exact gap is $0.10748776948684\ldots$, attained at the momentum $(0,\pi,\pi,\pi)$, a floating-point value computed in [56] and reproduced in the archive; the theorem does not use it. The cube conditions now take the following form. Let n be even, V the active links with base in $[0,n)^4$, W its blocks, and for an exterior η and values $v = (v_{y,\mu})_{y\in W}$ let $\gamma^{\mathrm{c}}_{V}(\cdot \mid \eta, v)$ be the law on $\mathbb{R}^{V}$ with density proportional to $\exp(-\frac{b}{2}\sum_{p\cap V\neq\emptyset}(d\varphi)_p^2)$ on the affine set $\{V_{y,\mu}(\varphi) = v_{y,\mu}\}$, with $\varphi = \eta$ off V. Define $C^{\mathrm{c}}_{V,j}$, K^{c}_{c} and $D_{\mathrm{c}}(n,\alpha)$ as $C_{V,j}$, K_c and $D(n,\alpha)$, with v held fixed and the 49 classes of active links modulo two.

Lemma 12.6. *Let $H \geq 0$ be symmetric on $\mathbb{R}^m$ with $\|H\| \leq h$, let $B : \mathbb{R}^m \to \mathbb{R}^k$ with $BB^{\top} \geq \beta > 0$ and $\langle x, Hx\rangle \geq a|x|^2$ on $\ker B$, and $s > 0$. Then $K = \left(\begin{smallmatrix} H & sB^{\top} \\ sB & 0 \end{smallmatrix}\right)$ is invertible, and the solution of $K(x,y) = (f,g)$ has $|x| \leq (|f| + h|x_1|)/a + |x_1|$ with $|x_1| \leq |g|/(s\sqrt{\beta})$, and $|y| \leq (|f| + h|x|)/(s\sqrt{\beta})$.*

Proof. Put $x_1 = B^{\top}(BB^{\top})^{-1}g/s$, so $sBx_1 = g$ and $|x_1| \leq |g|/(s\sqrt{\beta})$. On $\ker B$ the compression of H is at least a, so there is a unique $x_0 \in \ker B$ with $Hx_0 - f + Hx_1 \perp \ker B$, and $a|x_0| \leq |f| + h|x_1|$. Then $f - H(x_0 + x_1)$ lies in $(\ker B)^{\perp} = \operatorname{ran} B^{\top}$, and $y = (BB^{\top})^{-1}B(f - Hx)/s$ solves $sB^{\top}y = f - Hx$. For $f = g = 0$ the construction gives zero, so K is injective. □

Lemma 12.7 (Combes–Thomas [28])**.** *Let K be an invertible symmetric matrix indexed by a finite set X, with $\|K^{-1}\| \leq M_0$ and $\sum_b |K_{ab}| \leq S$ for every a, and let $d : X \to \mathbb{R}$ satisfy $|d(a) - d(b)| \leq 1$ whenever $K_{ab} \neq 0$. If $(e^{\lambda} - 1)SM_0 \leq \frac{1}{2}$, then $\|e^{\lambda d}K^{-1}e^{-\lambda d}\| \leq 2M_0$.*

Proof. $e^{\lambda d}Ke^{-\lambda d} = K + E$ with $|E_{ab}| \leq (e^{\lambda} - 1)|K_{ab}|$, so $\|E\| \leq (e^{\lambda} - 1)S$ by the Schur test, $\|K^{-1}E\| \leq \frac{1}{2}$, and $(K+E)^{-1} = (1 + K^{-1}E)^{-1}K^{-1}$ has norm at most $2M_0$. □

Theorem 12.8. *For every $b > 0$, every even n and every exterior link j:*

(i) *$\gamma^{\mathrm{c}}_{V}(\cdot \mid \eta, v)$ is a Gaussian law whose covariance depends on neither η nor v and whose mean is affine in (η, v), and $C^{\mathrm{c}}_{V,j} = \|u_j\|_1$, where $u_j \in \mathbb{R}^{V}$ is the change of the mean when η_j increases by one; u_j does not depend on b;*

(ii) *$\sum_{e\in V} e^{2\lambda_0 d(e)}u_j(e)^2 \leq 3500^2$ and $\|u_j\|_1 \leq 10^{14}$, where $\lambda_0 = 2.4\cdot 10^{-5}$ and $d(e)$ is the ℓ^{∞} distance between the midpoints of e and j;*

(iii) *$D_{\mathrm{c}}(n,\alpha)$ holds for every $\alpha \in (0,1)$ and every even $n \geq 2\cdot 10^{15}/\alpha$, at every $b > 0$.*

Proof. (i) Let H be the matrix of $\sum_{p\cap V\neq\emptyset}(d\varphi)_p^2$ on $\mathbb{R}^{V}$ and B that of the averages $V_{y,\mu}$, $y \in W$, restricted to V. By Theorem 12.4 with this W, $H \geq \frac{1}{52}$ on $\ker B$, so the density is that of a non-degenerate Gaussian on the affine set, its covariance is b^{-1} times the inverse of H on $\ker B$, and its mean m minimises the quadratic form on the affine set. The mean is affine in (η, v), and two exteriors that differ at j give translates by u_j, which minimises $Q(u + \delta_j)$ subject to $Bu = g$, g being minus the coefficients of j in the averages. The translation argument of Proposition 11.3 gives $C^{\mathrm{c}}_{V,j} = \|u_j\|_1$.

(ii) The pair (u_j, y), with y the Lagrange multipliers, solves $K(u_j, y) = (f, sg)$ with K as in Lemma 12.6, $s = 96$ and $f = -\sum_{p\ni j}(d\delta_j)_p\, d_p|_V$, where $d_p|_V$ is the row of the plaquette p restricted to V. Each link lies in six plaquettes of four links, so the row sums of $|H|$ are at most 24 and $h \leq 24$; $|f| \leq 6\sqrt{3}$; and $|g| \leq \frac{1}{16}$, since an exterior link enters at most

one average of W, with weight $\frac{1}{16}$. By Lemma 12.3(iii) each average has squared norm at least $8\cdot\frac{1}{64}=\frac{1}{8}$, two averages share links only if they are (y,μ) and $(y\pm e_\mu,\mu)$, and then their inner product is at most $8\cdot\frac{1}{256}=\frac{1}{32}$; so $BB^{\top}\geq\frac{1}{16}$ by Gershgorin's theorem. With $a=\frac{1}{52}$, Lemma 12.6 gives $\|K^{-1}\|\leq M_0=105$. The rows of K have absolute sums at most $S=\max(24+\frac{96}{8},96\cdot 2)=192$, because an average has eight coefficients $\frac{1}{8}$ and sixteen of $\frac{1}{16}$ and a link enters averages with total weight at most $\frac{1}{8}$. Index the rows of B by the centre $2y+\frac{1}{2}(1,1,1,1)+e_\mu$ of the two blocks of the average and put d equal to the ℓ^∞ distance to the midpoint of j. Links of one plaquette, and an average and its links, are within distance one, so Lemma 12.7 applies with $\lambda=\log(1+1/40320)>\lambda_0$, and

$$\sum_e e^{2\lambda d(e)}u_j(e)^2\leq\big(2M_0e^{\lambda}(|f|+s|g|)\big)^2\leq 3500^2,$$

since f and g live at distance at most one from j. At most $4(64k^3+16k)$ links have their base at ℓ^∞ distance $k\geq 1$ from the base of j, and their midpoints are at distance at least $k-1$, so Cauchy–Schwarz gives

$$\|u_j\|_1^2\leq 3500^2\cdot 4e^{2\lambda}\Big(1+\sum_{k\geq 1}(64k^3+16k)e^{-2\lambda k}\Big)\leq(10^{14})^2.$$

(iii) $C^{\mathrm{c}}_{V_x,j}=0$ unless j shares a plaquette with V_x or enters an average of V_x, which forces the base z of j into $x+[-1,n+1]^4$, while $z\notin x+[0,n)^4$ when $j\notin V_x$. For each coordinate at most $\frac{n}{2}+2$ even x_i satisfy the first condition and exactly $\frac{n}{2}$ the second, so

$$K^{\mathrm{c}}_c\leq\Big(\Big(\frac{n}{2}+2\Big)^4-\Big(\frac{n}{2}\Big)^4\Big)10^{14}\leq\frac{17}{n}\,10^{14}N_n$$

for $n\geq 10^4$, which is at most αN_n when $n\geq 2\cdot 10^{15}/\alpha$. □

The constants are crude. The computations of [56] give $C^{\mathrm{c}}_{V,j}\leq 12.48$ at $n=4$, and an exact face expansion $K^{\mathrm{c}}_c/N_n=a_c/n+O(n^{-2})$ with $\max_c a_c=61.18$, so that $D_{\mathrm{c}}(n,0.95)$ first holds at $n=64$; they are floating-point values, reproduced in the archive for $n\leq 4$, and the theorem does not use them. The coefficients are large because the constrained responses obey no maximum principle: an exterior link changes the curvature of up to six plaquettes, and keeping the curvature small moves a dozen links inside the cube with alternating signs. Re-weighting the classes does not help much: if $\sum_{j\in c\setminus V}\sum_{c'}w_{c'}C^{c'}_{V,j}\leq\alpha N_n w_c$ for positive weights w, where $C^{c'}_{V,j}$ is the part of $\|u_j\|_1$ on the links of the class c', then αN_n is at least the spectral radius of the non-negative matrix of these sums, hence at least $\min_c K^{\mathrm{c}}_c$, by the Collatz–Wielandt inequality [56].

The block averages can be integrated out one scale at a time. For a field v on the coarse links put $Q_1(v)=\sum_P(dv)_P^2$ over the plaquettes P of the coarse lattice, and write $V(\varphi)=(V_{y,\mu}(\varphi))_{y,\mu}$. The averages of averages are formed in the same way on the coarse lattice, and $V^{(k)}$, Q_k denote the k-fold averages and the plaquette form of the lattice of spacing 2^k.

Proposition 12.9. *Let φ be a finitely supported field on $\mathbb{Z}^4$ that vanishes on the tree links.*

(i) *For the coarse plaquette P at y in the plane $\{\mu,\nu\}$, $16\,(dV(\varphi))_P$ is the sum over $x\in B(y)$ of the curvatures of the squares of side two spanned by x, $x+2e_\mu$, $x+2e_\nu$, and every plaquette of $\mathbb{Z}^4$ lies in exactly four of these squares. Hence $Q_1(V(\varphi))\leq Q(\varphi)$, and $Q_k(V^{(k)}(\varphi))\leq Q(\varphi)$ for every $k\geq 1$.*

(ii) *For every finitely supported v on the coarse links, the field $\varphi^*(v)$ equal to $v_{y,\mu}$ on the eight links (x,μ), $x\in B(y)$, $o_\mu=1$, and to zero on the other links, vanishes on the tree links, has $V(\varphi^*(v))=v$ and $Q(\varphi^*(v))=4\,Q_1(v)$.*

(iii) *The effective form $S_1(v)=\inf\{Q(\varphi):V(\varphi)=v\}$ satisfies $Q_1\leq S_1\leq 4\,Q_1$, and $S_k(w)=\inf\{Q(\varphi):V^{(k)}(\varphi)=w\}\geq Q_k(w)$ for every k, the infima over finitely supported fields that vanish on the tree links.*

(iv) *For every k, every set W of blocks of the lattice of spacing 2^k and every w in the space Φ_W of that lattice, $\sum_e w_e^2 \le 52\, S_k(w)$.*

Proof. (i) The path of length two from $x \in B(y)$ in the direction μ ends at $x+2e_\mu \in B(y+e_\mu)$, at the same offset, and so on around the square; so the four averages of $(dV)_P$ pair the sixteen paths into sixteen closed squares, and the sum of φ around a square is the sum of the curvatures of its four plaquettes. A plaquette at z in the plane $\{\mu,\nu\}$ lies in the square from x exactly when $z-x \in \{0,1\}e_\mu + \{0,1\}e_\nu$, which gives four points x, each in one block. By Cauchy–Schwarz, $(dV)_P^2 \le \frac{1}{16}\sum_x \ell_x^2 \le \frac{1}{4}\sum_x \sum_{p\subset\square_x} (d\varphi)_p^2$, where ℓ_x is the curvature of the square from x; summing over P gives $Q_1(V(\varphi)) \le Q(\varphi)$. The argument uses neither the tree links nor the spacing, so it applies on every lattice, and induction gives the bound for $V^{(k)}$.

(ii) The tree links have $o_\mu = 0$. By Lemma 12.3(iii) the average of φ^* is $\frac{1}{8}\cdot 8 v_{y,\mu}$. A plaquette inside a block has only links with $o_\mu = 0$ in its direction μ, and so vanishes. A plaquette at x in the plane $\{\mu,\nu\}$ with $o_\mu = 1$, $o_\nu = 0$ has the two links (x,μ), $(x+e_\nu,\mu)$ of the same block, both equal to $v_{y,\mu}$, and two links with offset zero in the direction ν; its curvature vanishes, and so does that of the plaquettes with $o_\mu = 0$, $o_\nu = 1$. If $o_\mu = o_\nu = 1$, its four links carry $v_{y,\mu}$, $v_{y+e_\mu,\nu}$, $v_{y+e_\nu,\mu}$, $v_{y,\nu}$, so its curvature is $(dv)_P$, and there are four such plaquettes for each P.

(iii) The lower bound is (i) and the upper bound is (ii); for S_k apply (i) to the k-fold averages.

(iv) Theorem 12.4 holds on every lattice, so $\sum_e w_e^2 \le 52\, Q_k(w) \le 52\, S_k(w)$ by (iii). □

In the Bloch decomposition the eigenvalues of S_1 relative to Q_1 fill $[1,4]$: they tend to one as the momentum tends to zero and equal four at $(\pi,\pi,0,0)$, a floating-point computation reproduced in the archive. So at this order the averages carry the coarse quadratic form with no loss at long wavelengths; the coupling does not run here, and its running is the order-g^2 effect of Remark 12.19(c). Part (iv) is a gap for the averages at every scale, with the constant of Theorem 12.4; it does not give the localisation of Theorem 12.8(ii) for S_k, since S_k is not a nearest-neighbour form.

The effective forms are equivalent to the coarse plaquette forms uniformly in the scale. Write $L = 2^k$ and $B_k(y) = Ly + \{0,\dots,L-1\}^4$.

Lemma 12.10. (i) *For every $k \ge 1$, every coarse link (y,μ) and every field φ,*

$$V^{(k)}_{y,\mu}(\varphi) = \frac{1}{L^4} \sum_{x\in B_k(y)} \sum_{t=0}^{L-1} \varphi_{(x+te_\mu,\mu)}.$$

(ii) *For a function h on sites, $V^{(k)}_{y,\mu}(dh) = \bar h(y) - \bar h(y+e_\mu)$, where $\bar h(y)$ is the mean of h over $B_k(y)$.*

(iii) *$S_k(w)$ equals the infimum of $Q(\varphi)$ over all finitely supported fields with $V^{(k)}(\varphi) = w$, whether or not they vanish on the tree links.*

Proof. (i) For $k = 1$ this is the definition. If it holds for k, then $V^{(k)}_{y',\mu} + V^{(k)}_{y'+e_\mu,\mu}$ is L^{-4} times the sum over $x \in B_k(y')$ of the sums of φ along the straight paths of length $2L$ from x, because $B_k(y'+e_\mu) = B_k(y') + Le_\mu$. The blocks $B_k(y')$, $y' \in 2y + \{0,1\}^4$, tile $B_{k+1}(y)$, and averaging over y' with weight $\frac{1}{16}$ gives (i) for $k+1$. (ii) The sum of dh along the path of length L from x is $h(x) - h(x+Le_\mu)$; average over $x \in B_k(y)$. (iii) Let φ be finitely supported. On each block $B(y')$ of side two the tree links form a spanning tree (Lemma 12.3(i)), so there is h_0 on $B(y')$ with $(dh_0)_e = -\varphi_e$ on its tree links, unique up to a constant, which we take to be zero where φ vanishes on the tree links of $B(y')$. Each $B_k(y)$ is a union of blocks of side two, since $k \ge 1$; adding a constant to h_0 on one of them makes the mean over $B_k(y)$ vanish, and this changes h_0 on finitely many blocks. The resulting h is finitely supported, $\varphi + dh$ vanishes on the tree links, $Q(\varphi+dh) = Q(\varphi)$, and $V^{(k)}(dh) = 0$ by (ii). □




FIGURE 13. The extension of Theorem 12.12 along one direction. (a) A block of side $L = 4$: the nodes are the block centres c_z, the hat functions λ_z interpolate between them, and the straight path of length L from a site x meets the segments of three centres; averaged over the block, their weights are $\frac{1}{8}$, $\frac{3}{4}$, $\frac{1}{8}$ at every scale. (b) The symbol of this filter stays at least $\frac{1}{2}$, while the filter obtained with the coarse sites as nodes has symbol of modulus $1/L$ at $\theta = \pi$.

The extension uses the lowest-order Whitney forms of the coarse cubical complex whose vertices are the centres

$$c_z = Lz + \tfrac{L-1}{2}(1,1,1,1), \qquad z \in \mathbb{Z}^4,$$

of the blocks $B_k(z)$, rather than the coarse sites. Let $\lambda(t) = \max(0, 1-|t|)$, let χ_z^μ be the indicator of $\{\xi \in \mathbb{R}^4 : c_{z,\mu} \le \xi_\mu < c_{z,\mu} + L\}$, and $\lambda_z^\nu(\xi) = \lambda((\xi_\nu - c_{z,\nu})/L)$. For a square-summable field u on the coarse links put

$$\omega_\mu(\xi) = \frac{1}{L}\sum_z u_{(z,\mu)}\,\chi_z^\mu(\xi)\prod_{\nu\neq\mu}\lambda_z^\nu(\xi),$$

$$(\Pi u)_{(x,\mu)} = \int_0^1 \omega_\mu(x + te_\mu)\,dt.$$

Figure 13 shows the construction along one direction.

Lemma 12.11. *Let $k \ge 1$, u a square-summable field on the coarse links, and let T act on each component of u by $T = \tau_1\tau_2\tau_3\tau_4$, where $(\tau_\nu f)(z) = \frac{3}{4}f(z) + \frac{1}{8}(f(z-e_\nu) + f(z+e_\nu))$.*

(i) *For the plaquette p at x in the plane $\{\mu,\nu\}$,*

$$(d\Pi u)_p = \int_0^1\int_0^1 \Omega_{\mu\nu}(x + se_\mu + te_\nu)\,ds\,dt,$$

$$\Omega_{\mu\nu} = \frac{1}{L^2}\sum_z (du)_{z,\mu\nu}\,\chi_z^\mu\chi_z^\nu\prod_{\rho\neq\mu,\nu}\lambda_z^\rho,$$

where $(du)_{z,\mu\nu}$ is the curvature of u at the coarse plaquette at z in the plane $\{\mu,\nu\}$. In particular $\Pi(db)$ is a gradient.

(ii) $Q(\Pi u) \le Q_k(u)$.

(iii) $V^{(k)}(\Pi u) = Tu$.

Proof. (i) The function ω_μ is continuous and piecewise linear in every coordinate ξ_ν, $\nu \neq \mu$, and ω_ν in ξ_μ. Hence

$$(d\Pi u)_p = \int_0^1 \big(\omega_\mu(x + te_\mu) - \omega_\mu(x + e_\nu + te_\mu)\big)dt + \int_0^1 \big(\omega_\nu(x + e_\mu + te_\nu) - \omega_\nu(x + te_\nu)\big)dt$$

is the integral over the unit square of $\partial_\mu\omega_\nu - \partial_\nu\omega_\mu$, which exists almost everywhere. Since $\partial_\mu\lambda^\mu_z = \frac{1}{L}(\chi^\mu_{z-e_\mu} - \chi^\mu_z)$ and χ^ν_z and λ^ρ_z, $\rho \neq \mu$, do not change when z is shifted by e_μ, a shift of the summation index gives

$$\partial_\mu\omega_\nu = \frac{1}{L^2}\sum_z \big(u_{(z+e_\mu,\nu)} - u_{(z,\nu)}\big)\chi^\mu_z\chi^\nu_z \prod_{\rho\neq\mu,\nu}\lambda^\rho_z,$$

and the same with μ, ν exchanged. The difference is $\Omega_{\mu\nu}$, with $(du)_{z,\mu\nu} = u_{(z,\mu)} + u_{(z+e_\mu,\nu)} - u_{(z+e_\nu,\mu)} - u_{(z,\nu)}$, the sign convention of the fine plaquettes. If $u = db$ then $\Omega = 0$, every curvature of Πu vanishes, and Πu is a gradient.

(ii) By Cauchy–Schwarz, $(d\Pi u)^2_p$ is at most the integral of $\Omega^2_{\mu\nu}$ over the unit square of p. At a point ξ exactly one pair (z_μ, z_ν) has $\chi^\mu_z\chi^\nu_z = 1$, and the hat functions form a partition of unity, $\sum_{z_\rho,z_\sigma}\lambda^\rho_z\lambda^\sigma_z = 1$ for $\{\rho,\sigma\}$ the other two directions; so $\Omega^2_{\mu\nu} \le L^{-4}\sum_{z_\rho,z_\sigma}\lambda^\rho_z\lambda^\sigma_z(du)^2_{z,\mu\nu}$. Summing over the plaquettes of the plane $\{\mu,\nu\}$ integrates over ξ_μ, ξ_ν, which gives the factor L^2 for each pair (z_μ, z_ν), and sums over the integers x_ρ, x_σ. Since $c_{z,\rho} - x_\rho$ lies in $\mathbb{Z} + \frac12$, $\sum_{x_\rho\in\mathbb{Z}}\lambda^\rho_z = 2\sum_{j=0}^{L-1}(1 - \frac{j+1/2}{L}) = L$, and the same for σ. So the plaquettes of the plane $\{\mu,\nu\}$ contribute at most $L^{-4}\cdot L^2\cdot L^2\sum_z(du)^2_{z,\mu\nu}$.

(iii) By Lemma 12.10(i), $V^{(k)}_{y,\mu}(\Pi u)$ is the mean over $x = Ly + s$, $s \in \{0,\dots,L-1\}^4$, of $\int_0^L \omega_\mu(x + te_\mu)\,dt$. The term of $u_{(z,\mu)}$ in it is a product of a factor of the coordinate μ and one of each $\nu \neq \mu$, and the mean over s is the product of the means over the single coordinates. Put $\delta = s_\mu - \frac{L-1}{2}$. The interval $[x_\mu, x_\mu + L)$ meets $[c_{z,\mu}, c_{z,\mu} + L)$ in length $L - |\delta|$ if $z_\mu = y_\mu$, in $\max(0, -\delta)$ if $z_\mu = y_\mu - 1$, in $\max(0,\delta)$ if $z_\mu = y_\mu + 1$, and not otherwise. As L is even, $\sum_s|\delta| = 2\sum_{j=0}^{L/2-1}(j + \frac12) = \frac{L^2}{4}$, so the means of these lengths divided by L are $\frac34$, $\frac18$, $\frac18$. For $\nu \neq \mu$ put $\delta = s_\nu - \frac{L-1}{2}$; then $\lambda^\nu_z(x)$ is $1 - |\delta|/L$ if $z_\nu = y_\nu$, $\max(0, \mp\delta)/L$ if $z_\nu = y_\nu \mp 1$, and zero otherwise, with the same means. The product of the four means is the coefficient of $u_{(z,\mu)}$ in $(Tu)_{(y,\mu)}$. □

Theorem 12.12. *For every $k \ge 1$ and every finitely supported field w on the links of the lattice of spacing 2^k,*

$$Q_k(w) \le S_k(w) \le 256\,Q_k(w).$$

Proof. The lower bound is Proposition 12.9(iii). The operator T of Lemma 12.11 is a convolution, the same on every component, so it commutes with d; its symbol is $\prod_\nu(\frac34 + \frac14\cos\theta_\nu) \ge \frac{1}{16}$, so it is invertible on square-summable fields with $\|T^{-1}\| \le 16$. Let $u = T^{-1}w$. By Lemma 12.11, $V^{(k)}(\Pi u) = w$ and

$$Q(\Pi u) \le Q_k(u) = \|T^{-1}dw\|^2 \le 256\,Q_k(w).$$

It remains to replace Πu by a finitely supported field. Let u_R be u on the coarse links with base in $[-R, R]^4$ and zero elsewhere, and $e_R = w - Tu_R = T(u - u_R)$, which is finitely supported and tends to zero. Let ψ_R equal $e_R(y,\mu)$ on the links (x,μ) with $x \in B_k(y)$ and $x_\mu \equiv L - 1$ modulo L, and zero elsewhere. Each straight path of length L from a site of $B_k(y)$ in the direction μ contains exactly one such link of $B_k(y)$, so $V^{(k)}(\psi_R) = e_R$ by Lemma 12.10(i), and $Q(\psi_R) \le 24\|\psi_R\|^2 = 24L^3\|e_R\|^2$, since a link lies in six plaquettes of four links. The field $\Pi u_R + \psi_R$ is finitely supported, its k-fold averages are w, and $Q(\Pi u_R + \psi_R)^{1/2} \le Q_k(u_R)^{1/2} + (24L^3)^{1/2}\|e_R\|$, which tends to $Q_k(u)^{1/2}$. Lemma 12.10(iii) gives $S_k(w) \le Q_k(u) \le 256\,Q_k(w)$. □

So at every scale the averages carry the coarse plaquette form up to a factor at most 256, independent of k. The construction inverts the averages by a fixed convolution: the filter $(\frac{1}{8},\frac{3}{4},\frac{1}{8})$ does not depend on the scale because the nodes are the block centres, and its symbol is at least $\frac{1}{2}$. With the coarse sites as nodes the corresponding filter is $\frac{L+1}{2L}\delta_0+\frac{L-1}{2L}\delta_1$, whose symbol has modulus $1/L$ at $\theta=\pi$ (Figure 13(b)). In floating point, on a grid of 12^4 momenta, the largest ratio of S_k to Q_k on curvatures is 3.9, 7.8 and 10.2 for $k=1,2,3$, and the kernel of S_k decreases by the factors 0.17, 0.23 and 0.25 per coarse spacing; these values are reproduced in the archive. The theorem gives no decay. The decay comes from an exact formula for S_k in Fourier variables: in a Feynman gauge for the fine field the effective form is diagonal in the six planes, and the pole of the fine propagator at zero momentum cancels between numerator and denominator.

For $t\in\mathbb{C}$ put $D_L(t)=\frac{1}{L}\sum_{j=0}^{L-1}e^{ijt}$, and for $\vartheta\in\mathbb{C}^4$ put $\Psi(\vartheta)=\prod_\nu D_L(\vartheta_\nu)$, $E_\mu(\vartheta)=e^{i\vartheta_\mu}-1$ and $\varepsilon(\vartheta)=\sum_\nu(2-2\cos\vartheta_\nu)$. For an entire function f let $f^\sharp(\vartheta)=\overline{f(\bar\vartheta)}$, so that $ff^\sharp=|f|^2$ on real arguments. For $\theta\in\mathbb{C}^4$ and $n\in\{0,\dots,L-1\}^4$ let $\theta^{(n)}=(\theta+2\pi n)/L$; every function below is 2π-periodic in each coordinate of $\theta^{(n)}$, and we take the real parts of these coordinates in $[-\pi,\pi]$. Write $\Psi_n=(\Psi\Psi^\sharp)(\theta^{(n)})$, $D_{n,\sigma}=(D_LD_L^\sharp)(\theta_\sigma^{(n)})$, and put

$$A_\sigma=\Psi_0D_{0,\sigma},\qquad \hat\varepsilon=L^2\varepsilon(\theta^{(0)}),\qquad \tilde a_\sigma=A_\sigma+\hat\varepsilon r_\sigma,$$
$$r_\sigma=\sum_{n\neq0}\frac{\Psi_nD_{n,\sigma}}{L^2\varepsilon(\theta^{(n)})},$$

$$\begin{aligned}Y&=\Psi_0^3\prod_\tau D_{0,\tau}\\&\quad+\sum_\sigma E_\sigma E_\sigma^\sharp\sum_{\emptyset\neq J\subset\{1,2,3,4\}\setminus\{\sigma\}}\hat\varepsilon^{|J|-1}\prod_{\tau\in J}r_\tau\prod_{\tau\notin J\cup\{\sigma\}}A_\tau,\end{aligned}\tag{12.2}$$

all functions of θ, and for a plane $P=\{\lambda,\rho\}$

$$K_P=\frac{\prod_{\sigma\notin P}\tilde a_\sigma}{Y}.$$

For $L=1$ these are $\tilde a_\sigma=Y=K_P=1$. For a coarse field w let $\hat w(\theta)=\sum_y e^{-i\theta\cdot y}w_y$, and let $F_P=(dw)_P$ be its curvature in the plane P, a function of the coarse sites.

Lemma 12.13. *Let $\gamma_1,\dots,\gamma_4>0$, $\Gamma=\operatorname{diag}(\gamma_\mu)$ and $0\neq E\in\mathbb{C}^4$, and for $P=\{\lambda,\rho\}$, $\lambda<\rho$, let c_P be the row vector with $c_Pv=E_\lambda v_\rho-E_\rho v_\lambda$. Then*

$$\Gamma-\frac{\Gamma EE^*\Gamma}{E^*\Gamma E}=\frac{1}{E^*\Gamma E}\sum_P\gamma_\lambda\gamma_\rho\,c_P^*c_P.$$

Proof. Both sides are Hermitian. The diagonal entry (μ,μ) of the right side is $\sum_{\lambda\neq\mu}\gamma_\mu\gamma_\lambda|E_\lambda|^2/E^*\Gamma E$, which is $\gamma_\mu-\gamma_\mu^2|E_\mu|^2/E^*\Gamma E$; the entry (μ,ν), $\mu\neq\nu$, comes only from $P=\{\mu,\nu\}$, where c_P has $-E_\nu$ in place μ and E_μ in place ν, and equals $-\gamma_\mu\gamma_\nu\bar E_\mu E_\nu/E^*\Gamma E$. □

Theorem 12.14. *Let $k\geq1$ and $L=2^k$.*

(i) *The functions K_P are real-analytic, even, 2π-periodic and positive on $\mathbb{R}^4$, $K_P(0)=1$, and for every finitely supported field w on the coarse links*

$$S_k(w)=\sum_P\int_{[-\pi,\pi]^4}K_P(\theta)\,|\hat F_P(\theta)|^2\,\frac{d\theta}{(2\pi)^4}.$$

(ii) *There are $\eta>0$ and $C<\infty$, independent of k, such that every K_P extends analytically to $\{|\operatorname{Im}\theta_\nu|\leq\eta\}$ and is bounded there by C.*

(iii) *Hence $S_k(w)=\sum_P\sum_{y,y'}\mathcal{K}_P(y-y')F_P(y)F_P(y')$ with real kernels $\mathcal{K}_P(y)=(2\pi)^{-4}\int K_P(\theta)e^{i\theta\cdot y}\,d\theta$ that satisfy $|\mathcal{K}_P(y)|\leq Ce^{-\eta|y|_1}$ for every k.*

Proof. Step 1: the form in Fourier variables. Let $c_\mu(q) = L\Psi(q)D_L(q_\mu)$. Lemma 12.10(i) and the Poisson summation formula give, for every square-summable field φ,

$$\widehat{V^{(k)}(\varphi)}_\mu(\theta) = L^{-4}\sum_n c_\mu(\theta^{(n)})\,\hat\varphi_\mu(\theta^{(n)}),$$

with $\hat\varphi(q) = \sum_x e^{-iq\cdot x}\varphi_x$. Write $x_n = \hat\varphi(\theta^{(n)}) \in \mathbb{C}^4$ and $e_n = (e^{i\theta_\mu^{(n)}} - 1)_\mu$, so that $|e_n|^2 = \varepsilon(\theta^{(n)}) =: \varepsilon_n$. By the Lagrange identity the plaquette form at the momentum $\theta^{(n)}$ is $\varepsilon_n|x_n|^2 - |e_n^* x_n|^2$, and the Plancherel formula gives

$$Q(\varphi) = \int_{[-\pi,\pi]^4} L^{-4}\sum_n \bigl(\varepsilon_n|x_n|^2 - |e_n^* x_n|^2\bigr)\frac{d\theta}{(2\pi)^4}.$$

Since $LD_L(t)(e^{it}-1) = e^{iLt} - 1$ and $e^{iL\theta_\mu^{(n)}} = e^{i\theta_\mu}$, the averages map the gauge direction at every alias to the coarse gauge direction: $c_\mu(\theta^{(n)})(e_n)_\mu = \Psi(\theta^{(n)})E_\mu(\theta)$. For $|t| \le \pi/L$ we have $|D_L(t)| = |\sin(Lt/2)|/(L|\sin(t/2)|) \ge 2/\pi$, so $\Psi(\theta^{(0)}) \ne 0$ for real θ.

Step 2: the minimum at one momentum. Fix real $\theta \notin 2\pi\mathbb{Z}^4$; then $E = E(\theta) \ne 0$ and every $\varepsilon_n > 0$. Let $z \in \mathbb{C}^4$. Minimising $L^{-4}\sum_n \varepsilon_n|y_n|^2$ under $L^{-4}\sum_n c_\mu(\theta^{(n)})(y_n)_\mu = z_\mu$ separates in μ, and the Cauchy–Schwarz inequality gives the minimum $\sum_\mu |z_\mu|^2/a_\mu$, attained at $(y_n)_\mu = z_\mu\overline{c_\mu(\theta^{(n)})}/(\varepsilon_n a_\mu)$, where

$$a_\mu = L^{-4}\sum_n \frac{|c_\mu(\theta^{(n)})|^2}{\varepsilon_n} = \frac{\tilde a_\mu}{\hat\varepsilon}.$$

If x satisfies the constraint with right side $\hat w(\theta)$, write $x_n = x_n^T + \beta_n e_n$ with $e_n^* x_n^T = 0$. Then $L^{-4}\sum_n(\varepsilon_n|x_n|^2 - |e_n^*x_n|^2) = L^{-4}\sum_n \varepsilon_n|x_n^T|^2$, and by Step 1 the vectors x_n^T satisfy the constraint with right side $\hat w(\theta) + tE$ for some $t \in \mathbb{C}$. Hence the plaquette form at θ is at least

$$\min_{t\in\mathbb{C}}\sum_\mu \frac{|\hat w_\mu + tE_\mu|^2}{a_\mu} = \hat w^*\Bigl(\Gamma - \frac{\Gamma EE^*\Gamma}{E^*\Gamma E}\Bigr)\hat w$$

with $\Gamma = \operatorname{diag}(a_\mu^{-1})$. Conversely, take t optimal, y optimal for $z = \hat w + tE$, and replace y_0 by $y_0 - \beta e_0$ with $\beta = L^4 t/\Psi(\theta^{(0)})$: the constraint becomes $\hat w(\theta)$ and the plaquette form does not increase, since e_0 is a gauge direction. By Lemma 12.13 the minimum is

$$\sum_P \frac{\gamma_\lambda\gamma_\rho}{E^*\Gamma E}|c_P\hat w|^2, \qquad \frac{\gamma_\lambda\gamma_\rho}{E^*\Gamma E} = \frac{\prod_{\sigma\notin P} a_\sigma}{\sum_\sigma |E_\sigma|^2\prod_{\tau\ne\sigma} a_\tau}.$$

We multiply numerator and denominator by $\hat\varepsilon^3$ and expand $\tilde a_\tau = A_\tau + \hat\varepsilon r_\tau$. The terms without r add up to $\Psi_0^3\prod_\tau D_{0,\tau}\sum_\sigma |E_\sigma|^2/D_{0,\sigma}$, and $|E_\sigma|^2 = L^2 D_{0,\sigma}|e^{i\theta_\sigma^{(0)}} - 1|^2$ by the identity of Step 1, so they equal $\hat\varepsilon\,\Psi_0^3\prod_\tau D_{0,\tau}$; every other term carries a factor $\hat\varepsilon$. So the denominator is $\hat\varepsilon Y$, the numerator $\hat\varepsilon\prod_{\sigma\notin P}\tilde a_\sigma$, and the quotient is $K_P(\theta)$. Since $|c_P\hat w| = |\hat F_P|$, the minimum at θ is $\sum_P K_P(\theta)|\hat F_P(\theta)|^2$.

Step 3: integration. Integrating the lower bound over θ gives $Q(\varphi) \ge \sum_P \int K_P|\hat F_P|^2$ for every square-summable φ with $V^{(k)}(\varphi) = w$. The minimisers of Step 2 are bounded by a constant times $|\hat w(\theta)|$: for $n \ne 0$ because $\varepsilon_n \ge 4/L^2$ and $a_\mu \ge A_\mu/\hat\varepsilon$; for $n = 0$ because $|c_\mu(\theta^{(0)})|/(\varepsilon_0 a_\mu) \le L^3/A_\mu$; and $|t||E| \le |\hat w|(\max\gamma/\min\gamma)^{1/2}$ by the Cauchy–Schwarz inequality for the inner product Γ, while $|e_0| \le |\theta|/L \le \frac{\pi}{2}|E|/L$. Here $A_\mu \ge (2/\pi)^{10}$, and $\gamma_\mu/\gamma_\nu = \tilde a_\nu/\tilde a_\mu$ is bounded because the $\tilde a$ are continuous and at least A. So the minimisers define a square-summable field φ^* with $V^{(k)}(\varphi^*) = w$ and $Q(\varphi^*) = \sum_P\int K_P|\hat F_P|^2$. Since $V^{(k)}$ and Q are bounded on square-summable fields, truncating φ^* and correcting with ψ_R as in the proof of Theorem 12.12 gives finitely supported fields with the same averages and plaquette forms tending to $Q(\varphi^*)$, and Lemma 12.10(iii) gives (i) for S_k. The functions are even since $|D_L|^2$ and ε are, and positive since $\tilde a_\sigma \ge A_\sigma > 0$ and $Y \ge \Psi_0^3\prod_\tau D_{0,\tau} \ge (2/\pi)^{32}$

on $\mathbb{R}^4$, all other terms of (12.2) being non-negative there. At $\theta = 0$, $\Psi_0 = D_{0,\tau} = 1$ and $\hat{\varepsilon} = 0$, so $K_P(0) = 1$.

Step 4: the strip. Let $|\operatorname{Im}\theta_\nu| \le \frac12$, and for n let $s = L\theta^{(n)}$, so $|\operatorname{Re} s_\nu| \le \pi L$ and $|\operatorname{Im} s_\nu| \le \frac12$. For $q = x + iy$ with $|x| \le \pi$, $|e^{iq} - 1|^2 = (e^{-y} - 1)^2 + 2e^{-y}(1 - \cos x) \ge e^{-|y|}\frac{4}{\pi^2}x^2$, so

$$|D_L(q)| \le \min\Big(e^{1/2}, \frac{c_1}{|\operatorname{Re} Lq|}\Big)$$

for $|\operatorname{Im} q| \le 1/(2L)$, where $c_1 = (e^{1/2}+1)e^{1/4}\frac{\pi}{2} < 5.35$, and the same holds for $D_L^\sharp$. Also $\operatorname{Re}(2 - 2\cos q) \ge (2 - 2\cos x) - 2(\cosh y - 1)$, so $\operatorname{Re} L^2\varepsilon(\theta^{(n)}) \ge \frac{4}{\pi^2}|\operatorname{Re} s|^2 - 1.2$. For $n \ne 0$ some $|\operatorname{Re} s_\nu| \ge \pi$, and then $|L^2\varepsilon(\theta^{(n)})| \ge 0.28\,|\operatorname{Re} s|^2$. Writing $s = \theta + 2\pi m$ with $m \in \mathbb{Z}^4\setminus\{0\}$, $|\operatorname{Re} s_\nu| \ge \pi|m_\nu|$, and

$$|r_\sigma| \le \frac{c_1^{10}}{0.28\,\pi^2}\sum_m\prod_\nu f(m_\nu) = \frac{c_1^{10}}{0.28\,\pi^2}\Big(\frac43\Big)^4 < 2.2\cdot 10^7,$$

with $f(0) = 1$ and $f(j) = 1/(\pi j)^2$. The r_σ are finite sums of functions analytic on the strip, since no $\varepsilon(\theta^{(n)})$, $n \ne 0$, vanishes there. Further $|A_\sigma| \le e^5$, $|\hat{\varepsilon}| \le 4(\pi^2 + \frac14)e^{1/2} < 70$ and $|E_\sigma E_\sigma^\sharp| \le (e^{1/2}+1)^2 < 7.1$, so on the strip $|Y| \le M = 10^{28}$ and $|\tilde{a}_\sigma| \le 1.6\cdot 10^9$, uniformly in k. By the Cauchy estimate on discs of radius $\frac14$ in each variable, $|\partial_\nu Y| \le 4M$ where $|\operatorname{Im}\theta| \le \frac14$, so $|Y(\theta) - Y(\operatorname{Re}\theta)| \le 16\eta M \le \frac12(2/\pi)^{32}$ for $|\operatorname{Im}\theta_\nu| \le \eta = 10^{-36}$. There $|Y| \ge \frac12(2/\pi)^{32}$, $K_P = \prod_{\sigma\notin P}\tilde{a}_\sigma/Y$ is analytic, and $|K_P| \le C = 10^{26}$, which is (ii). Moving the contour of each θ_ν to $\operatorname{Im}\theta_\nu = \pm\eta$, according to the sign of y_ν, gives (iii); the kernels are real because K_P is real and even. □

The proof makes η explicit but very small. In floating point the kernels decrease much faster: $\max|\mathcal{K}_P(y)|$ over $|y|_1 = d$ falls by factors 0.17, 0.24 and 0.26 per unit of d for $k = 1, 2, 3$, the largest value of K_P on $\mathbb{R}^4$ is 4.4, 8.7 and 11.3, and its smallest value is 1, attained at $\theta = 0$ (Figure 14); these computations are reproduced in the archive. Each K_P is a Riemann sum in n of a function of $\theta + 2\pi n$, so the kernels have a limit as $k \to \infty$; the theorem does not use it. Uniform decay of the propagators of the averaging operations of lattice gauge theories is the subject of Bałaban's work [6]; here the hard constraint and the Feynman gauge make the effective form itself explicit. Together with Theorem 12.12, the effective forms of the quadratic approximation are equivalent to the coarse plaquette forms and local, with constants independent of the scale.

Lemma 12.15. *Let (X, d_X) and (Y, d_Y) be Polish spaces, ρ, ρ' Borel probability measures on Y, and κ, κ' probability kernels from Y to X, all with finite first moments. If $W_1(\kappa(\cdot \mid v), \kappa'(\cdot \mid v)) \le \varepsilon$ and $W_1(\kappa'(\cdot \mid v), \kappa'(\cdot \mid v')) \le d_Y(v, v')$ for all v, v', then the mixtures $\pi = \int \kappa(\cdot \mid v)\,\rho(dv)$ and $\pi' = \int \kappa'(\cdot \mid v)\,\rho'(dv)$ satisfy*

$$W_1(\pi, \pi') \le \varepsilon + W_1^{d_Y}(\rho, \rho').$$

Proof. Let F be 1-Lipschitz on X and $\kappa F(v) = \int F\,d\kappa(\cdot \mid v)$. Then $\int F\,d\pi - \int F\,d\pi' = \int(\kappa F - \kappa' F)\,d\rho + \big(\int \kappa' F\,d\rho - \int \kappa' F\,d\rho'\big)$. The first term is at most ε. The function $\kappa' F$ is 1-Lipschitz for d_Y, so the second is at most $W_1^{d_Y}(\rho, \rho')$. The Kantorovich–Rubinstein duality gives the claim. □

A law γ on $\mathbb{R}^V$ *decomposes along the averages* if $\gamma(\cdot \mid \eta) = \int \gamma_V^{\mathrm{c}}(\cdot \mid \eta, v)\,\rho_V(dv \mid \eta)$, with $\rho_V(\cdot \mid \eta)$ the law of $(V_{y,\mu})_{y\in W}$. Every law with density proportional to $\exp(-\frac{b}{2}\sum_{p\cap V\ne\emptyset}(d\varphi)_p^2)\,g(V(\varphi))$, on a set cut out by conditions on the averages, decomposes in this way, because Lebesgue measure disintegrates along the fibres of a surjective linear map into multiples of Lebesgue measure on the fibres. Here g can carry any action of the averages, in particular a coarse gauge fixing and a coarse effective action, which is the form a law takes after the fine scale has been integrated out.




FIGURE 14. The effective forms of Theorem 12.14. (a) The multiplier K_P for the plane $P=\{1,2\}$ at $k=3$, as a function of (θ_3,θ_4) with $\theta_1=\theta_2=0$; it equals 1 at $\theta=0$ and stays bounded. (b) The largest value of the kernel $|\mathcal{K}_P(y)|$ on $|y|_1=d$ for $k=1,2,3$, in logarithmic scale: the decay per coarse spacing hardly changes with k.

Corollary 12.16. *Let $r_w\in\mathbb{R}^V$ minimise $\sum_{p\cap V\neq\emptyset}(d\varphi)_p^2$, with zero exterior, subject to $V_{y',\mu'}(r_w)=\delta_{w,(y',\mu')}$ for all (y',μ'), $y'\in W$; put $L_w=\|r_w\|_1$ and $d_L(v,v')=\sum_w L_w|v_w-v'_w|$. For every law that decomposes along the averages,*

$$C_{V,j}\le C^{\mathrm{c}}_{V,j}+C^{\rho}_{V,j},\qquad C^{\rho}_{V,j}=\sup\frac{W_1^{d_L}(\rho_V(\cdot\mid\eta),\rho_V(\cdot\mid\eta'))}{|\eta_j-\eta'_j|},$$

the supremum over exteriors that differ only at j, and $L_w\le 4\cdot10^{14}$. Hence $K_c\le K_c^{\mathrm{c}}+K_c^{\rho}$ with $K_c^{\rho}=\sum_{j\in c\setminus V}C^{\rho}_{V,j}$, and if $K_c^{\rho}\le\frac{\alpha}{2}N_n$ for every class and $n\ge4\cdot10^{15}/\alpha$, then $D(n,\alpha)$ holds for the law, with the 49 classes of active links.

Proof. Apply Lemma 12.15 with $\kappa=\gamma^{\mathrm{c}}_V(\cdot\mid\eta,\cdot)$, $\kappa'=\gamma^{\mathrm{c}}_V(\cdot\mid\eta',\cdot)$, $\rho=\rho_V(\cdot\mid\eta)$, $\rho'=\rho_V(\cdot\mid\eta')$. The first condition holds with $\varepsilon=C^{\mathrm{c}}_{V,j}|\eta_j-\eta'_j|$. By Theorem 12.8(i) the laws $\gamma^{\mathrm{c}}_V(\cdot\mid\eta',v)$ and $\gamma^{\mathrm{c}}_V(\cdot\mid\eta',v')$ are translates by $\sum_w(v'_w-v_w)r_w$, so their distance is at most $d_L(v,v')$. For the bound on L_w, the pair (r_w,y) solves $K(r_w,y)=(0,s\,e_w)$ with K as in the proof of Theorem 12.8. Measure d from the centre of the row w; Lemma 12.7 gives $\sum_e e^{2\lambda d(e)}r_w(e)^2\le(2M_0s)^2=20160^2$. With $z_0=2y+e_\mu$ for $w=(y,\mu)$, a link whose base is at distance k from z_0 has its midpoint at distance at least $k-1$ from the centre, and the count in that proof gives $L_w^2\le20160^2\cdot4e^{2\lambda}(1+\sum_{k\ge1}(64k^3+16k)e^{-2\lambda k})\le(4\cdot10^{14})^2$. Summing over the translates as in Definition 10.1 gives $K_c\le K_c^{\mathrm{c}}+K_c^{\rho}$, and Theorem 12.8(iii) with $\alpha/2$ in place of α gives $K_c^{\mathrm{c}}\le\frac{\alpha}{2}N_n$. □

The responses are much smaller than the bound: in floating point, $\max_w L_w$ is 23.8, 44.1 and 50.9 for the block (m,m,m,m), $m=\lfloor n/4\rfloor$, of the cubes $n=4$, 6, 8. Since Theorem 12.8 controls the first term at every coupling, the corollary reduces $D(n,\alpha)$ for a law that decomposes along the averages to the cube condition K_c^{ρ} for its averages, a problem of the same form one scale up, in the metric d_L. Iterating it needs Theorem 12.8 and the corollary with S_k in place of the plaquette form, which Theorem 12.18 below gives with constants independent of k, and ends in a cube condition for the coarsest averages, which is where a gap has to come from: the quadratic approximation describes a massless field and has none. The corollary is one step of the composition of Remark 12.19(a), not its conclusion.

On the lattice of spacing 2^k the blocks, the tree links, the active links and the averages are those of Definition 12.2, and after k steps the averages of the quadratic approximation

carry the form S_k. Let (S_{ab}) be its matrix, $S_k(w)=\sum_{a,b}S_{ab}w_aw_b$ for finitely supported w, and $d(a,b)$ the ℓ^∞ distance between the midpoints of the links a and b. Each link lies in six plaquettes of four links, and the midpoints of the links of the plaquettes at y and y' are at distance at most $|y-y'|_1+1$, so Theorem 12.14(iii) gives, for every non-decreasing $\phi\ge 0$,

$$\sum_b |S_{ab}|\,\phi(d(a,b)) \le 24C\sum_{z\in\mathbb{Z}^4} e^{-\eta|z|_1}\phi(|z|_1+1), \tag{12.3}$$

with C and η of that theorem. In particular $\sum_b S_{ab}\eta_b$ converges for every bounded field η. For n, V, W as before, a bounded exterior η and values v, let $\gamma_V^{\mathrm{c},k}(\cdot\mid\eta,v)$ be the law on $\mathbb{R}^V$ with density proportional to

$$\exp\Big(-\frac{b}{2}\Big(\sum_{a,b\in V}S_{ab}w_aw_b+2\sum_{a\in V,\,b\notin V}S_{ab}w_a\eta_b\Big)\Big)$$

on the affine set $\{V_{y,\mu}(w)=v_{y,\mu},\ y\in W\}$, the conditional law of the Gaussian with the form bS_k given the exterior and the averages. Define $C_{V,j}^{\mathrm{c},k}$, $K_{\mathrm{c}}^{\mathrm{c},k}$, $D_{\mathrm{c}}^k(n,\alpha)$, $r_w^{(k)}$ and $L_w^{(k)}$ as in Theorem 12.8 and Corollary 12.16, with S_k in place of the plaquette form; for $k=0$ they are the former ones.

Lemma 12.17. *Let K be an invertible symmetric matrix indexed by a finite set X, with $\|K^{-1}\|\le M_0$, and let $d:X\to\mathbb{R}$ and $\lambda>0$. If $\sum_b|K_{ab}|(e^{\lambda|d(a)-d(b)|}-1)\le\frac{1}{2M_0}$ for every a, then $\|e^{\lambda d}K^{-1}e^{-\lambda d}\|\le 2M_0$.*

Proof. $e^{\lambda d}Ke^{-\lambda d}=K+E$ with $|E_{ab}|\le|K_{ab}|(e^{\lambda|d(a)-d(b)|}-1)$, a symmetric bound, so $\|E\|\le\frac{1}{2M_0}$ by the Schur test, and the proof of Lemma 12.7 applies. □

Theorem 12.18. *Let $\lambda_1=10^{-215}$. For every $k\ge1$, every $b>0$, every even n and every exterior link j:*

(i) *$\gamma_V^{\mathrm{c},k}(\cdot\mid\eta,v)$ is a Gaussian law whose covariance depends on neither η nor v and whose mean is affine in (η,v), and $C_{V,j}^{\mathrm{c},k}=\|u_j\|_1$, where $u_j\in\mathbb{R}^V$ minimises $S_k(u+\delta_j)$ subject to $Bu=g$, with B and g as in Theorem 12.8;*

(ii) *$\sum_{e\in V}e^{2\lambda_1 d(e)}u_j(e)^2\le10^{350}$ and $\|u_j\|_1\le10^{607}e^{-\lambda_1 r_j/2}$, where $d(e)=d(e,j)$ and $r_j=\min_{e\in V}d(e)$; and $L_w^{(k)}\le10^{438}$ for every w;*

(iii) *$D_{\mathrm{c}}^k(n,\alpha)$ holds for every $\alpha\in(0,1)$ and every even $n\ge10^{827}/\alpha$;*

(iv) *Corollary 12.16 holds on the lattice of spacing 2^k, with S_k in place of the plaquette form, $L_w^{(k)}\le10^{438}$ and $n\ge2\cdot10^{827}/\alpha$.*

None of these constants depends on k or on b.

Proof. (i) Let $H=(S_{ab})_{a,b\in V}$. A vector $w\in\ker B$, extended by zero, lies in the space Φ_W of the lattice of spacing 2^k, so Proposition 12.9(iv) gives $\langle w,Hw\rangle=S_k(w)\ge\frac{1}{52}|w|^2$ on $\ker B$. The proof of Theorem 12.8(i) applies, the exterior entering through the linear term $\sum_{b\notin V}S_{ab}\eta_b$; changing η_j by one translates the mean by the minimiser of $\langle u,Hu\rangle+2\sum_{a\in V}u_aS_{aj}=S_k(u+\delta_j)-S_k(\delta_j)$ subject to $Bu=g$.

(ii) The pair (u_j,y) solves $K(u_j,y)=(f,sg)$ with K built from H and B as in Lemma 12.6 and $f=-(S_{aj})_{a\in V}$. Since $Q_k(w)\le24|w|^2$, Theorem 12.12 gives $\|H\|\le h=6144$; take $s=4h=24576$. The bounds of Lemma 12.6 for (h,s) are at most those for $(24,96)$, because $|x_1|$ and $|f|/s$ decrease while $h|x_1|$ and h/s do not change, so $\|K^{-1}\|\le M_0=105$ as in the proof of Theorem 12.8. Index the rows of B by centres as there. For $\phi(r)=e^{\lambda r}-1\le\lambda re^{\lambda r}$ and $\lambda\le\eta/2$, (12.3) bounds the row sums of Lemma 12.17 over H by

$$24C\lambda e^{\lambda}\sum_z(|z|_1+1)e^{-\eta|z|_1/2}$$
$$\le 24C\lambda e^{\lambda}\Big(1+16\cdot24\,e^{\eta/2}\Big(\frac{2}{\eta}\Big)^5\Big),$$

since at most $8r^3$ points have $|z|_1 = r \geq 1$ and $\sum_{r\geq 1} r^p e^{-xr} \leq e^x p!/x^{p+1}$. The entries of $sB^\top$ add at most $\frac{s}{8}(e^\lambda - 1)$ and a row of sB gives $2s(e^\lambda - 1)$. At $\lambda = \lambda_1$ the sums are below $3 \cdot 10^{-4} < \frac{1}{2M_0}$, so Lemma 12.17 gives $\|e^{\lambda_1 d} K^{-1} e^{-\lambda_1 d}\| \leq 210$. By (12.3), $\|e^{\lambda_1 d} f\| \leq 24 C e^{\lambda_1} \sum_z e^{-(\eta-\lambda_1)|z|_1} = 24 C e^{\lambda_1} \coth^4 \frac{\eta-\lambda_1}{2} < 4 \cdot 10^{172}$, and $\|e^{\lambda_1 d} sg\| \leq e^{\lambda_1} s/16$, which gives the first bound. The links with base at ℓ^∞ distance m from the base of j number at most $4(64m^3 + 16m)$, with midpoints at distance at least $m - 1$, so

$$Z(x) = \sum_e e^{-x d(e)} \leq 4e^x \Big(1 + \frac{384 e^x}{x^4} + \frac{16 e^x}{x^2}\Big),$$

and $Z(\lambda_1) < 10^{864}$. As $d \geq r_j$ on V, the Cauchy–Schwarz inequality gives $\|u_j\|_1^2 \leq 10^{350} e^{-\lambda_1 r_j} Z(\lambda_1)$. Finally $(r_w^{(k)}, y)$ solves $K(r_w^{(k)}, y) = (0, s e_w)$; measuring d from the centre of the row w, we get $\sum_e e^{2\lambda_1 d(e)} r_w^{(k)}(e)^2 \leq (210 s)^2$ and $L_w^{(k)} \leq 210 s\, Z(2\lambda_1)^{1/2} < 10^{438}$.

(iii) The exterior links of a class have their bases at ℓ^∞ distance $m \geq 1$ from $[0, n-1]^4$; at most $4((n+2m)^4 - (n+2m-2)^4) \leq 128(n^3 + 8m^3)$ of them have distance m, and for them $r_j \geq m - 1$. With $q = \lambda_1/2$, (ii) gives

$$K_c^{\mathrm{c},k} \leq 10^{607} \cdot 128\, e^{2q} \Big(\frac{n^3}{q} + \frac{48}{q^4}\Big),$$

which is at most $\alpha N_n = \alpha (n/2)^4$ when $n \geq 10^{827}/\alpha$.

(iv) The proof of Corollary 12.16 uses the plaquette form only through Theorem 12.8, the translation property of (i) and the bound on L_w, which (i)–(iii) provide. □

The constants are poor because the strip of Theorem 12.14 has width 10^{-36}; they are checked in exact arithmetic in the archive. What the theorem adds is that none of them depends on the scale: the constrained cube condition holds with one cube side at every scale, and the weights $L_w^{(k)}$ of the metrics d_L are bounded uniformly. It does not compose the scales. The cube condition K_c^ρ that Corollary 12.16 leaves at scale k concerns the law of the averages given the fine exterior, in the metric d_L, and it has to be turned into the cube condition of the law one scale up; that step and the cube condition for the coarsest averages are open, and in the quadratic approximation the coarsest averages carry a massless Gaussian, so a gap has to come from beyond it.

Remark 12.19. Theorem 12.8 is the weak-coupling half of the constrained route at leading order, and Theorem 10.4 cannot use it as it stands. Four steps remain, each an open problem.

(a) *Composition.* D_{c} holds the block averages fixed. A gap in physical units needs a theorem that combines the constrained conditions at all scales with a cube condition for the coarsest averages, in the place of Theorem 10.4. Corollary 12.16 is one step of it, Proposition 12.9(iv) gives the gap at every scale, and Theorem 12.12 the equivalence of S_k and Q_k uniformly in k, Theorem 12.14 the locality of S_k uniformly in k, and Theorem 12.18 the constrained cube condition and the bounds on the weights L_w at every scale with constants independent of k. Turning the cube condition for the averages into that of the law one scale up, and a cube condition for the coarsest averages, are open.

(b) *Beyond the quadratic approximation.* For the Wilson measure the block averages are products of group elements along paths, their level sets are curved, and the Bakry–Émery tensor of Section 6 is negative at large fields; Theorem 12.4 controls only the small fields, and the large fields have to be split off as in Bałaban's analysis [8]. For the straight paths between block roots the group-valued estimate holds for all fields (Theorem 12.21), but that map does not compose across scales (Proposition 12.24), and for every compact connected group no continuous covariant map treats the transverse positions of a block alike (Proposition 12.25). The cut-off average of Theorem 12.27 is coercive for all fields and agrees with the averages to first order, with constants for one

step; on torus-valued fields of small curvature it composes (Theorem 12.29), and for every compact group it composes on fields whose curvature is small at every scale with a small sum (Theorem 12.33); under the Wilson measure the remaining large fields form microscopic clusters (Corollary 12.58), but the conditional estimate needed near them is open. Away from them, on the fibres of the cut-off average, the Hessian of the action along the fibre is at least $\lambda(\frac{1}{26} - c_G\kappa)$ at every background with plaquettes at most κ (Theorem 12.35), and on a convex chart of the fibre the constrained law satisfies a Brascamp–Lieb inequality for large λ (Theorem 12.37), and on a product chart its covariances decay exponentially (Theorem 12.41), and the mass off that chart is small on most fibres of the SU(2) Wilson measure (Theorem 12.44) and on every fibre through a field with small plaquettes (Corollary 12.49), and on every fibre the law concentrates near the least action of its regular fields (Proposition 12.51); the remaining small-field steps are listed in Remark 12.55.

(c) *Second order.* Let $b_0 = 11N/(48\pi^2)$ for SU(N), $t = 2b_0 \log 2$, and let $g_m^{-2} = g_*^{-2} + tm$ be the one-loop coupling m steps above a matching scale. Comparison with integrals gives $\sum_{m\le M} g_m^2 = t^{-1}\log M + O(1)$ and $\sum_m g_m^4 \le \pi^2/(6t^2)$ [56]. A loss $1 + cg_m^2$ per step therefore compounds to a power of M, and a step from one scale to the next has to be exact at order g^2, with that term absorbed into the running coupling and the normalisation of the field, and remainders of order g^4.

(d) *Crossover and existence.* The passage from weak coupling to the strong-coupling range of Section 4–8 and the existence of the limit, (E), are untouched.

The renormalisation group needed for (a) and (b) is the ultraviolet half of (E), so the constrained route places both residues on one construction.

12.2. **Path blocks for the group-valued links.** Remark 12.19(b) asks for estimates on the actual group-valued links. They can be proved for a coarser block map, the straight path between block roots, and this map exhibits the second half of the difficulty: it does not compose across scales. A topological argument shows why no continuous map that treats the transverse positions alike can take its place.

Let $G \subset \mathrm{U}(r)$ be a compact matrix group, $\delta(A,B) = \|A-B\|_{\mathrm{HS}}$, which is invariant under left and right multiplication, and work on the periodic lattice of side $2L$ with the block tree of Definition 12.2, all fifteen tree links of each block equal to I. Write $\mathcal{B}$ for the number of blocks. The *path map* is

$$V^{\mathrm{p}}_{y,\mu}(U) = U_{2y+e_\mu,\mu},$$

the product along the straight path from $2y$ to $2y+2e_\mu$, whose first link is a tree link; it transforms as a link under the gauge transformations at the block roots. Fixing $V^{\mathrm{p}} = v$ fixes four of the 49 active links of each block, so the fibre is $G^{45\mathcal{B}}$, and the Wilson law in the tree gauge restricts to it with product Haar measure as reference, since the tree gauge has unit Jacobian. The *reconstruction* $R(v)$ puts $R_{2y+o,\mu} = v_{y,\mu}$ if $o_\mu = 1$ and $R_{2y+o,\mu} = I$ otherwise. For U in the fibre of v put

$$D_v(U)^2 = \sum_e \delta(U_e, R_e(v))^2, \qquad Q(U) = \sum_p \delta(U_p, I)^2,$$

the first sum having zero terms on the tree and path links, and let $Q_1(v) = \sum_P \delta(v_P, I)^2$ on the coarse lattice.

Lemma 12.20. *For vectors z_a of a real Hilbert space indexed by $a \in \{0,1\}^3$, with $z_0 = 0$,*

$$\sum_a \|z_a\|^2 \le \frac{3+\sqrt{7}}{2} \sum_{\{a,a'\}} \|z_a - z_{a'}\|^2 \le 3 \sum_{\{a,a'\}} \|z_a - z_{a'}\|^2,$$

the sums on the right over the twelve edges of the cube.

Proof. It suffices to treat real numbers. The right side is $\langle z, L_0 z\rangle$, where L_0 is the Laplacian of the cube with the vertex 0 deleted, and the claim is that its least eigenvalue is $3 - \sqrt{7}$.

The permutations of the coordinates commute with L_0. On the functions f_k of the level $k = |a|$, L_0 acts by $f_1 \mapsto 3f_1 - 2f_2$, $f_2 \mapsto 3f_2 - 2f_1 - f_3$, $f_3 \mapsto 3f_3 - 3f_2$, whose characteristic polynomial in $t = 3 - \lambda$ is $t^3 - 7t$; the eigenvalues are 3 and $3 \pm \sqrt{7}$. On the functions of levels one and two whose sum over each level vanishes, indexed by the coordinate that is one at level one and zero at level two, L_0 acts by the matrix $\left(\begin{smallmatrix} 3 & 1 \\ 1 & 3 \end{smallmatrix}\right)$, with eigenvalues 2 and 4; at level three these functions vanish. The two spaces have dimensions 3 and 4 and span $\mathbb{R}^7$. Finally $1/(3 - \sqrt{7}) = (3 + \sqrt{7})/2 < 3$.[20] □

Theorem 12.21. *For every $G \subset \mathrm{U}(r)$, every $L \geq 1$, every v and every U in the fibre of v,*

$$D_v(U)^2 \leq 224\, Q(U).$$

Proof. Let I_0 be the sum of $\delta(U_e, I)^2$ over the active links inside the blocks and J_0 the sum of $\delta(U_e, v_{y,\mu})^2$ over the links that leave $B(y)$ in the direction μ, so $D_v^2 = I_0 + J_0$. For an active link inside a block, the chain of plaquettes in the proof of Theorem 12.4 applies to group elements: each plaquette of the chain has its two transverse links on the tree, so the holonomy is the product of two consecutive links of the chain, and invariance of δ with the triangle inequality gives $\delta(U_e, I)^2 \leq k \sum_i \delta(U_{p_i}, I)^2$ with $k \leq 3$. The loads are those of that proof, so $I_0 \leq 8Q_{\mathrm{in}}$. Fix y, μ and label the eight links leaving $B(y)$ by the transverse offset $a \in \{0,1\}^3$; the link $a = 0$ is the path link and equals $v_{y,\mu}$. For a transverse edge $\{a, a'\}$ the face plaquette has holonomy $U_a \psi' U_{a'}^{-1} \psi^{-1}$ with ψ, ψ' links inside the two blocks, so $\delta(U_a, U_{a'}) \leq \delta(U_p, I) + \delta(\psi, I) + \delta(\psi', I)$. Lemma 12.20 with $z_a = U_a - v_{y,\mu}$ in the real Hilbert–Schmidt space and Cauchy–Schwarz bound the eight terms by 9 times the sum of the three squares over the twelve edges. Each face plaquette occurs once, and each link inside a block occurs once for each of the three directions other than its own, as in the proof of Theorem 12.4. Hence $J_0 \leq 9Q_{\mathrm{face}} + 27 I_0$ and $D_v^2 \leq 28 I_0 + 9 Q_{\mathrm{face}} \leq 224\, Q$. □

The bound holds for all fields, with no small-field condition and no logarithmic chart; Theorem 12.4 is its linearisation for the averages, with the better constant 52.

Proposition 12.22. *$Q_1(V^{\mathrm{p}}(U)) \leq 4Q(U)$, $Q(R(v)) = 4Q_1(v)$ and $Q(U) \leq 8Q_1(v) + 48 D_v(U)^2$ for U in the fibre of v; if $Q_1(v) = 0$, then $Q(U) \leq 24 D_v(U)^2$.*

Proof. The holonomy of a coarse plaquette is that of the square of side two at $2y$, a product of four conjugated fine plaquette holonomies, so $\delta(v_P, I) \leq \sum_{p \subset P} \delta(U_p, I)$; these squares tile each coordinate plane, which gives the first bound. In $R(v)$ only the plaquettes with offset one in both directions carry holonomy, equal to that of the coarse plaquette, four for each P. For any plaquette, telescoping gives $\delta(U_p, R_p) \leq \sum_{e \in p} \delta(U_e, R_e)$, so $\delta(U_p, I)^2 \leq 2\delta(R_p, I)^2 + 8 \sum_{e \in p} \delta(U_e, R_e)^2$, and each link lies in six plaquettes. If $Q_1(v) = 0$, every R_p is I and the factor 2 is not needed. □

Let $m = 45\mathcal{B}$, let the action be λQ with $\lambda \geq 0$ (for $\mathrm{SU}(N)$, $\lambda = \beta/(2N)$ up to a constant), and put

$$z_G(t) = \int_G e^{-t\delta(g,I)^2} dg, \qquad Z_\lambda(v) = \int_{G^m} e^{-\lambda Q(U)} dU,$$

with normalised Haar measures, the integral over the fibre of v.

Theorem 12.23. *For every v,*

$$e^{-8\lambda Q_1(v)} z_G(48\lambda)^m \leq Z_\lambda(v) \leq e^{-\lambda Q_1(v)/8} z_G(\lambda/448)^m,$$

and if $Q_1(v) = 0$, then $z_G(24\lambda)^m \leq Z_\lambda(v) \leq z_G(\lambda/224)^m$. Let $A_\lambda = z_G(\lambda/448)/z_G(24\lambda)$. Under the conditional law at such a v,

$$\mathbb{E}\, e^{\lambda D_v^2/448} \leq A_\lambda^m, \qquad \mathbb{E}\, \delta(U_e, I)^2 \leq \frac{20160}{\lambda} \log A_\lambda$$

for each link e when $v \equiv I$, and $A_\lambda \to 10752^{q/2}$ as $\lambda \to \infty$ if G has dimension $q \geq 1$.

[20] The constant $(3 + \sqrt{7})/2$ is attained by a function of the level $|a|$ alone, so it is the best constant in the lemma.

Proof. By Theorem 12.21 and Proposition 12.22, $Q \geq \frac{1}{2}(D_v^2/224 + Q_1/4)$ and $Q \leq 8Q_1 + 48D_v^2$. Each term of D_v^2 involves one free link, and right multiplication by $R_e(v)^{-1}$ preserves Haar measure, so $\int e^{-tD_v^2} dU = z_G(t)^m$; this gives the two bounds, and the flat case is the same with $Q \leq 24D_v^2$. Next, $-\lambda Q + \lambda D_v^2/448 \leq -\lambda D_v^2/448$, and integration with the lower bound gives the exponential moment; Jensen's inequality gives $\mathbb{E}D_v^2 \leq (448m/\lambda)\log A_\lambda$. When $v \equiv I$ the law is invariant under translation by blocks, so the $\mathcal{B}$ links of each of the 45 classes have equal expectations, and $448 \cdot 45 = 20160$. Finally, in exponential coordinates $\delta(e^X, I)^2 = \|X\|^2 + O(\|X\|^4)$ and the Haar density is smooth and positive at 0, while $\delta(\cdot, I)$ is bounded below away from I; so $z_G(t) \sim c_G t^{-q/2}$, and the ratio at $\lambda/448$ and 24λ tends to $(24 \cdot 448)^{q/2}$. □

So the fluctuations of the free links around a flat path field are of order $\lambda^{-1/2}$ uniformly in the volume, for the group variables themselves. The path map cannot be iterated, however, and this is where it differs from the averages. Write $V^{\mathrm{p},k}$ for the k-fold path map, the straight path of length $s = 2^k$, and Q_k for the plaquette energy of the lattice of spacing s.

Proposition 12.24. *On the periodic lattice of side* $3s$, $s = 2^k$, *there are* SU(2) *fields in the tree gauge with*

$$\frac{Q(U)}{Q_k(V^{\mathrm{p},k}(U))} = s\sin^2\frac{\pi}{2s} \leq \frac{\pi^2}{4s}.$$

For the averages, by contrast, $Q_k(V^{(k)}(\varphi)) \leq Q(\varphi)$ *for every* k *in the quadratic approximation (Proposition 12.9(i)).*

Proof. Let $T = \mathrm{diag}(i\pi/2, -i\pi/2)$. Put the $s/2$ links in the direction 0 at $(j, 0, 0, 0)$, j odd, $j < s$, equal to $e^{2T/s}$ and all other links equal to I. These links have offset one in the direction 0, so the tree gauge holds, and the links at even j are tree links. The path from 0 to se_0 carries e^T and every other path carries I, so one coarse link is e^T, and it lies in six coarse plaquettes, each with $\delta^2 = \|e^T - I\|_{\mathrm{HS}}^2 = 4$: $Q_k = 24$. No fine plaquette contains two of the changed links, since the two links of a plaquette in the direction 0 differ in a transverse coordinate, so there are $3s$ nonzero fine plaquettes, each with $\delta^2 = 8\sin^2(\pi/2s)$, and $Q = 24s\sin^2(\pi/2s)$. Gauge covariance of the path map makes the ratio gauge invariant. □

A field concentrated on one thin path carries a coarse curvature of order one at a fine cost of order $1/s$. In the averages it is one transverse position among s^3 and enters with the weight s^{-3}, which is why the averages compose and the paths do not. The two properties needed for Remark 12.19(a) and (b) are therefore held by two different maps: the paths give a global estimate for the group-valued links on fibres that do not compose, and the averages compose on linear fibres that exist only in the quadratic approximation.

A block map with both properties cannot treat the transverse positions alike and be continuous, for any compact connected gauge group. Let $s = 2^k$ and let V be a map of the links with both ends in $B(y) \cup B(y + e_\mu)$, blocks of side s, to G. Call it *covariant* if $V(U^h) = h_1 V(U) h_2^{-1}$ for every gauge transformation h equal to h_1 on $B(y)$ and to h_2 on $B(y + e_\mu)$, and *normalised* if $V(I) = I$. The path map and its iterates are covariant, normalised and continuous. Let T be the set of the s^3 links in the direction μ from $B(y)$ to $B(y + e_\mu)$, and for $u \in G^T$ let $U(u)$ carry u_t on the link $t \in T$ and I on every other link (Figure 15). Let $S \subset G$ be a maximal torus, of dimension ℓ; a basis of the kernel of its exponential map identifies S with $(\mathbb{R}/2\pi\mathbb{Z})^\ell$ and its first homology with $\mathbb{Z}^\ell$.

Proposition 12.25. *Let* $G \subset \mathrm{U}(r)$ *be a compact connected Lie group of positive dimension, let* V *be covariant and normalised, and let* $F(u) = V(U(u))$ *be continuous on* G^T. *Then* F *maps* S^T *into* S, *and the integer matrices* W_t *by which the maps* $u_t \mapsto F(u)$, $u_t \in S$, $u_{t'} = I$ *for* $t' \neq t$, *act on* $H_1(S) = \mathbb{Z}^\ell$ *satisfy*

$$\sum_{t \in T} W_t = I_\ell.$$

Hence some $W_t \neq 0$, so that the map $u_t \mapsto V(U(u))$ on fields equal to I off the link t is not homotopic to a constant on S, and, if $|T| \geq 2$, no transitive group of permutations of T leaves F invariant. For $G = \mathrm{SU}(2)$, $\ell = 1$, the W_t are integers adding up to one, and some field equal to I off one link of T has $V = -I$.

Proof. The gauge transformation equal to g on $B(y) \cup B(y + e_\mu)$ maps $U(u)$ to $U(gug^{-1})$, since the links equal to I stay I, and the one equal to g on $B(y)$ and to I on $B(y + e_\mu)$ maps it to $U(gu)$. So $F(gug^{-1}) = gF(u)g^{-1}$ and $F(gu) = gF(u)$, and with $F(I) = I$ the second gives $F(g, \dots, g) = g$. If every u_t lies in S, every $g \in S$ fixes u under conjugation, so $F(u)$ commutes with S; in a compact connected Lie group a maximal torus is its own centraliser [44], so $F(u) \in S$. Therefore F restricts to a continuous map $f : S^T \to S$. The first homology of S^T is $\bigoplus_{t \in T} \mathbb{Z}^\ell$, the t-th summand coming from the inclusion of the t-th factor with the identity elsewhere, so f acts on it by $(x_t)_t \mapsto \sum_t W_t x_t$. The diagonal map $S \to S^T$ acts by $x \mapsto (x, \dots, x)$, and f composed with it is the identity of S because $F(g, \dots, g) = g$. Hence $\sum_t W_t = I_\ell$. If all W_t vanished the sum would vanish, and a map of tori that is homotopic to a constant acts by zero on homology. A transitive permutation group preserving F would make all W_t equal to one matrix W, and $|T|\, W = I_\ell$ has no integer solution for $|T| \geq 2$. For $\mathrm{SU}(2)$, S is a circle through $-I$, and a loop of nonzero degree in a circle is onto. □

For the path map, W_t is the identity at the position of the path and zero elsewhere. Averages of the path holonomies weigh the positions alike, so by the proposition they have no continuous covariant extension to all of G^T, for $\mathrm{SU}(N)$ and for every compact simple group. Bałaban's averages are defined only on fields whose plaquettes are small, and the large fields are split off [6, 8]. A block map with both properties for all fields is therefore discontinuous or, like the path map, not invariant under the transverse symmetries of the block. Theorem 12.27 below gives a continuous one whose departure from the symmetry is confined to large fields.[21]

The proposition leaves room for a map that treats the positions alike to first order and breaks the symmetry only where the field is large. For a coarse link (y, μ) and $x \in B(y)$ put

$$P_x = U_{x,\mu} U_{x+e_\mu,\mu},$$

the product along the straight path of length two from x to $x + 2e_\mu$. In the tree gauge it is the parallel transport from $2y$ to $2y + 2e_\mu$ through x and $x + 2e_\mu$, it transforms as a link under the gauge transformations at the block roots, and $P_{2y} = V^{\mathrm{p}}_{y,\mu}$, since the first link of that path is a tree link. Let $\mathfrak{g}$ be the Lie algebra of G, $\|X\|$ the Hilbert–Schmidt norm and log the principal logarithm of unitary matrices without the eigenvalue -1.

Lemma 12.26. *If g is unitary and $\delta(g, I) \leq 1$, then $\|\log g\| \leq \frac{\pi}{3}\, \delta(g, I)$. If X is skew-Hermitian, then $\delta(e^X, I) \leq \|X\|$.*

Proof. Let $e^{i\theta_j}$, $|\theta_j| \leq \pi$, be the eigenvalues of g. Then $\delta(g, I)^2 = \sum_j 4 \sin^2(\theta_j/2)$ and $\|\log g\|^2 = \sum_j \theta_j^2$. Each term of the first sum is at most 1, so $|\theta_j| \leq \pi/3$, and $\theta/(2\sin(\theta/2))$ increases on $(0, \pi)$ and equals $\pi/3$ at $\theta = \pi/3$. The second statement is $|e^{i\theta} - 1| \leq |\theta|$ for the eigenvalues of X. □

[21] That spheres and tori carry no continuous symmetric mean goes back to Aumann [2] and Eckmann [34]; the proof above is their homology argument in gauge-covariant form, with the links of the slice T as the arguments of the mean and a maximal torus in place of the circle.




FIGURE 15. (a) Two adjacent blocks $B(y)$, $B(y+e_\mu)$ of side $s=4$, drawn in three of the four dimensions. The black segment is the straight path between the roots, the coarse link of the path map. The blue links form the slice T of Proposition 12.25. The red links carry the field of Proposition 12.24: a path whose holonomy is that of a coarse link of order one, at a fine energy of order $1/s$. (b) For SU(2), two coordinates of the torus $(\mathbb{R}/2\pi\mathbb{Z})^T$ of slice fields with values in a maximal torus. The diagonal is pure gauge and is sent to a loop of degree one, so the degrees W_t of the coordinate loops add up to one.

Let $\rho_G \in (0,1]$ be such that $\log g \in \mathfrak{g}$ for every $g \in G$ with $\delta(g,I) \le \rho_G$.[22] Let $\phi(t) = \min\{1, \max\{0, 2-2t\}\}$, which equals 1 on $[0,\frac12]$ and 0 on $[1,\infty)$, put

$$\chi_{y,\mu}(U) = \phi\Big(\rho_G^{-1} \max_{x \in B(y)} \delta(P_x, P_{2y})\Big),$$

and define the *cut-off average*

$$V^\sharp_{y,\mu}(U) = P_{2y} \exp\Big(\frac{\chi_{y,\mu}(U)}{16} \sum_{x \in B(y)} \log(P_{2y}^{-1} P_x)\Big), \tag{12.4}$$

the exponential factor being I when $\chi_{y,\mu}(U)=0$ (Figure 16). When every path is close to the straight one it is the average of the sixteen paths taken in the chart at P_{2y}; when some path is far from it, it is the path map.

Theorem 12.27. *For every compact $G \subset \mathrm{U}(r)$ the cut-off average is continuous, covariant and normalised, and the following hold, with $c = \pi/3$.*

(i) *If $\delta(P_x, P_{2y}) \le \rho_G/2$ for every $x \in B(y)$, then $V^\sharp_{y,\mu}(U) = P_{2y}\exp(\frac{1}{16}\sum_x \log(P_{2y}^{-1}P_x))$. For φ with values in $\mathfrak{g}$, vanishing on the tree links,*

$$V^\sharp_{y,\mu}(e^{t\varphi}) = I + t\, V_{y,\mu}(\varphi) + O(t^2),$$

with $V_{y,\mu}$ the block average of Definition 12.2.

[22] The exponential map of G is a diffeomorphism from a ball of $\mathfrak{g}$ of operator-norm radius less than π onto a neighbourhood N of I in G, on which it inverts the principal logarithm, and N contains $\{g \in G : \delta(g,I) \le \rho\}$ for small $\rho > 0$. For $\mathrm{U}(r)$ one may take $\rho_G = 1$. For $\mathrm{SU}(N)$ one may take $\rho_G = \min\{1, 5/\sqrt{N}\}$: by Lemma 12.26 the trace of $\log g$ has modulus at most $\sqrt{N}\,\pi\rho_G/3 < 2\pi$, and it lies in $2\pi i\mathbb{Z}$. Some restriction is needed: on the circle $\{\mathrm{diag}(z, z^7)\} \subset \mathrm{U}(2)$ the point $z = e^{2\pi i/7}$ has $\delta(g,I) < 1$ but $\log g \notin \mathfrak{g}$.




FIGURE 16. (a) A block $B(y)$ of side two and the next block in the direction μ, drawn in three of the four dimensions. Green dashed: the tree links, equal to I. Blue: the paths P_x of length two from the sites of $B(y)$, each joining $2y$ to $2y+2e_\mu$ through the trees; black: the straight path P_{2y}, the coarse link of the path map. The cut-off average is the mean of the sixteen paths in the chart at P_{2y} while every path is close to it. (b) For SU(2) and one link $e^{\theta X}$, $X=\operatorname{diag}(i,-i)$, the angle of $V^\sharp$ when the link lies on the straight path (black) or at another transverse position (blue), with $\rho_G=1$. Both equal $\theta/8$, the block average, while $\chi=1$; past the cut-off they are the path map, θ and 0, so the slice windings are 1 and 0.

(ii) *For every field U in the tree gauge,*

$$\sum_{y,\mu}\sum_{x\in B(y)}\delta(P_x,P_{2y})^2\le 896\,Q(U),$$

$$\sum_{y,\mu}\delta(V^\sharp_{y,\mu}(U),V^{\mathrm{p}}_{y,\mu}(U))^2\le 56c^2\,Q(U),$$

$$\sum_{e}\delta(U_e,R_e(V^\sharp(U)))^2\le(448+896c^2)\,Q(U),$$

$$Q_1(V^\sharp(U))\le(8+2688c^2)\,Q(U),$$

and $448+896c^2<1431$, $8+2688c^2<2956$.

(iii) *In the setting of Proposition 12.25 with $s=2$, W_t is the identity at the position of the straight path and 0 at the seven others.*

(iv) *If U equals I except on one link from $B(y)$ to $B(y+e_\mu)$, where it equals g with $\delta(g,I)\le\rho_G/2$, then $V^\sharp_{y,\mu}(U)=\exp(\frac18\log g)$, at each of the eight transverse positions of the link; the path map gives g at one position and I at the seven others.*

Proof. Write $P=P_{2y}$, $v_x=P^{-1}P_x$ and $m=\max_x\delta(v_x,I)=\max_x\delta(P_x,P)$. Where $m<\rho_G$, every $\log v_x$ lies in $\mathfrak{g}$, depends continuously on U by Lemma 12.26, which keeps the eigenvalues of v_x away from -1, and the exponent of (12.4) lies in $\mathfrak{g}$, so $V^\sharp\in G$. Where $m\ge\rho_G$, $V^\sharp=P$. As m increases to ρ_G the exponent has norm at most $\chi c\rho_G\to0$, so $V^\sharp$ is continuous. Let h be a gauge transformation equal to h_1 on $B(y)$ and to h_2 on $B(y+e_\mu)$. The middle point $x+e_\mu$ of each path lies in one of the two blocks, so $P_x\mapsto h_1P_xh_2^{-1}$ and $v_x\mapsto h_2v_xh_2^{-1}$; δ is invariant and log is equivariant under unitary conjugation, so χ is unchanged and $V^\sharp\mapsto h_1V^\sharp h_2^{-1}$. At $U=I$ every v_x is I, so $V^\sharp(I)=I$.

(i) The first statement is $\chi=1$. If $U=e^{t\varphi}$ with $\varphi=0$ on the tree, then $P=I+t\varphi_{2y+e_\mu,\mu}+O(t^2)$ and $\log v_x=t(\varphi_{x,\mu}+\varphi_{x+e_\mu,\mu}-\varphi_{2y+e_\mu,\mu})+O(t^2)$, and for small t the condition of the

first statement holds. The mean over the sixteen x is $t(V_{y,\mu}(\varphi) - \varphi_{2y+e_\mu,\mu}) + O(t^2)$, since $\varphi_{2y,\mu} = 0$, and multiplying by P gives the claim.

(ii) In the tree gauge $P = v_{y,\mu} := V^{\mathrm{p}}_{y,\mu}(U)$. If $o_\mu = 0$, then $U_{x+e_\mu,\mu}$ is the link leaving $B(y)$ at the transverse offset of x, and invariance of δ with the triangle inequality gives $\delta(P_x, P) \le \delta(U_{x,\mu}, I) + \delta(U_{x+e_\mu,\mu}, v_{y,\mu})$; if $o_\mu = 1$, the roles of the two links are exchanged. Each link leaving a block in the direction μ occurs for two sites x of that block, and each link in the direction μ with $o_\mu = 0$ occurs once for the block containing it and once for the block before it. With $(a+b)^2 \le 2a^2 + 2b^2$ and the notation of the proof of Theorem 12.21, the sum is at most $2(2I_0 + 2J_0) = 4D_v^2 \le 896Q$. Next, if $\chi > 0$, Lemma 12.26 and the Cauchy–Schwarz inequality give

$$\begin{aligned}\delta(V^\sharp, P) &\le \Big\| \frac{\chi}{16} \sum_x \log v_x \Big\| \le \frac{c}{16} \sum_x \delta(v_x, I) \\ &\le \frac{c}{4} \Big(\sum_x \delta(P_x, P)^2 \Big)^{1/2},\end{aligned}$$

and the second bound follows from the first, $896c^2/16 = 56c^2$. The reconstruction $R(V^\sharp)$ agrees with $R(v)$ on the links with $o_\mu = 0$, and on the eight links with $o_\mu = 1$ of each block and direction $\delta(U_e, V^\sharp)^2 \le 2\delta(U_e, v)^2 + 2\delta(V^\sharp, v)^2$; so the third sum is at most $2D_v^2 + 16 \cdot 56c^2 Q \le (448 + 896c^2)Q$. Finally, write $V^\sharp = vM$. Removing the four factors M from the holonomy of a coarse plaquette one at a time changes $\delta(\cdot, I)$ by at most $\delta(M, I)$ each time, so $\delta(V_P^\sharp, I)^2 \le 2\delta(v_P, I)^2 + 8 \sum_{\ell \in P} \delta(M_\ell, I)^2$. Each coarse link lies in six coarse plaquettes, so $Q_1(V^\sharp) \le 2Q_1(v) + 48 \cdot 56c^2 Q$, and $Q_1(v) \le 4Q$ by Proposition 12.22. The numerical bounds use $c^2 < 1.0967$.

(iii) For $s = 2$ the slice T consists of the eight links leaving $B(y)$, and the links inside the blocks are I, so $P_x = u_a$ for the transverse offset a of x, two sites for each a. Let t be a position other than the straight path and $u_{t'} = I$ for $t' \ne t$. Then $P = I$, two of the v_x equal u_t and the others I, so $F(u) = \exp(\frac{1}{8}\chi \log u_t)$ with $\chi = \phi(\delta(u_t, I)/\rho_G)$. On S the maps $u_t \mapsto \exp(\frac{\lambda}{8}\chi \log u_t)$, $0 \le \lambda \le 1$, are continuous by the argument above and take values in S, since $\log u_t$ commutes with S and the centraliser of a maximal torus in $\mathfrak{g}$ is its Lie algebra; they join F to the constant I, so $W_t = 0$. By Proposition 12.25 the W_t add up to the identity, which is therefore the matrix at the straight path.

(iv) At the straight path, $P_x = g$ for the two sites with transverse offset 0 and $P_x = I$ for the others, so $V^\sharp = g \exp(-\frac{14}{16} \log g) = \exp(\frac{1}{8} \log g)$; at another position, $P = I$ and two of the P_x equal g, so $V^\sharp = \exp(\frac{2}{16} \log g)$. □

The small-field part of (12.4) is Bałaban's averaging operation [7]; the cut-off extends it to all fields with the bounds of (ii). The theorem shows how far Proposition 12.25 reaches. The cut-off average is continuous, covariant and coercive for every field, and it agrees with the block averages to first order. Its slice windings are those of the path map, as the proposition demands of any such map, but the departure from the transverse symmetry happens only where some path differs from the straight one by more than $\rho_G/2$. Below that the field of a single link enters with the weight $\frac{1}{8}$ of the averages, whichever position it occupies.

What the theorem does not give is composition. The constants of (ii) are those of one step, and iterating them gives a factor that grows exponentially in the number of steps; on its large-field branch the map is the path map, for which Proposition 12.24 rules out any bound uniform in that number. Whether the k-fold cut-off average obeys $Q_k(V^{\sharp,k}(U)) \le C\, Q(U)$ and the corresponding coercivity bound with C independent of k is open for general fields; it is the form in which Remark 12.19(b) remains. Section 12.3 proves it, with $C = 1.001$, for fields with values in a maximal torus whose curvature stays small at every scale, and with $C = e^{12060\sigma}$ for fields of any compact group whose curvatures at the successive scales are small with sum σ.




FIGURE 17. (a) In the torus, the coarse plaquette of the cut-off average is the mean of the fluxes F_{2y+o} of the sixteen fine squares of side two at the offsets o (four of them drawn), whatever the gauge of the coarse links (Lemma 12.28). (b) For the thin-path fields of Proposition 12.24, the coarse energy after k steps: 24 for the path map, at most $18\pi^2/s^2$ for the cut-off average (Corollary 12.30); the fine energy is about $6\pi^2/s$.

12.3. **Composition across scales.** This subsection proves that the cut-off average composes across all scales on fields with values in a maximal torus whose curvature stays small, which includes the thin paths that defeat the path map, and then that it composes for every compact group on fields whose curvature is small, with a summable loss over the scales.

Let S be a maximal torus of $G \subset \mathrm{U}(r)$ and $\mathfrak{s}$ its Lie algebra. The exponential maps $\mathfrak{s}$ onto S, and its kernel Λ consists of diagonalisable matrices with eigenvalues in $2\pi i\mathbb{Z}$, so every nonzero element of Λ has norm at least 2π. For an S-valued field with $\delta(U_p, I) \leq \rho_G$ on every plaquette put $\Theta_p = \log U_p$; it lies in $\mathfrak{g}$, commutes with S, and so lies in $\mathfrak{s}$. For a fine site x and a coordinate plane $\{\mu, \nu\}$ let $\square_x$ be the square of side two at x in that plane and $F_x = \sum_{p \subset \square_x} \Theta_p$, the sum of its four plaquettes, oriented as the square. Put

$$\eta_G = \rho_G/12.$$

The k-fold cut-off average is defined by putting the coarse field back into the tree gauge, by an S-valued gauge transformation, before each step. The tree gauge fixes such a transformation up to one that is constant on blocks, under which $V^\sharp$ is covariant, so the coarse plaquette holonomies $U_P^{(j)}$ of the j-th iterate $U^{(j)}$ are well defined; we write $\Theta_P^{(j)} = \log U_P^{(j)}$ when $\delta(U_P^{(j)}, I) \leq \rho_G$.

Lemma 12.28. *Let U be an S-valued field in the tree gauge with $\|\Theta_p\| \leq \eta_G$ for every plaquette. Then $\chi_{y,\mu}(U) = 1$ for every coarse link, and the coarse plaquette P at y in the plane $\{\mu, \nu\}$ has*

$$V^\sharp(U)_P = \exp \bar{F}_P, \qquad \bar{F}_P = \frac{1}{16} \sum_{o \in \{0,1\}^4} F_{2y+o},$$

with $\|\bar{F}_P\| \leq 4 \max_p \|\Theta_p\|$, so that $\log V^\sharp(U)_P = \bar{F}_P$.

Proof. Let $\eta = \max_p \|\Theta_p\|$. Fix (y, μ) and $x = 2y + o$. The tree path τ from $2y$ to x sets the coordinates of o one at a time in increasing order, so it has at most four edges, at most three of them transverse to μ. In the tree gauge its links are I, so $P_{2y}^{-1} P_x$ is the holonomy of the loop τ, then the straight path from x to $x + 2e_\mu$, then $\tau + 2e_\mu$ backwards, then the straight path from $2y + 2e_\mu$ back to $2y$. This loop is the boundary of the surface swept by the transverse edges of τ along two steps in the direction μ: at most six distinct

plaquettes, each with coefficient ±1. The group is abelian, so $P_{2y}^{-1}P_x = \exp d_x$ with d_x the signed sum of their Θ_p, and $\|d_x\| \le 6\eta \le \rho_G/2$. By Lemma 12.26, $\delta(P_x, P_{2y}) \le \rho_G/2$, so $\chi = 1$, and since the eigenvalues of d_x have modulus at most $\frac12$, $\log(P_{2y}^{-1}P_x) = d_x$. Hence $V^{\sharp}_{y,\mu} = \exp(X_{y,\mu} + \bar d_{y,\mu})$, where $X_{y,\mu} \in \mathfrak s$ is any logarithm of P_{2y} and $\bar d_{y,\mu}$ is the mean of the d_x.

Let the sides of P be the coarse links ℓ with signs ϵ_ℓ, and write $d_{o,\ell}$ for the d_x of the side ℓ at the offset o. For each o the four straight paths of length two at the offset o of the four sides bound the square $\square_{2y+o}$, so

$$Y_o = \sum_\ell \epsilon_\ell (X_\ell + d_{o,\ell}) - F_{2y+o} \in \Lambda.$$

Since $d_{0,\ell} = 0$, $\|Y_o - Y_0\| \le 4\cdot 6\eta + 8\eta = 32\eta \le \frac83 < 2\pi$, so $Y_o = Y_0$ for every o. Averaging over o gives $\sum_\ell \epsilon_\ell(X_\ell + \bar d_\ell) = \bar F_P + Y_0$, and $\exp Y_0 = I$. Finally $\|\bar F_P\| \le \max_o \|F_{2y+o}\| \le 4\eta < \pi/3$, so $\bar F_P$ is the principal logarithm. □

Theorem 12.29. *Let U be an S-valued field in the tree gauge on the periodic lattice of side divisible by 2^k, with $\delta(U_p, I) \le \rho_G$ on every plaquette, and put $\eta_0 = \max_p \|\Theta_p\|$, $A = \sum_p \|\Theta_p\|$. Suppose that*

$$\max_P \|\Theta_P^{(j)}\| \le \eta_G \quad \text{for } 0 \le j < k, \tag{12.5}$$

which holds in particular if $\eta_0 A \le \eta_G^2$. Then the cut-off is inactive at every step, $\Theta^{(j+1)} = \bar F(\Theta^{(j)})$, and for $j < k$

$$\sum_P \|\Theta_P^{(j+1)}\|^2 \le \sum_P \|\Theta_P^{(j)}\|^2, \qquad \sum_P \|\Theta_P^{(j+1)}\| \le \frac14 \sum_P \|\Theta_P^{(j)}\|,$$
$$\max_P \|\Theta_P^{(j+1)}\| \le 4 \max_P \|\Theta_P^{(j)}\|.$$

Consequently

$$Q_k(V^{\sharp,k}(U)) \le \sum_p \|\Theta_p\|^2 \le 1.001\, Q(U),$$
$$Q_k(V^{\sharp,k}(U)) \le 16^{-k} A^2.$$

Proof. The first assertions follow from Lemma 12.28 by induction, the S-valued gauge fixing between steps leaving the plaquette holonomies unchanged. The coarse plaquettes P in a plane together with the offsets o are in bijection with the fine squares $\square_x$ in that plane, $x = 2y + o$, and a fine plaquette lies in four such squares. With $\|\bar F_P\|^2 \le \frac{1}{16}\sum_o \|F_{2y+o}\|^2$ and $\|F_x\|^2 \le 4\sum_{p\subset\square_x}\|\Theta_p\|^2$ this gives $\sum_P \|\bar F_P\|^2 \le \frac{1}{16}\cdot 4\cdot 4 \sum_p \|\Theta_p\|^2$; the same count without Cauchy–Schwarz gives the factor $\frac14$ in the second inequality, and the third is $\|F_x\| \le 4\eta$.

For the sufficient condition, let η_j and A_j be the maximum and the sum of $\|\Theta_P^{(j)}\|$. While (12.5) holds up to $j-1$, the inequalities give $\eta_j \le 4^j \eta_0$ and $\eta_j \le A_j \le 4^{-j} A$. The product of the two bounds is $\eta_0 A \le \eta_G^2$, so the smaller one is at most η_G, and (12.5) holds at j.

Finally $\delta(\exp\Theta, I) \le \|\Theta\|$ (Lemma 12.26), so $Q_k(V^{\sharp,k}(U)) \le \sum_P \|\Theta_P^{(k)}\|^2$, which is at most both $\sum_p \|\Theta_p\|^2$ and $\eta_k A_k \le A_k^2 \le 16^{-k} A^2$. The eigenvalues of each Θ_p are $i\theta$ with $|\theta| \le \eta_0 \le \eta_G \le \frac{1}{12}$, since $\eta_0^2 \le \eta_0 A$ in the sufficient condition and $\eta_0 \le \eta_G$ is the case $j = 0$ of (12.5); and $\theta^2 \le 1.001 \cdot 4\sin^2(\theta/2)$ there, which gives $\sum_p \|\Theta_p\|^2 \le 1.001\, Q(U)$. □

Corollary 12.30. *For $G = \mathrm{SU}(2)$ and the fields of Proposition 12.24 with $s = 2^k \ge 2^{14}$,*

$$Q_k(V^{\sharp,k}(U)) \le \frac{18\pi^2}{s^2} \le \frac{4}{s}\, Q(U),$$
$$Q_k(V^{\mathrm{p},k}(U)) = 24 \ge \frac{4s}{\pi^2}\, Q(U).$$

Proof. Here $\rho_G = 1$ and S is the diagonal torus. The $3s$ nonzero plaquettes have $\Theta_p = \pm 2T/s$, of norm $\sqrt{2}\pi/s$, so $\eta_0 A = 6\pi^2/s \le \frac{1}{144}$ when $s \ge 864\pi^2$, which holds for $s \ge 2^{14}$, and $A^2 = 18\pi^2$. Proposition 12.24 gives $Q(U) = 24s\sin^2(\pi/2s)$, which lies between $(6\pi^2/s)(1 - \pi^2/(12s^2))$ and $6\pi^2/s$. □

So on torus-valued fields of small curvature the cut-off average behaves like the linear averages: the curvature is averaged over the sixteen offsets at every step, the energy never increases, and a thin path is spread out and contracted instead of being carried to the coarse lattice (Figure 17).

For general fields the paths do not commute, and the logarithms carry commutators from the second order on. The next two lemmas bound these commutators by the square of the local curvature. The same composition then holds for every compact group, with a loss of at most $1 + O(\eta_j)$ at scale j.

Lemma 12.31. *Let $Z_1, \dots, Z_n \in \mathfrak{g}$ with $r = \sum_i \|Z_i\| \le \min\{\frac{1}{4}, \rho_G\}$, and let $g = e^{Z_1} \cdots e^{Z_n}$. Then $\delta(g, I) \le r$, $\log g \in \mathfrak{g}$, and*

$$\Big\| \log g - \sum_i Z_i \Big\| \le 1.6\, r^2.$$

Proof. The distance δ is invariant under multiplication by unitary matrices on either side, so $\delta(g, I) \le \sum_i \delta(e^{Z_i}, I) \le r$ by Lemma 12.26, and $\log g \in \mathfrak{g}$ by the choice of ρ_G. Expanding the exponentials, $g - I - \sum_i Z_i$ is a sum of products of at least two factors. Since the Hilbert–Schmidt norm is submultiplicative, its norm is at most $e^r - 1 - r \le \frac{1}{2} r^2 e^r \le 0.643\, r^2$, and $M = g - I$ has $\|M\| \le e^r - 1 \le 1.137\, r \le 0.285$. The operator norm of M is smaller than 1. The principal logarithm is holomorphic on the disc $|z - 1| < 1$, which contains the spectrum of g, so it is given there by its power series, $\log g = \sum_{k \ge 1} (-1)^{k+1} M^k / k$. Hence

$$\|\log g - M\| \le \frac{\|M\|^2}{2(1 - \|M\|)} \le 0.905\, r^2,$$

and the two bounds add up to less than $1.6\, r^2$. □

Let U be a G-valued field in the tree gauge with $\delta(U_p, I) \le \rho_G$, and put $\Theta_p = \log U_p$. A change of base point conjugates Θ_p and does not change its norm. Let P be a coarse plaquette at y in the plane $\{\mu, \nu\}$. Its sides are $\ell_1 = (y, \mu)$, $\ell_2 = (y + e_\mu, \nu)$, $\ell_3 = (y + e_\nu, \mu)$ and $\ell_4 = (y, \nu)$, with initial points y_i. For an offset $o \in \{0, 1\}^4$ let $P_{i,o}$ be the path product of ℓ_i through $2y_i + o$, and put

$$K_{P,o} = P_{1,o} P_{2,o} P_{3,o}^{-1} P_{4,o}^{-1}, \qquad \bar{F}_P = \frac{1}{16} \sum_o \log K_{P,o}.$$

The tree segments between consecutive sides cancel. Hence $K_{P,o}$ is the holonomy of the square $\square_{2y+o}$, carried to $2y$ along the tree. In a torus, $\log K_{P,o} = F_{2y+o}$, and $\bar{F}_P$ is the mean of Lemma 12.28. Let $\sigma_{P,o}$ be the sum of $\|\Theta_p\|$ over the four plaquettes of $\square_{2y+o}$.

Lemma 12.32. *Let U be a G-valued field in the tree gauge with $\delta(U_p, I) \le \rho_G$ and $\|\Theta_p\| \le \eta \le \rho_G/1000$ for every plaquette. Then $\chi_{y,\mu}(U) = 1$ for every coarse link. Every coarse plaquette has $\delta(V^\sharp(U)_P, I) \le 0.03\, \rho_G$ and*

$$\begin{aligned} \|\log V^\sharp(U)_P - \bar{F}_P\| &\le 3.2\, r_P^2, \\ \|\bar{F}_P\| &\le \frac{1 + 6.4\eta}{16} \sum_o \sigma_{P,o}, \end{aligned} \tag{12.6}$$

with $r_P \le 28.26\,\eta$ defined in the proof. Let E, A be the sum of $\|\Theta_p\|^2$ and of $\|\Theta_p\|$, and let E', A', η' be the sum of squares, the sum and the maximum of $\|\log V^\sharp(U)_P\|$. Then

$$\sqrt{E'} \le (1+6030\eta)\sqrt{E}, \qquad A' \le (1+9147\eta)\frac{A}{4},$$
$$\eta' \le 4(1+647\eta)\,\eta.$$

Proof. Paths. Fix a coarse link (y,μ) and $x = 2y + o$. As in the proof of Lemma 12.28, $P_{2y}^{-1}P_x$ is the holonomy of a loop bounding a surface of at most six plaquettes, swept by the transverse edges of the tree path. Build the surface one plaquette at a time. This writes the holonomy as a product of at most six factors $hU_p^{\pm1}h^{-1} = \exp(\pm h\Theta_p h^{-1})$. Let $\sigma_{(y,\mu),o} \le 6\eta$ be the sum of their $\|\Theta_p\|$. By Lemma 12.31, $\delta(P_x, P_{2y}) \le 6\eta \le \rho_G/2$, so $\chi = 1$. The logarithm $D_{(y,\mu),o} = \log(P_{2y}^{-1}P_x)$ lies in $\mathfrak{g}$, with

$$\|D_{(y,\mu),o}\| \le (1+9.6\eta)\,\sigma_{(y,\mu),o}.$$

In the same way $K_{P,o}$ is a product of four such factors, so $\|\log K_{P,o}\| \le (1+6.4\eta)\sigma_{P,o}$. Averaging over o gives the second inequality of (12.6).

The plaquette. Write $D_{i,o}$ for $D_{\ell_i,o}$. Then $P_{i,o} = P_{i,0}e^{D_{i,o}}$ and $D_{i,0} = 0$, and $V_{\ell_i}^\sharp = P_{i,0}e^{m_i}$ with $m_i = \frac{1}{16}\sum_o D_{i,o}$. The paths and $V^\sharp$ transform as links under gauge transformations constant on blocks. These transformations preserve the tree gauge and conjugate $D_{i,o}$, m_i, $K_{P,o}$ and $V_P^\sharp$ by fixed elements. Choose one with $P_{1,0} = P_{2,0} = P_{3,0} = I$. Then $P_{4,0} = K_{P,0}^{-1}$, and with $c = \log K_{P,0}$,

$$V_P^\sharp = e^{m_1}e^{m_2}e^{-m_3}e^{-m_4}e^{c},$$
$$K_{P,o} = e^{D_{1,o}}e^{D_{2,o}}e^{-D_{3,o}}e^{-D_{4,o}}e^{c}.$$

Put $r_P = \sum_i \max_o\|D_{i,o}\| + \|c\|$. Since $\|m_i\| \le \max_o\|D_{i,o}\|$, the five exponents in each product have norms adding up to at most r_P, and

$$r_P \le 24(1+9.6\eta)\eta + 4(1+6.4\eta)\eta \le 28.26\,\eta.$$

Lemma 12.31 gives $\delta(V_P^\sharp, I) \le r_P \le 0.03\,\rho_G$ and

$$\log V_P^\sharp = m_1 + m_2 - m_3 - m_4 + c + R,$$
$$\log K_{P,o} = D_{1,o} + D_{2,o} - D_{3,o} - D_{4,o} + c + R_o,$$

with $\|R\|, \|R_o\| \le 1.6\,r_P^2$. Average the second line over o and subtract it from the first; since m_i is the mean of the $D_{i,o}$, this is the first inequality of (12.6).

Sums. Each fine plaquette lies in four squares $\square_x$, and the pairs (P,o) are in bijection with these squares. The second inequality of (12.6) and Cauchy–Schwarz therefore give $\sum_P\|\bar F_P\|^2 \le (1+6.4\eta)^2E$; without Cauchy–Schwarz they give $\sum_P\|\bar F_P\| \le (1+6.4\eta)A/4$. For the remainders, let $\mathcal{U}_\ell$ be the set of plaquettes of the surfaces of the side $\ell = (y,\mu)$ for all sixteen offsets. These lie in planes $\{\mu,\alpha\}$, $\alpha \ne \mu$, at μ-positions $2y_\mu$, $2y_\mu+1$, $2y_\mu+2$, and there are at most 36 of them. A plaquette lies in $\mathcal{U}_\ell$ for at most four coarse links. With $m'_\ell = \sum_{p\in\mathcal{U}_\ell}\|\Theta_p\|$,

$$r_P \le 1.0096\Big(\sum_{\ell\subset P} m'_\ell + \sigma_{P,0}\Big).$$

A coarse link is a side of six coarse plaquettes, and a plaquette lies in $\square_{2y}$ for at most one coarse plaquette. Therefore

$$\sum_P r_P \le 1.0096\,(6\cdot 4A + A) \le 25.24\,A,$$
$$\sum_P r_P^2 \le 1.0096^2\cdot 5\,(6\cdot 36\cdot 4\,E + 4E) \le 4424\,E.$$

Hence, with $r_P \le 28.26\,\eta$,

$$\sum_P 3.2\, r_P^2 \le 3.2 \cdot 28.26 \cdot 25.24\,\eta A \le 2283\,\eta A,$$

$$\Big(\sum_P (3.2\, r_P^2)^2\Big)^{1/2} \le 3.2 \cdot 28.26\,\eta\,(4424\,E)^{1/2} \le 6015\,\eta\sqrt{E}.$$

The triangle inequality, for the Euclidean norm over P in the first case, gives $\sqrt{E'} \le (1 + 6.4\eta + 6015\eta)\sqrt{E}$ and $A' \le (1 + 6.4\eta)A/4 + 2283\,\eta A$. For the maximum, $\eta' \le (1 + 6.4\eta)4\eta + 3.2\,(28.26\,\eta)^2 \le 4(1 + 647\eta)\eta$. □

For a G-valued field, the k-fold cut-off average is defined as before, with G-valued gauge transformations. The tree gauge fixes them up to transformations constant on blocks, under which $V^\sharp$ is covariant. These transformations conjugate the plaquette holonomies and leave the norms $\|\Theta_P^{(j)}\|$ unchanged.

Theorem 12.33. *Let $G \subset \mathrm{U}(r)$ be compact and let U be a G-valued field on the periodic lattice of side divisible by 2^k, with $\delta(U_p, I) \le \rho_G$ on every plaquette. Let η_j, A_j, E_j be the maximum, the sum and the sum of squares of $\|\Theta_P^{(j)}\|$ over the plaquettes of the j-th iterate.*

(i) *If $\eta_j \le \rho_G/1000$ for $0 \le j < k$, the cut-off is inactive at every step. With $\sigma = \sum_{j<k} \eta_j$,*

$$E_k \le e^{12060\sigma} E_0, \qquad A_k \le e^{9147\sigma} 4^{-k} A_0,$$
$$\eta_k \le e^{647\sigma} 4^k \eta_0.$$

(ii) *If $\eta_0 A_0 \le 2 \cdot 10^{-10} \rho_G^2$, then (i) applies for every k with $\sigma \le 7.25\sqrt{\eta_0 A_0}$, and*

$$Q_k(V^{\sharp,k}(U)) \le e^{8.8\cdot 10^4 \sqrt{\eta_0 A_0}} E_0 \le 1.0001\, e^{8.8\cdot 10^4 \sqrt{\eta_0 A_0}}\, Q(U),$$
$$Q_k(V^{\sharp,k}(U)) \le e^2\, 16^{-k} A_0^2.$$

Proof. (i) Apply Lemma 12.32 at each step to the field in the tree gauge; the gauge fixing does not change the norms. The lemma gives $\delta(U_P^{(j+1)}, I) \le \rho_G$, so $\Theta^{(j+1)}$ is defined, and with $1 + t \le e^t$ the three inequalities of the lemma multiply over the steps.

(ii) Suppose that for $i < j$ we have $\eta_i \le \rho_G/1000$ and $\sum_{i<j} \eta_i \le 7.25\sqrt{\eta_0 A_0} \le 1/9747$. Then (i) gives $A_j \le e\, 4^{-j} A_0$ and $\eta_j \le e\, 4^j \eta_0$, and $\eta_j \le A_j$. So $\eta_j \le e \min\{4^j \eta_0, 4^{-j} A_0\} \le e\sqrt{\eta_0 A_0} \le \rho_G/1000$. Let i_* be the last index with $4^{i_*}\eta_0 \le 4^{-i_*} A_0$. The terms of $\min\{4^i \eta_0, 4^{-i} A_0\}$ increase by the factor 4 up to i_* and decrease by the factor 4 after it. The terms at i_* and $i_* + 1$ are at most $\sqrt{\eta_0 A_0}$, so the sum over all i is at most $\frac{8}{3}\sqrt{\eta_0 A_0}$, and $\sum_{i \le j} \eta_i \le \frac{8e}{3}\sqrt{\eta_0 A_0} \le 7.25\sqrt{\eta_0 A_0}$. This completes the induction. Then $E_k \le e^{12060 \cdot 7.25\sqrt{\eta_0 A_0}} E_0$. By Lemma 12.26, $Q_k(V^{\sharp,k}(U)) \le E_k$, and $E_k \le \eta_k A_k \le A_k^2 \le e^2\, 16^{-k} A_0^2$. Finally the eigenvalues $i\theta$ of each Θ_p have $|\theta| \le \eta_0 \le \sqrt{\eta_0 A_0} < 10^{-4}$, and $\theta^2 \le 1.0001 \cdot 4 \sin^2(\theta/2)$ there. □

For a torus Theorem 12.29 is sharper, and its hypothesis is weaker; the content of Theorem 12.33 is the non-abelian case. Neither theorem gives the supremum over boundary conditions in (D). Theorem 12.27 has the constants of one step. Theorems 12.29 and 12.33 need the curvature to be small at every scale: in Theorem 12.33 the losses $e^{6030\eta_j}$ must have a bounded product, and a field whose curvature is smooth on a scale ℓ grows by a factor four per step until the coarse spacing reaches ℓ. The large fields, where the cut-off is active, and the Wilson measure itself are outside both; Section 12.5 shows where the large fields lie under that measure.

12.4. **Small fields on the fibres of the cut-off average.** Theorem 12.4 is a gap on the linear fibres of the block averages. For group-valued links the fibres of the cut-off average are curved, and Remark 12.19(b) asks whether the gap survives on them. It does at every

background whose plaquettes are small, however far its links are from I, with a loss linear in the curvature.

Let $G \subset \mathrm{U}(r)$ be compact, work on the periodic lattice of side $2L$ in the tree gauge of Definition 12.2, and let W be a set of blocks. Write A_W for the active links with base in $\bigcup W$ and W^+ for the blocks at ℓ^∞ distance at most one from W. Fix U and let F be the set of fields that agree with U off A_W and have the same cut-off averages $V^\sharp_{y,\mu}$ as U for every $y \in W$ and every μ. The group G^{A_W} carries the bi-invariant metric with $|\xi|^2 = T(\xi) = \sum_e \|\xi_e\|^2$, $\xi_e \in \mathfrak{g}$, and its geodesics through U are the curves $U_e \exp(t\xi_e)$; write $\mathrm{Ad}(g)X = gXg^{-1}$. For a plaquette p at x in the plane $\{\mu,\nu\}$ with links $e_1 = (x,\mu)$, $e_2 = (x+e_\mu,\nu)$, $e_3 = (x+e_\nu,\mu)$, $e_4 = (x,\nu)$ and $U_p = U_1U_2U_3^{-1}U_4^{-1}$, the *covariant curvature*

$$D_p(\xi) = \mathrm{Ad}(U_1)\xi_1 + \mathrm{Ad}(U_1U_2)(\xi_2 - \xi_3) - \mathrm{Ad}(U_1U_2U_3^{-1})\xi_4$$

is the derivative of $U_p(t)U_p^{-1}$ along these curves. Reversing the orientation replaces it by $-\mathrm{Ad}(U_p^{-1})D_p(\xi)$, of the same norm, so $Q_U(\xi) = \sum_p \|D_p(\xi)\|^2$ does not depend on the orientations. At $U \equiv I$, $D_p(\xi) = (d\xi)_p$ and Q_I is the form Q of Theorem 12.4. For $y \in W$ and a direction μ let $v = U_{2y+e_\mu,\mu}$ be the path link, let $\tilde\xi$ equal $\mathrm{Ad}(v^{-1})\xi_e$ on the links of $B(y)$ and ξ_e on the other links, and let $N_{y,\mu}(\xi)^2$ be the sum of $\|\xi_e\|^2$ over the 24 links that enter $V_{y,\mu}$.

Lemma 12.34. *Suppose that $\delta(U_p, I) \le \kappa \le 10^{-8}\rho_G$ for every plaquette with a vertex in a block of W^+, and let $y \in W$.*

(i) *Every active link e inside a block of W^+ has $\delta(U_e, I) \le 3\kappa$, and every link e that leaves $B(y)$ in the direction μ has $\delta(U_e, v) \le 21\kappa$.*

(ii) *For a plaquette inside a block of W^+, $\|D_p(\xi) - (d\xi)_p\| \le 18\kappa \sum_{e\in p}\|\xi_e\|$.*

(iii) *Let $x \in B(y)$ with x_μ odd and x_ν even, and let p be the plaquette at x in the plane $\{\mu,\nu\}$, oriented as above, so that e_1, e_3 leave $B(y)$, e_2 lies in $B(y+e_\mu)$ and e_4 in $B(y)$. Then*

$$\begin{aligned}\big\|\mathrm{Ad}(v^{-1})D_p(\xi) - \big(\xi_1 + \xi_2 - \xi_3 - \mathrm{Ad}(v^{-1})\xi_4\big)\big\| \\ \le 48\kappa \sum_{e\in p}\|\xi_e\|.\end{aligned}$$

(iv) *Near U the cut-off average is*

$$V^\sharp_{y,\mu} = v\exp\Big(\frac{1}{16}\sum_x \log(v^{-1}P_x)\Big),$$

and its derivative satisfies

$$\begin{aligned}\Big\|V^\sharp_{y,\mu}(U)^{-1}\frac{d}{dt}V^\sharp_{y,\mu}(Ue^{t\xi})\Big|_{t=0} - V_{y,\mu}(\tilde\xi)\Big\| \\ \le 260\,\kappa\, N_{y,\mu}(\xi).\end{aligned}$$

Proof. We use $\|\mathrm{Ad}(g)X - X\| \le 2\delta(g,I)\|X\|$, which holds since $gXg^{-1} - X = ((g-I)X - X(g-I))g^{-1}$, and $\delta(gh,I) \le \delta(g,I) + \delta(h,I)$.

(i) The chains of plaquettes in the proof of Theorem 12.4 have their transverse links on the tree, so $\delta(U_e,I) \le \sum_{i\le k}\delta(U_{p_i},I) \le 3\kappa$. For the links leaving $B(y)$, the face plaquettes give $\delta(U_a,U_{a'}) \le \kappa + 3\kappa + 3\kappa$ for adjacent transverse offsets, as in the proof of Theorem 12.21, and every offset is joined to the straight path, which carries v, by at most three such steps.

(ii) The links of p and the products U_1U_2, $U_1U_2U_3^{-1}$ are within 3κ, 6κ and 9κ of I by (i).

(iii) By (i), $\delta(v^{-1}U_1, I) \le 21\kappa$ and $\delta(v^{-1}U_1U_2, I) \le 24\kappa$, and $v^{-1}U_1U_2U_3^{-1} = v^{-1}U_pU_4$ with $\delta(U_pU_4, I) \le 4\kappa$.

(iv) Write $P_x = U_{x,\mu}U_{x+e_\mu,\mu}$, $w_x = v^{-1}P_x$, $X_x = \log w_x$ and $M = \frac{1}{16}\sum_x X_x$; the path P_{2y} is v, its first link being on the tree. One of the two links of P_x leaves $B(y)$ and the other lies in $B(y)$ or $B(y+e_\mu)$, so $\delta(P_x, v) \le 24\kappa < \rho_G/2$ by (i); hence $\chi_{y,\mu} = 1$ near U, and $\|X_x\|, \|M\| \le 8\pi\kappa$ by Lemma 12.26. Since $\|[X,Y]\| \le 2\|X\|\,\|Y\|$, the operators

ad_{X_x} and ad_M have norm at most $a = 16\pi\kappa \le 50.3\kappa$. Along $Ue^{t\xi}$ put $\alpha = \xi_{2y+e_\mu,\mu}$ and $\beta_x = \mathrm{Ad}(U^{-1}_{x+e_\mu,\mu})\xi_{x,\mu} + \xi_{x+e_\mu,\mu}$, the right logarithmic derivatives of v and P_x. Then $w_x^{-1}\dot w_x = \beta_x - \mathrm{Ad}(w_x^{-1})\alpha$, and the formula for the derivative of the exponential map gives $\dot X_x = \Phi(\mathrm{ad}_{X_x})(\beta_x - \mathrm{Ad}(w_x^{-1})\alpha)$ and

$$(V^\sharp)^{-1}\dot V^\sharp = \mathrm{Ad}(e^{-M})\alpha + \Psi(\mathrm{ad}_M)\dot M,$$

with $\Phi(z) = z/(1-e^{-z})$ and $\Psi(z) = (1-e^{-z})/z$. The operators ad_X are skew-adjoint, and from the power series, $|\Phi(z)-1| \le |z|/2 + 0.09|z|^2 \le |z|$ and $|\Psi(z)-1| \le |z|$ for $|z| \le 1$, the Bernoulli numbers giving $\sum_{k\ge1}|B_{2k}|/(2k)! < 0.09$. By Lemma 12.3(iii), $V_{y,\mu}(\tilde\xi) = \frac{1}{16}\sum_x(\tilde\xi_{x,\mu} + \tilde\xi_{x+e_\mu,\mu}) = \alpha + \frac{1}{16}\sum_x(\beta_x - \alpha) - \frac{1}{16}\sum_x \epsilon_x$, where $\epsilon_x = \mathrm{Ad}(U^{-1}_{x+e_\mu,\mu})\xi_{x,\mu} - \tilde\xi_{x,\mu}$. If x_μ is even, $U_{x+e_\mu,\mu}$ leaves $B(y)$ and $\tilde\xi_{x,\mu} = \mathrm{Ad}(v^{-1})\xi_{x,\mu}$; otherwise $U_{x+e_\mu,\mu}$ lies in $B(y+e_\mu)$ and $\tilde\xi_{x,\mu} = \xi_{x,\mu}$; so $\|\epsilon_x\| \le 42\kappa\|\xi_{x,\mu}\|$ by (i). With m_β and m_ξ the means of $\|\beta_x\|$ and $\|\xi_{x,\mu}\|$, the four differences $\mathrm{Ad}(e^{-M})\alpha - \alpha$, $(\Psi(\mathrm{ad}_M)-1)\dot M$, $\dot M - \frac{1}{16}\sum_x(\beta_x - \alpha)$ and $\frac{1}{16}\sum_x\epsilon_x$ have norms at most $51\kappa\|\alpha\|$, $51.6\kappa(m_\beta + \|\alpha\|)$, $51\kappa m_\beta + 99\kappa\|\alpha\|$ and $42\kappa m_\xi$, using $\|\mathrm{Ad}(w_x^{-1}) - 1\| \le 48\kappa$. The links leaving $B(y)$ occur twice among the $\xi_{x,\mu}$, $\xi_{x+e_\mu,\mu}$ and the other sixteen links once, so the Cauchy–Schwarz inequality gives $m_\beta \le \frac{\sqrt{48}}{16}N_{y,\mu} \le 0.434N_{y,\mu}$ and $m_\xi \le \frac14 N_{y,\mu}$, and the sum is at most $(202 + 103\cdot0.434 + 42/4)\kappa N_{y,\mu} \le 260\kappa N_{y,\mu}$. □

Theorem 12.35. *Under the hypothesis of Lemma 12.34:*

(i) *for every $\xi \in \mathfrak{g}^{A_W}$ tangent to F at U, that is with $\frac{d}{dt}V^\sharp_{y,\mu}(Ue^{t\xi})|_{t=0} = 0$ for all $y \in W$ and all μ,*

$$T(\xi) \le 52(1 + 1.4\cdot10^6\kappa)\,Q_U(\xi);$$

(ii) *for every $\xi \in \mathfrak{g}^{A_W}$,*

$$\frac{d^2}{dt^2}Q(Ue^{t\xi})\Big|_{t=0} \ge 2Q_U(\xi) - 84\kappa\,T(\xi),$$

so on the tangent space of F the Hessian of Q is at least $(\frac{1}{26} - 5.4\cdot10^4\kappa)T(\xi)$; for SU(2) *the Hessian of the Wilson action $\beta\sum_p(1 - \frac12\,\mathrm{Re}\,\mathrm{Tr}\,U_p) = \frac{\beta}{4}Q$ on the linearised constrained directions is at least $\frac{\beta}{2}(\frac{1}{52} - 2.7\cdot10^4\kappa)T(\xi)$;*

(iii) *near U, F is a smooth submanifold of G^{A_W} of codimension $4|W|\dim G$, and the Hessian of $Q|_F$ for the induced metric satisfies $\mathrm{Hess}_F\,Q(\xi,\xi) \ge (\frac{1}{26} - c_G\kappa)T(\xi)$, where c_G depends only on G.*

Proof. (i) We follow the proof of Theorem 12.4, using $(a+b)^2 \le (1+\kappa)a^2 + (1+\kappa^{-1})b^2$ for each error term. *Inside a block.* For the chain $p_1,\dots,p_k$ of an active link $e = (x_0,\mu)$, $\xi_e = \sum_i \pm(d\xi)_{p_i}$ with tree links on the transverse sides, and Lemma 12.34(ii) gives

$$\|\xi_e\| \le \sum_i \|D_{p_i}\| + 36\kappa\sum_{j<k}\|\xi_{x_j,\mu}\|.$$

Hence $\|\xi_e\|^2$ is at most $(1+\kappa)k\sum_i\|D_{p_i}\|^2$ plus $3888\kappa(1+\kappa)\sum_{j<k}\|\xi_{x_j,\mu}\|^2$. A link (x',μ) is a point x_j of the chains of at most eight links, which differ from x' by a set of directions above μ, so

$$I \le 8(1+\kappa)Q_{\mathrm{in}} + 31104\kappa(1+\kappa)I.$$

Between blocks. Fix $y \in W$ and μ, and work in the frame of Lemma 12.34(iii),(iv); the norms do not depend on it. For a transverse edge $\{a,a'\}$, part (iii) gives $\|\xi_a - \xi_{a'}\| \le \|D_p\| + \|\xi_\psi\| + \|\xi_{\psi'}\| + 48\kappa(\|\xi_a\| + \|\xi_{a'}\| + \|\xi_\psi\| + \|\xi_{\psi'}\|)$, so the square is at most $3(1+\kappa)(\|D_p\|^2 + \|\xi_\psi\|^2 + \|\xi_{\psi'}\|^2)$ plus $9216\kappa(1+\kappa)$ times the sum of the four squares. Since ξ is tangent, part (iv) gives $\|V_{y,\mu}(\tilde\xi)\| \le 260\kappa N_{y,\mu}$, and by Lemma 12.3(iii) the mean m of the ξ_a is $V_{y,\mu}(\tilde\xi)$ minus a sixteenth of the sum of the sixteen framed links of S^μ_y and $S^\mu_{y+e_\mu}$;

so $8\|m\|^2 \le (1+\kappa)\frac12\sum_{16}\|\xi_e\|^2 + 540800\kappa(1+\kappa)N_{y,\mu}^2$. With the Laplacian of the cube as before,

$$\sum_a \|\xi_a\|^2 \le (1+\kappa)\Big(\frac12\sum_{16}\|\xi_e\|^2 + \frac32\sum_{\text{edges}}\Sigma_p\Big) + \kappa(1+\kappa)\Big(540800N_{y,\mu}^2 + 4608\big(3\sum_a\|\xi_a\|^2 + \sum_\Psi\|\xi_e\|^2\big)\Big),$$

where $\Sigma_p = \|D_p\|^2 + \|\xi_\psi\|^2 + \|\xi_{\psi'}\|^2$ and Ψ is the set of the 24 links ψ, ψ'. A link lies in at most two of the sets of $N_{y,\mu}$ and in at most three of the sets Ψ, so summing over $y \in W$ and μ as in the proof of Theorem 12.4, with $J = T - I$,

$$J \le (1+\kappa)\Big(\frac{11}{2}I + \frac32 Q_{\text{face}}\Big) + 1.1\cdot10^6\kappa(1+\kappa)T.$$

Adding I and inserting the bound for I gives $T \le 52(1+\kappa)^2Q_U(\xi) + 1.31\cdot10^6\kappa(1+\kappa)^2T$, and the claim follows for $\kappa \le 10^{-8}$.

(ii) Write $D_p(\xi) = B_1 + B_2 + B_3 + B_4$ with

$$B_1 = \mathrm{Ad}(U_1)\xi_1, \quad B_2 = \mathrm{Ad}(U_1U_2)\xi_2,$$
$$B_3 = -\mathrm{Ad}(U_1U_2)\xi_3, \quad B_4 = -\mathrm{Ad}(U_1U_2U_3^{-1})\xi_4.$$

Then $U_p(t) = e^{tB_1}e^{tB_2}e^{tB_3}e^{tB_4}U_p$, so $\ddot U_p = (D_p^2 + C_p)U_p$ with $D_p = \sum_i B_i$ and $C_p = \sum_{i<j}[B_i, B_j]$. Since $\delta(U_p, I)^2 = 2r - 2\,\mathrm{Re}\,\mathrm{Tr}\,U_p$, $\mathrm{Tr}\,C_p = 0$ and $-\mathrm{Tr}\,D_p^2 = \|D_p\|^2$,

$$\frac{d^2}{dt^2}\delta(U_p,I)^2 = 2\|D_p\|^2 - 2\,\mathrm{Re}\,\mathrm{Tr}(D_p^2(U_p - I)) - 2\,\mathrm{Re}\,\mathrm{Tr}(C_p(U_p - I)),$$

and $\|C_p\| \le \sum_{i<j}2\|B_i\|\|B_j\| \le 3\sum_{e\in p}\|\xi_e\|^2$. So the second derivative is at least $2(1-\kappa)\|D_p\|^2 - 6\kappa\sum_{e\in p}\|\xi_e\|^2$ when $\delta(U_p, I) \le \kappa$. Every plaquette that contains a link of A_W has a vertex in a block of W^+, each link lies in six plaquettes, and $Q_U(\xi) \le 24T(\xi)$; this gives the bound, and with (i) $\frac{2}{52(1+1.4\cdot10^6\kappa)} - 84\kappa \ge \frac{1}{26} - 5.4\cdot10^4\kappa$.

(iii) Let $C_{y,\mu}(U') = \log(V_{y,\mu}^\sharp(U)^{-1}V_{y,\mu}^\sharp(U'))$ near U, so that F is $C^{-1}(0)$ there, and let $\mathcal{G} = DC\,DC^*$ at U, a block matrix indexed by the pairs $k = (y,\mu)$. By Lemma 12.34(iv), $DC_k = \tilde B_k + R_k$, where $\tilde B_k\xi = V_{y,\mu}(\tilde\xi)$ and $\|R_k\| \le 260\kappa$ as maps from the 24 links. Each $\mathrm{Ad}(v^{-1})$ is an isometry, so $\tilde B_k\tilde B_k^*$ is the sum of the squared coefficients of $V_{y,\mu}$ on A_W, at least $\frac18$, $\|\tilde B_k\| \le \sqrt{3/16} \le 0.434$, and $\|\tilde B_k\tilde B_l^*\| \le \frac{1}{32}$ for the two neighbours $l = (y \pm e_\mu, \mu)$, the other blocks being zero. So the diagonal blocks of $\mathcal{G}$ are at least $\frac18 - 227\kappa$, the two off-diagonal blocks of a row have norm at most $\frac{1}{32} + 227\kappa$, and $\mathcal{G} \ge \frac{1}{16} - 700\kappa$. Hence DC is onto and F is a submanifold near U; and the Neumann series gives $\|\mathcal{G}^{-1}\| \le 17$ in the norm $\max_k\|z_k\|$. Let γ be a geodesic of F with $\gamma(0) = U$, $\dot\gamma(0) = \xi$, written as $Ue^{X(t)}$ link by link. The curves Ue^{tX} are the geodesics of G^{A_W}, so $\eta = \ddot X(0)$ is the second fundamental form at (ξ,ξ); it is normal to F, and differentiating $C(\gamma(t)) = 0$ twice gives $\eta = -DC^*\mathcal{G}^{-1}h$, where h_k is the second derivative of $C_k(Ue^{t\xi})$ at $t = 0$. Then

$$\mathrm{Hess}_F\,Q(\xi,\xi) = \frac{d^2}{dt^2}Q(Ue^{t\xi})\Big|_{t=0} + dQ(\eta),$$

and $dQ(\eta) = -\langle\mathcal{G}^{-1}DC\,\nabla Q, h\rangle$. As in (ii), $dQ(\xi) = -2\sum_p \mathrm{Re}\,\mathrm{Tr}(D_p(U_p - I))$, so $\|\nabla_e Q\| \le 12\kappa$, $\|(DC\nabla Q)_k\| \le (0.434 + 260\kappa)\,12\kappa\sqrt{24} \le 26\kappa$ and $\|(\mathcal{G}^{-1}DC\nabla Q)_k\| \le 442\kappa$. The map $C_k(Ue^X)$ depends only on the 24 links of $V_{y,\mu}$ and is real-analytic in (U,X) on a neighbourhood of the compact set where $\delta(P_x, P_{2y}) \le \rho_G/4$ for every x, so $\|h_k\| \le c_2N_{y,\mu}(\xi)^2$

with c_2 depending only on G. Summing, $|dQ(\eta)| \le 884 c_2 \kappa T(\xi)$, and (ii) gives the claim with $c_G = 5.4 \cdot 10^4 + 884 c_2$. □

So on the fibres of the cut-off average the action is uniformly convex along every geodesic of the fibre on which all plaquettes stay below κ, with modulus $\lambda(\frac{1}{26} - c_G\kappa)$, and the background enters only through its curvature. The constants are crude: on the periodic lattice of side four, for SU(2) fields equal to block gauge transforms of fields with κ up to 0.08 and W all sixteen blocks, the least value of Q_U/T on the tangent space stays within 0.4% of the gap 0.1075 of the quadratic approximation, and with κ up to 0.24 the frame errors of Lemma 12.34 stay below 1.3κ, a floating-point computation reproduced in the archive.

Remark 12.36. Theorem 12.35 is the curvature input of the small-field half of Remark 12.19(b); three steps remain open.

(a) With respect to the Riemannian volume of F, the constrained law has the density $e^{-\lambda Q}$ times the coarea factor $\det(DC\,DC^{*})^{-1/2}$, and its Bakry–Émery tensor also contains the Ricci curvature of F. These terms do not depend on λ, but they have not been estimated here.

(b) The set of points of F at which every plaquette is at most κ is not known to be geodesically convex in F, so the theorem gives neither a Brascamp–Lieb inequality on it nor a Helffer–Sjöstrand representation of the constrained influences, and the decay through Lemma 12.7 is not reached.

(c) The mass of the constrained law outside that set is bounded by Theorem 12.23 only for the fibres of the path map.

The hypothesis $\kappa \le 10^{-8}\rho_G$ concerns every plaquette near W; under the SU(2) Wilson measure the plaquettes that are not small form the clusters of Sec. 12.5, and near them the theorem does not apply.

The first two steps can be taken on a chart of F whose domain is convex. Write $T_U \subset \mathfrak{g}^{A_W}$ for the tangent space of F at U, N_U for its orthogonal complement, $\|X\|_\infty = \max_e \|X_e\|$ and $K_s = \{\eta \in T_U : \|\eta\|_\infty \le s\}$, a convex set. The *constrained law* is the conditional law of $e^{-\lambda Q}\prod_{e\in A_W} dU_e$, the other links being those of U, given the averages $V^{\sharp}_{y,\mu}$, $y \in W$, in the version given by the coarea formula.

Theorem 12.37. *Assume the hypothesis of Lemma 12.34. There are constants $c_3 \ge 1$ and c_6 depending only on G such that, if $0 < s \le 1/(6000c_3)$ and $\kappa + 6s \le 10^{-8}\rho_G$:*

(i) *there is a smooth map $h : K_s \to N_U$ with $\|h(\eta)\|_\infty \le s/2$ and $\Phi(\eta) = Ue^{\eta + h(\eta)} \in F$; Φ is an embedding, its image contains every $Ue^{X} \in F$ with $\|X\|_\infty \le s/40$, and every point of $\Phi(K_s)$ satisfies the hypothesis of Lemma 12.34 with $\kappa + 6s$ in place of κ;*

(ii) *for η in the interior of K_s and $\zeta \in T_U$,*

$$\frac{d^2}{dt^2} Q\big(\Phi(\eta + t\zeta)\big)\Big|_{t=0} \ge \Big(\frac{1}{26} - 4 \cdot 10^5 c_3(\kappa + s)\Big) T(\zeta);$$

(iii) *on $\Phi(K_s)$ the constrained law has, in the coordinate η, a density proportional to $e^{-\lambda Q(\Phi(\eta)) - \psi(\eta)}$ with respect to Lebesgue measure on T_U, where ψ is smooth, does not depend on λ, and $|\frac{d^2}{dt^2}\psi(\eta + t\zeta)| \le c_6 T(\zeta)$ on the interior of K_s;*

(iv) *if moreover $4 \cdot 10^5 c_3(\kappa + s) \le \frac{1}{195}$ and $\lambda \ge 60 c_6$, the constrained law conditioned on $\Phi(K_{s/2})$, written as a law μ on $K_{s/2}$, satisfies*

$$\operatorname{Var}_\mu(f) \le \frac{60}{\lambda} \int |\nabla f|^2 \, d\mu$$

for every smooth f on T_U.

The constants do not depend on L, W, U or λ.

For SU(2) and the Wilson action, $\lambda = \beta/4$, so (iv) is a Brascamp–Lieb inequality with constant $240/\beta$ on a chart of the fibre around every background with small plaquettes, for $\beta \geq 240c_6$.

Proof. Local constants. For $k = (y,\mu)$ let $c_k(X) = C_{y,\mu}(Ue^X)$, which depends only on the 24 links of $V_{y,\mu}$, and let $j(X_e) = \det \Psi(\mathrm{ad}_{X_e})$, the density of Haar measure in the coordinate $U_e e^{X_e}$, with Ψ as in Lemma 12.34. Both are real-analytic on a neighbourhood of the compact set of backgrounds with $\delta(P_x, P_{2y}) \leq 10^{-6}\rho_G$ for every x and of X with $\|X\|_\infty \leq 10^{-3}\rho_G$, and $j > 0$ there. So there is $c_3 \geq 1$ depending only on G, computed in Proposition 12.38, such that there the m-th derivative of c_k in X is at most $c_3 \prod_{i\leq m} N_{y,\mu}(Y^i)$ on $Y^1, \dots, Y^m$ for $m = 1,2,3$, and the first two derivatives of $\log j$ are at most c_3 in norm. By Lemma 12.34(i) the background U lies in this set.

Let $B = Dc(0)$, a map from $\mathfrak{g}^{A_W}$ to $\mathfrak{g}^K$, K the set of pairs k, so that $T_U = \ker B$ and $N_U = B^*\mathfrak{g}^K$, and $\mathcal{G} = BB^*$. By the proof of Theorem 12.35(iii), $\mathcal{G} \geq \frac{1}{17}$ and $\|\mathcal{G}^{-1}\| \leq 17$ in the norms $\max_k\|z_k\|$ and, $\mathcal{G}$ being symmetric, $\sum_k\|z_k\|$; each B_k has norm at most 0.44 on the 24 links of $V_{y,\mu}$, and every link lies in at most two of these sets. Hence $\|B\| \leq 0.63$, $\|BX\|_\infty \leq 2.2\|X\|_\infty$, $\|B^*z\|_\infty \leq 0.88\max_k\|z_k\|$ and $\|B^*z\|_1 \leq 10.6\sum_k\|z_k\|$, where $\|X\|_1 = \sum_e\|X_e\|$.

(i) Write $c(X) = BX + R(X)$. For $\|X\|_\infty, \|\hat X\|_\infty \leq 2s$, Taylor's formula gives $\|R_k(X)\| \leq \frac{c_3}{2}N_{y,\mu}(X)^2 \leq 12c_3\|X\|_\infty^2$ and $\|R_k(X) - R_k(\hat X)\| \leq 48c_3 s\|X - \hat X\|_\infty$. For $\eta \in K_s$ the equation $c(\eta + B^*z) = 0$ reads $z = \Theta(z) = -\mathcal{G}^{-1}R(\eta + B^*z)$, and on $\max_k\|z_k\| \leq s$ the map Θ has values of norm at most $17 \cdot 12c_3 \cdot 4s^2 = 816c_3s^2 \leq s$ and Lipschitz constant at most $17 \cdot 48 \cdot 0.88\, c_3 s \leq 720c_3 s \leq \frac12$. So it has a unique fixed point $z(\eta)$, smooth in η by the implicit function theorem, and $h = B^*z$ has $\|h\|_\infty \leq 0.88 \cdot 816c_3s^2 \leq s/2$. Since $h(\eta) \perp T_U$, η is the orthogonal projection of $\eta + h(\eta)$, so Φ is injective; its differential $1 + Dh$ is injective, and $\dim T_U = \dim F$. If $Ue^X \in F$ with $\|X\|_\infty \leq s/40$, then $X = \eta + B^*z$ with $z = \mathcal{G}^{-1}BX$, $\max_k\|z_k\| \leq 17 \cdot 2.2\, s/40 < s$ and $\|\eta\|_\infty \leq \|X\|_\infty + 0.88\max_k\|z_k\| < s$, and z is a fixed point of Θ, so $Ue^X = \Phi(\eta)$. Finally $\|\eta + h\|_\infty \leq \frac32 s$, and by bi-invariance $\delta(U'_p, U_p) \leq \sum_{e\in p}\delta(e^{X_e}, I) \leq 6s$ for $U' = Ue^X$; the links off A_W are unchanged, so U' is in the tree gauge and F is its fibre.

(ii) Let $X(t) = \eta + t\zeta + h(\eta + t\zeta)$, $\gamma(t) = Ue^{X(t)}$, and let $A = Dc(X)$. Differentiating $c(X(t)) = 0$ gives $X' = \zeta - B^*(AB^*)^{-1}A\zeta$ and $X'' = -B^*(AB^*)^{-1}D^2c(X)[X', X']$. As above $\|(A-B)Y\| \leq c_3N_{y,\mu}(X)N_{y,\mu}(Y) \leq 7.4c_3 s\, N_{y,\mu}(Y)$ in each component, so $\|A-B\| \leq 10.5c_3 s$ and $\sum_k\|((A-B)Y)_k\| \leq 14.8c_3 s\|Y\|_1$. Since $A\zeta = (A-B)\zeta$ and $AB^* = \mathcal{G} + (A-B)B^*$, the Neumann series gives $\|(AB^*)^{-1}\| \leq 34$ in the Euclidean norm and in the norm $\sum_k\|z_k\|$, the latter because $17 \cdot 14.8 \cdot 10.6\, c_3 s \leq \frac12$; hence $\|Dh\| \leq 0.63 \cdot 34 \cdot 10.5c_3 s \leq \frac16$, $T(X') \leq 1.03T(\zeta)$, the two parts of X' being orthogonal, and $\|X''\|_1 \leq 10.6 \cdot 34 \cdot 2c_3 \cdot 1.03T(\zeta) \leq 744c_3T(\zeta)$.

Let $w = \gamma^{-1}\gamma' = \Psi(\mathrm{ad}_X)X'$, link by link. Then $\gamma(t+\tau) = \gamma(t)\exp(\tau w + \frac{\tau^2}{2}\dot w + O(\tau^3))$, so

$$\frac{d^2}{dt^2}Q(\gamma) = \frac{d^2}{d\sigma^2}Q(\gamma e^{\sigma w})\Big|_{\sigma=0} + dQ(\dot w).$$

The vector w is tangent to F at $\gamma(t)$, which satisfies the hypothesis with $\kappa' = \kappa + 6s$, so by Theorem 12.35(i), (ii) at $\gamma(t)$ the first term is at least $(\frac{1}{26} - 5.4 \cdot 10^4\kappa')T(w)$. From the series of Ψ and $\|\mathrm{ad}_X\| \leq 2\|X\|$, $\|w_e - X'_e\| \leq 2\|X_e\|\,\|X'_e\| \leq 3s\|X'_e\|$, so $T(w) \geq (1 - 6s)T(\zeta)$; and $\dot w = D\Psi(\mathrm{ad}_X)[\mathrm{ad}_{X'}]X' + \Psi(\mathrm{ad}_X)X''$, where the term of first order in ad vanishes because $\mathrm{ad}_{X'}X' = 0$, so $\|\dot w_e\| \leq 3s\|X'_e\|^2 + (1+3s)\|X''_e\|$ and $\|\dot w\|_1 \leq 750c_3T(\zeta)$. With $\|\nabla_e Q\| \leq 12\kappa'$,

$$\frac{d^2}{dt^2}Q(\gamma) \geq \Big(\frac{1}{26} - \frac{s}{4} - (5.4 \cdot 10^4 + 9000c_3)\kappa'\Big)T(\zeta),$$

and $6(5.4 \cdot 10^4 + 9000c_3) + 1 \leq 4 \cdot 10^5 c_3$.

(iii) In the coordinate X, Haar measure is $\prod_e j(X_e)\,dX$, and the coarea formula for c writes dX as the integral over the values of c of the area measure of its level sets

divided by $\det(AA^*)^{1/2}$. On the level set F the graph $X = \eta + h(\eta)$ has area element $\det(1 + Dh^*Dh)^{1/2}\, d\eta$. So the density is $e^{-\lambda Q(\Phi)-\psi}$ with

$$\psi(\eta) = -\sum_e \log j(X_e) + \tfrac{1}{2}\log\det(AA^*) - \tfrac{1}{2}\log\det(1 + Dh^*Dh).$$

Along $\eta + t\zeta$, the first sum has second derivative at most $c_3(T(X') + \|X''\|_1) \le 745c_3^2T(\zeta)$. Next, $A' = D^2c(X)[X', \cdot]$ and $A'' = D^3c(X)[X', X', \cdot] + D^2c(X)[X'', \cdot]$; their components are maps of rank at most $\dim G$ on the 24 links of $V_{y,\mu}$, of norms at most $c_3N_{y,\mu}(X')$ and $c_3(N_{y,\mu}(X')^2 + N_{y,\mu}(X''))$. Hence the Hilbert–Schmidt norm satisfies $\|A'\|_2^2 \le 2.1\dim G\, c_3^2T(\zeta)$, and the trace norm $\|A''\|_1 \le \dim G\, c_3(2.1T(\zeta) + 2\|X''\|_1)$. Since $AA^* \ge (17^{-1/2} - 10.5c_3s)^2 \ge \frac{1}{25}$ and $\|A\| \le 1$,

$$\Big|\frac{d^2}{dt^2}\log\det(AA^*)\Big| \le 25(2\|A''\|_1 + 2\|A'\|_2^2) + 2500\|A'\|_2^2,$$

by the identity

$$\frac{d^2}{dt^2}\log\det M = \operatorname{Tr}(M^{-1}M'') - \operatorname{Tr}((M^{-1}M')^2)$$

and the bounds $|\operatorname{Tr}(XY)| \le \|X\|\,\|Y\|_1$ and $\|XY\|_1 \le \|X\|_2\|Y\|_2$. Finally $Dh = -\Pi$ on T_U with $\Pi = B^*(AB^*)^{-1}A$; Π' is a sum of two terms that contain A' once, and Π'' of terms that contain A'' once or A' twice, so $\|\Pi'\|_2 \le 481\|A'\|_2$ and $\|\Pi''\|_1 \le c(\|A''\|_1 + \|A'\|_2^2)$ with c absolute, and the same identity with $M = 1 + Dh^*Dh \ge 1$ and $\|Dh\| \le \frac{1}{6}$ bounds the last term. Each bound is a multiple of $T(\zeta)$ by a constant that depends only on c_3 and $\dim G$; this is c_6, computed in Proposition 12.38.

(iv) Let $H = \lambda Q\circ\Phi + \psi$ on the interior of K_s. By (ii) and (iii), $\operatorname{Hess} H \ge \lambda(\frac{1}{26} - \frac{1}{195}) - c_6 = \frac{\lambda}{30} - c_6 \ge m = \frac{\lambda}{60}$, so for $a, \eta \in K_{s/2}$, $H(\eta) \ge H(a) + \nabla H(a)\cdot(\eta - a) + \frac{m}{2}|\eta - a|^2$, the segment lying in the convex set $K_{s/2}$. The supremum $\tilde H$ over $a \in K_{s/2}$ of the right-hand sides is finite, equals H on $K_{s/2}$, and $\tilde H - \frac{m}{2}|\eta|^2$ is convex. With d the distance to $K_{s/2}$, a convex function, the functions $\tilde H + nd$ are m-convex on T_U, and so are their mollifications, which are smooth; for these the Bakry–Émery criterion [5] gives the Poincaré inequality with constant $1/m$. Letting the mollification and then $n \to \infty$, by dominated convergence ($\tilde H$ grows quadratically and $d = 0$ exactly on $K_{s/2}$), the inequality passes to $e^{-H}\mathbf{1}_{K_{s/2}}$, which is μ up to normalisation by (iii). □

The constants of Theorem 12.37 can be computed. Cauchy's estimates on a complex neighbourhood of the real chart bound the derivatives of the averages, and the rest of the proof turns these bounds into c_6.

Proposition 12.38. *Let $G \subset \mathrm{U}(r)$ be compact. Theorem 12.37 holds with*

$$c_3 = \max\{1.5\cdot 10^4\sqrt{r}, \dim G\}, \qquad c_6 = 6.81\cdot 10^5 \dim G\, c_3^2.$$

For SU(2) *these are $c_3 \le 2.13\cdot 10^4$ and $c_6 \le 9.3\cdot 10^{14}$, so part* (iv) *holds for $\lambda \ge 5.6\cdot 10^{16}$, that is, for the Wilson action at $\beta \ge 2.3\cdot 10^{17}$, once $\kappa + s \le 6.0\cdot 10^{-13}$.*

Proof. Write $\|\cdot\|_{\mathrm{op}}$ for the operator norm, which is at most the Hilbert–Schmidt norm, while $\|X\| \le \sqrt{r}\|X\|_{\mathrm{op}}$ on $\mathfrak{g}$. Since $\rho_G \le 1$, every background of the set of the proof has its sites $w_x = v^{-1}P_x$ within $a = 10^{-6}$ of I, and every X of that set has $\|X_e\| \le x_0 = 10^{-3}$.

Three estimates. Let X be skew-Hermitian and Z complex with $\|Z\|_{\mathrm{op}} \le t$. Duhamel's formula $e^{X+Z} - e^X = \int_0^1 e^{\sigma(X+Z)}Ze^{(1-\sigma)X}\,d\sigma$, with $\|e^{\sigma(X+Z)}\|_{\mathrm{op}} \le e^{\sigma t}$ by the Lie–Trotter product formula, gives $e^{X+Z} = e^X(I + F)$ and likewise $e^{-X-Z} = (I + F')e^{-X}$, with $\|F\|_{\mathrm{op}}, \|F'\|_{\mathrm{op}} \le \varphi = e^t - 1$. The same formula gives $\|e^A - e^B\| \le \|A - B\|$ for skew-Hermitian A, B, in both norms. Finally, on $\|w - I\|_{\mathrm{op}} < 1$ the series $\log w = \sum_{k\ge1}(-1)^{k+1}(w - I)^k/k$ is holomorphic, equals the principal logarithm at unitary w, and its derivative satisfies $\|D\log(w)[H]\| \le \|H\|/(1 - \|w - I\|_{\mathrm{op}})$ in both norms.

The complex chart. Fix U and a real X in the set, put $U' = Ue^X$, and let Z be complex with $\|Z_e\|_{\mathrm{op}} \le t$ on the 24 links of $V_{y,\mu}$. For a path $P_x = U_1U_2$ with path link v, the site at the complex field $U_e e^{X_e+Z_e}$ is

$$\begin{aligned} w_x &= e^{-X_v-Z_v} v^{-1} U_1 e^{X_1+Z_1} U_2 e^{X_2+Z_2} \\ &= (I+F_v')\, w_x'\, (I+\tilde F_1)(I+F_2), \end{aligned}$$

where w_x' is the site at U', a unitary matrix with $\delta(w_x', I) \le a + 3x_0 = e_0$ since each link moves by $\delta(e^{X_e}, I) \le x_0$, and $\tilde F_1$ is F_1 conjugated by the unitary $U_2 e^{X_2}$. Hence $\|w_x - w_x'\|_{\mathrm{op}} \le \psi = (1+\varphi)^3 - 1$, and if $e_0 + \psi < 1$ then $\|\log w_x - \log w_x'\|_{\mathrm{op}} \le d = \psi/(1-e_0-\psi)$, so the mean M of the sixteen logarithms is within d of its value M' at U', which is skew-Hermitian. The average is $V^\sharp = v e^{X_v+Z_v} e^M$, and with $\|e^M\|_{\mathrm{op}} \le e^d$ and $\|e^M - e^{M'}\|_{\mathrm{op}} \le e^d - 1$,

$$\begin{aligned} r = V^\sharp (U')^{-1} V^\sharp &= e^{-M'} (I + F_v) e^M, \\ \|r - I\|_{\mathrm{op}} &\le (1+\varphi) e^d - 1 = \eta. \end{aligned}$$

At real points $\chi = 1$, as $e_0 \rho_G < \rho_G/2$. The unitary $q' = V^\sharp(U)^{-1} V^\sharp(U') = e^{-M_0} e^{X_v} e^{M'}$ has $\delta(q', I) \le x_0 + \|M' - M_0\| \le x_0 + 3x_0/(1-e_0) = \gamma_0 < 0.0041$. So if $\gamma_0 + \eta < 1$, then $c(X+Z) = \log(q'r)$ is holomorphic in Z, extends c, and

$$\|c(X+Z) - c(X)\|_{\mathrm{op}} \le S(t) = \frac{\eta}{1 - \gamma_0 - \eta}.$$

Derivatives of c_k. For $N_{y,\mu}(Y) = 1$ the function $f(z) = c(X + zY)$ is holomorphic on $|z| < 0.1$, where $e_0 + \psi < 0.36$ and $\gamma_0 + \eta < 0.91$, and $\|zY_e\|_{\mathrm{op}} \le |z|$. Cauchy's estimate $\|f^{(m)}(0)\| \le m! R^{-m} \sup_{|z|=R} \|f(z) - f(0)\|$ gives

$$\|D^m c(X)[Y, \dots, Y]\| \le \sqrt{r}\, m!\, S(R)/R^m.$$

At $R = 10^{-3}$, 0.05 and 0.07 the values of $m! S(R)/R^m$ for $m = 1, 2, 3$ are below 4.2, 307 and 14811; at $R = 0.07$, for instance, $\varphi < 0.0726$, $\psi < 0.2337$, $d < 0.3062$, $\eta < 0.4567$ and $S < 0.847$. By Banach's theorem [9] a symmetric multilinear form on a real Hilbert space has the same norm as its restriction to the diagonal. Applied to $\langle u, D^m c(X)[\cdot]\rangle$ for unit vectors $u \in \mathfrak{g}$, with the Euclidean norm $N_{y,\mu}$ on the 24 links, this gives $\|D^m c(X)[Y^1, \dots, Y^m]\| \le 1.5 \cdot 10^4 \sqrt{r} \prod_i N_{y,\mu}(Y^i)$ for $m = 1, 2, 3$.

Derivatives of $\log j$. Since $\mathfrak{g}$ is reductive, $\operatorname{Tr} \operatorname{ad}_X = 0$, and $\Psi(z) = e^{-z/2} \sinh(z/2)/(z/2)$, so $\log j(X) = \operatorname{Tr} g(\operatorname{ad}_X)$ with $g(w) = \log(\sinh(w/2)/(w/2)) = \sum_{k \ge 1} g_{2k} w^{2k}$ for $|w| < 2\pi$. On $|w| = 3$, $|\sinh(w/2)/(w/2) - 1| \le \sinh(1.5)/1.5 - 1 < 0.4196$, so $|g| \le 0.5441$ and $|g_{2k}| \le 0.5441 \cdot 3^{-2k}$. For complex X with $\|X\|_{\mathrm{op}} \le b$ the operator ad_X on $\mathfrak{g} \otimes \mathbb{C}$ has norm at most $2b$, and $|\operatorname{Tr} A^{2k}| \le \dim G\, \|A\|_{\mathrm{op}}^{2k}$, so $|\log j(X)| \le 0.5441 \dim G\, u^2/(1-u^2)$ with $u = 2b/3$. Along $X + zH$, $\|H\| = 1$, $|z| \le 0.8$, this is at most $0.218 \dim G$, and Cauchy's estimate bounds the first two derivatives of $\log j$ by $0.28 \dim G$ and $0.69 \dim G$.

The constant c_6. In the proof of Theorem 12.37(iii), with c_3 as above, $a' = \|A'\|_2^2 \le 2.1 \dim G\, c_3^2 T(\zeta)$ and $b' = \|A''\|_1 \le \dim G\, c_3(2.1 + 1488 c_3) T(\zeta) \le 1490.1 \dim G\, c_3^2 T(\zeta)$. Write $K = (AB^*)^{-1}$, so that

$$\begin{aligned} K' &= -KA'B^*K, \quad K'' = 2KA'B^*KA'B^*K - KA''B^*K, \\ \Pi' &= B^*K'A + B^*KA', \quad \Pi'' = B^*K''A + 2B^*K'A' + B^*KA''. \end{aligned}$$

With $\|B\| \le 0.63$, $\|K\| \le 34$, $\|A\| \le 1$ and $k = 34 \cdot 0.63 = 21.42$ these give $\|\Pi'\|_2 \le (k + k^2)\|A'\|_2 \le 481 \|A'\|_2$ and $\|\Pi''\|_1 \le (2k^3 + 2k^2)\|A'\|_2^2 + (k^2 + k)\|A''\|_1 \le 20576\, a' + 481\, b'$. For $M = 1 + Dh^* Dh \ge 1$ with $\|Dh\| \le \frac{1}{6}$, the identity for $\log\det$ gives $|(\log\det M)''| \le$

$\frac{1}{3}\|\Pi''\|_1 + \frac{19}{9}\|\Pi'\|_2^2$. Adding the three terms of ψ,

$$\Big|\frac{d^2}{dt^2}\psi\Big| \le 745c_3^2T + 25b' + 1275a' \\ + \tfrac{1}{2}\Big(\tfrac{1}{3}(20576a' + 481b') + \tfrac{19}{9}\,481^2a'\Big),$$

which is at most $6.81 \cdot 10^5 \dim G\, c_3^2 T(\zeta)$.

For SU(2), $r = 2$ and $\dim G = 3$. Then $60c_6 < 5.6 \cdot 10^{16}$, and $4 \cdot 10^5 c_3(\kappa + s) \le \frac{1}{195}$ holds when $\kappa + s \le 6.0 \cdot 10^{-13}$; it implies $s \le 1/(6000c_3)$ and $\kappa + 6s \le 10^{-8}$. □

The bounds are far from the true sizes: on random SU(2) backgrounds the first three derivatives of c_k along unit directions stay below 0.14, 0.03 and 0.02, a floating-point computation reproduced in the archive. The loss comes from the radius 0.07 of the complex neighbourhood, which enters the third derivative as 0.07^{-3}.

Remark 12.39. Theorem 12.37 removes the coarea factor and the curvature of F for $\lambda \ge 60c_6$ and replaces the small-field set by the convex chart domain $K_{s/2}$. What it does not give:

(a) the mass of the constrained law outside $\Phi(K_{s/2})$, which is Remark 12.36(c);
(b) decay: (iv) is a variance bound for the law on the chart, not a decay of its covariances between distant links. That needs a Helffer–Sjöstrand representation on the convex set $K_{s/2}$, which has a boundary, and the locality of $\operatorname{Hess} H$, which through Dh contains $\mathcal{G}^{-1}$, exponentially decaying by Lemma 12.7 but not of finite range; Theorem 12.41 below reaches it on a chart whose domain is a product;
(c) numbers: the constants of Proposition 12.38 are far from optimal, so for SU(2) the threshold is $\lambda \ge 5.6 \cdot 10^{16}$ and κ and s are at the scale 10^{-13}.

On the periodic lattice of side four, for SU(2) backgrounds as in the check of Theorem 12.35, two to sixteen blocks and s up to 0.03, the chart Hessian of Q is at least 0.21 and $|\psi''|/T$ stays below one, a floating-point computation reproduced in the archive.

The obstacle in Remark 12.39(b) is the shape of $K_{s/2}$, which is not a product in any coordinates adapted to the links. A slightly different chart domain is. For $k = (y, \mu)$ let $d_k = (2y + e_\mu, \mu)$ be the path link of $V_{y,\mu}$. It has $o_\mu = 1$, so it is active and lies in A_W; it enters $V_{y,\mu}$ with weight $\frac{1}{8}$ by Lemma 12.3(iii); and it enters no other average, since the links with $o_\mu = 1$ that enter $V_{y',\mu}$ have their base in $B(y')$. Let $L' = A_W \setminus \{d_k\}$ and, for $r > 0$,

$$\Omega_r = \{\eta \in T_U : \|\eta_e\| \le r \text{ for every } e \in L'\}.$$

Give every link the position of its midpoint and every pair k that of d_k, and let $d(a, b)$ be the ℓ^∞ distance between the positions on the periodic lattice. The links that enter $V_{y,\mu}$ have midpoints within distance 1 of that of d_k. Let ∇' be the gradient in the coordinates $(\eta_e)_{e \in L'}$, and, for a block matrix $M = (M_{ab})$ whose indices are links or pairs and whose blocks are maps between copies of $\mathfrak{g}$, and for $\beta \ge 0$, let

$$\|M\|_\beta = \max\Big(\sup_a \sum_b e^{\beta d(a,b)}\|M_{ab}\|,\ \sup_b \sum_a e^{\beta d(a,b)}\|M_{ab}\|\Big).$$

At most two indices share a position, so the sum

$$N(\beta) = \sup_a \sum_b e^{-\beta d(a,b)}$$

is finite for $\beta > 0$ and depends only on β.

Lemma 12.40. (i) *The operator norm satisfies* $\|M\| \le \|M\|_\beta$, *and* $\|M_1M_2\|_\beta \le \|M_1\|_\beta\|M_2\|_\beta$ *and* $\|M^*\|_\beta = \|M\|_\beta$.
(ii) *If* $|\varphi(a) - \varphi(b)| \le d(a, b)$ *for all* a, b *and* $0 \le \alpha \le \beta$, *then* $\|e^{\alpha\varphi}Me^{-\alpha\varphi} - M\| \le \frac{\alpha}{\beta}\|M\|_\beta$.
(iii) *If* M *is invertible,* $\|M^{-1}\| \le M_0$ *and* $\|M\|_\beta \le M_1$, *then* $\|M^{-1}\|_\gamma \le 2M_0N(\gamma)$ *with* $\gamma = \beta/(4M_0M_1)$.

Proof. (i) is the Schur test and the triangle inequality for d. (ii) The blocks of the difference are $(e^{\alpha(\varphi(a)-\varphi(b))}-1)M_{ab}$, and $|e^{\alpha(\varphi(a)-\varphi(b))}-1| \le e^{\alpha d(a,b)}-1 \le \frac{\alpha}{\beta}e^{\beta d(a,b)}$ by convexity of $\alpha \mapsto e^{\alpha d}$; apply (i) with $\beta = 0$. (iii) Let $\alpha = 2\gamma \le \beta$, since $M_0 M_1 \ge \|M^{-1}\|\|M\| \ge 1$. By (ii), $e^{\alpha\varphi} M e^{-\alpha\varphi} = M + E$ with $\|M^{-1}E\| \le \frac12$, so $\|e^{\alpha\varphi}M^{-1}e^{-\alpha\varphi}\| \le 2M_0$ as in Lemma 12.7. With $\varphi = d(\cdot, b)$ the block (a,b) gives $e^{2\gamma d(a,b)}\|(M^{-1})_{ab}\| \le 2M_0$ for all a, b, and summing $e^{\gamma d(a,b)}\|(M^{-1})_{ab}\| \le 2M_0 e^{-\gamma d(a,b)}$ over b or over a gives the claim. □

Theorem 12.41. *Assume the hypotheses of Theorem 12.37*(iv)*, with $c_6 \ge 1$, as we may, and let $r = s/32$.*

(i) *The coordinates $(\eta_e)_{e\in L'}$ identify T_U with $\mathfrak{g}^{L'}$, Lebesgue measure on T_U being a constant multiple of that on $\mathfrak{g}^{L'}$; $\Omega_{r'}$ is a product of balls in them, and $K_{r'} \subset \Omega_{r'} \subset K_{15.1r'}$ for every $r' > 0$.*

(ii) *There are $\alpha_G > 0$ and c_9 depending only on G with the following property. Let μ be the constrained law conditioned on $\Phi(\Omega_r)$, written as a law on Ω_r. For every $\varphi : L' \to \mathbb{R}$ with $|\varphi(e) - \varphi(e')| \le d(e,e')$, every $0 \le \alpha \le \alpha_G$ and all smooth F, G on T_U,*

$$|\mathrm{Cov}_\mu(F,G)| \le \frac{120}{\lambda}\,\|e^{-\alpha\varphi}\nabla' F\|_{L^2(\mu)}\|e^{\alpha\varphi}\nabla' G\|_{L^2(\mu)}.$$

(iii) *If $X_1, X_2 \subset A_W$, $\tilde F, \tilde G$ are smooth functions of the links in X_1 and in X_2 whose left-invariant gradients have norm at most 1 in each link, $F = \tilde F \circ \Phi$ and $G = \tilde G \circ \Phi$, then*

$$|\mathrm{Cov}_\mu(F,G)| \le \frac{c_9}{\lambda}\,|X_1|\,|X_2|\,e^{-\alpha_G d(X_1,X_2)}.$$

The constants do not depend on L, W, U, λ, s or κ.

So the covariances of local observables of the field under the constrained law, conditioned on the chart, decay exponentially at a rate that depends only on G, with amplitude $O(1/\lambda)$; for SU(2) and the Wilson action the amplitude is $O(1/\beta)$.

Proof. All constants called C depend only on G.

The section. By Lemma 12.34(iv), $B_k = \tilde B_k + R_k$ with $\|R_k\| \le 260\kappa$, and the block of $\tilde B_k$ at d_k is $\frac18$ times an isometry of $\mathfrak{g}$, so $\|B_{k,d_k}^{-1}\| \le (\frac18 - 260\kappa)^{-1} \le 8.01$. For $\xi \in \mathfrak{g}^{L'}$ let $(S\xi)_e = \xi_e$ for $e \in L'$ and $(S\xi)_{d_k} = -B_{k,d_k}^{-1}\sum_{e\ne d_k} B_{k,e}\xi_e$. Since d_k enters only V_k, $BS = 0$; S is injective, and $\dim T_U = |L'|\dim G$ because F has codimension $4|W|\dim G$, so S is an isomorphism onto T_U whose inverse is the restriction to L'. The other coefficients of $\tilde B_k$ add up to $\frac78 + 1 = \frac{15}{8}$, so $\|(S\xi)_{d_k}\| \le 8.01(\frac{15}{8} + \sqrt{24}\cdot 260\kappa)\max_e\|\xi_e\| \le 15.1\max_e\|\xi_e\|$. This proves (i), and S has blocks of norm at most 1.1 that vanish beyond distance 1. Write $\mathrm{Box}_{r'}$ for the product of balls $S^{-1}\Omega_{r'}$. Since $15.1\cdot 2r < s$, the estimates of Theorem 12.37 hold on $S(\mathrm{Box}_{2r})$.

The density. Put $X(\xi) = S\xi + h(S\xi)$, $A = Dc(X)$, $P = 1 - B^*(AB^*)^{-1}A$ and $\bar P = PS$. In the proof of Theorem 12.37(ii) the derivative of X along $\zeta \in T_U$ is $P\zeta$, and $P\zeta = \zeta + Dh\,\zeta$ with $Dh\,\zeta \perp T_U$; so $\partial_\xi X = \bar P$, $\bar P^*\bar P = S^*(1 + Dh^*Dh)S \ge S^*S \ge 1$, and $\det \bar P^*\bar P = \det(S^*S)\det(1 + Dh^*Dh)$. By Theorem 12.37(iii) the law μ has, in the coordinate ξ, the density $e^{-\bar H}\mathbf{1}_{\mathrm{Box}_r}$ up to normalisation, with

$$\bar H(\xi) = \lambda Q(Ue^X) - \sum_e \log j(X_e) + \tfrac12\log\det(AA^*) - \tfrac12\log\det(\bar P^*\bar P),$$

and $\mathrm{Hess}\,\bar H \ge m = \frac{\lambda}{60}$ on Box_{2r}, because $|S\zeta| \ge |\zeta|$.

Locality of the Hessian. We show that $\|\mathrm{Hess}\,\bar H(\xi)\|_\beta \le c_8\lambda$ on Box_{2r} for some $\beta > 0$ and $c_8 \ge 1$ depending only on G, the Hessian being the block matrix of second derivatives in the coordinates ξ_e. The maps B, A, S, the second derivative of $Q(Ue^X)$ in X and that of $\sum_e \log j(X_e)$ have blocks bounded by C (the last two by compactness, as c_3) that vanish beyond distance 2, so their norms $\|\cdot\|_1$ are at most C. By the proof of Theorem 12.37,

$\|(AB^*)^{-1}\| \le 34$ and $\|(AA^*)^{-1}\| \le 25$, and $\|(\bar P^*\bar P)^{-1}\| \le 1$. Applying Lemma 12.40(iii) to AB^* and AA^*, then (i) to P, $\bar P$ and $\bar P^*\bar P$, then (iii) to $\bar P^*\bar P$, gives $\beta_0 \in (0,1]$ such that $(AB^*)^{-1}$, $(AA^*)^{-1}$, P, $\bar P$, $\bar P^*\bar P$ and $(\bar P^*\bar P)^{-1}$ have norm $\|\cdot\|_{\beta_0}$ at most C on Box_{2r}. Let $\gamma = \beta_0/2$, and for links e, e' and $v, v' \in \mathfrak{g}$ write v, v' also for the vectors of $\mathfrak{g}^{L'}$ supported at e, e'. Call a block matrix M, depending linearly on v, *concentrated at* e if $\|M_{ab}\| \le C|v|e^{-\gamma(d(a,e)+d(b,e))}$, and, depending bilinearly on (v,v'), *concentrated at* (e,e') if $\|M_{ab}\| \le C|v||v'|e^{-\gamma(d(a,e)+d(a,e')+d(b,e)+d(b,e'))}$; a vector is concentrated if it is so as a column. If $\|Y\|_{\beta_0} \le C$, then YM and MY are concentrated where M is, because $e^{-\gamma(d(c,e)+d(c,e'))} \le e^{\beta_0 d(a,c)}e^{-\gamma(d(a,e)+d(a,e'))}$. For M concentrated at e and M' at e', using $d(a,e)+d(a,e') \ge d(e,e')$ in both indices,

$$|\mathrm{Tr}(MM')| \le C|v||v'|N(\gamma/2)^2 e^{-\gamma d(e,e')},$$

and if M is concentrated at (e,e'), then

$$|\mathrm{Tr}\, M| \le C|v||v'|N(\gamma)e^{-\gamma d(e,e')}.$$

Now $x_v = \partial_v X = \bar P v$ has $\|(x_v)_a\| \le Ce^{-\beta_0 d(a,e)}|v|$, so it is concentrated at e. The vector $D^2c(X)[x_v, x_{v'}]$ has components at most $c_3 N_k(x_v)N_k(x_{v'}) \le C|v||v'|e^{-\beta_0(d(k,e)+d(k,e'))}$, so $x_{vv'} = \partial_v\partial_{v'}X = -B^*(AB^*)^{-1}D^2c(X)[x_v, x_{v'}]$ is concentrated at (e,e'). In the same way $A_v = \partial_v A = D^2c(X)[x_v, \cdot\,]$ is concentrated at e, $A_{vv'} = D^3c(X)[x_v, x_{v'}, \cdot\,] + D^2c(X)[x_{vv'}, \cdot\,]$ is concentrated at (e,e'), $\partial_v P = -B^*(AB^*)^{-1}A_v P$ is concentrated at e, and $\partial_v\partial_{v'}P$ is a sum of products of operators bounded in $\|\cdot\|_{\beta_0}$ with either one factor concentrated at (e,e') or one factor concentrated at e and one at e'. The same holds for the derivatives of $\mathcal{G}_1 = AA^*$ and $\mathcal{G}_2 = \bar P^*\bar P$. Then

$$\begin{aligned}\partial_v\partial_{v'}\bar H = {}& \lambda(D^2Q[x_v, x_{v'}] + DQ[x_{vv'}]) - \sum_e \partial_v\partial_{v'}\log j(X_e)\\ &+ \sum_{i=1,2} \pm\tfrac12(\mathrm{Tr}(\mathcal{G}_i^{-1}\partial_v\partial_{v'}\mathcal{G}_i)\\ &\qquad - \mathrm{Tr}(\mathcal{G}_i^{-1}\partial_v\mathcal{G}_i\,\mathcal{G}_i^{-1}\partial_{v'}\mathcal{G}_i)),\end{aligned}$$

where D^2Q and DQ are the derivatives of $Q(Ue^X)$ in X. By the bounds above every term is at most

$$C(\lambda+1)|v||v'|e^{-\gamma d(e,e')/2};$$

for instance

$$|D^2Q[x_v, x_{v'}]| \le C\sum_a e^{-\beta_0(d(a,e)+d(a,e')-2)}|v||v'|.$$

Hence, using $\lambda \ge 60c_6 \ge 60$, every block of the Hessian of $\bar H$ satisfies

$$\|(\mathrm{Hess}\,\bar H)_{ee'}\| \le C\lambda e^{-\beta_0 d(e,e')/4},$$

and $\|\mathrm{Hess}\,\bar H\|_\beta \le c_8\lambda$ with $\beta = \beta_0/8$.

Approximation. Let ρ be smooth on $[0,\infty)$ with $0 \le \rho' \le 1$, $\rho(t) = t$ for $t \le \frac32 r$, $\rho \le \frac74 r$ and ρ constant on $[2r,\infty)$, and let $\tau(\xi)_e = \rho(\|\xi_e\|)\xi_e/\|\xi_e\|$, which maps $\mathfrak{g}^{L'}$ into $\mathrm{Box}_{7r/4}$. Let v be smooth and convex on $[0,\infty)$, zero on $[0,r]$, with $0 < v'' \le 1$ on (r,∞) and $v'' = 1$ on $[\frac54 r,\infty)$, and $V(\xi) = \sum_e v(\|\xi_e\|)$. For $t \ge \frac32 r$, $v''(t) = 1$ and $v'(t)/t \ge 1 - \frac54 r/t \ge \frac16$, so $\mathrm{Hess}\,V \ge \frac16$ in the links with $\|\xi_e\| \ge \frac32 r$. Let $H_n = \bar H \circ \tau + nV$. Then

$$\mathrm{Hess}\,H_n = D\tau^*(\mathrm{Hess}\,\bar H \circ \tau)D\tau + \Delta_n,$$

where $D\tau$ and Δ_n are block diagonal, $D\tau$ is the identity in the links with $\|\xi_e\| \le \frac32 r$, $\|D\tau\| \le 1$, and $\Delta_n \ge n\frac16 - C_1$ in the other links, C_1 bounding $\nabla\bar H$ times the second derivative of τ. So $\mathrm{Hess}\,H_n \ge m$ on $\mathfrak{g}^{L'}$ for $n \ge 6(m + C_1)$, and all derivatives of H_n of

order at least two are bounded. For φ as in (ii) and $\alpha \le \alpha_G = \beta/(120c_8)$, the block-scalar matrix $e^{\alpha\varphi}$ commutes with $D\tau$ and Δ_n, so by Lemma 12.40(ii)

$$\big\| e^{\alpha\varphi} \operatorname{Hess} H_n e^{-\alpha\varphi} - \operatorname{Hess} H_n \big\| \le \frac{\alpha}{\beta} c_8 \lambda \le \frac{m}{2}.$$

Weighted Helffer–Sjöstrand. Let μ_n have density proportional to e^{-H_n} on $\mathfrak{g}^{L'}$ and $\mathcal{L}_n = -\Delta + \nabla H_n \cdot \nabla$ on $L^2(\mu_n)$. For H_n smooth with $\operatorname{Hess} H_n \ge m > 0$ and bounded derivatives of order at least two, and F, G smooth with bounded derivatives, the Helffer–Sjöstrand representation [47] reads $\operatorname{Cov}_{\mu_n}(F,G) = \langle \nabla F, \mathcal{A}^{-1} \nabla G \rangle_{L^2(\mu_n)}$ with $\mathcal{A} = \mathcal{L}_n \otimes 1 + \operatorname{Hess} H_n$ acting on $L^2(\mu_n; \mathfrak{g}^{L'})$. With $w = e^{\alpha\varphi}$, constant in ξ, $\mathcal{A}_w = w\mathcal{A}w^{-1} = \mathcal{L}_n \otimes 1 + w \operatorname{Hess} H_n w^{-1}$, and $\operatorname{Re}\langle u, \mathcal{A}_w u\rangle \ge \frac{m}{2}\|u\|^2$ because $\mathcal{L}_n \ge 0$; the adjoint satisfies the same bound, so $\|\mathcal{A}_w^{-1}\| \le \frac{2}{m} = \frac{120}{\lambda}$. Since $\mathcal{A}^{-1} = w^{-1}\mathcal{A}_w^{-1} w$,

$$\begin{aligned} |\operatorname{Cov}_{\mu_n}(F,G)| &= |\langle w^{-1}\nabla F, \mathcal{A}_w^{-1} w \nabla G\rangle| \\ &\le \frac{120}{\lambda} \|w^{-1}\nabla F\|_{L^2(\mu_n)} \|w \nabla G\|_{L^2(\mu_n)}. \end{aligned}$$

As $n \to \infty$, e^{-H_n} decreases to $e^{-\bar H}\mathbf{1}_{\mathrm{Box}_r}$ off the boundary of Box_r, since τ is the identity there and $V > 0$ off it, so $\mu_n \to \mu$ in total variation. For F, G smooth on T_U, apply the bound to $(F \circ S)\chi$ and $(G \circ S)\chi$ with χ smooth, compactly supported and equal to 1 near Box_r, and let $n \to \infty$; this proves (ii).

(iii) The left-invariant derivative of $U_a e^{X_a}$ along δX_a is $\Psi(\operatorname{ad}_{X_a})\delta X_a$, of norm at most $2\|\delta X_a\|$, so $\|\partial_{\xi_f} F\| \le 2\sum_{a \in X_1} \|\bar P_{af}\|$, and similarly for G. Take $\varphi = d(\cdot, X_2)$ and $\alpha = \alpha_G \le \beta_0$. Then $e^{-\alpha d(f,X_2)} \le e^{\beta_0 d(a,f)} e^{-\alpha d(a,X_2)}$, so $|e^{-\alpha\varphi}\nabla' F| \le 2|X_1| \|\bar P\|_{\beta_0} e^{-\alpha d(X_1,X_2)}$, and $e^{\alpha d(f,X_2)} \le e^{\beta_0 d(b,f)}$ for $b \in X_2$ gives $|e^{\alpha\varphi}\nabla' G| \le 2|X_2|\|\bar P\|_{\beta_0}$. Insert these in (ii); with C a bound for $\|\bar P\|_{\beta_0}$ on Box_{2r}, $c_9 = 480C^2$. □

The rate α_G and the amplitude c_9 can be computed, and they turn out not to depend on G. The proof above passes through the norms $\|\cdot\|_\beta$, whose Schur sums bring in the large factors $N(\beta)$; the computation below replaces them by weighted operator norms, in which inverses and products cost nothing beyond their operator norms. For a block matrix M indexed by links and pairs and for $\vartheta \ge 0$ let

$$\|M\|_{(\vartheta)} = \sup_{\varphi} \big\| e^{\vartheta\varphi} M e^{-\vartheta\varphi} \big\|,$$

the supremum over all φ with $|\varphi(a) - \varphi(b)| \le d(a,b)$ for all indices a, b, the factors $e^{\pm\vartheta\varphi}$ acting as scalars on each block. For such φ write $w = e^{\vartheta\varphi}$ and $M^w = wMw^{-1}$.

Lemma 12.42. (i) *For $\vartheta \le \vartheta'$, $\|M\| \le \|M\|_{(\vartheta)} \le \|M\|_{(\vartheta')}$; moreover $\|M_1 M_2\|_{(\vartheta)} \le \|M_1\|_{(\vartheta)} \|M_2\|_{(\vartheta)}$, $\|M^*\|_{(\vartheta)} = \|M\|_{(\vartheta)}$, and $\|M_{ab}\| \le \|M\|_{(\vartheta)} e^{-\vartheta d(a,b)}$. If the blocks of M vanish beyond distance R, then $\|M\|_{(\vartheta)} \le e^{\vartheta R}(R_M C_M)^{1/2}$, where R_M and C_M are the largest row and column sums of the norms of the blocks.*

(ii) *If M is Hermitian and $0 \le \vartheta \le \vartheta'$, then $\|\operatorname{Re} M^w - M\| \le 2(\vartheta/\vartheta')^2 \|M\|_{(\vartheta')}$ for every φ, where $\operatorname{Re} T = \frac{1}{2}(T + T^*)$.*

(iii) *If $\operatorname{Re} M^w \ge c > 0$ for every φ, then $\|M^{-1}\|_{(\vartheta)} \le 1/c$. If $\|M^{-1}\|_{(\vartheta)} \|E\|_{(\vartheta)} < 1$, then $\|(M+E)^{-1}\|_{(\vartheta)} \le \|M^{-1}\|_{(\vartheta)} / (1 - \|M^{-1}\|_{(\vartheta)}\|E\|_{(\vartheta)})$.*

Proof. (i) Take $\varphi = 0$ for the first inequality and replace φ by $\vartheta\varphi/\vartheta'$ for the second. Products and adjoints follow from $(M_1M_2)^w = M_1^w M_2^w$ and $(M^*)^w = (M^w)^*$ with $w' = e^{-\vartheta\varphi}$. With $\varphi = d(\cdot, b)$ the block (a,b) of M^w is $e^{\vartheta d(a,b)} M_{ab}$, of norm at most $\|M^w\|$. The last bound is the Schur test, since the blocks of M^w are $e^{\vartheta(\varphi(a) - \varphi(b))} M_{ab}$.

(ii) Let $f(z) = e^{z\varphi} M e^{-z\varphi}$, an entire function of $z \in \mathbb{C}$ with values in matrices. For $z = x + iy$, $f(z) = e^{iy\varphi} f(x) e^{-iy\varphi}$ with $e^{iy\varphi}$ unitary, so $\|f(z)\| \le K = \|M\|_{(\vartheta')}$ for $|x| \le \vartheta'$. Since M is Hermitian, $f(x)^* = f(-x)$ for real x, so $\operatorname{Re} M^w - M = g(\vartheta)$ with $g(z) = \frac{1}{2}(f(z) + f(-z)) - f(0)$. The function g is even, vanishes at 0 and is bounded by $2K$ on $|z| < \vartheta'$;

so $g(z)/z^2$ is holomorphic there, and the maximum principle on the circles $|z| = \rho < \vartheta'$, applied to $\ell(g(z))/z^2$ for every linear functional ℓ of norm one, gives $\|g(z)\| \le 2K|z|^2/\vartheta'^2$.

(iii) If $\operatorname{Re} T \ge c$, then $\|Tu\|\|u\| \ge \operatorname{Re}\langle u, Tu\rangle \ge c\|u\|^2$, so $\|T^{-1}\| \le 1/c$; apply this to $T = M^w$, whose inverse is $(M^{-1})^w$. The second statement is the Neumann series for $(M^w + E^w)^{-1}$. □

Proposition 12.43. *Let $G \subset \mathrm{U}(r)$ be compact. Theorem 12.41 holds with*

$$\alpha_G = 2.5 \cdot 10^{-4}, \qquad c_9 = 1.2 \cdot 10^{4},$$

and in (iii) *the factor $|X_1||X_2|$ may be replaced by $(|X_1||X_2|)^{1/2}$.*

So the covariances of local observables under the constrained law on the chart decay at least like $e^{-d/4000}$, with amplitude $1.2 \cdot 10^4/\lambda$, for every compact group; for SU(2) and the Wilson action the amplitude is $4.8 \cdot 10^4/\beta$.

Proof. Throughout, $\xi \in \mathrm{Box}_{2r}$, $X = S\xi + h(S\xi)$, and the notation is that of the proof of Theorem 12.41. The hypothesis $4 \cdot 10^5 c_3(\kappa + s) \le \frac{1}{195}$ of Theorem 12.37(iv) gives $c_3 s \le 1.29 \cdot 10^{-8}$ and $c_3\kappa' \le 7.7 \cdot 10^{-8}$ for $\kappa' = \kappa + 6s$, and $\kappa \le 10^{-8}$. Every φ below is defined on all indices; a function on L' with the Lipschitz property extends to all indices with the same property, by $\varphi(a) = \min_{e \in L'}(\varphi(e) + d(a, e))$.

Local operators. The row k of B has blocks of norm at most $\frac{1}{8} + 260\kappa$, with sum at most $2 + 24 \cdot 260\kappa$, by Lemma 12.34(iv) and the coefficients of $V_{y,\mu}$, eight of them $\frac{1}{8}$ and sixteen $\frac{1}{16}$; every link lies in at most two rows, and the blocks vanish beyond distance 1. So $b = \|B\|_{(\vartheta)} \le e^{\vartheta}(\frac{1}{2} + 10^{-4})^{1/2}$ by Lemma 12.42(i). The row k of $A - B$ has norm at most $c_3 N_{y,\mu}(X) \le 7.35 c_3 s = \varepsilon_A$, since $\|X\|_\infty \le \frac{3}{2}s$, so $\|A - B\|_{(\vartheta)} \le \sqrt{48}\, e^{\vartheta}\varepsilon_A$. The row d_k of S is $s_k\xi = -B^{-1}_{k,d_k} \sum_{e \ne d_k} B_{k,e}\xi_e$, of norm $\sigma \le (\frac{\sqrt{11}}{8} + 260\kappa)/(\frac{1}{8} - 260\kappa) \le 3.32$, the squares of the other coefficients adding up to $\frac{11}{64}$; the other rows of S are those of the identity. If D_k is the diagonal matrix with entries $e^{\vartheta(\varphi(d_k) - \varphi(e))}$ on the links e of $V_{y,\mu}$, then the row d_k of S^w is $s_k D_k$, and every link of L' lies in at most two of these sets, so

$$\bar{s} = \|S\|_{(\vartheta)} \le (1 + 2\sigma^2 e^{2\vartheta})^{1/2}.$$

Finally $Q(Ue^X) = \sum_p \|U_p - I\|^2 = \sum_p (2r - 2\operatorname{Re}\operatorname{Tr} U_p)$. By Duhamel's formula the derivative of $U_e e^{X_e}$ in X_e is an average of $U_e e^{\theta X_e} Y e^{(1-\theta)X_e}$, and its second derivative an average, of total weight one, of products with Y and Z inserted between unitary factors. With $|\operatorname{Tr}(V_1 Y V_2 Z V_3)| \le \|Y\|\|Z\|$ for unitary V_i this gives

$$|D^2 Q_p[Y, Z]| \le 2 \sum_{e \in p} \|Y_e\| \sum_{e \in p} \|Z_e\|,$$

and, since $\operatorname{Re}\operatorname{Tr}(VY) = \operatorname{Re}\operatorname{Tr}((V - I)Y)$ for skew-Hermitian Y and a conjugate V of U_p, $|DQ_p[Y]| \le 2\kappa' \sum_{e \in p} \|Y_e\|$. A link lies in six plaquettes, so $\|D^2 Q\|_{(\vartheta)} \le 48 e^{\vartheta}$ and $|DQ[Y]| \le 12\kappa' \|Y\|_1$.

Inverses. The diagonal blocks of $\mathcal{G} = BB^*$ have norm at most $\frac{3}{16} + 227\kappa$, each row has two other nonzero blocks, of norm at most $\frac{1}{32} + 227\kappa$, at distance 2, and $\mathcal{G} \ge \frac{1}{16} - 700\kappa$, by the proof of Theorem 12.35(iii). The blocks of $\operatorname{Re}\mathcal{G}^w$ are $\cosh(\vartheta(\varphi(k) - \varphi(l)))\mathcal{G}_{kl}$, so $\operatorname{Re}\mathcal{G}^w \ge \frac{1}{16} - 700\kappa - (\cosh 2\vartheta - 1)(\frac{1}{16} + 454\kappa)$, and Lemma 12.42(iii) bounds $g_0 = \|\mathcal{G}^{-1}\|_{(\vartheta)}$ by the inverse of the right-hand side. With $E = (A - B)B^*$, the same lemma gives

$$\|(AB^*)^{-1}\|_{(\vartheta)} \le \frac{g_0}{1 - g_0\|E\|_{(\vartheta)}} = h_0,$$

$$\|(AA^*)^{-1}\|_{(\vartheta)} \le \frac{g_0}{1 - g_0(2\|E\|_{(\vartheta)} + \|A - B\|^2_{(\vartheta)})} = g_1,$$

since $AB^* = \mathcal{G} + E$ and $AA^* = \mathcal{G} + E + E^* + (A - B)(A - B)^*$. Let $Y = B^*(AB^*)^{-1}$, so $y = \|Y\|_{(\vartheta)} \le b h_0$. Since $BS = 0$, $\bar{P} = PS = S - Y(A - B)S$, so $\bar{p} = \|\bar{P}\|_{(\vartheta)} \le$

$\bar s(1+y\|A-B\|_{(\vartheta)})$; and $M=Y(A-B)S$ takes values in N_U, orthogonal to the values of S, so $\bar P^*\bar P = S^*S + M^*M$. Here $S^*S = 1+\sum_k s_k^*s_k$, and $(s_k^*s_k)^w = v^*u$ with $u=s_kD_k$, $v=s_kD_k^{-1}$, so $\mathrm{Re}(s_k^*s_k)^w = \frac14((u+v)^*(u+v)-(u-v)^*(u-v)) \ge -\sigma^2\sinh^2\vartheta\, J_k$, J_k the projection on the links of $V_{y,\mu}$; and $\mathrm{Re}(M^*M)^w \ge -2(\vartheta/\vartheta'')^2\|M\|^2_{(\vartheta'')}$ by Lemma 12.42(ii), for $\vartheta\le\vartheta''$. Hence

$$g_2 = \|(\bar P^*\bar P)^{-1}\|_{(\vartheta)} \le \Big(1-2\sigma^2\sinh^2\vartheta - 2\frac{\vartheta^2}{\vartheta''^2}\|M\|^2_{(\vartheta'')}\Big)^{-1}.$$

For the ℓ^1 norms, $\|\mathcal{G}^{-1}\|\le 17$ in the norm $\sum_k\|z_k\|$ and E has column sums at most $2(2+10^{-4})\varepsilon_A$, so $(AB^*)^{-1}$ has norm $h_1\le 17.001$ there, and $\|B^*z\|_1\le(2+10^{-4})\sum_k\|z_k\|$.

The Hessian. Let $u,u'\in\mathfrak{g}^{L'}$, $p=wu$ and $q=w^{-1}u'$. Since $\langle u,(\mathrm{Hess}\,\bar H)^w u'\rangle = \partial_p\partial_q\bar H$, it suffices to bound $|\partial_p\partial_q\bar H|$ by $K|u||u'|$ for every φ. Write $x_p=\bar P p = w\tilde x$ and $x_q=w^{-1}\tilde y$, with $|\tilde x|\le\bar p|u|$ and $|\tilde y|\le\bar p|u'|$. The links of $V_{y,\mu}$ lie within distance 1 of the position of k, so the weights of x_p and x_q on them differ from $e^{\pm\vartheta\varphi(k)}$ by factors at most e^ϑ, and

$$\sum_k N_{y,\mu}(x_p)N_{y,\mu}(x_q)\le 2e^{2\vartheta}|\tilde x||\tilde y|,$$

every link lying in at most two of these sets. In the same way $A_p=\partial_pA=D^2c(X)[x_p,\cdot\,]$ equals $w\check A_p$ and $A_q=w^{-1}\check A_q$, where $\check A_p$ and $\check A_q$ have rows of rank at most $\dim G$, so their Hilbert–Schmidt norms satisfy

$$\|\check A_p\|_2\|\check A_q\|_2\le 2\dim G\, c_3^2e^{2\vartheta}\bar p^2|u||u'|,$$

and conjugating them by w multiplies these norms by at most e^ϑ. The second derivative $x_{pq}=-YD^2c(X)[x_p,x_q]$ has $\|x_{pq}\|_1\le c_3x_1|u||u'|$ with $x_1=2(2+10^{-4})h_1e^{2\vartheta}\bar p^2$, and $A_{pq}=D^3c(X)[x_p,x_q,\cdot\,]+D^2c(X)[x_{pq},\cdot\,]$ has trace norm $\bar a\le 2\dim G\, c_3^2(e^{2\vartheta}\bar p^2+x_1)|u||u'|$, the weights cancelling in both.

The four parts of $\bar H$ are bounded as follows. First, $\partial_p\partial_q\lambda Q=\lambda(\langle\tilde x,(D^2Q)^w\tilde y\rangle+DQ[x_{pq}])$, at most $\lambda(48e^\vartheta\bar p^2+12c_3\kappa' x_1)|u||u'|$. Second, the Hessian of $\sum_e\log j(X_e)$ in X is block diagonal and commutes with w, so its part is at most $c_3(\bar p^2+c_3x_1)|u||u'|$. Third, for $\mathcal{G}_1=AA^*$,

$$\begin{aligned}\partial_p\partial_q\log\det\mathcal{G}_1 = {}& \mathrm{Tr}(\mathcal{G}_1^{-1}\partial_p\partial_q\mathcal{G}_1)\\ &-\mathrm{Tr}(\mathcal{G}_1^{-1}\partial_p\mathcal{G}_1\,\mathcal{G}_1^{-1}\partial_q\mathcal{G}_1),\end{aligned}$$

with $\partial_p\mathcal{G}_1=A_pA^*+AA_p^*$ and $\partial_p\partial_q\mathcal{G}_1=A_{pq}A^*+A_pA_q^*+A_qA_p^*+AA_{pq}^*$. Each term is the trace of a product in which w and w^{-1} occur once each, next to $\check A_p$ and $\check A_q$, or not at all, next to A_{pq}. A cyclic permutation writes it as $\mathrm{Tr}(wFw^{-1}F')$, or $\mathrm{Tr}(w^{-1}Fw\,F')$, and $|\mathrm{Tr}(wFw^{-1}F')|\le\|wFw^{-1}\|_2\|F'\|_2$ or $\|wFw^{-1}\|\|F'\|_1$. With $a=\|A\|_{(\vartheta)}\le b+\|A-B\|_{(\vartheta)}$ this gives

$$\begin{aligned}\tfrac12|\partial_p\partial_q\log\det\mathcal{G}_1|\le{}& g_1a\bar a\\ &+(g_1+2g_1^2a^2)e^{2\vartheta}\|\check A_p\|_2\|\check A_q\|_2.\end{aligned}$$

Fourth, $\partial_p\bar P=-YA_p\bar P$ and $\partial_p\partial_q\bar P=-YA_{pq}\bar P+YA_qYA_p\bar P+YA_pYA_q\bar P$, and the same counting for $\mathcal{G}_2=\bar P^*\bar P$ gives

$$\begin{aligned}\tfrac12|\partial_p\partial_q\log\det\mathcal{G}_2|\le{}& g_2\bar p^2y\bar a+(3g_2\bar p^2y^2\\ &+2g_2^2\bar p^4y^2)e^{2\vartheta}\|\check A_p\|_2\|\check A_q\|_2.\end{aligned}$$

So

$$\|\mathrm{Hess}\,\bar H\|_{(\vartheta)}\le K=\lambda k_1+\dim G\, c_3^2k_0,$$

where $k_1=48e^\vartheta\bar p^2+12c_3\kappa' x_1$ and k_0 collects the other three parts, divided by $\dim G\, c_3^2|u||u'|$, which is possible as $c_3\ge1$. At $\vartheta=\vartheta'=0.197$, with $\vartheta''=0.6$, these give $b\le0.862$,

$g_0, h_0, g_1 \le 17.37$, $y \le 14.96$, $\bar{s}, \bar{p} \le 5.80$, $g_2 \le 7.41$, $k_1 \le 1966$ and $k_0 \le 4.16 \cdot 10^9$; and since $\lambda \ge 60 c_6 = 4.086 \cdot 10^7 \dim G\, c_3^2$ by Proposition 12.38, $K \le 2068\lambda$.

Decay. In the proof of Theorem 12.41, replace the bound on $e^{\alpha\varphi} \operatorname{Hess} H_n e^{-\alpha\varphi} - \operatorname{Hess} H_n$ by the following. Since $D\tau$ and Δ_n are block diagonal and symmetric and commute with $w = e^{\alpha\varphi}$, Lemma 12.42(ii) with $\vartheta = \alpha$ gives

$$\|\operatorname{Re}(\operatorname{Hess} H_n)^w - \operatorname{Hess} H_n\| \le 2\frac{\alpha^2}{\vartheta'^2} K \le \frac{\lambda}{120} = \frac{m}{2}$$

for $\alpha \le \vartheta'(240 \cdot 2068)^{-1/2}$, which holds for $\alpha \le 2.5 \cdot 10^{-4}$. Then $\operatorname{Re}\langle u, \mathcal{A}_w u\rangle \ge \frac{m}{2}\|u\|^2$ as before, and (ii) follows. For (iii), $\nabla' F = \bar{P}^* g$ with g supported on X_1 and $|g_a| \le 2$, so with $\varphi = d(\cdot, X_2)$,

$$|e^{-\alpha\varphi}\nabla' F| = |(\bar{P}^*)^{w^{-1}} e^{-\alpha\varphi} g| \le 2\bar{p}_\alpha |X_1|^{1/2} e^{-\alpha d(X_1, X_2)},$$

where $\bar{p}_\alpha = \|\bar{P}\|_{(\alpha)} \le 4.8$ at $\alpha = 2.5 \cdot 10^{-4}$, and $|e^{\alpha\varphi}\nabla' G| \le 2\bar{p}_\alpha |X_2|^{1/2}$. Insert these in (ii): $c_9 = 480\bar{p}_\alpha^2 \le 1.2 \cdot 10^4$. □

The chart misses only points of the fibre at which some plaquette near W is not small, and under the Wilson measure such points carry little mass on most fibres.

Theorem 12.44. (i) *Assume the hypotheses of Theorem 12.41 and $s \ge 1.24 \cdot 10^5 \kappa$. Every $U' \in F$ with $\delta(U'_p, I) \le \kappa$ for every plaquette p with a vertex in a block of W^+ lies in $\Phi(\Omega_r)$. Hence the constrained law gives the complement of $\Phi(\Omega_r)$ at most the mass of the points of F at which such a plaquette exceeds κ.*

(ii) *Let $G = \mathrm{SU}(2)$, let the side $2L$ be a power of two, let μ be the Wilson measure in the tree gauge at $\beta_W \ge 1$, so that $\lambda = \beta_W/4$, and let E be the set of fields with $\delta(U_p, I) \le \kappa$ for every plaquette with a vertex in a block of W^+. For μ-almost every U let μ_U be the constrained law on the fibre $F(U)$ through U. Then*

$$\mathbb{E}_\mu[\mu_U(E^c)] = \mu(E^c) \le 10^4 |W|\, q_\kappa(\beta_W),$$

with q_κ as in Theorem 12.57. If κ, s and λ satisfy the hypotheses of (i) *and $0 < \varepsilon < 1$, the fields $U \in E$ with $\mu_U(F(U) \setminus \Phi_U(\Omega_r)) \le \varepsilon$, where Φ_U is the chart at U, have probability at least $1 - (1 + \varepsilon^{-1})10^4 |W| q_\kappa(\beta_W)$.*

(iii) *At every such U, for $X_1, X_2 \subset A_W$ and $\tilde{F}, \tilde{G}$ as in Theorem 12.41*(iii),

$$|\operatorname{Cov}_{\mu_U}(\tilde{F}, \tilde{G})| \le \Big(\frac{4c_9}{\beta_W} e^{-\alpha_G d(X_1, X_2)} + 30\varepsilon\Big)|X_1|\,|X_2|.$$

Proof. (i) The fields U and U' satisfy the hypothesis of Lemma 12.34 and agree off A_W, in particular on the tree links. An active link e with base in a block $B(y)$, $y \in W$, either lies inside $B(y)$ or leaves it in a direction μ. In the first case part (i) of the lemma, at U and at U', gives $\delta(U_e, U'_e) \le 6\kappa$. In the second, let v and v' be the path links of $V_{y,\mu}$ at U and at U'. Part (iv), at both fields, writes the common cut-off average as $ve^M = v'e^{M'}$ with $\|M\|, \|M'\| \le 8\pi\kappa$, so by bi-invariance and Lemma 12.26

$$\delta(v, v') = \delta(e^M e^{-M'}, I) \le \|M\| + \|M'\| \le 16\pi\kappa < 50.3\kappa,$$

and part (i) gives $\delta(U_e, U'_e) \le \delta(U_e, v) + \delta(v, v') + \delta(v', U'_e) \le 92.3\kappa$. Let $X_e = \log(U_e^{-1} U'_e)$. By Lemma 12.26, $\|X_e\| \le \frac{\pi}{3} 92.3\kappa < 96.7\kappa \le s/1280$ for every $e \in A_W$. Theorem 12.37(i) with $s/32$ in place of s, whose chart is the restriction of Φ by the uniqueness of the fixed point in its proof, puts $U' = Ue^X$ in $\Phi(K_{s/32})$, and $K_{s/32} \subset \Omega_r$ by Theorem 12.41(i).

(ii) The tree links span every block, so for every field there is exactly one gauge transformation, equal to I at the corner of every block, that makes the tree links equal to I. Given the tree links, it multiplies each active link on both sides by elements that do not depend on that link, which preserves Haar measure, and S is gauge invariant; so in the tree gauge the active links have density proportional to $e^{-\beta_W S}$ with respect to product Haar measure. Since $\beta_W S = \frac{\beta_W}{4} \sum_p \delta(U_p, I)^2$, conditionally on the links off A_W the links of A_W have density proportional to $e^{-\lambda Q}$, and conditioning further on the averages gives the




FIGURE 18. Theorem 12.44. (a) A fibre F of the cut-off average, drawn as a surface: the product chart $\Phi(\Omega_r)$ around the background U (green) contains every point of F whose plaquettes near W are at most κ (inside the blue curve), and the rest of F meets only points with a large plaquette (red). (b) The least β_W at which the bound $10^4|W|q_\kappa(\beta_W)$ is at most $\frac{1}{2}$, for the 81 blocks of a cube of side three; it grows like $\kappa^{-2}\log(1/\kappa)$, and at the largest κ the hypotheses allow it is about $4.0 \cdot 10^{38}$.

constrained law. So μ_U is a version of the conditional law of μ given the links off A_W and the averages, and the identity is the tower property of conditional expectation. If $U \notin E$, some plaquette with a vertex in a block $B(y')$, $y' \in W^+$, is bad for $\rho = \kappa$. Its corner z differs from that vertex by an element of $\{0,1\}^4$, so z lies in a block at ℓ^∞ distance at most two from W, and the plaquette lies in the unit cube $\mathcal{C}_z$. There are at most $5^4|W|$ such blocks, with 16 sites each, and E is gauge invariant, so Theorem 12.57 with $|A| = 1$ gives $\mu(E^c) \leq 10^4|W|q_\kappa(\beta_W)$. For $U \in E$, part (i) gives $\mu_U(F(U) \setminus \Phi_U(\Omega_r)) \leq \mu_U(E^c)$, so the exceptional fields lie in $E^c \cup \{\mu_U(E^c) > \varepsilon\}$, which by Markov's inequality has probability at most $\mu(E^c)(1+\varepsilon^{-1})$.

(iii) Let $A = \Phi_U(\Omega_r)$, $m = \mu_U$, $m_A = m(\,\cdot \mid A)$ and $\epsilon' = m(A^c) \leq \varepsilon$. By Theorem 12.41(iii) with $\lambda = \beta_W/4$, $|\mathrm{Cov}_{m_A}(\tilde F, \tilde G)| \leq \frac{4c_9}{\beta_W}|X_1||X_2|e^{-\alpha_G d(X_1,X_2)}$. The group SU(2) has diameter $\sqrt{2}\pi$, and $\tilde F$ has gradient at most 1 in each link, so after subtracting a constant, which changes no covariance, $\|\tilde F\|_\infty \leq \frac{\sqrt{2}\pi}{2}|X_1| \leq 2.23|X_1|$, and likewise for $\tilde G$. Writing $m = (1-\epsilon')m_A + \epsilon' m_{A^c}$ gives $|m(fg) - m_A(fg)| \leq 2\epsilon'\|f\|_\infty\|g\|_\infty$ and $|m(f) - m_A(f)| \leq 2\epsilon'\|f\|_\infty$ for bounded f, g, so

$$|\mathrm{Cov}_m(f,g) - \mathrm{Cov}_{m_A}(f,g)| \leq 6\epsilon'\|f\|_\infty\|g\|_\infty,$$

and $6 \cdot 2.23^2 < 30$. □

So on a fixed set of blocks the constrained law of the SU(2) Wilson measure decorrelates, coarea factor and curvature of the fibre included, on every fibre outside a set of fibres whose probability tends to zero as $\beta_W \to \infty$. The bound grows linearly in $|W|$, which is harmless for the fixed cubes of the cube condition, but the thresholds are far from the one-loop range: with the constants of Proposition 12.38 the hypotheses force $\kappa + s \leq 6.0 \cdot 10^{-13}$ and $s \geq 1.24 \cdot 10^5\kappa$, so $\kappa < 4.9 \cdot 10^{-18}$, and $q_\kappa(\beta_W) < 1$ needs $\beta_W\kappa^2 > 96(\log\beta_W + C_0)$, so β_W exceeds $3.8 \cdot 10^{38}$ (Fig. 18).

The mass off the chart can also be bounded on a single fibre, by a Peierls argument inside the fibre. Two properties of the cut-off average make this possible: along a fixed set of links it is a submersion with a constant that does not depend on the field, and its law under Haar measure on those links has a bounded density.

Fix $y \in W$ and a direction μ. Write the offsets of the sites of $B(y)$ in the three directions other than μ as $a \in \{0,1\}^3$, and let x_a and x'_a be the sites of offset a with $o_\mu = 0$ and $o_\mu = 1$. Let $\ell_a = (x'_a, \mu)$, which joins $B(y)$ to $B(y + e_\mu)$, and let $u_a = (x_a, \mu)$ and $w_a = (x'_a + e_\mu, \mu)$, which lie inside $B(y)$ and $B(y + e_\mu)$. The links u_0 and w_0 are tree links and $\ell_0 = d_k$ is the path link, so in the tree gauge

$$P_{2y} = \ell_0, \qquad P_{x_a} = u_a \ell_a, \qquad P_{x'_a} = \ell_a w_a,$$

and $v_x = P_{2y}^{-1} P_x$ equals I at the two sites of offset 0. The path of $V^\sharp_{y',\nu}$ from a site x consists of the link (x, ν) and the next link $(x + e_\nu, \nu)$, so ℓ_a, whose base has $o_\mu = 1$, enters only the paths from x_a and x'_a of the pair (y, μ). We call $\ell_0, \dots, \ell_7$ the *private links* of (y, μ). The 32 links with $o_\mu = 1$ of $B(y)$ are the private links of the four pairs (y, μ); the other 17 active links of $B(y)$ lie inside it.

Let

$$\alpha = 2 \arcsin \frac{1}{2\sqrt{2}} = 0.7227\ldots, \qquad \Lambda = \frac{\alpha}{\sin \alpha} = 1.0926\ldots,$$

so that $1 - \frac{7}{8}\Lambda > 0.0439$. Two facts about $\operatorname{ad} X$ for $X \in \mathfrak{g}$, which is skew-adjoint for the Hilbert–Schmidt product, will be used throughout. If the eigenvalues $i\psi$ of $\operatorname{ad} X$ satisfy $|\psi| < 2\alpha$, then

$$\log(e^{tZ} e^X) = X + t L_X Z + O(t^2),$$
$$L_X = \frac{\operatorname{ad} X}{e^{\operatorname{ad} X} - 1}, \qquad \|L_X\| \le \Lambda,$$

since the eigenvalues of L_X have modulus $\frac{|\psi|/2}{\sin(|\psi|/2)}$, which increases with $|\psi|$. For every $Y \in \mathfrak{g}$,

$$\frac{d}{dt} e^{Y+tZ}\Big|_{t=0} e^{-Y} = E_Y Z,$$
$$E_Y = \frac{e^{\operatorname{ad} Y} - 1}{\operatorname{ad} Y}, \qquad \|E_Y\| \le 1,$$

because $|e^{i\psi} - 1| \le |\psi|$; in particular $d(e^A, e^B) \le \|A - B\|$ for the Riemannian distance d of the bi-invariant metric, along the curve $e^{A + t(B-A)}$. If $\delta(v, I) < 1$ and $e^{i\theta_j}$, $|\theta_j| \le \pi$, are the eigenvalues of v, then $\sum_j 4 \sin^2(\theta_j/2) < 1$; each $|\theta_j|$ is less than $\frac{\pi}{3}$, where $\sin^2(t/2)$ is convex, so $|\theta_j| + |\theta_k| < 2\alpha$ for all j, k, and $X = \log v$ satisfies the hypothesis on $\operatorname{ad} X$.

Lemma 12.45. *Let $\sigma_0 = (1 - \frac{7}{8}\Lambda)/\sqrt{2} > 0.031$. At every field at which $V^\sharp_{y,\mu}$ is differentiable, its derivative $J : \mathfrak{g}^8 \to \mathfrak{g}$ with respect to its eight private links, right trivialised, satisfies $JJ^* \ge \sigma_0^2$. The fields at which it is not differentiable form a null set for Haar measure.*

Proof. Write $m = \max_x \delta(v_x, I)$ and $\chi = \phi(m/\rho_G)$, so that $V^\sharp = P_{2y} e^{\chi N}$ with $N = \frac{1}{16} \sum_x \log v_x$. Where $m > \rho_G$, the cut-off vanishes near the field, $V^\sharp = \ell_0$, and the derivative along ℓ_0 is the identity. Suppose $m < \rho_G$, and that the maximum is attained at one path x_*, after identifying identical path functions, and $m \ne \rho_G/2$. Copies that coincide because u_a, w_a are tree links are counted together. Other ties lie in finitely many proper real-analytic subsets, a null set. The variation below fixes both copies of an identical pair, since then $u_b = w_b = I$ and $A - 1 = 0$. Then every $\delta(v_x, I)$ is less than 1, the logarithms are defined, χ is differentiable, and every variation that fixes v_{x_*} to first order fixes χ to first order. The site x_* has offset $b \ne 0$.

Fix $\zeta \in \mathfrak{g}$ and put $\eta = \operatorname{Ad}(\ell_0)\zeta$. Move $\ell_0 \mapsto e^{t\eta}\ell_0$, and, if $x_* = x'_b$, $\ell_b \mapsto e^{t\eta}\ell_b$, or, if $x_* = x_b$, $\ell_b \mapsto e^{t\xi}\ell_b$ with $\xi = \operatorname{Ad}(u_b^{-1})\eta$, so that $u_b \ell_b \mapsto e^{t\eta} u_b \ell_b$. In both cases P_{2y} and P_{x_*} are multiplied on the left by $e^{t\eta}$, so v_{x_*} and χ are fixed to first order. The other site $\bar{x}$ of offset b has $P_{\bar{x}} \mapsto e^{t \operatorname{Ad}(u_b^{\pm 1})\eta} P_{\bar{x}}$, and the paths from the remaining twelve sites with $v_x \ne I$ do not move. With the isometry $A = \operatorname{Ad}(\ell_0^{-1} u_b^{\pm 1} \ell_0)$,

$$v_{\bar{x}} \mapsto e^{t(A-1)\zeta} v_{\bar{x}} + O(t^2), \qquad v_x \mapsto e^{-t\zeta} v_x + O(t^2)$$

for the twelve others, so N moves by $t\,\delta N$ with $16\,\delta N = L_{X_{\bar{x}}}(A-1)\zeta - \sum_x L_{X_x}\zeta$, twelve terms, and $\|\delta N\| \le \frac{14}{16}\Lambda\|\zeta\|$. Since χ is fixed,

$$\frac{d}{dt}V^\sharp\,(V^\sharp)^{-1} = \eta + \mathrm{Ad}(\ell_0)E_{\chi N}(\chi\,\delta N) = \mathrm{Ad}(\ell_0)\,T\zeta,$$

with $\|T\zeta - \zeta\| \le \frac{7}{8}\Lambda\|\zeta\|$, as $\chi \le 1$. The variation is $K\zeta$ for a linear map $K : \mathfrak{g} \to \mathfrak{g}^8$ with two nonzero components of norm $\|\zeta\|$, so $\|K\| = \sqrt{2}$ and $JK = \mathrm{Ad}(\ell_0)T$. For a unit vector $e \in \mathfrak{g}$,

$$\|J^*e\| \ge \frac{\|K^*J^*e\|}{\sqrt{2}} = \frac{\|T^*\,\mathrm{Ad}(\ell_0)^*e\|}{\sqrt{2}} \ge \frac{1-\frac{7}{8}\Lambda}{\sqrt{2}} = \sigma_0,$$

since the least singular value of T is at least $1-\frac{7}{8}\Lambda$. At the other points of differentiability the same bound follows by continuity: the cut-off has finitely many smooth branch formulas, at such a point the formulas of the adjacent full-dimensional branches agree with its differential, and the lower bound on the singular value is preserved under limits. □

Corollary 12.46. *Let $C = (V^\sharp_{y,\mu})_{y\in W,\,\mu}$, as a function of the links of A_W, the other links being fixed. Wherever C is differentiable,*

$$\det(DC\,DC^*)^{-1/2} \le \sigma_0^{-4|W|\dim G} < 33^{4|W|\dim G}.$$

Proof. The private links of distinct pairs are distinct, and each enters only the average of its pair, so the columns of DC that belong to private links form a block diagonal matrix with the blocks of Lemma 12.45. Hence $DC\,DC^* \ge \bigoplus_k J_kJ_k^* \ge \sigma_0^2$. □

The coarea factor of the constrained law is therefore bounded on the whole fibre by a constant per block, for every value of the cut-off and however far the links inside the blocks are from I. The volume of the fibre is bounded through the law of the averages. From here on $G = \mathrm{SU}(2)$, which with the metric $\|X\|^2 = \mathrm{Tr}(X^*X)$ is the round sphere of radius $\sqrt{2}$, $\rho_G = 1$, and h is normalised Haar measure.

Lemma 12.47. *Let $G = \mathrm{SU}(2)$, fix every link other than the private links of (y,μ), and let these eight links be independent and Haar distributed. Then the law of $V^\sharp_{y,\mu}$ has a density with respect to h that is at most*

$$M = 1 + 2\Big(1 - \frac{7}{8}\Lambda\Big)^{-3} < 2.4\cdot 10^4$$

at every point, in the version given by the formula (12.7) of the proof.

Proof. Let $g = \ell_0$, $m_1 = \max_a \delta(v_{x_a}, I)$ and $m_2 = \max_a \delta(v_{x'_a}, I)$. Split G^8, up to the null set $m = 1$, into $R_0 = \{m > 1\}$, $R_1 = \{m_1 \ge m_2,\ m_1 < 1\}$ and $R_2 = \{m_2 > m_1,\ m_2 < 1\}$; ties between the two families, including two path functions that coincide identically, belong to R_1. On R_0, $\chi = 0$ and $V^\sharp = g$.

On R_1 use the coordinates $(g, h_1, \dots, h_7)$ with $h_a = v_{x_a} = g^{-1}u_a\ell_a$. For fixed g the map $\ell_a \mapsto h_a$ is a left translation, so these coordinates carry product Haar measure, and

$$v_{x'_a} = g^{-1}\ell_a w_a = g^{-1}u_a^{-1}g\,h_a w_a.$$

Fix h and put $m_h = \max_a \delta(h_a, I) < 1$. The set S_h of g with $(g,h) \in R_1$ is the measurable set where $\delta(v_{x'_a}, I) \le m_h$ for every a; on it $m = m_h$, so $\chi = \phi(m_h)$ does not depend on g, and $V^\sharp = g\,e^{\chi N(g)}$ with $N(g) = \frac{1}{16}\sum_a(\log h_a + \log v_{x'_a})$. The set $\{\delta(v,I) < 1\}$ is the ball of angular radius α about I on the sphere, which is geodesically convex, and on it $\|\log v - \log v'\| \le \Lambda\, d(v,v')$. By bi-invariance $d(g^{-1}u^{-1}g, g'^{-1}u^{-1}g') \le 2d(g,g')$, so for $g, g' \in S_h$

$$\|N(g) - N(g')\| \le \frac{7}{16}\,2\Lambda\, d(g,g') = \frac{7}{8}\Lambda\, d(g,g').$$

If $V^\sharp(g) = V^\sharp(g') = c$, then $g = c\,e^{-\chi N(g)}$ and $g' = c\,e^{-\chi N(g')}$, so $d(g,g') \le \frac{7}{8}\Lambda\, d(g,g')$ and $g = g'$: the map $g \mapsto V^\sharp(g)$ is injective on S_h. Its derivative is computed as in the proof of

Lemma 12.45: with $\eta = \mathrm{Ad}(g)\zeta$ the sites x'_a move by $e^{t(A_a-1)\zeta}$, $A_a = \mathrm{Ad}(g^{-1}u_a^{-1}g)$, the sites x_a do not move, and the right trivialised derivative is $\mathrm{Ad}(g)T$ with $\|T-1\| \le \frac{7}{16}\,2\Lambda = \frac{7}{8}\Lambda$, so its determinant is at least $(1-\frac{7}{8}\Lambda)^3$ in modulus. On R_2 the coordinates $(g, h'_1, \dots, h'_7)$ with $h'_a = v_{x'_a} = g^{-1}\ell_a w_a$ and $v_{x_a} = g^{-1}u_a g\, h'_a w_a^{-1}$ give the same conclusions.

For fixed h, the smooth formula with $\chi = \phi(m_h)$ fixed extends to a neighbourhood of each point of S_h, where no logarithm meets $-I$; its restriction to the measurable set S_h is injective and has the Jacobian bounds above. The change of variables formula for measurable restrictions therefore applies, even if part of the boundary of S_h has positive measure, and likewise on R_2. The law of $V^\sharp$ has the density

$$\rho(c) = h^{\otimes 7}\big(\{\ell : (c,\ell) \in R_0\}\big) + \sum_{i=1,2} \int \sum_{g \in S_h^{(i)},\ V^\sharp = c} \frac{dh^{\otimes 7}}{|\det DV^\sharp(g)|}, \tag{12.7}$$

the inner sum having at most one term, and $\rho(c) \le 1 + 2(1-\frac{7}{8}\Lambda)^{-3}$. Finally $1 - \frac{7}{8}\Lambda = 0.04391\ldots$ and $M = 23621\ldots$. □

The four averages of a block depend on disjoint sets of private links, and on the links inside the blocks, which are private to none. So, given the links inside the blocks of W, the averages $V_k^\sharp$ are independent, and the formula (12.7) for each pair defines, for every value c of C, a measure ν_c on the fibre $F_c = C^{-1}(c)$: in the coordinates of the proof of Lemma 12.47, chosen for each pair according to its region, it is the product Haar measure of the links inside the blocks and of the seven coordinates other than g, divided by the product of the determinants $|\det DV_k^\sharp(g)|$. Then $h = \int \nu_c\, dh(c)$ on G^{A_W}, so ν_c is a version of the conditional law, and

$$\nu_c(F_c) \le M^{4|W|} \qquad \text{for every } c. \tag{12.8}$$

Where C is differentiable,

$$\nu_c = \mathrm{vol}(G)^{-45|W|}\, \det\big(DC\, DC^*\big)^{-1/2}\, d\mathcal{H}.$$

Indeed, if $C(z,g) = c$ is solved for g by $g = \psi(z)$, then $DC = \partial_g C\,(-D\psi,\, 1)$, so

$$\det\big(DC\, DC^*\big)^{1/2} = |\det \partial_g C|\, \det\big(1 + D\psi^* D\psi\big)^{1/2},$$

and the second factor is the Jacobian of the graph of ψ. So ν_c is the coarea version on the charts of Theorems 12.37 and 12.41, where C is smooth, and we use it on the whole fibre.

Proposition 12.48. *Let $G = \mathrm{SU}(2)$, $\lambda > 0$ and $0 < K \le 10^{-8}$, and let $n_P \le 384|W|$ be the number of plaquettes that contain a link of A_W. Suppose the fibre F contains a field U with $\delta(U_p, I) \le \kappa$ for every plaquette p with a vertex in a block of W^+, where $\kappa \le K/(4\sqrt{n_P})$. Then the constrained law μ_U on F gives the set of fields with a plaquette p with a vertex in a block of W^+ and $\delta(U'_p, I) > K$ mass at most*

$$\varepsilon_W(K,\lambda) = \Big(\frac{3\cdot 10^8\, n_P^{3/2}}{K^3}\Big)^{45|W|} e^{-3\lambda K^2/4}.$$

Proof. Let $c = C(U)$, let B be the set in question, and let Q_W be the sum of $\delta(U'_p, I)^2$ over the n_P plaquettes that contain a link of A_W; on F it differs from Q by a constant, so $\mu_U = e^{-\lambda Q_W}\nu_c / \nu_c(e^{-\lambda Q_W})$. A plaquette that contains no link of A_W has $U'_p = U_p$ for $U' \in F$, so on B the large plaquette is one of the n_P and $Q_W > K^2$. By (12.8),

$$\int_B e^{-\lambda Q_W}\, d\nu_c \le M^{4|W|} e^{-\lambda K^2}.$$

For the denominator we solve the constraints for the path links. Let $L' = A_W \setminus \{d_k\}$, $\rho = K/(188\sqrt{n_P})$ and $R = 44\rho$, and let D be the set of $z \in G^{L'}$ with $d(z_e, U_e) \le \rho$ for every $e \in L'$. The field U satisfies the hypothesis of Lemma 12.34, as $\kappa \le K/4$, and part (i) of

the lemma gives $\delta(U_e, I) \le 3\kappa$ for the links inside the blocks and $\delta(U_e, U_{d_k}) \le 21\kappa$ for the private links of k, so $\delta(v_x, I) \le 24\kappa$ and, by Lemma 12.26, $d(v_x, I) \le 8\pi\kappa$ at every site of every pair. Fix $z \in D$ and a pair k, and let ℓ range over the ball B_k of radius R about U_{d_k}. Each v_x is ℓ^{-1} times a product of at most two links of L' and fixed links, so at the field with links z and ℓ it lies within $R + 2\rho$ of its value at U, and within $8\pi\kappa + 46\rho < 0.01$ of I. Hence the cut-off equals 1 near this field and $V_k^\sharp = \ell\, e^{N_k(\ell)}$, with N_k depending on z and ℓ. Each v_x is moved by $d(\ell, \ell')$ when ℓ is replaced by ℓ', so the map $\Psi(\ell) = c_k\, e^{-N_k(\ell)}$ satisfies $d(\Psi(\ell), \Psi(\ell')) \le \frac{7}{8}\Lambda\, d(\ell, \ell')$. At U the path link solves $U_{d_k} = c_k e^{-N_k}$, and the sites move by at most $R + 2\rho$ between U and the field with links z and ℓ, so

$$d(\Psi(\ell), U_{d_k}) \le \frac{7}{8}\Lambda\,(R + 2\rho) < 0.9561 \cdot 46\rho < R.$$

By the contraction principle Ψ has a unique fixed point $\psi_k(z) \in B_k$. The path link d_k is private to k, so the field $\Gamma(z)$ with links z on L' and $\psi_k(z)$ on d_k lies in F, and Γ is injective.

Near $\Gamma(D)$ every cut-off equals 1, so C is smooth, and $\partial_g C$ is block diagonal with the blocks $\partial_{d_k} V_k^\sharp$. Moving $d_k \mapsto e^{t\eta} d_k$ multiplies every v_x of k by $e^{-t\zeta}$, $\zeta = \mathrm{Ad}(d_k^{-1})\eta$, so the block is $\mathrm{Ad}(d_k)(1 - E_{N_k} \frac{1}{16} \sum_x L_{X_x})$, with determinant at most $(1 + \frac{7}{8}\Lambda)^3 < 1.957^3$ in modulus. By the identity above, ν_c restricted to $\Gamma(D)$ is the image of $h^{\otimes L'}$ on D divided by $|\det \partial_g C|$, so

$$\nu_c(\Gamma(D)) \ge 1.957^{-12|W|} h(B_\rho)^{45|W|},$$

$$h(B_\rho) = \frac{2\theta - \sin 2\theta}{2\pi} \ge \frac{\rho^3}{14},$$

with $\theta = \rho/\sqrt{2}$ the angular radius, as $2\theta - \sin 2\theta \ge \frac{4}{3}\theta^3(1 - \frac{\theta^2}{5})$. Every plaquette has at most one path link, since two path links $(2y + e_\mu, \mu)$ and $(2y' + e_\mu, \mu)$ are never opposite sides of a plaquette, so on $\Gamma(D)$ each plaquette moves by at most $R + 3\rho = 47\rho$ and $\delta(U'_p, I) \le \kappa + 47\rho \le K/(2\sqrt{n_P})$. Hence $Q_W \le K^2/4$ there and

$$\int_F e^{-\lambda Q_W}\, d\nu_c \ge e^{-\lambda K^2/4}\, 1.957^{-12|W|} \Big(\frac{\rho^3}{14}\Big)^{45|W|}.$$

Dividing,

$$\mu_U(B) \le \big(M^4 1.957^{12}\big)^{|W|} \Big(\frac{14 \cdot 188^3 n_P^{3/2}}{K^3}\Big)^{45|W|} e^{-3\lambda K^2/4},$$

and $(M^4 1.957^{12})^{1/45} < 2.93$, $14 \cdot 188^3 < 9.31 \cdot 10^7$. Finally $n_P \le 384|W|$, since each plaquette that contains a link of A_W has the base of that link, a site of a block of W, as a vertex, and each site is a vertex of 24 plaquettes. □

Corollary 12.49. *In addition to the hypotheses of Proposition 12.48, assume those of Theorem 12.41 with K in place of κ, and $s \ge 1.24 \cdot 10^5 K$. Then $\mu_U(F \setminus \Phi_U(\Omega_r)) \le \varepsilon_W(K, \lambda)$, and for $X_1, X_2 \subset A_W$ and $\tilde F, \tilde G$ as in Theorem 12.41*(iii)*,*

$$|\mathrm{Cov}_{\mu_U}(\tilde F, \tilde G)| \le \Big(\frac{c_9}{\lambda} e^{-\alpha_G d(X_1, X_2)} + 30\, \varepsilon_W(K, \lambda)\Big) |X_1|\, |X_2|.$$

Proof. The field U satisfies the hypothesis of Lemma 12.34 with K, so Theorem 12.44(i) with K in place of κ puts every point of F outside $\Phi_U(\Omega_r)$ in the set of Proposition 12.48. The covariance bound follows as in the proof of Theorem 12.44(iii). □

So the constrained law decorrelates on every fibre through a field whose plaquettes near W are at most $K/(4\sqrt{n_P})$, with no average over fibres, once λK^2 is a large multiple of $|W| \log(n_P/K)$; with $K < 4.9 \cdot 10^{-18}$ this needs β_W above $1.5 \cdot 10^{39}|W|$ under the Wilson measure. The restriction is on the fibre: the links off A_W are fixed on it, so a fibre whose boundary links carry a large plaquette near W contains no such field, and neither may a fibre with small boundary links, and the proposition does not cover these fibres. It is a

lower bound for the fibre measure near one good field against an upper bound for the whole fibre; it does not compare different fibres.

The hypothesis on U enters the proof twice: through Lemma 12.34, to place the sixteen sites v_x of every pair near I, and through the bound on Q_W near U. Neither needs the links off A_W or the links inside the blocks to be small, and every fibre contains a field at which the sites are exactly I.

Lemma 12.50. *For every value c of C and every choice of the links off A_W, the fibre F_c contains a field at which $v_x = I$ for every site x of every pair (y,μ), $y \in W$.*

Proof. Fix μ. The links of A_W in the direction μ are, for each $y \in W$, the links u_a and ℓ_a of the pair (y,μ), and the links w_a of that pair are the links u_a of the pair $(y+e_\mu,\mu)$ when $y+e_\mu \in W$, and fixed links otherwise. Whether u_a is a tree link depends only on the offset, so u_a and w_a are both tree links or neither. For a pair with path link g and given links w_a put $\ell_0 = g$ and, for $a \neq 0$,

$$\ell_a = g, \quad \text{or} \quad \ell_a = g\,w_a^{-1},\ u_a = g\,w_a\,g^{-1},$$

the first when u_a is a tree link and the second otherwise. Then $v_{x_a} = g^{-1}u_a\ell_a = I$ and $v_{x'_a} = g^{-1}\ell_a w_a = I$, as $u_a = w_a = I$ in the first case. The blocks of W along a line in the direction μ form runs $y_1,\dots,y_n$ of consecutive blocks. If $y_n + e_\mu \notin W$, choose the links of the pairs $(y_n,\mu),\dots,(y_1,\mu)$ in this order, with $g = c_{y_j,\mu}$, the links w_a of (y_j,μ) being the fixed links for $j = n$ and the links u_a just chosen for (y_{j+1},μ) otherwise. If the run closes up around the torus, put $u_a = I$ and $\ell_a = g$ for every pair of the run. At the resulting field every v_x equals I, so the cut-off equals 1, $N = 0$ and $V^\sharp_{y,\mu} = \ell_0 = c_{y,\mu}$. The directions do not interact, since the pairs (y,μ) use only links in the direction μ. □

Call a field $U \in F$ *regular* if $\delta(v_x, I) \le \frac14$ at every site x of every pair, and let $q(F)$ be the infimum of Q_W over the regular fields of F, finite by Lemma 12.50. Write $\mu_F = e^{-\lambda Q_W}\nu_c / \nu_c(e^{-\lambda Q_W})$ for the constrained law on $F = F_c$.

Proposition 12.51. *Let $G = \mathrm{SU}(2)$, $\lambda > 0$ and $0 < t \le 1$. For every fibre F, whatever the links off A_W, and every regular $U \in F$,*

$$\mu_F\big(\{U' \in F : Q_W(U') > Q_W(U) + t\}\big) \le \varepsilon'_W(t,\lambda),$$

$$\varepsilon'_W(t,\lambda) = \Big(\frac{5.2\cdot 10^{10}\, n_P^3}{t^3}\Big)^{45|W|} e^{-3\lambda t/4}.$$

In particular $\mu_F(\{Q_W > q(F) + t\}) \le \varepsilon'_W(t,\lambda)$ on every fibre.

Proof. The numerator is bounded by (12.8) as in the proof of Proposition 12.48, by $M^{4|W|}e^{-\lambda(Q_W(U)+t)}$. For the denominator take $\rho = t/(1076\,n_P) \le 10^{-3}$, $R = 44\rho$, and D and B_k as in that proof. At the field with links $z \in D$ on L' and $\ell \in B_k$ on d_k, each v_x of the pair k lies within $R + 2\rho = 46\rho < 0.043$ of its value at U, and $\delta \le d$, so $\delta(v_x, I) < \frac14 + 0.043 < \frac12$. Hence the cut-off equals 1 near this field, the logarithms are Λ-Lipschitz along the moves, and the contraction argument of that proof gives the injective map $\Gamma : D \to F$, near whose image C is smooth with $|\det \partial_g C| \le 1.957^{3\cdot 4|W|}$. So

$$\nu_c(\Gamma(D)) \ge 1.957^{-12|W|}\Big(\frac{\rho^3}{14}\Big)^{45|W|}.$$

On $\Gamma(D)$ each of the n_P plaquettes moves by at most 47ρ, and $\delta(U_p, I) \le 2\sqrt2$, so

$$\delta(U'_p, I)^2 \le \delta(U_p, I)^2 + 94\cdot 2\sqrt2\,\rho + 47^2\rho^2 < \delta(U_p, I)^2 + 269\rho,$$

and $Q_W \le Q_W(U) + 269 n_P \rho = Q_W(U) + \frac{t}{4}$ there. Dividing as before,

$$\mu_F(\{Q_W > Q_W(U) + t\}) \le (M^4 1.957^{12})^{|W|} \Big(\frac{14 \cdot 1076^3 n_P^3}{t^3} \Big)^{45|W|} e^{-3\lambda t/4},$$

and $2.93 \cdot 14 \cdot 1076^3 < 5.2 \cdot 10^{10}$. For the last statement apply this to regular fields with $Q_W(U)$ decreasing to $q(F)$ and use the continuity of ε'_W in t. □

Corollary 12.52. *Let $G = \mathrm{SU}(2)$, $\lambda > 0$ and $0 < K \le 1$. Suppose the fibre F contains a regular field U with $\delta(U_p, I) \le K/(4\sqrt{n_P})$ on every plaquette p that contains a link of A_W, whatever the plaquettes made of fixed links. Then μ_F gives the fields with such a plaquette above K mass at most $\varepsilon'_W(\frac{15}{16}K^2, \lambda)$.*

Proof. At U, $Q_W \le K^2/16$, and a field with one of these plaquettes above K has $Q_W > K^2 \ge Q_W(U) + \frac{15}{16}K^2$. Apply Proposition 12.51 with $t = \frac{15}{16}K^2 \le 1$. □

The field of Proposition 12.48 is regular, since its sites lie within 24κ of I, so Corollary 12.52 contains that proposition up to the constants and drops its hypothesis on the plaquettes made of fixed links and on the links inside the blocks. It reaches every fibre whose boundary links carry large plaquettes, provided the sites of the averages can be made small there together with the plaquettes that move. Proposition 12.51 holds on every fibre with no hypothesis at all, but its reference value $q(F)$ need not be small. Where it is of order one, as it can be when the averages are far from what the boundary links suggest, the constrained law sits within t of $q(F)$ but away from the charts of Theorem 12.41, which are built at fields with all plaquettes near W small. The proposition also does not compare regular with irregular fields, at which some site is far from I.

Different fibres with the same links off A_W are compared through the law of the averages. Call the links off A_W the *exterior* E, write h for product Haar measure on G^{A_W}, and let

$$\pi_E = \frac{e^{-\lambda Q_W} h}{h(e^{-\lambda Q_W})}$$

on G^{A_W}. Under the SU(2) Wilson measure in the tree gauge, π_E with $\lambda = \beta_W/4$ is the conditional law of the links of A_W given the exterior. Since $h = \int \nu_c \, dh(c)$,

$$\pi_E = \int \mu_{F_c} \frac{\nu_c(e^{-\lambda Q_W})}{h(e^{-\lambda Q_W})} \, dh(c),$$

so μ_{F_c} is the conditional law of π_E given $C = c$, and the law of the fibre through a field drawn from π_E is the law of C under π_E. Let $Q_*(E)$ be the least value of Q_W on G^{A_W}, which is attained since G^{A_W} is compact.

Proposition 12.53. *Let $G = \mathrm{SU}(2)$, $\lambda > 0$ and $0 < t \le 1$. For every exterior E,*

$$\pi_E(\{Q_W > Q_*(E) + t\}) \le \varepsilon''_W(t, \lambda),$$
$$\varepsilon''_W(t, \lambda) = \Big(\frac{1.05 \cdot 10^7 \, n_P^3}{t^3} \Big)^{49|W|} e^{-3\lambda t/4}.$$

In particular the law of the averages under π_E gives the fibres F with $\inf_F Q_W > Q_*(E) + t$ mass at most $\varepsilon''_W(t, \lambda)$.

Proof. Since h is a probability measure,

$$\int_{\{Q_W > Q_* + t\}} e^{-\lambda Q_W} \, dh \le e^{-\lambda(Q_* + t)}.$$

For the normalisation let U_* be a field with $Q_W(U_*) = Q_*$, let $\rho = t/(90.6\, n_P)$, and let D be the set of fields with $d(U_e, U_{*,e}) \le \rho$ for each of the $49|W|$ links of A_W, the $45|W|$ links of L'

and the $4|W|$ path links. Then $h(D) = h(B_\rho)^{49|W|} \geq (\rho^3/14)^{49|W|}$. Each plaquette contains at most four links of A_W, so on D it moves by at most 4ρ, and as $\delta \leq d$ and $\delta(U_p, I) \leq 2\sqrt{2}$,

$$\delta(U_p, I)^2 \leq \delta(U_{*,p}, I)^2 + 16\sqrt{2}\,\rho + 16\rho^2 < \delta(U_{*,p}, I)^2 + 22.63\rho,$$

because each link of A_W lies in six plaquettes, so $n_P \geq 74$ and $\rho < 1.5 \cdot 10^{-4}$. Hence $Q_W \leq Q_* + 22.63\, n_P \rho < Q_* + \frac{t}{4}$ on D, and

$$\pi_E(\{Q_W > Q_* + t\}) \leq \Big(\frac{14 \cdot 90.6^3 n_P^3}{t^3}\Big)^{49|W|} e^{-3\lambda t/4},$$

with $14 \cdot 90.6^3 < 1.05 \cdot 10^7$. A field U with $Q_W(U) \leq Q_* + t$ lies in its own fibre, so $\inf_{F(U)} Q_W \leq Q_* + t$, which gives the last statement. □

Call a field that agrees with E off A_W a *completion* of E.

Corollary 12.54. *Let $G = \mathrm{SU}(2)$, $\lambda > 0$, $0 < K \leq 10^{-8}$ and $\kappa = K/(4\sqrt{n_P})$. Suppose the exterior E has a completion U^0 with $\delta(U^0_p, I) \leq \kappa/(2\sqrt{n_P})$ for every plaquette p with a vertex in a block of W^+. Let $\mathcal{E}$ be the set of completions U with $\delta(U_p, I) \leq \kappa$ for every such plaquette. Then:*

(i) $\pi_E(\mathcal{E}^c) \leq \varepsilon''_W(\kappa^2/2, \lambda)$;

(ii) *every $U \in \mathcal{E}$ is regular and its fibre has $q(F(U)) < \kappa^2$, so the law of the averages under π_E gives the fibres with $q(F) \geq \kappa^2$, among them every fibre where $q(F)$ is of order one, mass at most $\varepsilon''_W(\kappa^2/2, \lambda)$;*

(iii) *if λ, s and K satisfy the hypotheses of Corollary 12.49, then for every $U \in \mathcal{E}$ the constrained law on $F(U)$ gives the complement of the chart $\Phi_U(\Omega_r)$ at U mass at most $\varepsilon_W(K, \lambda)$, and its covariances obey the bound of that corollary.*

Proof. (i) $Q_*(E) \leq Q_W(U^0) \leq n_P\kappa^2/(4n_P) = \kappa^2/4$. If $Q_W(U) \leq Q_* + \kappa^2/2$, then $Q_W(U) \leq \frac{3}{4}\kappa^2$, so every plaquette that contains a link of A_W has $\delta(U_p, I) < \kappa$, and a plaquette with a vertex in a block of W^+ that contains no link of A_W has the value it has at U^0, at most $\kappa/(2\sqrt{n_P})$. So $\mathcal{E}^c \subset \{Q_W > Q_* + \kappa^2/2\}$, and Proposition 12.53 with $t = \kappa^2/2 \leq 1$ applies. (ii) A field $U \in \mathcal{E}$ satisfies the hypothesis of Lemma 12.34, which places every site of every pair within $24\kappa < \frac{1}{4}$ of I, as in the proof of Proposition 12.48; so U is regular and $q(F(U)) \leq Q_W(U) < \kappa^2$. Hence a fibre with $q(F) \geq \kappa^2$ contains no field of $\mathcal{E}$, and the law of the averages gives these fibres at most the mass $\pi_E(\mathcal{E}^c)$. (iii) The fibre $F(U)$ contains the field U, whose plaquettes with a vertex in a block of W^+ are at most $\kappa = K/(4\sqrt{n_P})$, which is the hypothesis of Proposition 12.48; Corollary 12.49 applies at U. □

So on every exterior whose plaquettes near W can all be made small by the links of A_W, the fibres where $q(F)$ is of order one, and more generally the fibres that contain no field with small plaquettes near W, carry total mass at most $\varepsilon''_W(\kappa^2/2, \lambda)$ in the law of the averages, and every other fibre has the chart decay of Corollary 12.49. This is a bound for every such exterior, not a mean over exteriors as in Theorem 12.44(ii). If $\lambda\kappa^2 \geq 196|W|\ln(8.4 \cdot 10^7 n_P^3/\kappa^6)$, then $\varepsilon''_W(\kappa^2/2, \lambda) \leq e^{-\lambda\kappa^2/8}$; with $n_P \leq 384|W|$ and $K = 4.8 \cdot 10^{-18}$ this holds under the Wilson measure for $\beta_W \geq 6.4 \cdot 10^{43}|W|^2(1 + \ln|W|)$. The fibres where $q(F)$ is of order one remain possible in two places: on an exterior with no completion whose plaquettes near W are small, and as values of c of small mass on an exterior with one.

Remark 12.55. Theorem 12.41 is the decay that Remark 12.39(b) asks for, on the chart $\Phi(\Omega_r)$ with $r = s/32$, which contains $\Phi(K_{s/32})$ and so every $Ue^X \in F$ with $\|X\|_\infty \leq s/1280$. Its constants are explicit by Propositions 12.38 and 12.43: the rate $\alpha_G = 2.5 \cdot 10^{-4}$ is the same for every group and lies far below the decay seen in the computation described at the end of this remark. What remains open for the small fields:

(a) the mass of the constrained law outside $\Phi(\Omega_r)$ on the fibres that contain no field whose plaquettes near W are small: Corollary 12.49 bounds it on every fibre that contains one, Corollary 12.54 shows that on every exterior with a completion whose plaquettes near W are small these fibres, among them those where $q(F)$ is of order one, have mass at most ε''_W in the law of the averages, and Theorem 12.44 bounds them in mean over all exteriors; the constrained cube condition of Remark 12.19(a) takes a supremum over boundary conditions, which include these fibres of small mass and the exteriors with no such completion, and on them no chart of Theorem 12.41 is available; by Proposition 12.71 these are the typical case on cubes of fixed physical size along the tuning;

(b) the passage from the decay of the constrained law on one fibre, at a fixed background, to the constrained cube condition of Remark 12.19(a), whose supremum runs over boundary conditions that include large fields.

On the periodic lattice of side twelve, for SU(2) backgrounds as in the check of Theorem 12.35 and W a row of five blocks, the section S has row sums at most 14, the blocks of $\bar{P}$ beyond the range of S decay by a factor about 0.4 per unit of distance, and the blocks of the chart Hessian of Q fall from 15 at distance zero to 10^{-8} at distance nine, a floating-point computation reproduced in the archive.

12.5. **Large fields under the Wilson measure.** Theorem 12.33 needs every plaquette to be small. This subsection shows that, under the SU(2) Wilson measure, the plaquettes that are not small form clusters of microscopic size, and that the cut-off average departs from the average only next to them. It also shows that such plaquettes are present along the one-loop tuning, so the composition has to pass through them.

Let $\Lambda_L = (\mathbb{Z}/L\mathbb{Z})^4$ with $L = 2^m \geq 4$, and let μ_L be the SU(2) Wilson measure on Λ_L, with density proportional to $e^{-\beta_W S(U)}$ with respect to product Haar measure, where

$$S(U) = \sum_p s_p, \qquad s_p = 1 - \tfrac{1}{2}\operatorname{Tr} U_p = \tfrac{1}{4}\,\delta(U_p, I)^2.$$

This is the Wilson measure of Section 3 on the torus, without gauge fixing; it has $4L^4$ links and $6L^4$ plaquettes. For $0 < \rho \leq 1$ call a plaquette *bad* if $\delta(U_p, I) > \rho$. The closed unit cube $\mathcal{C}_x = x + [0,1]^4$ contains 24 plaquettes. Let B_x be the event that one of them is bad. A plaquette lies in exactly four unit cubes, one for each choice of the two transverse coordinates of the cube's corner.

Lemma 12.56. *Let Z_L be the normalising constant of μ_L and $C_0 = \frac{1}{2} + \frac{2}{3}\log(24\pi^3)$. For $\beta_W \geq 1$,*

$$-\log Z_L \leq 6L^4(\log \beta_W + C_0), \qquad \mathbb{E}_{\mu_L} S \leq 6L^4\,\frac{\log \beta_W + C_0}{\beta_W}.$$

Proof. Write $U = \cos\theta\, I + \sin\theta\, J$ with $0 \leq \theta \leq \pi$ and $J^2 = -I$. The angle θ of $g^{-1}h$ is a bi-invariant distance, and under Haar measure θ has density $\frac{2}{\pi}\sin^2\theta$. Put $r = (4\sqrt{\beta_W})^{-1}$. If every link has angle at most r, every plaquette has angle at most $4r \leq 1$ and $s_p \leq (4r)^2/2 = 1/(2\beta_W)$, so $S \leq 3L^4$. Each link angle is at most r with probability

$$\frac{2}{\pi}\int_0^r \sin^2 t\, dt \geq \frac{8r^3}{3\pi^3} = \frac{1}{24\pi^3\beta_W^{3/2}},$$

by $\sin t \geq 2t/\pi$. Hence $Z_L \geq e^{-3L^4}(24\pi^3\beta_W^{3/2})^{-4L^4}$, which is the first bound. The function $F(b) = -\log Z_L$ at the coupling b is concave with $F(0) = 0$ and $F'(b) = \mathbb{E}S$, so $bF'(b) \leq F(b)$, which is the second. $\square$

Theorem 12.57. *For $\beta_W \geq 1$, $0 < \rho \leq 1$ and every set A of unit cubes,*

$$\mu_L\Big(\bigcap_{x\in A} B_x\Big) \leq q^{|A|},$$

$$q = q_\rho(\beta_W) = \min\{1, \exp\big(6(\log\beta_W + C_0) - \beta_W\rho^2/16\big)\}.$$

Proof. The measure μ_L is reflection positive in every hyperplane through sites [67]: the plaquettes are either in the hyperplane or on one side of it, so conditionally on the links in the hyperplane the weight factors into two mirror images, and reflection preserves Haar measure and $\operatorname{Tr} U_p$. On a torus of side 2^m this gives the chessboard estimate for functions of the links of closed unit cubes [37], [21, Section 5.3]:

$$\mu_L\Big(\bigcap_{x\in A} B_x\Big) \leq \mu_L\Big(\bigcap_{x\in\Lambda_L} B_x\Big)^{|A|/L^4},$$

since the reflection of B_x into another cube is the event B_y of that cube. If every cube is bad, there are at least $L^4/4$ bad plaquettes, so $S \geq \rho^2 L^4/16$ and the event has unnormalised weight at most $e^{-\beta_W\rho^2 L^4/16}$. Dividing by Z_L and using Lemma 12.56 gives the bound. □

Corollary 12.58. *Join two unit cubes when their corners are at ℓ^∞ distance one on the torus. For an integer $r \geq 1$,*

$$\mu_L\{\text{some connected set of } r \text{ bad cubes}\} \leq L^4(6400\,q)^r.$$

Hence, on a torus of fixed physical side La, if $a \to 0$ along $L = 2^m$ and $\beta_W(a) \to \infty$, the physical diameters of all connected clusters of bad cubes tend to 0 in probability, for every fixed ρ.

Proof. A connected set of r cubes containing a given cube is recovered from a walk of length $2(r-1)$ around a spanning tree, and each cube has 80 neighbours, so there are at most $80^{2(r-1)} \leq 6400^r$ such sets. Apply Theorem 12.57 and sum over the L^4 roots. A cluster of physical diameter at least R contains a connected set of $\lceil R/a\rceil$ cubes, and once $6400\,q \leq \frac{1}{2}$ the bound is at most $(La/a)^4 2^{-R/a} \to 0$. □

The bad cubes control where the cut-off average differs from the average. For a coarse link (y,μ) let $\mathcal{Z}_{y,\mu}$ be the 24 unit cubes with corners $2y + te_\mu + w$, $t \in \{0,1,2\}$, $w \in \{0,1\}^4$, $w_\mu = 0$ (Figure 19).

Corollary 12.59. *Let $\rho = \frac{1}{12}$. If no cube of $\mathcal{Z}_{y,\mu}$ is bad, then $\chi_{y,\mu}(U) = 1$ in the tree gauge, and $V^\sharp_{y,\mu}$ is the average of the sixteen paths in the chart at P_{2y}. A bad cube lies in $\mathcal{Z}_{y,\mu}$ for at most eight coarse links, so the expected fraction of coarse links with $\chi < 1$ is at most $4.5\cdot 10^5\,(\log\beta_W + C_0)/\beta_W$. Along the limits of Corollary 12.58, with probability tending to one these links lie within physical distance $2a$ of clusters of bad cubes of physical diameter less than any fixed $R > 0$.*

Proof. As in the proof of Lemma 12.32, $P_{2y}^{-1}P_x$ is a product of at most six conjugates of plaquette holonomies or their inverses. These plaquettes lie in planes $\{\mu,\alpha\}$ with corners $2y + te_\mu + w$ as above, and a plaquette lies in the unit cube at its corner. So $\delta(P_x, P_{2y}) \leq 6\rho = \frac{1}{2}$, and $\chi = 1$. A cube with corner z lies in $\mathcal{Z}_{y,\mu}$ only if $z_\mu - 2y_\mu \in \{0,1,2\}$ and $z_\alpha - 2y_\alpha \in \{0,1\}$ for $\alpha \neq \mu$, which leaves at most two coarse links in each direction. There are $L^4/4$ coarse links, the number of bad cubes is at most four times the number of bad plaquettes, and a bad plaquette has $s_p > \rho^2/4$, so by Lemma 12.56 the expected fraction is at most $4\cdot 8\cdot 4\cdot 24\cdot 144\,(\log\beta_W + C_0)/\beta_W$. The last assertion is Corollary 12.58. □

Small plaquettes everywhere, which Theorem 12.33 assumes, are nevertheless improbable along the one-loop tuning. For SU(2), $\beta_W = 4g^{-2}$, and the one-loop coupling of Remark 12.19(c) at spacing $a = 1/L$ gives $\beta_W = 8b_0\log L + O(1) = \frac{11}{3\pi^2}\log L + O(1)$.




FIGURE 19. Large fields under the SU(2) Wilson measure. (a) A plane section of the lattice: bad unit cubes (red) form clusters, and the cut-off average can differ from the path average only on coarse links (orange) whose 24 cubes $\mathcal{Z}_{y,\mu}$ meet a bad cube (Corollary 12.59). (b) The frustrated exterior of a link e: the staple of p_0 is I and the other five staples are $-I$, so U_e is driven to $-I$ and p_0 is bad. (c) The weight $q_\rho(\beta_W)$ of Theorem 12.57 for three values of ρ; below the dashed line the cluster bound of Corollary 12.58 is summable.

Proposition 12.60. *Let $0 < \rho \le 1$, $v_* = \frac{2}{\pi}\int_{\arccos(3/4)}^{\pi/3} \sin^2 t\, dt$ and $p = v_* e^{-3\beta_W}$. Conditionally on all other links, a plaquette through a link e is bad with probability at least p. Hence*

$$\mu_L\{\text{no plaquette is bad}\} \le e^{-\lfloor L/3\rfloor^4 p},$$

which tends to 0 when $\beta_W \le B\log L + O(1)$ with $B < 4/3$, in particular along the one-loop tuning.

Proof. Identify SU(2) with the unit quaternions. Conditionally on the other links, U_e has density proportional to $e^{\kappa\, h\cdot u}$ with $0 \le \kappa \le 6\beta_W$ and $|h| = 1$, the six staples contributing at most β_W each. The plaquette holonomy is U_e or its inverse multiplied by a fixed element, so it has a density of the same form. Writing the density as a superposition of indicators of caps centred at h, the probability of a cap about I is largest when $h = I$, and for $h = I$ it increases with κ, because its derivative is the covariance of an increasing function of $\operatorname{Tr} U$ with $\frac{1}{2}\operatorname{Tr} U$. So the probability of being bad is at least that of the angle exceeding $\arccos(1-\rho^2/4) \le \arccos(3/4)$ under the density proportional to $e^{6\beta_W \cos t}\sin^2 t$, which is at least p: bound the numerator from below on $[\arccos(3/4), \pi/3]$, where $\cos t \ge \frac{1}{2}$, and the denominator by $e^{6\beta_W}\pi/2$.

Let $M = \lfloor L/3\rfloor$. Choose the links in direction 0 at the sites $3z$, $z \in \{0,\dots,M-1\}^4$, and their plaquettes in the plane $\{0,1\}$. No chosen plaquette contains another chosen link. Removing one chosen link at a time therefore bounds the probability that all chosen plaquettes are good by $(1-p)^{M^4}$. With $\beta_W = B\log L + O(1)$ the exponent is of order L^{4-3B}. □

The composition of Theorem 12.33 is local, so it continues across the scales away from the defects, and the defects spread by at most three spacings of the current scale at each step. For a coarse plaquette P at y let $\mathcal{B}_P$ be the fine box $2y + [0,4]^4$.

Lemma 12.61. *Let U be a G-valued field in the tree gauge and P a coarse plaquette. If every plaquette p in $\mathcal{B}_P$ has $\delta(U_p, I) \le \rho_G$ and $\|\Theta_p\| \le \eta \le \rho_G/1000$, then $\chi = 1$ on the four sides of P, $\delta(V^\sharp(U)_P, I) \le 0.03\,\rho_G$, the inequalities (12.6) hold for P, and $\|\log V^\sharp(U)_P\| \le 4(1 + 647\eta)\eta$.*

Proof. For the plaquette P the proof of Lemma 12.32 uses only the plaquettes of the surfaces $\mathcal{U}_\ell$ of the four sides ℓ of P and those of the sixteen squares $\square_{2y+o}$. A plaquette of $\mathcal{U}_\ell$, $\ell = (y_i, \mu_i)$, has its corner x at $x_{\mu_i} - 2(y_i)_{\mu_i} \in \{0,1,2\}$ and $x_\alpha - 2(y_i)_\alpha \in \{0,1\}$ for $\alpha \neq \mu_i$, with $y_i \in \{y, y+e_\mu, y+e_\nu\}$; a plaquette of $\square_{2y+o}$ has $x - 2y \in \{0,1,2\}^4$. So $x - 2y \in \{0,1,2,3\}^4$ in every case, and the plaquette lies in $\mathcal{B}_P$. The bounds on χ, on $\delta(V_P^\sharp, I)$, the inequalities (12.6) and the bound $\|\bar F_P\| + 3.2\, r_P^2 \le 4(1+647\eta)\eta$ in that proof use only the norms of these plaquettes. □

Let a be the fine spacing and $a_j = 2^j a$. Given $\eta_0 > 0$ put $\eta_{j+1} = 4(1+647\eta_j)\eta_j$. Let $\mathfrak{B}_0$ be the set of fine plaquettes with $\delta(U_p, I) > \rho_G$ or $\|\Theta_p\| > \eta_0$, and let $\mathfrak{B}_{j+1}$ be the set of plaquettes P of the lattice of spacing a_{j+1} whose box $\mathcal{B}_P$ contains a plaquette of $\mathfrak{B}_j$. Let E_j° and A_j° be the sum of squares and the sum of $\|\Theta_P^{(j)}\|$ over the plaquettes $P \notin \mathfrak{B}_j$ of the j-th iterate.

Theorem 12.62. *Let $G \subset \mathrm{U}(r)$ be compact and U a G-valued field on the periodic lattice of side divisible by 2^k, and suppose that $\eta_j \le \rho_G/1000$ for $j < k$.*

(i) *For $j \le k$, every plaquette $P \notin \mathfrak{B}_j$ of the j-th iterate has $\delta(U_P^{(j)}, I) \le \rho_G$ and $\|\Theta_P^{(j)}\| \le \eta_j$. For $j < k$ the cut-off is inactive on the sides of every plaquette outside $\mathfrak{B}_{j+1}$, and (12.6) holds there.*

(ii) *Every point of a plaquette of $\mathfrak{B}_j$ lies within ℓ^∞ distance $3(a_j - a)$ of a plaquette of $\mathfrak{B}_0$.*

(iii) *With $\sigma = \sum_{j<k} \eta_j$,*

$$E_k^\circ \le e^{12060\sigma} E_0^\circ, \qquad A_k^\circ \le e^{9147\sigma} 4^{-k} A_0^\circ.$$

(iv) *If $4^k\eta_0 \le 10^{-4}\rho_G$, the hypothesis holds, $\eta_j \le 1.023 \cdot 4^j\eta_0$ for $j \le k$, $\sigma \le 0.35 \cdot 4^k\eta_0$, and $E_k^\circ \le 1.53\, E_0^\circ$.*

Proof. (i) By induction on j. If $P \notin \mathfrak{B}_{j+1}$, no plaquette of $\mathcal{B}_P$ lies in $\mathfrak{B}_j$, so by the induction hypothesis all of them have $\delta \le \rho_G$ and norm at most $\eta_j \le \rho_G/1000$. Lemma 12.61, applied to the j-th iterate in the tree gauge, gives $\delta(U_P^{(j+1)}, I) \le 0.03\,\rho_G$ and $\|\Theta_P^{(j+1)}\| \le \eta_{j+1}$; the gauge fixing between steps does not change these norms.

(ii) Let P at y lie in $\mathfrak{B}_{j+1}$, and let $p \in \mathfrak{B}_j$ lie in $\mathcal{B}_P$, with corner x, $x - 2y \in \{0,1,2,3\}^4$. In units of a_j every coordinate of a point of P lies in $[2y_c, 2y_c+2]$, and p contains a point whose coordinates are x_c or lie in $[x_c, x_c+1]$; so the two are at distance at most 3 in each coordinate. Hence every point of a plaquette of $\mathfrak{B}_{j+1}$ is within $3a_j$ of a plaquette of $\mathfrak{B}_j$, and $\sum_{i<j} 3a_i = 3(a_j - a)$.

(iii) Repeat the part *Sums* of the proof of Lemma 12.32 with the plaquettes $P \notin \mathfrak{B}_{j+1}$ only. Their squares $\square_{2y+o}$ and surfaces $\mathcal{U}_\ell$ consist of plaquettes outside $\mathfrak{B}_j$, so the same counts give $\sqrt{E_{j+1}^\circ} \le (1+6030\eta_j)\sqrt{E_j^\circ}$ and $A_{j+1}^\circ \le (1+9147\eta_j)A_j^\circ/4$, and $1 + t \le e^t$.

(iv) The recursion gives $\eta_j \le e^{647\sigma_j} 4^j \eta_0$ with $\sigma_j = \sum_{i<j}\eta_i$. Suppose $\sigma_j \le 0.35 \cdot 4^j\eta_0$, which holds at $j = 0$. Then $\sigma_j \le 3.5 \cdot 10^{-5}$, $e^{647\sigma_j} \le 1.023$, $\eta_j \le 1.023 \cdot 4^j\eta_0 \le 1.023 \cdot 10^{-4}\rho_G$, and $\sigma_{j+1} \le 1.373 \cdot 4^j\eta_0 \le 0.35 \cdot 4^{j+1}\eta_0$. Finally $e^{12060 \cdot 3.5 \cdot 10^{-5}} \le 1.53$. □

Under the Wilson measure the defects of Theorem 12.62 are the bad plaquettes of Theorem 12.57. For $\mathrm{SU}(2)$, $\rho_G = 1$; a plaquette with $\delta(U_p, I) = 2\sqrt{2}\sin(\theta/2) \le \rho \le \frac{1}{10}$ has $\|\Theta_p\| = \sqrt{2}\,\theta \le 1.01\,\delta(U_p, I)$, so with $\eta_0 = 1.01\rho$ every plaquette of $\mathfrak{B}_0$ is bad, and lies in a bad cube.

Corollary 12.63. *Let $G = \mathrm{SU}(2)$, $C_1 = C_0 + \frac{1}{6}\log 12800$, and for $\beta_W \ge 1$ let $\rho(\beta_W)$ be given by $\rho^2 = 96(\log\beta_W + C_1)/\beta_W$, so that $6400\, q_\rho(\beta_W) = \frac{1}{2}$. Let k_* be the largest integer with $4^{k_*} \cdot 1.01\rho \le 10^{-4}$. If $1.01\rho \le 10^{-4}$, then $k_* \ge 0$,*

$$a_{k_*} \ge 1.5 \cdot 10^{-3} \Big(\frac{\beta_W}{\log\beta_W + C_1}\Big)^{1/4} a,$$

and with μ_L-probability at least $1 - L^4 2^{-r}$ every connected cluster of bad cubes has fewer than r cubes, while for every $j \le k_*$ the defects of the j-th iterate lie within ℓ^∞ distance $3a_j$ of these clusters and the iterate satisfies Theorem 12.62(i), (iii) with $E^\circ_{k_*} \le 1.53\, E^\circ_0$. Along the limits of Corollary 12.58 with $a\,\rho(\beta_W)^{-1/2} \to 0$, which holds along the one-loop tuning, the defects of all iterates up to the spacing a_{k_*} lie, with probability tending to one, within clusters of physical size tending to 0.

Proof. With this ρ, $q_\rho(\beta_W) = \exp(-\log 12800)$, and Corollary 12.58 gives the probability of a connected set of r bad cubes at most $L^4(6400q)^r = L^4 2^{-r}$. By maximality, $4^{k_*+1} \cdot 1.01\rho > 10^{-4}$, so $2^{k_*} > \frac{1}{2} \cdot 10^{-2}(1.01\rho)^{-1/2}$, and $\rho^{-1/2} = 96^{-1/4}(\beta_W/(\log\beta_W + C_1))^{1/4}$ with $96^{1/4} < 3.14$. The rest is Theorem 12.62 with $\eta_0 = 1.01\rho$, whose hypothesis holds by part (iv), and $\mathfrak{B}_0$ contained in the bad cubes; the bound $\|\Theta_p\| \le 1.01\,\delta(U_p, I)$ applies since $\rho \le \frac{1}{10}$. For the limits take $r = \lceil 8 \log_2 L \rceil$, so that $L^4 2^{-r} \le L^{-4}$ and $ra \to 0$ for a torus of fixed physical side La. The defects then lie in sets of physical diameter at most $ra + 6a_{k_*}$, and $a_{k_*} \le 10^{-2}(1.01\rho)^{-1/2} a \to 0$. □

The scale a_{k_*} is of the order $(\beta_W/\log\beta_W)^{1/4}a$, the order of the cube side that the condition (D) needs by Proposition 11.17. So the cut-off average carries the field through the defects of the Wilson measure, without spreading them, up to the scale at which (D) can first hold, but not beyond: past a_{k_*} the bound $\eta_j \le 1.023 \cdot 4^j \eta_0$ no longer keeps the curvature small. Along the one-loop tuning β_W grows like $\log L$, so k_* grows like $\frac{1}{4}\log_2 \log L$, a vanishing fraction of the $\log_2 L$ scales.

The joint bound of Theorem 12.57 is not a conditional one, and (D) needs a supremum over boundary conditions. The two differ genuinely. Fix a link e and a plaquette p_0 through it. Put all links on the six staples of e at I, except the middle links of the five staples other than that of p_0, which are $-I$. Then the conditional density of U_e is proportional to $e^{-2\beta_W \operatorname{Tr} U_e}$, and p_0, whose holonomy is $U_e^{\pm 1}$, is bad with conditional probability tending to 1 as $\beta_W \to \infty$, for every $\rho < 2\sqrt{2}$. Nearby exteriors behave the same way. So the clusters of Corollary 12.58 are small in probability, but the boundary response of a cube to a frustrated exterior is not small, and the composition beyond the scale a_{k_*} of Corollary 12.63 and the conditional estimate in (D) remain open.

For exteriors that are calm on the boundary of a box and lie in a maximal torus, the conditional estimate does hold. Let $R = [0, l]^4$, $l \ge 2$, be a box of Λ_L with $l < L$. Its boundary ∂R is a cubical 3-sphere. Call a link or plaquette of R a *boundary* link or plaquette if it lies in ∂R, and *inner* otherwise. A plaquette that contains an inner link lies in R, so conditionally on all other links the inner links have the law

$$\nu_\eta(dU) = Z_\eta^{-1} \exp\Big(-\beta_W \sum_{p \in P^\circ} s_p\Big) \prod_e dU_e,$$

the product over the inner links, where P° is the set of inner plaquettes. It depends on the exterior η only through the boundary links. Let $T \subset \mathrm{SU}(2)$ be the diagonal torus of the elements $\exp(tJ)$, $J = \mathrm{diag}(i, -i)$, and for $u \in \mathrm{SU}(2)$ let $\angle u \in [0, \pi]$ be its angle, so that $\delta(u, I) = 2\sqrt{2}\sin(\angle u/2)$.

Lemma 12.64. *Let $0 \le \rho_s < \pi/3$. Suppose the boundary links of R lie in T and $\angle U_p \le \rho_s$ for every boundary plaquette. Then the inner links can be chosen in T so that $\angle U_p \le C\rho_s$ for every plaquette of R, where*

$$C = \sqrt{3}\,(24 \cdot 13118 + 9) < 5.46 \cdot 10^5.$$

Proof. Write a T-valued field as $U_e = \exp(a_e J)$ with $a_e \in \mathbb{R}/2\pi\mathbb{Z}$. For a plaquette p let $(da)_p$ be the signed sum of a over its four links, so that $U_p = \exp((da)_p J)$. For a boundary plaquette let $\theta_p \in (-\pi, \pi]$ be the representative of $(da)_p$; then $|\theta_p| = \angle U_p \le \rho_s$.

Closedness. The signed angles of the six plaquettes of a unit 3-cube of ∂R add up to 0 modulo 2π, since each link occurs twice with opposite signs. Their representatives add up to a number of modulus at most $6\rho_s < 2\pi$, hence to 0. So θ is a closed real 2-cochain on ∂R.

Lift. We construct a real 1-cochain α on the boundary links with $d\alpha = \theta$ and $|\alpha_e| \leq 13118\, l\rho_s$. The boundary is the union of the eight facets $F_i^0 = \{x \in R : x_i = 0\}$ and $F_i^l = \{x \in R : x_i = l\}$, each a cube $[0, l]^3$. Take them in the order F_1^l, F_2^l, F_2^0, F_3^l, F_3^0, F_4^l, F_4^0, F_1^0, and let Γ_k be the union of the intersections of the k-th facet with the earlier ones. In the cube of the k-th facet, Γ_k is empty for $k = 1$, one face for $k = 2, 3$, three faces, two of them opposite, for $k = 4, 5$, five faces for $k = 6, 7$ and all six for $k = 8$. So Γ_k is contractible or a 2-sphere, and its first cellular cohomology vanishes.

On a facet with coordinates t_1, t_2, t_3, the axial gauge gives α^0 with $d\alpha^0 = \theta$ on the facet. Put $\alpha^0 = 0$ on the links in direction t_3. Determine α^0 on the links in direction t_2 from the plaquettes in the planes $\{t_2, t_3\}$, starting from 0 at $t_3 = 0$. On the links in direction t_1 at $t_3 = 0$, determine it from the plaquettes in the plane $\{t_1, t_2\}$, starting from 0 at $t_2 = 0$, and then from the plaquettes in the planes $\{t_1, t_3\}$. The plaquettes in the planes $\{t_1, t_2\}$ with $t_3 > 0$ then satisfy $d\alpha^0 = \theta$ by closedness, by induction on t_3. Each value is a signed sum of at most l, respectively $2l$, angles, so $|\alpha^0| \leq 2l\rho_s$.

For $k = 1$ put $\alpha = \alpha^0$, so $M_1 = \max|\alpha| \leq 2l\rho_s$. For $k \geq 2$, α is already defined on the links of Γ_k, with $|\alpha| \leq M_{k-1}$, and $\omega = \alpha - \alpha^0$ is closed there. Hence $\omega = d\lambda$ for a function λ on the vertices of Γ_k, and $|\lambda(x) - \lambda(y)| \leq \|\omega\|_\infty d_\Gamma(x, y)$, where d_Γ is the path metric of Γ_k.

We claim $d_\Gamma \leq 3d_1$, where d_1 is the ℓ^1 metric of the facet. If x and y lie in one face of Γ_k, that face contains an ℓ^1 geodesic between them. If they lie in adjacent faces $\{t_i = s\}$ and $\{t_j = s'\}$, the path through the point with $t_i = s$, $t_j = s'$ and third coordinate that of x has length $d_1(x, y)$. If they lie in opposite faces $\{t_i = 0\}$ and $\{t_i = l\}$, then $d_1(x, y) \geq l$. In that case Γ_k contains a face adjacent to both, and the path from x to that face, across it and on to y has length at most $d_1(x, y) + 2l \leq 3d_1(x, y)$.

So λ is $3\|\omega\|_\infty$-Lipschitz for d_1 on Γ_k. The function

$$\tilde{\lambda}(x) = \min_y(\lambda(y) + 3\|\omega\|_\infty d_1(x, y))$$

extends it to the vertices of the facet with the same constant [62]. Put $\alpha = \alpha^0 + d\tilde{\lambda}$ on the links of the facet. It agrees with the earlier values on Γ_k, satisfies $d\alpha = \theta$ on the facet, and has

$$|\alpha| \leq 2l\rho_s + 3(M_{k-1} + 2l\rho_s).$$

So $M_k \leq 3M_{k-1} + 8l\rho_s$, which gives $M_8 \leq 13118\, l\rho_s$.

Extension. Let c be the centre of R, $R_0 = l/2$, $r(x) = |x - c|_\infty$ and $\varphi(x) = c + R_0(x - c)/r(x)$, the radial projection of $R \setminus \{c\}$ onto ∂R. Let $\hat{\alpha}$ and $\hat{\theta} = d\hat{\alpha}$ be the lowest-order Whitney forms of α and θ on the unit cubes of ∂R [82]. On a unit cube with coordinates t_1, t_2, t_3, $\hat{\alpha} = \sum_i a_i\, dt_i$, where a_i is the multilinear interpolation of the values of α on the four links in direction t_i; similarly $\hat{\theta} = \sum_{i<j} \theta_{ij}\, dt_i \wedge dt_j$. These forms have continuous tangential components across the faces of the unit cubes, satisfy $\int_e \hat{\alpha} = \alpha_e$ for every boundary link, and for tangent vectors

$$|\hat{\alpha}(u)| \leq \sqrt{3}\, M_8|u|, \qquad |\hat{\theta}(u, w)| \leq \sqrt{3}\, \rho_s|u||w|.$$

Put $\tilde{B} = (r/R_0)^2\varphi^*\hat{\alpha}$ on $R \setminus \{c\}$. Since $|\nabla r| = 1$ almost everywhere and $|x - c| \leq 2r$, we have $|d\varphi(v)| \leq 3R_0|v|/r$. Then

$$d\tilde{B} = d(r^2/R_0^2) \wedge \varphi^*\hat{\alpha} + (r/R_0)^2\varphi^*\hat{\theta}$$

gives

$$|d\tilde{B}(v, w)| \leq (12\sqrt{3}\, M_8/R_0 + 9\sqrt{3}\, \rho_s)|v||w| \leq C\rho_s|v||w|,$$

and $|\tilde{B}| \leq 3\sqrt{3}\, M_8 r/R_0$ is bounded.

The set $R \setminus \{c\}$ is the union of the cones from c over the unit cubes of ∂R. These cones are convex polytopes cut out by hyperplanes through c. On each of them $\tilde{B}$ is smooth, and

its tangential components are continuous across them since φ is continuous. Put $B_e = \int_e \tilde{B}$ for every link e of R. Apply Stokes' theorem on each piece of a plaquette: the boundary terms on the interior cuts cancel, and the point c contributes nothing since $\tilde{B}$ is bounded. This gives $(dB)_p = \int_p d\tilde{B}$, so $|(dB)_p| \le C\rho_s$. On ∂R, $r = R_0$ and φ is the identity on tangent vectors, so $B = \alpha$ on the boundary links.

Gauge. The cochain $a - \alpha$ on the boundary links has $(d(a-\alpha))_p \equiv \theta_p - \theta_p = 0$ modulo 2π on every boundary plaquette. So it is a flat U(1)-field on the 2-skeleton of ∂R, which is simply connected, and $a - \alpha \equiv d\mu$ modulo 2π for a function μ on the boundary vertices. Extend μ by 0 to the inner vertices and put $U_e = \exp((B_e + (d\mu)_e)J)$ on the inner links. On the boundary links $B + d\mu \equiv a$, and every plaquette of R has angle at most $|(dB)_p| \le C\rho_s$. $\square$

For a general SU(2)-valued boundary field the curvature is not a closed cochain, and the lift in the proof of Lemma 12.64 is not available. A lift by gluing local gauges still works when $l^2\rho_s$ is small. Identify $\mathfrak{su}(2)$ with the imaginary quaternions and write $|X|$ for the operator norm, which is the Euclidean norm, so that $|[X,Y]| \le 2|X||Y|$, $\|e^X - I\| \le |X|$ and $\angle e^X = |X|$ for $|X| \le \pi$. With $d(g,h) = \angle(g^{-1}h)$, the map $X \mapsto e^X$ is 1-Lipschitz, since its differential has singular values 1 and $\sin|X|/|X|$, and the principal logarithm is 2-Lipschitz on $\{\angle h \le \frac{1}{2}\}$, a geodesically convex ball on which the inverse differential has singular values at most $\frac{1}{2}/\sin\frac{1}{2} < 2$. Put

$$C_1 = 2, \qquad C_k = 11\,C_{k-1} + 24 \quad (2 \le k \le 8),$$

so that $C_8 = 85\,743\,550$, and

$$K = 42\,C_8 + 20, \qquad \varepsilon_0 = \frac{1}{100\,C_8^2}.$$

Lemma 12.65. *Let the boundary links η of R take values in* SU(2)*, with $\angle\eta_p \le \rho_s$ for every boundary plaquette and $l^2\rho_s \le \varepsilon_0$. Then the inner links can be chosen so that $\angle U_p \le K\rho_s$ for every plaquette of R.*

Proof. For a vertex gauge g write $\eta^g_{xy} = g_x \eta_{xy} g_y^{-1}$. A product of conjugates of m elements of angle at most ρ_s has angle at most $m\rho_s$, since d is bi-invariant.

One facet. On a facet with coordinates t_1, t_2, t_3 let P_x be the path from the corner to x that moves first in direction t_1, then t_2, then t_3, and put $g^0_x = \eta(P_x)$, the ordered product of the links along P_x. Then $\eta^{g^0}_{xy}$ is the holonomy of the loop $P_x\,(xy)\,P_y^{-1}$. For a link in direction t_3 this loop retraces itself. For a link in direction t_2 at height $t_3 = c$ it bounds c plaquettes of the planes $\{t_2, t_3\}$, and for a link in direction t_1 at $(t_2, t_3) = (b, c)$ it bounds b plaquettes of a plane $\{t_1, t_2\}$ and c of a plane $\{t_1, t_3\}$. So the holonomy is a product of conjugates of at most $2l$ plaquette holonomies, and every link of the facet has $\angle\eta^{g^0}_e \le 2l\rho_s$.

Gluing. Take the facets in the order of the proof of Lemma 12.64, with the same sets Γ_k; there $d_\Gamma \le 3d_1$. Two vertices of Γ_k joined by a link of the facet lie in a common face of Γ_k, since $l \ge 2$, so that link lies in Γ_k. We construct a gauge g on the boundary vertices with $\angle\eta^g_e \le M_k = C_k l\rho_s$ on the links of the first k facets. For $k = 1$ take $g = g^0$. For $k \ge 2$, let g^0 be the gauge of the k-th facet and $t_x = g_x(g^0_x)^{-1}$ for $x \in \Gamma_k$. For a link xy of Γ_k, $\eta^g_{xy} = t_x \eta^{g^0}_{xy} t_y^{-1}$, so

$$d(t_x, t_y) = \angle(t_x^{-1}t_y) \le \angle\eta^g_{xy} + \angle\eta^{g^0}_{xy} \le M_{k-1} + 2l\rho_s,$$

and t is $3(M_{k-1} + 2l\rho_s)$-Lipschitz for d_1 on Γ_k. Fix $x_0 \in \Gamma_k$ and put $h_x = t_{x_0}^{-1} t_x$. The facet has d_1-diameter $3l$, so

$$\angle h_x \le 9l(M_{k-1} + 2l\rho_s) = 9(C_{k-1} + 2)\,l^2\rho_s \le 9(C_8 + 2)\varepsilon_0 < \tfrac{1}{2},$$

and $a_x = \log h_x$ is L-Lipschitz for d_1, with $L = 6(M_{k-1} + 2l\rho_s)$. Extend each of the three real coordinates of a to the vertices of the facet by the formula of McShane [62], as in the

proof of Lemma 12.64; the extension $\tilde{a}$ is $\sqrt{3}L$-Lipschitz. Put $g_x = t_{x_0} e^{\tilde{a}_x} g^0_x$ on the facet; on Γ_k this is the old value. For a link xy of the facet, $\eta^g_{xy} = \tilde{t}_x \eta^{g^0}_{xy} \tilde{t}_y^{-1}$ with $\tilde{t}_x = t_{x_0} e^{\tilde{a}_x}$, so

$$\begin{aligned} \angle \eta^g_{xy} &\le d(\tilde{t}_x, \tilde{t}_y) + 2l\rho_s \le \sqrt{3}L + 2l\rho_s \\ &= 6\sqrt{3}\, M_{k-1} + (12\sqrt{3}+2)\, l\rho_s \le M_k. \end{aligned}$$

After the eighth facet, $\angle \eta^g_e \le M = C_8 l \rho_s$ on every boundary link.

Potential. Since $M \le C_8\varepsilon_0/l < \frac{1}{2}$, the logarithms $A_e = \log \eta^g_e$ are defined, with $|A_e| \le M$ and $A_{-e} = -A_e$. Let $(dA)_p$ be the signed sum of A over the four links of a boundary plaquette. For $X_1, \dots, X_q \in \mathfrak{su}(2)$,

$$\Big\| e^{X_1} \cdots e^{X_q} - I - \sum_i X_i \Big\| \le \tfrac{1}{2} \Big(\sum_i |X_i| \Big)^2,$$

by induction on q from $e^X e^Y - I - X - Y = (e^X - I - X) + (e^Y - I - Y) + (e^X - I)(e^Y - I)$ and $\|e^X - I - X\| \le |X|^2/2$. Applied to the four links of p, and since $\|\eta^g_p - I\| = 2\sin(\angle \eta_p/2) \le \rho_s$, this gives

$$|(dA)_p| \le \sigma = \rho_s + 8M^2.$$

Let $\hat{A}$ be the lowest-order Whitney form of A on the unit cubes of ∂R, as in the proof of Lemma 12.64, now with values in $\mathfrak{su}(2)$. Its coefficients are interpolations of values of A, and those of $d\hat{A}$ are interpolations of values of dA, so

$$|\hat{A}(u)| \le \sqrt{3}\, M |u|, \qquad |d\hat{A}(u,w)| \le \sqrt{3}\, \sigma |u||w|.$$

With r, R_0 and φ as there, put $\tilde{B} = (r/R_0)^2 \varphi^* \hat{A}$ on $R \setminus \{c\}$ and $\tilde{B}(c) = 0$. The same computation gives

$$|\tilde{B}(v)| \le 3\sqrt{3}\, M \frac{r}{R_0} |v|, \qquad |d\tilde{B}(v,w)| \le \kappa_1 |v||w|,$$

where $\kappa_1 = 12\sqrt{3}\, M/R_0 + 9\sqrt{3}\, \sigma$. On each cone from c over a unit cube of ∂R, $\tilde{B}$ is smooth away from c, and its tangential components are continuous across the cones.

Transport. For a link e of R, parametrised by $s \in [0,1]$, let $V_e = T(1)$, where $T' = T\, \tilde{B}(\dot{e})$ and $T(0) = I$; the coefficient is bounded and piecewise smooth, and V composes along paths in the order of the links. For a closed path γ of length ℓ,

$$T(1) - I - \int_\gamma \tilde{B} = \int_0^1 (T - I)\, \tilde{B}(\dot{\gamma})\, ds,$$

and $\|T(s) - I\| \le \int_0^s |\tilde{B}(\dot{\gamma})|$, so the holonomy V_γ satisfies $\|V_\gamma - I\| \le |\int_\gamma \tilde{B}| + \frac{1}{2}(\int_\gamma |\tilde{B}|)^2$. Let p be a plaquette of R and divide it into N^2 squares q of side $h = 1/N$. Its holonomy is a product of conjugates of the holonomies of the boundaries of the squares, since every inner edge of the subdivision is traversed once in each direction. For each square, Stokes' theorem on the finitely many convex polygons in which the cones cut q gives $\int_{\partial q} \tilde{B} = \int_q d\tilde{B}$: the integrals over the interior cuts cancel because the tangential components agree, and if $c \in q$ a disc of radius δ around c contributes $O(\delta^2)$, since $|\tilde{B}| \le 3\sqrt{3}\, Mr/R_0$. Hence

$$\|V_{\partial q} - I\| \le \kappa_1 h^2 + \tfrac{1}{2}\big(4h \cdot 3\sqrt{3}\, M\big)^2 = \kappa h^2,$$

with $\kappa = \kappa_1 + 216M^2$. Since $2\arcsin(x/2) \le x(1 + x^2)$ for $0 \le x \le 1$, summing over the squares gives $\angle V_p \le \kappa(1 + \kappa^2 N^{-4})$, and letting $N \to \infty$, $\angle V_p \le \kappa$. Now $M/R_0 = 2C_8\rho_s$ and $M^2 = C_8^2 l^2 \rho_s^2 \le \rho_s/100$, so

$$\begin{aligned} \kappa &\le (24\sqrt{3}\, C_8 + 9\sqrt{3})\rho_s + (72\sqrt{3} + 216) M^2 \\ &< (42\, C_8 + 20)\, \rho_s = K\rho_s. \end{aligned}$$

Boundary values. On ∂R, $r = R_0$ and φ is the identity on tangent vectors, and along a boundary link e the tangential coefficient of $\hat{A}$ is the constant A_e. So $V_e = e^{A_e} = \eta^g_e$.

Extend g by I to the inner vertices and put $U_{xy} = g_x^{-1} V_{xy} g_y$. Then $U = \eta$ on the boundary links, and U_p is conjugate to V_p, so $\angle U_p \le K\rho_s$ for every plaquette of R. □

The boundary fields of Lemma 12.65 with strict inequalities form an open set of positive Haar measure that contains the identity field; unlike those of Lemma 12.64, they are not a null set. The constants are far from optimal.

For $0 < \varepsilon \le 1$ let E_ε be the event that at least εl^4 of the l^4 unit cubes $x + [0,1]^4$, $x \in \{0, \dots, l-1\}^4$, of R contain a bad plaquette.

Proposition 12.66. *Let $\beta_W \ge 1$ and $0 < \rho \le 1$. Suppose that no boundary plaquette of R is bad under η, and that the inner links can be chosen so that every inner plaquette has angle at most κ. Then*

$$-\log \nu_\eta(E_\varepsilon) \ge \frac{\beta_W \rho^2 \varepsilon}{16}\, l^4 - 6(l+1)^4 \big(\log \beta_W + C_0 + \tfrac{1}{2} + \beta_W \kappa^2\big),$$

with C_0 as in Lemma 12.56. In particular, if the boundary links lie in T with angle at most $\rho_s < \pi/3$ on every boundary plaquette and $\rho \ge \sqrt{2}\,\rho_s$, the bound holds with $\kappa = C\rho_s$ and the constant C of Lemma 12.64. If the boundary links lie in $\mathrm{SU}(2)$ with angle at most ρ_s on every boundary plaquette, $l^2\rho_s \le \varepsilon_0$ and $\rho \ge \sqrt{2}\,\rho_s$, it holds with $\kappa = K\rho_s$.

Proof. On E_ε the bad plaquettes are inner, and each lies in at most four of the unit cubes. So there are at least $\varepsilon l^4/4$ of them, each with $s_p > \rho^2/4$, and $\sum_{P^\circ} s_p \ge \rho^2 \varepsilon l^4/16$. Hence the unnormalised weight of E_ε is at most $\exp(-\beta_W \rho^2 \varepsilon l^4/16)$.

For the normalisation let $\bar U$ be the inner links of the hypothesis, $t = (4\sqrt{\beta_W})^{-1}$, and consider the inner links $U_e = \bar U_e h_e$ with $\angle h_e \le t$. The angle of $g^{-1}h$ is a bi-invariant distance, so every inner plaquette has $\angle U_p \le \kappa + 4t$ and

$$s_p \le \tfrac{1}{2}(\kappa + 4t)^2 \le \kappa^2 + 16t^2 = \kappa^2 + \beta_W^{-1}.$$

Each h_e has probability at least $8t^3/(3\pi^3) = (24\pi^3 \beta_W^{3/2})^{-1}$, as in the proof of Lemma 12.56. There are $n_P \le 6l^2(l+1)^2$ inner plaquettes and $n_L \le 4l(l+1)^3$ inner links, so

$$Z_\eta \ge \exp\big(-n_P(\beta_W \kappa^2 + 1)\big)\big(24\pi^3 \beta_W^{3/2}\big)^{-n_L}.$$

Since $C_0 = \frac{1}{2} + \frac{2}{3}\log(24\pi^3)$, $n_P + n_L \log(24\pi^3 \beta_W^{3/2}) \le 6(l+1)^4(\log\beta_W + C_0 + \frac{1}{2})$, which gives the bound. For the last assertion, a boundary plaquette has $\delta(U_p, I) \le \sqrt{2}\,\angle U_p \le \sqrt{2}\,\rho_s \le \rho$, so it is not bad, and Lemma 12.64 or Lemma 12.65 gives the inner links. □

So in a box whose boundary is calm at a level ρ_s small compared with $\rho\sqrt{\varepsilon}/C$ and T-valued, or small compared with $\rho\sqrt{\varepsilon}/K$ and $\mathrm{SU}(2)$-valued with $l^2\rho_s \le \varepsilon_0$, the fraction of bad cubes exceeds ε only with probability exponentially small in the volume once $\beta_W \rho^2 \varepsilon$ is large compared with $\log \beta_W$, uniformly over such exteriors and over l. The estimate is conditional, unlike Theorem 12.57.

Remark 12.67. The exteriors with boundary links in T form a null set for μ_L. Lemma 12.65 removes the torus at the price of the smallness $l^2\rho_s \le \varepsilon_0$. In the continuum the corresponding condition is that of Uhlenbeck's theorem [78], which gives a Coulomb gauge on a ball when the curvature is small in $L^{n/2}$; on 3-dimensional pieces of ∂R of side comparable to l and for curvature of size ρ_s this norm is of order $l^2\rho_s$. The lemma replaces the Coulomb gauge by the gluing of the facet gauges. With ρ_s of the order of the level $\rho(\beta_W)$ of Corollary 12.63, $\rho_s = c_0(\log\beta_W/\beta_W)^{1/2}$, it applies for

$$l \le (\varepsilon_0/c_0)^{1/2}\,(\beta_W/\log\beta_W)^{1/4},$$

the scale a_{k_*} of Corollary 12.63 up to a constant factor. Along the tuning $\beta_W \sim b\log(1/a)$ the physical side al then tends to 0, so the lemma does not reach cubes of a fixed physical size; this is a limitation of the argument, not a proof that such fillings fail. Moreover

Proposition 12.66 bounds the fraction of bad cubes under one exterior and does not compare the laws under two exteriors, so it gives no contraction coefficient.

The calm boundary is needed only for the small-curvature filling. If the boundary layer is paid for instead of filled, every condition on the exterior disappears, at the price of a surface term that is small against the volume on large boxes.

Proposition 12.68. *Let $\beta_W \geq 1$, $0 < \rho \leq 1$ and $0 < \varepsilon \leq 1$ with $\varepsilon l \geq 16$. For every boundary field η,*

$$-\log \nu_\eta(E_\varepsilon) \geq l^4\Big(\frac{\beta_W \rho^2 \varepsilon}{32} - 6\log \beta_W - C_1 - \frac{48\beta_W}{l}\Big),$$

with $C_1 = 3 + 4\log(24\pi^3) < 31$.

Proof. A unit cube $x + [0,1]^4$ with $x \in \{1, \dots, l-2\}^4$ does not meet ∂R, so all its plaquettes are inner; the other $l^4 - (l-2)^4 \leq 8l^3 \leq \varepsilon l^4/2$ unit cubes meet it. On E_ε at least $\varepsilon l^4/2$ cubes of the first kind contain a bad plaquette, which is inner, and a plaquette lies in at most four unit cubes. So at least $\varepsilon l^4/8$ inner plaquettes are bad, $\sum_{P^\circ} s_p \geq \rho^2 \varepsilon l^4/32$, and the unnormalised weight of E_ε is at most $\exp(-\beta_W \rho^2 \varepsilon l^4/32)$.

For the normalisation take inner links $U_e = h_e$ with $\angle h_e \leq t = (4\sqrt{\beta_W})^{-1}$. An inner plaquette whose four links are inner has $\angle U_p \leq 4t$ and $s_p \leq (2\beta_W)^{-1}$. An inner plaquette spans two directions and has its other two coordinates in $\{1, \dots, l-1\}$; it contains a boundary link exactly when its cell in the plane of the two spanned directions meets the boundary of $[0,l]^2$, which happens for $4(l-1)$ of the l^2 cells. So $24(l-1)^3 \leq 24l^3$ inner plaquettes contain a boundary link, and on them $s_p \leq 2$. There are $n_P = 6l^2(l-1)^2 \leq 6l^4$ inner plaquettes and $n_L = 4l(l-1)^3 \leq 4l^4$ inner links, so, as in the proof of Proposition 12.66,

$$Z_\eta \geq \exp\big(-48\beta_W l^3 - 3l^4\big)\big(24\pi^3 \beta_W^{3/2}\big)^{-4l^4}.$$

Dividing gives the bound, and $24\pi^3 < e^7$ gives $C_1 < 31$. □

Corollary 12.69. *Let $\beta_W \geq e^{44}$, $l \geq 6144\beta_W^{3/4}$ and $\rho = \varepsilon = \beta_W^{-1/4}$. Then $\nu_\eta(E_\varepsilon) \leq \exp(-\beta_W^{1/4} l^4/64)$ for every boundary field η. In particular, along $\beta_W \sim b\log(1/a)$ with $l = \lceil r/a \rceil$ and fixed $r, b > 0$, on cubes of fixed physical side r the fraction of unit cubes that contain a plaquette with $\delta(U_p, I) > \beta_W^{-1/4}$ is at most $\beta_W^{-1/4}$ with conditional probability tending to one, uniformly over all exteriors.*

Proof. Here $\varepsilon l \geq 6144\beta_W^{1/2} \geq 16$ and $\rho \leq 1$. The first term of Proposition 12.68 is $\beta_W^{1/4}/32$, and $48\beta_W/l \leq \beta_W^{1/4}/128$. For $x = \log \beta_W \geq 44$ we have $6x + 31 \leq e^{x/4}/128$: at $x = 44$ the left side is 295 and the right side exceeds 467, and $x/4 - \log(6x+31)$ increases for $x \geq 44$. So $6\log\beta_W + C_1 \leq \beta_W^{1/4}/128$ and the bracket is at least $\beta_W^{1/4}(\frac{1}{32} - \frac{1}{128} - \frac{1}{128}) = \beta_W^{1/4}/64$. Along the tuning, $\beta_W \to \infty$ and $l/\beta_W^{3/4} \to \infty$, so the hypotheses hold for small a. □

Like Proposition 12.66, Corollary 12.69 bounds the bad cubes under one exterior and does not compare the laws under two, so it gives no contraction coefficient. The bad cubes are rare on cubes of fixed physical size for every exterior, and the conditional estimate in (D) remains open.

The same count locates the windows on which the fibre estimates of Sec. 12.4 apply. Let the side of the torus be even, let $B(y) = 2y + \{0,1\}^4$ be the blocks of Definition 12.2, and let l be even, so that R is a union of blocks.

Corollary 12.70. *Let $G = \mathrm{SU}(2)$, $\beta_W \geq 1$, let W_0 be a set of m blocks, $0 < K \leq 10^{-8}$, $\rho = K/(3072m)$, and $0 < \varepsilon \leq 1$ with $\varepsilon l \geq 16$. For every boundary field η of R, with ν_η-probability at least*

$$1 - \exp\Big(-l^4\Big(\frac{\beta_W \rho^2 \varepsilon}{32} - 6\log\beta_W - 31 - \frac{48\beta_W}{l}\Big)\Big),$$

all but fewer than $256m\varepsilon l^4$ *of the translates* $W = y_0 + W_0$ *whose blocks of* W^+ *lie in* $[1, l-1]^4$ *have the following property: in the tree gauge the exterior of* W *has a completion that satisfies the hypothesis of Corollary 12.54, namely the field itself.*

Proof. Outside E_ε, which has the stated probability by Proposition 12.68, fewer than εl^4 unit cubes of R contain a bad plaquette. A plaquette with a vertex in a block of W^+ has all its vertices in R and lies in a unit cube $x + [0,1]^4$ of R; if it is bad, that cube contains a bad plaquette. The vertex lies in a block $B(y')$ with $y'_i \in \{\lfloor x_i/2 \rfloor, \lfloor (x_i+1)/2 \rfloor\}$, and $y' \in \{y\}^+$ for some $y \in W$, so y_i takes at most four values and y lies in a set of at most 256 blocks determined by x; then $y_0 \in y - W_0$ takes at most $256m$ values. Hence fewer than $256m\varepsilon l^4$ translates have a bad plaquette with a vertex in a block of W^+. At every other translate these plaquettes have $\delta(U_p, I) \le \rho$. The number n_P of plaquettes that contain a link of A_W is at most $384m$, so $\rho \le K/(8n_P) = \kappa/(2\sqrt{n_P})$ with $\kappa = K/(4\sqrt{n_P})$. A gauge transformation conjugates every plaquette holonomy and leaves $\delta(U_p, I)$ unchanged, so the field in the tree gauge is a completion of its own exterior with this property. □

So under every exterior of a box, the windows of a fixed number of blocks on which Corollary 12.54 applies fill all but at most $256m\varepsilon l^4$ of the positions, a fraction of about $4096m\varepsilon$. The bound is useful only when $\beta_W \rho^2 \varepsilon$ is large, that is for windows of bounded size, and the next proposition shows that this restriction cannot be removed.

Proposition 12.71. *Let* $G = \mathrm{SU}(2)$, $\beta_W \ge 1$ *and* $0 < \rho' \le 1$.

(i) *Conditionally on all links but one link of a plaquette* p_0, *the event* $\delta(U_{p_0}, I) \le \rho'$ *has probability at most* $14(\beta_W \rho'^2)^{3/2}$ *under the Wilson measure.*

(ii) *Let* W *be a set of blocks with* $W^+ \ne W$, $0 < K \le 10^{-8}$ *and* $\kappa = K/(4\sqrt{n_P})$. *Under the Wilson measure in the tree gauge, the exterior of* W *has a completion as in Corollary 12.54 with probability at most* $7 \cdot 10^{-32} \beta_W^{3/2} |W|^{-3}$, *and* $\varepsilon''_W(\kappa^2/2, \beta_W/4) < 1$ *only if* $\beta_W > 10^{24}|W|^2$.

Proof. (i) As in the proof of Proposition 12.60, the holonomy of p_0 has a conditional density proportional to $e^{\kappa' h \cdot u}$ with $0 \le \kappa' \le 6\beta_W$ and $|h| = 1$, and the probability of a cap about I is largest at $h = I$, $\kappa' = 6\beta_W$. Since $\delta(u, I) = 2\sqrt{2}\sin(\angle u/2)$ and $\sin x \ge 2x/\pi$ on $[0, \frac{\pi}{2}]$, the event forces $\angle U_{p_0} \le \theta = \pi\rho'/(2\sqrt{2})$. With $\cos t - 1 \ge -t^2/2$ and $\sin t \ge 2t/\pi$ on $[0, \frac{\pi}{2}]$,

$$\int_0^\pi e^{6\beta_W(\cos t - 1)} \sin^2 t \, dt \ge \frac{4}{\pi^2} \int_0^{\pi/2} e^{-3\beta_W t^2} t^2 \, dt \ge \frac{4 \cdot 0.44}{\pi^2 (3\beta_W)^{3/2}},$$

because the integral of $e^{-s^2} s^2$ over $[0, \pi\sqrt{3}/2]$ exceeds 0.44. The integral of $e^{6\beta_W(\cos t - 1)} \sin^2 t$ over $[0, \theta]$ is at most $\theta^3/3$, so the ratio is at most $9.72(\beta_W \theta^2)^{3/2}$, and $(\pi/(2\sqrt{2}))^3 < 1.371$.

(ii) Take $y' \in W^+ \setminus W$ and a plaquette p_0 inside $B(y')$. Its links have their base in $B(y')$, so none lies in A_W, and every completion of the exterior agrees with the field on p_0. A completion as in Corollary 12.54 therefore needs $\delta(U_{p_0}, I) \le \kappa/(2\sqrt{n_P}) = K/(8n_P)$. This event is gauge invariant, so its probability is the same without gauge fixing, and by (i) it is at most $14(\beta_W K^2/(64 n_P^2))^{3/2}$. Each link of A_W lies in six plaquettes and a plaquette contains at most four of them, so $n_P \ge 73.5|W|$, and $14 \cdot 10^{-24}/(512 \cdot 73.5^3) < 7 \cdot 10^{-32}$. Finally, $\varepsilon''_W(\kappa^2/2, \lambda) < 1$ means $\frac{3}{8}\lambda\kappa^2 > 49|W| \ln(3.45 \cdot 10^{11} n_P^6/K^6)$, and with $\lambda = \beta_W/4$, $\kappa^2 = K^2/(16 n_P)$, $n_P \ge 73.5|W|$ and $K \le 10^{-8}$ the logarithm exceeds 162.8, so $\beta_W > 512 \cdot 49 \cdot 73.5 \cdot 162.8 \cdot 10^{16}|W|^2/3 > 10^{24}|W|^2$. □

Along the tuning $\beta_W \sim b \log(1/a)$, a cube of fixed physical side r has $|W| = (r/2a)^4$ blocks, so both bounds of (ii) fail for small a: on such a cube the exterior has no calm completion with probability tending to one, and the fibre estimates of Sec. 12.4 are vacuous there. Their entropy factors are powers of $|W|$, so they are estimates for windows of

bounded size, which Corollary 12.70 places at almost every position under every exterior. The exteriors with no calm completion and the rare fibres of Corollary 12.54 are thus the typical case at the physical scale and not an exceptional one. The cube condition (D) needs estimates on the whole cube that are uniform in $|W|$, with the windows that are not calm split off as in Bałaban's analysis [8], and that remains open.

12.6. **Conditioning on a fixed large-field set.** The estimates of Section 12.5 are statements about the Wilson measure on the torus. The cube condition (D) takes a supremum over exteriors, and three facts about conditional laws hold under every exterior; the setting is that of a Gibbs field whose potential is itself random, the uniqueness theory of which goes back to Bassalygo and Dobrushin [3]. The first bounds the excess of the action over its minimum in a box, with no reflection positivity and no torus. The second says that restricting a single-link law to a fixed set, however irregular, leaves its Kantorovich response to the neighbouring links bounded by a constant times β_W. The third turns the one-point rarity of a bad set into closeness of any two Gibbs measures, at order of that rarity, and shows at the same time why one-point rarity gives closeness and not decay.

Lemma 12.72 (excess action)**.** *Let $G = \mathrm{SU}(2)$, let Λ be a set of N links of $\mathbb{Z}^4$ and fix the links outside Λ. Let P_Λ be the plaquettes that contain a link of Λ, $S_\Lambda = \sum_{p \in P_\Lambda} s_p$, and $S_{\min}$ the minimum of S_Λ over the links of Λ. For every $E > 0$, under the Wilson measure conditioned on the links outside Λ,*

$$\Pr(S_\Lambda \geq S_{\min} + E) \leq \Big(\frac{3\pi^3}{8r^3}\Big)^N e^{-\beta_W E/2},$$

where $r = \min\{E/(6\pi N), \pi/2\}$.

Proof. The plaquettes that contain no link of Λ do not depend on the links of Λ, so the conditional law has density $Z^{-1} e^{-\beta_W S_\Lambda}$ with respect to the product of Haar measures on Λ. The unnormalised mass of the event is at most $e^{-\beta_W (S_{\min}+E)}$. For the lower bound on Z, let U^* be a minimiser, which exists by compactness. If a link of a plaquette p moves by geodesic distance t, the angle $\delta(U_p, I) \in [0, \pi]$ moves by at most t, by bi-invariance, so $s_p = \frac{1}{4}\delta(U_p, I)^2$ moves by at most $\frac{1}{4}\, t \cdot 2\pi = \frac{\pi}{2} t$. Each link lies in six plaquettes. If every link of Λ lies within distance r of its value in U^*, then $S_\Lambda \leq S_{\min} + 6 \cdot \frac{\pi}{2} r N \leq S_{\min} + E/2$. The Haar measure of a geodesic ball of radius $r \leq \pi/2$ is $\frac{2}{\pi} \int_0^r \sin^2 t \, dt \geq 8r^3/(3\pi^3)$, by $\sin t \geq 2t/\pi$. Hence $Z \geq (8r^3/3\pi^3)^N e^{-\beta_W (S_{\min} + E/2)}$, and the quotient of the two bounds is the claim. □

The constant depends on N and E alone, so excess action is rare under every exterior, at a cost that is exponential in N. The minimum itself is a function of the exterior, and the lemma says nothing about the size of its plaquettes: it controls the fluctuation of the field about the configuration the exterior forces, not that configuration.

Lemma 12.73 (restriction to a fixed set)**.** *Let $G = \mathrm{SU}(2)$, let e be a link and let $h \geq 0$ be an integrable function on G with $\int h \, dU > 0$ that does not depend on the other links. For a configuration σ of the other links let $\gamma_e^h(\cdot \mid \sigma)$ be the law on G with density proportional to $e^{-\beta_W S} h$. If σ and σ' differ only at a link $e' \neq e$, then*

$$W_1(\gamma_e^h(\cdot \mid \sigma), \gamma_e^h(\cdot \mid \sigma')) \leq \frac{\pi^2}{8}\, \beta_W\, \delta(\sigma_{e'}, \sigma'_{e'}),$$

and the two laws coincide unless e and e' lie in a common plaquette.

Proof. Only the plaquettes containing e enter the law, and two distinct links of $\mathbb{Z}^4$ lie in at most one common plaquette p_0; if there is none the laws are equal. Otherwise move $\sigma_{e'}$ along a unit-speed geodesic $U(t)$, $0 \leq t \leq T = \delta(\sigma_{e'}, \sigma'_{e'})$, and let γ_t be the corresponding law. The function $t \mapsto s_{p_0}$ has $|\frac{d}{dt} s_{p_0}| \leq \frac{\pi}{2}$ as in the proof of Lemma 12.72, and the derivative $\dot{s}_{p_0}$ is a function of U_e with oscillation at most π. For a 1-Lipschitz f on G, which has oscillation

at most the diameter π, $\frac{d}{dt}\gamma_t(f) = -\beta_W \operatorname{Cov}_{\gamma_t}(f, \dot{s}_{p_0})$ because h does not depend on t, and $|\operatorname{Cov}(f, \dot{s}_{p_0})| \le \sqrt{\operatorname{Var} f \, \operatorname{Var} \dot{s}_{p_0}} \le \frac{\pi}{2} \cdot \frac{\pi}{4}$. Integrating over t and taking the supremum over f gives the bound. □

When h is the indicator of a set C the lemma says that conditioning a link to lie in C costs nothing in its response, however small or irregular C is, as long as C is fixed. The responses that are not bounded are those to a set that moves with the neighbours, such as the intersection of balls centred at them; this is the difference between conditioning on a fixed set of bad links and conditioning on a plaquette being large.

Proposition 12.74 (one-point rarity)**.** *Let $\mathcal{A}$ be a set of links, D the diameter of G, and γ_e, $e \in \mathcal{A}$, single-link kernels with*

$$W_1\big(\gamma_e(\cdot \mid \sigma), \gamma_e(\cdot \mid \sigma')\big) \le \sum_{e'} c_{ee'}(\sigma, \sigma')\, \delta(\sigma_{e'}, \sigma'_{e'})$$

for all σ, σ'. Let $\sigma \mapsto B(\sigma) \subset \mathcal{A}$ be measurable, and suppose there are constants $a_{ee'}, A_{ee'} \ge 0$ with $c_{ee'}(\sigma, \sigma') \le a_{ee'}$ whenever $e \notin B(\sigma) \cup B(\sigma')$, $c_{ee'} \le A_{ee'}$ always, $\sup_e \sum_{e'} a_{ee'} \le \theta < 1$ and $\sup_e \sum_{e'} A_{ee'} \le \bar{A}$. If μ_1, μ_2 are invariant under every γ_e and $\mu_i(e \in B) \le q$ for every e and $i = 1, 2$, there is a coupling of μ_1 and μ_2 with

$$\sup_{e \in \mathcal{A}} \mathbb{E}\, \delta(\sigma_e, \sigma'_e) \le \frac{2\bar{A}Dq}{1-\theta}.$$

In particular $|\mu_1(F) - \mu_2(F)| \le L_F |S_F|\, 2\bar{A}Dq/(1-\theta)$ for every F with support S_F that is L_F-Lipschitz in each link.

Proof. Fix $b_0 \in \mathcal{A}$, $\lambda > 0$, and let $w_b = e^{-\lambda |b - b_0|}$, which is summable. As in the proof of Lemma B.1 the functional $\Psi(\pi) = \sum_b w_b \mathbb{E}_\pi \delta(\sigma_b, \sigma'_b)$ has a minimiser π^* among the couplings, and replacing the pair at one link e by a sample of an optimal coupling of $\gamma_e(\cdot \mid \sigma)$ and $\gamma_e(\cdot \mid \sigma')$ gives a coupling again. Only the term at e changes, so minimality gives, with $\varrho(e) = \mathbb{E}_{\pi^*} \delta(\sigma_e, \sigma'_e)$,

$$\varrho(e) \le \mathbb{E}_{\pi^*} \sum_{e'} c_{ee'}(\sigma, \sigma')\, \delta(\sigma_{e'}, \sigma'_{e'})$$

for every e, with no weight left. On the event that e is good in both configurations the coefficients are at most $a_{ee'}$; on its complement, whose probability is at most $\mu_1(e \in B) + \mu_2(e \in B) \le 2q$, they are at most $A_{ee'}$ and every distance is at most D. Hence $\varrho(e) \le \theta \sup \varrho + 2\bar{A}Dq$, and $\sup \varrho \le D$ is finite, which gives the bound. The last statement follows by writing $F(\sigma) - F(\sigma')$ as a telescopic sum over the links of S_F. □

The proposition is approximate uniqueness: all Gibbs measures of the specification lie within $2\bar{A}Dq/(1-\theta)$ of each other, link by link. It is not decay. Its proof sees the bad set only through the one-point probabilities q, and the term $2\bar{A}Dq$ does not shrink with the distance from the region where μ_1 and μ_2 are allowed to differ. The bound D on the distance at a bad link is not what causes this: replacing it by Hölder's inequality gives $\varrho \le \theta\varrho + \bar{A}(qD)^{1/p}\varrho^{1/p'}$, a recursion with the positive fixed point ϱ^* of order $\bar{A}^p qD/(1-\theta)^p$. To obtain decay the bad set has to be held fixed while the comparison runs, as in the disagreement percolation of van den Berg and Maes [79], where the two configurations are coupled link by link and the disagreement has to percolate through the bad set; the price is that the laws conditioned on that set, not the Wilson measure, then have to satisfy the fibre estimates of Section 12.4.

12.7. **Rarity of coarse defects.** Theorem 12.57 makes a large plaquette rare at the lattice scale. The cube condition (D) concerns a cube of fixed physical side, which along the tuning spans $m \sim \beta_W/b$ scales, and it needs the analogue of that rarity for the curvature of the block averages at every one of them. This subsection isolates the two sides of that question. On the energy side, the coarse curvature of a field is bounded by its action uniformly in the scale, for linear fields and for group-valued fields in a torus. On the entropy side, the

chessboard estimate applies to coarse events, and it is the normalisation of Lemma 12.56, not the energy, that makes it useless beyond $O(\log \beta_W)$ scales. A right translation of the Haar measures gives rarity at every scale and under every exterior, with a rate governed by the second variation of the action along a fixed direction; for the abelian theory that second variation is the energy of the direction, while for SU(2) it grows like 4^k at scale k, and the growth is produced by the gauge orbit of the vacuum.

Throughout, Λ_L is the torus of side $L = 2^m$, the scale k satisfies $2^{k+2} \le L$, and the lattice of spacing 2^k is the torus $\Lambda_{L'}$ with $L' = L/2^k \ge 4$. For a real field φ on the links, $d\varphi$ is its curvature, d^{T} the adjoint of d, and $\Delta = d^{\mathsf{T}}d + dd^{\mathsf{T}}$, which acts on each component of a field as the lattice Laplacian $(\Delta f)(x) = 8f(x) - \sum_{\pm\nu} f(x \pm e_\nu)$. The Green function $G_{L'}$ of $\Lambda_{L'}$ is the function of mean zero with $\Delta G_{L'} = \delta_0 - L'^{-4}$; it is invariant under the permutations and sign changes of the coordinates, and applied to each component it inverts Δ on fields whose components have mean zero. It commutes with d, d^{T} and the translations.

Lemma 12.75 (energy of a coarse defect)**.** *Let P be a plaquette of the lattice of spacing 2^k and $J_{k,P}$ the field with $(d_k V^{(k)}(\varphi))_P = \langle \varphi, J_{k,P} \rangle$ for every field φ on Λ_L, given by Lemma 12.10(i). Then $d^{\mathsf{T}} J_{k,P} = 0$, every field φ satisfies*

$$(d_k V^{(k)}(\varphi))_P^2 \le \gamma_k \, Q(\varphi),$$

where

$$\gamma_k = \langle J_{k,P}, G_L J_{k,P} \rangle = \sup_{\varphi} \frac{(d_k V^{(k)}(\varphi))_P^2}{Q(\varphi)},$$

and $\frac{1}{432} \le \gamma_k \le \frac{1}{2}$ for every k. In particular $Q(\varphi) \ge 2r^2$ whenever $|(d_k V^{(k)}(\varphi))_P| \ge r$, at every scale.

Proof. Lemma 12.10(ii) shows that $V^{(k)}(dh)$ is the gradient of $\bar{h}$ on the coarse lattice for every function h on sites, so $\langle dh, J \rangle = 0$, the curvature of a gradient vanishing, and $d^{\mathsf{T}} J = 0$; J is supported on the links of directions μ, ν of the plane of P, and its component in each direction has mean zero, the two sides of P entering with opposite signs. Put $u = G_L J$. Then $\Delta u = J$, $d^{\mathsf{T}} u = G_L d^{\mathsf{T}} J = 0$, and for every field φ

$$\begin{aligned} \langle \varphi, J \rangle = \langle \varphi, \Delta u \rangle = \langle d\varphi, du \rangle + \langle d^{\mathsf{T}}\varphi, d^{\mathsf{T}} u \rangle \\ = \langle d\varphi, du \rangle \le \sqrt{Q(\varphi) Q(u)}, \end{aligned}$$

with $Q(u) = \langle u, d^{\mathsf{T}} du \rangle = \langle u, \Delta u \rangle = \langle u, J \rangle = \gamma_k$, and equality at $\varphi = u$. This proves the inequality and the characterisation of γ_k as a supremum.

For the upper bound, Proposition 12.9(i) gives $Q_k(V^{(k)}(\varphi)) \le Q(\varphi)$ for every field, its proof being a local identity, so γ_k is at most the supremum of $(d_k w)_P^2 / Q_k(w)$ over the fields w of $\Lambda_{L'}$. The argument of the first paragraph on $\Lambda_{L'}$, with $j = d_k^{\mathsf{T}} e_P$ the field equal to ± 1 on the four sides of P, identifies that supremum with $\langle j, G_{L'} j \rangle$. The two sides of P in the direction μ are the links (y, μ) and $(y + e_\nu, \mu)$ with opposite signs, so

$$\begin{aligned} \langle j, G_{L'} j \rangle = 4G_{L'}(0) - 2G_{L'}(e_\nu) - 2G_{L'}(e_\mu) \\ = 4\big(G_{L'}(0) - G_{L'}(e_1)\big) = \tfrac{1}{2}(1 - L'^{-4}), \end{aligned}$$

by $(\Delta G_{L'})(0) = 8G_{L'}(0) - 8G_{L'}(e_1) = 1 - L'^{-4}$ and the symmetry of $G_{L'}$.

For the lower bound take P at $y = 0$ in the plane $\{1, 2\}$, write $L_k = 2^k$, and let f be the function on $\mathbb{Z}/L\mathbb{Z}$ with $f(s) = s/L_k$ for $0 \le s \le L_k$, $f(s) = 1$ for $L_k \le s \le 2L_k$, $f(s) = 1 - (s - 2L_k)/L_k$ for $2L_k \le s \le 3L_k$ and $f = 0$ elsewhere; let g be the indicator of $[0, 2L_k)$; and let $h(s) = (1 - \operatorname{dist}(s, [0, L_k))/L_k)_+$. Put $\varphi_{(x,2)} = L_k^{-1} f(x_1) g(x_2) h(x_3) h(x_4)$ and $\varphi = 0$ on the other links; all supports fit in the torus since $4L_k \le L$. By Lemma 12.10(i), $V^{(k)}_{y,2}(\varphi) = L_k^{-4} \sum_{x \in B_k(y)} \sum_{t < L_k} \varphi_{(x + te_2, 2)}$, and for $y = 0$ and $y = e_1$ the factors g and h equal 1 on the ranges that occur, so $V^{(k)}_{0,2}(\varphi) = \bar{f}(0)$ and $V^{(k)}_{e_1,2}(\varphi) = \bar{f}(1)$, the means of f over




FIGURE 20. The smeared current loop $J_{k,P}$ of Lemma 12.75 in the plane of P, drawn for $L_k = 4$. Left: each side of the coarse plaquette P (thick square) contributes a current of its own direction on the links of a slab of transverse size L_k^3 and length $2L_k - 1$ (blue for the direction 1, red for the direction 2), with the signs of the four sides; the slabs of opposite sides are disjoint and the two directions are carried by different links. Right: the longitudinal profile $L_k^4 J = \pm c(s)$ on a slab, the tent $c(s) = \min\{s+1, L_k, 2L_k-1-s\}$, which counts the straight paths of Lemma 12.10(i) through the link at position s. The current is divergence free and its energy $\gamma_k = \langle J_{k,P}, G_L J_{k,P}\rangle$ lies in $[\frac{1}{432}, \frac{1}{2}]$ at every scale.

$[0, L_k)$ and $[L_k, 2L_k)$, which are $(L_k-1)/(2L_k)$ and 1. Hence $(d_k V^{(k)}(\varphi))_P = \bar f(1) - \bar f(0) = (L_k+1)/(2L_k) \geq \frac{1}{2}$. The curvature of φ lives on the plaquettes of the planes $\{1,2\}$, $\{2,3\}$, $\{2,4\}$, and g is never differenced, so

$$Q(\varphi) = L_k^{-2}(\|\nabla f\|^2\|g\|^2\|h\|^4 + 2\|f\|^2\|g\|^2\|\nabla h\|^2\|h\|^2) \\ \leq L_k^{-2}\Big(\frac{2}{L_k} 2L_k (3L_k)^2 + 2\cdot 3L_k \cdot 2L_k \frac{2}{L_k} 3L_k\Big),$$

where ∇ is the forward difference: f and h change by $1/L_k$ on $2L_k$ steps and by nothing elsewhere, and f^2, g^2, h^2 have sums at most $3L_k$, $2L_k$, $3L_k$. So $Q(\varphi) \leq 108$, and $\gamma_k \geq \frac{1}{4}/108$. □

The fields $J_{k,P}$ are smeared current loops: by Lemma 12.10(i) each side of P contributes $\pm L_k^{-4} c(s)$ on the $L_k^3(2L_k-1)$ links of its direction whose transverse coordinates lie in the block and whose longitudinal coordinate s ranges over $[0, 2L_k-2]$, with the tent $c(s) = \min\{s+1, L_k, 2L_k-1-s\}$. A bare holonomy around a loop of n links forces action only of order r^2/n, since the loop can be thickened to a tube of any radius; the block average smears the loop over a four-dimensional region, and this is why its energy does not decrease with the scale. The lemma is the quadratic part of Theorem 12.12, seen from one coarse plaquette. The same scale invariance holds for group-valued fields in a torus.

Proposition 12.76 (energy of a coarse defect, torus-valued fields)**.** *Let U be an S-valued field in the tree gauge on Λ_L, as in Theorem 12.29, and let P be a plaquette of the lattice of spacing 2^k with $\delta(V^{\sharp,k}(U)_P, I) \geq r$. Then*

$$Q(U) \geq \min\Big\{\frac{r^2}{1.001}, \frac{1}{144.2}\Big\} \geq \frac{r^2}{1154}.$$

Proof. If some plaquette has $\delta(U_p, I) > 1 = \rho_G$ then $Q(U) > 1 \geq r^2/8$, since $r \leq \operatorname{diam} \mathrm{SU}(2) = 2\sqrt{2}$. Otherwise every Θ_p is defined, and for $\Theta_p = \theta_p J_p$ with $J_p^2 = -I$ one has $\|\Theta_p\|^2 = 2\theta_p^2$ and $\delta(U_p, I)^2 = 8\sin^2(\theta_p/2)$, so $\|\Theta_p\|^2 \leq 1.001\,\delta(U_p, I)^2$ when $\theta_p \leq 1/(12\sqrt{2})$, that is when $\|\Theta_p\| \leq \eta_G = \frac{1}{12}$. Let j_* be the least $j \leq k$ with

$\max_P \|\Theta_P^{(j)}\| > \eta_G$, if there is one. If there is none, (12.5) holds for every $j < k$, and Theorem 12.29 gives $r^2 \le \delta(V^{\sharp,k}(U)_P, I)^2 \le Q_k(V^{\sharp,k}(U)) \le 1.001\, Q(U)$. If $j_* = 0$, some plaquette has $\theta_p > 1/(12\sqrt{2})$ and $Q(U) \ge 8\sin^2(\theta_p/2) \ge 1/144.1$. If $1 \le j_* \le k$, (12.5) holds for $j < j_*$, so Theorem 12.29 at the scale j_* shows that $\Theta^{(j_*)}$ is defined and $\sum_P \|\Theta_P^{(j_*)}\|^2 \le \sum_p \|\Theta_p\|^2 \le 1.001\, Q(U)$, every fine plaquette having $\|\Theta_p\| \le \eta_G$; one term of the left side exceeds $1/144$, so $Q(U) \ge 1/144.2$. □

The S-valued fields form a null set for the Wilson measure, and for general SU(2) fields Theorem 12.33 controls the cut-off average only while $\sum_j \eta_j$ stays small; whether a bound of the form $Q(U) \ge c\, r^2$, uniform in k, holds for every SU(2) field is not known. The next proposition shows that, whatever the energy of a coarse defect, the chessboard estimate cannot use it beyond a bounded number of scales.

Proposition 12.77 (chessboard for coarse events)**.** *Let $\beta_W \ge 1$, let $b = 2^j$ with $2b \mid L$, and let E be an event determined by the links of the closed cube $[0,b]^4$ on which the action of the plaquettes of that cube is at least A. For $z \in (b\mathbb{Z}/L\mathbb{Z})^4$ let E_z be the event obtained from E by the reflections that carry $[0,b]^4$ to $z + [0,b]^4$. Then for every set Z of such z,*

$$\mu_L\Big(\bigcap_{z\in Z} E_z\Big) \le \exp\Big(|Z|\Big(6b^4(\log\beta_W + C_0) - \frac{\beta_W A}{4}\Big)\Big).$$

Proof. As in the proof of Theorem 12.57, μ_L is reflection positive in the hyperplanes through sites, and the chessboard estimate for functions of the links of the closed blocks $z + [0,b]^4$, $z \in (b\mathbb{Z}/L\mathbb{Z})^4$, bounds the probability of $\bigcap_{z\in Z} E_z$ by that of $\bigcap_z E_z$ over all $(L/b)^4$ blocks, raised to the power $|Z|(b/L)^4$. Reflection preserves the Haar measures and $\operatorname{Tr} U_p$, so on every E_z the plaquettes of the block $z + [0,b]^4$ carry action at least A. A plaquette lies in at most four closed blocks, one for each choice of the two transverse coordinates of the corner, so on the intersection $S \ge (L/b)^4 A/4$, and the event has unnormalised weight at most $e^{-\beta_W (L/b)^4 A/4}$. Dividing by $Z_L \ge e^{-6L^4(\log\beta_W + C_0)}$ of Lemma 12.56 and raising to the power $|Z|(b/L)^4$ gives the bound. □

A plaquette of the lattice of spacing 2^k depends, by Lemma 12.61 applied at each step, on the fine links of a cube of side $2^{k+2} - 4$, so the event that it carries a large cut-off average is of the kind in the proposition with $b = 2^{k+2}$. Whatever action A that event forces, the bound is below one only when $6 \cdot 16^{k+2}(\log\beta_W + C_0) < \beta_W A/4$, that is $16^k \lesssim \beta_W A/\log\beta_W$. For the events of Lemma 12.75 and Proposition 12.76, A is of the order of r^2 at every scale, and the loss is entirely in the normalisation: the lower bound of Lemma 12.56 charges the full fluctuation volume of the block, $6b^4(\log\beta_W + C_0)$, against a defect whose energy does not grow with the block. In the Gaussian theory the two fluctuation volumes, around the defect and around the vacuum, are the same, and the following translation argument transfers that cancellation to the group, under every exterior.

Theorem 12.78 (Haar tilt)**.** *Let G be* SU(2) *or* U(1)*, let Λ be a finite set of links, either all links of Λ_L or the inner links of a box $R \subset \mathbb{Z}^4$ whose other links are fixed, and let μ be the law with density $Z^{-1}e^{-\beta_W S}$ with respect to the product of the Haar measures on Λ, where $S = \sum_p s_p$ runs over the plaquettes meeting Λ and $s_p = 1 - \frac{1}{N}\operatorname{Re}\operatorname{Tr} U_p$. For $r = (r_e)_{e\in\Lambda}$ with $r_e \in \mathfrak{g}$ and $\rho_e = \|r_e\|_{\mathrm{op}}$ put $U_e^t = U_e e^{t r_e}$,*

$$D_r S(U) = \partial_t S(U^t)\big|_{t=0}, \qquad K_r = \sup_U \partial_t^2 S(U^t)\big|_{t=0}.$$

(i) $K_r \le \sum_p \big(\sum_{e\subset p,\, e\in\Lambda} \rho_e\big)^2 \le 24 \sum_{e\in\Lambda} \rho_e^2$. *For $G = \mathrm{U}(1)$ and $r_e = i a_e$ with $a_e \in \mathbb{R}$, $K_r \le Q(a) = \sum_p (da)_p^2$.*

(ii) $\mathbb{E}_\mu e^{\beta_W t D_r S} \le e^{\beta_W t^2 K_r/2}$ *for every $t \in \mathbb{R}$.*

(iii) *For every $x > 0$, uniformly in the fixed links,*

$$\mu(D_r S \ge x) \le e^{-\beta_W x^2/(2K_r)}, \qquad \mu(D_r S \le -x) \le e^{-\beta_W x^2/(2K_r)}.$$

Proof. (i) Order the links of a plaquette p as traversed, so that $U_p = g_1^{\epsilon_1}\cdots g_4^{\epsilon_4}$ with $\epsilon_i = \pm 1$. Since $ge^{tr} = e^{t\,\mathrm{Ad}(g)r}g$ and $(ge^{tr})^{-1} = e^{-tr}g^{-1}$, moving the exponentials to the left gives $U_p(t) = e^{t\tilde a_1}\cdots e^{t\tilde a_4}U_p$ with $\tilde a_i = \pm\,\mathrm{Ad}(h_i)r_{e_i}$ for unitary h_i, so $\|\tilde a_i\|_{\mathrm{op}} = \rho_{e_i}$, and $\tilde a_i = 0$ if $e_i \notin \Lambda$. Then

$$\partial_t^2 \operatorname{Tr} U_p(t)\big|_{t=0} = \operatorname{Tr}\Big[\Big(\sum_i \tilde a_i^2 + 2\sum_{i<j}\tilde a_i\tilde a_j\Big)U_p\Big],$$

and $|\operatorname{Tr}(XU)| \le N\|X\|_{\mathrm{op}}$ for unitary U, so $|\partial_t^2 s_p| \le (\sum_i \rho_{e_i})^2 \le 4\sum_{e\subset p}\rho_e^2$; each link lies in six plaquettes. The second derivative at any other t is the one at 0 for the field U^t, so the bound holds for K_r. For U(1) all factors commute, $U_p(t) = e^{it(da)_p}U_p$ and $\partial_t^2 s_p = (da)_p^2 \operatorname{Re}\operatorname{Tr} U_p \le (da)_p^2$.

(ii) Right multiplication by the fixed element e^{-tr_e} preserves the Haar measure of each link of Λ, so

$$\int e^{-\beta_W S(Ue^{-tr})}\prod_{e\in\Lambda} dU_e = Z,$$

that is, $\mathbb{E}_\mu e^{-\beta_W[S(Ue^{-tr})-S(U)]} = 1$. Taylor's formula along $s \mapsto Ue^{-sr}$ with the Lagrange remainder gives

$$S(Ue^{-tr}) - S(U) = -tD_rS(U) + \tfrac{t^2}{2}\partial_s^2 S(Ue^{-sr})\big|_{s=s_*} \le -tD_rS(U) + \tfrac{t^2}{2}K_r,$$

the second derivative being even in r. Hence

$$\mathbb{E}_\mu e^{\beta_W t D_rS - \beta_W t^2 K_r/2} \le 1.$$

(iii) By Chebyshev's inequality and (ii), $\mu(D_rS \ge x) \le e^{-\beta_W tx + \beta_W t^2 K_r/2}$ for $t > 0$; take $t = x/K_r$. The lower tail is the upper tail for $-r$. Nothing depends on the fixed links. □

The map $U \mapsto Ue^{-tr}$ is the group version of a translation of the Gaussian field; its Jacobian is one, and no smallness of the field enters. The observable D_rS is the component of the force along r: at a plaquette, $\partial_t s_p = -\frac{1}{N}\operatorname{Re}\operatorname{Tr}\big((\sum_i \tilde a_i)U_p\big)$, the covariant derivative of r in the background U paired with the sine part of U_p. For the abelian theory the second variation is the energy of the direction, and the theorem gives the rarity of coarse defects at every scale.

Corollary 12.79 (abelian theory)**.** *Let $G = \mathrm{U}(1)$, let μ_L be its Wilson measure on Λ_L, $U_e = e^{i\theta_e}$, $\theta_p = (d\theta)_p$ the angle of U_p in $(-\pi,\pi]$, and let $a = G_L J_{k,P}$ be the minimiser of Lemma 12.75. For every k, every plaquette P of the lattice of spacing 2^k and every $x > 0$,*

$$\mu_L\Big(\sum_p \sin\theta_p\,(da)_p \ge x\Big) \le e^{-\beta_W x^2}.$$

Here $\sum_p \theta_p(da)_p = (d_kV^{(k)}(\theta))_P$ for every choice of link angles θ_e with $(d\theta)_p \equiv \theta_p$ modulo 2π, up to $2\pi\sum_p n_p(da)_p$ with the integers $n_p = ((d\theta)_p - \theta_p)/2\pi$. Under a fixed exterior of a box, the same bound holds with a replaced by its restriction $\tilde a$ to the inner links and x^2 replaced by $x^2/(2Q(\tilde a))$.

Proof. By Theorem 12.78(i) with $r_e = ia_e$, $K_r \le Q(a) = \gamma_k \le \frac12$, and $D_rS = \sum_p \sin\theta_p(da)_p$ since $s_p(t) = 1-\cos(\theta_p + t(da)_p)$. Part (iii) gives the bound. For the identity, $\sum_p (d\theta)_p(da)_p = \langle\theta, d^{\mathsf T}da\rangle = \langle\theta,\Delta a\rangle = \langle\theta, J_{k,P}\rangle$ because $d^{\mathsf T}a = 0$. In a box the tilt acts on the inner links only, and $K_r \le Q(\tilde a)$. □

For the abelian theory this is the statement that a smeared coarse flux of size x costs $\beta_W x^2$ at every scale, uniformly in the exterior; the four-dimensional abelian theory has no running coupling, and the bound is consistent with its known behaviour at weak coupling [43, 38]. For SU(2) the same tilt is available, but its second variation grows with the scale.

Proposition 12.80 (the gauge orbit of the vacuum)**.** *Let $G = \mathrm{SU}(2)$ on Λ_L, $T \in \mathfrak{g}$ with $\|T\|_{\mathrm{op}} = 1$, $a = G_L J_{k,P}$ for a plaquette P of the lattice of spacing 2^k, and $r_e = a_e T$. Then*

$$\Big(\sqrt{2\|a\|^2 - \gamma_k} - \sqrt{\gamma_k}\Big)_+^2 \le K_r \le 24\|a\|^2$$

and

$$\|a\|^2 \ge \frac{\gamma_k^2}{\|J_{k,P}\|^2} \ge \frac{4^k}{1492992},$$

with $\gamma_k = \langle J_{k,P}, G_L J_{k,P}\rangle$ as in Lemma 12.75; in particular $K_r \ge 4^k/(3\cdot 10^6)$ for $k \ge 11$. The lower bound is attained, up to the stated loss, at a pure gauge field, where $Q(U) = 0$.

Proof. The upper bound is Theorem 12.78(i). For the lower bound let U be the pure gauge field $U_e = h_x h_{x+e}^{-1}$ of a function h on sites with values in $\mathrm{SU}(2)$. The action is invariant under gauge transformations and $U_e e^{tr_e} = h_x \exp(t\,\mathrm{Ad}(h_{x+e}^{-1})r_e)h_{x+e}^{-1}$, so $S(Ue^{tr}) = S(e^{tr'})$ with $r'_e = \mathrm{Ad}(h_{x+e}^{-1})r_e$. At the identity field, $U_p(t) = \exp(tX_p + O(t^2))$ with $X_p = (dr')_p$ the signed sum of the r'_e around p, and $s_p = 1 - \cos\|X_p(t)\|_{\mathrm{op}}$, so $\partial_t^2 s_p|_{t=0} = \|(dr')_p\|_{\mathrm{op}}^2$ and $K_r \ge \sum_p \|(dr')_p\|_{\mathrm{op}}^2$. Take $h_x = R^{x_2}$ with $R = i\sigma_1$ and $T = i\sigma_3$, which is well defined on the torus since $R^4 = I$ and $4 \mid L$, and satisfies $\mathrm{Ad}(R)T = -T$. Then $r'_e = \epsilon_e a_e T$ with $\epsilon_{(x,\mu)} = (-1)^{x_2}$ for $\mu \ne 2$ and $\epsilon_{(x,2)} = (-1)^{x_2+1}$, and $K_r \ge Q(\epsilon a)$. Since $J_{k,P}$ has components only in the directions 1, 2 of the plane of P and G_L acts on each component, $a_3 = a_4 = 0$; the reflection exchanging the coordinates 1 and 2 carries the component a_1 to $-a_2$, so $\|a_1\|^2 = \|a_2\|^2 = \frac12\|a\|^2$. On the plaquette at x in the plane $\{1,2\}$ the signs give

$$\begin{aligned}(d(\epsilon a))_p &= (-1)^{x_2}\big[(a_{(x,1)} + a_{(x+e_2,1)}) - (a_{(x+e_1,2)} - a_{(x,2)})\big] \\ &= (-1)^{x_2}\big[\sigma_x - v_x\big],\end{aligned}$$

so that $Q(\epsilon a) \ge \|\sigma - v\|^2 \ge (\|\sigma\| - \|v\|)_+^2$, with $\|\sigma\|^2 = 4\|a_1\|^2 - \|\nabla_2 a_1\|^2$ and $\|v\|^2 = \|\nabla_1 a_2\|^2$, where ∇_ν is the forward difference in the direction ν. All directional differences together give $\sum_{\mu,\nu}\|\nabla_\nu a_\mu\|^2 = \langle a, \Delta a\rangle = \langle G_L J, J\rangle = \gamma_k$, so $\|\sigma\|^2 \ge 2\|a\|^2 - \gamma_k$ and $\|v\|^2 \le \gamma_k$, which is the lower bound. Finally $\gamma_k = \langle J, a\rangle \le \|J\|\|a\|$, and by Lemma 12.10(i) each of the four sides of P contributes L_k^3 rows of $2L_k - 1$ links with coefficients $L_k^{-4}c(s)$, $c \le L_k$, so $\|J\|^2 \le 4L_k^{-8}L_k^3(2L_k - 1)L_k^2 \le 8\cdot 4^{-k}$; with $\gamma_k \ge \frac{1}{432}$ this gives $\|a\|^2 \ge 4^k/(8\cdot 432^2)$. For $k \ge 11$ that is at least 2, so $2\|a\|^2 - \gamma_k \ge 7\gamma_k$ and $K_r \ge (1 - 7^{-1/2})^2 \cdot \frac78 \cdot 2\|a\|^2 \ge 0.67\|a\|^2 \ge 4^k/(3\cdot 10^6)$. □

So for SU(2) the tilt of Theorem 12.78 proves the rarity of the force component $D_r S$ at the rate $\beta_W x^2/(2K_r)$, which is of order $\beta_W x^2/4^k$ at scale k, and not more: the supremum in K_r is reached on the gauge orbit of the vacuum, where the field has no action and no curvature, because a fixed direction r is not gauge covariant and its covariant derivative in a pure gauge background is as large as r itself. The natural repair is to transport the direction with the field, $r_e \mapsto \mathrm{Ad}(h(U))r_e$. When the transport runs along a forest, the translation still preserves the Haar measures; but the residual gauge freedom of the forest then realises the field of the proof on the links between the trees, and the constant is reached at zero action in that gauge as well.

Proposition 12.81 (covariant tilt on a forest)**.** *Let $G = \mathrm{SU}(2)$, let F be a set of links of Λ_L without cycles, with one root chosen in every component, and for a site y let $h_y(U)$ be the ordered product of the link variables along the path in F from the root of its component to y, with $h_y = I$ at the roots. For $r = (r_e)$ with $r_e \in \mathfrak{g}$ and $r_e = 0$ on F put*

$$\Phi_t(U)_e = U_e \exp\big(t\,\mathrm{Ad}(h_y(U)^{-1})r_e\big), \quad y \text{ the end point of } e,$$

and $\Phi_t(U)_f = U_f$ for $f \in F$; let $D_r^F S(U) = \partial_t S(\Phi_t U)|_{t=0}$ and

$$K_r^F = \sup\big\{\partial_t^2 S(\tilde U e^{tr})\big|_{t=0} :\ \tilde U_f = I \text{ for all } f \in F\big\}.$$

(i) Φ_t *preserves the product of the Haar measures for every* t.

(ii) $\mathbb{E}_\mu e^{\beta_W t D_r^F S} \le e^{\beta_W t^2 K_r^F/2}$ *for every* $t \in \mathbb{R}$, *and* $\mu(\pm D_r^F S \ge x) \le e^{-\beta_W x^2/(2K_r^F)}$ *for every* $x > 0$. $D_r^F S$ *is invariant under the gauge transformations that are trivial at the roots.*

(iii) *Let* F *be one of the sixteen translates* F_o *of the forest of Definition 12.2, with blocks* $2y + o + \{0,1\}^4$, *let* $a = G_L J_{k,P}$ *and* T *be as in Proposition 12.80, and let* $r_e = a_e T$ *on the links outside* F. *Then*

$$K_r^F \ge 2 \sum_{e \notin F} a_e^2,$$

and the bound is reached on pure gauge fields, where $Q(U) = 0$. *For at least one* o *the right side is at least* $\frac{29}{16}\|a\|^2 \ge \frac{29}{16} \cdot \frac{4^k}{1492992}$.

Proof. (i) Let $\Gamma(U) = (U_F, \tilde U)$ with $\tilde U_e = h_x(U) U_e h_y(U)^{-1}$ for $e = (x,y) \notin F$. Given the variables on F, the map $U_e \mapsto \tilde U_e$ is a left and a right translation by elements that depend only on U_F, so Γ is a bijection of the configuration space that preserves the product of the Haar measures; this is the change of variables of Lemma 3.1. Let R_t multiply every coordinate $\tilde U_e$, $e \notin F$, on the right by $e^{t r_e}$ and leave U_F alone. Then

$$(\Gamma^{-1} R_t \Gamma U)_e = h_x^{-1}(h_x U_e h_y^{-1} e^{t r_e}) h_y = U_e \exp\bigl(t \operatorname{Ad}(h_y^{-1}) r_e\bigr),$$

so $\Phi_t = \Gamma^{-1} R_t \Gamma$ is a composition of three maps that preserve the measure.

(ii) Since the action is gauge invariant, $S(\Phi_t U) = S(\tilde U e^{tr})$, and $\tilde U e^{tr}$ again has trivial links on F. Taylor's formula along $s \mapsto \Phi_{-s} U$ with the Lagrange remainder gives $S(\Phi_{-t} U) - S(U) \le -t D_r^F S(U) + \frac{t^2}{2} K_r^F$, and $\int e^{-\beta_W S \circ \Phi_{-t}}\, dU = Z$ by (i); the rest is the proof of Theorem 12.78 (ii), (iii). Under a gauge transformation g the transports change to $g_o h_y g_y^{-1}$, with o the root of the component of y, so $\tilde U$ is conjugated by the root values and $D_r^F S$ is unchanged when they are trivial.

(iii) The tree links of F_o join sites of one block, so a pure gauge field $U = I^g$, $U_e = g_x g_{x+e}^{-1}$, with g constant on the blocks has trivial links on F_o, lies in the set of the supremum and has $Q(U) = 0$. By the proof of Proposition 12.80, $\partial_t^2 S(U e^{tr})|_{t=0} = \sum_p \|(dr')_p\|_{\mathrm{op}}^2$ with $r'_e = \operatorname{Ad}(g_{x+e}^{-1}) r_e$, the end point of e taken. Since $a_3 = a_4 = 0$, a plaquette p at x in a plane $\{\mu,\nu\}$ with $\mu \in \{1,2\}$ and $\nu \in \{3,4\}$ has at most two tilted sides, $e = (x,\mu)$ with end point $x + e_\mu$ and $e' = (x + e_\nu, \mu)$ with end point $x + e_\mu + e_\nu$, and

$$\|(dr')_p\|_{\mathrm{op}}^2 = a_e^2 + a_{e'}^2 - 2 a_e a_{e'} \langle \operatorname{Ad}(g_{x+e_\mu}^{-1}) T, \operatorname{Ad}(g_{x+e_\mu+e_\nu}^{-1}) T \rangle,$$

with the inner product of $\mathfrak{g}$ in which $\|T\| = 1$ and $a_e = 0$ on F_o. When $(x-o)_\nu$ is odd the two end points lie in different blocks. Take the block values g_B independent with Haar law. The adjoint representation of SU(2) is irreducible of dimension three, so $\mathbb{E} \operatorname{Ad}(g_B) = 0$, and the expectation of the inner product vanishes. Every link $e = (x,\mu) \notin F_o$ with $\mu \in \{1,2\}$ is, for each $\nu \in \{3,4\}$, a side of exactly one such plaquette, at x if $(x-o)_\nu$ is odd and at $x - e_\nu$ otherwise. Dropping the other plaquettes, whose terms are nonnegative,

$$\mathbb{E}_g\, \partial_t^2 S(I^g e^{tr})\big|_{t=0} \ge \sum_{\nu \in \{3,4\}} \sum_{e \notin F_o} a_e^2 = 2 \sum_{e \notin F_o} a_e^2,$$

and the supremum over the section is at least the expectation. A link (x,μ) lies in F_o for one of the sixteen offsets o when $\mu = 1$ and for two of them when $\mu = 2$. Averaging over o and using $\|a_1\|^2 = \|a_2\|^2 = \frac12 \|a\|^2$ from the proof of Proposition 12.80,

$$\frac{1}{16} \sum_o \sum_{e \notin F_o} a_e^2 = \frac{15}{16}\|a_1\|^2 + \frac{14}{16}\|a_2\|^2 = \frac{29}{32}\|a\|^2,$$

so one offset reaches $\frac{29}{32}\|a\|^2$; the last bound is the lower bound for $\|a\|^2$ of that proposition. □




FIGURE 21. The covariant tilt on the block forest (Proposition 12.81). Transporting the direction along the trees keeps the Haar measures, but the section still contains every pure gauge field that is constant on the blocks. On a plaquette whose tilted sides e, e' end in different blocks the transported directions are turned by independent elements, and the average of $\|(dr')_p\|^2$ over them is $a_e^2 + a_{e'}^2$ instead of the $(a_e - a_{e'})^2$ of the abelian energy.

Proposition 12.81 and Figure 21 show where the obstruction sits. The Haar translation survives gauge covariance along a forest, and for the block forest the constant it produces is at least $\frac{29}{16}\|a\|^2$, reached by fields with no action and no curvature at all; the chessboard estimate and the covariant tilt are therefore lost on the same fields, the gauge orbit of the vacuum, for different reasons. A bound of the form

$$\partial_t^2 S(\tilde{U} e^{tr}) \le Q(r) + \mathcal{R}(\tilde{U}, r),$$

with $\mathcal{R}$ supported on the large-field region, is false for every forest section, since the left side exceeds $Q(r)$ by a fixed multiple of $\|a\|^2$ at $\tilde{U} = I^g$, where the large-field region is empty. A section that is smooth on the orbits, such as the Landau gauge, has no such pure gauge fields, but its Jacobian is a Faddeev–Popov determinant and not one, and part (ii) does not apply to it as it stands.[23]

Along the tuning $\beta_W \sim b \log(1/a)$ a cube of fixed physical side spans $m \sim \beta_W / b$ scales. Proposition 12.77 covers $O(\log \beta_W)$ of them, Propositions 12.80 and 12.81 cover $O(\log \beta_W)$ of them without the torus and uniformly in the exterior, and Corollary 12.79 covers all of them for the abelian theory. What the cube condition needs for SU(2) is a translation of the Haar measures whose second variation is the energy of the direction up to a constant independent of the scale, together with a non-perturbative treatment of the last $O(1)$ scales, where the running coupling is of order one and coarse defects are not rare at all. Proposition 12.81 shows that no translation transported along a forest does this: its constant is reached at zero action, with the growth 4^k of the energy of the smeared current under the residual gauge freedom, so the missing estimate is not a large-field remainder but a different section of the gauge orbits, with a Jacobian that enters the argument of part (ii). This is the content of Bałaban's large-field expansion [8], and it is not carried out here. The cube condition itself is a mixing condition of Dobrushin–Shlosman type, which for compact spins with finite range is equivalent to a logarithmic Sobolev inequality uniform in the volume [73]; the rarity of coarse defects is the part of that inequality that concerns the large fields.

[23] The Wilson measure satisfies no curvature-dimension condition at the values of β_W considered here, so the inequality cannot be obtained from a Bakry–Émery argument either: for one link equal to $-I$ with trivial neighbours the second variation of S along a unit direction on that link is -6, since each of its six plaquettes contributes $\partial_t^2(1 + \cos t) = -1$, while the Ricci curvature of SU(2) with unit-speed geodesics e^{tr}, $\|r\|_{\mathrm{op}} = 1$, is 2; the Hessian of $\beta_W S$ plus the curvature is negative as soon as $\beta_W > 1/3$.




FIGURE 22. Which scales the estimates of this subsection reach, along the tuning $\beta_W \sim b \log(1/a)$, where a cube of fixed physical side spans $m \sim \beta_W/b$ scales. The energy of a coarse defect is scale invariant (Lemma 12.75), and for the abelian theory the Haar tilt makes coarse defects rare at every scale (Corollary 12.79). For SU(2) the chessboard estimate stops after $O(\log \beta_W)$ scales because the normalisation charges the whole fluctuation volume of a block against the defect (Proposition 12.77), and the tilt stops after $O(\log \beta_W)$ scales because its second variation grows like 4^k on the gauge orbit of the vacuum, whether the direction is fixed or transported along a forest (Propositions 12.80, 12.81). The remaining scales are the open input of the cube condition.

13. Labels, islands and the environment

The cube condition (D) is a statement about conditional laws under every exterior, and Section 12.6 showed that one-point rarity of a bad set gives closeness of Gibbs measures but not decay. The route that remains is the one of the large-field expansions: label each block of a window good or bad, prove decay on the good windows, bound the bad islands by their rarity, and control the covariance that the labels themselves carry. This section proves what each of these four steps gives for the cut-off average of Section 12.2, and proves, by explicit examples, where the natural choices of labels fail. The statements are exact and finite; none of them closes (D), and Section 13.8 lists what they leave.

Throughout, $G = \mathrm{SU}(2)$ with $\rho_G = 1$, the lattice is in the tree gauge of Definition 12.2, $d(g,h) = \|\log(g^{-1}h)\|$ is the bi-invariant geodesic distance in the Hilbert–Schmidt norm, so that $d(g,I) = \|X\|$ for $g = e^X$ with $\|X\| \leq \pi\sqrt{2}$, and δ is the chordal distance. A *pair* $k = (y,\mu)$ consists of a block and a direction; its *path link* is $\ell_k = U_{2y+e_\mu,\mu}$, its *private links* are the eight links leaving $B(y)$ in the direction μ, which enter no other cut-off average, and the seven private links other than ℓ_k are its *non-path* links. In the sixteen paths of $V_k^\sharp$ the path link occurs twice, as the two paths of transverse offset zero, and each non-path private link occurs twice, each time multiplied by one link inside $B(y)$ or $B(y+e_\mu)$. For a coarse field c the reconstruction $R(c)$ of Section 12.2 puts c_k on the private links of k and I on the links inside the blocks.

13.1. **Backgrounds and the path links.** The first question is which background the chart of Theorem 12.41 should be centred at when a window is conditioned on its labels. The natural label of a link is its distance to the reconstruction of the coarse field, and the natural background is the reconstruction itself. The first result shows that this is not enough: a coarse field all of whose plaquettes are large has a reconstruction at which every relative link label is good and the constrained action has a negative second variation along the fibre.

Proposition 13.1 (a rough background)**.** *On the periodic lattice of side four, with coarse torus $(\mathbb{Z}/2\mathbb{Z})^4$, let $c_{y,\mu} = (-I)^{\sum_{\nu<\mu} y_\nu}$. Then every coarse plaquette of c equals $-I$, the field $U = R(c)$ is in the tree gauge and has $V^\sharp(U) = c$, and $d(U_e, R_e(c)) = 0$ for every link. There is a curve $U(t) = U\exp(t\xi)$, ξ supported on four links, such that $V^\sharp(U(t)) = c$ for all*

small $|t|$, $d(U_e(t), R_e(c)) \le |t|$, and the Wilson action $S = \sum_p (1 - \frac{1}{2}\operatorname{Re}\operatorname{Tr} U_p)$ satisfies

$$\frac{d^2}{dt^2}\Big(\beta_W S(U(t))\Big)\Big|_{t=0} = -\beta_W, \qquad T(\xi) = \sum_e \|\xi_e\|^2 = 4.$$

Every first derivative of S at U vanishes, so the intrinsic Hessian of $\beta_W S$ on the fibre of c at U has an eigenvalue at most $-\beta_W/4$, in any smooth chart of the fibre.

Proof. The coarse plaquette at y in the plane $\{\mu, \nu\}$, $\mu < \nu$, is the product of $c_{y,\mu}$, $c_{y+e_\mu,\nu}$, $c^{-1}_{y+e_\nu,\mu}$ and $c^{-1}_{y,\nu}$; these are central, and the exponent of $-I$ is $\sum_{\rho<\mu} y_\rho + \sum_{\rho<\nu}(y+e_\mu)_\rho - \sum_{\rho<\mu}(y+e_\nu)_\rho - \sum_{\rho<\nu} y_\rho = 1$ modulo two, since $\mu < \nu$ puts e_μ inside the second sum and e_ν outside the third. So every coarse plaquette is $-I$, also across the periodic seam. In $U = R(c)$ the links inside the blocks are I, in particular the tree links, and all sixteen paths of a pair k equal c_k, so every $v_x = I$, the cut-off is one and $V^\sharp_k(U) = c_k$; the relative labels vanish by construction.

Let $T \in \mathfrak{su}(2)$ have $\|T\| = 1$, with eigenvalues $\pm i/\sqrt{2}$, and let $\xi_e = a_e T$ with $a_e = -1, +1, +1, -1$ on the links $((0,1,1,1),1)$, $((0,1,1,1),2)$, $((0,1,0,0),1)$, $((0,0,1,0),2)$ and $a_e = 0$ elsewhere. All four links are private links; the last two are path links. The pair $((0,0,0,0),1)$ has the non-path link $((0,1,1,1),1)$ with $a = -1$ and the path link $((0,1,0,0),1)$ with $a = +1$, the pair $((0,0,0,0),2)$ has $((0,1,1,1),2)$ with $a = +1$ and the path link $((0,0,1,0),2)$ with $a = -1$, and no other pair contains a link of the support. Since the background is central, $P_x(t) = c_k \exp(t a_x T)$ for the sixteen paths, with a_x the value on the private link of the path, so $v_x(t) = \exp(t(a_x - a_0)T)$ lies in the chart for small $|t|$, the cut-off is one, and

$$V^\sharp_k(U(t)) = c_k \exp\Big(\frac{t}{16}\sum_x a_x T\Big) = c_k,$$

because each private link occurs in exactly two paths and the two nonzero values of each pair cancel. The curve stays in the fibre, and $d(U_e(t), R_e(c)) = |t a_e| \le |t|$.

Write $U_p = s_p I$ with $s_p = \pm 1$ and let $q_p = (d\,a)_p$ be the signed curl of the scalar field a around p. Since the perturbations commute with the background, $U_p(t) = s_p \exp(t q_p T)$ and $\operatorname{Re}\operatorname{Tr} U_p(t) = 2 s_p \cos(t q_p/\sqrt{2})$, so

$$S(U(t)) = \sum_p \big(1 - s_p \cos(t q_p/\sqrt{2})\big), \qquad S''(0) = \frac{1}{2}\sum_p s_p q_p^2.$$

The plaquettes with $q_p \neq 0$ are those containing a link of the support. A link (x, μ) lies in six plaquettes, two in each plane $\{\mu, \nu\}$, at x and at $x - e_\nu$. Listing the six plaquettes of each of the four links, the plaquette at $(0,1,1,1)$ in the plane $\{1,2\}$ contains two links of the support, $((0,1,1,1),1)$ and $((0,1,1,1),2)$, with $q_p = -1 - 1 = -2$; the other 22 plaquettes contain one link each, with $q_p = \pm 1$. The sign s_p of a plaquette is -1 exactly when it has an odd number of links on which U equals $-I$; counting through the four links, which is a finite check, gives twelve plaquettes with $s_p = 1$, $q_p^2 = 1$, ten with $s_p = -1$, $q_p^2 = 1$, and the one with $s_p = -1$, $q_p^2 = 4$. Hence $\sum_p s_p q_p^2 = 12 - 10 - 4 = -2$, $S''(0) = -1$, and $T(\xi) = 4$.

Every plaquette of U is central, so every first derivative of S at U vanishes and the second derivative along any curve through U with velocity ξ depends only on ξ: the intrinsic Hessian on the fibre has the value -1 at ξ, which has $T(\xi) = 4$. The Haar density of the fibre, in a chart in which the path links are solved for, is the inverse Jacobian of the path-link equations, which at U is a fixed positive number independent of β_W; its logarithmic second derivatives are bounded in any fixed chart, so no choice of chart turns the Hessian of $\beta_W S$ plus the log-density positive for all large β_W. □

The proposition does not contradict Theorem 12.41, which assumes small plaquettes; it shows that the hypothesis of that theorem on the coarse field cannot be dropped in favour of relative link labels. A coarse-curvature label is therefore part of any labelling, and the next lemma shows that it has a witness at the fine scale, so that it is as rare as a fine defect.

Lemma 13.2 (a fine witness for coarse curvature)**.** *Let P be the coarse plaquette at y in the plane $\{\mu,\nu\}$ and suppose that $\delta(U_p, I) \le \kappa \le 10^{-8}$ for every fine plaquette p with a vertex in one of the blocks at ℓ^∞ distance at most one from $B(y)$, $B(y+e_\mu)$, $B(y+e_\nu)$ or $B(y+e_\mu+e_\nu)$. Then $c = V^\sharp(U)$ satisfies*

$$\delta(c_P, I) \le (4 + 28\pi)\kappa < 92\kappa.$$

Proof. For a pair k among the four sides of P, Lemma 12.34(i) gives $\delta(U_e, I) \le 3\kappa$ for the link of each path inside a block and $\delta(U_e, \ell_k) \le 21\kappa$ for its private link, so $\delta(P_x, \ell_k) \le 24\kappa$ for every path, and by Lemma 12.26 $\|\log(\ell_k^{-1} P_x)\| \le 8\pi\kappa$. The two paths of offset zero contribute nothing, so $c_k = \ell_k M_k$ with $\|\log M_k\| \le \frac{14}{16}\, 8\pi\kappa = 7\pi\kappa$ and $\delta(c_k, \ell_k) \le 7\pi\kappa$. The four path links around P are the holonomy of the square of side two at $2y$: the other four links of its sides are the first links of the straight paths, which are tree links and equal I. That square is tiled by four fine plaquettes, and inserting them with their parallel transports gives $\delta(\ell_P, I) \le 4\kappa$, where ℓ_P is that holonomy. Replacing the four ℓ_k by c_k one at a time, bi-invariance of δ and the triangle inequality add $4 \cdot 7\pi\kappa$. □

With the coarse field calm, the chart has a centre. The next lemma shows that for a pair whose non-path links are near the reconstruction, the constraint $V_k^\sharp = c_k$ determines the path link, on every branch of the cut-off, and the fibre of the window over its non-path links is a single graph.

Lemma 13.3 (the path link is determined)**.** *Let k be a pair with cut-off average c_k, and suppose that each of its seven non-path private links and each link inside $B(y)$ and $B(y+e_\mu)$ is within geodesic distance r of its reconstruction value. Then every path other than the two copies of ℓ_k has $d(P_x, c_k) \le 2r$, and every ℓ_k with $V_k^\sharp = c_k$ satisfies*

$$d(\ell_k, c_k) \le 14r, \qquad d(v_x, I) \le 16r \text{ for every path.}$$

If moreover $r \le 10^{-4}$, the cut-off equals one at every such ℓ_k, and ℓ_k is unique: it is the only fixed point of

$$X \mapsto F(X) = -\frac{1}{16} \sum_x \log(e^{-X} c_k^{-1} P_x), \qquad \ell_k = c_k e^X,$$

in the ball $\|X\| \le 14r$, where the sum runs over the fourteen paths of nonzero offset, and F is a contraction of that ball with constant less than 0.9*.*

Proof. A path of nonzero offset is the product of a link inside a block, within r of I, and a non-path private link, within r of c_k, in one of the two orders; bi-invariance of d and the triangle inequality give $d(P_x, c_k) \le 2r$. If the cut-off is zero, $V_k^\sharp = \ell_k$, so $\ell_k = c_k$. Otherwise every $v_x = \ell_k^{-1} P_x$ has $\delta(v_x, I) < 1$, its principal logarithm is defined and $\|\log v_x\| = d(v_x, I) = d(\ell_k, P_x)$. Put $D = d(\ell_k, c_k)$ and $N = \frac{1}{16}\sum_x \log v_x$, the two terms of offset zero being zero. Then $c_k = \ell_k \exp(\chi N)$ and

$$D = d(e^{\chi N}, I) \le \chi \|N\| \le \frac{14}{16} \max_x d(\ell_k, P_x) \le \frac{7}{8}(D + 2r),$$

so $D \le 14r$, and $d(v_x, I) \le D + 2r \le 16r$. For $r \le 10^{-4}$ this is less than $\frac{1}{2}$, so $\chi = 1$, and $V_k^\sharp = c_k$ reads $X = F(X)$ for $\ell_k = c_k e^X$ with $\|X\| \le 14r$. On that ball each $A_x = c_k^{-1} P_x$ has $d(A_x, I) \le 2r$, so $d(e^{-X} A_x, I) \le 16r$ and $\|F(X)\| \le \frac{14}{16} \cdot 16r = 14r$: F maps the ball into itself. Writing $e^{-X} A_x = I + M$ with $\|M\| \le 16r$, the series of the logarithm gives $\|D \log(I + M)\| \le 1/(1 - 16r)$, and the differential of $X \mapsto e^{-X} A_x$ has norm at most e^{14r}; hence $\|DF\| \le \frac{7}{8} e^{14r}/(1 - 16r) < 0.9$, and the fixed point is unique by the contraction principle. □

Proposition 13.4 (good windows are graphs)**.** *Let W be a finite set of blocks, let c be a coarse field, let K_W be the set of pairs based in W or entering W from a block outside it, and let Z_W be the set of non-path active links with base in $\bigcup W$. Free the links of A_W and*

the path links of the pairs entering W, fix every other link, and condition on $V_k^\sharp = c_k$ for every $k \in K_W$ and on the event that every link of Z_W and every non-path link of a pair entering W is within $r \le 10^{-4}$ of its reconstruction value. Then the fibre of these constraints is the graph of a smooth map Γ from the product of the r-balls of the links of Z_W to the path links of the pairs of K_W, and the conditional law is the Wilson weight times the Haar disintegration of this graph, whose density with respect to the product Haar measure on Z_W is the product over $k \in K_W$ of the inverse Jacobians of $X \mapsto X - F(X)$ at the fixed point.

Proof. The constraint of a pair $k \in K_W$ involves its own private links and the links inside two blocks, so it involves only its own path link, links of Z_W, and fixed links that are within r of their reconstruction values; no other constraint involves a freed link, since the other pairs have their supports off the free set. Lemma 13.3 applies to every $k \in K_W$ with the same r and gives one solution for each pair, a smooth function of the links of Z_W by the implicit function theorem since $I - DF$ is invertible. The fibre is therefore the graph of Γ, and the coarea formula for the submersion $U \mapsto (V_k^\sharp)_k$ restricted to the product of the balls gives the stated density, since the derivative of the constraint of k in its own path link is $I - DF$ in the chart at c_k and the cross derivatives between different pairs vanish. □

The proposition says that, once the coarse field is calm and the labels are placed on the non-path links, the good windows carry the product structure that the proof of Theorem 12.41 needs, with no branch of the cut-off to discard; the Hessian estimate of that theorem then applies on the graph. Two further examples show that the two obvious alternatives fail: labelling all physical links, including the path links, produces a nonconvex domain, and a bad pair can have several path links.

Proposition 13.5 (the all-link label domain is not convex)**.** *Let $c \equiv I$, freeze every link outside the eight private links of one pair at I, and write the seven non-path links as e^{z_j} and the path link as e^X. For every sufficiently small $r > 0$ the set of $(z_1, \dots, z_7)$ for which all eight links lie in the geodesic ball of radius r at I is not convex.*

Proof. With the cut-off equal to one, the constraint is $H(X, z) = X + \frac{1}{8}\sum_{j=1}^{7} \log(e^{-X} e^{z_j}) = 0$, since each non-path link occurs in two of the sixteen paths and the two paths of offset zero contribute nothing. Take $A, B \in \mathfrak{su}(2)$ orthonormal for the Hilbert–Schmidt norm, so that $[A,[A,B]] = -2B$ and $[B,[B,A]] = -2A$, and put $z_1 = -\frac{a}{7}A + hB$, $z_2 = -\frac{a}{7}A - hB$, $z_3 = \cdots = z_7 = -\frac{a}{7}A$ with $a > 0$ small. The inner automorphism of SU(2) that fixes A and negates B and $[A,B]$ exchanges the logarithms of $e^{-xA}e^{z_1}$ and $e^{-xA}e^{z_2}$, so their sum lies along A when $X = xA$, and the equation reduces to its A-component; by the implicit function theorem the solution is $X = x(a,h)A$ with x analytic, even in h by the same symmetry, and $x(a,0) = a$ since $\log(e^{-xA}e^{bA}) = (b-x)A$ gives $x + \frac{7}{8}(b-x) = 0$ at $h = 0$. Let $L(x,b,h)$ be the A-component of $\log(e^{-xA}e^{bA+hB})$. The Baker–Campbell–Hausdorff series through degree three has one term of degree two in h along A, namely $\frac{1}{12}[hB,[hB,-xA]] = \frac{xh^2}{6}A$, so $\partial_h^2 L(x,b,0) = \frac{x}{3} + O((|x|+|b|)^2)$. The A-component of H is $x + \frac{1}{8}(2L(x,-\frac{a}{7},h) + 5(-\frac{a}{7} - x))$, and $\partial_x L(x,b,0) = -1$; differentiating $H = 0$ twice in h at $h = 0$ gives $\frac{1}{8}\partial_h^2 x + \frac{1}{4}\partial_h^2 L = 0$, that is

$$\partial_h^2 x(a,0) = -2\,\partial_h^2 L(a,-\tfrac{a}{7},0) = -\frac{2a}{3} + O(a^2) < 0.$$

Fix r small and η small, and take a with $r < a < r(1 + \frac{\eta^2}{4})$. The points $z^\pm$ with $h = \pm\eta$ and their midpoint z^0 with $h = 0$ have all seven non-path links within $\sqrt{a^2/49 + \eta^2} < r$ of I; the path link of $z^\pm$ is at distance $x(a,\pm\eta) = a - \frac{a}{3}\eta^2 + O(a^2\eta^2 + \eta^4) < r$ from I, while that of z^0 is at distance $x(a,0) = a > r$. So the endpoints lie in the domain and the midpoint does not. □

Proposition 13.6 (a bad pair with three path links)**.** *Let $c \equiv I$, freeze every link outside the eight private links of one pair at I, and take the private links in the one-parameter*

subgroup $\operatorname{diag}(e^{iq}, e^{-iq})$*: one non-path link at the angle* $-\frac{1}{2}$*, the other six at the angle* $\frac{49}{100}$*. Then there are at least three values of the path-link angle* q *with the same cut-off average, and the derivative of the cut-off average with respect to* q *vanishes at some* $q \in (0, \frac{1}{5})$*.*

Proof. On the subgroup all logarithms are scalar and the cut-off average has the angle $f(q) = q + \frac{\chi(q)}{8}(-\frac{1}{2} + 6\cdot\frac{49}{100} - 7q)$, where $\chi(q) = \phi(m(q))$, $m(q) = \max_j 2\sqrt{2}\,|\sin((\theta_j - q)/2)|$ and $\theta_1 = -\frac{1}{2}$, $\theta_2 = \cdots = \theta_7 = \frac{49}{100}$, the factor $2\sqrt{2}|\sin(\theta/2)|$ being the chordal distance of $\operatorname{diag}(e^{i\theta}, e^{-i\theta})$ from I. At $q = 0$ the maximum is attained at θ_1 alone, with $m(q) = 2\sqrt{2}\sin((q + \frac{1}{2})/2)$ near 0 and $\frac{1}{2} < m(0) < 1$, so $\chi = 2 - 2m$ is smooth near 0 with $\chi'(0) = -2\sqrt{2}\cos\frac{1}{4}$, and

$$\begin{aligned} f'(0) &= 1 - \frac{7}{8}\chi(0) + \frac{\chi'(0)}{8}\Big(-\frac{1}{2} + \frac{294}{100}\Big) \\ &= -\frac{3}{4} + \sqrt{2}\Big(\frac{7}{2}\sin\frac{1}{4} - \frac{61}{100}\cos\frac{1}{4}\Big) \end{aligned}$$

after inserting $\chi(0) = 2 - 4\sqrt{2}\sin\frac{1}{4}$. With $\sin\frac{1}{4} < \frac{1}{4}$, $\cos\frac{1}{4} > \frac{31}{32}$ and $1 < \sqrt{2} < \frac{3}{2}$ on the positive and the negative term separately, $f'(0) < -\frac{91}{3200} < 0$. Put $\gamma = f(0)$, which lies in $(0, 1)$. For $|q| = 1$ the maximal chordal distance exceeds one, so $\chi = \pm 1$ gives $f(\pm 1) = \pm 1$. Since f decreases through γ at 0 and $f(-1) < \gamma < f(1)$, the intermediate value theorem gives one preimage of γ in $(-1, 0)$, one in $(0, 1)$, and $q = 0$. On $[0, \frac{1}{5}]$ the maximiser stays θ_1 and $\frac{9}{10} < m < 1$ at $q = \frac{1}{5}$, so $\chi(\frac{1}{5}) < \frac{1}{5}$ and $f'(\frac{1}{5}) > 1 - \frac{7}{40} - 3\cdot\frac{13}{100} > 0$; with $f'(0) < 0$, continuity gives a zero of f' in $(0, \frac{1}{5})$. □

Hence the Jacobian of the path-link constraint, which enters the density of Proposition 13.4, has no positive lower bound on the bad side, and a bad pair contributes a sum over branches to any estimate of the conditional law. On the good side none of this happens: the labels of Proposition 13.4 are placed on the non-path links only.

13.2. **Counting bad labels.** A label is bad when some fine plaquette in a fixed neighbourhood is bad: for the relative link labels this is the content of Theorem 12.27 and Lemma 12.34, and for the coarse-curvature labels it is Lemma 13.2. The chessboard estimate of Theorem 12.57 bounds the probability that a set of fine plaquettes is simultaneously bad by a product. The following lemma transfers such a bound to the labels, and the proposition after it shows that the loss it incurs is the true loss of this kind of argument.

Lemma 13.7 (coverage counting)**.** *Let a family of events* E_y*, indexed by a finite set, be such that each* E_y *forces at least one of a specified set* Q_y *of at most* M *sites to be bad, and suppose that each site lies in at most* H *of the sets* Q_y *and that* $\Pr(\text{all sites of } T \text{ bad}) \le p^{|T|}$ *for every finite set* T *of sites. Then for every nonempty set* S *of* n *labels and* $k = \lceil n/H \rceil$*,*

$$\Pr\Big(\bigcap_{y\in S} E_y\Big) \le \binom{Mn}{k} p^k \le (eMHp)^k,$$

and if $A = eMHp < 1$ *then* $\Pr(\bigcap_{y\in S} E_y) \le A^{n/H}$*.*

Proof. On the intersection, the union of the witness sets Q_y, $y \in S$, contains at least k distinct bad sites, since one site witnesses at most H labels; the union has at most Mn sites. A union bound over its k-subsets and the product bound for each give the first inequality, and $\binom{Mn}{k} \le (eMn/k)^k \le (eMH)^k$ since $n/k \le H$. No choice of witness is required to be measurable, and no independence is used. □

Proposition 13.8 (the exponent $1/H$ cannot be improved)**.** *In the setting of Lemma 13.7 let* H *labels share one site and let each be bad exactly when that site is bad, with probability* p*. Then* $\Pr(\text{all } H \text{ labels bad}) = p$*, so every bound of the form* $q^{|S|}$ *valid for this class of families has* $q \ge p^{1/H}$*.*

Proof. The event that all H labels are bad is the event that the one site is bad. □

For the labels of Section 13.1 the witness sets are boxes of side 7 for the relative link labels, $M = 7^4$, $H = 4^4$, and of side 11 when the coarse-curvature labels are included, $M = 11^4$, $H = 6^4$, with p the chessboard bound of Theorem 12.57 at the threshold of Lemma 13.2. Lemma 13.7 then charges a bad island of n blocks at the rate $A^{n/H}$; the proposition shows that the denominator H is not an artefact of the union bound but the price of knowing only that each label has a witness, and that a better exponent needs joint information on the gauge field.

13.3. **Islands and the environment.** Two bad islands are separated by good blocks, and the question is how the weight of one island depends on the state of the other. The following lemma answers it in terms of a single quantity: the cross-ratio of a kernel on the good layer between them. It integrates arbitrary measures on the two islands, including hard constraints and sums over branches, and their masses do not enter.

Lemma 13.9 (integration through a common kernel)**.** *Let $K(x, y) > 0$ be measurable, let a_0, a_1 and b_0, b_1 be finite positive measures on the spaces of x and of y, and suppose that $Z_{ij} = \iint K\, da_i\, db_j$ is finite and positive for the four pairs $i, j \in \{0, 1\}$. If*

$$e^{-\Delta} \le \frac{K(x_1, y_1)K(x_0, y_0)}{K(x_1, y_0)K(x_0, y_1)} \le e^{\Delta}$$

for all x_i in the support of a_i and y_j in the support of b_j, then

$$\bigl|\log Z_{11} + \log Z_{00} - \log Z_{10} - \log Z_{01}\bigr| \le \Delta.$$

Proof. Multiply the upper inequality by the product $K(x_1, y_0)K(x_0, y_1)$ and integrate against $da_1(x_1)\, da_0(x_0)\, db_1(y_1)\, db_0(y_0)$. By Tonelli the left side integrates to $Z_{11}Z_{00}$ and the right side to $e^{\Delta}Z_{10}Z_{01}$. The lower inequality gives the reverse bound, and the four numbers are positive, so logarithms may be taken. □

Corollary 13.10 (two labels)**.** *Let (A, B) be a pair of binary labels with joint law $\Pr(A = i, B = j) = Z_{ij}/\sum Z$, all $Z_{ij} > 0$, and let Δ bound the connected normaliser as in Lemma 13.9. Then*

$$\bigl|\Pr(B = 1 \mid A = 1) - \Pr(B = 1 \mid A = 0)\bigr| \le \tanh\frac{\Delta}{4},$$

$$|\operatorname{Cov}(A, B)| \le \frac{1}{4}\tanh\frac{\Delta}{4}.$$

Proof. The conditional odds of $B = 1$ given $A = 1$ and given $A = 0$ are Z_{11}/Z_{10} and Z_{01}/Z_{00}, whose logarithms differ by $\log(Z_{11}Z_{00}/Z_{10}Z_{01})$, of modulus at most Δ. For logistic probabilities $\sigma(u) = (1 + e^{-u})^{-1}$ the difference $\sigma(u + s) - \sigma(u)$ is largest at $u = -s/2$ by the symmetry $\sigma(-u) = 1 - \sigma(u)$, where it equals $\sigma(s/2) - \sigma(-s/2) = \tanh(s/4)$, and it increases in s. The covariance is $\Pr(A = 1)\Pr(A = 0)$ times the difference of the conditional probabilities, and $\Pr(A = 1)\Pr(A = 0) \le \frac{1}{4}$. □

When the kernel is the partition function of a calm layer, its cross-ratio is a connected second derivative of a log-normaliser, and that derivative is a covariance under the law of the layer.

Lemma 13.11 (connected normalisers)**.** *Let D be a fixed product of balls in the coordinates z of a window, let $\mathcal{U}(z; \eta, \zeta)$ be smooth in all variables, with η and ζ exterior parameters, and let $Z(\eta, \zeta) = \int_D e^{-\mathcal{U}} dz$. Then for directions u in η and v in ζ,*

$$\partial_u \partial_v \log Z = -\mathbb{E}\bigl[\partial_u \partial_v \mathcal{U}\bigr] + \operatorname{Cov}\bigl(\partial_u \mathcal{U}, \partial_v \mathcal{U}\bigr),$$

expectation and covariance under $e^{-\mathcal{U}} dz/Z$ on D. If $\mathcal{U}$ is a sum of local terms and the supports of η and ζ are farther apart than the range, the first term vanishes; and if the law

on D satisfies

$$|\operatorname{Cov}(F,H)| \le \mathcal{A}\,\|\nabla F\|\,\|\nabla H\|\,e^{-\alpha\,\operatorname{dist}(\operatorname{supp}\nabla F,\operatorname{supp}\nabla H)}$$

for all smooth F and H, then for two segments $\eta_0 \to \eta_1$ and $\zeta_0 \to \zeta_1$ along which the domain D and the bounds persist,

$$\begin{aligned}&\bigl|\log Z(\eta_1,\zeta_1) - \log Z(\eta_1,\zeta_0) - \log Z(\eta_0,\zeta_1) + \log Z(\eta_0,\zeta_0)\bigr| \\ &\qquad \le \mathcal{A} L_\eta L_\zeta\,\|\eta_1-\eta_0\|\,\|\zeta_1-\zeta_0\|\,e^{-\alpha d'},\end{aligned}$$

where L_η, L_ζ bound the gradients in z of $\partial_u\mathcal{U}$ and $\partial_v\mathcal{U}$ per unit of u, v, and d' is the distance between the supports of those gradients.

Proof. Differentiating under the integral sign on the fixed domain D gives $\partial_v \log Z = -\mathbb{E}[\partial_v\mathcal{U}]$ and then the identity, the covariance arising from the derivative of the normalised density. The integrated bound is the identity, the covariance hypothesis, and the fundamental theorem of calculus along the two segments. □

On a good window the law of Proposition 13.4, with the Hessian bound of Theorem 12.41 on the graph, supplies a covariance bound of the required form, so Lemma 13.11 controls the connected normaliser for continuous changes of the sources inside the chart, and Lemma 13.9 then integrates the hard island weights on the two sides. What the three statements do not supply is the kernel bound when a label change moves the domain D, enters an island, or adds a branch of Proposition 13.6; that is the open comparison of Section 13.8. The last result of this subsection shows that the comparison cannot be replaced by rarity, however strong.

Proposition 13.12 (rarity does not control the environment)**.** *Fix $q \in (0,1)$. Let S be a fair bit and, given S, let the labels B_x be independent Bernoulli variables with parameter $q/2$ if $S = 0$ and q if $S = 1$. Then $\Pr(B_x = 1\ \forall x \in T) \le q^{|T|}$ for every finite T, and for all distinct x, y,*

$$\operatorname{Cov}(B_x, B_y) = \frac{q^2}{16}.$$

If the configuration is a function of the labels, every covariance under the law conditioned on all labels vanishes, and the covariance of two observables is the covariance of their conditional expectations. More generally, for a binary label B with $p = \Pr(B = 1 \mid \xi) \in (0,1)$ given the other labels ξ and $\Delta_F = \mu(F \mid B = 1, \xi) - \mu(F \mid B = 0, \xi)$,

$$\operatorname{Cov}(F, \mathbf{1}_B \mid \xi) = p(1-p)\Delta_F,$$

$$\operatorname{Cov}(\mathbb{E}[F \mid B, \xi], \mathbb{E}[H \mid B, \xi] \mid \xi) = p(1-p)\Delta_F\Delta_H.$$

Proof. Given S the labels are independent, so $\Pr(B_x = 1\ \forall x \in T) = \frac{1}{2}((q/2)^{|T|} + q^{|T|}) \le q^{|T|}$, and $\operatorname{Cov}(B_x, B_y) = \operatorname{Var}(\Pr(B_x = 1 \mid S)) = \frac{1}{4}(q - q/2)^2 = q^2/16$ by the conditional covariance formula. The two identities follow by expanding the expectations over the two values of B. □

The example satisfies every finite-intersection rarity bound at every order, yet its labels have a covariance with a positive floor at all distances. A labelling argument must therefore bound $\Delta_F\Delta_H$, the change of the conditional expectations under a label flip, and not only the probability of the flip: small p does not help when $\Delta_F\Delta_H$ does not decay, and normalising by p can cancel it. That bound is the connected normaliser of Lemma 13.9, which is where the island comparison enters.

13.4. **Coupling cost and the comparison lemma.** The proofs of Lemmas 5.4 and B.1 take a coupling that minimises a weighted sum of expected displacements and derive, from its minimality, one inequality per site. The following lemma isolates this step; it needs neither stationarity of the optimal coupling under the local updates nor any property of the updates beyond the two stated ones, and its conclusion is exactly the recurrence that Lemma B.1 iterates.

Lemma 13.13 (minimality gives the recurrence)**.** *Let I be a finite index set, let Π be a set of couplings of two probability measures on a product of metric spaces indexed by I, let $d_i(\pi) = \mathbb{E}_\pi \operatorname{dist}(x_i, x_i')$, and let $w_i > 0$. Suppose $\pi \in \Pi$ minimises $\sum_i w_i d_i$ over Π, and that for each $i \notin S$ there is $\pi^{(i)} \in \Pi$ with $d_j(\pi^{(i)}) = d_j(\pi)$ for $j \neq i$ and $d_i(\pi^{(i)}) \le \sum_j C_{ij} d_j(\pi)$. Then $d_i(\pi) \le \sum_j C_{ij} d_j(\pi)$ for every $i \notin S$. If π is minimal only up to ε_i against $\pi^{(i)}$, then $d_i(\pi) \le \sum_j C_{ij} d_j(\pi) + \varepsilon_i / w_i$. Consequently, if $C_{ij} \ge 0$, $\sum_i w_i C_{ij} \le \theta w_j$ with $\theta < 1$ and $d_i(\pi) \le D$ for $i \in S$, then $\sum_{i \notin S} w_i d_i(\pi) \le \frac{\theta D}{1-\theta} \sum_{j \in S} w_j$.*

Proof. Since $\pi^{(i)}$ changes only the i-th displacement, minimality gives $w_i d_i(\pi) \le w_i d_i(\pi^{(i)}) + \varepsilon_i$, and the bound on $d_i(\pi^{(i)})$ gives the recurrence. The last statement is the arithmetic of Lemma B.1: with $X = \sum_{i \notin S} w_i d_i$ and $Y = \sum_{i \in S} w_i d_i \le D \sum_{i \in S} w_i$, the recurrence and the column condition give $X \le \sum_j d_j \sum_i w_i C_{ij} \le \theta (X + Y)$, so $X \le \theta Y / (1 - \theta)$. □

Lemma 13.14 (minimal couplings exist)**.** *Let μ, ν be probability measures on a compact metric space Ω and c a continuous cost on $\Omega \times \Omega$. The set of couplings of μ and ν is nonempty and compact in the weak topology, and c attains its minimum on it.*

Proof. The product $\mu \otimes \nu$ is a coupling; the marginal conditions are closed in the weak topology, and the probability measures on the compact space $\Omega \times \Omega$ form a compact set, so the couplings form a compact set on which the continuous functional $\pi \mapsto \int c \, d\pi$ attains its minimum. □

The displacements d_i are continuous costs on the compact product of copies of SU(2), so the minimal couplings of Lemma 5.4 and of the block comparison exist by Lemma 13.14, and the coupled update of one site is the choice of $\pi^{(i)}$ in Lemma 13.13. The recurrence, the column-weighted and row-weighted comparisons of Lemma B.1, the decay with a floor of Lemma B.3, and the two lemmas above are also written in Lean 4 in the archive, with Mathlib as the library; the specification of SU(2) enters only through the hypotheses of Lemma 13.13, which the proofs of Sections 5 and 10 supply by hand.

The block response that Lemma 5.4 bounds can also be computed exactly in one direction, which measures the room that the coupling argument leaves.

Proposition 13.15 (the block response at zero field)**.** *Let μ_0 be the law of three unit quaternions (y_1, y_2, y_3) with density proportional to $\exp(\beta_W \operatorname{Re}(y_1 y_2 y_3))$ with respect to Haar measure, and h a unit quaternion. Then $f = \langle y_1, h \rangle + \langle \overline{y_2 y_3}, h \rangle$ is 1-Lipschitz for the sum of the three chordal distances and*

$$\begin{aligned} \operatorname{Cov}_{\mu_0}(f, \langle y_1, h \rangle) &= \frac{1}{4}\Big(1 + \frac{I_2(\beta_W)}{I_1(\beta_W)}\Big) \\ &= \frac{1}{4}\Big(1 + \frac{\beta_W}{4} - \frac{\beta_W^3}{96} + O(\beta_W^5)\Big), \end{aligned}$$

which equals 0.271826 at $\beta_W = 0.351$. Hence the Kantorovich modulus of the block in the field at y_1 is at least this number, and $10\beta_W$ times it exceeds one for $\beta_W > 0.3666$.

Proof. Given y_1, the product $w = y_2 y_3$ is Haar distributed a priori, and the weight is $\exp(\beta_W \langle y_1, \bar{w} \rangle)$ since $\operatorname{Re}(pq) = \langle p, \bar{q} \rangle$ for unit quaternions. So $\bar{w}$ given y_1 follows the von Mises–Fisher law on S^3 with mean direction y_1, whose mean resultant is $\mathbb{E}[\bar{w} \mid y_1] = A(\beta_W) y_1$ with $A = I_2 / I_1$, the ratio of the Bessel functions of orders $\frac{p}{2}$ and $\frac{p}{2} - 1$ for $p = 4$, and the marginal of y_1 is Haar. Multiplication and conjugation preserve the chordal distance, so f changes by at most the sum of the three displacements. Since y_1 is uniform, $\operatorname{Var}\langle y_1, h \rangle = \frac{1}{4}$ and $\operatorname{Cov}(\langle \bar{w}, h \rangle, \langle y_1, h \rangle) = \mathbb{E}[A \langle y_1, h \rangle^2] = A/4$. The series of A follows from $I_\nu(x) = \sum_m (x/2)^{2m+\nu} / (m! \, \Gamma(m + \nu + 1))$, and the derivative of $\mu_b(f)$ in the field b_1 at $b = 0$ in the direction h is the covariance, which bounds the modulus below; the crossing is the root of $\frac{5}{2} \beta_W (1 + A(\beta_W)) = 1$. □

13.5. **From one scale to the next.** A multiscale version of Section 12.3 would eliminate the fine coordinates of a window scale by scale. The following statements say what one elimination step preserves and what it does not, for the quadratic forms that Theorem 12.41 controls at one scale. They are finite matrix lemmas; no all-scale construction is made here.

Lemma 13.16 (weighted Schur complements)**.** *Let $H = \left(\begin{smallmatrix} A & B \\ B^* & D \end{smallmatrix}\right)$ be positive definite in retained and eliminated coordinates, with $S = A - BD^{-1}B^*$ its constrained minimum form on the retained ones. Give both index sets a metric, and for a 1-Lipschitz function ϕ and $\alpha \geq 0$ write $\|M\|_\alpha$ for the supremum over ϕ of the operator norms of $e^{\alpha\phi} M e^{-\alpha\phi}$. If $\|A\|_\alpha \leq a$, $\|B\|_\alpha \leq b$ and $\|D^{-1}\|_\alpha \leq g$, then $\|S\|_\alpha \leq a + b^2 g$, and $|S_{ij}| \leq (a+b^2 g) e^{-\alpha \operatorname{dist}(i,j)}$. If H_0 has blocks A_0, B_0, D_0 with the same bounds and $\|A - A_0\|_\alpha, \|B - B_0\|_\alpha, \|D - D_0\|_\alpha \leq \varepsilon$, then*

$$\|S - S_0\|_\alpha \leq \varepsilon (1 + bg)^2.$$

Proof. Conjugation by $e^{\alpha\phi}$ respects products and inverses, so the weighted norm is submultiplicative on compatible blocks, and $\|B^*\|_\alpha = \|B\|_\alpha$ because $-\phi$ is admissible; the bound on S follows from its formula. For the entry bound take $\phi(x) = \operatorname{dist}(x, j)$. For the difference write $BD^{-1}B^* - B_0 D_0^{-1} B_0^* = (B - B_0) D^{-1} B^* + B_0 D^{-1} (B - B_0)^* + B_0 (D^{-1} - D_0^{-1}) B_0^*$ with $D^{-1} - D_0^{-1} = -D^{-1}(D - D_0) D_0^{-1}$; the four terms are bounded by ε, εbg, εbg and $\varepsilon b^2 g^2$. □

Proposition 13.17 (what form comparison preserves)**.** *Let H_0, H be positive definite with $(1 - \varepsilon) H_0 \leq H \leq (1 + \varepsilon) H_0$, $0 \leq \varepsilon < 1$, and for a surjective linear map Q let $S(v) = \inf_{Qx = v} x^{\mathsf{T}} H x$ and S_0 likewise. Then $(1 - \varepsilon) S_0 \leq S \leq (1 + \varepsilon) S_0$. Form comparison does not preserve locality: on a path of n sites, $H_0 = I$ and $H = I + \varepsilon u u^{\mathsf{T}}$ with $u = (1, \dots, 1)/\sqrt{n}$ satisfy the comparison, while $(H^{-1})_{1n} = -\varepsilon/((1 + \varepsilon) n)$ decays in no exponential way. If instead, for $W = e^{\alpha\phi}$, $\|W H_0^{-1} W^{-1}\| \leq G_0$ and $\|W (H - H_0) W^{-1}\| \leq \varepsilon / G_0$, then $\|W H^{-1} W^{-1}\| \leq G_0/(1 - \varepsilon)$.*

Proof. The comparison of forms passes to infima over the same affine subspaces. The rank-one inverse is $H^{-1} = I - \frac{\varepsilon}{1+\varepsilon} u u^{\mathsf{T}}$, whose off-diagonal entries all equal $-\varepsilon/((1 + \varepsilon) n)$. The last statement is the Neumann series $H^{-1} = \sum_m (-H_0^{-1}(H - H_0))^m H_0^{-1}$ in the weighted norm. □

Repeating the comparison over scales multiplies the factors $(1 \pm \varepsilon_k)$, so the one-step errors must be summable, and Lemma 13.16 identifies the one-step amplification $(1 + b_k g_k)^2$ of the locality norm. Both are conditions that an effective action built from Theorem 12.41 at every scale would have to meet; the theorem at one scale provides the bounds a, b, g for the first step only.

13.6. **Spectral weight at positive time.** Theorem 10.7 makes the nontriviality of the continuum limit equivalent to a ratio of lattice two-point functions staying away from zero. The following two lemmas show that a ratio at a fixed physical time controls the spectral weight of the renormalised observable, so that the ratio criterion is a tightness condition on its spectral measure and not on its unsmeared moments, and the example after them shows that a mass gap by itself does not give nontriviality.

Lemma 13.18 (tightness at positive time)**.** *Let σ_a be positive measures on $[0, \infty)$ with Laplace transforms $C_a(t) = \int e^{-tE} d\sigma_a(E)$, let $s > 0$ be fixed, and suppose $0 < C_a(s/2) < \infty$ and $C_a(s)/C_a(s/2) \geq c > 0$ uniformly in a. Let $d\nu_a = e^{-sE} d\sigma_a / C_a(s)$. Then*

$$\int E \, d\nu_a \leq M_s = \frac{2}{esc}, \qquad \nu_a([R, \infty)) \leq \frac{M_s}{R},$$

$$\frac{C_a(s + t)}{C_a(s)} \geq e^{-M_s t} \quad (t \geq 0).$$

Proof. Since $ue^{-u} \le 1/e$, $Ee^{-sE} \le \frac{2}{es} e^{-sE/2}$; integrating and dividing by $C_a(s)$ gives the moment bound. Markov's inequality gives the tail, and Jensen's inequality for the probability ν_a gives $C_a(s+t)/C_a(s) = \int e^{-tE} d\nu_a \ge \exp(-t \int E \, d\nu_a)$. □

When σ_a is the spectral measure of a centred observable O_a for a nonnegative Hamiltonian, ν_a is the spectral law of the unit vector $e^{-sH_a/2} O_a \Omega_a / C_a(s)^{1/2}$; the lemma keeps its spectral mass from escaping to infinite physical energy. The ratio at the two separations $s/2$ and s is the quantity that Proposition 10.10 bounds by slice correlations.

Lemma 13.19 (nonvanishing limits)**.** *If $\sigma_a([0,\infty)) \ge v > 0$ and $\int E \, d\sigma_a \le M$ uniformly in a, then $C_a(t) \ge v e^{-Mt/v}$ for all $t \ge 0$, and every limit of the $C_a(t)$ is positive.*

Proof. With $v_a = \sigma_a([0,\infty))$, Jensen's inequality for the probability σ_a / v_a gives

$$C_a(t) \ge v_a \exp\Big(-\frac{t}{v_a} \int E \, d\sigma_a\Big) \ge v_a e^{-Mt/v_a},$$

and $x \mapsto x e^{-Mt/x}$ is increasing. □

Proposition 13.20 (a gap without a limit)**.** *Let X be the symmetric two-state Markov process with jump rate $m/2$ in each direction, sampled at times na, and let $O_a = a\,X$. The transfer matrix has eigenvalues 1 and e^{-ma}, so the gap in physical units is m for every a, while $C_a(t) = a^2 e^{-mt} \to 0$ and every moment of O_a tends to zero. The sampled process is reflection positive, and the spectral measure σ_a is the atom $a^2 \delta_m$, whose Laplace transform at $s/2$ and s has the ratio $e^{-ms/2}$, uniform in a; Lemma 13.18 applies, and the limit vanishes because $\sigma_a([0,\infty)) = a^2 \to 0$, which Lemma 13.19 excludes by its hypothesis on the mass.*

Proof. The correlation $\mathbb{E}[X_0 X_t] = e^{-mt}$ for the two-state process, and O_a scales it by a^2; the transfer matrix of the sampled chain is the matrix of its transition probabilities, with the stated eigenvalues. Reflection positivity of a stationary reversible Markov chain is standard. The rest is read off. □

The example separates the two conditions that Theorem 10.7 keeps apart: a uniform gap and a nonvanishing ratio. For a sequence of measures whose mass does not vanish, Lemma 13.19 converts a bound on the first moment at positive time, which Lemma 13.18 derives from one ratio, into a positive lower bound for the limit; the mass itself is the normalisation of Theorem 10.7.

13.7. **The island threshold at one scale.** The counting of Lemma 13.7 and the collar of Proposition 13.4 determine a coupling past which the bad blocks of one scale form islands whose entropy is beaten by their rarity. This subsection computes that coupling. Fix a radius r for the relative link labels and a radius η for the coarse plaquettes: a block y is *bad* when some non-path active link of a pair based at y is at distance more than r from its reconstruction value, or some coarse plaquette with corner y has $\delta(c_P, I) > \eta$, and *good* otherwise. Fix an integer collar $s \ge 1$: a window W is conditioned as in Proposition 13.4 when every block at ℓ^∞ distance at most s from W is good. Join two blocks when they are at ℓ^∞ distance at most $2s+1$, and call a maximal connected set of bad blocks an *island*; two islands are then separated by a layer of good blocks of width at least $2s+1$, so that every block of the layer is at distance more than s from both, and the good layer of Lemma 13.9 is available between them. Each block has $\Delta = (4s+3)^4 - 1$ neighbours in this graph, and the witness sets of Lemma 13.7 are the boxes of side 11, $M = 11^4$, $H = 6^4$.

Theorem 13.21 (the island threshold)**.** *Let $G = \mathrm{SU}(2)$, let $\rho = \min\{r/43, \eta/92\} \le 10^{-8}$, and put*

$$\log q^* = -H\big(\log 2 + 2 \log \Delta\big) - \log(eMH),$$

$$f(\beta) = 6(\log \beta + C_0) - \frac{\beta \rho^2}{16},$$

with C_0 as in Lemma 12.56. The equation $f(\beta) = \log q^*$ has a largest solution $\beta^* = \beta^*(\rho, s) > 96/\rho^2$, and for every $\beta_W \geq \beta^*$:

(i) *every block is bad with probability at most $A = eMH\, q_\rho(\beta_W) \leq (2\Delta^2)^{-H}$;*

(ii) *for every block y and every $n \geq 1$,*

$$\mu_L\{y \text{ lies in an island of at least } n \text{ blocks}\} \leq 2^{1-n}\Delta^{-2} \leq 2^{-n},$$

and the expected number of islands of at least n blocks on a torus of L^4 blocks is at most $L^4\, 2^{-n}/n$;

(iii) *if an estimate charges every bad block a factor $\Gamma \geq 1$ and is summed over the islands through a given block, the sum $\sum_{n\geq 1} \Delta^{2(n-1)}(\Gamma A^{1/H})^n$ is at most 1 as soon as $\beta_W \geq \beta^*(\rho, s, \Gamma)$, the largest solution of $f(\beta) = \log q^* - H \log \Gamma$.*

Numerically, $\beta^*(10^{-8}, 1) = 3.427 \cdot 10^{21}$, $\beta^*(10^{-8}, 5) = 5.401 \cdot 10^{21}$, $\beta^*(10^{-9}, 1) = 3.431 \cdot 10^{23}$, $\beta^*(10^{-9}, 5) = 5.405 \cdot 10^{23}$ and $\beta^*(10^{-8}, 5, 10^{30}) = 1.973 \cdot 10^{22}$; in every case $\beta^* = 16|\log q^*|\rho^{-2}(1+\epsilon)$ with $0 < \epsilon < 2 \cdot 10^{-2}$, and $|\log q^*| = 21090.2$ for $s = 1$ and 33424.9 for $s = 5$.

Proof. Witnesses. Let y be bad because of a non-path link e of the pair $k = (y, \mu)$, and let κ be the largest defect $\delta(U_p, I)$ over the plaquettes with a vertex in a block at ℓ^∞ distance at most one from $B(y)$ or $B(y + e_\mu)$. If $\kappa \leq 10^{-8}$, Lemma 12.34(i) with $W = \{y, y + e_\mu\}$ gives $\delta(U_e, I) \leq 3\kappa$ when e lies inside a block, where $R_e(c) = I$, and $\delta(U_e, \ell_k) \leq 21\kappa$ when e leaves $B(y)$, where $R_e(c) = c_k$ and ℓ_k is the path link; the proof of Lemma 13.2 gives $\delta(c_k, \ell_k) \leq 7\pi\kappa$, so $\delta(U_e, R_e(c)) \leq (21 + 7\pi)\kappa < 43\kappa$. Since the label is bad, either $\kappa > 10^{-8} \geq \rho$ or $43\kappa > r$, and in both cases some plaquette of that set has defect more than ρ. If y is bad because of a coarse plaquette with corner y, Lemma 13.2 gives in the same way a plaquette with defect more than $\min\{10^{-8}, \eta/92\} \geq \rho$ among those with a vertex in a block at distance at most one from the four corner blocks. All these plaquettes lie in cubes whose corners are at ℓ^∞ distance at most five from $2y$, the box of side 11 of the fine lattice, and a cube lies in the box of at most 6^4 blocks. With the sites of Lemma 13.7 the unit cubes and $p = q_\rho(\beta_W)$, Theorem 12.57 gives the product bound, and the lemma gives $\mu_L(\text{all blocks of } S \text{ bad}) \leq A^{n/H}$ for every set S of n blocks when $A = eMHp < 1$, and $\leq A$ for one block.

The threshold. The function f is concave on $(0, \infty)$ with its maximum at $\beta = 96/\rho^2$, where $f = 6\log(96/\rho^2) + 6C_0 - 6 > 0 > \log q^*$; since $f(\beta) \to -\infty$, the equation $f = \log q^*$ has exactly two solutions and the larger one, β^*, exceeds $96/\rho^2$, with $f \leq \log q^*$ on $[\beta^*, \infty)$. For $\beta_W \geq \beta^*$ this gives $q_\rho(\beta_W) \leq e^{f(\beta_W)} \leq q^*$, hence $A \leq eMHq^* = (2\Delta^2)^{-H}$ and $\Delta^2 A^{1/H} \leq \frac{1}{2}$, which is (i).

Islands. A connected set of n blocks containing y in the graph of the collar is recovered from a walk of length $2(n-1)$ around a spanning tree, each step to one of Δ neighbours, so there are at most $\Delta^{2(n-1)}$ of them, and the island of y has at least n blocks only if some connected set of exactly n blocks containing y consists of bad blocks. The probability that a given set of n blocks is bad is at most $A^{n/H}$, so

$$\mu_L\{y \text{ in an island of } n \text{ blocks}\} \leq \Delta^{2(n-1)} A^{n/H} \leq \Delta^{-2}\, 2^{-n},$$

and summing over the sizes $m \geq n$ of the island gives $2^{1-n}\Delta^{-2}$. An island of at least n blocks contains at least n blocks that lie in islands of at least n blocks, so the number of such islands is at most $1/n$ times the number of such blocks, whose expectation is at most $L^4\, 2^{1-n}\Delta^{-2} \leq L^4 2^{-n}$; on a torus the number of neighbours of a block is at most Δ, which is all the count uses. For (iii), replace A by $\Gamma^H A$ in the threshold: then $\Gamma\Delta^2 A^{1/H} \leq \frac{1}{2}$ and the sum is at most $\Delta^{-2}\sum_n 2^{-n} \leq 1$. The numerical values are the roots of $f = \log q^*$ computed to thirty digits, and the two-term expression follows from $f(\beta) = \log q^*$ with $6(\log\beta + C_0) \leq 0.02\,|\log q^*|$ at these values. □

The theorem gives the value asked for in item (iii) of the earlier versions of this section, but only at the block scale and in lattice units: at $\beta_W = 5 \cdot 10^{21}$ the lattice spacing is

far below every physical length, the correlation length of the theory in lattice units is astronomically larger than the side of a block, and the rarity of the islands at that scale says nothing yet about the islands of the effective theory at the scale of the cube, which is what condition (D) needs. The threshold therefore closes the entropy count of one scale, and leaves the comparison between scales to the effective action of Proposition 13.17 and Lemma 13.16. The numbers also show where the loss sits: the factor $16/\rho^2$ comes from the witness radius $\rho \le 10^{-8}$ of Lemma 12.34, and the factor $|\log q^*| \approx 2H \log \Delta$ from the exponent $1/H$, which Proposition 13.8 shows cannot be improved without joint information on the gauge field.

13.8. **What remains.** The results of this section leave the following statements, which together with Theorem 12.41 and Section 12.7 are what the cube condition (D) needs for SU(2) past the Dobrushin ceiling.

(i) *The bad side.* For a pair whose non-path links are bad, a bound on the Kantorovich response of the conditional law of its path links that includes all branches of Proposition 13.6 and the disintegration density, uniform in the exterior.

(ii) *The island comparison.* The hypothesis of Lemma 13.9: that the weight of two bad islands factors through one kernel on the good layer between them, with a cross-ratio controlled by Lemma 13.11 on a domain that the labels of the islands do not move. Proposition 13.12 shows that no rarity bound replaces it.

(iii) *The island threshold.* Theorem 13.21 gives the explicit coupling $\beta^*(\rho, s)$ past which the islands of bad blocks at one scale are rare enough to beat their entropy, at the exponent $1/H$ of Proposition 13.8 and the collar of Proposition 13.4. What is not given is the same statement at the scale of the cube: β^* is of order 10^{21}, far beyond the range in which the chessboard estimate reaches that scale, so that (iv) is needed before the threshold bears on the gap.

(iv) *The scales.* An effective action at every scale whose one-step errors are summable in the sense of Proposition 13.17 and whose locality norms obey Lemma 13.16, down to the last $O(1)$ scales where the running coupling is of order one; this is the large-field expansion of Bałaban, and it is not carried out here.


Items (i) and (ii) are the single-scale part of (D) that remains open; item (iii) is settled at one scale by Theorem 13.21; item (iv) is the multiscale part. On the continuum side, Lemmas 13.18 and 13.19 reduce the nontriviality of the limit to a lower bound on one ratio of lattice two-point functions at a fixed physical time, uniform along the tuning, which is the open input of Theorem 10.7.

Acknowledgements

This paper rests on the work of many earlier scientists. Lattice gauge theory is due to Wilson [83] and its Hamiltonian form to Kogut and Susskind [54]; reflection positivity and the reconstruction of the transfer matrix to Osterwalder and Schrader [65, 66], and their lattice form, with the first proof of a gap at strong coupling, to Osterwalder and Seiler [67, 72]. The uniqueness method is Dobrushin's [30, 31, 32], the curvature method is that of Bakry and Émery [4, 5] in the form of Shen, Zhu and Zhu [74], the gauge fixing on a forest is Creutz's [27], and the curvature of Lie groups is Milnor's [63]; the localisation estimate of Section 12.1 is that of Combes and Thomas [28]. The analysis of the single-link law uses results of Barta [10], Biernacki and Krzyż [20], McShane [62], Stein and Weiss [77] and Watson [81], and the continuum problem is the one formulated by Jaffe and Witten [50] and studied by Bałaban [6, 7, 8].

Declarations

Data and code availability. The proofs are analytic except for the evaluations in Propositions 7.8 and 7.9 and Lemma 8.3 and the numerical bounds of Lemma 5.5, which



label	witness and result
non-path link off $R(c)$	fine defect, Thm. 12.27
coarse plaquette $> \eta$	fine defect, Lem. 13.2
path link	none: determined, Lem. 13.3
all links	nonconvex, Prop. 13.5
good window	graph law, Prop. 13.4
n bad labels	$A^{n/H}$, Lem. 13.7
two islands	$\frac{1}{4}\tanh(\Delta/4)$, Cor. 13.10
rarity alone	floor $q^2/16$, Prop. 13.12
island of n blocks	2^{-n} past β^*, Thm. 13.21

FIGURE 23. Left: a window W of coarse blocks (solid) with its collar (dashed), good blocks in green and bad islands in red. On the good blocks the conditional law is the graph law of Proposition 13.4; the weight of two islands is compared through a kernel on the good layer between them, whose cross-ratio bounds their covariance (Lemma 13.9, Corollary 13.10). Right: the labels considered in Section 13, their fine witnesses, and the estimate or the obstruction that each produces.

are computations in interval arithmetic with rigorous error bounds, and for Proposition 12.5 and Theorem L.5, whose certificates are checked in exact arithmetic. The programs of these steps, in Python with python-flint (FLINT and Arb) and a C kernel for products of Taylor polynomials, their outputs and the list of certified boxes are in the repository `https://github.com/creelie/4Y-M`, release v4.32.0 (doi:10.5281/zenodo.23933856), together with a Lean 4 certificate of the line field, Lean 4 files for the comparison arithmetic and the coupling lemmas of Section 13.4, independent exact or numerical checks of the other finite statements, our checks of the certificate of [69], and the certificate of Proposition 12.5 with its exact check. The code was written by Deep Bhattacharjee, James Montgomery Lockwood and Tristan Romanov.

The computations, the checks and parts of the drafting were done with AI assistance (Claude Opus 5.5, Claude Fable 5.1, GPT-6.1 Sol Ultra and GPT-6 Astra Ultra) under the direction of Deep Bhattacharjee. The combination of the forest gauge with the sharp quarter behind Theorem 1.1 was first assembled by Anaxia, the reasoning engine of the BLLM system [13, 14]; the line field, the free-boundary limit, the ceiling, the curvature bound, the SU(3) part, the plaquette blocks and the certified constants are the authors' own. All four authors approved the final text.

Funding. None.

Competing interests. None.

Ethics approval. Not applicable: the work involves no human participants, animals or personal data.

APPENDIX A. THE LINE FIELD

Table 3 lists, for each residue class of x modulo two, the direction $\sigma(x)$ and the permutation $\pi_x(\mu) = \sigma(x + e_\mu)$. In every row π_x fixes $\sigma(x)$ and is a three-cycle on the other directions, which is the condition of Lemma 2.1.

APPENDIX B. THE COMPARISON LEMMA

Let (G, δ) be a compact group with a bi-invariant metric of diameter D, $\Omega = G^A$, and $C_{ee'}$ the coefficients of Section 3, with row and column sums at most $\alpha < 1$ and $C_{ee'} = 0$

TABLE 3. The line field: x modulo two (digits $x_0x_1x_2x_3$), the direction $\sigma(x)$ and the values $\sigma(x+e_\mu)$ for $\mu=0,1,2,3$.

x	$\sigma(x)$	$\mu=0$	$\mu=1$	$\mu=2$	$\mu=3$
0000	0	0	2	3	1
1000	0	0	3	1	2
0100	2	3	0	2	1
1100	3	2	0	1	3
0010	3	1	2	0	3
1010	1	3	1	0	2
0110	2	1	3	2	0
1110	1	2	1	3	0
0001	1	2	1	3	0
1001	2	1	3	2	0
0101	1	3	1	0	2
1101	3	1	2	0	3
0011	3	2	0	1	3
1011	2	3	0	2	1
0111	0	0	3	1	2
1111	0	0	2	3	1

unless e, e' share a plaquette, so that $|e-e'|\le 1$ for the distance of link midpoints. Fix $\lambda>0$ with $\theta=\alpha e^{\lambda}<1$.

Lemma B.1. *Let $S\subset A$, possibly infinite, and let μ_1,μ_2 be probability measures on Ω invariant under γ_e^F for every $e\notin S$. Let $T\subset A\setminus S$ be finite, $w_b=e^{-\lambda\,\mathrm{dist}(b,T)}$, and suppose $\sum_{z\in S}w_z<\infty$. There is a coupling of μ_1,μ_2 whose link distances $\varrho(b)=\mathbb{E}\,\delta(\sigma_b,\sigma'_b)$ satisfy*

$$\sum_{b\notin S} w_b\varrho(b)\le \frac{\theta D}{1-\theta}\sum_{z\in S} w_z.$$

If $S=\emptyset$ then $\mu_1=\mu_2$.

Proof. The couplings form a weakly compact set and $\Psi(\pi)=\sum_b w_b\mathbb{E}_\pi\delta(\sigma_b,\sigma'_b)$ is weakly continuous because $\sum_b w_b<\infty$; let π^* be a minimiser. For $e\notin S$ replace the pair (σ_e,σ'_e) by a sample of an optimal coupling of $\gamma_e^F(\cdot\mid\sigma)$ and $\gamma_e^F(\cdot\mid\sigma')$, chosen measurably [80, Cor. 5.22]. Both μ_i are invariant, so the result is again a coupling, and changing the finitely many links that enter γ_e^F one at a time gives $W_1\le\sum_{e'}C_{ee'}\delta(\sigma_{e'},\sigma'_{e'})$. Minimality gives $\varrho(e)\le\sum_{e'}C_{ee'}\varrho(e')$. Multiply by w_e and sum over $e\notin S$; where $C_{ee'}\neq0$, $w_e\le e^{\lambda}w_{e'}$, so

$$\sum_{e\notin S} w_e\varrho(e)\le\theta\Big[\sum_{e'\notin S} w_{e'}\varrho(e')+\sum_{z\in S} w_z\varrho(z)\Big],$$

all sums are finite, and $\varrho\le D$ on S. If $S=\emptyset$, $\varrho\equiv0$. □

Corollary B.2. *Let μ be the Gibbs measure of γ^F, F a local function with support S_F, L_F-Lipschitz in each link, and H bounded with finite support S_H disjoint from S_F. Then*

$$|\operatorname{Cov}_\mu(F,H)|\le\|H\|_\infty L_F\frac{\theta D}{1-\theta}|S_H|\,e^{-\lambda\,\mathrm{dist}(S_F,S_H)}.$$

Proof. Condition on the links of S_H [53, Ch. 3]; the conditional measures $\mu^\sigma=\mu(\cdot\mid\sigma_{S_H})$ are invariant under γ_e^F for $e\notin S_H$. Lemma B.1 with $S=S_H$ and $T=S_F$, for which $w_b=1$ on S_F and $w_z\le e^{-\lambda\,\mathrm{dist}(S_F,S_H)}$ on S_H, gives $|\mu^\sigma(F)-\mu^{\sigma'}(F)|\le L_F\sum_{b\in S_F}\varrho(b)\le L_F\frac{\theta D}{1-\theta}|S_H|e^{-\lambda\,\mathrm{dist}(S_F,S_H)}$ for any two boundary values. Since H depends only on σ_{S_H}, $\operatorname{Cov}_\mu(F,H)=\iint H(\sigma)(\mu^\sigma(F)-\mu^{\sigma'}(F))\,d\mu(\sigma)\,d\mu(\sigma')$, and the bound follows. □

Lemma B.3. *Let the blocks partition A, let C_{Bj} be the block coefficients of Section 5, with $\sum_B C_{Bj} \le \alpha < 1$ for every j and $C_{Bj} = 0$ unless every link of B lies within distance 2 of j, and fix $\lambda > 0$ with $\theta = \alpha e^{2\lambda} < 1$. Let $S \subset A$ be a union of blocks and let μ_1, μ_2 be invariant under $\gamma_B^{\mathcal{B}}$ for every block $B \not\subset S$. With T, w_b as in Lemma B.1 and $\sum_{z\in S} w_z < \infty$, there is a coupling with $\sum_{b\not\subset S} w_b \varrho(b) \le \frac{\theta D}{1-\theta} \sum_{z\in S} w_z$, and $\mu_1 = \mu_2$ if $S = \emptyset$.*

Proof. Take a minimiser π^* of Ψ as before. For a block $B \not\subset S$ replace (σ_B, σ'_B) by an optimal coupling, for the metric d, of the two block laws; it is again a coupling, and changing the boundary links one at a time bounds the expected new d-distance by $\sum_j C_{Bj}\varrho(j)$. Put $\bar w_B = \max_{i\in B} w_i$. Minimality gives

$$\sum_{i\in B} w_i \varrho(i) \le \bar w_B \sum_j C_{Bj}\varrho(j),$$

and $\bar w_B \le e^{2\lambda} w_j$ whenever $C_{Bj} \ne 0$. Summing over the blocks outside S and using the column sums,

$$\sum_{b\not\subset S} w_b \varrho(b) \le \theta\Big[\sum_{j\not\in S} w_j \varrho(j) + \sum_{z\in S} w_z \varrho(z)\Big],$$

which gives the claim as in Lemma B.1. □

Lemma B.4. *Let $n \ge 2$ be even, let V_x, $x \in 2\mathbb{Z}^4$, be the translates of Section 10, and suppose that $D(n,\alpha)$ holds.*

(a) *If a probability measure μ on G^A is invariant under γ_e^F for every e in a finite set V, it is invariant under γ_V.*

(b) *Fix $\lambda > 0$ with $\theta = \alpha e^{\lambda\rho_n} < 1$. Let $S \subset A$, and let μ_1, μ_2 be invariant under γ_{V_x} for every x with $V_x \cap S = \emptyset$. With T, w_b as in Lemma B.1 and $\sum_{z\in S} w_z < \infty$, there is a coupling with*

$$\sum_{b\not\subset S} w_b \varrho(b) \le \frac{D(1 + c_n e^{\lambda\rho_n})}{1-\theta} \sum_{z\in S} w_z,$$

and $\mu_1 = \mu_2$ if $S = \emptyset$.

Proof. (a) Let ν_η be the conditional law of Q_V given the links η outside V. For bounded f on G^V and bounded g of η, $\gamma_e^F(fg) = g\,\gamma_e^F f$ for $e \in V$, so invariance gives $\int g(\eta)\big(\nu_\eta(\gamma_e^F f) - \nu_\eta(f)\big)\,d\mu = 0$ for all g; taking f in a countable dense set of continuous functions, ν_η is invariant under every $\gamma_e^F(\cdot \mid \cdot, \eta)$, $e \in V$, for μ-almost every η. So is $\gamma_V(\cdot \mid \eta)$, whose conditional laws of single links are the γ_e^F. The composition P_η of the kernels γ_e^F, $e \in V$, in a fixed order has a density with respect to Haar measure on G^V bounded below by a positive constant, because each single-link density is at least $e^{-12N^2\beta}$ times its maximum over the group. A Markov kernel with such a lower bound has at most one invariant probability measure (Doeblin's condition), so $\nu_\eta = \gamma_V(\cdot \mid \eta)$.

(b) Let π^* minimise Ψ as in Lemma B.1 and call x admissible if $V_x \cap S = \emptyset$. For admissible x replace $(\sigma_{V_x}, \sigma'_{V_x})$ by an optimal coupling, for d_{V_x}, of the two block laws; both μ_i are invariant, so the result is a coupling, and changing the boundary links one at a time bounds the new expected d_{V_x}-distance by $\sum_j C_{V_x,j}\varrho(j)$. With $\bar w_x = \max_{i\in V_x} w_i$, minimality gives

$$\sum_{i\in V_x} w_i \varrho(i) \le \bar w_x \sum_{j\not\in V_x} C_{V_x,j}\varrho(j),$$

and $\bar w_x \le e^{\lambda\rho_n} w_j$ whenever $C_{V_x,j} \ne 0$. Let S' be the set of links outside S that lie in a translate meeting S; each is within distance ρ_n of a link of S, so $\sum_{i\in S'} w_i \le c_n e^{\lambda\rho_n} \sum_{z\in S} w_z$. A link outside $S \cup S'$ lies in N_n translates, all admissible. Summing over admissible x and using $\sum_{x:\, j\not\in V_x} C_{V_x,j} = K_{c(j)} \le \alpha N_n$,

$$N_n \sum_{i\not\in S\cup S'} w_i \varrho(i) \le \theta N_n \Big[\sum_{i\not\in S\cup S'} w_i \varrho(i) + \sum_{i\in S\cup S'} w_i \varrho(i)\Big].$$

All sums are finite and $\varrho \le D$, so $\sum_{i\notin S\cup S'} w_i\varrho(i) \le \frac{\theta D}{1-\theta}(1 + c_n e^{\lambda\rho_n})\sum_{z\in S} w_z$; adding $\sum_{i\in S'} w_i\varrho(i) \le Dc_n e^{\lambda\rho_n}\sum_{z\in S} w_z$ gives the bound. If $S = \emptyset$ then $S' = \emptyset$ and $\varrho \equiv 0$. □

The covariance bound of Corollary B.2 follows from part (b) in the same way, with $\theta D/(1-\theta)$ replaced by $D(1 + c_n e^{\lambda\rho_n})/(1-\theta)$, since the conditional measures μ^σ there are invariant under γ_{V_x} for $V_x \cap S_H = \emptyset$ by part (a).

Appendix C. The plaquette blocks

Table 4 lists the block plaquettes of Section 5 modulo $2\mathbb{Z}^4$: the plaquette at x in the plane $\{\mu,\nu\}$ is a block plaquette exactly when x modulo two and $\{\mu,\nu\}$ appear in the table. Eight lie in the plane $\{0,1\}$ and eight in $\{2,3\}$; together they contain the 48 active links of a cell of side two, each once.

Table 4. The sixteen block plaquettes modulo two: the base point x (digits $x_0x_1x_2x_3$ modulo two) and the plane.

x	plane	x	plane
0000	$\{0,1\}$	0000	$\{2,3\}$
0001	$\{0,1\}$	0011	$\{2,3\}$
0010	$\{0,1\}$	0100	$\{2,3\}$
0011	$\{0,1\}$	0111	$\{2,3\}$
1100	$\{0,1\}$	1000	$\{2,3\}$
1101	$\{0,1\}$	1011	$\{2,3\}$
1110	$\{0,1\}$	1100	$\{2,3\}$
1111	$\{0,1\}$	1111	$\{2,3\}$

In closed form, the plaquette at x in $\{0,1\}$ is a block plaquette when $x_0 \equiv x_1$ and the plaquette at x in $\{2,3\}$ when $x_2 \equiv x_3$; the blocks do not depend on the line field beyond the fact that every plaquette contains one fixed link. The bounds of Lemma 5.5 are evaluated in ball arithmetic at 128 bits: the series of Z_0 and of its first three derivatives in $s = r^2$ is summed to forty-eight terms, and the tail is bounded for $s \le 5$ by twice the first omitted term, because consecutive terms decrease by a factor below one half. The program also writes the constants, rounded outward to 10^{-12}, for a check of the chain (5.2) at $\beta_W = \frac{1}{3}$, 0.345, 0.35, 0.351 in exact rational arithmetic by the Lean kernel (`gauge-fixed/blocks/sharp`). The constants of (5.2) and α are then evaluated at a given β_W from exact input.

For SU(3) the bounds of Lemma 8.3 are evaluated on the 841 boxes of side $\frac{3}{16}$ at $r = \frac{3}{2}$ and the 784 boxes of side $\frac{19}{100}$ at $r = \frac{38}{25}$ and the 826 boxes of side $\frac{77}{400}$ at $r = \frac{77}{50}$ with the moments of Appendix E, and the constants of (8.2) and α_3 are evaluated from these bounds and the cover of Proposition 7.8 or Proposition 7.9 in ball arithmetic at 120 bits. The cover of Proposition 7.9 is computed as in Appendix E with $r = \frac{38}{25}$ and the threshold 1.03, and with $r = \frac{77}{50}$ and the threshold 1.0275.

Appendix D. The blocks of Lemma 6.2

Diagonal blocks. Let $w_a = e^{i\phi}$. On the diagonal part, $f_p = -\cos\phi\,(y_1 + y_2 - y_3 - y_4)^2$ and $c_p = 2(y_1y_4 - y_1y_2 + y_2y_3 - y_3y_4)$, so

$$\mathcal{F} = \sum_i \alpha_i y_i^2 + \cos\phi\,(y_1 + y_2 - y_3 - y_4)^2 - \tfrac{9}{5}(y_1y_4 - y_1y_2 + y_2y_3 - y_3y_4)$$

with $y_k = 0$ and weights $\alpha_{k\pm1} = \frac{21}{10}$, $\alpha_{k+2} = \frac{5}{4}$. It is affine in $\cos\phi$, so positivity at $\cos\phi = \pm1$ suffices. For $k = 1$ and $\cos\phi = 1$ the matrix in (y_2, y_3, y_4) is

$$\begin{pmatrix} \frac{31}{10} & -\frac{19}{10} & -1 \\ -\frac{19}{10} & \frac{9}{4} & \frac{19}{10} \\ -1 & \frac{19}{10} & \frac{31}{10} \end{pmatrix},$$

and Table 5 lists the leading principal minors for every position; all are positive.

TABLE 5. Leading principal minors of the diagonal blocks of Lemma 6.2.

Fixed link	$\cos\phi = 1$			$\cos\phi = -1$		
e_1, e_4	$\frac{31}{10}$,	$\frac{673}{200}$,	$\frac{8421}{2000}$	$\frac{11}{10}$,	$\frac{53}{200}$,	$\frac{21}{2000}$
e_2	$\frac{31}{10}$,	$\frac{861}{100}$,	$\frac{8421}{2000}$	$\frac{11}{10}$,	$\frac{21}{100}$,	$\frac{21}{2000}$
e_3	$\frac{9}{4}$,	$\frac{673}{200}$,	$\frac{8421}{2000}$	$\frac{1}{4}$,	$\frac{53}{200}$,	$\frac{21}{2000}$

Pair blocks. Let $w_a = e^{i(\chi+\vartheta)}$ and $w_b = e^{i(\chi-\vartheta)}$. The terms of f_p carry $w_a + \bar{w}_b = 2\cos\chi\, e^{i\vartheta}$ or its conjugate, and c_p carries $w_a\bar{w}_b = e^{2i\vartheta}$, so the Hermitian matrix $M(m,\vartheta)$ of $\mathcal{F}$, $m = \cos\chi$, is affine in m with $M(-1,\vartheta) = M(1,\vartheta+\pi)$, and $m = 1$ suffices. For the fixed link e_2, in the variables (z_1, z_3, z_4) and with $c = \cos\vartheta$,

$$M = \begin{pmatrix} \frac{21}{5} + 2c & -2e^{i\vartheta} & -\frac{9}{5} - 2e^{i\vartheta} \\ -2e^{-i\vartheta} & \frac{21}{5} + 2c & \frac{9}{5} + 2e^{-i\vartheta} \\ -\frac{9}{5} - 2e^{-i\vartheta} & \frac{9}{5} + 2e^{i\vartheta} & \frac{5}{2} + 2c \end{pmatrix}.$$

With $u = \cos^2(\vartheta/2) \in [0,1]$ its leading principal minors are

$$\tfrac{20u+11}{5}, \quad \tfrac{(20u+1)(20u+21)}{25}, \quad \tfrac{400u(40u^2-23u+4)+21}{250},$$

and $40u^2 - 23u + 4$ has negative discriminant $529 - 640$. The other positions give the same determinant and the leading minors $\frac{20u+11}{5}$ or $\frac{8u+1}{2}$ and $\frac{800u^2-180u+53}{50}$, whose discriminant $32400 - 169600$ is negative. Every block is positive definite.

The inequality is not tight. In floating point the least eigenvalue of the whole form over $W \in \mathrm{U}(N)$, $N \le 4$, is 0.0149, which is also the least eigenvalue of the exact blocks; this slack is what the choice of the weights $\frac{21}{10}$, $\frac{5}{4}$, $\frac{9}{10}$ trades against the constant $\frac{29}{2}$.

APPENDIX E. THE CERTIFIED COMPUTATION

Truncation. For a corner b of a box the moments of Lemma 7.6 are derivatives of Z at $(b, \bar{b})$. The truncated series is $\mathbb{E}_H[A_t(UD)B_t(\bar{U}D')]$ with A_t the part of $\sum_\lambda c_\lambda s_\lambda(UD)$ over partitions with $\lambda_1 - \lambda_3 \le 34$ and $\lambda_3 \le 11$, and B_t likewise. Every omitted partition has $|\lambda| > 34$, because $3 \cdot 12 > 34$, and the omitted part is at most $T(\|D\|) = \sum_{n>34}(\frac{3}{2}\|D\|)^n/n!$ in modulus. On the polydisc $|D_{ij} - b_{ij}| \le \rho$, $\|D\| \le \|b\| + 3\rho$, and Cauchy's estimate bounds the α-th derivative of the omitted part by $\alpha!\rho^{-|\alpha|}T(\|b\| + 3\rho)$. The derivatives of the full exponential are at most $2^{-|\alpha|}e^{v/2}$ with $v = \sum_k |b_k|$, since $|\operatorname{Tr}(Ub)| \le \sum_k |b_k|$ for diagonal b. With $\varepsilon_\alpha = 2^{|\alpha|}\alpha!\rho^{-|\alpha|}T(\|b\| + 3\rho)$ the truncated moment differs from the true one by at most $\varepsilon_\alpha e^{v/2} + (e^{v/2} + \varepsilon_\alpha)\varepsilon_\beta$, and ρ is chosen to minimise $\rho^{-6}T$. The largest error bound of an unnormalised moment at any corner is 1.7×10^{-11}.

Weight zero. Conjugation by the diagonal torus T preserves Haar measure and $\operatorname{Re}\operatorname{Tr}(Ub)$ for diagonal b, and multiplies u_{ij} by $t_i t_j^{-1}$; a moment of a monomial on which T acts by a nontrivial character is therefore 0, and only 8278 of the 134596 monomials of degree at most six have the trivial character. The functions $\operatorname{Tr}(X)$, $e_2(X)$, $\det X$ of $X = DD'^{\top}$ and $\det D$, $\det D'$ are invariant under the same action on the variables, so after these are formed in the full ring of Taylor polynomials of degree six in the 18 variables, the series is computed in the subring of weight-zero monomials.

Rounding. The truncated series is a Taylor polynomial of total degree six in the 36 real directions of $(D-b, D'-\bar{b})$, computed in double precision. Every operation is repeated on absolute values and carries an operation count n, so that the computed coefficient differs from the exact one by at most $\gamma_n = nu/(1-nu)$, $u = 2^{-53}$, times the absolute-value coefficient [45, Sec. 3.1]. With C in $2^{-30}\mathbb{Z}$, the polynomials $2^{31}q_h$, $X_a(2^{31}q_h)$, $12{\cdot}2^{31}(\varphi_h{+}\Delta q_h)$ and $12\cdot 2^{31}\langle\nabla\varphi_{F_k},\nabla q_h\rangle$ for the coordinate vectors h have coefficients in $\mathbb{Z}[i]$, and every partial sum in their computation is bounded, through the same operations on absolute values, by less than 2^{50}; they are therefore computed exactly. Normalisation by Z, the products with the corner coordinates, the quadratic forms and their error matrices are formed with the same bookkeeping, and the largest eigenvalues are bounded in ball arithmetic [51].

Separation. The functional c of a discarded box is the dual solution of a linear program for the gauge of the centre of the box with respect to a 72×72 grid of T. Only the final inequality is used: $\max_T \ell$ is bounded by its maximum over a 512×512 grid plus $\frac{1}{2}H\delta^2/2$, with $\delta = 2\pi/512$ and $H = |c_1| + |c_2| + 2|c_3|$ a bound for its Hessian on the torus, and the inequality is checked with relative margin 10^{-12}.

Appendix F. The transfer matrix

This appendix recalls the Osterwalder–Schrader reconstruction for the lattice [65, 66, 67] in the form used for the gaps, following our earlier paper [15].

Let ω be a state on the gauge-invariant local functions of $\mathbb{Z}^4$ that is invariant under translations and the hypercubic group and reflection positive for reflections in planes through sites and in planes bisecting links; by Proposition 3.2(iii)–(iv) and its analogues in the other theorems, the states of this paper are of this kind. The reconstruction gives a Hilbert space with a vector Ω, a map $F\mapsto\Psi_F$ with dense range, and a self-adjoint $\mathbb{T}$ with $0\le\mathbb{T}\le 1$, $\mathbb{T}\Omega=\Omega$ and $\langle\Psi_F,\mathbb{T}^t\Psi_H\rangle = \omega(\vartheta(\bar{F})\tau_t H)$, where ϑ is the time reflection and τ_t the time translation.[24]

Proposition F.1. *For $m>0$ the following are equivalent:*

(a) *the spectrum of $\mathbb{T}$ on $\Omega^\perp$ lies in $[0,e^{-m}]$;*

(b) *for every F in a class of gauge-invariant local functions supported at times in a bounded interval and dense in $L^2(\omega)$ on each such support, there is $C_F<\infty$ with $0\le\omega(\vartheta(\bar{F})\tau_t F)-|\omega(F)|^2\le C_F e^{-mt}$ for all $t\ge 0$.*

Under (a) *one may take $C_F=\|F\|_\infty^2$ for every bounded F.*

Proof. If (a) holds, then

$$\omega(\vartheta(\bar{F})\tau_t F)-|\omega(F)|^2 = \langle\Psi'_F,\mathbb{T}^t\Psi'_F\rangle$$

with $\Psi'_F=\Psi_F-\langle\Omega,\Psi_F\rangle\Omega\in\Omega^\perp$; this number lies in $[0,e^{-mt}\|\Psi_F\|^2]$, and $\|\Psi_F\|^2\le\|F\|_\infty^2$. Conversely, let σ be the spectral measure of $\mathbb{T}$ at Ψ'_F and let $\delta>0$. Then

$$\sigma\big((e^{-m+\delta},1]\big)e^{(-m+\delta)t}\le C_F e^{-mt}$$

for all t, so $\sigma((e^{-m+\delta},1])=0$. The spectral projection of $(e^{-m},1]$ restricted to $\Omega^\perp$ therefore annihilates the vectors Ψ'_F of the class, which are dense by the footnote above, and it vanishes. □

The class in (b) cannot be replaced by one observable. On $\mathbb{C}^3$ let $\mathbb{T}=\operatorname{diag}(1,e^{-m},e^{-m/2})$, $\Omega=e_1$ and $\Psi'_F=e_2$: the function in (b) is e^{-mt}, while the spectrum on $\Omega^\perp$ reaches $e^{-m/2}$. The decay of one two-point function places only the spectral measure of Ψ'_F in $[0,e^{-m}]$, and the gap follows when such vectors are dense in $\Omega^\perp$, as in (b) and in Theorem 10.5.

[24] Ψ_F is the class of F in the completion of the positive-time local functions for the form $\langle F,H\rangle = \omega(\vartheta(\bar{F})H)$, which is positive semidefinite by reflection positivity. Since $|\omega(\vartheta(\bar{F})F)|\le\omega(|F|^2)$ by the Cauchy–Schwarz inequality and the invariance of ω under ϑ, functions that are close in $L^2(\omega)$ give close vectors.

Proposition 8.8 of [35] draws the gap on $\Omega^{\perp}$ from the two-point function of one fixed observable.

Appendix G. Spectral measures and Laplace transforms

This appendix collects the facts on spectral measures that the continuum step uses, with proofs.

Lemma G.1. *Let T be a self-adjoint operator on a Hilbert space with $0 \le T \le 1$ and let ψ be a unit vector. There is a unique probability measure ν on $[0,1]$ with $\langle \psi, h(T)\psi\rangle = \int h\, d\nu$ for every bounded Borel function h on $[0,1]$. If $T\psi' = \psi'$ implies $\langle \psi', \psi\rangle = 0$, then $\nu(\{1\}) = 0$. If μ is the image of ν under $\lambda \mapsto -c^{-1}\log\lambda \in [0,\infty]$ for some $c > 0$, then $\langle \psi, T^{\sigma/c}\psi\rangle = \int_{[0,\infty]} e^{-\sigma E} d\mu(E)$ for $\sigma > 0$, with the integrand 0 at $E = \infty$.*

Proof. The spectral theorem [68, Sec. VII.2] gives the measure ν, carried by the spectrum of T, which lies in $[0,1]$; uniqueness holds because the moments $\int \lambda^n d\nu = \langle \psi, T^n\psi\rangle$ determine a measure on a compact interval, polynomials being dense in $C[0,1]$. The spectral projection $P = \mathbf{1}_{\{1\}}(T)$ projects onto the eigenvectors of T with eigenvalue 1, so $\nu(\{1\}) = \|P\psi\|^2 = \langle P\psi, \psi\rangle = 0$ under the hypothesis. The last identity is the change of variables $\lambda^{\sigma/c} = e^{-\sigma E}$ for $\lambda = e^{-cE}$, with $\lambda = 0$ corresponding to $E = \infty$. □

Lemma G.2. *Every sequence of probability measures on a compact metric space K has a subsequence that converges weakly, that is, against every continuous function, to a probability measure on K.*

Proof. $C(K)$ is separable; let (h_j) be dense. The numbers $\int h_j\, d\mu_k$ are bounded by $\|h_j\|_\infty$, so a diagonal subsequence makes all of them converge. By density the limit $\Lambda(h) = \lim \int h\, d\mu_k$ exists for every $h \in C(K)$; Λ is linear, positive and $\Lambda(1) = 1$, so by the Riesz representation theorem it is integration against a probability measure. □

Lemma G.3. *Let μ be a probability measure on $[0,\infty]$ and $\rho(\sigma) = \int e^{-\sigma E} d\mu(E)$ for $\sigma > 0$.*

(a) *ρ is nonincreasing and log-convex: $\rho(\sigma)^2 \le \rho(\sigma - s)\rho(\sigma + s)$ for $0 < s < \sigma$.*
(b) *$\rho(\sigma) > 0$ for one $\sigma > 0$ if and only if $\mu([0,\infty)) > 0$, and then $\rho(\sigma) > 0$ for every $\sigma > 0$.*
(c) *$\rho(\sigma) \to \mu([0,\infty))$ as $\sigma \downarrow 0$.*
(d) *If μ is carried by $[m,\infty]$, then $\rho(\sigma) \le e^{-m\sigma}$.*
(e) *For $M > 0$, $\mu([0,M]) \ge \rho(\sigma) - e^{-\sigma M}$.*
(f) *If $\mu_k \to \mu$ weakly on $[0,\infty]$, then $\int e^{-\sigma E} d\mu_k \to \rho(\sigma)$ for every $\sigma > 0$.*

Proof. (a) The integrand decreases in σ. By the Cauchy–Schwarz inequality,

$$\Big(\int e^{-\sigma E} d\mu\Big)^2 \le \int e^{-(\sigma-s)E} d\mu \int e^{-(\sigma+s)E} d\mu.$$

(b) The integrand is positive exactly on $[0,\infty)$. (c) The integrand increases to $\mathbf{1}_{[0,\infty)}$ as $\sigma \downarrow 0$, and monotone convergence applies. (d) The integrand is at most $e^{-m\sigma}$ on $[m,\infty]$. (e) Split the integral at M: it is at most $\mu([0,M]) + e^{-\sigma M}$. (f) For $\sigma > 0$ the function $E \mapsto e^{-\sigma E}$, extended by 0 at ∞, is continuous on $[0,\infty]$. □

By (b) and (e), a lower bound $\rho(\sigma) \ge c > 0$ at one σ places mass at least $c/2$ on the bounded interval $[0, \sigma^{-1}\log(2/c)]$; for the measures μ_k of Theorem 10.7 a bound uniform in k therefore makes them tight on $[0,\infty)$ with a fixed fraction of their mass.

Part (b) also excludes a positive Laplace representation for a spectral cutoff of compact support. If $\chi(\lambda) = \int_{[0,\infty)} e^{-s\lambda} d\nu(s)$ with ν a positive measure and $\chi(0) = \nu([0,\infty)) > 0$, then (b), applied to $\nu/\chi(0)$, gives $\chi(\lambda) > 0$ for every $\lambda > 0$, so χ vanishes nowhere. The cutoff χ_σ of [35, Sec. 9, Eqs. (9.11), (9.12)] equals 1 on $[0,\sigma]$ and 0 on $[2\sigma,\infty)$ and is asserted to have such a representation.

Appendix H. The character expansion

Let G be a compact connected Lie group with normalised Haar measure dU. For an irreducible unitary representation ρ_λ of dimension d_λ write $\chi_\lambda = \operatorname{Tr}\rho_\lambda$. The characters form an orthonormal basis of the square-integrable class functions (Peter–Weyl and Schur orthogonality [23, Chs. 2–4]).

Lemma H.1. *Let $w \in L^1(G)$ be a class function with $w(U^{-1}) = w(U)$, and put*

$$\hat{c}_\lambda = d_\lambda^{-1} \int w\,\chi_\lambda\,dU.$$

Then $\int w(U)\rho_\lambda(U)\,dU = \hat{c}_\lambda I$, and $\int w(U)\chi_\lambda(U^{-1}V)\,dU = \hat{c}_\lambda\chi_\lambda(V)$.

Proof. $M = \int w(U)\rho_\lambda(U)dU$ commutes with every $\rho_\lambda(V)$, since $w(VUV^{-1}) = w(U)$ and Haar measure is invariant; by Schur's lemma $M = cI$, and taking traces gives $c = \hat{c}_\lambda$. Then $\int w(U)\chi_\lambda(U^{-1}V)dU = \operatorname{Tr}(\int w(U)\rho_\lambda(U^{-1})dU\,\rho_\lambda(V))$, and the substitution $U \mapsto U^{-1}$ with $w(U^{-1}) = w(U)$ turns the integral into M. □

Lemma H.2. *Let $G = \mathrm{SU}(N)$ and $\beta > 0$, and let*

$$w_\beta(U) = e^{\frac{\beta}{N}\operatorname{Re}\operatorname{Tr}U} \Big/ \int e^{\frac{\beta}{N}\operatorname{Re}\operatorname{Tr}V} dV.$$

Then $\hat{c}_\lambda(\beta) = d_\lambda^{-1}\int w_\beta\chi_\lambda\,dU$ is real, $0 \le \hat{c}_\lambda(\beta) \le 1$, and $\hat{c}_\lambda(\beta) < 1$ for $\lambda \neq 0$. Moreover

$$\hat{c}_\lambda(\beta) = 1 - \frac{N\,C_2(\lambda)}{\beta} + O(\beta^{-2}) \qquad (\beta \to \infty),$$

where $-\sum_a \rho_\lambda(T_a)^2 = 2C_2(\lambda)I$ for a basis (T_a) of $\mathfrak{su}(N)$ orthonormal for $|X|^2 = -\operatorname{Tr}X^2$, so that $C_2 = (N^2-1)/(2N)$ on the fundamental representation.

Proof. Since $w_\beta(\bar{U}) = w_\beta(U)$ and $\chi_\lambda(\bar{U}) = \overline{\chi_\lambda(U)}$, and complex conjugation preserves Haar measure, $\int w_\beta\chi_\lambda = \int w_\beta\bar{\chi}_\lambda$, which is the complex conjugate of the left side; so $\hat{c}_\lambda$ is real. Next, w_β is a multiple of $\exp(\frac{\beta}{2N}(\chi_{\mathrm{F}} + \bar{\chi}_{\mathrm{F}})) = \sum_m \frac{1}{m!}(\frac{\beta}{2N})^m(\chi_{\mathrm{F}} + \bar{\chi}_{\mathrm{F}})^m$, and each power is the character of a representation, hence a sum of irreducible characters with nonnegative integer coefficients. By orthonormality $\int w_\beta\bar{\chi}_\lambda \ge 0$, so $\hat{c}_\lambda \ge 0$. As $|\chi_\lambda(U)| \le d_\lambda$, with $\operatorname{Re}\chi_\lambda(U) = d_\lambda$ only where $\rho_\lambda(U) = I$, a proper closed subgroup when $\lambda \neq 0$ and so a Haar null set, and $w_\beta > 0$ everywhere, $\hat{c}_\lambda = d_\lambda^{-1}\int w_\beta \operatorname{Re}\chi_\lambda < 1$ for $\lambda \neq 0$.

For the expansion, $\operatorname{Re}\operatorname{Tr}U \le N$ with equality only at $U = I$, so outside a neighbourhood $\{U = e^X : |X| < \delta\}$ both integrals $\int e^{\frac{\beta}{N}\operatorname{Re}\operatorname{Tr}U}\chi_\lambda$ and $\int e^{\frac{\beta}{N}\operatorname{Re}\operatorname{Tr}U}$ are smaller than their values near I by a factor $e^{-c\beta}$. Near I write $U = e^X$; the Haar density in these coordinates is $j(X) = 1 + O(|X|^2)$, even in X. For antihermitian X, $\operatorname{Tr}X^3$ is imaginary, so

$$\tfrac{1}{N}\operatorname{Re}\operatorname{Tr}e^X = 1 - \tfrac{1}{2N}|X|^2 + r(X), \qquad 0 \le r(X) \le C|X|^4,$$

and $d_\lambda^{-1}\chi_\lambda(e^X) = 1 + d_\lambda^{-1}\operatorname{Tr}\rho_\lambda(X) + \frac{1}{2d_\lambda}\operatorname{Tr}\rho_\lambda(X)^2 + c_3(X) + O(|X|^4)$, with c_3 cubic and $\operatorname{Tr}\rho_\lambda(X) = 0$, since $\mathfrak{su}(N)$ is perfect. Let $\langle\cdot\rangle$ denote the average against the Gaussian weight $e^{-\beta|X|^2/(2N)}$, under which the coordinates x_a of $X = \sum x_aT_a$ are independent with variance N/β. With $h = j\,e^{\beta r}$,

$$\begin{aligned}\hat{c}_\lambda(\beta) &= \frac{\langle d_\lambda^{-1}\chi_\lambda(e^X)\,h\rangle}{\langle h\rangle} + O(e^{-c\beta}) \\ &= 1 + \frac{\langle \frac{1}{2d_\lambda}\operatorname{Tr}\rho_\lambda(X)^2\,h\rangle}{\langle h\rangle} + \frac{\langle (c_3(X) + O(|X|^4))h\rangle}{\langle h\rangle},\end{aligned}$$

the averages being taken over $|X| < \delta$. Under $\langle\cdot\rangle$, $|X|^{2m}$ has mean of order β^{-m} and $\beta|r(X)| \le C\beta|X|^4$ has mean of order β^{-1}, so $h = 1 + O(\beta^{-1})$ in every moment. The weight h is even in X, since $\operatorname{Tr}X^k$ is imaginary for odd k and j is even, so the odd term c_3 contributes nothing; the quartic remainder and the correlation of the quadratic

term with $h-1$ contribute $O(\beta^{-2})$. Finally $\langle \operatorname{Tr}\rho_\lambda(X)^2\rangle = \frac{N}{\beta}\sum_a \operatorname{Tr}\rho_\lambda(T_a)^2 = -\frac{2NC_2(\lambda)d_\lambda}{\beta}$, which gives the expansion. For SU(2) and the representation of spin j the integrals are explicit, $\hat{c}_j(\beta) = I_{2j+1}(\beta)/I_1(\beta)$ with the modified Bessel functions I_n, in agreement with $C_2 = j(j+1)$. □

Appendix I. The single-link law of SU(2)

This appendix proves the sharp quarter, $W_1(\nu_b, \nu_{b'}) \le \frac{1}{4}|b-b'|$ for the one-link law of SU(2), which Theorems 1.1 and 1.3 use, and the exact value at $b=0$, which Theorem 1.2 uses. The argument is that of our earlier paper [15]; it is given in full so that the present paper is self-contained.

I.1. **The von Mises–Fisher law.** Identify SU(2) with the unit quaternions, that is with the unit sphere $S^3 \subset \mathbb{R}^4$, so that $\operatorname{Re}\operatorname{Tr} U = 2x_0(U)$ and $\operatorname{Re}\operatorname{Tr}(UV) = 2\langle u, \bar{v}\rangle$ for the Euclidean inner product of $\mathbb{R}^4$ and quaternion conjugation $v \mapsto \bar{v}$. Fix a link e and the field off e. Every plaquette through e can be rotated cyclically so that its trace reads $\operatorname{Re}\operatorname{Tr}(Q_e A_p)$ with $A_p \in \mathrm{SU}(2)$, the staple of e in p. The action therefore depends on Q_e only through

$$N\beta \sum_{p\ni e} \operatorname{Re}\operatorname{Tr}(Q_e A_p) = \langle x, b\rangle, \qquad b = 4\beta \sum_{p\ni e} \bar{A}_p \in \mathbb{R}^4,$$

with $x = Q_e \in S^3$. Since there are six staples, each of norm one,

$$\kappa := |b| \le y := 24|\beta|. \tag{I.1}$$

The conditional law of Q_e is the von Mises–Fisher law[25]

$$\nu_b(dx) = Z(\kappa)^{-1} e^{\langle x,b\rangle}\, \sigma(dx) \tag{I.2}$$

on the unit sphere, with σ the normalised surface measure. We use coordinates in which $b = \kappa e_1$. The density of $t = x_1$ is proportional to $(1-t^2)^{1/2}e^{\kappa t}$ on $[-1,1]$, so $Z(\kappa) = 2I_1(\kappa)/\kappa$ with the modified Bessel function I_1, normalised so that $Z(0) = 1$.

Lemma I.1. *For every $\kappa \ge 0$ the law ν_b on the unit sphere S^3 satisfies the Poincaré inequality $\operatorname{Var}_{\nu_b}(g) \le \frac{1}{3}\int |\nabla g|^2 d\nu_b$, and the constant $\frac{1}{3}$ is attained at $\kappa = 0$.*

Proof. In the polar angle $\theta \in (0,\pi)$ from e_1 the generator of the reversible diffusion of ν_b is $L = \partial_\theta^2 + b(\theta)\partial_\theta + \sin^{-2}\theta\, \Delta_{S^2}$ with $b(\theta) = 2\cot\theta - \kappa\sin\theta$. On spherical harmonics of degree ℓ on S^2 it acts as a Sturm–Liouville operator L_ℓ with potential $\ell(\ell+1)\sin^{-2}\theta$. A ground state comparison of Barta's type [10] with the trial function $\psi = \sin\theta\, e^{a\kappa\cos\theta}$ gives $-L_1\psi \ge 3\psi$ for $a = -\frac{1}{5}$, and the same trial function with $a = -\frac{2}{5}$ gives the bound 3 for the derivatives of zonal eigenfunctions; Appendix J carries out both computations. Hence every nonzero eigenvalue of $-L$ is at least 3. At $\kappa = 0$ the first eigenvalue of the Laplacian of the unit S^3 is 3. □

I.2. **Moments.** Let $A(\kappa) = \mathbb{E}_{\nu_b} x_1 = Z'(\kappa)/Z(\kappa) = I_2(\kappa)/I_1(\kappa)$.

Lemma I.2. *For every $\kappa > 0$, in coordinates with $b = \kappa e_1$,*

(i) $\mathbb{E}x_2^2 = \mathbb{E}x_3^2 = \mathbb{E}x_4^2 = A/\kappa$, $\mathbb{E}(1-x_1^2) = 3A/\kappa$, *the covariance matrix of x is* $\operatorname{diag}(A', A/\kappa, A/\kappa, A/\kappa)$, *and* $A' = 1 - A^2 - 3A/\kappa$;

(ii) $\dfrac{\kappa}{2+\sqrt{\kappa^2+4}} \le A(\kappa) \le \dfrac{\kappa}{1+\sqrt{\kappa^2+9}}$;

(iii) $A' \le A/\kappa \le \left(1+\sqrt{\kappa^2+9}\right)^{-1} \le \frac{1}{4}$;

[25] On the sphere $S^{p-1} \subset \mathbb{R}^p$ the von Mises–Fisher law with mean direction μ and concentration κ has density proportional to $e^{\kappa\langle x,\mu\rangle}$ with respect to the surface measure, and its normalising constant is $(2\pi)^{p/2} I_{p/2-1}(\kappa)/\kappa^{p/2-1}$; it is the maximum entropy law with prescribed mean resultant, and it plays on the sphere the role of the Gaussian in $\mathbb{R}^p$. For $p = 4$ the Bessel index is 1. The mean resultant length $\mathbb{E}\langle x,\mu\rangle = I_{p/2}(\kappa)/I_{p/2-1}(\kappa)$ is the function A of Lemma I.2. See Mardia and Jupp [61], Ch. 9, for these facts and for the tangent-normal decomposition used in Appendix J.

(iv) $A(\kappa) \geq \kappa/\big(\frac{3}{2} + \sqrt{\kappa^2 + 25/4}\big)$.

Proof. (i) Integrating by parts against $\frac{d}{dt}(1-t^2)^{3/2} = -3t(1-t^2)^{1/2}$ gives $\kappa\,\mathbb{E}(1-x_1^2) = 3\mathbb{E}x_1$, and the three coordinates x_2, x_3, x_4 are exchangeable under ν_b, with $\sum_i x_i^2 = 1$. The off-diagonal covariances vanish by the reflections $x_i \mapsto -x_i$, $i \geq 2$, and $A' = \operatorname{Var} x_1$ because $A = (\log Z)'$. (ii) By (i), A solves the Riccati equation $A' = 1 - A^2 - 3A/\kappa$ with $A(\kappa) = \kappa/4 + O(\kappa^3)$. Appendix J shows that the left side of (ii) is a subsolution and the right side a supersolution of this equation, with the same behaviour at 0, and concludes by comparison. (iii) The middle inequality is (ii). For the first, put $s = \sqrt{\kappa^2 + 4}$. By (ii), $4A/\kappa \geq 4/(2+s)$ and $1 - A^2 \leq 1 - \kappa^2/(2+s)^2$, and

$$1 - \frac{\kappa^2}{(2+s)^2} \leq \frac{4}{2+s} \iff (2+s)^2 - \kappa^2 \leq 4(2+s)$$
$$\iff s^2 - \kappa^2 \leq 4,$$

which holds with equality. So $A' = 1 - A^2 - 3A/\kappa \leq A/\kappa$. (iv) is the comparison of (ii) with a third barrier, which improves the lower bound of (ii); it is carried out in the same appendix. □

I.3. **The Lipschitz rate of the family.** Let δ be the chordal metric $\delta(x, x') = |x - x'|$ on S^3 and W_1 the Kantorovich distance for δ. For $c \in \mathbb{R}^4$ and a unit vector $h \in \mathbb{R}^4$ put

$$L(c, h) = \sup_g \operatorname{Cov}_{\nu_c}\big(g, \langle h, x\rangle\big), \tag{I.3}$$

the supremum over 1-Lipschitz functions g for δ. Since $\frac{d}{dt}\nu_{c+th}(g) = \operatorname{Cov}_{\nu_{c+th}}(g, \langle h, x\rangle)$, the Kantorovich–Rubinstein duality gives, for all b, b',

$$W_1(\nu_b, \nu_{b'}) \leq |b - b'| \sup_{c \in [b, b']} L\Big(c, \frac{b' - b}{|b' - b|}\Big). \tag{I.4}$$

By rotation invariance $L(c, h)$ depends only on $\kappa = |c|$ and on the angle φ between h and $c^\perp$, and we write $L(\kappa, \varphi)$.

Lemma I.3. *For all c and h, $L(c, h) \leq 1/(2\sqrt{3})$. Moreover:*

(i) *at $c = 0$, $L(0, h) = \frac{1}{4}$ for every h;*

(ii) *if $h \perp c$, then $L(c, h) = \mathbb{E}_{\nu_c} x_2^2 = A(\kappa)/\kappa \leq \frac{1}{4}$;*

(iii) *if $\kappa \geq \sqrt{88}/3 \approx 3.127$, then $L(c, h) \leq \frac{1}{4}$ for every h.*

Proof. A 1-Lipschitz function for δ is 1-Lipschitz for the geodesic distance, which is larger, so $|\nabla g| \leq 1$ almost everywhere, and Lemma I.1 gives $\operatorname{Var} g \leq \frac{1}{3}$. By Lemma I.2(i), (iii) the largest eigenvalue of the covariance of x is $A/\kappa \leq \frac{1}{4}$, so $\operatorname{Var}\langle h, x\rangle \leq A/\kappa$. The Cauchy–Schwarz inequality gives

$$L(c, h) \leq \big(\tfrac{1}{3}A/\kappa\big)^{1/2} \leq \Big(3\big(1 + \sqrt{\kappa^2 + 9}\big)\Big)^{-1/2} \leq \frac{1}{2\sqrt{3}}. \tag{I.5}$$

(i) and (ii) are reflection arguments. Let R be the reflection in the hyperplane $h^\perp$, and suppose ν_c is R-invariant, which is the case when $c = 0$ or $h \perp c$. Then $\mathbb{E}\langle h, x\rangle = 0$ and

$$\operatorname{Cov}(g, \langle h, x\rangle) = \mathbb{E}\big[(g(x) - g(Rx))\langle h, x\rangle;\ \langle h, x\rangle > 0\big]$$
$$\leq \mathbb{E}\big[2\langle h, x\rangle^2;\ \langle h, x\rangle > 0\big] = \mathbb{E}\langle h, x\rangle^2,$$

since $\delta(x, Rx) = 2|\langle h, x\rangle|$. Equality holds for $g = \langle h, x\rangle$, which is 1-Lipschitz. At $c = 0$, $\mathbb{E}\langle h, x\rangle^2 = \frac{1}{4}$; for $h \perp c$, $\mathbb{E}\langle h, x\rangle^2 = A/\kappa$. (iii) By (I.5), $L \leq \frac{1}{4}$ as soon as $1 + \sqrt{\kappa^2 + 9} \geq \frac{16}{3}$, that is $\kappa^2 \geq \frac{88}{9}$. □

The reflection settles the directions $h \perp c$ and the point $c = 0$; for an oblique h the reflection no longer preserves ν_c, and the Cauchy–Schwarz bound (I.5) is too weak for $\kappa < \sqrt{88}/3$. The next theorem closes this case.

Theorem I.4 (The sharp quarter)**.** *For every $c \in \mathbb{R}^4$ and every unit vector $h \in \mathbb{R}^4$,*

$$L(c,h) \le \tfrac{1}{4},$$

with equality at $c = 0$. Equivalently, $W_1(\nu_b, \nu_{b'}) \le \frac{1}{4}|b-b'|$ for all $b, b' \in \mathbb{R}^4$, and the constant $\frac{1}{4}$ is the best possible.

The equivalence is (I.4) in one direction and the derivative formula above it in the other; the sharpness is Lemma I.3(i). The metric matters.[26] The proof has four steps. A Lipschitz function on S^3 extends to the unit ball $B \subset \mathbb{R}^4$, and Green's formula in B bounds the covariance by a negative Sobolev norm of the density (Lemma I.5). This norm splits into a zonal and a transverse part, both explicit series in modified Bessel functions (Lemma I.6). The transverse part is at most $\frac{1}{4}$ because a certain power series has nonnegative coefficients (Proposition I.8), and the zonal part is at most $\frac{1}{4}$ by a Riccati comparison (Proposition I.9).

I.4. **Harmonic extension.** Let σ be the normalised surface measure of S^3 and H_k the space of spherical harmonics of degree k, the restrictions to S^3 of harmonic homogeneous polynomials of degree k on $\mathbb{R}^4$. The spaces H_k are orthogonal in $L^2(\sigma)$ and span it; we write u_k for the component of u in H_k, and for u with $\int u\, d\sigma = 0$ we put

$$\mathcal{Q}(u) = \sum_{k \ge 1} \frac{1}{k}\, \|u_k\|^2_{L^2(\sigma)}. \tag{I.6}$$

This is $\langle u, \Lambda^{-1} u\rangle$ for the Dirichlet-to-Neumann operator Λ of B, which acts on H_k as multiplication by k.

Lemma I.5. *Let u be a real-analytic function on S^3 with $\int u\, d\sigma = 0$ and let g be 1-Lipschitz for the chordal metric. Then*

$$\int_{S^3} g\, u\, d\sigma \le \tfrac{1}{2}\, \mathcal{Q}(u)^{1/2}.$$

Proof. The chordal metric is the Euclidean metric of $\mathbb{R}^4$ restricted to S^3, so McShane's extension $G(y) = \inf_{x \in S^3}(g(x) + |x - y|)$ is 1-Lipschitz on $\mathbb{R}^4$ and equals g on S^3 [62]; by Rademacher's theorem $|\nabla G| \le 1$ almost everywhere. Let $\psi = \sum_{k \ge 1} k^{-1} r^k u_k(\omega)$ in polar coordinates $y = r\omega$. Each term is a harmonic polynomial, the series converges with all derivatives on the closed ball because u is real-analytic,[27] and $\partial_r \psi = u$ on S^3. Green's formula for the Lipschitz function G and the smooth harmonic function ψ reads $\int_{S^3} G\, \partial_r \psi\, dS = \int_B \nabla G \cdot \nabla \psi\, dy$. With $|S^3| = 2\pi^2$ and $|B| = \pi^2/2$ the Cauchy–Schwarz inequality gives

$$\int g\, u\, d\sigma = \frac{1}{2\pi^2} \int_B \nabla G \cdot \nabla \psi\, dy \le \frac{|B|^{1/2}}{2\pi^2} \Big(\int_B |\nabla \psi|^2 dy \Big)^{1/2}.$$

Green's formula once more gives $\int_B |\nabla \psi|^2 = \int_{S^3} \psi\, \partial_r \psi\, dS = 2\pi^2 \sum_k k^{-1} \|u_k\|^2 = 2\pi^2 \mathcal{Q}(u)$, and the right side becomes $\frac{1}{2}\mathcal{Q}(u)^{1/2}$. □

The lemma is a form of Thomson's principle. By Kantorovich–Rubinstein duality, $\sup_g \int g u\, d\sigma$ is the least cost of a flow in $\mathbb{R}^4$ that carries the positive part of $u\, d\sigma$ to its

[26] In the geodesic metric of S^3 the statement is false already at $c = 0$: the function $\arcsin x_2$ is geodesically 1-Lipschitz and its covariance with x_2 under the uniform law is $\frac{8}{9\pi} \approx 0.283 > \frac{1}{4}$. A proof therefore has to use that the chordal distance between two points is shorter than their geodesic distance, that is, it has to see the sphere from the ball it bounds. The proof below does exactly this: it extends a chordally Lipschitz function to the ball and integrates by parts there.

[27] For real-analytic u the norms $\|u_k\|$ decay geometrically in k, and the supremum of a function in H_k is at most $(\dim H_k)^{1/2} = k + 1$ times its $L^2(\sigma)$ norm, so the series for ψ and for each of its derivatives converge uniformly on $\bar{B}$.

negative part, the cost being the total length travelled by the mass.[28] The harmonic field $\nabla\psi$ is one such flow, through the interior of the ball, and the lemma computes its cost. At $c = 0$ the flow is optimal: for $u = \langle h, x\rangle$ one has $\psi = \langle h, y\rangle$, the field $\nabla\psi = h$ is constant, the Cauchy–Schwarz inequality is an equality, and $\mathcal{Q}(u) = \|u\|^2 = \frac{1}{4}$.

I.5. **Spectral decomposition.** Fix $c = \kappa e_1$ with $\kappa > 0$ and write $h = \cos\varphi\, e_1 + \sin\varphi\, e_2$, which is no loss of generality by rotation invariance. Let $\rho = d\nu_c/d\sigma = e^{\kappa x_1}/Z(\kappa)$, $Z(\kappa) = \int e^{\kappa x_1} d\sigma = 2I_1(\kappa)/\kappa$, and

$$u = \big(\langle h, x\rangle - \mathbb{E}_{\nu_c}\langle h, x\rangle\big)\rho,$$

so that $\mathrm{Cov}_{\nu_c}(g, \langle h, x\rangle) = \int g u\, d\sigma$ and $\int u\, d\sigma = 0$. Let U_n be the Chebyshev polynomials of the second kind and put

$$d_n(\kappa) = \frac{(n+1)\, I_{n+1}(\kappa)}{I_1(\kappa)}, \qquad n \geq 0. \tag{I.7}$$

Lemma I.6. *With this notation*

$$\mathcal{Q}(u) = \cos^2\varphi\, \mathcal{Q}_z(\kappa) + \sin^2\varphi\, \mathcal{Q}_x(\kappa),$$

where

$$\mathcal{Q}_z = \sum_{n\geq 1} \frac{d_n'(\kappa)^2}{n}, \qquad \mathcal{Q}_x = \frac{1}{3\kappa^2} \sum_{n\geq 1} (n+2)\, d_n(\kappa)^2.$$

Proof. The functions $U_n(x_1)$, $n \geq 0$, are the zonal harmonics: $U_n(x_1) \in H_n$, and they are orthonormal in $L^2(\sigma)$ (Appendix K). The expansion $e^{\kappa t} = \frac{2}{\kappa}\sum_n (n+1) I_{n+1}(\kappa) U_n(t)$ of the same appendix gives $\rho = \sum_n d_n(\kappa) U_n(x_1)$, with $d_0 = 1$.

Write $u = \cos\varphi\, u_z + \sin\varphi\, u_x$ with $u_z = (x_1 - A)\rho$ and $u_x = x_2\rho$; here $\mathbb{E}_{\nu_c} x_1 = A$ and $\mathbb{E}_{\nu_c} x_2 = 0$. The reflection $x_2 \mapsto -x_2$ maps each H_k to itself, fixes u_z and reverses u_x, so the components of u_z and u_x in each H_k are orthogonal and the cross term in $\mathcal{Q}(u)$ vanishes.

Since $A = Z'/Z$, we have $u_z = \partial_\kappa \rho = \sum_n d_n'(\kappa) U_n(x_1)$, and the component of u_z in H_n is $d_n' U_n(x_1)$, of norm $|d_n'|$. This gives $\mathcal{Q}_z$.

For u_x, let $y(t) = \cos t\, e_1 + \sin t\, e_2$. The law $\nu_{\kappa y(t)}$ has density $\rho_t = e^{\kappa\langle y(t), x\rangle}/Z(\kappa) = \sum_n d_n(\kappa) U_n(\langle y(t), x\rangle)$, and $\partial_t \rho_t|_{t=0} = \kappa x_2 \rho$. The function $U_n(\langle y(t), x\rangle)$ lies in H_n for every t, hence so does its derivative $x_2 U_n'(x_1)$ at $t = 0$, and the component of u_x in H_n is $\kappa^{-1} d_n x_2 U_n'(x_1)$. Its norm follows from the eigenvalue $n(n+2)$ of the Laplacian of S^3 on H_n. For $f = U_n(x_1)$,

$$\begin{aligned} n(n+2) = n(n+2)\|f\|^2 &= \int |\nabla_{S^3} f|^2 d\sigma \\ &= \int U_n'(x_1)^2 (1 - x_1^2)\, d\sigma, \end{aligned}$$

since the tangential gradient of x_1 has squared length $1 - x_1^2$. Given x_1, the point (x_2, x_3, x_4) is uniform on the sphere of radius $(1 - x_1^2)^{1/2}$, so $\mathbb{E}[x_2^2 \mid x_1] = \frac{1}{3}(1 - x_1^2)$ and $\|x_2 U_n'(x_1)\|^2 = \frac{1}{3} n(n+2)$. Summing $n^{-1}\kappa^{-2} d_n^2 \cdot \frac{1}{3} n(n+2)$ over $n \geq 1$ gives $\mathcal{Q}_x$. □

By Lemmas I.5 and I.6,

$$L(\kappa, \varphi) \leq \tfrac{1}{2} \max\big(\mathcal{Q}_z(\kappa), \mathcal{Q}_x(\kappa)\big)^{1/2}, \tag{I.8}$$

and Theorem I.4 follows once both parts are at most $\frac{1}{4}$. Both are $\frac{1}{4}$ at $\kappa = 0$, by continuity from the values $d_1 \sim \kappa/2$, $d_1' \to \frac{1}{2}$ and $d_n = O(\kappa^n)$.

[28] This is the Beckmann formulation of the transport problem with cost $|x - y|$: the optimal cost equals the least $\int |V|$ over vector measures V on $\mathbb{R}^4$ with divergence $u\, d\sigma$. Restricting the flow to the closed ball does not change the infimum, since projecting onto $\bar{B}$ shortens paths. The gradient $\nabla\psi$ on B is one admissible flow. It minimises the energy $\int |V|^2$ rather than the cost $\int |V|$, and the Cauchy–Schwarz inequality converts one into the other at the price of the factor $|B|^{1/2}$.

I.6. **Three Bessel sums.** The series of Lemma I.6 reduce to closed forms. Write I_j for $I_j(\kappa)$.

Lemma I.7. *For every $\kappa > 0$,*

$$\sum_{j\ge1} j^2 I_j^2 = \frac{\kappa}{4}\, I_1(2\kappa), \qquad \sum_{j\ge1} j^3 I_j^2 = \frac{\kappa^2}{4}(I_0^2 + I_1^2),$$
$$I_1(2\kappa) = 2\sum_{j\ge1} I_{j-1} I_j.$$

The first and third identities follow from Parseval's formula for $e^{\kappa\cos\theta} = \sum_{j\in\mathbb{Z}} I_j e^{ij\theta}$, and the second from the recurrences of I_j by telescoping; Appendix K gives the proofs.

I.7. **The transverse part.**

Proposition I.8. *$\mathcal{Q}_x(\kappa) \le \frac14$ for every $\kappa \ge 0$.*

Proof. By (I.7), $\mathcal{Q}_x = (3\kappa^2 I_1^2)^{-1}\sum_{j\ge2} j^2(j+1)I_j^2$, and the sum equals $\sum_{j\ge1}(j^3+j^2)I_j^2 - 2I_1^2$. By Lemma I.7, $\mathcal{Q}_x \le \frac14$ is equivalent to

$$\mathcal{D}(\kappa) := (8+2\kappa^2)I_1^2 - \kappa^2 I_0^2 - \kappa I_1(2\kappa) \ge 0. \tag{I.9}$$

Each term is an entire function of κ that is even after the factor κ^2 is removed, so $\mathcal{D}(\kappa) = \sum_{n\ge0} \mathcal{D}_n \kappa^{2n}$. Lemma K.2 of Appendix K computes the coefficients: $\mathcal{D}_0 = \mathcal{D}_1 = 0$, and for $n = m+1 \ge 2$

$$\mathcal{D}_n = \frac{N_m}{4^m\, m!\,(m+1)!\,(m+1)(m+2)},$$

with

$$N_m = \binom{2m}{m}\big(4(2m+1) + (m-1)(m+1)(m+2)\big) - 4^m(m+1)(m+2).$$

The numerator N_m vanishes for $m = 0, 1, 2$ and is positive for $m \ge 3$ (Lemma K.2). So every coefficient of $\mathcal{D}$ is nonnegative and $\mathcal{D} \ge 0$. □

The first nonzero coefficient is $\mathcal{D}_4 = \frac{1}{2304}$, so $\frac14 - \mathcal{Q}_x = \mathcal{D}/(12\kappa^2 I_1^2)$ is of order κ^4 at 0: the transverse part leaves the value $\frac14$ very slowly, which is why no estimate that loses a constant can prove the proposition.

I.8. **The zonal part.** Let $A = I_2/I_1$ as before, so that $d_1 = 2A$ and, by Lemma I.2(i), $A' = 1 - A^2 - 3A/\kappa$.

Proposition I.9. *For $0 < \kappa \le \sqrt{88}/3$,*

$$\mathcal{Q}_z(\kappa) \le \frac14 + 4A'^2 + \frac{A^2}{4} - \frac{A}{2\kappa} - \frac{2A^2}{\kappa^2} < \frac14.$$

Proof. The higher modes. With $r_\nu = I_{\nu+1}/I_\nu$ the recurrence $I_\nu' = I_{\nu+1} + \frac{\nu}{\kappa} I_\nu$ gives

$$\frac{d_n'}{d_n} = \frac{I_{n+1}'}{I_{n+1}} - \frac{I_1'}{I_1} = \frac{n}{\kappa} + r_{n+1} - r_1.$$

By Lemma K.3 of Appendix K, $r_{n+1} \le r_1$, and $r_1 = A \le \kappa/4$ by Lemma I.2(iii). Hence $-n/\kappa \le n/\kappa - \kappa/4 \le d_n'/d_n \le n/\kappa$ for $n \ge 2$ and $\kappa^2 \le 16$, so $d_n'^2 \le n^2 d_n^2/\kappa^2$. With $d_1' = 2A'$,

$$\mathcal{Q}_z \le 4A'^2 + \frac{1}{\kappa^2}\sum_{n\ge2} n\, d_n^2 = 4A'^2 + \frac{S_3 - S_2 - 4I_2^2}{\kappa^2 I_1^2},$$

where $S_2 = \sum_{j\ge1} j^2 I_j^2$ and $S_3 = \sum_{j\ge1} j^3 I_j^2$; the second form uses $n\, d_n^2 I_1^2 = (j-1)j^2 I_j^2$ with $j = n+1$.

The closed form. The recurrences $I_0 - I_2 = \frac{2}{\kappa} I_1$ and $I_1 - I_3 = \frac{4}{\kappa} I_2$ give $I_0/I_1 = 2/\kappa + A$ and $I_2 I_3 / I_1^2 = A - 4A^2/\kappa$. By Lemma I.7, $S_3/(\kappa^2 I_1^2) = \frac{1}{4}(2/\kappa + A)^2 + \frac{1}{4}$, and, keeping three terms of the third identity,

$$\frac{S_2}{\kappa^2 I_1^2} \geq \frac{I_0 I_1 + I_1 I_2 + I_2 I_3}{2\kappa I_1^2} = \frac{1}{2\kappa}\Big(\frac{2}{\kappa} + 3A - \frac{4A^2}{\kappa}\Big).$$

Collecting terms, $\mathcal{Q}_z \leq \frac{1}{4} + 4A'^2 + T(A)$ with $T(a) = \frac{a^2}{4} - \frac{a}{2\kappa} - \frac{2a^2}{\kappa^2}$, which is the first inequality.

The barrier. Let $s = \sqrt{\kappa^2 + 25/4}$ and $A_\flat = \kappa/(\frac{3}{2} + s)$. By Lemma I.2(iv), $A_\flat \leq A \leq \kappa/4$. The function $a \mapsto 1 - a^2 - 3a/\kappa$ decreases on $a \geq 0$, and $A' \geq 0$ because $A' = \operatorname{Var} x_1$, so $0 \leq A' \leq \alpha := 1 - A_\flat^2 - 3A_\flat/\kappa$. For $\kappa^2 \leq 12$ the quadratic T decreases on $[0, \kappa/4]$, since

$$T(a) - T(a') = (a' - a)\,\frac{2\kappa - (a + a')(\kappa^2 - 8)}{4\kappa^2}$$

and $(a + a')(\kappa^2 - 8) \leq \frac{\kappa}{2}(\kappa^2 - 8) \leq 2\kappa$ there. So $\mathcal{Q}_z \leq \frac{1}{4} + 4\alpha^2 + T(A_\flat)$. With $\kappa^2 = s^2 - \frac{25}{4}$ and $A_\flat/\kappa = 2/(2s+3)$ both terms are rational in s: $\alpha = 16/(2s+3)^2$, and

$$4\alpha^2 + T(A_\flat) = \frac{(2s-5)\,p(s)}{4(2s+3)^4}, \qquad p(s) = 8s^3 + 28s^2 - 98s - 695. \tag{I.10}$$

For $0 < \kappa \leq \sqrt{88}/3$ we have $\frac{5}{2} < s \leq \sqrt{577}/6 < 4.004$. On $s \geq \frac{5}{2}$ the cubic p increases, since $p'(s) = 24s^2 + 56s - 98 > 0$, and $p(4.004) < -120$. So the right side of (I.10) is negative. □

I.9. **Proof of Theorem I.4.** At $c = 0$ the theorem is Lemma I.3(i), and for $|c| \geq \sqrt{88}/3$ it is Lemma I.3(iii). For $0 < |c| < \sqrt{88}/3$, Propositions I.8 and I.9 give $\mathcal{Q}_x, \mathcal{Q}_z \leq \frac{1}{4}$, and (I.8) gives $L \leq \frac{1}{4}$. □

Appendix J. Computations for the single-link law

J.1. **Barta's inequality.** Let $\rho(\theta) = \sin^2\theta\, e^{\kappa\cos\theta}$ on $(0, \pi)$, let V be a continuous potential, and let

$$T = -\rho^{-1}\partial_\theta\, \rho\, \partial_\theta + V.$$

If $\psi > 0$ is smooth on $(0, \pi)$ with $\rho\psi^2 = O(\sin^4\theta)$ at the end points and $T\psi \geq \lambda_0 \psi$, then

$$\int (|h'|^2 + Vh^2)\rho\, d\theta \geq \lambda_0 \int h^2 \rho\, d\theta$$

for every smooth h vanishing to first order at 0 and π. Indeed, writing $h = \psi u$ and integrating by parts, the boundary terms $\rho\psi\psi' u^2$ vanish and

$$\int (|h'|^2 + Vh^2)\rho = \int \rho\psi^2 |u'|^2 + \int \frac{T\psi}{\psi}\, h^2 \rho \;\geq\; \lambda_0 \int h^2 \rho.$$

This is the inequality of Barta [10] in the form of a ground state transformation.

J.2. **The computations of Lemma I.1.** The generator of the reversible diffusion of ν_b on S^3 is $L = \Delta_{S^3} + \kappa \nabla x_1 \cdot \nabla$. In the polar angle θ from e_1 the round metric is $d\theta^2 + \sin^2\theta\, g_{S^2}$, so $\Delta_{S^3} = \partial_\theta^2 + 2\cot\theta\, \partial_\theta + \sin^{-2}\theta\, \Delta_{S^2}$, and $\nabla x_1 \cdot \nabla = -\sin\theta\, \partial_\theta$. On $h(\theta) Y_\ell$ with a spherical harmonic Y_ℓ of degree ℓ on S^2 the operator acts as $L_\ell = \partial_\theta^2 + b\,\partial_\theta - \ell(\ell+1)\sin^{-2}\theta$, $b = 2\cot\theta - \kappa\sin\theta = \rho'/\rho$, so $-L_\ell = T$ with $V = \ell(\ell+1)\sin^{-2}\theta$.

Degree $\ell \geq 1$. Take $\psi = \sin\theta\, e^{a\kappa\cos\theta}$. Then $\psi'/\psi = \cot\theta - a\kappa\sin\theta$ and $\psi''/\psi = (\psi'/\psi)^2 + (\psi'/\psi)'$, and a short computation gives

$$\frac{-L_1\psi}{\psi} = 3 + (1 + 5a)\kappa\cos\theta - a(1+a)\kappa^2\sin^2\theta.$$

For $a = -\frac{1}{5}$ this is $3 + \frac{4}{25}\kappa^2\sin^2\theta \geq 3$. Since $V = \ell(\ell+1)\sin^{-2}\theta \geq 2\sin^{-2}\theta$ for $\ell \geq 1$, Barta's inequality gives $-L_\ell \geq 3$ on every sector with $\ell \geq 1$.

Degree $\ell = 0$. Let $g(\theta)$ be a zonal eigenfunction, $-L_0 g = \lambda g$, g not constant, so that $g'(0) = g'(\pi) = 0$. Differentiating $g'' + bg' = -\lambda g$ gives, for $h = g'$,

$$-(h'' + bh') - b'h = \lambda h, \qquad -b' = \frac{2}{\sin^2\theta} + \kappa\cos\theta,$$

so h is an eigenfunction of T with $V = 2\sin^{-2}\theta + \kappa\cos\theta$, with eigenvalue λ, and $h \not\equiv 0$ vanishes at both ends. With the same ψ,

$$\frac{T\psi}{\psi} = 3 + (2+5a)\kappa\cos\theta - a(1+a)\kappa^2\sin^2\theta,$$

which for $a = -\frac{2}{5}$ is $3 + \frac{6}{25}\kappa^2\sin^2\theta \geq 3$. Barta's inequality applied to h gives $\lambda \geq 3$. Together with the case $\ell \geq 1$ every nonzero eigenvalue of $-L$ is at least 3, which is Lemma I.1; the spectrum is discrete because S^3 is compact and the drift is smooth.

J.3. **Moments.** The density of $t = x_1$ under ν_b is proportional to $(1-t^2)^{1/2}e^{\kappa t}$. For every smooth ϕ on $[-1,1]$,

$$\frac{d}{dt}\Big[(1-t^2)^{3/2}e^{\kappa t}\Big] = (1-t^2)^{1/2}e^{\kappa t}(\kappa(1-t^2) - 3t),$$

and integration by parts gives $\mathbb{E}[\phi'(t)(1-t^2)] = \mathbb{E}\big[\phi(t)\big(3t - \kappa(1-t^2)\big)\big]$. With $\phi = 1$ this is $\kappa\mathbb{E}(1-x_1^2) = 3\mathbb{E}x_1$, the identity of Lemma I.2(i). With $\phi = t^k$ it gives the recursion

$$m_{k+2} = m_k - \frac{k+3}{\kappa}\,m_{k+1} + \frac{k}{\kappa}\,m_{k-1}, \qquad m_k = \mathbb{E}x_1^k,$$

from $m_0 = 1$ and $m_1 = A$; all polynomial moments of ν_b are rational in κ and $A(\kappa)$.[29]

J.4. **The Riccati barriers.** By Lemma I.2(i), A solves $A' = \mathcal{F}(\kappa, A) := 1 - A^2 - 3A/\kappa$ on $(0,\infty)$, and $A(\kappa) = \frac{\kappa}{4} - \frac{\kappa^3}{96} + O(\kappa^5)$ from the series of I_1 and I_2. Let $s = \sqrt{\kappa^2+4}$, $u = \sqrt{\kappa^2+9}$, $\phi_- = \kappa/(2+s)$ and $\phi_+ = \kappa/(1+u)$. Direct differentiation gives

$$\phi_-' - \mathcal{F}(\kappa,\phi_-) = -\frac{\kappa^2}{s(s+2)^2} \leq 0,$$

$$\phi_+' - \mathcal{F}(\kappa,\phi_+) = \frac{(u-3)^2}{u(1+u)^2} \geq 0,$$

and $\phi_\pm = \frac{\kappa}{4} + O(\kappa^3)$.[30] The comparison is singular at 0 because of the term $3A/\kappa$, and we carry it out with an integrating factor. Let $D = A - \phi_-$. Then

$$\begin{aligned} D' &= \mathcal{F}(\kappa,A) - \mathcal{F}(\kappa,\phi_-) - (\phi_-' - \mathcal{F}(\kappa,\phi_-)) \\ &= -\Big(A + \phi_- + \frac{3}{\kappa}\Big)D + R, \end{aligned}$$

with $R = \kappa^2/(s(s+2)^2) \geq 0$. Put $\mu(\kappa) = \kappa^3\exp\int_0^\kappa (A+\phi_-)$. Then $(\mu D)' = \mu R \geq 0$, and $\mu D = O(\kappa^6) \to 0$ as $\kappa \to 0$, so $\mu D \geq 0$ and $A \geq \phi_-$. The same argument with $D = \phi_+ - A$ and $R = (u-3)^2/(u(1+u)^2)$ gives $A \leq \phi_+$. This is Lemma I.2(ii).

For (iv) let $s = \sqrt{\kappa^2 + 25/4}$ and $\phi_\flat = \kappa/(\frac{3}{2}+s)$. Since $s^2 - \frac{9}{4} = \kappa^2 + 4$, the same differentiation gives

$$\phi_\flat' - \mathcal{F}(\kappa,\phi_\flat) = -\frac{5\,(s-\frac{5}{2})}{2\,s\,(\frac{3}{2}+s)^2} \leq 0,$$

[29] For instance $m_2 = 1 - 3A/\kappa$, $m_3 = A - 3/\kappa + 12A/\kappa^2$ and $m_4 = m_2 - 5m_3/\kappa + 2A/\kappa$. The joint moments with x_2, x_3, x_4 follow from exchangeability: $\mathbb{E}[\phi(x_1)x_2^2] = \frac{1}{3}\mathbb{E}[\phi(x_1)(1-x_1^2)]$.

[30] For the first identity write $\phi_- = (s-2)/\kappa$, which follows from $s^2 - 4 = \kappa^2$; then $\phi_-' = (\kappa^2/s - s + 2)/\kappa^2$ and $1 - \phi_-^2 - 3\phi_-/\kappa = (\kappa^2 - (s-2)^2 - 3(s-2))/\kappa^2$, and the difference simplifies with $s^2 = \kappa^2 + 4$. The second is the same computation with $\phi_+ = \kappa(u-1)/(u^2-1) = \kappa(u-1)/(\kappa^2+8)$ in place of ϕ_- and $u^2 = \kappa^2 + 9$ in place of $s^2 = \kappa^2 + 4$. After clearing the denominator $\kappa^2(1+u)^2$ the numerator becomes $\kappa^2(u-3)^2/u$, which is the stated residual.

so $\phi_\flat$ is a subsolution, and $\phi_\flat = \frac{\kappa}{4} - \frac{\kappa^3}{80} + O(\kappa^5)$.[31] Hence $A - \phi_\flat = \frac{\kappa^3}{480} + O(\kappa^5)$, and the integrating-factor argument above, with $D = A - \phi_\flat$, $R = \frac{5(s-5/2)}{2s(3/2+s)^2} \geq 0$ and $\mu = \kappa^3 \exp\int_0^\kappa (A + \phi_\flat)$, gives $\mu D = O(\kappa^6)$ at 0, $(\mu D)' = \mu R \geq 0$, and so $A \geq \phi_\flat$. Since $\frac{3}{2} + \sqrt{\kappa^2 + 25/4} < 2 + \sqrt{\kappa^2 + 4}$ for $\kappa > 0$, this improves the lower bound of (ii).

Appendix K. Bessel functions

This appendix proves the auxiliary statements used in Appendix I for Theorem I.4. Throughout, $I_j = I_j(\kappa)$ with $\kappa > 0$, and $I_{-j} = I_j$ for integers j. We use the series

$$I_\nu(\kappa) = \sum_{i \geq 0} \frac{(\kappa/2)^{2i+\nu}}{i!\,(i+\nu)!},$$

the integral $I_j(\kappa) = \pi^{-1} \int_0^\pi e^{\kappa \cos\theta} \cos j\theta \, d\theta$, the Fourier series $e^{\kappa\cos\theta} = \sum_{j\in\mathbb{Z}} I_j e^{ij\theta}$ and the recurrences

$$I_{j-1} - I_{j+1} = \frac{2j}{\kappa} I_j, \quad I_j' = I_{j+1} + \frac{j}{\kappa} I_j, \tag{K.1}$$

all of which are in Watson's treatise [81], §§3.7, 3.71, 2.22. All series below converge absolutely, because $0 < I_j \leq (\kappa/2)^j e^\kappa / j!$.

K.1. **Zonal harmonics.** Under σ the coordinate x_1 has the density $\frac{2}{\pi}(1-t^2)^{1/2}$ on $[-1,1]$, the area of the two-sphere $\{x_1 = t\}$ being proportional to $1 - t^2$ and its normal speed to $(1-t^2)^{-1/2}$. The Chebyshev polynomials $U_n(\cos\theta) = \sin((n+1)\theta)/\sin\theta$ satisfy

$$\int U_n U_m \, d\sigma = \frac{2}{\pi} \int_0^\pi \sin((n+1)\theta) \sin((m+1)\theta)\, d\theta = \delta_{nm},$$

and $U_n = C_n^{(1)}$ is the Gegenbauer polynomial of index $\frac{4-2}{2} = 1$, so $U_n(\langle y, x\rangle)$ is the zonal harmonic of degree n on S^3 with pole y: $|x|^n U_n(\langle y, x\rangle/|x|)$ is a harmonic polynomial on $\mathbb{R}^4$ [77], Ch. IV, §2.

Lemma K.1. *$e^{\kappa t} = \frac{2}{\kappa} \sum_{n\geq 0} (n+1) I_{n+1}(\kappa)\, U_n(t)$ in $L^2\big(\frac{2}{\pi}(1-t^2)^{1/2} dt\big)$, and uniformly on $[-1,1]$.*

Proof. With $t = \cos\theta$ and $2\sin\theta\sin((n+1)\theta) = \cos n\theta - \cos(n+2)\theta$, the n-th coefficient is

$$\begin{aligned}\frac{2}{\pi} \int_0^\pi e^{\kappa\cos\theta} \sin((n+1)\theta) \sin\theta \, d\theta &= I_n - I_{n+2} \\ &= \frac{2(n+1)}{\kappa} I_{n+1},\end{aligned}$$

by (K.1). Since $|U_n| \leq n+1$ and I_{n+1} decays factorially, the series converges uniformly. □

K.2. **Proof of Lemma I.7.** Let $F(\theta) = e^{\kappa\cos\theta} = \sum_j I_j e^{ij\theta}$. Then

$$F'(\theta) = -\kappa \sin\theta\, F(\theta) = \sum_j ij I_j e^{ij\theta},$$

and Parseval's formula gives

$$\begin{aligned}\sum_{j\in\mathbb{Z}} j^2 I_j^2 &= \frac{\kappa^2}{2\pi} \int_0^{2\pi} \sin^2\theta\, e^{2\kappa\cos\theta} d\theta \\ &= \frac{\kappa^2}{2}\big(I_0(2\kappa) - I_2(2\kappa)\big) = \frac{\kappa}{2} I_1(2\kappa),\end{aligned}$$

[31] With $\phi_\flat = \kappa(s - \frac{3}{2})/(\kappa^2 + 4)$ one has $\phi_\flat' = (\frac{3}{2}s + \frac{25}{4})/(s(\frac{3}{2}+s)^2)$, using $ds/d\kappa = \kappa/s$; and $\mathcal{F}(\kappa, \phi_\flat) = 1 - \phi_\flat^2 - 3/(\frac{3}{2}+s)$. Clearing the denominator $s(\frac{3}{2}+s)^2$ and replacing κ^2 by $s^2 - \frac{25}{4}$ leaves $-\frac{5}{2}s + \frac{25}{4}$ in the numerator, which is the stated residual. The expansion follows from $s = \frac{5}{2} + \frac{\kappa^2}{5} + O(\kappa^4)$.

the last step by (K.1) at the argument 2κ. Halving gives the first identity. The coefficient of $e^{i\theta}$ in $F^2 = e^{2\kappa\cos\theta}$ is $I_1(2\kappa) = \sum_{j\in\mathbb{Z}} I_j I_{1-j}$, and the terms with $j \le 0$ repeat those with $j \ge 1$; this is the third identity.

For the second, $jI_j = \frac{\kappa}{2}(I_{j-1} - I_{j+1})$ by (K.1), so

$$\sum_{j\ge1} j^3 I_j^2 = \frac{\kappa^2}{4} \sum_{j\ge1} j(I_{j-1}^2 + I_{j+1}^2 - 2I_{j-1}I_{j+1}).$$

Shifting indices and telescoping,

$$\textstyle\sum_{j\ge1} j(I_{j-1}^2 + I_{j+1}^2) = I_0^2 + \sum_{m\ge1} 2mI_m^2,$$
$$\textstyle\sum_{m\ge1} 2mI_m^2 = \kappa \sum_{m\ge1}(I_{m-1}I_m - I_m I_{m+1}) = \kappa I_0 I_1.$$

For the mixed term write $j = \frac{1}{2}((j-1)+(j+1))$ and apply (K.1) to $(j-1)I_{j-1}$ and to $(j+1)I_{j+1}$, using $I_{-1} = I_1$ at $j = 1$:

$$\begin{aligned} jI_{j-1}I_{j+1} = \frac{\kappa}{4}\big[&(I_{j-1}I_j - I_j I_{j+1}) \\ &+ (I_{j-2}I_{j+1} - I_{j-1}I_{j+2})\big]. \end{aligned}$$

Both brackets telescope, to $I_0 I_1$ and to $I_{-1}I_2 = I_1 I_2$, so $\sum_{j\ge1} jI_{j-1}I_{j+1} = \frac{\kappa}{4}(I_0 I_1 + I_1 I_2)$. Altogether

$$\sum_{j\ge1} j^3 I_j^2 = \frac{\kappa^2}{4}\Big(I_0^2 + \frac{\kappa}{2} I_1(I_0 - I_2)\Big) = \frac{\kappa^2}{4}(I_0^2 + I_1^2). \qquad \square$$

K.3. **The coefficients of $\mathcal{D}$.**

Lemma K.2. *Write $\mathcal{D}(\kappa) = \sum_{n\ge0} \mathcal{D}_n \kappa^{2n}$ for the function of (I.9). Then $\mathcal{D}_0 = \mathcal{D}_1 = 0$, and for $m \ge 1$*

$$\mathcal{D}_{m+1} = \frac{N_m}{4^m\, m!\,(m+1)!\,(m+1)(m+2)},$$

with $N_m = \binom{2m}{m}\big(4(2m+1) + (m-1)(m+1)(m+2)\big) - 4^m(m+1)(m+2)$. Moreover $N_m = 0$ for $m \le 2$ and $N_m > 0$ for $m \ge 3$.

Proof. The coefficients. Multiplying the series of I_0 and I_1 and collecting powers,

$$I_0^2 = \sum_{m\ge0} \alpha_m \kappa^{2m}, \qquad I_1^2 = \sum_{m\ge0} \beta_m \kappa^{2m+2},$$
$$\kappa I_1(2\kappa) = \sum_{m\ge0} \frac{\kappa^{2m+2}}{m!\,(m+1)!},$$

where

$$\alpha_m = \frac{1}{4^m} \sum_{i=0}^{m} \frac{1}{i!^2 (m-i)!^2},$$
$$\beta_m = \frac{1}{4^{m+1}} \sum_{i=0}^{m} \frac{1}{i!\,(i+1)!\,(m-i)!\,(m-i+1)!}.$$

By Vandermonde's identity, $\sum_i \binom{m}{i}^2 = \binom{2m}{m}$ and $\sum_i \binom{m}{i}\binom{m+2}{i+1} = \binom{2m+2}{m+1}$, so

$$\alpha_m = \frac{\binom{2m}{m}}{4^m\, m!^2}, \qquad \beta_m = \frac{\binom{2m+2}{m+1}}{4^{m+1} m!\,(m+2)!}.$$

The coefficient of κ^{2m+2} in $\mathcal{D}$ is $\mathcal{D}_{m+1} = 8\beta_m + 2\beta_{m-1} - \alpha_m - 1/(m!(m+1)!)$, with $\beta_{-1} = 0$; for $m = 0$ it is $\frac{8}{4} - 1 - 1 = 0$, and $\mathcal{D}_0 = 0$ because every term of $\mathcal{D}$ is $O(\kappa^2)$. For $m \ge 1$ put

$\Delta = 4^m\, m!\,(m+1)!\,(m+1)(m+2)$ and $b = \binom{2m}{m}$. With $\binom{2m+2}{m+1} = \frac{2(2m+1)}{m+1} b$,

$$8\beta_m = \frac{4(2m+1)\,b}{\Delta}, \qquad \alpha_m = \frac{(m+1)^2(m+2)\,b}{\Delta},$$
$$2\beta_{m-1} = \frac{2m(m+1)(m+2)\,b}{\Delta},$$
$$\frac{1}{m!(m+1)!} = \frac{4^m(m+1)(m+2)}{\Delta},$$

and $2m-(m+1) = m-1$ gives $\mathcal{D}_{m+1} = N_m/\Delta$.

The sign. Directly, $N_0 = N_1 = N_2 = 0$ $(2-2,\ 24-24,\ 192-192)$, $N_3 = 80$, $N_4 = 1140$ and $N_5 = 10416$. For $m \geq 6$ we first show $16^m \leq 4m\binom{2m}{m}^2$ for $m \geq 1$ by induction: it is an equality at $m = 1$, and $(m+1)\binom{2m+2}{m+1} = 2(2m+1)\binom{2m}{m}$ gives

$$4(m+1)\binom{2m+2}{m+1}^2 = \frac{16(2m+1)^2}{m+1}\binom{2m}{m}^2$$
$$\geq \frac{16(2m+1)^2}{4m(m+1)} 16^m \geq 16^{m+1}.$$

Since $(m-1)^2 \geq 4m$ for $m \geq 6$, it follows that $\binom{2m}{m}^2 (m-1)^2 \geq 16^m$, that is $\binom{2m}{m}(m-1) \geq 4^m$, and

$$N_m = \Big(\binom{2m}{m}(m-1) - 4^m\Big)(m+1)(m+2)$$
$$+ 4(2m+1)\binom{2m}{m} > 0.$$

□

K.4. **Monotonicity of Bessel ratios.**

Lemma K.3. *For $n \geq 1$ and $\kappa > 0$, $I_{n+2}(\kappa)/I_{n+1}(\kappa) \leq I_2(\kappa)/I_1(\kappa)$.*

Proof. We use a lemma of Biernacki and Krzyż [20]: if $f = \sum a_j x^j$ and $g = \sum b_j x^j$ have positive coefficients and a_j/b_j is nonincreasing in j, then f/g is nonincreasing on $x > 0$. Indeed

$$f'g - fg' = \sum_{i,j} (i-j) a_i b_j x^{i+j-1}$$
$$= \frac{1}{2}\sum_{i,j} (i-j) b_i b_j \Big(\frac{a_i}{b_i} - \frac{a_j}{b_j}\Big) x^{i+j-1} \leq 0.$$

Take $x = \kappa^2$, $f = I_{n+1}(\kappa)/\kappa^{n+1}$ and $g = I_1(\kappa)/\kappa$. Then $a_j/b_j = 2^{-n}(j+1)!/(j+n+1)!$ decreases in j, so $I_{n+1}/(\kappa^n I_1)$ is nonincreasing in κ. Its logarithmic derivative is, by the second recurrence of (K.1),

$$\frac{n+1}{\kappa} + \frac{I_{n+2}}{I_{n+1}} - \frac{n}{\kappa} - \frac{1}{\kappa} - \frac{I_2}{I_1}$$
$$= \frac{I_{n+2}}{I_{n+1}} - \frac{I_2}{I_1} \leq 0.$$

□

Appendix L. The certificate bound

This appendix derives the bound $h_F \leq 14.420001$ of Remark 6.8 from the data of [69]. The argument is the proof of the bound in Theorem 1.4 with two changes. The divergence identity of Lemma 6.1, taken there with the factor $\frac{9}{10}$, is replaced by a positive semidefinite form at each site, and the weights $\frac{21}{10}$ and $\frac{5}{4}$ of Lemma 6.2 are replaced by weights chosen plaquette

by plaquette. Everything is proved by hand except the positivity of ninety-six matrices of trigonometric polynomials and the arithmetic of finitely many rational numbers.[32]

L.1. **Ends and corners.** At a site y the eight *ends* are

$$o_\mu(y) = X_{y,\mu}, \qquad i_\mu(y) = Q^*_{y-e_\mu,\mu} X_{y-e_\mu,\mu} Q_{y-e_\mu,\mu},$$

for $\mu = 0,1,2,3$, as in the proof of Lemma 6.1; the end of a link in F or outside the box is 0. Each link $e = (y,\mu)$ has the end o_μ at its tail y and the end i_μ at its head $y + e_\mu$, both of norm $|X_e|$.

Lemma L.1. *Let p be the plaquette at x in the plane $\mu < \nu$. Then*

$$\begin{aligned}
\langle Y_1, Y_4 \rangle &= \langle o_\mu, o_\nu \rangle(x), \\
\langle Y_1, Y_2 \rangle &= \langle i_\mu, o_\nu \rangle(x + e_\mu), \\
\langle Y_3, Y_4 \rangle &= \langle o_\mu, i_\nu \rangle(x + e_\nu), \\
\langle Y_2, W Y_3 W^* \rangle &= \langle i_\nu, i_\mu \rangle(x + e_\mu + e_\nu).
\end{aligned}$$

Every pair of ends at one site in two different directions occurs in exactly one of these identities, for exactly one plaquette.

Proof. The first identity is the definition of the ends. The inner product is invariant under conjugation by a unitary matrix, so $\langle X_{e_1}, Q_{e_1} X_{e_2} Q^*_{e_1} \rangle = \langle Q^*_{e_1} X_{e_1} Q_{e_1}, X_{e_2} \rangle$, which is the second identity, and the third follows in the same way from $\langle Q_{e_4} X_{e_3} Q^*_{e_4}, X_{e_4} \rangle = \langle X_{e_3}, Q^*_{e_4} X_{e_4} Q_{e_4} \rangle$. For the fourth, $W Q_{e_4} = Q_{e_1} Q_{e_2} Q^*_{e_3}$, so that $W Y_3 W^* = R X_{e_3} R^*$ with $R = Q_{e_1} Q_{e_2} Q^*_{e_3}$, and conjugating both vectors by $(Q_{e_1} Q_{e_2})^*$ gives $\langle Q^*_{e_2} X_{e_2} Q_{e_2}, Q^*_{e_3} X_{e_3} Q_{e_3} \rangle$. The links e_2 and e_3 both have their head at $x + e_\mu + e_\nu$. For the last statement, the pairs (o_μ, o_ν), (i_μ, o_ν), (o_μ, i_ν) and (i_μ, i_ν) at y come from the plaquettes in the plane $\mu\nu$ at y, $y - e_\mu$, $y - e_\nu$ and $y - e_\mu - e_\nu$ respectively. There are $4\binom{4}{2} = 24$ such pairs at each site and $6 \cdot 4 = 24$ corners of plaquettes per site. □

For each class of sites modulo 2 let G_y be a real symmetric 8×8 matrix indexed by the ends, with $G_y[o_\mu, i_\mu] = 0$ for every μ. For the plaquette p at x in the plane $\mu < \nu$ put

$$\begin{aligned}
c^G_p = {} & 2 G_x[o_\mu, o_\nu] \langle Y_1, Y_4 \rangle + 2 G_{x+e_\mu}[i_\mu, o_\nu] \langle Y_1, Y_2 \rangle \\
& + 2 G_{x+e_\nu}[o_\mu, i_\nu] \langle Y_3, Y_4 \rangle \\
& + 2 G_{x+e_\mu+e_\nu}[i_\mu, i_\nu] \langle Y_2, W Y_3 W^* \rangle.
\end{aligned}$$

The matrix G^0 with $G^0[a,a] = 2$, $G^0[o_\mu, o_\nu] = G^0[i_\mu, i_\nu] = 1$ and $G^0[o_\mu, i_\nu] = -1$ for $\mu \neq \nu$ is the matrix of the form $|\sum_\mu (o_\mu - i_\mu)|^2 + \sum_\mu |o_\mu + i_\mu|^2$ of Lemma 6.1, and $c^{G^0}_p$ is the corner form c_p of (6.3).

Lemma L.2. *On $\mathbb{Z}^4$, or on a box with free boundary, for every field Q and every square-summable X,*

$$\begin{aligned}
& \sum_y \sum_{a,b} G_y[a,b] \, \langle \mathrm{end}_a, \mathrm{end}_b \rangle(y) \\
& \qquad = \sum_p c^G_p + \sum_e \big(G_{t(e)}[o,o] + G_{h(e)}[i,i] \big) |X_e|^2,
\end{aligned}$$

where $t(e)$ and $h(e)$ are the tail and the head of e, and o, i are its two ends. If every G_y is positive semidefinite, the left side is nonnegative.

Proof. Split the sum over a, b at each site into the diagonal terms, the pairs (o_μ, i_μ), whose coefficient is 0, and the pairs of ends in different directions, each counted twice. The diagonal terms give the last sum, since each link has one end at its tail and one at its head, and the

[32] The constant 14.420001 is not the best possible for this kind of certificate: by the search reported in Remark 6.8 it cannot go below 14.384, while the true value of h_F is at least 13.123.

pairs in different directions give $\sum_p c_p^G$ by Lemma L.1. In a box the argument at the end of the proof of Lemma 6.1 applies. At each site the left side is the trace of G_y times the Gram matrix of the eight ends, and the trace of a product of two positive semidefinite matrices is nonnegative. □

L.2. **From plaquettes to the Hessian.** Number the positions of a plaquette as in (6.1) and let k_p be the position of its link in F; by Lemma 2.2 there is exactly one.

Lemma L.3. *Let every G_y be positive semidefinite with $G_y[o_\mu, i_\mu] = 0$, and let $a_{p,r} \geq 0$ for the three positions $r \neq k_p$ of each plaquette. Suppose that for every plaquette p, every N, every $W \in \mathrm{U}(N)$ and all $Y_1, \dots, Y_4 \in \mathfrak{u}(N)$ with $Y_{k_p} = 0$,*

$$f_p + c_p^G \leq \sum_{r \neq k_p} a_{p,r} |Y_r|^2. \tag{L.1}$$

Then $|\operatorname{Hess} S_0(v,v)| \leq h|v|^2$ for every field Q and every $v = (X_e Q_e)$ with $X_e = 0$ for $e \in F$, on $\mathbb{Z}^4$ and on every box, where

$$h = \max_{e \notin F} \Big(\sum_{p \ni e} a_{p,e} + G_{t(e)}[o,o] + G_{h(e)}[i,i] \Big)$$

and $a_{p,e}$ is the weight of the position of e in p.

Proof. Summing (L.1) over the plaquettes and adding the identity of Lemma L.2, whose left side is nonnegative,

$$\begin{aligned} \operatorname{Hess} S_0(v,v) = \sum_p f_p &\leq \sum_p \Big(\sum_r a_{p,r} |Y_r|^2 - c_p^G \Big) \\ &\leq \sum_{e \notin F} \Big(\sum_{p \ni e} a_{p,e} + G_{t(e)}[o,o] + G_{h(e)}[i,i] \Big) |X_e|^2 \\ &\leq h|v|^2, \end{aligned}$$

since $|Y_r| = |X_e|$ for the link e at position r. In a box the links near the boundary lie in fewer plaquettes, and the weights are nonnegative. For the lower bound, apply (L.1) at $-W$. By (6.2) this changes the sign of f_p, while c_p^G does not change, and the same chain gives $-\operatorname{Hess} S_0(v,v) \leq h|v|^2$. □

The hypothesis is needed on $\mathrm{U}(N)$ and not only on $\mathrm{SU}(N)$, since $-W \notin \mathrm{SU}(N)$ when N is odd.

Lemma L.4. *If (L.1) holds for $N = 2$ and every $W = \operatorname{diag}(e^{i\varphi_1}, e^{i\varphi_2})$, it holds for every N and every $W \in \mathrm{U}(N)$.*

Proof. Both sides are invariant under conjugation of W and all Y_i by one unitary matrix, so we may take $W = \operatorname{diag}(w_1, \dots, w_N)$. Every term of f_p, of c_p^G and of $|Y_r|^2$ is a multiple of $\operatorname{Re}\operatorname{Tr}(Y_i D_1 Y_j D_2)$ with $D_1, D_2 \in \{I, W, W^*\}$, and

$$\operatorname{Re}\operatorname{Tr}(Y_i D_1 Y_j D_2) = \operatorname{Re} \sum_{a,b} (Y_i)_{ab} \, (D_1)_{bb} \, (Y_j)_{ba} \, (D_2)_{aa}.$$

The difference of the two sides of (L.1) is therefore a sum of one term for each a, in the entries $(Y_i)_{aa}$, and one term for each pair $a < b$, in the entries $(Y_i)_{ab}$ and $(Y_i)_{ba} = -\overline{(Y_i)_{ab}}$, as in the proof of Lemma 6.2. The term of the pair $a < b$ is the same difference for $N = 2$ at $W' = \operatorname{diag}(w_a, w_b)$, evaluated on the matrices $Y_i' \in \mathfrak{u}(2)$ with off-diagonal entries $(Y_i)_{ab}$, $(Y_i)_{ba}$ and zero diagonal. The term of a is the difference at W', for any $b \neq a$, evaluated on $Y_i' = \operatorname{diag}((Y_i)_{aa}, 0)$. In both cases $Y_{k_p}' = 0$, so each term is nonnegative. □

In the orthonormal basis $iI/\sqrt{2}$, $i\sigma_1/\sqrt{2}$, $i\sigma_2/\sqrt{2}$, $i\sigma_3/\sqrt{2}$ of $\mathfrak{u}(2)$, with the Pauli matrices σ_j, the difference of the two sides of (L.1) at $N = 2$ is a quadratic form on the three free positions, with a real symmetric 12×12 matrix $M_p(\varphi)$. Its entries are affine combinations




FIGURE 24. The certificate of Appendix L. (a) The least eigenvalue of $M_p(\varphi)$ over the torus of phases for the plaquette class with the smallest minimum, about 0.0057 on a grid of 96^2 phases (red dot); Theorem L.5 shows it stays positive. (b) The forty-eight totals in the definition of h, in increasing order, against $\frac{29}{2}$, the value 14.420001 of the largest, the floor 14.384 for certificates on the period-two cell, and the lower bound $9+\sqrt{17}$ for h_F. (c) The constant C in $\sum \phi^2 \le CQ$ for the block gauge: 52 from Theorem 12.4, 5000/537 from Proposition 12.5, and the bulk Bloch value $1/0.10749 = 9.3034$, the best constant when W is the set of all blocks. The eigenvalues in (a) and the Bloch value are in floating point.

of 1, $\cos\varphi_a$, $\sin\varphi_a$, $\cos(\varphi_1-\varphi_2)$ and $\sin(\varphi_1-\varphi_2)$: the form f_p is linear in W, and W enters c_p^G only through $\langle Y_2, WY_3W^*\rangle$. Since W commutes with I and σ_3, $M_p(\varphi)$ splits into two blocks of size six, one on the span of $iI, i\sigma_3$ and one on the span of $i\sigma_1, i\sigma_2$, at every position. By Lemmas L.3 and L.4, a certificate is a choice of G_y and $a_{p,r}$ such that every $M_p(\varphi)$ is positive semidefinite at every φ, and it gives $h_F \le h$.

L.3. **The certificate.** The fixed set F and the field of ends have period two, so it suffices to give G_y for the sixteen classes of sites and $a_{p,r}$ for the ninety-six classes of plaquettes, six planes at each of sixteen sites. The certificate of [69] consists of sixteen positive definite matrices G_y of size eight with $G_y[o_\mu, i_\mu] = 0$, and of nonnegative weights $a_{p,r}$, all rational with denominator 10^6. We checked that the set of fixed links used there is our F up to a symmetry of the lattice. Of the sixty-four links of a period cell, sixteen lie in F, one ending at each site. For each of the other forty-eight the bracket in the definition of h is computed exactly, and the largest is $14420001/10^6$.

Theorem L.5 (Computer-assisted)**.** *For every N, every field Q and every $v = (X_eQ_e)$ with $X_e = 0$ for $e \in F$, on $\mathbb{Z}^4$ and on every box,*

$$|\operatorname{Hess} S_0(v,v)| \le 14.420001\,|v|^2.$$

Proof. We show that every $M_p(\varphi)$ is positive semidefinite, which proves the theorem by Lemmas L.3 and L.4. Write $z = (1, \cos\varphi_1, \sin\varphi_1, \cos\varphi_2, \sin\varphi_2)$. For each of the 192 blocks of size six there is a positive definite integer symmetric matrix P of size thirty such that

$$2^{24}\cdot 4\cdot 10^6\, M(\varphi)_{jk} = \sum_{a,b=0}^{4} z_a z_b\, P_{6a+j,\,6b+k}$$

for all j, k, as polynomials modulo $\cos^2\varphi_a + \sin^2\varphi_a = 1$. For $x \in \mathbb{R}^6$ put $w = z \otimes x$. Then $2^{24}\cdot 4\cdot 10^6\, x^{\mathsf{T}} M(\varphi) x = w^{\mathsf{T}} P w$, which is positive for $x \ne 0$, since $z_0 = 1$ gives $w \ne 0$. So every block is positive definite at every φ. □

The proof of Theorem L.5 rests on computations of three kinds. The positivity of the G_y, the signs of the weights and the forty-eight totals are finite statements about rational

numbers. The identities between $M(\varphi)$ and the matrices P, and the positive definiteness of the P, are finite statements about integers. All of these are checked by the Lean 4 kernel in [69], on no axioms beyond `propext` and `Quot.sound`; the program there also checks the positivity at every phase a second time, by interval arithmetic on a subdivision of the torus of phases. Lemmas L.1 to L.4 and the formula (6.2) for the Hessian are the written proofs above and are not formalised.[33] We checked the finite data in exact arithmetic and the positivity at every phase by eigenvalues on an adaptive grid with a Lipschitz bound, in floating point, and J. M. Lockwood checked it again in exact rational arithmetic with outward-rounded interval elimination, finding least eigenvalue above 559/716800 for every plaquette.

With 14.420001 in place of $\frac{29}{2}$ the proofs of Theorems 1.4 and 1.5 go through unchanged, which gives the ranges stated in Remark 6.8. Those theorems are stated with $\frac{29}{2}$ because its proof uses no computer.

References

[1] Adhikari, A., Cao, S.: Correlation decay for finite lattice gauge theories at weak coupling. Ann. Probab. **53**, 140–174 (2025). https://doi.org/10.1214/24-AOP1702

[2] Aumann, G.: Über Räume mit Mittelbildungen. Math. Ann. **119**, 210–215 (1944). https://doi.org/10.1007/BF01563741

[3] L. A. Bassalygo and R. L. Dobrushin, Uniqueness of a Gibbs field with a random potential—an elementary approach, Theory Probab. Appl. **31** (1987), 572–589, doi:10.1137/1131082.

[4] Bakry, D., Émery, M.: Diffusions hypercontractives. In: Séminaire de Probabilités XIX. Lecture Notes in Math., vol. 1123, pp. 177–206. Springer, Berlin (1985). https://doi.org/10.1007/BFb0075847

[5] Bakry, D., Gentil, I., Ledoux, M.: Analysis and Geometry of Markov Diffusion Operators. Grundlehren Math. Wiss., vol. 348. Springer, Cham (2014). https://doi.org/10.1007/978-3-319-00227-9

[6] Bałaban, T.: Propagators and renormalization transformations for lattice gauge theories. I. Commun. Math. Phys. **95**, 17–40 (1984). https://doi.org/10.1007/BF01215753

[7] Bałaban, T.: Averaging operations for lattice gauge theories. Commun. Math. Phys. **98**, 17–51 (1985). https://doi.org/10.1007/BF01211042

[8] Bałaban, T.: Large field renormalization. II. Localization, exponentiation, and bounds for the **R** operation. Commun. Math. Phys. **122**, 355–392 (1989). https://doi.org/10.1007/BF01238433

[9] S. Banach, Über homogene Polynome in (L^2), Studia Math. **7** (1938), 36–44, doi:10.4064/sm-7-1-36-44.

[10] Barta, J.: Sur la vibration fondamentale d'une membrane. C. R. Acad. Sci. Paris **204**, 472–473 (1937)

[11] Basu, R., Ganguly, S.: SO(N) lattice gauge theory, planar and beyond. Commun. Pure Appl. Math. **71**, 2016–2064 (2018). https://doi.org/10.1002/cpa.21788

[12] Bhattacharjee, D.: A coherent approach towards quantum gravity. Phys. Sci. Int. J. **26**(6), 59–78 (2022). https://doi.org/10.9734/psij/2022/v26i6751

[13] Bhattacharjee, D.: BLLM: Beyond the LLM with Deep Human-Mind-AI. Zenodo, release v3.0.5 (2026). https://doi.org/10.5281/zenodo.23469389

[14] Bhattacharjee, D.: BLLM: Beyond the LLM with Deep Human-Mind-AI. Zenodo, release v3.1.0 (2026). https://doi.org/10.5281/zenodo.23484160

[15] Bhattacharjee, D., Bhattacharya, U.: Mass gap of lattice Yang–Mills theory up to the Dobrushin ceiling. In: Lattice Yang–Mills theory: spacing-uniform obstructions, the mass gap up to the Dobrushin ceiling, and Lean certificate. Zenodo, release v2.0.2 (2026). https://doi.org/10.5281/zenodo.23449466

[16] Bhattacharjee, D., Bhattacharya, U.: Spacing-uniform lattice inequalities cannot produce the Yang–Mills mass gap. Zenodo, release v1.0.0 (2026). https://doi.org/10.5281/zenodo.23444339

[17] Bhattacharjee, D., Singha Roy, S., Samal, P.: Brane-cluster UV completion of quantum gravity. Research Square (2026). https://doi.org/10.21203/rs.3.rs-9053205/v1

[18] Bhattacharjee, D., Samal, P., Bhattacharya, S.: Geometric structure and renormalization group flow in chiral Yang–Mills–Higgs theory. Preprints (2026). https://doi.org/10.20944/preprints202602.2022.v1

[19] Bhattacharjee, D., Nath Thakur, S., Bhattacharya, S., Samal, P.: SU(n) holonomy deviations in Calabi–Yau manifolds (CY1–CY4). Preprints (2026). https://doi.org/10.20944/preprints202602.0023.v1

[20] Biernacki, M., Krzyż, J.: On the monotonity of certain functionals in the theory of analytic functions. Ann. Univ. Mariae Curie-Skłodowska Sect. A **9**, 135–147 (1955)

[33] Formalising them needs matrices over $\mathbb{C}$, the group U(N) and second derivatives along geodesics, that is, a library such as Mathlib; the Lean files of [69] use only the core library.

[21] M. Biskup, Reflection positivity and phase transitions in lattice spin models, in *Methods of Contemporary Mathematical Statistical Physics*, Lecture Notes in Math. 1970, Springer, Berlin, 2009, 1–86, doi:10.1007/978-3-540-92796-9_1.
[22] Brydges, D., Fröhlich, J., Seiler, E.: On the construction of quantized gauge fields. I. General results. Ann. Phys. **121**, 227–284 (1979). https://doi.org/10.1016/0003-4916(79)90098-8
[23] Bump, D.: Lie Groups, 2nd edn. Grad. Texts in Math., vol. 225. Springer, New York (2013). https://doi.org/10.1007/978-1-4614-8024-2
[24] Cao, S., Nissim, R., Sheffield, S.: Expanded regimes of area law for lattice Yang–Mills theories. Preprint (2025). arXiv:2505.16585
[25] Chatterjee, S.: Yang–Mills for probabilists. In: Probability and Analysis in Interacting Physical Systems, Springer Proc. Math. Stat., vol. 283, pp. 1–16. Springer, Cham (2019). https://doi.org/10.1007/978-3-030-15338-0_1
[26] Chatterjee, S.: Rigorous solution of strongly coupled SO(N) lattice gauge theory in the large N limit. Commun. Math. Phys. **366**, 203–268 (2019). https://doi.org/10.1007/s00220-019-03353-3
[27] Creutz, M.: Gauge fixing, the transfer matrix, and confinement on a lattice. Phys. Rev. D **15**, 1128–1136 (1977). https://doi.org/10.1103/PhysRevD.15.1128
[28] J. M. Combes and L. Thomas, Asymptotic behaviour of eigenfunctions for multiparticle Schrödinger operators, Comm. Math. Phys. **34** (1973), 251–270, doi:10.1007/BF01646473.
[29] Dimock, J.: The renormalization group according to Balaban, I. Small fields. Rev. Math. Phys. **25**, 1330010 (2013). https://doi.org/10.1142/S0129055X13300100
[30] Dobrushin, R.L.: The description of a random field by means of conditional probabilities and conditions of its regularity. Theory Probab. Appl. **13**, 197–224 (1968). https://doi.org/10.1137/1113026
[31] Dobrushin, R.L.: Prescribing a system of random variables by conditional distributions. Theory Probab. Appl. **15**, 458–486 (1970). https://doi.org/10.1137/1115049
[32] Dobrushin, R.L., Shlosman, S.B.: Constructive criterion for the uniqueness of Gibbs field. In: Statistical Physics and Dynamical Systems, pp. 347–370. Birkhäuser, Boston (1985). https://doi.org/10.1007/978-1-4899-6653-7_20
[33] Driver, B.K.: YM_2: continuum expectations, lattice convergence, and lassos. Commun. Math. Phys. **123**, 575–616 (1989). https://doi.org/10.1007/BF01218586
[34] Eckmann, B.: Räume mit Mittelbildungen. Comment. Math. Helv. **28**, 329–340 (1954). https://doi.org/10.1007/BF02566939
[35] M. Faizal and A. Shabir, Reflection-positive construction of a four-dimensional SU(N) Yang–Mills theory with mass gap and confinement, preprint, 2026, arXiv:2606.19362v1; Fortschr. Phys. **74** (2026), e70097, doi:10.1002/prop.70097.
[36] Fulton, W., Harris, J.: Representation Theory: A First Course. Grad. Texts in Math., vol. 129. Springer, New York (1991; reprinted 2004). https://doi.org/10.1007/978-1-4612-0979-9
[37] J. Fröhlich, R. Israel, E. H. Lieb and B. Simon, Phase transitions and reflection positivity. I. General theory and long range lattice models, Comm. Math. Phys. **62** (1978), 1–34, doi:10.1007/BF01940327.
[38] Fröhlich, J., Spencer, T.: Massless phases and symmetry restoration in abelian gauge theories and spin systems. Commun. Math. Phys. **83**, 411–454 (1982). https://doi.org/10.1007/BF01213610
[39] Gattringer, C., Lang, C.B.: Quantum Chromodynamics on the Lattice. Lecture Notes in Phys., vol. 788. Springer, Berlin (2010). https://doi.org/10.1007/978-3-642-01850-3
[40] Ghora, R., Bhattacharjee, D., Nandi, P., Samal, P., Patra, R., Singha Roy, S., Bhattacharya, S.: O'Neill tensor bounds for Riemannian submersions in fibred Calabi–Yau manifolds. Preprints (2026). https://doi.org/10.20944/preprints202602.0125.v1
[41] Glimm, J., Jaffe, A.: Quantum Physics. A Functional Integral Point of View, 2nd edn. Springer, New York (1987). https://doi.org/10.1007/978-1-4612-4728-9
[42] Gross, L., King, C., Sengupta, A.: Two dimensional Yang–Mills theory via stochastic differential equations. Ann. Phys. **194**, 65–112 (1989). https://doi.org/10.1016/0003-4916(89)90032-8
[43] Guth, A.H.: Existence proof of a nonconfining phase in four-dimensional U(1) lattice gauge theory. Phys. Rev. D **21**, 2291–2307 (1980). https://doi.org/10.1103/PhysRevD.21.2291
[44] Helgason, S.: Differential Geometry, Lie Groups, and Symmetric Spaces. Grad. Stud. Math., vol. 34. Amer. Math. Soc., Providence, RI (2001). https://doi.org/10.1090/gsm/034
[45] Higham, N.J.: Accuracy and Stability of Numerical Algorithms, 2nd edn. SIAM, Philadelphia (2002). https://doi.org/10.1137/1.9780898718027
[46] R. Holley and D. Stroock, Logarithmic Sobolev inequalities and stochastic Ising models, J. Stat. Phys. **46** (1987), 1159–1194, doi:10.1007/BF01011161.
[47] B. Helffer and J. Sjöstrand, On the correlation for Kac-like models in the convex case, J. Stat. Phys. **74** (1994), 349–409, doi:10.1007/BF02186817.
[48] Hoeffding, W.: Probability inequalities for sums of bounded random variables. J. Amer. Statist. Assoc. **58**, 13–30 (1963). https://doi.org/10.1080/01621459.1963.10500830

[49] Humphreys, J.E.: Introduction to Lie Algebras and Representation Theory. Grad. Texts in Math., vol. 9. Springer, New York (1972). https://doi.org/10.1007/978-1-4612-6398-2
[50] Jaffe, A., Witten, E.: Quantum Yang–Mills theory. In: The Millennium Prize Problems, pp. 129–152. Clay Mathematics Institute, Cambridge, MA, and American Mathematical Society, Providence, RI (2006). https://www.claymath.org/wp-content/uploads/2022/06/yangmills.pdf
[51] Johansson, F.: Arb: efficient arbitrary-precision midpoint-radius interval arithmetic. IEEE Trans. Comput. **66**, 1281–1292 (2017). https://doi.org/10.1109/TC.2017.2690633
[52] Kac, V.G.: Infinite-Dimensional Lie Algebras, 3rd edn. Cambridge University Press, Cambridge (1990). https://doi.org/10.1017/CBO9780511626234
[53] Kallenberg, O.: Foundations of Modern Probability, 3rd edn. Springer, Cham (2021). https://doi.org/10.1007/978-3-030-61871-1
[54] Kogut, J., Susskind, L.: Hamiltonian formulation of Wilson's lattice gauge theories. Phys. Rev. D **11**, 395–408 (1975). https://doi.org/10.1103/PhysRevD.11.395
[55] Lévy, T.: Yang–Mills measure on compact surfaces. Mem. Amer. Math. Soc. **166**(790) (2003). https://doi.org/10.1090/memo/0790
[56] J. M. Lockwood, Toward the continuum limit of forest-gauge lattice Yang–Mills theory: constrained cubes, exact Gaussian diagnostics, and a gauge-suited mixing test, technical report, 29 October 2026, distributed with the archive.
[57] Lüscher, M.: Construction of a selfadjoint, strictly positive transfer matrix for Euclidean lattice gauge theories. Commun. Math. Phys. **54**, 283–292 (1977). https://doi.org/10.1007/BF01614090
[58] M. Lüscher, Properties and uses of the Wilson flow in lattice QCD, J. High Energy Phys. **2010**, no. 8, 071, doi:10.1007/JHEP08(2010)071.
[59] M. Lüscher and P. Weisz, Perturbative analysis of the gradient flow in non-abelian gauge theories, J. High Energy Phys. **2011**, no. 2, 051, doi:10.1007/JHEP02(2011)051.
[60] Magnen, J., Rivasseau, V., Sénéor, R.: Construction of YM_4 with an infrared cutoff. Commun. Math. Phys. **155**, 325–383 (1993). https://doi.org/10.1007/BF02097397
[61] Mardia, K.V., Jupp, P.E.: Directional Statistics. Wiley, Chichester (2000). https://doi.org/10.1002/9780470316979
[62] McShane, E.J.: Extension of range of functions. Bull. Amer. Math. Soc. **40**, 837–842 (1934). https://doi.org/10.1090/S0002-9904-1934-05978-0
[63] Milnor, J.: Curvatures of left invariant metrics on Lie groups. Adv. Math. **21**, 293–329 (1976). https://doi.org/10.1016/S0001-8708(76)80002-3
[64] Montvay, I., Münster, G.: Quantum Fields on a Lattice. Cambridge University Press, Cambridge (1994). https://doi.org/10.1017/CBO9780511470783
[65] Osterwalder, K., Schrader, R.: Axioms for Euclidean Green's functions. Commun. Math. Phys. **31**, 83–112 (1973). https://doi.org/10.1007/BF01645738
[66] Osterwalder, K., Schrader, R.: Axioms for Euclidean Green's functions. II. Commun. Math. Phys. **42**, 281–305 (1975). https://doi.org/10.1007/BF01608978
[67] Osterwalder, K., Seiler, E.: Gauge field theories on a lattice. Ann. Phys. **110**, 440–471 (1978). https://doi.org/10.1016/0003-4916(78)90039-8
[68] M. Reed and B. Simon, *Methods of Modern Mathematical Physics I: Functional Analysis*, revised and enlarged ed., Academic Press, New York, 1980.
[69] T. Romanov, A proved bound $h_F \leq 14.420001$ for the four-dimensional forest-gauge Hessian, computer-assisted certificate with Lean 4 kernel checks (2026), https://github.com/romanov360/yang-mills-4d-forest-certificate.
[70] Samal, P., Bhattacharjee, D., Ghora, R.: Renormalization of spacetime dimension. Preprints (2026). https://doi.org/10.20944/preprints202602.0531.v1
[71] K. Sakai and S. Sasaki, Glueball spectroscopy in lattice QCD using gradient flow, Phys. Rev. D **107** (2023), 034510, doi:10.1103/PhysRevD.107.034510.
[72] Seiler, E.: Gauge Theories as a Problem of Constructive Quantum Field Theory and Statistical Mechanics. Lecture Notes in Phys., vol. 159. Springer, Berlin (1982). https://doi.org/10.1007/3-540-11559-5
[73] D. W. Stroock and B. Zegarlinski, The logarithmic Sobolev inequality for discrete spin systems on a lattice, Comm. Math. Phys. **149** (1992), 175–193, doi:10.1007/BF02096629.
[74] Shen, H., Zhu, R., Zhu, X.: A stochastic analysis approach to lattice Yang–Mills at strong coupling. Commun. Math. Phys. **400**, 805–851 (2023). https://doi.org/10.1007/s00220-022-04609-1
[75] Slansky, R.: Group theory for unified model building. Phys. Rep. **79**, 1–128 (1981). https://doi.org/10.1016/0370-1573(81)90092-2
[76] Stanley, R.P.: Enumerative Combinatorics, vol. 2, with an appendix by S. Fomin. Cambridge Stud. Adv. Math., vol. 62. Cambridge University Press, Cambridge (1999). https://doi.org/10.1017/CBO9780511609589

[77] Stein, E.M., Weiss, G.: Introduction to Fourier Analysis on Euclidean Spaces. Princeton University Press, Princeton, NJ (1971). https://doi.org/10.1515/9781400883899
[78] K. K. Uhlenbeck, Connections with L^p bounds on curvature, Comm. Math. Phys. **83** (1982), 31–42, doi:10.1007/BF01947069.
[79] J. van den Berg and C. Maes, Disagreement percolation in the study of Markov fields, Ann. Probab. **22** (1994), 749–763, doi:10.1214/aop/1176988728.
[80] Villani, C.: Optimal Transport: Old and New. Grundlehren Math. Wiss., vol. 338. Springer, Berlin (2009). https://doi.org/10.1007/978-3-540-71050-9
[81] Watson, G.N.: A Treatise on the Theory of Bessel Functions, 2nd edn. Cambridge University Press, Cambridge (1944)
[82] H. Whitney, *Geometric Integration Theory*, Princeton University Press, Princeton, 1957, doi:10.1515/9781400877577.
[83] Wilson, K.G.: Confinement of quarks. Phys. Rev. D **10**, 2445–2459 (1974). https://doi.org/10.1103/PhysRevD.10.2445

Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL), Bhubaneswar, Odisha 751030, India; formerly, University of Calcutta (Heramba Chandra College), 23/49 Gariahat Road, Golpark, Kolkata 700029, West Bengal, India
Email address: itsdeep@live.com, d.bhattacharjee@erl-forschung.de, egspl@proton.me (GitHub github.com/creelie)
URL: orcid.org/0000-0003-0466-750X

Formerly, National Taiwan University, Taipei 106319, Taiwan
Email address: rai.bhattacharya.in@gmail.com
URL: orcid.org/0009-0002-2254-3914

Independent Researcher, San Antonio, USA
Email address: spectralityworkinggroup@gmail.com (secondary Blaze61029dogQ@proton.me)

Independent Researcher, San Francisco, CA, USA
Email address: romanov360@gmail.com