THE BOOK RAMSEY NUMBER $R(B_{57}, B_{58}) = 231$

DEEP BHATTACHARJEE

Abstract. We construct a graph on 230 vertices with no book B_{57} whose complement contains no B_{58}. With the upper bound of Rousseau and Sheehan this gives $R(B_{57}, B_{58}) = 231$, the smallest case of Wesley's conjecture $R(B_{n-1}, B_n) = 4n - 1$ that was open. The graph is a two-block circulant graph over $\mathbb{Z}_{115}$, and the proof reduces to 57 inequalities.

1. Introduction

The book B_k is the graph $K_2 + \overline{K_k}$: k triangles sharing one edge, its spine. For graphs G and H, the Ramsey number $R(G, H)$ is the least N such that every graph on N vertices contains G or has a complement containing H. Rousseau and Sheehan [9] proved, from Goodman's count of monochromatic triangles [6], that

$$R(B_{n-1}, B_n) \le 4n - 1 \qquad (n \ge 2). \tag{1.1}$$

Wesley [12] proved equality for $n \le 20$ and whenever $2n - 1$ is a prime power congruent to 1 modulo 4, and conjectured equality for every n; the conjecture is one of the open problems of Epoch AI's FrontierMath collection [5]. Turturean [11] constructed extremal graphs for every $n \le 57$ and for fourteen further values below 100, partly through a cyclotomic construction of Szekeres [10]; Chung, Du and Wesley [3] found three further infinite families and settled $n = 67$ by a finite certificate [4], for which [2] gives a proof checkable by hand; and Price obtained $n = 70$ [5]. Before the present note the smallest open case was $n = 58$.[1]

Theorem. *$R(B_{57}, B_{58}) = 231$.*

The upper bound is (1.1), and the lower bound is a graph on 230 vertices with no B_{57} whose complement has no B_{58}. For odd n the graph of [2] and the Paley graphs of [12] are regular two-block circulant graphs over $\mathbb{Z}_{2n-1}$ whose two blocks are complementary circulants of degree $n - 1$. When n is even such a graph does not exist, because a symmetric subset of $\mathbb{Z}_{2n-1} \setminus \{0\}$ has even size. Turturean's examples for even n [11] enlarge the first block by one element and shrink the second by one, and our graph has that shape. Section 2 reduces the two book conditions for it to one inequality (2.1) on the connection sets, a reduction that underlies the search of [11] and that we prove here in full. Section 3 gives the connection sets for $n = 58$, and Table 1 verifies (2.1) at the 57 values of d that matter. The graph has also been checked by two independent programs and in Lean 4 (see the end of the paper).

2020 *Mathematics Subject Classification.* Primary 05C55; Secondary 05C25, 05B10.

Key words and phrases. Ramsey number, book graph, block-circulant graph.

Corresponding author: d.bhattacharjee@erl-forschung.de; ORCID 0000-0003-0466-750X.

[1] The values $n \le 100$ that none of the constructions cited here covers were 58, 59, 60, 64, 65, 68, 72, 73, 76, 78, 80, 81, 84, 88, 89, 92, 93, 94, 96 and 100. This list is our reading of [3, 4, 5, 11, 12] as of 15 October 2026; it agrees with the ledger in [4, Section 6.1].

2. Two-block circulant graphs of even order

Throughout, $n \geq 4$ is even, $v = 2n-1$ and $G = \mathbb{Z}_v$. For $X, Y \subseteq G$ and $d \in G$ put

$$\mathrm{N}_{X,Y}(d) = |X \cap (Y+d)| = \#\{y \in Y : \ y+d \in X\}.$$

Translating by $-d$ shows $\mathrm{N}_{X,X}(-d) = \mathrm{N}_{X,X}(d)$, and $\sum_{d\in G} \mathrm{N}_{X,X}(d) = |X|^2$.

Let $A \subseteq G \setminus \{0\}$ with $A = -A$ and $|A| = n$, let $C = G \setminus (A \cup \{0\})$, so that $C = -C$ and $|C| = n-2$, and let $B \subseteq G$ with $|B| = n-1$. The graph $\Gamma = \Gamma(A,B)$ has vertex set $G \times \{0,1\}$, and

$$(x,0) \sim (y,0) \iff y-x \in A, \qquad (x,1) \sim (y,1) \iff y-x \in C,$$

and $(x,0) \sim (y,1)$ if and only if $y - x \in B$. Since A and C are symmetric and avoid 0, Γ is a simple graph on $4n-2$ vertices. A vertex $(x,0)$ has $|A|+|B| = 2n-1$ neighbours and a vertex $(x,1)$ has $|C|+|B| = 2n-3$; the translations $(x,i) \mapsto (x+t,i)$ are automorphisms. Write $\lambda(p,q)$ for the number of common neighbours of two vertices, $\mu(p,q)$ for the number of vertices other than p and q adjacent to neither, $[\,P\,]$ for the indicator of a statement P, and

$$f(d) = \mathrm{N}_{A,A}(d) + \mathrm{N}_{B,B}(d) \qquad (d \in G).$$

Lemma 2.1. *For $d \neq 0$,*

$$\lambda\big((0,0),(d,0)\big) = f(d), \qquad \lambda\big((0,1),(d,1)\big) = f(d) - 3 + 2[\,d \in A\,],$$

and for every $d \in G$, $\lambda\big((0,0),(d,1)\big) = n - 1 - [\,d \in B\,]$.

Proof. A vertex $(z,0)$ is a common neighbour of $(0,0)$ and $(d,0)$ when $z \in A$ and $z-d \in A$, and $(z,1)$ is one when $z \in B$ and $z-d \in B$; this gives $\mathrm{N}_{A,A}(d) + \mathrm{N}_{B,B}(d)$.

For $(0,1)$ and $(d,1)$, the common neighbours $(z,1)$ are counted by $\mathrm{N}_{C,C}(d)$. A vertex $(z,0)$ is adjacent to $(0,1)$ when $-z \in B$ and to $(d,1)$ when $d-z \in B$; with $w = -z$ these say $w \in B$ and $w+d \in B$, so there are $\mathrm{N}_{B,B}(-d) = \mathrm{N}_{B,B}(d)$ of them. To compute $\mathrm{N}_{C,C}(d)$, let $S = A \cup \{0\}$, so that $|S| = n+1$ and $C = G \setminus S$. Then $|C \cap (C+d)| = v - |S \cup (S+d)| = v - 2(n+1) + |S \cap (S+d)|$. The set $S \cap (S+d)$ is the disjoint union of $A \cap (A+d)$, of $\{0\}$ when $-d \in A$, and of $\{d\}$ when $d \in A$; neither 0 nor d lies in $A \cap (A+d)$, because $0 \notin A$. As $A = -A$ and $v - 2(n+1) = -3$, we get $\mathrm{N}_{C,C}(d) = \mathrm{N}_{A,A}(d) - 3 + 2[\,d \in A\,]$.

For $(0,0)$ and $(d,1)$, a vertex $(z,0)$ is a common neighbour when $z \in A$ and $d - z \in B$, that is, $z = d - b$ with $b \in B$ and $d - b \in A$; since $A = -A$ the last condition reads $b - d \in A$. A vertex $(z,1)$ is a common neighbour when $z \in B$ and $z - d \in C$. Together, $\lambda = \#\{b \in B : \ b - d \in A \cup C\} = \#\{b \in B : \ b \neq d\} = n - 1 - [\,d \in B\,]$. □

Lemma 2.2. *Let p, q be distinct non-adjacent vertices of Γ. Then $\mu(p,q) = \lambda(p,q) - 2$ if both lie in $G \times \{0\}$, $\mu(p,q) = \lambda(p,q) + 2$ if both lie in $G \times \{1\}$, and $\mu(p,q) = \lambda(p,q)$ otherwise.*

Proof. The neighbourhoods of p and q lie among the other $4n-4$ vertices, and their union has $\deg p + \deg q - \lambda(p,q)$ elements, so $\mu(p,q) = 4n - 4 - \deg p - \deg q + \lambda(p,q)$. Insert the degrees $2n-1$ and $2n-3$. □

Proposition 2.3. *$\Gamma(A,B)$ contains no B_{n-1} and its complement contains no B_n if and only if*

$$f(d) \leq n + 1 - 3\,[\,d \in A\,] \qquad \text{for every } d \in G \setminus \{0\}. \tag{2.1}$$

In that case $R(B_{n-1}, B_n) = 4n-1$.

Proof. An edge lying in k triangles is the spine of a B_k, and a non-edge pq is the spine of a B_k in the complement exactly when $\mu(p,q) \geq k$. So the two conditions say that $\lambda(p,q) \leq n-2$ for

every edge and $\mu(p,q) \le n-1$ for every pair of distinct non-adjacent vertices. By translation it suffices to take $p = (0,i)$, and we use Lemmas 2.1 and 2.2.

For $p = (0,0)$ and $q = (d,0)$ with $d \neq 0$, the pair is an edge if and only if $d \in A$. For $d \in A$ the condition is $f(d) \le n-2$, and for $d \notin A$ it is $f(d) - 2 \le n-1$; together they are (2.1). For $p = (0,1)$ and $q = (d,1)$, the pair is an edge if and only if $d \in C$, that is, $d \notin A$. For $d \notin A$ the condition is $f(d) - 3 \le n-2$, and for $d \in A$ it is $(f(d)-1)+2 \le n-1$; together they are again (2.1). For $p = (0,0)$ and $q = (d,1)$ the pair is an edge if and only if $d \in B$; then $\lambda = n-2$, and otherwise $\mu = \lambda = n-1$, so the condition holds with equality in both cases. The pairs $(0,1), (d,0)$ are translates of these. The last sentence follows from (1.1), since Γ has $4n-2$ vertices. □

Remark 2.4. Summing over $d \neq 0$ gives $\sum f(d) = n(n-1) + (n-1)(n-2) = 2(n-1)^2$ and $\sum (n+1-3[\,d \in A\,]) = (n+1)(2n-2) - 3n = (n-2)(2n+1)$. So when (2.1) holds, the slacks $s(d) = n+1-3[\,d \in A\,] - f(d)$ are non-negative integers with sum $n-4$, and (2.1) is an equality for at least $n+2$ of the $2n-2$ values of d. For the regular graphs used when n is odd [2, 12] the corresponding sum is $n-1$, so the even case is tighter. The cross pairs never matter: B enters only through its autocorrelation $\mathrm{N}_{B,B}$. Since $f(-d) = f(d)$ and $[\,-d \in A\,] = [\,d \in A\,]$, it suffices to check (2.1) for $1 \le d \le n-1$.

3. The case $n = 58$

Let $n = 58$ and $v = 115 = 5 \cdot 23$. Let $A = \pm A_+$ and let B be the sets

$$\begin{aligned} A_+ = \{&3, 10, 11, 14, 16, 18, 21, 23, 24, 25, 26, 27, 28, 29, 30, 32, 36, 37, 38, 43, 44, 46, 47,\\ &48, 49, 52, 54, 55, 56\},\\ B = \{&0, 2, 5, 6, 7, 9, 10, 11, 15, 17, 20, 21, 22, 23, 24, 26, 30, 34, 38, 43, 44, 46, 47, 50, 52,\\ &53, 54, 56, 59, 65, 66, 67, 69, 71, 72, 73, 74, 75, 80, 81, 83, 84, 85, 88, 90, 92, 93, 97,\\ &100, 102, 105, 106, 107, 109, 110, 111, 114\}. \end{aligned}$$

The 29 elements of A_+ lie in $\{1, \dots, 57\}$, so $A = A_+ \cup (-A_+)$ has 58 elements, avoids 0 and satisfies $A = -A$; and $|B| = 57$. Written out,

$$\begin{aligned} A = \{&3, 10, 11, 14, 16, 18, 21, 23, 24, 25, 26, 27, 28, 29, 30, 32, 36, 37, 38, 43, 44, 46, 47,\\ &48, 49, 52, 54, 55, 56, 59, 60, 61, 63, 66, 67, 68, 69, 71, 72, 77, 78, 79, 83, 85, 86, 87,\\ &88, 89, 90, 91, 92, 94, 97, 99, 101, 104, 105, 112\}. \end{aligned}$$

Figure 1 draws both sets on the torus $\mathbb{Z}_5 \times \mathbb{Z}_{23}$.

Theorem 3.1. *The sets A and B above satisfy (2.1) for $n = 58$. Consequently $\Gamma(A,B)$ is a graph on 230 vertices, with degrees 115 and 113, that contains no B_{57} and whose complement contains no B_{58}, and $R(B_{57}, B_{58}) = 231$.*

Proof. By Remark 2.4 it suffices to check (2.1) for $1 \le d \le 57$, and Table 1 lists $\mathrm{N}_{A,A}(d)$ and $\mathrm{N}_{B,B}(d)$ there; each entry is the number of elements of A, or of B, whose translate by d stays in the set.[2] In every row $f(d) \le 59 - 3[\,d \in A\,]$. Proposition 2.3 gives the rest. □

[2]For instance $\mathrm{N}_{A,A}(1) = 32$ is the number of pairs $a, a+1$ of consecutive residues in A: the runs $10, 11$; $23, \dots, 30$; $36, 37, 38$; $43, 44$; $46, \dots, 49$; $54, 55, 56$; $59, 60, 61$; $66, \dots, 69$; $71, 72$; $77, 78, 79$; $85, \dots, 92$; $104, 105$ contribute $1+7+2+1+3+2+2+3+1+2+7+1 = 32$, and neither 0 nor 114 lies in A.




FIGURE 1. The sets A and B for $n = 58$. The residue $x \in \mathbb{Z}_{115}$ sits at angle $2\pi x/5$ around the tube and $2\pi x/23$ around the axis.




FIGURE 2. The slack $s(d) = 59 - 3[\,d \in A\,] - f(d)$ for $n = 58$ at every $d \neq 0$, with d placed by its residues modulo 5 and 23; the black cell is $d = 0$, and a dot marks $d \in A$. Equality holds for 70 values of d, the slack is 1 for 34 and 2 for 10.

4. REMARKS

Remark 4.1. The function f is not constant on A or on C (Figure 2): on the 58 elements of A it takes the values 54, 55 and 56, for 4, 18 and 36 of them, and on the 56 elements of C the values 57, 58 and 59, for 6, 16 and 34 of them. So the example is not a difference-set construction of the kind behind the infinite families in [10, 11, 12].

Remark 4.2. Unlike the graph for $n = 67$ in [2], this one has no multiplier symmetry beyond $d \mapsto -d$: the only units u of $\mathbb{Z}_{115}$ with $uA = A$ are ± 1, the only one with $uB = B$ is 1, and B meets $-B$ in 29 elements. So the 57 rows of Table 1 cannot be grouped into orbits as in [2].

Remark 4.3. We found the example by a computer search following the two-phase scheme of Turturean [11]. The first phase chooses A so that the bounds $\mathrm{N}_{B,B}(d) \leq n + 1 - 3[\,d \in A\,] - \mathrm{N}_{A,A}(d)$ leave a large margin in Fourier space, where they become linear conditions on the power spectrum of B. The second phase fixes A and searches for B by local search over exchanges of one element of B with one outside it, guided by weights on the inequalities (2.1); when no exchange lowers the weighted excess, the weights of the violated inequalities are raised, as in the breakout method of Morris [8]. No step of the proof above depends on the search.

TABLE 1. The inequality (2.1) for $n = 58$ at $1 \le d \le 57$, with $b(d) = 59 - 3[\,d \in A\,]$ and the slack $s(d) = b(d) - f(d)$ of Remark 2.4. The slacks add up to 27, half of $n - 4 = 54$.

d	$[\,d \in A\,]$	$\mathrm{N}_{A,A}$	$\mathrm{N}_{B,B}$	f	b	s	d	$[\,d \in A\,]$	$\mathrm{N}_{A,A}$	$\mathrm{N}_{B,B}$	f	b	s
1	0	32	27	59	59	0	30	1	26	28	54	56	2
2	0	32	27	59	59	0	31	0	30	29	59	59	0
3	1	29	27	56	56	0	32	1	27	28	55	56	1
4	0	28	30	58	59	1	33	0	30	28	58	59	1
5	0	31	27	58	59	1	34	0	30	27	57	59	2
6	0	31	27	58	59	1	35	0	30	27	57	59	2
7	0	33	26	59	59	0	36	1	27	29	56	56	0
8	0	32	27	59	59	0	37	1	26	29	55	56	1
9	0	28	31	59	59	0	38	1	28	27	55	56	1
10	1	26	29	55	56	1	39	0	31	28	59	59	0
11	1	35	21	56	56	0	40	0	32	26	58	59	1
12	0	30	29	59	59	0	41	0	31	28	59	59	0
13	0	30	29	59	59	0	42	0	31	27	58	59	1
14	1	30	26	56	56	0	43	1	29	27	56	56	0
15	0	26	32	58	59	1	44	1	26	28	54	56	2
16	1	28	27	55	56	1	45	0	30	28	58	59	1
17	0	29	30	59	59	0	46	1	29	27	56	56	0
18	1	30	26	56	56	0	47	1	26	29	55	56	1
19	0	29	30	59	59	0	48	1	27	28	55	56	1
20	0	31	26	57	59	2	49	1	28	28	56	56	0
21	1	25	31	56	56	0	50	0	29	30	59	59	0
22	0	31	28	59	59	0	51	0	29	30	59	59	0
23	1	29	27	56	56	0	52	1	27	29	56	56	0
24	1	28	28	56	56	0	53	0	32	27	59	59	0
25	1	28	28	56	56	0	54	1	27	29	56	56	0
26	1	28	27	55	56	1	55	1	27	28	55	56	1
27	1	27	29	56	56	0	56	1	27	29	56	56	0
28	1	27	29	56	56	0	57	0	31	28	59	59	0
29	1	27	29	56	56	0							

Question 4.4. The smallest open case of $R(B_{n-1}, B_n) = 4n - 1$ is now $n = 59$, with $v = 117$; there n is odd, and the regular two-block circulant graphs of [2, 12] are available. Is there a two-block circulant graph over $\mathbb{Z}_{117}$, or over $\mathbb{Z}_3^2 \times \mathbb{Z}_{13}$, that settles it?

Formal verification. Theorem 3.1 is also proved in Lean 4 [7], without any library beyond the Lean core. The development defines the graph $\Gamma(A, B)$ on the vertices $0, \ldots, 229$ by the two bit masks of A and B, and the kernel evaluates the common neighbours and common non-neighbours of all 26,335 pairs of vertices and checks that the adjacency is symmetric. The final theorem states that every edge has at most 56 common neighbours and every pair of distinct non-adjacent vertices has at most 57 common non-neighbours, and it depends only on the standard axioms `propext`, `Classical.choice` and `Quot.sound`. The graph is also checked by two independent programs, in Python and in C, which read the sets A and B and test all pairs directly. The development, the programs and the search are archived in [1].

Use of generative AI. The author used Claude (Anthropic) to search for the example, to check it by computer and in Lean, and to draft the text; the author verified every step and is responsible for the content.

References

[1] D. Bhattacharjee, *Book Ramsey numbers $R(B_{57}, B_{58}) = 231$ and $R(B_{66}, B_{67}) = 267$: data, checks, search and Lean 4 proofs*, Zenodo, 2026, all versions doi:10.5281/zenodo.23373628. Source code at https://github.com/creelie/Breakthrough-problems-2

[2] D. Bhattacharjee, *A hand-checkable proof that $R(B_{66}, B_{67}) = 267$*, preprint, 2026, archived in [1].

[3] S. Chung, W. Du and W. J. Wesley, *Autonomous mathematical discovery in an open-world multi-agent environment*, preprint, 2026. arXiv:2608.23691

[4] S. Chung, W. Du and W. J. Wesley, *Station data v2*, book Ramsey verification notebook and certificates, first public version 24 August 2026. https://github.com/dualverse-ai/station_data_v2, accessed 14 October 2026.

[5] Epoch AI, *FrontierMath open problems: Ramsey numbers for book graphs*, 2026. https://epoch.ai/frontiermath/open-problems/ramsey-book-graphs, accessed 15 October 2026.

[6] A. W. Goodman, *On sets of acquaintances and strangers at any party*, Amer. Math. Monthly **66** (1959), no. 9, 778–783. doi:10.1080/00029890.1959.11989408

[7] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in: Automated Deduction – CADE 28, Lecture Notes in Comput. Sci. **12699**, Springer, Cham, 2021, 625–635. doi:10.1007/978-3-030-79876-5_37

[8] P. Morris, *The breakout method for escaping from local minima*, in: Proceedings of the Eleventh National Conference on Artificial Intelligence (AAAI-93), AAAI Press, 1993, 40–45. https://cdn.aaai.org/AAAI/1993/AAAI93-007.pdf

[9] C. C. Rousseau and J. Sheehan, *On Ramsey numbers for books*, J. Graph Theory **2** (1978), 77–87. doi:10.1002/jgt.3190020110

[10] G. Szekeres, *Tournaments and Hadamard matrices*, Enseign. Math. (2) **15** (1969), 269–278.

[11] D. Turturean, *Summary of new results on the Ramsey numbers for book graphs open problem*, 2026. https://docs.google.com/document/d/1VinXOiMov2v-Y27GTJwoAECywv2a34MH4E8qa712zJY

[12] W. J. Wesley, *Lower bounds for book Ramsey numbers*, Discrete Math. **349** (2026), no. 5, 114913. doi:10.1016/j.disc.2025.114913

Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL),
Bhubaneswar, Odisha 751030, India

Email address: d.bhattacharjee@erl-forschung.de

Email address: itsdeep@live.com

URL: https://orcid.org/0000-0003-0466-750X