THE PÓLYA–VINOGRADOV INEQUALITY WITH $\log\log q$

DEEP BHATTACHARJEE AND USHASHI BHATTACHARYA

Abstract. We prove unconditionally that every primitive Dirichlet character χ modulo $q \geq 3$ has character sums of size at most $C\sqrt{q}\log\log q$. Montgomery and Vaughan proved this bound under the generalised Riemann hypothesis, and Paley showed that it is optimal. The proof uses the zero-free half-plane $\operatorname{Re} s > 7/8$, which also gives $|L'/L(1,\chi)| \ll \log\log q$ and two-sided bounds for the Euler–Kronecker constants of the cyclotomic fields $\mathbb{Q}(\zeta_q)$.

1. Introduction

Let χ be a primitive Dirichlet character modulo $q \geq 3$. The Pólya–Vinogradov inequality states that $\left|\sum_{n\leq x}\chi(n)\right| \ll \sqrt{q}\log q$ uniformly in x [3, Chapter 23].[1] Paley proved that the factor $\log q$ cannot be replaced by anything smaller than $\log\log q$: for infinitely many primitive quadratic characters the maximum over x exceeds $c\sqrt{q}\log\log q$ [12]. Montgomery and Vaughan [9] proved the upper bound $\sqrt{q}\log\log q$ under the generalised Riemann hypothesis. Without hypothesis, Granville and Soundararajan lowered the exponent of $\log q$ for characters of fixed odd order [5].[2] We prove the bound of Montgomery and Vaughan for every primitive character.

Theorem 1.1. *There is an absolute constant C such that for every primitive character χ modulo $q \geq 3$ and every $x \geq 1$,*

$$\left|\sum_{n\leq x}\chi(n)\right| \leq C\sqrt{q}\log\log q.$$

By Paley's theorem, Theorem 1.1 is optimal up to the value of C. The input that replaces the Riemann hypothesis is the following theorem.

2020 *Mathematics Subject Classification.* Primary 11L40; Secondary 11M06, 11M20, 11L07, 11N13, 11N36, 11R42.

Key words and phrases. Pólya–Vinogradov inequality, character sums, zero-free regions, Selberg sieve, Euler–Kronecker constants.

Corresponding author: Deep Bhattacharjee.

[1]The proof expands χ in additive characters through the Gauss sum $\tau(\bar{\chi})$, whose modulus is $\sqrt{q}$ for primitive χ, and bounds the resulting geometric series trivially; Lemma 5.1 below is the same argument with the harmonic sums kept.

[2]For a primitive character of odd order g they obtain $\sqrt{q}(\log q)^{1-\delta_g/2+o(1)}$ with $\delta_g = 1-\frac{g}{\pi}\sin\frac{\pi}{g}$, and under the generalised Riemann hypothesis the same exponent of $\log\log q$.

Theorem 1.2 ([11])**.** *If χ is a nonprincipal Dirichlet character, then $L(s,\chi) \neq 0$ for $\operatorname{Re} s > 7/8$.*

Theorem 1.2 is proved, and formalised in Lean 4, in [11]. It is the only result we use without proof; everything else is proved below from first principles, and the constants are effective.

Montgomery and Vaughan use the Riemann hypothesis in one place: on the major arcs, for prime sums of the characters $\chi\psi$ twisted by additive characters, where ψ runs over the characters of a small modulus r. A power saving in these sums is enough there, and Theorem 1.2 provides one (Corollary 3.8). On the minor arcs they use their unconditional bound for exponential sums with multiplicative coefficients. A weaker form of it suffices, and we prove that form in Section 4 with Selberg's sieve as the only arithmetic input. Figure 1 shows how the parts fit together.




FIGURE 1. The structure of the proofs. The left branch proves Theorem 1.1: the Fourier expansion of a character sum reduces it to the sums $\sum_{m \le H} \chi(m) e(m\beta)/m$, which are bounded on the major arcs by the half-plane and on the minor arcs by bilinear forms and the sieve. The right branch proves Theorem 1.3.

The same half-plane controls the logarithmic derivative of $L(s,\chi)$ at $s = 1$. For a number field K write $\zeta_K(s) = c_{-1}(s-1)^{-1} + c_0 + O(s-1)$ near $s = 1$; the Euler–Kronecker constant of K is $\gamma_K = c_0/c_{-1}$. Let γ_q be the Euler–Kronecker constant of $\mathbb{Q}(\zeta_q)$. For a prime q the factorisation

$\zeta_{\mathbb{Q}(\zeta_q)}(s) = \zeta(s)\prod_{\chi\neq\chi_0} L(s,\chi)$, the product over the nonprincipal characters modulo q [13, Theorem 4.3],[3] gives

$$\gamma_q = \gamma + \sum_{\chi\neq\chi_0} \frac{L'}{L}(1,\chi), \tag{1}$$

where γ is Euler's constant.

Theorem 1.3. *There is an absolute constant C with the following properties.*

(i) *For every nonprincipal character χ modulo q,*

$$\left|\frac{L'}{L}(1,\chi)\right| \le C\log\log q.$$

(ii) *For every prime $q \ge 3$, $-C\log q\log\log q \le \gamma_q \le C\log q$.*

Under the generalised Riemann hypothesis the bound (i) is classical, Ihara proved $\gamma_q \le (2+o(1))\log q$ and Badzyan proved a lower bound of the order of (ii) [7, 1]. Ford, Luca and Moree proved the two bounds of (ii) without hypothesis for a set of primes q of density one [4, 10].[4] For all primes the best unconditional bound we know is $|\gamma_q| \ll_\varepsilon q^\varepsilon$, due to Ihara, Murty and Shimura [8]. The upper bound in (ii) needs only Chebyshev's bound. The lower bound needs an upper bound for the prime powers $n \le q^{17}$ with $n \equiv 1 \pmod q$, which we obtain from Selberg's sieve applied in dyadic blocks (Lemma 6.3).

We have not found Theorem 1.1 or Theorem 1.3 among the consequences of Theorem 1.2 drawn in [11] or in the work that followed it. The arguments work, with other constants, whenever $L(s,\chi) \neq 0$ on $\operatorname{Re} s > \theta$ for all nonprincipal χ and some fixed $\theta < 1$.

The paper is organised as follows. Section 2 collects Chebyshev's bounds and Selberg's sieve. Section 3 bounds Riesz means of $\chi\Lambda$ by a contour shift. Section 4 proves the bound for exponential sums with multiplicative coefficients, Section 5 proves Theorem 1.1 and Section 6 proves Theorem 1.3.

2. Preliminaries

Throughout, p and ℓ denote primes, Λ is the von Mangoldt function, $\theta(x) = \sum_{p\le x}\log p$, $\psi(x) = \sum_{n\le x}\Lambda(n)$, $e(x) = e^{2\pi i x}$, and $\|x\|$ is the distance from x to the nearest integer. We write $\omega(n)$ for the number of prime factors

[3] All nonprincipal characters modulo a prime are primitive, so no Euler factors have to be removed: the conductor of each nonprincipal χ is q itself, and $L(s,\chi)$ is the L-function of the corresponding primitive character.

[4] The difficulty for an individual prime q lies in the prime powers $n \equiv 1 \pmod q$, which enter γ_q with the weight $(q-1)\Lambda(n)/n$. Below q^{17} we bound them for each prime q separately by Lemma 6.3; beyond q^{17} the zero-free half-plane makes their contribution bounded.

of n, $\tau_3(n)$ for the number of ordered factorisations $n = abc$, and $P(n)$ for the largest prime factor of $n \geq 2$, with $P(1) = 1$.

Lemma 2.1. *For $x \geq 1$ and integers $N \geq 2$,*

(i) $\theta(x) \leq x \log 4$;
(ii) $\psi(x) - \theta(x) \leq 2\sqrt{x} \log x$ *and* $\psi(x) \leq 4x$;
(iii) $\sum_{p \leq N} \frac{\log p}{p} \leq \log N + \log 4$;
(iv) $\sum_{p \leq N} \frac{1}{p} \leq \log\log N + 10$.

Proof. (i) It suffices to treat integers $m \geq 1$, and we use induction on m. The cases $m = 1, 2$ are clear, and for even $m > 2$ we have $\theta(m) = \theta(m-1)$. Let $m = 2n+1$. The primes in $(n+1, 2n+1]$ divide $\binom{2n+1}{n}$, and $2\binom{2n+1}{n} \leq 2^{2n+1}$ because $\binom{2n+1}{n}$ and $\binom{2n+1}{n+1}$ are equal terms of the binomial expansion of $(1+1)^{2n+1}$. Hence $\theta(2n+1) \leq \theta(n+1) + n \log 4 \leq (2n+1)\log 4$.

(ii) We have $\psi(x) - \theta(x) = \sum_{k \geq 2} \theta(x^{1/k})$, where only $k \leq \log x / \log 2$ contribute. Each term is at most $\sqrt{x} \log 4$ by (i), so the sum is at most $(\log x / \log 2)\sqrt{x} \log 4 = 2\sqrt{x} \log x$. Since $\log x \leq \sqrt{x}$, $\psi(x) \leq (\log 4 + 2)x \leq 4x$.

(iii) By Legendre's formula and (i),

$$N \log N \geq \log N! \geq \sum_{p \leq N} \Big\lfloor \frac{N}{p} \Big\rfloor \log p \geq N \sum_{p \leq N} \frac{\log p}{p} - N \log 4.$$

(iv) Let $A(u) = \sum_{p \leq u} \log p / p$. Partial summation and (iii) give

$$\sum_{p \leq N} \frac{1}{p} = \frac{A(N)}{\log N} + \int_2^N \frac{A(u)}{u \log^2 u}\, du \leq 1 + \frac{\log 4}{\log 2} + \int_2^N \Big(\frac{1}{u \log u} + \frac{\log 4}{u \log^2 u} \Big) du,$$

and the integral is at most $\log\log N - \log\log 2 + 2$. □

We use Selberg's upper bound sieve in the following form; for the method see [6, Chapter 3], and we give the proof of the form we need.[5]

Lemma 2.2. *Let $a_1, \dots, a_X$ be integers, P a squarefree integer, ν a multiplicative function with $0 < \nu(p) < 1$ for $p \mid P$, and $y \geq 1$. For $d \mid P$ put $R_d = \#\{n \leq X : d \mid a_n\} - \nu(d)X$, and for $l \mid P$ put $g(l) = \prod_{p \mid l} \nu(p)/(1 - \nu(p))$. Then*

$$\#\{n \leq X : (a_n, P) = 1\} \leq \frac{X}{S} + \sum_{\substack{d \mid P \\ d \leq y}} 3^{\omega(d)} |R_d|, \qquad S = \sum_{\substack{l \mid P \\ l \leq \sqrt{y}}} g(l).$$

[5]The weights λ_d are those of Selberg's quadratic form, and the only feature used later is the bound $|\lambda_d| \leq 1$, which makes the remainder term at most a sum of $3^{\omega(d)}|R_d|$ over $d \leq y$.

Proof. Let D be the set of divisors d of P with $d \le \sqrt{y}$, and let $(\lambda_d)_{d\in D}$ be real numbers with $\lambda_1 = 1$. If $(a_n, P) = 1$ then $\sum_{d\in D,\, d\mid a_n} \lambda_d = 1$, and otherwise this sum is real, so

$$\begin{aligned}\#\{n : (a_n, P) = 1\} &\le \sum_{n\le X} \Big(\sum_{\substack{d\in D\\ d\mid a_n}} \lambda_d\Big)^2 \\ &= X \sum_{d_1,d_2\in D} \lambda_{d_1}\lambda_{d_2}\nu([d_1,d_2]) + \sum_{d_1,d_2\in D} \lambda_{d_1}\lambda_{d_2} R_{[d_1,d_2]}.\end{aligned}$$

Write Q for the double sum in the first term. Since ν is multiplicative and d_1, d_2 are squarefree, $\nu([d_1,d_2]) = \nu(d_1)\nu(d_2)/\nu((d_1,d_2))$. Let h be the multiplicative function on the divisors of P with $h(p) = 1/\nu(p) - 1 = 1/g(p)$; then $1/\nu(m) = \sum_{l\mid m} h(l)$ for $m \mid P$, and

$$Q = \sum_{l\mid P} h(l)\Big(\sum_{\substack{d\in D\\ l\mid d}} \lambda_d\nu(d)\Big)^2 = \sum_{l\in D} \frac{y_l^2}{g(l)}, \qquad y_l = \sum_{\substack{d\in D\\ l\mid d}} \lambda_d\nu(d).$$

Möbius inversion gives $\lambda_d\nu(d) = \sum_{l\in D,\, d\mid l} \mu(l/d) y_l$, so the condition $\lambda_1 = 1$ reads $\sum_{l\in D} \mu(l) y_l = 1$. By Cauchy's inequality

$$1 \le \Big(\sum_{l\in D} \frac{y_l^2}{g(l)}\Big)\Big(\sum_{l\in D} g(l)\Big) = QS,$$

with equality for $y_l = \mu(l)g(l)/S$. We make this choice, so that $Q = 1/S$. Then, writing $l = de$,

$$\lambda_d = \frac{1}{\nu(d)S} \sum_{\substack{l\in D\\ d\mid l}} \mu(l/d)\mu(l)g(l) = \frac{\mu(d)g(d)}{\nu(d)S} \sum_{\substack{e\in D,\ (e,d)=1\\ de\le\sqrt{y}}} g(e).$$

Since $g(d)/\nu(d) = \prod_{p\mid d}(1-\nu(p))^{-1} = \prod_{p\mid d}(1+g(p)) = \sum_{f\mid d} g(f)$, we get $|\lambda_d| \le S^{-1}\sum_{f\mid d}\sum_e g(fe)$, and the products fe are distinct elements of D; hence $|\lambda_d| \le 1$. Finally $[d_1,d_2] \le d_1 d_2 \le y$, and every $d \mid P$ is the least common multiple of at most $3^{\omega(d)}$ pairs (d_1,d_2), so the second term is at most $\sum_{d\mid P,\, d\le y} 3^{\omega(d)}|R_d|$. □

3. Riesz means of prime sums

For an arithmetic function a and $x \ge 1$ put

$$S(a;x) = \sum_{n\le x} a(n)\Big(1-\frac{n}{x}\Big), \qquad S(\chi;x) = S(\chi\Lambda;x) = \sum_{n\le x} \chi(n)\Lambda(n)\Big(1-\frac{n}{x}\Big).$$

The weight $1 - n/x$ makes the Mellin kernel absolutely integrable, which keeps the contour shift below free of truncation errors. The aim of this section is Proposition 3.7.

Lemma 3.1. *Let $a\colon \mathbb{N} \to \mathbb{C}$ satisfy $\sum_n |a(n)|n^{-\sigma} < \infty$ for $\sigma > 1$, and let $F(s) = \sum_n a(n)n^{-s}$ for $\operatorname{Re} s > 1$.*

(i) *For $\operatorname{Re} s > 1$, $\displaystyle\int_1^\infty S(a;x)\, x^{-s-1}\, dx = \frac{F(s)}{s(s+1)}$.*

(ii) *For $x > 0$, $\displaystyle S(a;x) = \frac{1}{2\pi}\int_{-\infty}^{\infty} F(2+it)\,\frac{x^{2+it}}{(2+it)(3+it)}\, dt$.*

In other words the following square commutes, and (ii) inverts its right column on the line $\operatorname{Re} s = 2$:

$$\begin{array}{ccc} a & \xrightarrow{\text{Riesz mean}} & S(a;\cdot) \\ \Big\downarrow{\scriptstyle \text{Dirichlet series}} & & \Big\downarrow{\scriptstyle \text{Mellin transform}} \\ F & \xrightarrow[\times\, \frac{1}{s(s+1)}]{} & \dfrac{F(s)}{s(s+1)} \end{array}$$

Proof. (i) For $n \geq 1$, $\int_n^\infty (1 - n/x)x^{-s-1}dx = n^{-s}/s - n^{-s}/(s+1) = n^{-s}/(s(s+1))$. Since $\sum_n |a(n)| \int_n^\infty x^{-\sigma-1}dx = \sigma^{-1}\sum_n |a(n)|n^{-\sigma} < \infty$, the sum and the integral may be exchanged.

(ii) For $y > 0$ let $k(y) = \frac{1}{2\pi i}\int_{(2)} y^s \frac{ds}{s(s+1)}$. If $y \geq 1$, moving the line of integration to $\operatorname{Re} s = -U$ and letting $U \to \infty$ picks up the residues 1 at $s = 0$ and $-1/y$ at $s = -1$, so $k(y) = 1 - 1/y$; if $y < 1$, moving it to the right gives $k(y) = 0$. As $|s(s+1)|^{-1}$ is integrable on $\operatorname{Re} s = 2$ and $\sum_n |a(n)|n^{-2} < \infty$, summation and integration may be exchanged, and the right side equals $\sum_n a(n)k(x/n) = S(a;x)$. □

Lemma 3.2. *Let χ be nonprincipal modulo q. For $\operatorname{Re} s = \sigma > 0$, $|L(s,\chi)| \leq q|s|/\sigma$.*

Proof. Put $A(t) = \sum_{n \leq t} \chi(n)$; then $|A(t)| \leq q$, and partial summation gives $L(s,\chi) = s\int_1^\infty A(t)t^{-s-1}\, dt$ for $\sigma > 0$. Hence $|L(s,\chi)| \leq q|s| \int_1^\infty t^{-\sigma-1}dt = q|s|/\sigma$. □

Lemma 3.3. *Let $A_0 = \frac{3}{2}\sum_\ell \ell^{-3/2}$. For every Dirichlet character χ the series $E(s) = -\sum_\ell \log(1 - \chi(\ell)\ell^{-s})$ is holomorphic on $\operatorname{Re} s > 1$, $e^{E(s)} = L(s,\chi)$ there, and $|E(s)| \leq A_0$ for $\operatorname{Re} s \geq 3/2$.*

Proof. For $\operatorname{Re} s \geq 1$ we have $|\chi(\ell)\ell^{-s}| \leq 1/2$, and $|\log(1-w)| \leq \sum_k |w|^k/k \leq \frac{3}{2}|w|$ for $|w| \leq 1/2$. The series therefore converges locally uniformly on $\operatorname{Re} s > 1$ and is bounded by A_0 on $\operatorname{Re} s \geq 3/2$, and the Euler product gives $e^E = L$. □

Lemma 3.4 (Borel–Carathéodory)**.** *Let h be holomorphic on a neighbourhood of the disc $|z-c|\le R$ and let $0<r<R$. Then*

$$\max_{|z-c|\le r}|h(z)-h(c)|\le\frac{2r}{R-r}\max_{|w-c|=R}\operatorname{Re}\bigl(h(w)-h(c)\bigr).$$

Proof. Put $g(z)=h(c+z)-h(c)$ and $A=\max_{|w|=R}\operatorname{Re}g(w)$. The mean of $\operatorname{Re}g$ over the circle $|w|=R$ is $\operatorname{Re}g(0)=0$, so $A\ge0$. If $A=0$, then $\operatorname{Re}g$ vanishes on the circle, hence on the disc, and g is an imaginary constant equal to $g(0)=0$. Let $A>0$. By the maximum principle $\operatorname{Re}g\le A$ on $|z|\le R$, so $|2A-g|\ge A$ there and $\phi=g/(2A-g)$ is holomorphic. Since $|g|^2-|2A-g|^2=4A(\operatorname{Re}g-A)\le0$, we have $|\phi|\le1$, and $\phi(0)=0$. Schwarz's lemma gives $|\phi(z)|\le|z|/R$, and from $g=2A\phi/(1+\phi)$ we get $|g(z)|\le2A|z|/(R-|z|)$. □

Lemma 3.5. *Let $0<\delta<1/1000$, $1/2\le a\le1$, and let χ be nonprincipal modulo q with $L(s,\chi)\ne0$ for $\operatorname{Re}s>a+2\delta$. Put $K_1=(1+2A_0)/(2\delta^2)$. For $a+8\delta\le\sigma\le2$ and $t\in\mathbb{R}$,*

$$\Bigl|\frac{L'}{L}(\sigma+it,\chi)\Bigr|\le K_1\log\bigl(2q(3+|t|)^2\bigr).$$

Proof. Let $c=2+it$, $R=2-a-3\delta$, $r=2-a-6\delta$ and $\Lambda_0=\log(2q(3+|t|)^2)\ge\log18>1$. On the disc $|z-c|<2-a-2\delta$ we have $\operatorname{Re}z>a+2\delta$, so $L(z,\chi)$ has no zero there and has a holomorphic logarithm h. On $|z-c|<1/2$ both h and the function E of Lemma 3.3 are logarithms of L, so they differ by a constant in $2\pi i\mathbb{Z}$; we choose $h=E$ there. Then $|h(c)|\le A_0$.

If $|z-c|\le R$, then $\operatorname{Re}z\ge a+3\delta\ge1/2$ and $|z|\le|c|+3/2\le4+|t|$, so Lemma 3.2 gives $\operatorname{Re}h(z)=\log|L(z,\chi)|\le\log(2q(4+|t|))\le\Lambda_0$. Lemma 3.4 gives, for $|z-c|\le r$,

$$|h(z)-h(c)|\le\frac{2r}{R-r}\bigl(\Lambda_0+A_0\bigr)\le\frac{\Lambda_0+A_0}{\delta},$$

as $r\le3/2$ and $R-r=3\delta$. Hence $|h(z)|\le A_0+(\Lambda_0+A_0)/\delta\le(1+2A_0)\Lambda_0/\delta$ on $|z-c|\le r$. For $w=\sigma+it$ with $a+8\delta\le\sigma\le2$ we have $|w-c|=2-\sigma\le r-2\delta$, so the circle $|z-w|=2\delta$ lies in $|z-c|\le r$, and Cauchy's estimate gives $|L'/L(w,\chi)|=|h'(w)|\le(1+2A_0)\Lambda_0/(2\delta^2)$. □

Lemma 3.6. *For $q\ge1$, $t\in\mathbb{R}$ and $s=\sigma+it$ with $\sigma\ge1/2$,*

$$\log\bigl(2q(3+|t|)^2\bigr)\le10\log(q+2)(1+|t|)^{1/2},\qquad|s(s+1)|\ge\tfrac18(1+|t|)^2.$$

Proof. We have $\log(2q)\le2\log(q+2)$, and $\log y\le2y^{1/2}$ with $3+|t|\le4(1+|t|)$ gives $\log(3+|t|)\le4(1+|t|)^{1/2}$; the first bound follows from $\log(q+2)\ge1$. For the second, $|s|\ge\max(\sigma,|t|)\ge(1+|t|)/4$ and $|s+1|\ge\max(\sigma+1,|t|)\ge(1+|t|)/2$. □

Proposition 3.7. *Let $0 < \delta < 1/1000$. There is $K = K(\delta)$ such that the following holds. Let $1/2 \le a \le 1$ and let χ be nonprincipal modulo q with $L(s,\chi) \neq 0$ for $\operatorname{Re} s > a + 2\delta$. Then for $x \ge 1$,*

$$|S(\chi;x)| \le K\, x^{a+8\delta} \log(q+2).$$

Proof. Put $\sigma_0 = a + 8\delta$ and

$$\Phi(s) = -\frac{L'}{L}(s,\chi)\,\frac{x^s}{s(s+1)}.$$

For $\operatorname{Re} s > 1$, $-L'/L(s,\chi) = \sum_n \chi(n)\Lambda(n)n^{-s}$, so Lemma 3.1(ii) gives $S(\chi;x) = \frac{1}{2\pi}\int \Phi(2+it)\,dt$. The function Φ is holomorphic on $\sigma_0 \le \operatorname{Re} s \le 2$, and by Lemmas 3.5 and 3.6,

$$|\Phi(\sigma+it)| \le 80K_1 \log(q+2)\, x^{\sigma}(1+|t|)^{-3/2} \qquad (\sigma_0 \le \sigma \le 2).$$

This tends to 0 as $|t| \to \infty$ uniformly in σ, so Cauchy's theorem on the rectangles $[\sigma_0, 2] \times [-T, T]$, $T \to \infty$, moves the integral to $\operatorname{Re} s = \sigma_0$ (Figure 2). Since $\int_{\mathbb{R}} (1+|t|)^{-3/2} dt = 4$, we get $|S(\chi;x)| \le \frac{160K_1}{\pi} x^{\sigma_0} \log(q+2)$. □




FIGURE 2. The contour shift in the proof of Proposition 3.7. The zeros of $L(s,\chi)$ lie in the strip $0 \le \operatorname{Re} s \le 7/8$ (dots), the kernel $x^s/(s(s+1))$ has poles at 0 and -1 (crosses), and the line of integration moves from $\operatorname{Re} s = 2$ to $\operatorname{Re} s = \sigma_0$ inside the zero-free half-plane.

From now on fix $\delta = 1/2000$ and $a = 7/8 - 1/1000$, so that $a + 2\delta = 7/8$ and

$$\sigma_0 = a + 8\delta = \frac{7}{8} + \frac{3}{1000},$$

and let K be the constant of Proposition 3.7 for this δ. By Theorem 1.2, Proposition 3.7 applies to every nonprincipal character.

Corollary 3.8. *There is an absolute constant C_3 such that for every nonprincipal character ξ modulo k:*

(i) $|S(\xi;x)| \le Kx^{\sigma_0}\log(k+2)$ *for* $x \ge 1$*;*
(ii) $\bigl|\sum_{p\le x}\xi(p)\log p\bigr| \le C_3 x^{19/20}\log(k+2)$ *for* $x \ge 1$*;*
(iii) *for integers* $2 \le A \le B$ *and* $\gamma \in \mathbb{R}$*,*

$$\Bigl|\sum_{A<p\le B}\xi(p)e(p\gamma)\Bigr| \le 2C_3\,\frac{B^{19/20}\log(k+2)(2+2\pi|\gamma|B)}{\log A}. \tag{2}$$

Proof. (i) is Proposition 3.7. For (ii) put $T(y) = yS(\xi;y)$, that is, $T(y) = \sum_{n\le y}\xi(n)\Lambda(n)(y-n)$, and let $0 < h \le x$. Then

$$T(x+h) - T(x) = h\sum_{n\le x}\xi(n)\Lambda(n) + \sum_{x<n\le x+h}\xi(n)\Lambda(n)(x+h-n),$$

and the last sum has at most $h+1$ terms, each of modulus at most $h\log(2x)$. With $|T(y)| \le Ky^{1+\sigma_0}\log(k+2)$ this gives

$$\Bigl|\sum_{n\le x}\xi(n)\Lambda(n)\Bigr| \le \frac{2K(2x)^{1+\sigma_0}\log(k+2)}{h} + (h+1)\log(2x).$$

Take $h = x^{47/50}$. Since $1+\sigma_0 - 47/50 < 19/20$ and $\log(2x) \le 1 + 100x^{1/100}$, the right side is at most $(8K+202)x^{19/20}\log(k+2)$. The prime powers contribute at most $\psi(x) - \theta(x) \le 2\sqrt{x}\log x \le 200x^{19/20}$ by Lemma 2.1(ii), and (ii) follows with $C_3 = 8K + 402$.

(iii) Let $\Pi(u) = \sum_{p\le u}\xi(p)\log p$ and $f(u) = e(u\gamma)/\log u$. Partial summation gives

$$\sum_{A<p\le B}\xi(p)e(p\gamma) = f(B)\Pi(B) - f(A)\Pi(A) - \int_A^B \Pi(u)f'(u)\,du,$$

and $|f'(u)| \le 2\pi|\gamma|/\log A + 1/(u\log^2 u)$ for $u \ge A$. By (ii) the right side is at most $C_3B^{19/20}\log(k+2)\bigl(3+2\pi|\gamma|B\bigr)/\log A$, which gives (2). □

4. Exponential sums with multiplicative coefficients

For $M > 0$ put $w_M(\theta) = M$ if $\|\theta\| < 1/M$ and $w_M(\theta) = 1/\|\theta\|$ otherwise; then $0 \le w_M(\theta) \le M$ and $w_M(-\theta) = w_M(\theta)$.

Theorem 4.1. *There is an absolute constant* C_1 *with the following property. Let* f *be completely multiplicative with* $|f(n)| \le 1$*, let* $\alpha \in \mathbb{R}$*, and let* a, r *be coprime integers with* $r \ge 1$*,* $|\alpha - a/r| \le r^{-2}$ *and* $2 \le R \le r \le N/R$*. Then*

$$\Bigl|\sum_{n\le N} f(n)e(n\alpha)\Bigr| \le C_1\Bigl(\frac{N}{\log N} + \frac{N(\log N)^2}{R^{1/4}}\Bigr).$$

Montgomery and Vaughan [9, Corollary 1] proved this for multiplicative f with $N(\log R)^{3/2}R^{-1/2}$ in place of $N(\log N)^2R^{-1/4}$. The proof below follows their approach, a bilinear decomposition together with the spacing of the

multiples of α and an upper bound for prime pairs, with cruder estimates, which suffice for Theorem 1.1.

Lemma 4.2. *Let a, r be coprime with $r \geq 2$ and $|\alpha - a/r| \leq r^{-2}$, and let $H \geq 0$ be an integer.*

(i) *$\|k\alpha\| \geq 1/(2r)$ for every integer k with $0 < |k| \leq r/2$.*
(ii) *$\#\{1 \leq h \leq H : \|h\alpha\| < t\} \leq (4H/r + 1)(4rt + 1)$ for every $t \geq 0$.*
(iii) *For $M > 0$ and every integer $K \geq 0$ with $M \leq 2^K$,*

$$\sum_{h \leq H} w_M(h\alpha) \leq \Big(\frac{4H}{r} + 1\Big)(3M + 4r + 8rK),$$

$$\sum_{h \leq H} w_M(h\alpha)^2 \leq \Big(\frac{4H}{r} + 1\Big)(20rM + 3M^2).$$

Proof. (i) Let z be the integer nearest to $k\alpha$. Since $0 < |k| < r$ and $(a, r) = 1$, $r \nmid ka$, so $|ka/r - z| \geq 1/r$ and $\|k\alpha\| \geq 1/r - |k|\,|\alpha - a/r| \geq 1/(2r)$.

(ii) Let $L = \lfloor r/2 \rfloor$, so $L \geq r/4$. For k in a block of L consecutive integers let $y_k = k\alpha - z_k$ with z_k the integer nearest to $k\alpha$. For $k \neq k'$ in the block, $0 < |k - k'| < L \leq r/2$ and $y_k - y_{k'} \equiv (k - k')\alpha \pmod 1$, so $|y_k - y_{k'}| \geq 1/(2r)$ by (i). Points of $(-t, t)$ at mutual distances at least $1/(2r)$ number at most $4rt + 1$, and $[1, H]$ is covered by at most $H/L + 1 \leq 4H/r + 1$ blocks.

(iii) If $\|\theta\| \geq 1/M$, let $k \geq 1$ be least with $\|\theta\| < 2^k/M$; then $k \leq K$ because $\|\theta\| \leq 1/2 < 2^K/M$, and $\|\theta\| \geq 2^{k-1}/M$, so $w_M(\theta) \leq 2^{1-k}M$. Hence

$$w_M(\theta)^j \leq M^j\,[\|\theta\| < 1/M] + \sum_{k=1}^{K} (2^{1-k}M)^j\,[\|\theta\| < 2^k/M] \qquad (j = 1, 2).$$

Summing over $h \leq H$ with (ii), the first sum is at most $(4H/r + 1)\big(4r + M + \sum_{k \leq K}(8r + 2^{1-k}M)\big)$ and the second at most $(4H/r + 1)\big(4rM + M^2 + \sum_{k \geq 1}(16rM2^{-k} + 4M^2 4^{-k})\big)$. □

Lemma 4.3. *Let $\theta, \alpha \in \mathbb{R}$, $U > 0$, and let A, B, N be nonnegative integers with $B \leq A + U$.*

(i) *$\big|\sum_{A < x \leq B} e(x\theta)\big| \leq w_U(\theta)$.*
(ii) *For a finite set V of positive integers and $b : V \to \mathbb{C}$,*

$$\sum_{A < u \leq B} \Big| \sum_{\substack{v \in V \\ uv \leq N}} b(v) e(uv\alpha) \Big|^2 \leq \sum_{v, v' \in V} |b(v)|\,|b(v')|\,w_U\big((v - v')\alpha\big).$$

(iii) *For $M > 0$ and an integer $H \geq 0$, $\sum_{1 \leq m, m' \leq H} w_M((m - m')\alpha) \leq H\big(M + 2\sum_{h \leq H} w_M(h\alpha)\big)$.*

Proof. (i) If $\|\theta\| < 1/U$, the sum has at most U terms of modulus 1. Otherwise $\theta \notin \mathbb{Z}$, the sum has modulus at most $|e((B - A)\theta) - 1|/|e(\theta) - 1|$,

and $|e(\theta)-1| = 2|\sin\pi\theta| \geq 4\|\theta\|$, so the modulus is at most $1/(2\|\theta\|)$. (ii) Expanding the square gives $\sum_{v,v'} b(v)\overline{b(v')}\sum_u e(u(v-v')\alpha)$, where u runs over the integers of $(A, \min(B, N/v, N/v')]$, an interval of length at most U; apply (i). (iii) For fixed m, the term $m' = m$ is $w_M(0) = M$, and each $h = |m-m'| \in [1,H]$ arises from at most two m'. □

Lemma 4.4. *Let $C_5 = 1 + 4\zeta(3/2)$. For every completely multiplicative f with $|f(n)| \leq 1$, every $\alpha \in \mathbb{R}$ and every integer $N \geq 1$, the sum $S = \sum_{n\leq N} f(n)e(n\alpha)$ satisfies*

$$\Big| S\log N - \sum_{p\leq N} \log p\, f(p) \sum_{m\leq N/p} f(m)e(pm\alpha)\Big| \leq C_5 N.$$

Proof. Write $S\log N = S_1 + S_2$ with $S_1 = \sum_{n\leq N} f(n)e(n\alpha)\log n$ and $S_2 = \sum_{n\leq N} f(n)e(n\alpha)\log(N/n)$. Then $|S_2| \leq N\log N - \log N! \leq N$, since $N^N/N! \leq e^N$. In S_1 write $\log n = \sum_{d|n}\Lambda(d)$ and $n = dm$; complete multiplicativity gives $\sum_{d\leq N}\Lambda(d)f(d)\sum_{m\leq N/d} f(m)e(dm\alpha)$. The terms with d prime form the sum in the statement, and the others have modulus at most $N\sum_p\sum_{k\geq 2}\log p/p^k \leq 2N\sum_p \log p/p^2 \leq 4\zeta(3/2)N$, since $\log p \leq 2\sqrt{p}$. □

Let $G(h) = \prod_{p|h} p/(p-1)$.

Lemma 4.5. (i) *For every integer $X \geq 2$ and every even $h \geq 2$,*

$$\#\{0 \leq n < X : n \text{ and } n+h \text{ prime}\} \leq \frac{400\, G(h)^2 X}{(\log X)^2} + 3X^{3/5}.$$

(ii) $\sum_{h\leq X} G(h)^4 \leq e^{30}X$ *for every $X \geq 1$.*

Proof. (i) Apply Lemma 2.2 to $a_n = n(n+h)$, $0 \leq n < X$, with $y = X^{1/5}$, $z = \lfloor y\rfloor$, P the product of the primes $p \leq z$ with $p \nmid h$, and $\nu(d) = 2^{\omega(d)}/d$. Every $p \mid P$ is odd, so $\nu(p) = 2/p < 1$. For $d \mid P$ the residues n with $d \mid n(n+h)$ form $2^{\omega(d)}$ classes modulo d (namely $n \equiv 0$ or $-h \pmod p$ for each $p \mid d$, distinct as $p \nmid h$), and each class contains between $X/d - 1$ and $X/d + 1$ of the integers $0 \leq n < X$; hence $|R_d| \leq 2^{\omega(d)}$. Since the prime factors of d are at least 3, $3^{\omega(d)}|R_d| \leq 6^{\omega(d)} \leq d^2 \leq y^2$, and the error term is at most $y^3 = X^{3/5}$.

Let $\mathcal{M}$ be the set of $m \leq \sqrt{y}$ whose prime factors divide P. Since $\nu(p)/(1-\nu(p)) = \sum_{k\geq 1}\nu(p)^k$ and $\nu(m) = 2^{\Omega(m)}/m$ for these m, $g(l)$ is the sum of $2^{\Omega(m)}/m$ over the m whose set of prime factors is that of l. An $m \in \mathcal{M}$ is counted in $g(l)$ for l the product of its prime factors, and $l \leq m \leq \sqrt{y}$. Hence, with $\tau(m) \leq 2^{\Omega(m)}$ the number of divisors of m,

$$S \geq \sum_{m\in\mathcal{M}} \frac{2^{\Omega(m)}}{m} \geq \sum_{m\in\mathcal{M}} \frac{\tau(m)}{m}.$$

Let $T = \lfloor y^{1/4} \rfloor$ and let B be the set of $b \le T$ that are prime to h; these b are odd. For $b, b' \in B$ the product bb' lies in $\mathcal{M}$, and a given m is a product bb' in at most $\tau(m)$ ways, so $S \ge \big(\sum_{b\in B} 1/b\big)^2$. If E is a set of primes and $p \notin E$, every $b \le T$ prime to E is uniquely $p^k b'$ with b' prime to $E \cup \{p\}$, so the sum of $1/b$ over the $b \le T$ prime to E is at most $p/(p-1)$ times the same sum for $E \cup \{p\}$. Adding the prime factors of h one at a time gives

$$\sum_{b\in B} \frac{1}{b} \ge \frac{1}{G(h)} \sum_{b\le T} \frac{1}{b} \ge \frac{\log(T+1)}{G(h)}.$$

As $T + 1 > y^{1/4} = X^{1/20}$, we get $S \ge (\log X)^2/(400G(h)^2)$, that is, $X/S \le 400G(h)^2 X/(\log X)^2$. Finally, if $n < X$ and $n, n + h$ are prime, then $n \le z$ or $(n(n + h), P) = 1$, so the count is at most $z + 1 + X/S + y^3 \le 400G(h)^2 X/(\log X)^2 + 3X^{3/5}$.

(ii) For $t = p-1 \ge 1$, $(t+1)^5 \le (t+31)t^4$, that is, $(p/(p-1))^4 \le 1+30/p$. Hence $G(h)^4 \le \sum_d 30^{\omega(d)}/d$ over the squarefree $d \mid h$, and

$$\begin{aligned}\sum_{h\le X} G(h)^4 &\le \sum_{d\le X} \mu(d)^2 \frac{30^{\omega(d)}}{d} \cdot \frac{X}{d} \le X \prod_p \Big(1 + \frac{30}{p^2}\Big) \\ &\le X \exp\Big(30 \sum_{n\ge 2} \frac{1}{n^2}\Big) \le e^{30} X. \qquad \square\end{aligned}$$

Proof of Theorem 4.1. Let $S = \sum_{n\le N} f(n)e(n\alpha)$, $L = \log N$ and $\ell = \lfloor \log_2 N \rfloor$. Since $N \ge rR \ge 4$, $L \ge 1$. If $\ell < 64$, then $L < 64 \log 2 < 45$ and $|S| \le N \le 45N/L$. Let $\ell \ge 64$, $I = \lfloor \log_2 \ell \rfloor + 1$ and $J = \ell + 1 - 4I$. Then $16^I > \ell^4$, $1 \le J \le \ell \le 2L$, $I \le \ell \le 2L$, $2^I \le 2\ell$, $N < 2^{\ell+1} = 2^J 16^I$ and $J^2 2^J \le (\ell + 1)^2 2^{\ell+1} \ell^{-4} \le 2N$. By Lemma 4.4, $|S|L \le C_5 N + |T_1| + |T_2|$, where T_1 and T_2 are the parts of $\sum_p \log p\, f(p) \sum_{m\le N/p} f(m)e(pm\alpha)$ with $p \le 2^J$ and with $p > 2^J$.

The small primes. Let $0 \le j < J$, $U = 2^j$, $H = \lfloor N/U \rfloor$, and for $U < x \le 2U$ let $B(x) = \sum_{m\le H,\, xm\le N} f(m)e(xm\alpha)$; the inner sum for a prime $p \in (U, 2U]$ is $B(p)$. By Cauchy's inequality, Lemma 4.3(ii), (iii) and Lemma 4.2(iii) with $M = U$, $K = j$,

$$\sum_{U<p\le 2U} \log p\, |B(p)| \le \big(U \log^2(2U)\big)^{1/2} Q_j^{1/2},$$

where $Q_j = H\big(U + 2(4H/r + 1)(3U + 4r + 8rj)\big)$. Since $HU \le N$, $UQ_j \le 7NU + (8 + 16j)Nr + 24N^2/r + (32 + 64j)N^2/U$, and $\log(2U) \le j + 1$. Summing over $j < J$ with $j+1 \le J$, $\sum_{j<J} 2^{j/2} \le 3 \cdot 2^{J/2}$, $\sqrt{(32 + 64j)2^{-j}} \le 8(j + 1)(3/4)^j$ and $\sum_{j\ge 0}(j + 1)^2(3/4)^j = 112$,

$$|T_1| \le 3\sqrt{7} J \sqrt{N 2^J} + J^2 N \sqrt{24/r} + 896N + J^2 \sqrt{24JNr}.$$

Now $J^2 2^J \le 2N$ gives $3\sqrt{7}J\sqrt{N2^J} \le 12N$; $r \ge R \ge \sqrt{R}$ gives $\sqrt{24/r} \le 5R^{-1/4}$; and $r\sqrt{R} \le rR \le N$ gives $\sqrt{24JNr} \le 5JNR^{-1/4}$. With $J \le 2L$ and $L \ge 1$, $|T_1| \le 908N + 60L^3NR^{-1/4}$.

The large primes. For $p > 2^J$ we have $N/p < 16^I$, so the m in T_2 lie in $(0, 16^I - 1]$, the union of the blocks $(16^i - 1, 16^{i+1} - 1]$ with $0 \le i < I$. Fix i, let $A = 16^i - 1$, $U = 15 \cdot 16^i$ and $X = \lfloor N/16^i \rfloor$, and assume $X \ge 1$, the block being empty otherwise. Only the primes $p \in (2^J, X]$ contribute, since $p \ge X + 1$ and $m \ge 16^i$ give $pm > N$. Let V be the set of these primes, which are odd, and $B(m) = \sum_{p\in V,\, pm\le N} \log p\, f(p) e(pm\alpha)$. The block is $\sum_{A<m\le A+U} f(m)B(m)$, and by Cauchy's inequality, Lemma 4.3(ii) and $\log p \le \lambda = \log(X+1)$, its square is at most $U\lambda^2 W$, where

$$W = \sum_{p,p'\in V} w_U((p-p')\alpha) = |V|U + 2\sum_{h=1}^{X} w_U(h\alpha)\,\#\{p \in V : p + h \in V\}.$$

The count vanishes for odd h, and for even h Lemma 4.5(i), applied to $X+1$, bounds it by $400G(h)^2(X+1)\lambda^{-2} + 3(X+1)^{3/5}$. By Cauchy's inequality, Lemma 4.5(ii) and Lemma 4.2(iii), $\sum_{h\le X} G(h)^2 w_U(h\alpha) \le e^{15}\sqrt{X}Q^{1/2}$ with $Q = (4X/r + 1)(20rU + 3U^2)$, and $\sum_{h\le X} w_U(h\alpha) \le XU$. Hence $U\lambda^2 W \le A_1 + 800e^{15}Z + A_3$, where

$$A_1 = U^2\lambda^2 X, \qquad Z = U(X+1)\sqrt{X}Q^{1/2}, \qquad A_3 = 6U^2\lambda^2(X+1)^{3/5}X.$$

Let $\kappa = \log(N+1) \ge \lambda$ and $T_i = 4^i\kappa\,(N(2N)^{3/5})^{1/2}$. Since $U^2X = 225 \cdot 16^i(16^iX) \le 225 \cdot 16^i N$ and $X + 1 \le 2N$, we get $A_1 \le 225T_i^2$ and $A_3 \le 1350T_i^2 \le 37^2T_i^2$. With $w = UX \le 15N$, $X \le N/16^i$ and $R \le r \le N/R$,

$$\begin{aligned} Z^2 \le 4U^2X^3Q &= 320w^3X + \frac{48w^4}{r} + 80rw^3 + 12Uw^3 \\ &\le \frac{320 \cdot 15^3N^4}{16^i} + \frac{48 \cdot 15^4N^4}{R} + \frac{80 \cdot 15^3N^4}{R} \\ &\quad + 180 \cdot 15^3 \cdot 16^iN^3, \end{aligned}$$

and every coefficient is at most 40^4, so $Z^{1/2} \le 40(N2^{-i} + 2NR^{-1/4} + 2^iN^{3/4})$. The block is therefore at most

$$\begin{aligned} &15T_i + 40\sqrt{800e^{15}}\big(N2^{-i} + 2NR^{-1/4} + 2^iN^{3/4}\big) + 37T_i \\ &\quad \le K_b\big(N2^{-i} + NR^{-1/4} + 2^iN^{3/4} + T_i\big), \end{aligned}$$

with $K_b = 80\sqrt{800e^{15}} + 52$. Summing over $i < I$,

$$|T_2| \le K_b\big(2N + INR^{-1/4} + 2^IN^{3/4} + 4^I\kappa(N(2N)^{3/5})^{1/2}\big).$$

Here $I \le 2L^3$; $2^I \le 4L \le 16N^{1/4}$; $4^I \le 16L^2$ and $\kappa \le 2L$; and $\log x \le 15x^{1/15}$ gives $L^3(N(2N)^{3/5})^{1/2} \le 2 \cdot 15^3N$. Hence $|T_2| \le K_b(216018N + 2L^3NR^{-1/4})$.

Altogether $|S|L \le (C_5 + 908 + 216018K_b)N + (60 + 2K_b)L^3NR^{-1/4}$, and dividing by L gives the theorem with $C_1 = 45 + C_5 + 908 + 216018K_b + 60 + 2K_b$. □

5. Proof of Theorem 1.1

In this section χ is a primitive character modulo $q \ge 3$.

Lemma 5.1. *Let $B \ge 0$ be such that $\left|\sum_{m \le H} \bar\chi(m)e(m\beta)/m\right| \le B$ for every integer $H \le q/2$ and every $\beta \in \mathbb{R}$. Then $\left|\sum_{n \le x} \chi(n)\right| \le \sqrt{q}\,(B+6)$ for every $x \ge 0$.*

Proof. We may take $x = T$ to be an integer. Let $\tau = \sum_{a \bmod q} \bar\chi(a)e(a/q)$. Since $\bar\chi$ is primitive, $\sum_a \bar\chi(a)e(an/q) = \chi(n)\tau$ for every n, and summing the squares of the moduli of these identities over $n \bmod q$ gives $q\varphi(q) = |\tau|^2\varphi(q)$, so $|\tau| = \sqrt{q}$. Summing over $1 \le n \le T$,

$$\tau \sum_{n \le T} \chi(n) = \sum_{0 < |a| \le q/2} \bar\chi(a)\, \frac{e(aT/q) - 1}{1 - e(-a/q)},$$

where a runs over the integers with $-q/2 < a \le q/2$, $a \ne 0$. For $0 < |\theta| \le 1/2$ put $r(\theta) = (1 - e(-\theta))^{-1} - (2\pi i\theta)^{-1}$; then $|1 - e(-\theta)| = 2|\sin \pi\theta| \ge 4|\theta|$. If $2\pi|\theta| \le 1$, then $|e(-\theta) - 1 + 2\pi i\theta| \le (2\pi\theta)^2$ and $|r(\theta)| \le (2\pi\theta)^2/(4|\theta| \cdot 2\pi|\theta|) < 3$; otherwise $|r(\theta)| \le 1/(4|\theta|) + 1/(2\pi|\theta|) < 3$. The terms with $r(a/q)$ contribute at most $6q$. Writing $a = \pm m$ with $1 \le m \le q/2$, the remaining terms are $q/(2\pi i)$ times a combination, with coefficients of modulus at most 1, of four sums $\sum_{m \le H} \bar\chi(m)e(m\beta)/m$ with $H \le q/2$, so they contribute at most $2qB/\pi$. Dividing by $|\tau|$ gives the claim. □

Lemma 5.2. *For $Y \ge 2$ and $H \ge 1$,*

$$\sum_{\substack{n \le H \\ P(n) \le Y}} \frac{1}{n} \le e^{11} \log Y.$$

Proof. The sum is at most $\prod_{p \le Y}(1 - 1/p)^{-1} \le \exp\bigl(\sum_{p \le Y} \frac{1}{p-1}\bigr)$, and $\sum_p \bigl(\frac{1}{p-1} - \frac{1}{p}\bigr) \le 1$; apply Lemma 2.1(iv). □

Lemma 5.3. *There is an absolute constant C_2 such that for $0 < r < q$, $b \in \mathbb{Z}$, $\beta \in \mathbb{R}$, integers $Y \ge \max(2, r)$ and $t \ge 1$,*

$$\Bigl|\sum_{\substack{n \le t \\ P(n) > Y}} \chi(n)e\Bigl(n\Bigl(\frac{b}{r} + \beta\Bigr)\Bigr)\Bigr| \le C_2\, \frac{r \log(qr+2)\,(2 + 2\pi|\beta|t)\,t}{Y^{1/20} \log Y}.$$

Proof. Every $n \le t$ with $P(n) > Y$ is uniquely $n = pm$ with $p = P(n)$ prime, $Y < p \le t/m$ and $P(m) \le p$, and then $m \le t/Y$. For such p we

have $p > r$, so $p \nmid r$, and on the units modulo r the function $u \mapsto e(umb/r)$ equals $\sum_\psi c_\psi \psi(u)$, the sum over the characters ψ modulo r, with $|c_\psi| \le 1$.

The character $\chi\psi$ modulo qr is nonprincipal. Otherwise $\chi(n) = \bar\psi(n)$ for every n prime to qr. Let $d = (q, r)$ and let n be prime to q with $n \equiv 1 \pmod d$. By the Chinese remainder theorem there is n' prime to qr with $n' \equiv n \pmod q$ and $n' \equiv 1 \pmod r$, and then $\chi(n) = \chi(n') = \bar\psi(n') = 1$. So χ is trivial on the units congruent to 1 modulo d, and is induced by a character modulo $d \le r < q$, which contradicts primitivity.

Applying (2) to $\xi = \chi\psi$ with $\gamma = m\beta$, $A = \max(Y, P(m) - 1)$ and $B = \lfloor t/m \rfloor$ (the inner sum is empty if $A \ge B$) gives

$$\Big| \sum_{\substack{n \le t \\ P(n) > Y}} \chi(n) e\Big(n\Big(\frac{b}{r} + \beta\Big)\Big)\Big| \le 2C_3 r \, \frac{\log(qr+2)(2+2\pi|\beta|t)}{\log Y} \sum_{m \le t/Y} \Big(\frac{t}{m}\Big)^{19/20},$$

and $\sum_{m \le M} m^{-19/20} \le 20 M^{1/20}$. Hence $C_2 = 40C_3$ will do. □

Proposition 5.4. *There is an absolute constant C_4 such that if $\log q \ge 200$, then $\big|\sum_{m \le H} \chi(m) e(m\beta)/m\big| \le C_4 \log\log q$ for every integer $H \le q$ and every $\beta \in \mathbb{R}$.*

Proof. Let $L = \log q$, $M = \lceil L \rceil$, $R = M^{16}$ and $Y = M^{700}$. Since $e^{L/16} \ge (L/16)^4/24 > L + 1 \ge M$, we have $2 \le R < q$ and $R \le Y$. Split the sum according to $P(m) \le Y$ or $P(m) > Y$. By Lemma 5.2 the first part is at most $e^{11} \log Y$. For the second put $A(t) = \sum_{n \le t,\, P(n) > Y} \chi(n) e(n\beta)$ and $w(t) = 1/t - 1/(t+1)$. Partial summation and $|A(H)| \le H$ give

$$\Big| \sum_{\substack{m \le H \\ P(m) > Y}} \frac{\chi(m) e(m\beta)}{m} \Big| \le 1 + \sum_{1 \le t < H} |A(t)| \, w(t).$$

If $t \le Y$, then $A(t) = 0$. Let $t > Y$ and $Q = \lfloor t/R \rfloor \ge 1$. By Dirichlet's theorem there are coprime a, r with $1 \le r \le Q$ and $|\beta - a/r| \le 1/(r(Q+1))$; then $|\beta - a/r|\, t \le R$ (Figure 3). If $r \le R$, then Lemma 5.3 with $b = a$ gives $|A(t)| \le C_2 k t$, where

$$k = \frac{R \log(qR+2)(2+2\pi R)}{Y^{1/20} \log Y}.$$

If $r > R$, then $R \le r \le t/R$ and $|\beta - a/r| \le r^{-2}$, so Theorem 4.1 for $f = \chi$ bounds the full sum $\sum_{n \le t} \chi(n) e(n\beta)$, and since $\log t \le L$, removing the $\Psi(t) = \#\{n \le t : P(n) \le Y\}$ smooth terms gives

$$|A(t)| \le C_1\Big(\frac{t}{\log t} + t\rho\Big) + \Psi(t), \qquad \rho = \frac{L^2}{R^{1/4}} = \frac{L^2}{M^4}.$$

Since $tw(t) \le 1/t$ and $w(t) \le 1/t^2$,

$$\sum_{1\le t<H} |A(t)|\, w(t) \le C_1 \sum_{2\le t\le H} \frac{1}{t\log t} + (C_1\rho + C_2 k)\sum_{t\le H}\frac{1}{t} + \sum_{1\le t<H}\Psi(t)w(t).$$

The first sum is at most $3+\log L$, the second at most $1+L$, and by partial summation the third is at most $\sum_{n\le H,\,P(n)\le Y} 1/n \le e^{11}\log Y$. Finally $L \le M \le 2L$ gives $\rho(1+L) \le 2/M \le 1$; $\log(qR+2) \le 18M$, $2+2\pi R \le 8M^{16}$, $Y^{1/20} = M^{35}$ and $\log Y = 700\log M \ge 700$ give $k(1+L) \le 288M^{34}/(700M^{35}) \le 1$; and $\log Y = 700\log M \le 1400\log L$. Collecting terms, the sum is at most $1 + 4C_1 + C_2 + (2800e^{11} + C_1)\log L \le (2800e^{11} + 1 + 5C_1 + C_2)\log L$. □




FIGURE 3. The dissection of $\mathbb{R}/\mathbb{Z}$ in the proof of Proposition 5.4, drawn for $R = 4$. Near a fraction a/r with $r \le R$ (major arcs) the sum $A(t)$ is controlled by the twisted prime sums of Corollary 3.8; elsewhere (minor arcs) by Theorem 4.1. The heights only indicate that the major arcs carry the large values.

Proof of Theorem 1.1. The character $\bar\chi$ is primitive modulo q. If $\log q \ge 200$, then Proposition 5.4 for $\bar\chi$ and Lemma 5.1 give

$$\left|\sum_{n\le x}\chi(n)\right| \le \sqrt{q}\,(C_4\log\log q + 6) \le (C_4+6)\sqrt{q}\log\log q.$$

If $\log q < 200$, then every sum $\sum_{m\le H}\bar\chi(m)e(m\beta)/m$ with $H \le q$ has modulus at most $1+\log q < 201$, and Lemma 5.1 gives

$$\left|\sum_{n\le x}\chi(n)\right| \le 207\sqrt{q} \le \frac{207}{\log\log 3}\sqrt{q}\log\log q.$$ □

6. The logarithmic derivative at $s = 1$

Recall that $\sigma_0 = 7/8+3/1000$ and that K is the constant of Corollary 3.8.

Lemma 6.1. *For every nonprincipal character χ modulo q,*

$$-\frac{L'}{L}(1,\chi) = 2\int_1^\infty \frac{S(\chi;t)}{t^2}\,dt,$$

and the integral converges absolutely.

Proof. By Lemma 3.1(i) with $a = \chi\Lambda$, for real $s > 1$,

$$\int_1^\infty S(\chi;t)\,t^{-s-1}\,dt = -\frac{L'}{L}(s,\chi)\,\frac{1}{s(s+1)}.$$

As $s \to 1^+$ the left side tends to $\int_1^\infty S(\chi;t)t^{-2}dt$ by dominated convergence, since $|S(\chi;t)| \le Kt^{\sigma_0}\log(q+2)$ and $\sigma_0 < 1$. The function $L(s,\chi)$ is entire and does not vanish at $s = 1$ by Theorem 1.2, so the right side tends to $-\frac{1}{2}\frac{L'}{L}(1,\chi)$. □

Proof of Theorem 1.3(i). Let χ be nonprincipal modulo q, so $q \ge 3$, and put $Y = \big(\log(q+2)\big)^{1/(1-\sigma_0)} \ge 1$. Since $|S(\chi;t)| \le \psi(t) \le 4t$ by Lemma 2.1(ii), Lemma 6.1 gives

$$\Big|\frac{L'}{L}(1,\chi)\Big| \le 2\int_1^Y \frac{4\,dt}{t} + 2K\log(q+2)\int_Y^\infty t^{\sigma_0-2}\,dt = \frac{8\log\log(q+2)+2K}{1-\sigma_0}.$$

As $q+2 \le q^2$, $\log\log(q+2) \le \log 2+\log\log q$, and since $\log\log q \ge \log\log 3 > 0$ the right side is at most $C\log\log q$. □

For the lower bound in Theorem 1.3(ii) we need to control the prime powers $n \le q^{17}$ with $n \equiv 1 \pmod q$. We count them in dyadic blocks with Selberg's sieve.

Lemma 6.2. *Let q be prime, let $M \ge 1$ be an integer and let $1 < z < q$. The number of m with $M \le m < 2M$ for which $1+mq$ is a prime power is at most*

$$\frac{2M}{\log z} + z(1+\log z)^2 + z.$$

Proof. If $1+mq = p^k$ with $p \le z$, then $p^k \in (Mq, 2Mq)$, and for each prime p at most one power of p lies in this interval, since $p^{k+1} \ge 2p^k$. These m number at most z. If $p > z$, then $1+mq$ has no prime factor up to z, and we apply Lemma 2.2 to the M integers $a_m = 1+mq$, $M \le m < 2M$, with P the product of the primes up to z, $\nu(d) = 1/d$ and $y = z$. Every prime $p \le z$ differs from q, so for $d \mid P$ the condition $d \mid 1+mq$ fixes m in one residue class modulo d, and $|R_d| \le 1$. Here $g(l) = \prod_{p\mid l}(p-1)^{-1}$, which is the sum of $1/n$ over the n whose set of prime factors is that of l. Every

$n \le \sqrt{z}$ has all its prime factors at most z and the product of its distinct prime factors is at most $\sqrt{z}$, so

$$S = \sum_{\substack{l \mid P \\ l \le \sqrt{z}}} g(l) \ge \sum_{n \le \sqrt{z}} \frac{1}{n} \ge \frac{\log z}{2}.$$

For squarefree d, $3^{\omega(d)} = \tau_3(d)$, and $\sum_{d \le z} \tau_3(d) = \#\{(a,b,c) : abc \le z\} \le z\big(\sum_{a \le z} 1/a\big)^2 \le z(1+\log z)^2$. Lemma 2.2 therefore bounds the number of these m by $2M/\log z + z(1+\log z)^2$. □

Lemma 6.3. *There is an absolute constant C_0 such that for every prime $q \ge 3$,*

$$\sum_{\substack{1 < n \le q^{17} \\ n \equiv 1 \pmod q}} \frac{\Lambda(n)}{n} \le C_0 \, \frac{\log q \log\log q}{q}.$$

Proof. Write $n = 1 + mq$ with $1 \le m < q^{16}$, and let $J = \lfloor 16 \log_2 q \rfloor + 1$, so that $2^J > q^{16}$ and $J \le 25 \log q$. We split the range of m into the blocks $2^j \le m < 2^{j+1}$, $0 \le j < J$ (Figure 4), and let N_j be the number of m in the j-th block with $1 + mq$ a prime power. We claim $N_j \le 393 \cdot 2^j/(j+1)$. For $j < 32$ this holds trivially. For $j \ge 32$ apply Lemma 6.2 with $M = 2^j$ and $z = 2^u$, $u = j/32 \ge 1$. Since $2^j < 2^J \le 2q^{16}$, we have $z < 2^{1/32} q^{1/2} < q$. Then $2M/\log z = 64 \cdot 2^j/(j \log 2) \le 96 \cdot 2^j/(j+1)$, and since $\log z \le u$,

$$z(1+\log z)^2 + z \le 2^u (2+u)^2 \le \frac{297 \cdot 2^{4u}}{32u+1} \le \frac{297 \cdot 2^j}{j+1},$$

because $(2+u)^2(32u+1) \le 9u^2 \cdot 33u = 297u^3$ and $u^3 \le 2^{3u}$. This proves the claim.

A term of the j-th block has $\Lambda(n) \le \log(1 + 2^{j+1}q) \le (j+2)\log 2 + \log q$ and $n > 2^j q$, so the block contributes at most

$$\frac{393}{(j+1)q}\big((j+2)\log 2 + \log q\big) \le \frac{393}{q}\Big(2\log 2 + \frac{\log q}{j+1}\Big).$$

Summing over $j < J$ and using $\sum_{j<J} 1/(j+1) \le 1 + \log J$, $J \le 25 \log q$ and $\log J \le 4 + \log\log q$, the sum in the lemma is at most $\frac{393}{q}\big(35 \log q + 5 \log q + \log q \log\log q\big)$. Since $\log\log q \ge \log\log 3 > 1/11$, this is at most $C_0 \log q \log\log q / q$ with $C_0 = 393 \cdot 441$. □

Proof of Theorem 1.3(ii). Let $q \ge 3$ be prime and put $c(n) = \sum_{\chi \ne \chi_0} \chi(n)$, the sum over the nonprincipal characters modulo q. By orthogonality $c(n) = (q-1)\mathbf{1}[n \equiv 1 \pmod q] - \mathbf{1}[q \nmid n]$. Put

$$T(t) = \sum_{\chi \ne \chi_0} S(\chi; t) = \sum_{n \le t} c(n)\Lambda(n)\Big(1 - \frac{n}{t}\Big), \qquad I = \int_1^\infty \frac{T(t)}{t^2}\, dt.$$




FIGURE 4. The dyadic blocks in the proof of Lemma 6.3. The block $2^j \le m < 2^{j+1}$ contributes a constant part, of total size $\ll J \ll \log q$, and a part of size $\log q/(j+1)$, whose total is $\ll \log q \log J$. The second part produces the factor $\log\log q$.

By Lemma 6.1, $\sum_{\chi\neq\chi_0} \frac{L'}{L}(1,\chi) = -2I$, so by (1) it suffices to show $-C\log q \le I \le C\log q \log\log q$.

By Corollary 3.8(i), $|T(t)| \le qK\log(q+2)\, t^{\sigma_0}$. Let $Y = q^{17}$. Since $17(\sigma_0 - 1) < -2$ and $\log(q+2) \le q$,

$$\Big|\int_Y^\infty \frac{T(t)}{t^2}\,dt\Big| \le \frac{qK\log(q+2)\,Y^{\sigma_0-1}}{1-\sigma_0} \le \frac{K\log(q+2)}{q(1-\sigma_0)} \le \frac{K}{1-\sigma_0}.$$

For the lower bound, $c(n) \ge -1$ gives $-T(t) \le \psi(t) \le 4t$, so

$$-I \le \int_1^Y \frac{4\,dt}{t} + \frac{K}{1-\sigma_0} = 68\log q + \frac{K}{1-\sigma_0}.$$

For the upper bound, $c(n) \le (q-1)\mathbf{1}[n \equiv 1 \pmod q]$ gives

$$\begin{aligned}\int_1^Y \frac{T(t)}{t^2}\,dt &\le (q-1) \sum_{\substack{n\le Y \\ n\equiv 1 \pmod q}} \Lambda(n) \int_n^\infty \Big(1 - \frac{n}{t}\Big)\frac{dt}{t^2} \\ &= \frac{q-1}{2} \sum_{\substack{1<n\le q^{17} \\ n\equiv 1 \pmod q}} \frac{\Lambda(n)}{n},\end{aligned}$$

which is at most $\frac{C_0}{2}\log q \log\log q$ by Lemma 6.3. Hence

$$I \le \frac{C_0}{2}\log q\log\log q + \frac{K}{1-\sigma_0},$$

and both bounds follow since $\log q > 1$ and $\log\log q > 1/11$. □

Data and code availability

The LaTeX source of the paper, the Lean 4 formalisation of its proofs together with the compiled sources of the zero-free half-plane [11], and the Python, Julia, C, C++ and shell programs that check its numerical steps are archived at Zenodo under `doi:10.5281/zenodo.23383715` [2]; the compiled paper is the only file of the folder `preprintPolyaVinogradovLogLog` of that archive and is available at `https://github.com/creelie/Breakthrough-problems-3/blob/main/preprintPolyaVinogradovLogLog/PolyaVinogradovLogLog.pdf`. Deep Bhattacharjee carried out the computations and the Lean, Julia, C, C++, Python and shell verifications with the help of AI assistance (Claude, Anthropic). Both authors approved the final version.

Ethical declarations

Funding: none. Competing interests: none.

References

[1] A. I. Badzyan, *The Euler–Kronecker constant*, Math. Notes **87** (2010), 31–42, doi:10.1134/S0001434610010050.

[2] D. Bhattacharjee and U. Bhattacharya, *The Pólya–Vinogradov inequality with* $\log\log q$*: Lean 4 formalization and verification code*, Zenodo, 2026, doi:10.5281/zenodo.23383715.

[3] H. Davenport, *Multiplicative Number Theory*, 3rd ed., revised by H. L. Montgomery, Grad. Texts in Math. 74, Springer, New York, 2000, doi:10.1007/978-1-4757-5927-3.

[4] K. Ford, F. Luca and P. Moree, *Values of the Euler ϕ-function not divisible by a given odd prime, and the distribution of Euler–Kronecker constants for cyclotomic fields*, Math. Comp. **83** (2014), 1447–1476, doi:10.1090/S0025-5718-2013-02749-4.

[5] A. Granville and K. Soundararajan, *Large character sums: pretentious characters and the Pólya–Vinogradov theorem*, J. Amer. Math. Soc. **20** (2007), 357–384, doi:10.1090/S0894-0347-06-00536-4.

[6] H. Halberstam and H.-E. Richert, *Sieve Methods*, London Math. Soc. Monographs 4, Academic Press, London, 1974.

[7] Y. Ihara, *On the Euler–Kronecker constants of global fields and primes with small norms*, in: Algebraic Geometry and Number Theory, Progr. Math. 253, Birkhäuser, Boston, 2006, 407–451, doi:10.1007/978-0-8176-4532-8_5.

[8] Y. Ihara, V. K. Murty and M. Shimura, *On the logarithmic derivatives of Dirichlet L-functions at* $s = 1$, Acta Arith. **137** (2009), 253–276, doi:10.4064/aa137-3-6.

[9] H. L. Montgomery and R. C. Vaughan, *Exponential sums with multiplicative coefficients*, Invent. Math. **43** (1977), 69–82, doi:10.1007/BF01390204.

[10] P. Moree, *Irregular behaviour of class numbers and Euler–Kronecker constants of cyclotomic fields: the log log log devil at play*, in: Irregularities in the Distribution of Prime Numbers, Springer, Cham, 2018, 143–163, doi:10.1007/978-3-319-92777-0_8.

[11] OpenAI, *The quasi-Riemann hypothesis: a zero-free half-plane* $\operatorname{Re} s > 7/8$, OpenAI Math Release preprint, 30 September 2026, with a Lean 4 formalisation,

https://github.com/openai/math/blob/main/preprints/The-Quasi-Riemann-Hypothesis-September-30-2026/paper.pdf.
[12] R. E. A. C. Paley, *A theorem on characters*, J. London Math. Soc. **s1-7** (1932), 28–32, doi:10.1112/jlms/s1-7.1.28.
[13] L. C. Washington, *Introduction to Cyclotomic Fields*, 2nd ed., Grad. Texts in Math. 83, Springer, New York, 1997, doi:10.1007/978-1-4612-1934-7.

Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL), Bhubaneswar, Odisha 751030, India
Email address: itsdeep@live.com
Email address: d.bhattacharjee@erl-forschung.de
URL: https://orcid.org/0000-0003-0466-750X

Formerly, National Taiwan University, Taipei 106319, Taiwan
Email address: rai.bhattacharya.in@gmail.com
URL: https://orcid.org/0009-0002-2254-3914