




$T_0X = T_1 \oplus T_2$: the eigenplanes of F_0 .

T_pC' lies in at most one of them; here in T_1 .

$T_2 \cap T_pC' = 0$, so π_2 has nonzero image in

$\bigwedge^2 N$, $N = T_pX/T_pC'$, and the germ of

$\text{ev}_{E'}(x_2)$ at p is that image:

$x_2 \lrcorner \text{ch}(E') = 0$ and $\text{ev}_{E'}(x_2) \neq 0$.

pairs $\{k, l\}$ with $a_{\partial_k} a_{\partial_l} \neq 0$

- (i) $\mathcal{O}_C$, a line bundle on a smooth curve
- (ii) $k(p)$, a skyscraper
- (iii) a curve ideal in a smooth divisor
- (iv) a point ideal in a smooth divisor
- (v) $\mathcal{O}_D$, a smooth divisor
- (vi) two planes meeting at a point
- (vii) maximal Cohen–Macaulay on a node
- (viii) the glued germ of Example 11.2.7
- (ix) reflexive, non-Cohen–Macaulay, node
- (x) $\mathcal{O}_C^2$, rank two on a smooth curve

	12	13	14	23	24	34	pd
(i)	■	■	□	■	□	□	3
(ii)	■	■	■	■	■	■	4
(iii)	■	■	□	□	□	□	2
(iv)	■	■	■	■	■	■	3
(v)	□	□	□	□	□	□	1
(vi)	□	■	■	■	■	□	2
(vii)	□	□	□	□	□	□	1
(viii)	□	□	□	■	■	■	3
(ix)	■	□	■	■	□	■	2
(x)	■	■	□	■	□	□	3

coordinates adapted to the germ, as in item (LXVIII): a filled cell is a nonzero Yoneda square in the local Ext^2 ; at a point of C' the pairs of normal directions are filled, and one of π_1, π_2 projects onto them.