CONJECTURES OF ASHOKE SEN: SCOPE, EXACT TESTS AND ANALYTIC LIMITS

DEEP BHATTACHARJEE

Abstract. *We examine thirty-six selected statements associated with Ashoke Sen's work on duality, tachyon condensation, black hole indices and string field theory. The purpose is to distinguish mathematical results within a specified framework, physical conjectures, and extensions beyond the assumptions used in a derivation. Exact product expansions reproduce known negative dyon indices in toroidal orbifolds; with the charge cone transcribed correctly, a finite scan contains thirty negative-index charge triples. These numbers concern the full index, whose sign need not agree with a horizon index after hair removal. For tachyon dust we give the characteristic flow, the corrected Hessian sign criterion, and precise hypotheses under which the first focusing time is attained. We also separate the D7 divisor on the F-theory base from its singular pullback to the orientifold cover and give a local weak-coupling family with an I_0^* divisor. These calculations delimit claims; they do not disprove the homogeneous rolling solution or Sen's positivity argument under its stated assumptions. The accompanying formal certificates check specified finite computations and algebraic identities, rather than the catalogue's physical interpretations. Established results and later corrections are credited to their sources.*

Note. This paper was first made public on 25 October 2026, as release `v1.0.0` of the repository `https://github.com/creelie/paper_11` (concept DOI 10.5281/zenodo.23652472, version DOI 10.5281/zenodo.23652473); it was revised there on 26 October 2026 as `v1.0.1` (10.5281/zenodo.23652641) and `v1.1.0` (10.5281/zenodo.23656435). The first version counted two of the thirty-six statements as false and twenty-one negative charge triples; it used a charge cone with the discriminant in place of the linear inequality of Sen's equation (3.9), and its dust theorem replaced a supremum by a maximum without an attainment hypothesis. The later versions corrected the cone, extended the scan to thirty triples, restated the two verdicts as limits of scope, since the negative indices are full indices that the hair analysis of Chattopadhyaya and David already accounts for and the caustic concerns the pressureless effective equations rather than the homogeneous solution, and added the infrared appendices. This version is the same mathematics in a single-column form for submission; its source is release `v2.0.0` of the same repository, archived as 10.5281/zenodo.23669248, with the exact program that computes every table, the Lean 4 certificates of the finite statements, and the scripts that check the explicit constants of the appendices.

1. Introduction

Ashoke Sen has published conjectures for more than thirty years, and most of them were made in the same way: a computation in one regime, a symmetry that carries it to a regime where nothing can be computed, and a statement about that second regime sharp enough to be refuted. The statement that the heterotic string on a six-torus is invariant under $SL(2,\mathbb{Z})$ required a bound state of two monopoles that nobody had seen [115, 116]. The statement

Date: 26 October 2026.

2020 *Mathematics Subject Classification.* 81T30; 81T13; 83C57; 11F46; 35L67; 53C26.

Key words and phrases. S-duality; tachyon condensation; black hole entropy; Siegel modular forms.

Corresponding author: Deep Bhattacharjee. Email: d.bhattacharjee@erl-forschung.de, itsdeep@live.com, egspl@proton.me; ORCID 0000-0003-0466-750X. Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL), Bhubaneswar, Odisha 751030, India, and formerly, University of Calcutta, Senate House, 87/1 College Street, Kolkata 700073, West Bengal, India.

that the tachyon of an unstable D-brane rolls to a vacuum without open strings, at an energy equal to the tension of the brane [119, 125], required an exact solution of open string field theory that did not exist for seven years. The statement that the Fourier coefficients of a Siegel modular form have a definite sign on a cone of charges [100] required an argument about the angular momentum of a black hole. These are predictions in the strict sense, and they can be checked.

This paper examines a selected catalogue, not the complete publications of one author. For each entry we separate the original hypotheses, the result actually derived, and subsequent tests or proofs. An unproved conjecture is not a false theorem. A counterexample to a statement with an assumption removed is not a counterexample to the statement with that assumption retained.

The labels in the catalogue describe scope. PROVED means established in the mathematical or perturbative framework identified in the corresponding section; it does not assert a nonperturbative construction of string theory. CONFIRMED means supported by the cited tests without a general proof supplied here. SCOPE LIMITED means that an unrestricted restatement loses a necessary hypothesis or exceeds the approximation. The two HEURISTIC entries are proposals rather than theorem statements. The catalogue is a reading guide, not a numerical ranking of scientific merit. Claims already settled or qualified in the cited literature are included as context, not presented as newly discovered errors. Burnt orange marks an explicitly unsupported extension; yellow marks a hypothesis or unresolved step. Neither colour attributes misconduct or a false proof to an author.

Two exact tests receive detailed treatment. For the freely acting toroidal orbifolds, the negative full dyon indices were already found by Chattopadhyaya and David [16]; their later hair analysis [17] explains why the horizon index is a different quantity. Our contribution is a reproducible finite computation with the linear charge-cone inequality of [100, Eq. (3.9)] retained. The corrected scan contains thirty negative triples in the stated range. The original twenty-one listed triples have a separate finite Lean certificate; the larger scan is checked by the supplied exact Python program. No inference from thirty charges to infinitely many charges is made without the published Fourier–Jacobi argument.

For tachyon matter, the characteristic calculation applies to the pressureless effective equations. Focusing is controlled by a positive eigenvalue of the Hessian of the initial tachyon profile on the branch $\partial_t T > 0$. Concave profiles have no forward local focusing. Bounded nonconstant profiles have positive curvature somewhere and hence a finite focal time on some characteristic. The first time is a minimum only under an attainment hypothesis. This describes the limit of a classical effective description, not a counterexample to the homogeneous solution in [88, 87]. Indeed [86] also gives static dust with arbitrary nonconstant density, a useful example that must be retained.

The F-theory calculation similarly requires two spaces to be kept distinct: the base, where $\eta^2 - h\chi = 0$ can be smooth, and the orientifold double cover, where its pullback has a double locus. Non-Higgsable gauge symmetry alone is not an obstruction to weak coupling: I_0^* is an exception, as the explicit family below illustrates. Small-black-hole mismatches and the localisation issues that Sen himself discussed are historical qualifications, not new criticisms.

Plan. Section 2 is the catalogue, Tables 1, 2, 3 and 4, with the verdict of every claim and the place where it is treated. Sections 3 and 4 treat duality: the monopole conjectures and the counting of dyons, with the exact verification of the wall-crossing formula. Section 5 treats the positivity conjecture and its domain of applicability. Sections 6 to 8 treat the tachyon conjectures, the rolling tachyon with the caustic theorem, and the non-BPS branes. Section 9 treats the Sen limit, Sections 10 and 11 the entropy function, the small black holes and the quantum entropy function, Section 12 string field theory, D-instantons, the soft theorems and matrix theory. Section 13 records the remaining questions. Appendix A describes the computation of the Fourier coefficients and Appendix B completes the proof of the caustic theorem.

Claim	Source	Verdict	Where
Duality			
1. The heterotic string on T^6 is invariant under $SL(2,\mathbb{Z})$ acting on the axion–dilaton	[115]	confirmed	§3.1
2. The Atiyah–Hitchin manifold carries exactly one L^2 anti-self-dual harmonic two-form, the two-monopole bound state	[116]	proved	§3.2, Thm 3.2
3. The L^2 cohomology of the moduli space of k monopoles, for every k, is as S-duality requires	[116, 84]	confirmed	§3.3
4. A manifestly duality-symmetric action is equivalent to the Maxwell and heterotic actions	[83]	proved	§3.4
5. The Kerr–Sen fields exactly solve the specified leading-order heterotic equations	[114]	proved	§3.4
Quarter-BPS dyons			
6. The CHL product has the stated modularity; its physical counting interpretation uses duality	[56, 34]	proved	§4.1
7. Across a wall of marginal stability the index jumps by $(-1)^{Q\cdot P+1}\lvert Q\cdot P\rvert\, d_h(Q)\, d_h(P)$	[97, 96]	proved	§4.2, Thm 4.2
8. Negative-discriminant indices in the stated sectors are accounted for by two-centred sums	[99]	proved	§4.3
9. A single-centred index is positive when the stated horizon and hair assumptions hold	[100]	scope limited	§5, Thm 5.4
10. The hair of a quarter-BPS black hole can be identified and removed, leaving a horizon index	[12]	confirmed	§5.3

TABLE 1. The catalogue, claims 1 to 10. The labels refer to the frameworks and assumptions discussed in the text.

The primary-source register in Appendix C identifies the passages used for the principal scope corrections. Appendix E constructs an explicit finite-energy soft model. Appendix F derives a quantitative subtracted tree-emission bound and identifies the uniform loop estimates still needed for transfer to the complete theory. Literature-status statements concern the sources examined, rather than a claim to have read every citing paper.

2. The catalogue

Tables 1, 2, 3 and 4 list the claims, and Figure 1 shows them by verdict. The first column gives the claim in one line, as a concise restatement, with hypotheses supplied in the corresponding section; the second the paper in which it was made; the third the verdict; the fourth the section of this paper where the claim is examined and, where one exists, the theorem that settles it. The column of verdicts is the summary of the paper, and every entry in it is justified in the section named.[1]

The list records selected claims from the earlier manuscript. The short entries must be read with the hypotheses in the indicated section. A result established within an effective action is not automatically a theorem about the full string theory. Later proofs and qualifications remain credited in those sections; they are excluded from the list of proposed new objections.

[1]The format of the table, one claim per line with its source, its verdict and the place of the argument, is the barrier ledger of the engine Anaxia (`github.com/creelie/eBLLM`, release 3.1.0, `doi:10.5281/zenodo.23484160`), described in [11] and used in the same way for the records of [38, 130]. The engine was not run for this paper; only its ledger format is used.

Claim	Source	Verdict	Where
Tachyons on unstable branes			
11. The tachyon potential of a D-brane is universal, independent of the background	[125]	proved	§6.1
12. The tachyon vacuum has energy density $-\mathcal{T}_p$ relative to the brane	[119, 124]	proved	§6.2
13. Lower-dimensional D-branes are lump solutions of the tachyon effective theory	[119, 124]	proved	§6.2
14. There are no open string excitations around the tachyon vacuum	[124, 78]	proved	§6.2
15. The three conjectures hold in the superstring field theory of Berkovits	[15]	confirmed	§6.2
16. The homogeneous rolling tachyon conserves energy and its pressure vanishes exponentially	[88]	proved	§7.1
17. Homogeneous rolling has a pressureless late-time limit; inhomogeneous dust can focus	[87, 86]	scope limited	§7.2, Thm 7.4
18. Open–closed duality at tree level: the open string description of the decay encodes the closed strings emitted	[90]	confirmed	§7.3
19. The kink on a non-BPS Dp-brane is the BPS D$(p-1)$-brane (descent relations)	[122, 123]	proved	§8
20. Type I string theory has a stable non-BPS D-particle, the $SO(32)$ spinor	[122]	proved	§8
21. The world-volume theory of the kink and the vortex is the Dirac–Born–Infeld action of the lower brane	[89]	proved	§8

TABLE 2. The catalogue, claims 11 to 21. Claim 17 requires separate treatment of the effective dust limit.

Claim	Source	Verdict	Where
F-theory			
22. F-theory on an elliptic K3 has a weak coupling limit that is the type IIB orientifold on $T^2/\mathbb{Z}_2$	[117]	proved	§9
23. The Sen family gives a formal orientifold limit; smoothness and global weak coupling need separate hypotheses	[118]	scope limited	§9
Black hole entropy			
24. The entropy function, extremised over the near-horizon data, gives the Wald entropy of an extremal black hole	[93]	proved	§10.1
25. The entropy function extends to rotating extremal black holes	[1]	proved	§10.1
26. A Gauss–Bonnet truncation stretches small horizons; its entropy coefficient agrees in $D = 4, 5$	[92, 94]	scope limited	§10.2
27. The quantum entropy function, a path integral over AdS_2, equals the microscopic index	[98]	confirmed	§11.1
28. The logarithmic correction to the entropy is 0 for $\mathcal{N} = 4$ and $-4 \ln A_H$ for $\mathcal{N} = 8$	[7]	proved	§11.2
29. For $\mathcal{N} = 2$ the correction is $\frac{1}{12}(23 + n_H - n_V) \ln(A_H/G_N)$	[101]	confirmed	§11.2

TABLE 3. The catalogue, claims 22 to 29. Claims 23 and 26 record geometric and truncation qualifications already present in the literature.

Claim	Source	Verdict	Where
String field theory and other claims			
30. Closed string field theory is background independent under infinitesimal deformations	[127]	proved	§12.1
31. Superstring field theory has a gauge-invariant 1PI effective action and a BV master action, and is unitary	[104, 105, 106, 107]	proved	§12.1
32. The normalisation of D-instanton amplitudes is fixed by string field theory	[109, 112, 111]	confirmed	§12.2
33. The classical logarithmic soft graviton terms are universal	[64, 80]	proved	§12.3
34. The quantum logarithmic soft terms have the conjectured universal form	[80]	confirmed	§12.3
35. A network of (p, q) strings is a BPS configuration of type IIB	[121]	proved	§3.4
36. Every microstate of a single-centred BPS black hole carries zero angular momentum after the fermion zero modes are removed	[21]	confirmed	§5.3
H1. M-theory on T^n is the large-N limit of D0-branes on T^n for every n	[120]	heuristic	§12.4
H2. For $n \geq 6$ the matrix description is a theory without an independent definition	[120]	heuristic	§12.4

TABLE 4. The catalogue, claims 30 to 36 and the two heuristic statements, which receive no verdict.




FIGURE 1. The thirty-six claims of Tables 1, 2, 3 and 4, one block per claim, grouped by the section of the paper that treats them and coloured by verdict: established in the stated framework, supported by cited tests, and scope limited. Colours describe statements, not authors.

3. Duality and the monopole conjectures

3.1. **The $SL(2,\mathbb{Z})$ conjecture.** The conjecture of [115] (claim 1 of Table 1) states that the heterotic string compactified on T^6, whose low energy theory is $\mathcal{N}=4$ supergravity coupled to 22 vector multiplets, is invariant under the group $SL(2,\mathbb{Z})$ generated by the shift of the axion and by the inversion $\lambda \mapsto -1/\lambda$ of the complex coupling $\lambda = a + ie^{-\phi}$, acting on the electric and magnetic charge vectors (Q,P) of the $O(6,22)$ lattice by the same matrix.[2] The inversion exchanges weak and strong coupling, so no perturbative computation can test it directly. Sen reduced the test to the spectrum of states that saturate the Bogomolnyi bound, whose masses and degeneracies are protected by supersymmetry: the conjecture requires that for every pair of coprime integers (p,q) there exist states carrying p units of electric and q units of magnetic charge, in the same supermultiplet as the elementary string states with $(1,0)$.

For $q=1$ these are the dyons of the $\mathcal{N}=4$ theory and they were known to exist [73]. For $(p,q)=(1,2)$ the conjecture requires a bound state of two monopoles and one unit of electric charge, and nothing in the classical theory produces such a state: the static BPS monopoles have no net force, but a normalisable quantum bound state is not supplied by a static separated configuration and there is no classical bound state. The prediction is therefore a genuine one, and it was tested in [116], as described in the next subsection. The subsequent tests of the conjecture are the three-string junction states [121, 95], the counting of quarter-BPS dyons (Section 4), whose partition function is a Siegel form invariant under the duality group, the match of the black hole entropy with the index (Section 10), and the equivalent statement for $\mathcal{N}=4$ super-Yang–Mills theory, where the conjecture of Montonen and Olive [68] is tested by the same bound states [116, 75, 48, 65], by the modular properties of partition functions on four-manifolds [128], and by its embedding into the geometric Langlands programme [57].

Verdict 3.1 (claim 1)**.** Confirmed. Every protected quantity that has been computed is consistent with the $SL(2,\mathbb{Z})$ symmetry, and the strongest test, the exact dyon index of Section 4, is a Siegel form of the duality group. A general nonperturbative proof is not supplied here; which has no definition at strong coupling other than through such dualities; what a proof would require is a non-perturbative definition of the heterotic string on T^6, and none exists. The tests cited here do not constitute such a construction.

One point about the statement itself should be made precisely, because it is the kind of place where a conjecture can be "wrong" without its author being wrong. For gauge theories with a non-simply-laced gauge group the duality group of the $\mathcal{N}=4$ theory is not $SL(2,\mathbb{Z})$: the inversion maps the gauge group G to its Langlands dual $G^\vee$, and the group generated by the shift and the inversion is a Hecke group $\Gamma_0(q)$-type subgroup, with $q=2$ or 3 [49, 31, 2]. This is not a correction of [115] or of [116]: both papers treat $SU(2)$ and the heterotic string, where the duality group is $SL(2,\mathbb{Z})$, and [116] is explicit that the test is for $SU(2)$. We mention it because the literature sometimes attributes an $SL(2,\mathbb{Z})$ conjecture "for all groups" to the 1994 papers; they contain no such statement.

3.2. **The two-monopole bound state.** The moduli space of two $SU(2)$ monopoles of total charge two is $\mathbb{R}^3 \times (S^1 \times \mathcal{M}_0)/\mathbb{Z}_2$, where $\mathcal{M}_0$ is the Atiyah–Hitchin manifold [3], a complete hyperkähler four-manifold with an $SO(3)$ action and a metric known in closed form (for the hypercomplex structures underlying such metrics see also [9]). The low energy dynamics of two monopoles is the supersymmetric quantum mechanics on this moduli space, and a bound state of two monopoles with one unit of electric charge and the right supermultiplet

[2] The conjecture was made in two steps. Schwarz and Sen [82] showed that the equations of motion of the low energy theory, not its action, are invariant under $SL(2,\mathbb{R})$, and that the charge lattice reduces the symmetry to $SL(2,\mathbb{Z})$; [115] then proposed that the full string theory, at every order and beyond perturbation theory, has this symmetry. The $SL(2,\mathbb{R})$ of the equations of motion is a theorem of the supergravity theory; it is the lift to the full string theory that is the conjecture.

structure exists if and only if the Atiyah–Hitchin manifold carries a harmonic two-form that is square integrable, anti-self-dual (with the orientation fixed by the complex structures), odd under the $\mathbb{Z}_2$, and unique.[3] Sen wrote down the form. In the coordinates of [3], with the metric $ds^2 = f^2 d\rho^2 + a^2\sigma_1^2 + b^2\sigma_2^2 + c^2\sigma_3^2$ on left-invariant forms with $d\sigma_i = \frac{1}{2}\epsilon_{ijk}\sigma_j \wedge \sigma_k$, the form is

$$\begin{aligned} \omega &= F(\rho)\,\Big(d\sigma_1 - \frac{fa}{bc}\,d\rho \wedge \sigma_1\Big), \\ F(\rho) &= F_0 \exp\Big(-\int_\pi^\rho \frac{fa}{bc}\,d\rho'\Big), \end{aligned} \tag{1}$$

which is anti-self-dual by the choice of the coefficient of $d\rho \wedge \sigma_1$, closed because $dF/d\rho = -(fa/bc)F$, and square integrable because $fa/(bc) \to \frac{1}{2}$ as $\rho \to \infty$ (where $f \simeq -1$, $a \simeq b \simeq \rho$, $c \simeq -2$) makes F decay as $e^{-\rho/2}$ while the volume grows polynomially; near the bolt $\rho \to \pi$, in the coordinates adapted to it, the form is smooth.[4]

Theorem 3.2 (Sen 1994; Hitchin 2000)**.** *The space of L^2 harmonic forms on the Atiyah–Hitchin manifold is one-dimensional. It is spanned by the two-form* (1)*, which is anti-self-dual and odd under the $\mathbb{Z}_2$ that acts on the moduli space of two monopoles.*

Proof. Existence, anti-self-duality, square integrability and the $\mathbb{Z}_2$ parity are the computation of [116] summarised above; we have checked each step. The uniqueness result is taken from [53]. We do not infer it merely by counting a circle fixed set: orientation and the particular cover of the monopole moduli space matter. □

Verdict 3.3 (claim 2)**.** Proved. The bound state predicted by $SL(2,\mathbb{Z})$ exists, and it is the only one in the two-monopole sector, which is also what the duality predicts: the $(1,2)$ state must be a single short multiplet.

3.3. **The Sen conjecture on L^2 cohomology.** The argument of [116] applies to any charge. Let $\mathcal{M}_k^0$ denote the reduced moduli space of k monopoles, of real dimension $4k-4$, and let $\mathcal{M}_k = \mathbb{R}^3 \times (S^1 \times \mathcal{M}_k^0)/\mathbb{Z}_k$. The states with k units of magnetic charge and p units of electric charge, with $\gcd(p,k) = 1$, correspond to L^2 harmonic forms on $\mathcal{M}_k^0$ transforming under the $\mathbb{Z}_k$ by $e^{2\pi i p/k}$; S-duality requires exactly one such form, in the middle degree $2k-2$, for every p coprime to k, and none for other p (which would give states not in the duality orbit of the elementary string states). This is the *Sen conjecture*; Segal and Selby [84] proved that the image of the compactly supported cohomology in the ordinary cohomology, which is a lower bound for the L^2 cohomology, has exactly the predicted dimension in the middle degree and vanishes elsewhere. The conjecture is thus reduced to the statement that every L^2 harmonic form comes from compactly supported cohomology.

Theorem 3.4 (published partial results and their hypotheses)**.** *The Segal–Selby image has the dimensions predicted by the conjecture* [84]*. Hitchin's middle-degree result and the charge-two case are established in* [53]*. Fritzsch–Kottke–Singer* [45, Theorem 1.2 and Section 6] *state the coprime electric/magnetic charge case for arbitrary magnetic charge, using their metric compactification. Their introduction announces a separate complete proof of the asymptotic metric description. Kottke–Rochon* [60, Theorem A] *give the charge-three result provided the*

[3]The electric charge of the state is the momentum along the S^1 factor, and the $\mathbb{Z}_2$ acts on it; the state with one unit of electric charge requires the form on $\mathcal{M}_0$ to be odd under the $\mathbb{Z}_2$ which combines with the half shift on S^1. The multiplet structure is fixed by the fermion zero modes: the state must be a single short multiplet of $\mathcal{N} = 4$, which is the statement that the harmonic form is unique and of a definite duality type. Sen's paper carries this out in detail and we do not repeat it.

[4]Equations (16) to (19) of [116]. The form is invariant under the identification $(\rho,\theta,\phi,\psi) \sim (\rho,\pi-\theta,\pi+\phi,-\psi)$ that defines the Atiyah–Hitchin manifold and changes sign under $\psi \mapsto \psi + \pi$, which is the $\mathbb{Z}_2$ of the two-monopole moduli space; this is the parity required for odd electric charge. The uniqueness of the form among L^2 harmonic two-forms was argued in [116] from the symmetries; the complete statement that the space of L^2 harmonic forms of $\mathcal{M}_0$ is one-dimensional is Hitchin's theorem below. Sen notes that the same form had appeared in work of Gibbons and Ruback for another purpose.

monopole metric has the announced quasi-fibered boundary structure. These hypotheses and dependencies are part of the statements, not additional results proved here.

Proof. This paragraph records the cited results with their own hypotheses. In particular it does not reduce all progress for $k \geq 4$ to a lower bound from compactly supported cohomology: the coprime case is explicitly treated in Section 6 of [45]. Conversely, quoting Theorem A of [60] without its metric qualification would overstate that theorem. □

Verdict 3.5 (claim 3)**.** Supported, with substantial established and conditional cases. No proof of the complete statement in every character is given here. The non-coprime sector and the cited metric construction dependencies must be checked separately before asserting a general unconditional closure. These are mathematical questions rather than evidence that the original conjecture is false.

3.4. **Duality-symmetric actions, the Kerr–Sen black hole and string networks.** Three further claims of this period are theorems and we record them briefly.

Proposition 3.6 (claims 4, 5 and 35)**.** *(i) Schwarz and Sen* [83] *wrote an action with two gauge potentials and a first-order Lagrangian; it is invariant under the duality rotation and has equations of motion equivalent to Maxwell's equations; the same construction gives a duality-symmetric form of the heterotic effective action in four dimensions.*

(ii) The metric, dilaton, gauge field and axion of [114] *solve the equations of motion of the low energy heterotic string in four dimensions; the solution is obtained from the Kerr solution by the $O(1,1)$ twisting transformation of the string background, which maps solutions to solutions.*

(iii) A planar network of (p,q)-strings of type IIB with the charges conserved at each junction and the angles fixed by the tensions is supersymmetric, preserving one quarter of the supersymmetry. The proof is in [121] *itself, from the supersymmetry algebra of the (p,q)-string; the three-string junction was shown to be a BPS state by* [76] *and, lifted to a holomorphic membrane in M-theory, by* [67]*. The statement was a theorem when it was made, and the catalogue records it as proved.*

Proof. (i) is a direct computation in the cited paper: the two potentials $A^{(1)}, A^{(2)}$ enter through a first-order Lagrangian whose equations of motion are $F^{(2)} = *F^{(1)}$ up to gauge, so that eliminating $A^{(2)}$ recovers the Maxwell equations; the Lorentz invariance is not manifest but is verified. (ii) is the statement that the twisting transformation of [114], which acts on backgrounds with a Killing direction, is a symmetry of the equations of motion; this was proved for the full string effective action at lowest order in α', and the Kerr–Sen metric is the image of Kerr. (iii) follows from the supersymmetry algebra: the BPS condition for a network is linear in the charge vectors at each junction, and the tension balance at a junction is the statement that the central charges add; [121] verifies this and [95] uses the same networks to count quarter-BPS dyons. For (iii), "proved" refers to the effective BPS configuration; existence and completeness of the quantum spectrum at finite coupling are separate physical questions. □

4. Quarter-BPS dyons

4.1. **The partition function and its product formula.** Dijkgraaf, Verlinde and Verlinde [37] proposed that the index of quarter-BPS dyons of the heterotic string on T^6, with electric charge Q and magnetic charge P in the lattice $\Gamma^{6,22}$, is a Fourier coefficient of the inverse of the Igusa cusp form Φ_{10} of weight ten on $Sp(2,\mathbb{Z})$:

$$d(Q,P) = (-1)^{Q\cdot P+1} \oint_{\mathcal{C}} d\rho\, d\sigma\, dv\, \frac{e^{-\pi i(\rho Q^2+\sigma P^2+2v\, Q\cdot P)}}{\Phi_{10}(\rho,\sigma,v)}, \tag{2}$$

the contour $\mathcal{C}$ being a three-torus at fixed, large imaginary parts of (ρ,σ,v), with the sign of $\operatorname{Im} v$ determined by the asymptotic moduli [97].[5] Jatkar and Sen [56] proposed the analogous formula for the CHL orbifolds $\mathbb{Z}_N$, $N=2,3,5,7$, with Φ_{10} replaced by a Siegel form Φ_k of weight $k=24/(N+1)-2$ for a subgroup of $Sp(2,\mathbb{Z})$, and David, Jatkar and Sen [34, 35] derived the product representation

$$\widetilde{\Phi}(\rho,\sigma,v)=e^{2\pi i(\rho+\sigma/N+v)}\prod_{b=0}^{1}\prod_{r=0}^{N-1}\prod_{\substack{k'\in\mathbb{Z}+r/N,\ l\in\mathbb{Z},\ j\in 2\mathbb{Z}+b\\ k',l\geq 0;\ j<0 \text{ if } k'=l=0}}\left(1-e^{2\pi i(k'\sigma+l\rho+jv)}\right)^{c(k',l,j)},$$
$$c(k',l,j)=\sum_{s=0}^{N-1}e^{2\pi isl/N}c_b^{(r,s)}(4k'l-j^2), \tag{3}$$

from the twisted elliptic genera

$$F^{(r,s)}(\tau,z)=\sum_b\sum_{j\in 2\mathbb{Z}+b}\sum_{n\in\mathbb{Z}/N}c_b^{(r,s)}(4n-j^2)\,q^n y^j$$

of K3, by the method of [37]: the dyons are counted in the type IIB frame on $K3\times T^2$ by the D1–D5 system bound to a Kaluza–Klein monopole, and the three factors of the partition function (the motion of the centre of mass, the Kaluza–Klein monopole and the D1–D5 world-volume theory) each contribute a product. For $N=1$, (3) with the K3 elliptic genus $8A(\tau,z)$ is the Borcherds product for Φ_{10}.

Theorem 4.1 (claim 6)**.** *For $N\in\{2,3,5,7\}$ the function $\widetilde{\Phi}_k$ defined by (3) is a Siegel modular form of weight k for the subgroup $\widetilde{G}\subset Sp(2,\mathbb{Z})$ of [56], and it coincides with the form constructed by Borcherds lifting in [27] and with the genus-two partition function of [26].*

Proof. The modularity of the product and its identification with the additive lift are [27, Section 3], who show that (3) is the Borcherds lift of the twisted elliptic genus and that the lift is the Siegel form of [56]; the identification with the genus-two partition function of the heterotic string is [26]. For $N=1$ the product is the classical Borcherds product of Φ_{10} [50]. We have checked (3) numerically for $N=1,2$ by reproducing the tables of [100], see Appendix A. □

The physical content of the conjecture, that this Siegel form counts the quarter-BPS states of the string theory, is not a theorem and cannot be: it is a statement about the strong coupling regime, tested through the duality orbit. The tests are the agreement with the black hole entropy to the first two orders in the large charge expansion [19, 56, 28] and at the logarithmic order (Section 11), the wall-crossing behaviour described next, and the agreement at low charges with the string-junction counting [95].

4.2. **Walls of marginal stability and the two-centred states.** The Fourier coefficient (2) depends on the contour because $1/\widetilde{\Phi}$ has double poles on the divisors $v=0$ and its images under the duality group. Sen [97] identified the poles with the walls of marginal stability of the quarter-BPS dyon, on which it can decay into a pair of half-BPS states with charges $(Q,0)$ and $(0,P)$ (and the images under duality), and computed the jump of the index across the wall $v=0$ from the residue of the double pole. Near $v=0$,

$$\widetilde{\Phi}(\rho,\sigma,v)=-4\pi^2v^2f(N\rho)\,g(\sigma/N)+O(v^4), \tag{4}$$

[5] The sign $(-1)^{Q\cdot P+1}$ converts the Fourier coefficient into the helicity trace index B_6 of the four-dimensional theory; it was identified in [97, 96] (footnote 2 of [96] records that it was absent from the earlier papers) and traced to the change of statistics between the five- and four-dimensional descriptions in the presence of a Kaluza–Klein monopole. The index d of this paper is $-B_6$ in the notation of [100].

with $f(\tau)^{-1}$ and $g(\tau)^{-1}$ the generating functions of the half-BPS electric and magnetic states; for $N = 1$, $f = g = \eta^{24}$ and $d_h(Q) :=$ coefficient of $q^{Q^2/2}$ in $q\,\eta(\tau)^{-24}$, so that $d_h = 1, 24, 324, 3200, 25650, \ldots$ for $Q^2/2 = -1, 0, 1, 2, 3, \ldots$.

Theorem 4.2 (claim 7; Sen 2007)**.** *Let $N = 1$ and let $d_>(Q,P)$, $d_<(Q,P)$ denote the Fourier coefficients of (2) without the sign $(-1)^{Q\cdot P+1}$, taken over the contours with $\operatorname{Im} v < 0$ and $\operatorname{Im} v > 0$ respectively. Then*

$$d_>(Q,P) - d_<(Q,P) = -\,Q\cdot P\; d_h(Q)\, d_h(P). \tag{5}$$

In particular the index $d = (-1)^{Q\cdot P+1} d_{\gtrless}$ changes across the wall by $(-1)^{Q\cdot P}\, Q\cdot P\, d_h(Q) d_h(P)$, whose absolute value $|Q\cdot P|\, d_h(Q) d_h(P)$ is the index of a two-centred configuration of a half-BPS electric and a half-BPS magnetic centre with angular momentum $(|Q\cdot P| - 1)/2$.

Proof. The difference of the two contour integrals is the residue at $v = 0$. From (4),

$$1/\widetilde{\Phi} = -(4\pi^2 v^2)^{-1} f(\rho)^{-1} g(\sigma)^{-1} (1 + O(v^2)),$$

and the residue of $e^{-2\pi i v Q\cdot P} v^{-2}$ at $v = 0$ is $-2\pi i\, Q\cdot P$; the remaining integrals over ρ and σ pick out $d_h(Q)$ and $d_h(P)$, and the factors of $2\pi i$ cancel against the normalisation of the contour. This is [96, eq. (2.13)]. The sign and the factor $|Q\cdot P|$ are those of a bound state of two centres carrying angular momentum $J = (|Q\cdot P| - 1)/2$ from the electromagnetic field, with $2J + 1 = |Q\cdot P|$ states [96, 30].

We have also verified (5) independently of the residue computation, by expanding the product (3) for $N = 1$ in exact arithmetic on both sides of the wall and comparing the coefficients: for every $Q^2, P^2 \in \{2, 4, 6\}$ and $|Q\cdot P| \le 4$ the two expansions differ by exactly $-Q\cdot P\, d_h(Q) d_h(P)$ (Appendix A, Table 6). □

Verdict 4.3 (claim 7)**.** Proved. The formula is a residue computation, and the identification of the jump with the two-centred bound state is confirmed by the exact match of the degeneracy $|Q\cdot P| d_h(Q) d_h(P)$ with the quantisation of the two-centred solution [96], and by the general wall-crossing formulas of Denef and Moore [30] and of Kontsevich and Soibelman [62], of which (5) is the primitive case.

4.3. **Negative discriminant states.** The discriminant $\Delta = Q^2P^2 - (Q\cdot P)^2$ must be positive for the regular classical single-centred solutions in the stated sector. The treatment of negative discriminant states uses chamber-dependent sums over two-centred decompositions and bound-state metamorphosis [99, 24]. A single primitive contribution has the form

$$(-1)^{\langle \Gamma_1, \Gamma_2\rangle + 1} |\langle \Gamma_1, \Gamma_2\rangle|\, d_h(\Gamma_1) d_h(\Gamma_2). \tag{6}$$

The full answer requires the allowed decompositions and their identifications. For example $(Q^2, P^2, Q\cdot P) = (2, 2, -2)$ has $\Delta = 0$, not negative discriminant; it must not be used as a negative-discriminant example. The identity $209952 - 648 = 209304$ is an arithmetic wall-crossing check, not a proof of a proposed microscopic decomposition. Similarly $327 = 3\cdot 109$ does not identify 109 as a half-BPS degeneracy.

Theorem 4.4 (claim 8)**.** *Every Fourier coefficient of $1/\Phi_{10}$ with negative discriminant is given by a finite sum of two-centred contributions of the form (6) over the duality images of the charges. The CHL generalisations require the separate hypotheses and formulas of [99]; no CHL formula is deduced solely from the Φ_{10} theorem.*

Proof. For Φ_{10} this is the main theorem of Chowdhury, Kidambi, Murthy, Reys and Wrase [24], who derive the exact formula for all polar (negative discriminant) coefficients from the mock modular decomposition of [36] and check it against 1650 coefficients; its direct scope is the $K3\times T^2$ theory associated with Φ_{10}, and it is not cited as a proof for every CHL form. The physical interpretation, that the states are two-centred, is [99]. □

5. The positivity conjecture

5.1. **The statement and its hypothesis.** A single-centred supersymmetric black hole in four dimensions has a spherically symmetric horizon and carries no angular momentum. The helicity trace index B_6 of a quarter-BPS state is $-\frac{1}{6!}\,\mathrm{Tr}\,[(-1)^{2h}(2h)^6]$, and for a configuration whose only degrees of freedom beyond the horizon are the twelve fermion zero modes of the broken supersymmetries, the trace evaluates to -1 times the number of horizon states; the sign is fixed and the index is negative, that is, $d = -B_6 > 0$. Sen [100] turned this into a conjecture about Fourier coefficients. In the region $\mathcal{R}$ of the axion–dilaton moduli space in which the attractor point of the charges (Q, P) lies, bounded by walls of marginal stability, the intended contribution is isolated under the exclusions described in the original argument, and the Fourier coefficient (2) taken with the contour of that region should be positive. For the heterotic string on T^6 the region is cut out by

$$\begin{gathered} Q\cdot P\ge 0, \qquad Q\cdot P\le Q^2, \qquad Q\cdot P\le P^2,\\ Q^2>0, \qquad P^2>0, \qquad Q^2P^2-(Q\cdot P)^2>0, \end{gathered} \tag{7}$$

(equation (3.7) of [100]), and for the $\mathbb{Z}_2$ CHL model, whose duality group is $\Gamma_1(2)$, by

$$\begin{gathered} Q^2>0, \qquad P^2>0, \qquad Q^2P^2>(Q\cdot P)^2,\\ 2Q^2+P^2\ge 3(Q\cdot P), \qquad 2Q^2\ge Q\cdot P,\\ P^2\ge Q\cdot P, \qquad Q\cdot P\ge 0 \end{gathered} \tag{8}$$

(equation (3.9) there, with Q^2 normalised so that $NQ^2/2\in\mathbb{Z}$). Sen verified the positivity of $d(Q,P)$ on these cones for the heterotic string and for the CHL models $N=2,3,5,7$ in the range of charges of his Tables 1 to 5, and stated the conjecture for all charges in the cone. We have reproduced his Tables 1 and 2 exactly (Appendix A).

The hypothesis of the argument is explicit in [100]: the conclusion requires "some duality frame in which only single centered black hole solution, whose only hair are the fermion zero modes associated with broken supersymmetry, contributes" to the index. In the heterotic frame the quarter-BPS black hole does have hair beyond the fermion zero modes (the hair modes identified in [12]), but there is a frame, the type IIB frame on $K3\times T^2$, in which the hair reduces to the fermion zero modes; this is what makes the conjecture consistent there, and what was proved later.

Theorem 5.1 (Bringmann–Murthy 2013)**.** *For type II string theory on $K3\times T^2$ (equivalently the heterotic string on T^6, the form $1/\Phi_{10}$) and on T^6 (the $\mathcal{N}=8$ theory), the single-centred indices in the attractor region are positive for the classes of charges treated in* [13]*; the proof uses the mock modular decomposition of the Fourier–Jacobi coefficients of $1/\Phi_{10}$* [36].

The restriction to classes of charges was removed by Rosselló [77], who bounds the tail of the Rademacher expansion of the mock Jacobi forms ψ_m^F of [36] against its leading term.

Theorem 5.2 (Rosselló 2024)**.** *For type II string theory on $K3\times T^2$, the single-centred index $d^*(m,n,\ell)$, with $m=P^2/2$, $n=Q^2/2$, $\ell=Q\cdot P$, is positive for every $m,n\ge 1$ with $4mn-\ell^2>0$: for $n\ge m$ and $0\le\ell\le m$ one has $(-1)^{\ell+1}c_m^F(n,\ell)>0$, and the general case follows by the S-duality invariance of the single-centred index.*

Verdict 5.3 (claim 9, heterotic and CHL)**.** Proved for the heterotic string on T^6, by Theorems 5.1 and 5.2; our scan of the single-centred cone up to $Q^2\le 12$, $P^2\le 6$, $Q\cdot P\le 4$ (Appendix A) agrees. For the CHL orbifolds no counterexample is known, our scan for $N=2$ finds only positive indices, while a general CHL proof is not supplied here: the Rademacher bound of [77] has not been carried out for the mock Jacobi forms of $1/\widetilde{\Phi}_k$, and [77] names this as the next case. The conjecture is proved in the model in which it was first stated and confirmed in the CHL models.

5.2. **Full and horizon indices in the toroidal orbifolds.** Sen's paper states the conjecture as a consequence of the black hole argument for "the $\mathcal{N}=4$ supersymmetric string theories", and the subsequent literature [16, 24] tested it in the full list of $\mathcal{N}=4$ compactifications with known dyon partition functions. Chattopadhyaya and David [16] found that it fails in two of them: the freely acting $\mathbb{Z}_2$ and $\mathbb{Z}_3$ orbifolds of type IIB on T^6, in which the orbifold acts by a reflection of four coordinates of T^4 (or a rotation by $2\pi/3$ of two planes) together with a shift of half (or a third) of a circle of the remaining T^2. The $\mathbb{Z}_2$ model has the twisted elliptic genera

$$\begin{aligned} F^{(0,0)} &= 0, & F^{(0,1)} &= \tfrac{8}{3}A - \tfrac{4}{3}B\,E_2(\tau), \\ F^{(1,0)} &= \tfrac{8}{3}A + \tfrac{2}{3}B\,E_2(\tfrac{\tau}{2}), & F^{(1,1)} &= \tfrac{8}{3}A + \tfrac{2}{3}B\,E_2(\tfrac{\tau+1}{2}), \end{aligned} \tag{9}$$

where $A = \sum_{i=2}^{4}\theta_i(\tau,z)^2/\theta_i(\tau,0)^2$, $B = \theta_1(\tau,z)^2/\eta(\tau)^6$, and $E_2(\tau) := 2E(2\tau) - E(\tau)$ is the weight-two form of $\Gamma_0(2)$ built from the Eisenstein series $E(\tau) = 1 - 24\sum\sigma_1(n)q^n$ (this is the E_N of [34]; it is not the quasi-modular Eisenstein series, and the two are easily confused). The Siegel form is $\widetilde{\Phi}_2 = \widetilde{\Phi}_6^2/\widetilde{\Phi}_{10}$, of weight two, with the prefactor $e^{2\pi i(\rho+v)}$ in place of $e^{2\pi i(\rho+\sigma/N+v)}$ in (3), and the index is given by (2) with $Q^2 \in \mathbb{Z}$, $P^2 \in 2\mathbb{Z}$. The single-centred region is again cut out by (8), since the duality group is again $\Gamma_1(2)$.

Theorem 5.4 (known negative full indices; finite exact test)**.** *In the $\mathbb{Z}_2$ toroidal orbifold of type IIB on T^6, the index $d(Q,P)$ computed from (2) with the contour of the attractor region is negative for infinitely many charges satisfying the single-centred conditions (8). In the range $Q^2 \le 12$, $P^2 \le 8$, $Q\cdot P \le 4$ there are exactly thirty such charge triples, listed in Table 5; the first ones are*

$$\begin{aligned} d(1,2,0) &= -224, \quad & d(1,4,0) &= -1248, \\ d(2,2,1) &= -256, \quad & d(3,2,0) &= -3392, \\ & & d(3,4,0) &= -10320, \end{aligned}$$

in the notation $d(Q^2,P^2,Q\cdot P)$. In the first column of Table 7 ($P^2=0$, outside the cone) the sign is $(-1)^{Q^2}$ for $Q^2 \le 12$; Figure 2 draws the table.

Proof. We expanded the product (3) for the genera (9) in exact rational arithmetic, with the Fourier coefficients of the theta functions, of η^{-6} and of the Eisenstein series computed from their definitions, and read off the coefficients in the convention $q_\rho = e^{2\pi i\rho}$, $q_\sigma = e^{2\pi i\sigma/N}$ and $y = e^{2\pi iv}$, keyed by $(NQ^2/2, P^2/2, Q\cdot P)$ in the expansion in which the factor $(1-e^{-2\pi iv})^{-2}$ at $k' = l = 0$ is expanded in powers of $e^{-2\pi iv}$ (the attractor region for $Q\cdot P \ge 0$). The resulting tables agree entry by entry with Tables 17, 18 and 19 of [16] in their common range. The thirty charge triples of Table 5 satisfy each inequality of (8) by inspection and have negative index. For the infinitude: the values at $P^2 = 2$, $Q\cdot P = 0$ are

$$64,\ -224,\ 1152,\ -3392,\ 11520,\ -30336,\ 83968,$$
$$-202560,\ 496512,\ -1118496,\ 2521600,\ -5374656$$

for $Q^2 = 0,1,2,\dots,11$; every odd Q^2 with $P^2=2$, $Q\cdot P=0$ lies in the cone (8), and [16, Section 4.1] show from the Fourier–Jacobi coefficient ψ_1 of $1/\widetilde{\Phi}_2$ at $P^2=2$ that the finite part of ψ_1, which is what the single-centred index is, has coefficients of alternating sign in Q^2; the alternation is visible in our table for $Q^2 \le 12$. The analogous statement for the $\mathbb{Z}_3$ orbifold is [16, Section 4.2]. □

5.3. **Which hypothesis fails.** The sign argument requires assumptions about horizon states, hair and other contributions to the trace. The toroidal models of [16] are examples where a negative full index calls for a separate hair analysis. The proposed hair factors and finite positive horizon-index tests of [17] address this distinction. They are not used here as a proof that all horizon indices in all models are positive.




FIGURE 2. The full index $d(Q^2, P^2, 0)$ of the $\mathbb{Z}_2$ toroidal orbifold from Table 7, drawn on the scale $\mathrm{sign}(d)\log_{10}(1+|d|)$. Every bar in burnt orange is a negative full index; along $P^2 = 2$ the sign alternates with Q^2, and every odd Q^2 lies inside the cone (8) of Theorem 5.4. A horizon index requires the separate hair analysis.

Q^2	P^2	$Q\cdot P$	d	Q^2	P^2	$Q\cdot P$	d
1	2	0	-224	6	2	1	-33536
1	4	0	-1248	7	2	0	-202560
1	6	2	-224	7	2	2	-30336
1	8	2	-1248	7	4	0	-179022
2	2	1	-256	7	4	4	-10320
2	6	3	-256	8	2	1	-220672
3	2	0	-3392	9	2	0	-1118496
3	2	2	-224	9	2	2	-202560
3	4	0	-10320	9	4	0	-155232
3	6	4	-224	9	4	4	-55424
4	2	1	-3840	10	2	1	-1204992
5	2	0	-30336	11	2	0	-5374656
5	2	2	-3392	11	2	2	-1118496
5	4	0	-55424	11	4	4	-179022
5	4	4	-1248	12	2	1	-5741824

TABLE 5. The thirty negative full-index triples in the linear cone (8), for $0 < Q^2 \le 12$, $0 < P^2 \le 8$ and $0 \le Q\cdot P \le 4$. The exact Python scan covers this entire box. The separate Lean certificate covers the original twenty-one listed triples with $P^2 \le 4$ and $Q\cdot P \le 2$; it does not certify all thirty entries.

Proposition 5.5 (conditional scope of the sign argument)**.** *For a charge with negative full index, the conjunction of the assumptions leading to a positive full index cannot hold. If the horizon degeneracy is positive and the only external contribution is the universal fermion-zero-mode factor with its prescribed sign, positivity follows. A negative full index therefore rules out this conjunction; by itself it does not identify which physical assumption fails or prove the nonexistence of every conceivable duality frame.*

Proof. The last implication is the contrapositive of the sign argument. Removing hair changes the generating function, not a numerical overall convention. In the toroidal models the proposed hair partition functions and finite positive horizon-index tests are given in [17]. Their interpretation is evidence for nontrivial hair, rather than a classification of all frames proved here. □

Verdict 5.6 (claim 9, generalisation across models)**.** Scope limited. Negative full indices in the toroidal orbifolds are known results of [16]. They test an unrestricted extension of positivity, while [100] explicitly states its hair and single-centre assumptions. The later analysis [17] already addresses the horizon/full-index distinction. We therefore do not count this as a newly discovered false proof by Sen. The computations here reproduce and extend a finite table; the physical identification of horizon states remains subject to the cited assumptions.

Verdict 5.7 (claim 10)**.** Confirmed. The hair modes of the quarter-BPS black hole were identified in the type IIB frame for $K3\times T^2$ in [12] and for every CHL model and both toroidal orbifolds in [17], and in each case the remaining horizon index is positive. The identification rests on the classification of the supersymmetric deformations of the black hole solution that are supported outside the horizon, within the effective descriptions used in those papers; a full string-spectrum classification is not established by the finite computations here.

Verdict 5.8 (claim 36)**.** Confirmed. The companion of the positivity conjecture is the statement of [21] that every microstate of a single-centred BPS black hole, once the fermion zero modes of broken supersymmetry are removed, carries zero angular momentum, so that the index counts the states with sign $+1$ and positivity follows. It was tested in [21] on the quarter-BPS states of type IIB on T^6 at low charges by an explicit enumeration of the supersymmetric configurations of D-branes, and in [51] on multi-centred configurations with pure D-brane charges, and holds in every case computed. It is not proved, and the toroidal orbifolds show what its proof must contain: there the index of the full dyon is negative, so either some microstates carry angular momentum or the hair modes do, and [17] locates the sign in the hair. Later tests at higher charges [22, 23] reproduce the index of the pure D-brane system by algebraic geometry and find no state of nonzero angular momentum. Positivity of the index, now a theorem for $K3\times T^2$ (Theorem 5.2), does not imply the statement, as [77] notes: an index that is positive can still count fermionic states. The sources examined here do not supply a general proof; the statement for the horizon states remains open and consistent with every test.

Remark 5.9. The numbers of Theorem 5.4 are the index $d=-B_6$ with the sign $(-1)^{Q\cdot P+1}$ included and without the factor $1/N$ written in front of the contour integral in [16]: the integral over σ runs over a period of length N and produces a factor N that cancels it. The same normalisation reproduces Sen's Table 2 for the CHL model, which fixes it.

6. The tachyon conjectures

6.1. **Universality.** An unstable D-brane of bosonic string theory, or a brane–antibrane pair or non-BPS brane of type II, carries an open string tachyon. Sen's conjectures about it [119, 124] are three: the tachyon potential has a minimum at which the negative energy density exactly cancels the tension of the brane; the lump and kink solutions of the tachyon field are the D-branes of lower dimension; and around the minimum there are no perturbative open string states, since there is no brane for them to end on. The first step towards testing them was the observation [125] that the tachyon potential is universal.

Theorem 6.1 (claim 11; Sen 1999)**.** *In the cubic open string field theory of Witten on a D-brane described by a boundary conformal field theory* BCFT_0*, the subspace* $\mathcal{H}_{\mathrm{univ}}$ *of the string field spanned by the states obtained from the* $SL(2,\mathbb{R})$*-invariant vacuum by the matter Virasoro generators and the ghost oscillators is closed under the star product and under the BRST operator, and the action restricted to it is* $\mathcal{T}_p V_{\mathrm{univ}}$*, where* $\mathcal{T}_p$ *is the tension of the brane and* V_{univ} *does not depend on* BCFT_0*. The same holds for the Berkovits field theory of the brane–antibrane pair and of the non-BPS brane, with the GSO-odd and GSO-even sectors.*

Proof. This is [125, Sections 2 and 3]. The matter part of the action involves only correlators of the matter Virasoro generators and the identity on the disc, which are fixed by the central charge $c = 26$ (or $c = 15$) and the conformal Ward identities, and the ghost part is independent of the background; the normalisation of the disc partition function is the tension. Closure of $\mathcal{H}_{\mathrm{univ}}$ under the star product follows from the fact that the star product is a conformal map of three half-discs into one, which maps Virasoro descendants of the identity to Virasoro descendants of the identity. The equations of motion restricted to $\mathcal{H}_{\mathrm{univ}}$ are therefore the equations of motion in it, and a solution in $\mathcal{H}_{\mathrm{univ}}$ is a solution of the full theory. □

Verdict 6.2 (claim 11)**.** Proved. The universality is used in every subsequent analytic solution: Schnabl's tachyon vacuum lies in $\mathcal{H}_{\mathrm{univ}}$, and so does the solution of Erler in the superstring.

6.2. **The three conjectures.** Let Ψ be the open string field of the cubic theory on a Dp-brane with action $S = -\frac{1}{g^2}\big[\frac{1}{2}\langle\Psi, Q_B\Psi\rangle + \frac{1}{3}\langle\Psi, \Psi*\Psi\rangle\big]$, normalised so that the tension of the brane is $\mathcal{T}_p = 1/(2\pi^2 g^2)$.

Theorem 6.3 (claims 12, 13, 14)**.** *(i) There is a solution* $\Psi_0 \in \mathcal{H}_{\mathrm{univ}}$ *of the equation of motion* $Q_B\Psi + \Psi*\Psi = 0$ *whose action is* $S[\Psi_0] = \mathcal{T}_p \,\mathrm{Vol}$*, that is, whose energy density is* $-\mathcal{T}_p$ [81].

(ii) The kinetic operator around Ψ_0*,*

$$Q_{\Psi_0}\Phi = Q_B\Phi + \Psi_0*\Phi - (-1)^{|\Phi|}\Phi*\Psi_0,$$

has trivial cohomology in every ghost number: there is a string field A *with* $Q_{\Psi_0}A = 1$*, the identity of the star algebra* [39].

(iii) For every boundary conformal field theory BCFT_* *with the same bulk theory as* BCFT_0 *and admitting boundary condition changing operators between the two of weight zero with regular operator products, there is a solution* Ψ_* *of the equation of motion on the Dp-brane whose energy is that of* BCFT_* *and whose fluctuation spectrum is that of* BCFT_**; the tachyon lump describing a* $D(p-1)$*-brane at any compactification radius is of this form* [41].

Proof. (i) is the solution of [81] in the gauge $\mathcal{B}_0\Psi = 0$, with the energy computed there and confirmed by [72] and by [46], who checked that the equation of motion holds when contracted with the solution itself; the simpler solution of [42], $\Psi = \frac{1}{\sqrt{1+K}}\, c(1+K)Bc\, \frac{1}{\sqrt{1+K}}$ in the KBc algebra, has the same energy by a two-line computation. (ii) is [39]: for the Erler–Schnabl solution the homotopy operator is $A = B/(1+K)$, and $Q_\Psi A = 1$ is an identity of the KBc algebra. (iii) is [41, Section 4]: the solution is $\Psi_* = \Psi_{\mathrm{tv}} - \Sigma\,\Psi_{\mathrm{tv}}\,\bar\Sigma$ with $\Sigma, \bar\Sigma$ built from the boundary condition changing operators and a timelike Wilson line, and $\bar\Sigma\Sigma = 1$; the energy is the disc partition function of BCFT_* and the cohomology of Q_{Ψ_*} is isomorphic to that of BCFT_*. The hypothesis on the boundary condition changing operators is satisfied by the lump at every radius, by the Wilson line construction of the same paper; the numerical lumps of [63] agree. □

Verdict 6.4 (claims 12, 13, 14)**.** Proved. Each statement is a theorem of cubic open string field theory; the sense in which it is a theorem about string theory depends on cubic string

field theory being a complete description of the open string, which is the framework of the conjectures and is not in question in them.

Theorem 6.5 (claim 15; Erler 2013)**.** *In the open superstring field theory of Berkovits on a non-BPS D-brane there is an analytic solution for the tachyon vacuum, built from 2×2 matrices in two GL_2 subgroups of the star algebra, whose action equals the tension of the non-BPS brane, so that the energy density of the vacuum is $-\mathcal{T}_p$.*

Proof. [40]. The energy is computed in closed form from the nonpolynomial action. □

Verdict 6.6 (claim 15)**.** Proved for the energy. The absence of open string states around the superstring vacuum follows from the homotopy operator of the cubic reformulation of the same solution, and the lower-dimensional branes in the superstring are not yet covered by an analytic solution; the numerical results of [15], the level truncation to level $(2,4)$ with 85% of the tension, were the state of the conjecture before [40]. The energy, fluctuation cohomology, and lower-brane solutions are separate assertions; no comprehensive negative claim about all later constructions is made here.

7. The rolling tachyon and tachyon matter

7.1. **Homogeneous decay.** Sen [88] described the time-dependent decay of an unstable Dp-brane by the boundary conformal field theory with the marginal deformation $\lambda\int\cosh X^0$ of the free boson X^0 on the boundary of the world-sheet, and computed the energy-momentum tensor from the boundary state. In the bosonic string, for $0<\lambda\le\frac12$,

$$\begin{aligned} T_{00} &= \tfrac12\mathcal{T}_p(1+\cos 2\pi\lambda), \qquad T_{ij} = -\delta_{ij}\,\mathcal{T}_p\,f(x^0),\\ f(x^0) &= \frac{1}{1+e^{x^0}\sin\pi\lambda}+\frac{1}{1+e^{-x^0}\sin\pi\lambda}-1, \end{aligned} \tag{10}$$

and in the superstring the same with x^0 replaced by $\sqrt{2}x^0$ and $\sin\pi\lambda$ by $\sin^2\pi\lambda$.

Theorem 7.1 (claim 16; Sen 2002)**.** *The energy density* (10) *is conserved, and the pressure $-\mathcal{T}_p f(x^0)$ tends to zero as $x^0\to\pm\infty$, as $e^{-|x^0|}$ in the bosonic string and $e^{-\sqrt{2}|x^0|}$ in the superstring. The endpoint $\lambda=0$ is the undeformed brane with constant nonzero pressure and is excluded from the decay assertion. For $\lambda=\frac12$ the energy density vanishes and the solution is the tachyon vacuum at all times.*

Proof. The boundary state is the product of the Dirichlet boundary state for the spatial directions, the $c=1$ boundary state of the $\cosh X^0$ deformation, obtained by Wick rotation of the $SU(2)$ rotation of the Neumann boundary state at the self-dual radius, and the ghost boundary state; $T_{\mu\nu}$ is read off from the coefficient of the closed string graviton. As $x^0\to+\infty$ the first term of f is $O(e^{-x^0})$ and the second is $1-\sin\pi\lambda\,e^{-x^0}+O(e^{-2x^0})$, so $f=O(e^{-x^0})$; the case $x^0\to-\infty$ is symmetric. Conservation is $\partial_0 T_{00}=0$, immediate from the form of T_{00}. This is [88, Section 3]; the computation is exact in α' and at tree level in the string coupling. □

Verdict 7.2 (claim 16)**.** Proved. The statement is a theorem of boundary conformal field theory. The open question is its physical scope, treated in Section 7.3.

7.2. **Tachyon matter and the caustic theorem.** Sen [87] called the asymptotic state of (10), with constant energy density and vanishing pressure, *tachyon matter*, and [86] proposed an effective field theory for the tachyon near the minimum of its potential whose solutions are this matter:

$$\begin{aligned} S &= -\int d^{p+1}x\; V(T)\sqrt{1+\eta^{\mu\nu}\partial_\mu T\partial_\nu T},\\ \eta &= \mathrm{diag}(-1,1,\dots,1), \qquad V(T)\to 0 \text{ as } T\to\infty. \end{aligned} \tag{11}$$

The late-time pressure statement in [87] concerns the homogeneous rolling solution. The field theory in [86] also has the exact dust solution $T = t$, $\Pi = f(x)$, with arbitrary time-independent nonnegative density f. Thus a nonconstant density alone does not imply focusing. Caustics arise from a compressive velocity gradient. The calculation below determines when that happens in the pressureless equations; it does not refute the existence or the homogeneous asymptotics claimed in those papers.

Near the minimum, $V \to 0$, the equation of motion of (11) forces $\eta^{\mu\nu}\partial_\mu T\partial_\nu T \to -1$ wherever the energy density stays finite [86, Section 2], and the energy-momentum tensor takes the form

$$T^{\mu\nu} = \rho\, u^\mu u^\nu, \qquad u^\mu = -\eta^{\mu\nu}\partial_\nu T, \qquad u^\mu u_\mu = -1, \tag{12}$$

of a pressureless fluid with velocity u^μ and rest density $\rho = V(T)/\sqrt{1+\partial T\cdot\partial T} \geq 0$. We write $W = \partial_t T = \sqrt{1+|\nabla T|^2}$ for the branch in which T increases with time (the other branch is obtained by $T \mapsto -T$), $v = -\nabla T/W$ for the three-velocity u^i/u^0 of the fluid, and $\varepsilon = T^{00} = \rho W^2$ for the energy density.

Lemma 7.3. *Let T be a C^2 solution of $\eta^{\mu\nu}\partial_\mu T\partial_\nu T = -1$ on $[0,t_1)\times\mathbb{R}^p$ with $\partial_t T > 0$, and let $\rho \geq 0$ be C^1 with $\partial_\mu(\rho u^\mu) = 0$. Then:*

(i) $u^\mu\partial_\mu u^\nu = 0$: the integral curves of u, equivalently the curves $\dot X = v(t,X)$, are straight lines $X(t;x_0) = x_0 + t\,v_0(x_0)$, $v_0 = v(0,\cdot)$, along which v is constant;

(ii) $\partial_t\varepsilon + \partial_i(\varepsilon v^i) = 0$, and along the lines of (i)

$$\varepsilon\big(t, X(t;x_0)\big)\det\big(I + t\,Dv_0(x_0)\big) = \varepsilon(0,x_0);$$

(iii) $Dv_0 = -W^{-1}P\,H$ with $H = \nabla^2 T(0,\cdot)$ the Hessian of the initial profile and $P = I - \nabla T\nabla T^{\mathsf T}/W^2$ symmetric and positive definite; the eigenvalues of $Dv_0(x_0)$ are real, and $Dv_0(x_0)$ has a negative eigenvalue if and only if $H(x_0)$ has a positive eigenvalue.

Proof. (i) Since $u_\mu = -\partial_\mu T$ is a gradient,

$$u^\mu\partial_\mu u_\nu = u^\mu\partial_\nu u_\mu = \tfrac12\partial_\nu(u^\mu u_\mu) = 0.$$

In components, with $u^0 = W$ and $u^i = Wv^i$, the equations $u^\mu\partial_\mu W = 0$ and $u^\mu\partial_\mu(Wv^i) = 0$ give $(\partial_t + v\cdot\nabla)v = 0$, so v is constant along the curves $\dot X = v$, which are therefore straight. (ii) $\partial_\mu(\rho u^\mu u^0) = u^0\partial_\mu(\rho u^\mu) + \rho u^\mu\partial_\mu u^0 = 0$ is the continuity equation for $\varepsilon = \rho W^2$ with flux εv. Writing it as $(\partial_t + v\cdot\nabla)\varepsilon = -\varepsilon\,\nabla\cdot v$ and using

$$\nabla\cdot v(t, X(t;x_0)) = \tfrac{d}{dt}\ln\det\big(I + tDv_0(x_0)\big),$$

which holds because the Jacobian $D_{x_0}X = I + tDv_0$ satisfies $\frac{d}{dt}D_{x_0}X = (Dv)(D_{x_0}X)$, gives

$$\tfrac{d}{dt}\big[\varepsilon\det(I + tDv_0)\big] = 0$$

along the line. (iii) Differentiating $v^i = -\partial_i T/W$ and $W^2 = 1 + |\nabla T|^2$ gives $\partial_j v^i = -W^{-1}(\delta_{ik} - \partial_i T\partial_k T/W^2)\partial_k\partial_j T$. The matrix P has eigenvalue 1 on the orthogonal complement of ∇T and $1 - |\nabla T|^2/W^2 = W^{-2} > 0$ along it. PH is similar to $P^{1/2}HP^{1/2}$, which is symmetric, so its eigenvalues are real, and Sylvester's law gives the same inertia as H. The additional negative scalar in $Dv_0 = -W^{-1}PH$ reverses positive and negative signs. □

Theorem 7.4 (forward focusing in the dust equations)**.** *Let $T_0 \in C^3(\mathbb{R}^p)$, $v_0 = -\nabla T_0/\sqrt{1+|\nabla T_0|^2}$, and ε_0 be a positive C^1 initial density. Put $A(x) = Dv_0(x)$ and assume A is bounded. Its eigenvalues are real. Define*

$$M = \sup_x \max\big(\{0\}\cup\{-\lambda : \lambda\in\operatorname{spec}A(x)\}\big),$$
$$t_* = \begin{cases} 1/M & M > 0,\\ +\infty & M = 0. \end{cases} \tag{13}$$

Then the following statements hold.




FIGURE 3. Exact relativistic dust characteristics and density for $T_0 = T_* + a\cos kx$, $a = 1/2$, $k = 1$, before $t_c = 1/(ak^2)$. The velocity is $ak\sin(kx)/\sqrt{1 + a^2k^2\sin^2(kx)}$. The surface is capped for display; the formula is not capped.

(i) The characteristic map is $X_t(x) = x + tv_0(x)$, with $DX_t = I + tA(x)$. Before its first loss of invertibility,

$$\varepsilon(t, X_t(x)) = \frac{\varepsilon_0(x)}{\det(I + tA(x))}.$$

For $0 \le t < t_*$ the map is a global diffeomorphism.

(ii) If T_0 is concave, all eigenvalues of A are nonnegative; no forward focal point occurs and the density along each characteristic is nonincreasing. If T_0 has a positive Hessian eigenvalue at some point, a forward focal time is finite on that characteristic.

(iii) If $0 < M < \infty$ is attained at x_*, the first singular Jacobian occurs at $t_* = 1/M$. Its determinant tends to zero from above along x_* and $\varepsilon(t, X_t(x_*)) \to +\infty$. Without attainment, t_* is only the infimum of the local focal times; a singular characteristic at exactly t_* and a density blow-up there are not asserted.

(iv) In one dimension, $v_0' = -T_0''/(1 + T_0'^2)^{3/2}$. Hence compression occurs precisely where $T_0'' > 0$. Every bounded nonconstant C^3 profile on $\mathbb{R}^p$ has a positive Hessian eigenvalue somewhere and therefore a finite local focal time. A periodic profile also attains its finite maximum compression.

Proof. Lemma 7.3 gives (i) locally. The similarity of PH to the symmetric matrix $P^{1/2}HP^{1/2}$ and the factor $-W^{-1}$ show that a negative eigenvalue of A corresponds to a positive eigenvalue of H, proving (ii). For $t < t_*$ every eigenvalue factor $1 + t\lambda$ is positive. Also $|v_0| \le 1$, so $|X_t(x)| \ge |x| - t$ and X_t is proper. A proper local diffeomorphism $\mathbb{R}^p \to \mathbb{R}^p$ is a covering onto the connected target: its image is both open and closed. Since the target is simply

connected and the source connected, it is a global diffeomorphism. This proves the global assertion in (i).

At an attained maximiser, some $\lambda = -M$ and all remaining factors are positive for $t < t_*$. Their product approaches zero, while the numerator stays positive, proving (iii). If no maximiser exists, this argument only gives a sequence of local focal times approaching t_*; passing to a fixed singular characteristic would be unjustified. For (iv), differentiate the displayed velocity. A C^2 function with no positive Hessian eigenvalue anywhere is concave. A concave function on all of $\mathbb{R}^p$ bounded below is constant, by restriction to lines; hence a bounded nonconstant profile cannot have that property. Periodicity supplies compactness for attainment. The dust theory loses a classical single-valued velocity field at focusing; higher-derivative or string effects at that point are outside this theorem. □

Verdict 7.5 (claim 17)**.** Scope limited. Homogeneous rolling has the pressureless asymptotic in the original papers. Inhomogeneous effective dust may develop caustics, as already analysed in [44, 43, 4]. The theorem above states the sign criterion and attainment conditions precisely. It neither gives the string-theoretic continuation through a caustic nor disproves Sen's homogeneous result. The stationary solution $T = t$, $\Pi = f(x)$ remains a counterexample to any assertion that every spatially varying density must focus.

7.3. **Closed string emission and open–closed duality.** The decaying brane couples to closed strings, and the question whether the classical open string description of Theorem 7.1 survives their emission was raised in [88] and answered in part by Lambert, Liu and Maldacena [66]: the energy radiated into closed strings by a decaying Dp-brane is infinite at leading order for $p \leq 2$ and finite for $p > 2$; for $p = 0$ the divergence, cut off at energies of order $1/g_s$, carries away an energy of the order of the mass of the brane, which signals the need to control backreaction before inferring an exact energy partition. Sen [90] turned this into the conjecture of *open–closed duality at tree level*: the tree-level open string theory on the decaying brane, with its tachyon matter, is a dual description of the closed string state produced by the decay, so that the two are not to be added.

Verdict 7.6 (claim 18)**.** Confirmed. The conjecture is supported by the computation of [66], in which the closed string state produced is determined by the open string boundary state and its energy, for $p = 0$, is the energy of the brane; by the matching of the energy-momentum tensor of the emitted closed strings with that of tachyon matter in the regime where both can be computed [90]; and by the two-dimensional string theory, where the decay of the D0-brane is the motion of a single fermion of the matrix model and the open and closed descriptions are provably the same [91]. A general proof in critical string theory is not supplied by the sources examined here, and the missing step is a control of the backreaction of the emitted closed strings on the open string dynamics beyond leading order in g_s. The conjecture is stated for homogeneous decay; after Theorem 7.4 the inhomogeneous case has no classical open string side to be dual to beyond t_c.

8. Non-BPS branes and the descent relations

The tachyon conjectures of Section 6 have a static counterpart: the solitons of the tachyon field are themselves D-branes. Sen [119] identified the vortex on a Dp–$\overline{\text{D}}p$ pair of type II with the BPS D$(p-2)$-brane, [122] identified the kink on a D1–$\overline{\text{D}}$1 pair of type I with a stable non-BPS D-particle transforming as a spinor of $SO(32)$, the state required by the duality with the heterotic string, and [123] organised these into the *descent relations*: the non-BPS Dp-brane of type II is the orbifold of the Dp–$\overline{\text{D}}p$ pair by $(-1)^{F_L}$, and the BPS D$(p-1)$-brane is the kink on the non-BPS Dp-brane.

Theorem 8.1 (claims 19, 20)**.** *(i) At the critical radius $R = \sqrt{2}$ (in units $\alpha' = 1$) of a circle transverse to the brane pair, the tachyon kink on a Dp–$\overline{\text{D}}p$ pair of type II wrapped on the circle is an exactly marginal boundary deformation, and the boundary conformal*

field theory at the end of the deformation is that of a non-BPS $D(p-1)$-brane; the kink on the non-BPS Dp-brane at $R=1$ is, in the same way, the BPS $D(p-1)$-brane.

(ii) The conserved charges of D-branes and of their tachyon solitons are classified by the K-theory of spacetime: $K(X)$ for type IIB, $K^{-1}(X)$ for type IIA, $KO(X)$ for type I; the kink on the type I D1–$\overline{\text{D}}$1 pair carries the class generating $KO^{-1}(\text{pt})=\mathbb{Z}_2$ and is the $SO(32)$ spinor, stable and non-BPS.

Proof. (i) is the construction of [122, 123]: the tachyon of the wrapped pair at the critical radius has a vertex operator $\cos(X/\sqrt{2})\otimes\sigma_1$ which is exactly marginal, the deformation by it can be followed to its end by fermionising the boson X, and the boundary state at the end is the boundary state of the lower brane with Dirichlet condition on X. (ii) is Witten's theorem [129] (for equivariant classes of this kind in a group-theoretic setting see [8]), which starts from Sen's identification of the brane–antibrane system with a pair of bundles (E,F) and the tachyon with a map between them and identifies the charge with the class of (E,F) in K-theory; the type I computation $KO^{-1}(\text{pt})=\mathbb{Z}_2$ gives the $\mathbb{Z}_2$ charge of the spinor, which prevents its decay. □

Verdict 8.2 (claims 19, 20)**.** Proved. Both statements are theorems of boundary conformal field theory and of K-theory respectively; the existence of the spinor as a stable non-BPS state was confirmed by its mass at weak coupling, which matches the heterotic spinor's at strong coupling up to the expected corrections, and by the identification of the stable non-BPS states of other orbifolds and orientifolds by the same method [124].

Proposition 8.3 (claim 21; Sen 2003)**.** *In the effective action (11) on a non-BPS Dp-brane, the kink solution $T(x)=\lim_{a\to\infty} a\,x$ (a step of infinite slope) is singular as a field and regular as a source: its energy-momentum tensor is $\mathcal{T}_{p-1}$ times a delta function at $x=0$, and the effective action of its collective coordinates and world-volume gauge field is exactly the Dirac–Born–Infeld action of a BPS $D(p-1)$-brane, to all orders in the fields and for every potential V with $\int V\,dT$ finite. The same holds for the vortex on the brane–antibrane pair.*

Proof. [89], Sections 3 and 4: for the ansatz $T=f(a(x-\xi(y)))$ with f odd and increasing, in the limit the action evaluated on the ansatz is independent of the profile f and of the scale a, by the change of variables $T=f(\cdot)$, and equals $-\mathcal{T}_{p-1}\int\sqrt{-\det(g_{\alpha\beta}+F_{\alpha\beta})}$ with the induced metric of the embedding ξ; the equations of motion of the ansatz follow from the full equations because the action is stationary in f and a. □

Verdict 8.4 (claim 21)**.** Proved, within the effective action. The statement is about (11) and is exact there; it says nothing about the string theory corrections to (11), which are not controlled on the kink, where second derivatives are large. The agreement of the tension and of the form of the world-volume action with those of the $\text{D}(p-1)$-brane is the content of the claim and holds.

9. The Sen limit of F-theory

Sen [117] showed that F-theory on an elliptic K3 surface, at the point of its moduli space where the Weierstrass model $y^2=x^3+f(z)x+g(z)$ has constant modulus τ and four D_4 singular fibres, is the type IIB orientifold on $T^2/\mathbb{Z}_2$ with four orientifold seven-planes and four D7-branes on each, with gauge group $SO(8)^4$ and constant coupling; and that moving away from this point, in a one-parameter family $f=-3h^2+\varepsilon\eta$, $g=-2h^3+\varepsilon h\eta-\frac{1}{12}\varepsilon^2\chi$ with $\varepsilon\to0$, the coupling becomes weak away from the loci $h=0$, which are the orientifold planes, while the discriminant $\Delta=-9\varepsilon^2h^2(\eta^2-h\chi)+O(\varepsilon^3)$ places the D7-branes on $\eta^2=h\chi$. In [118] the same expansion was applied to F-theory on elliptic Calabi–Yau threefolds and fourfolds over a base B, with h,η,χ sections of $K_B^{-2},K_B^{-4},K_B^{-6}$: the statement is that F-theory on such a Calabi–Yau has a weak coupling limit which is type IIB on the double

cover of B branched along $h = 0$, with the orientifold projection, O7-planes on $h = 0$ and D7-branes on $\eta^2 = h\chi$.

Theorem 9.1 (scope of the weak-coupling expansion)**.** *For $f = -3h^2 + \varepsilon\eta$ and $g = -2h^3 + \varepsilon h\eta - \varepsilon^2\chi/12$, one has*

$$\begin{aligned} 4f^3 + 27g^2 = & -9\varepsilon^2 h^2(\eta^2 - h\chi) \\ & + \varepsilon^3(4\eta^3 - \tfrac{9}{2}h\eta\chi) + \tfrac{3}{16}\varepsilon^4\chi^2. \end{aligned} \tag{14}$$

At points with $h(\eta^2 - h\chi) \neq 0$, the modular invariant $j \propto f^3/(4f^3 + 27g^2)$ diverges as ε^{-2}. This is a generic-point limit, not a smooth elliptic fibration at $\varepsilon = 0$: its discriminant vanishes identically there. The perturbative orientifold and D7 interpretation is the construction in [117, 118]*. The generic O7 monodromy is $-T^{-4}$ in the convention of* [118]*; the enhanced O7 with four coincident D7-branes has combined monodromy $-I$.*

Proof. The discriminant identity follows by expansion; the leading nonzero coefficient gives the asserted growth. The two monodromies concern different configurations: adding the four D7 monodromies T^4 cancels the T^{-4} factor. A Kodaira type cannot be read from the everywhere-degenerate central fibre as though it were an ordinary smooth family. □

Verdict 9.2 (claim 22)**.** Established in the stated geometric and physical framework. The constant-coupling $SO(8)^4$ model and the generic weak-coupling family are related constructions, not identical fibre configurations.

Proposition 9.3 (base divisor and double-cover divisor)**.** *Suppose h, η are local coordinates near $h = \eta = 0$ on a smooth base and χ is nonzero there. The divisor $D_B = \{\eta^2 - h\chi = 0\}$ on the base is smooth at that point. On the smooth orientifold double cover $X = \{\xi^2 = h\}$ its pullback is*

$$D_X = \{\eta^2 - \xi^2\chi = 0\},$$

which is singular along $\xi = \eta = 0$. Where χ is a unit it has two local branches. At a transverse triple locus $\xi = \eta = \chi = 0$ in a three-dimensional cover, its local equation is the Whitney pinch form $u^2 = v^2 w$.

Proof. For $F = \eta^2 - h\chi$, at $h = \eta = 0$ one has $dF = -\chi\, dh \neq 0$. On the cover, the three derivatives of $G = \eta^2 - \xi^2\chi$ in these coordinates are $2\eta, -2\xi\chi, -\xi^2$, all zero on the stated double locus. A unit χ has a local analytic square root, so $G = (\eta - \xi\sqrt{\chi})(\eta + \xi\sqrt{\chi})$. The transverse triple locus gives the pinch form. This is the distinction used in the D7 analysis of [18]. □

Proposition 9.4 (a non-Higgsable divisor is not a sufficient obstruction)**.** *In a local chart let $s = 0$ be a divisor and take*

$$h = sh_1, \qquad \eta = s^2\eta_2, \qquad \chi = s^3\chi_3.$$

Then $f = s^2 F_\varepsilon$, $g = s^3 G_\varepsilon$, with

$$F_\varepsilon = -3h_1^2 + \varepsilon\eta_2, \quad G_\varepsilon = -2h_1^3 + \varepsilon h_1\eta_2 - \varepsilon^2\chi_3/12.$$

If the coefficients are generic units, the divisor has vanishing orders $(2, 3, 6)$ for nonzero ε and j on it grows as ε^{-2} when $h_1(\eta_2^2 - h_1\chi_3) \neq 0$. Thus an I_0^* divisor does not itself force strong coupling in the limit.

Proof. The common factor s^6 cancels in $f^3/(4f^3 + 27g^2)$, leaving Theorem 9.1 with h_1, η_2, χ_3. This is a local calculation, not a global existence theorem on an arbitrary base. The allowed I_0^* case is explicitly distinguished from exceptional-fibre obstructions in [131]. □

Verdict 9.5 (claim 23)**.** Scope limited, with no false smoothness claim attributed to Sen here. The original paper calls the orientifold reduction formal and discusses singular auxiliary spaces. Its formula must not be restated as a theorem that every base admits a smooth

global perturbative orientifold. The singularity in Proposition 9.3 is on the cover, and non-Higgsability alone is insufficient to exclude weak coupling. The global obstructions of [131] require their additional fibre and section conditions. These known qualifications replace an unrestricted statement; they are not a newly discovered contradiction in Sen's expansion.

10. Black hole entropy: the entropy function and small black holes

10.1. **The entropy function.** For an extremal black hole in a theory of gravity with covariant higher-derivative terms in the framework specified below, whose near-horizon geometry has the symmetry $SO(2,1)\times SO(D-1)$ of $AdS_2\times S^{D-2}$ with electric fields e_i and magnetic charges p_a, Sen [93] defined the entropy function

$$\begin{aligned}\mathcal{E}(\vec{u},\vec{v},\vec{e},\vec{q},\vec{p}) &= 2\pi(e_i q_i - f(\vec{u},\vec{v},\vec{e},\vec{p})),\\ f &= \int_{S^{D-2}} \sqrt{-\det g}\,\mathcal{L},\end{aligned} \tag{15}$$

where f is the Lagrangian density integrated over the sphere on the near-horizon ansatz with radii v_1, v_2 of AdS_2 and S^{D-2} and constant scalars u_s.

Theorem 10.1 (claims 24, 25)**.** *(i) The equations of motion of the near-horizon geometry are the conditions*

$$\partial\mathcal{E}/\partial u_s = \partial\mathcal{E}/\partial v_{1,2} = 0, \qquad \partial\mathcal{E}/\partial e_i = 0,$$

the last being the definition of the charges q_i, and the Wald entropy of the black hole is the value of $\mathcal{E}$ at the extremum [93].

(ii) For rotating extremal black holes, whose near-horizon geometry has the symmetry $SO(2,1)\times U(1)$ and depends on the polar angle through functions $\Omega(\theta),\psi(\theta),\dots$, the same holds with $\mathcal{E}$ a functional of these functions and with an additional term $2\pi\alpha J$ for the angular momentum, the extremisation being over the functions as well as the constants [1].

Proof. The derivation, including the normalization and curvature-variation convention, is given in [93, Section 2] and its rotating extension in [1]. For a covariant Lagrangian with no derivatives of curvature, the Wald expression is $-2\pi\int_H(\partial\mathcal{L}/\partial R_{\mu\nu\rho\sigma})\epsilon_{\mu\nu}\epsilon_{\rho\sigma}$ with the induced volume factor and $\epsilon_{\mu\nu}\epsilon^{\mu\nu}=-2$. Curvature derivatives require the corresponding variational derivative, and Chern–Simons terms require separate treatment. The near-horizon equations and the Legendre transform give the entropy function in this framework. Local extremisation does not by itself construct a regular solution extending to prescribed asymptotic data. □

Verdict 10.2 (claims 24, 25)**.** Proved. The theorem is a derivation within classical higher-derivative gravity with the stated symmetry; the attractor mechanism, that the entropy is independent of the asymptotic moduli, is a corollary whenever the extremum is isolated in the scalars. The results agree with the earlier supergravity computations of Cardoso, de Wit and Mohaupt where both apply [93], and the formalism has been used in every later comparison of macroscopic and microscopic entropies.

10.2. **Small black holes.** A half-BPS state of the heterotic string on T^6 with winding w and momentum n on a circle, with $nw>0$, has $d_h(nw)\sim e^{4\pi\sqrt{nw}}$ states, while the two-derivative supergravity solution with these charges has a singular horizon of zero area. Sen [92] argued that the higher-derivative terms stretch the horizon to string scale and that the entropy function with the Gauss–Bonnet term, $\mathcal{L}\supset\frac{1}{16\pi G}\phi(S)(R^2_{\mu\nu\rho\sigma}-4R^2_{\mu\nu}+R^2)$ with the one-loop coupling $\phi(S)$, gives

$$S_{\mathrm{BH}} = 4\pi\sqrt{nw} \tag{16}$$

in four dimensions, in agreement with the statistical entropy to leading order; [94] extended the computation to the heterotic string in D dimensions.

Proposition 10.3 (claim 26)**.** *With the Gauss–Bonnet term as the only higher-derivative correction, the entropy function gives a finite entropy for the small black hole in every dimension $D \geq 4$, proportional to $\sqrt{nw}$. The coefficient equals the statistical one, 4π, for $D = 4$ and $D = 5$, and differs from it for every $D \geq 6$* [94, Section 4]*. In four dimensions the agreement extends to all orders in $1/\sqrt{nw}$ when the full supersymmetric F-term corrections are used in place of the Gauss–Bonnet term* [25, 29]*.*

Proof. The near-horizon solution of [94] has v_1, v_2 and the dilaton fixed by the extremisation with the Gauss–Bonnet coupling, and the entropy at the extremum is $c_D\sqrt{nw}$ with c_D computed there: $c_4 = c_5 = 4\pi$ and $c_D \neq 4\pi$ for $D \geq 6$ (the paper tabulates the ratio). For $D = 4$, [25] evaluated the entropy with the full F-term, and [29] matched the result to the asymptotic expansion of d_h to all orders in $1/\sqrt{nw}$ up to the non-perturbative terms; the Gauss–Bonnet term alone reproduces the leading term because the four-dimensional F-term reduces to it on the near-horizon geometry at this order. □

Verdict 10.4 (claim 26)**.** Scope limited and already qualified by Sen. The Gauss–Bonnet truncation gives the stated match in $D = 4, 5$ and a mismatch in higher dimensions. The abstract and calculation of [94] explicitly identify this limitation and the absence of a full supersymmetric completion in the truncation. It is therefore excluded from proposed new errors. Agreement within a truncated action does not establish control of the full string-scale derivative expansion.

11. The quantum entropy function and the logarithmic corrections

11.1. **The quantum entropy function.** Sen [98] proposed that the exact degeneracy of the horizon states of an extremal black hole with charges $\vec{q}$ is the finite part of the Euclidean string theory path integral on the near-horizon AdS_2,

$$d_{\mathrm{hor}}(\vec{q}) = \Big\langle \exp\Big[-iq_i \oint A_\theta^{(i)}\, d\theta \Big] \Big\rangle_{AdS_2}^{\text{finite}}, \tag{17}$$

with a Wilson line insertion at the boundary carrying the charges, the boundary conditions fixing the charges rather than the chemical potentials, and the "finite part" defined by subtracting the terms proportional to the length of the boundary; and that this quantity matches the appropriately signed horizon index for quarter-BPS black holes after hair removal and the helicity-trace convention are fixed (Section 5.3).

Verdict 11.1 (claim 27)**.** Confirmed. The proposal passes every test that can be computed: the classical limit of (17) is the exponential of the Wald entropy; the one-loop correction is the logarithmic term, which matches the microscopic index exactly in $\mathcal{N} = 4$ and $\mathcal{N} = 8$ (Theorem 11.2); and the localisation of the path integral in $\mathcal{N} = 2$ supergravity, proposed in [6] and carried out in [32, 33], reproduces the exact Rademacher expansion of the index of $\mathcal{N} = 8$ black holes and the leading Bessel function of the $\mathcal{N} = 4$ one [71, 54, 55]. The last of these results is weaker than its statement in the literature, and Sen's own later assessment [113] says so: the claim that supersymmetric localisation of the AdS_2 path integral reproduces the Rademacher expansion from first principles rests on three steps that are not established. The Killing spinors of the near-horizon geometry grow at the boundary of AdS_2 and do not act on the space of normalisable fluctuations over which the path integral is defined, so the localisation argument has no proved domain; the measure on the localisation manifold used in [32, 71] is postulated and not derived, and a different measure changes every subleading term; and the one-loop determinants over the transverse directions depend on an ultraviolet cut-off that the computation sets, without justification, at the four-dimensional Planck scale. Sen himself discusses these qualifications in [113]; they are not counted here as errors left unacknowledged by him. There is no proof: the path integral is not defined beyond the semiclassical expansion, the localisation argument assumes an off-shell supersymmetric completion of the higher-derivative terms, and the identification of the subtracted terms with the hair is made case by case. The missing step is a definition of the right-hand side of (17)

that does not rely on a saddle point expansion; the conjecture is, in this sense, a proposal for what that definition should give.

11.2. **The logarithmic corrections.** The one-loop contribution to (17) from the massless fields on $AdS_2 \times S^2$ gives a term $c \ln A_H$ in the entropy, with c determined by the spectrum and the two-derivative action alone. Banerjee, Gupta, Mandal and Sen [7] computed it for quarter-BPS black holes in $\mathcal{N} = 4$ and eighth-BPS black holes in $\mathcal{N} = 8$ string theory, and Sen [101] for half-BPS black holes in $\mathcal{N} = 2$ theories with n_V vector and n_H hypermultiplets.

Theorem 11.2 (claim 28)**.** *The logarithmic correction to the entropy of a quarter-BPS black hole in any $\mathcal{N} = 4$ string compactification is zero, and that of an eighth-BPS black hole in $\mathcal{N} = 8$ string theory on T^6 is $-4 \ln A_H$. Both agree with the microscopic indices: the asymptotic expansion of the Fourier coefficients of $1/\Phi_{10}$ and its CHL analogues has no logarithmic term, and that of the $\mathcal{N} = 8$ index, a Fourier coefficient of $\vartheta_1(\tau, z)^2/\eta(\tau)^6$, has the term $-2 \ln \Delta$ with $\Delta \propto A_H^2$, that is, $-2 \ln \Delta = -4 \ln A_H$* [7].

Proof. The macroscopic computation is the heat kernel of the quadratic fluctuation operator of all massless fields on $AdS_2 \times S^2$ with the near-horizon flux, including the graviton, the gravitini and the vector fields, with the zero modes of the asymptotic symmetries treated separately; the coefficient is the s^0 term of the heat kernel together with the zero-mode counting, $-(c_{\text{local}} + c_{\text{zm}}) \ln a$ with a the radius. In $\mathcal{N} = 4$ the contributions cancel between the gravity multiplet and the vector multiplets [7, Section 7]; in $\mathcal{N} = 8$ the net result is $-8 \ln a = -4 \ln A_H$ [7, Section 8]. The microscopic expansions are in [7, Section 2], from the saddle point expansion of the inverse Siegel form [28] and of the Jacobi form. The computation was rederived from the on-shell action of the Euclidean solution and from the AdS_2 Schwarzian in [54, 55], with the same coefficients. □

Verdict 11.3 (claim 28)**.** Proved, as a match between two computations each of which is complete: the macroscopic coefficient is a theorem of one-loop quantum field theory on the near-horizon geometry and the microscopic one of the asymptotics of modular forms. The rotating black holes in four and five dimensions [102] and the extremal Kerr and Reissner–Nordström black holes without supersymmetry [101, 103], for which no microscopic count exists, are predictions and remain confirmed in the sense of claim 29.

Proposition 11.4 (claim 29; Sen 2012)**.** *For a half-BPS black hole in an $\mathcal{N} = 2$ supergravity with n_V vector and n_H hypermultiplets, the logarithmic correction is $\frac{1}{12}(23 + n_H - n_V) \ln A_H$, equal to $\left(2 - \frac{\chi}{24}\right) \ln A_H$ with $\chi = 2(n_V - n_H + 1)$ the Euler number of the Calabi–Yau threefold in the type IIA convention, with the universal hypermultiplet included; the type IIB Euler-number relation has the opposite sign.*

Proof. [101], by the same heat kernel method, the only difference from $\mathcal{N} = 4$ being the absence of the cancellation; the result was rederived by [59] from the heat kernel of $\mathcal{N} = 2$ supergravity on the same background. □

Verdict 11.5 (claim 29)**.** Confirmed. A universal comparison for the full class of $\mathcal{N} = 2$ theories is not established here; this does not deny microscopic indices in particular $\mathcal{N} = 2$ compactifications. The comparison made in [101] is with the measure in the OSV formula, and agrees with the measure proposed by Denef and Moore if their result is extrapolated to weak topological string coupling. A model-independent microscopic comparison in a fixed ensemble is not established here. The general framework and the ensemble must be fixed before comparing a logarithmic coefficient.

11.3. **A check of the cutoff-scale preprint.** The current preprint [133] separates the extended-supergravity cancellation from an unresolved nonzero-Euler-number case. There is a local notation problem in its Eq. (2.7): a contribution to $\Delta \ln Z_n$ is printed as the multiplicative factor $(g_s^{-1} \lambda_S^2)^2$. In the logarithm it must instead read

$$\Delta \ln Z_{\text{rot}} = \ln[(g_s^{-1} \lambda_S^2)^2] = -2 \ln g_s + 4 \ln \lambda_S. \tag{18}$$

This is a notation correction, not a refutation of the later result. Combining this expression with the three translational modes, the fermionic modes, two-form modes and the ensemble terms gives the stated coefficients $(\alpha_{NZ} + 7 - n)$ and $(n_{NS} - 4)$ in its Eq. (2.10). The elementary coefficient sums are checked in the analytic supplement. The current arXiv PDF was inspected for this equation; the INSPIRE record checked for this revision supplied no journal-version or erratum metadata. That search result is not a proof that no correction exists elsewhere.

The cutoff comparison in [133] keeps other inputs fixed. Its Section 4 explicitly allows coordinated changes of duality frame and measure, and its Section 5 leaves the $\chi \neq 0$ discrepancy unresolved. These qualifications prevent the conditional cutoff calculation from being restated as a theorem excluding every other ultraviolet completion. We do not claim to solve its Calabi–Yau discrepancy here.

12. String field theory, D-instantons, soft theorems and matrix theory

12.1. **String field theory.** Two claims of Sen about string field theory itself are in the catalogue. The first, with Zwiebach [127], is that closed string field theory is background independent under infinitesimal deformations of the conformal field theory: the string field theories formulated around two nearby conformal backgrounds are related by a field redefinition. The second is the construction of superstring field theory [104, 105, 106] and the proof of its unitarity [107]: a gauge-invariant one-particle-irreducible effective action for heterotic and type II strings, free of the divergences of the picture-changing operators of earlier formulations through the use of a second, free string field of picture number $-3/2$; a Batalin–Vilkovisky master action; and the cutting rules, hence unitarity, of the resulting amplitudes.

Verdict 12.1 (claims 30, 31)**.** Proved. Claim 30 is proved in [127] for marginal deformations to first order (and to second order in the companion paper), by constructing the field redefinition from the connection on the space of conformal field theories; its extension to finite deformations is a theorem only within the formal perturbation theory of the string field and is not a claim of the paper. Claim 31 is a construction and is proved in the papers that make it: the 1PI action is gauge invariant by the main identity of the vertices, the master action satisfies the master equation, and the Cutkosky rules are derived in [107] with the $i\epsilon$ prescription of [74]. The reader who wants the construction in one place has the review [126]. None of this depends on a conjecture; the open questions of superstring field theory are not about these statements.

Corrected versions and infrared scope. The published erratum to [74] and the corrected arXiv v3 must be used for the cutting argument; the correction is excluded from the list of unresolved errors. Section 4 of [107] explicitly notes the infrared problem for $D \leq 4$ and that the inclusive-cross-section construction is not carried out there. This is an acknowledged limitation, not a hidden failure of its perturbative cutting proof. A simple leading-soft cancellation and a counterexample to inferring exact unitarity from a formal series are given in Appendix D. Appendix E constructs the finite-energy limit in the explicit positive soft model and states the additional string-amplitude estimate needed for transfer. Appendix F derives a uniform subtracted tree-emission bound from Sen's soft theorem and demonstrates why overlapping limits require additional estimates. None of these is a construction of the full infrared-finite string S-matrix.

12.2. **D-instantons.** The contribution of D-instantons to string amplitudes has an overall normalisation that the world-sheet theory does not fix, because the open string path integral on the instanton has zero modes, bosonic and fermionic, whose measure is ambiguous. Sen [109, 112, 111] showed that open string field theory fixes it: the zero modes are treated by changing variables from the string field to the collective coordinates with a Jacobian computed in the field theory, the gauge zero mode of the $U(1)$ on the instanton is handled

by a gauge fixing that the field theory provides, and the resulting normalisation is finite and unambiguous.

Verdict 12.2 (claim 32)**.** Confirmed. In two-dimensional string theory the normalisation computed this way [109, 110] agrees with the matrix model to the order compared, including the two-instanton contribution [14]; in type IIB string theory in ten dimensions the normalisation of the D-instanton correction to the four-graviton amplitude [111] agrees with the coefficient predicted by $SL(2,\mathbb{Z})$ invariance, the Green–Gutperle Eisenstein series. The claim that string field theory fixes the normalisation in every background is confirmed by these cases and not proved in general: the construction requires the string field theory of the instanton to be well defined, which it is in these examples, and a general statement needs a general treatment of the zero modes of the Ramond sector, which is not established by the examples discussed here.

12.3. **Soft theorems.** Laddha and Sen [64] proved the classical soft graviton theorem in dimensions $D > 4$, the statement that the low-frequency gravitational radiation from a scattering process is determined by the momenta and angular momenta of the incoming and outgoing objects, including the sub-subleading term, directly from the classical equations [108]; and Sahoo and Sen [80] found that in $D = 4$ the soft expansion acquires logarithmic terms, $\ln\omega$ at the subleading order, whose coefficients are again universal, and conjectured the form of the corresponding quantum terms.

Verdict 12.3 (claims 33, 34)**.** Claim 33 is proved: the classical theorem in $D > 4$ is derived in [64], and the classical logarithmic terms in $D = 4$ in [80], from the classical equations of motion with the long-range forces included. Claim 34 is confirmed: the quantum logarithmic terms were computed at one loop in [80] for scalar matter coupled to gravity and to electromagnetism and found to have the conjectured form, with the classical term and a quantum correction that depends only on the asymptotic momenta; the universality of the one-loop $\ln\omega$ terms was extended to spinning particles with the non-minimal couplings allowed by gauge and coordinate invariance by Krishna and Sahoo [61], and the classical soft factors have been pushed to all orders in the electromagnetic case [58]. The remaining conjecture is that the quantum $\ln\omega$ terms are one-loop exact, and the missing step is a proof that the two-loop and higher infrared divergences exponentiate in the way the conjecture requires; [61] states this as open, a complete higher-loop argument is not supplied here.

A recent scope check. The all-orders classical analysis of [132] distinguishes the Newtonian result from its proposed relativistic extension. It also finds that a separate all-orders conjecture of Alessio, Di Vecchia and Heissenberg, consistent with two-particle scattering, does not hold for general initial or final states with at least three particles. This is not a counterexample to Sen's earlier subleading quantum result. The extension in [132] is explicitly conjectural and is recorded as an open construction, not a wrong proof.

12.4. **Matrix theory.** Sen [120] argued, independently of Seiberg [85], that the matrix theory description of M-theory in the infinite momentum frame follows from the compactification of M-theory on a light-like circle, taken as a limit of a small space-like circle, and extended it to tori T^n: the theory of N D0-branes on T^n in a limit in which the torus and the string scale shrink. For $n \leq 3$ the limit is a maximally supersymmetric Yang–Mills theory in $n+1$ dimensions; for $n = 4, 5$ it is the $(2,0)$ theory on T^5 and the little string theory on T^5 respectively; and for $n \geq 6$ the limit is, as Sen states, "a theory without an independent definition", the degrees of freedom of the D0-branes not decoupling from gravity. The two statements H1 and H2 of the catalogue are these, and they are recorded without a verdict: the first is the Seiberg–Sen argument, which is a derivation within the assumptions of the infinite momentum frame and has no independent test for $n \geq 4$; the second is a statement about a theory that is not defined, and cannot be tested in any regime. We mention them

because the paper is cited as a derivation of matrix theory and the reader should know that its content for $n \geq 6$ is a limit of a non-decoupled system, as its author says.

13. Conclusion

The selected catalogue contains results proved within specified frameworks and physical proposals with different degrees of support. The exact computations and analytic tests do not justify a count of "fatal errors" in an author's work. The toroidal negative full indices and their subsequent hair interpretation are already in the literature. The caustic calculation concerns the classical dust approximation and leaves the homogeneous rolling solution intact. The divisor singularity belongs to the orientifold cover, while a local I_0^* family shows why non-Higgsability alone does not prohibit weak coupling.

The remaining questions are mathematical and physical construction problems: the full monopole L^2 statement beyond the cited cases, general hair identification, a controlled definition of the quantum entropy function, and extensions of the stated soft and string-field results. They are not resolved by the finite tables. A meaningful new criticism would require a precise original statement with all its hypotheses retained and a counterexample within those hypotheses. No new false proof by Sen meeting that criterion is established in this revision.

A reader who wants to repeat the computations of this paper will find them in the archive [10], together with the Lean formalisation of the finite statements: the Fourier coefficients of Tables 6 and 7, the wall-crossing identities of Theorem 4.2 and the twenty-one negative indices of Theorem 5.4, each computed inside the Lean kernel from the definitions of the theta functions. The catalogue and physical interpretations are not formalised by these certificates. Mathematical parts of such subjects can be formalised, but that work must not be inferred from a finite arithmetic check.

Appendix A. The computation of the Fourier coefficients

The tables of this paper are Fourier coefficients of the inverse of the product (3), computed in exact rational arithmetic. We record the steps so that the numbers can be reproduced without the program, which is in the archive [10].

The genera. The twisted elliptic genera are expanded in $q = e^{2\pi i\tau}$ and $y = e^{2\pi iz}$ from the theta functions $\theta_i(\tau, z)$ and $\eta(\tau)$ as q-series with coefficients in $\mathbb{Z}[y, y^{-1}]$, and the coefficients $c_b^{(r,s)}(4n - j^2)$ are read off from $\sum_{n,j} c_b^{(r,s)}(4n - j^2)\, q^n y^j$. For K3, $F^{(0,0)} = 8A$ with $A = \sum_{i=2}^{4} \theta_i(\tau, z)^2/\theta_i(\tau, 0)^2$; for the $\mathbb{Z}_2$ CHL model the genera are $F^{(0,0)} = 4A$, $F^{(0,1)} = \frac{4}{3}A - \frac{2}{3}B\,E_2(\tau)$, $F^{(1,0)} = \frac{4}{3}A + \frac{1}{3}B\,E_2(\tau/2)$, $F^{(1,1)} = \frac{4}{3}A + \frac{1}{3}B\,E_2((\tau+1)/2)$, with $B = \theta_1(\tau, z)^2/\eta(\tau)^6$ and $E_2(\tau) = 2E(2\tau) - E(\tau)$, where E is the Eisenstein series of weight two normalised to $1 - 24\sum_{n\geq 1}\sigma_1(n)q^n$; for the $\mathbb{Z}_2$ toroidal orbifold they are (9). The arguments $\tau/2$ and $(\tau+1)/2$ are handled by expanding in $q^{1/2}$ with the sign $(-1)^n$ on the odd powers, so every series is a Laurent series in $x = q^{1/8}$, which is the smallest fractional power that occurs.

The product. Each factor $(1 - e^{2\pi i(k'\sigma + l\rho + jv)})^{-c}$ of $1/\widetilde{\Phi}$ is expanded as a binomial series in the monomial $e^{2\pi i(k'\sigma+l\rho+jv)}$ to the order needed, and the factors are multiplied as polynomials in the three variables $e^{2\pi i\rho}$, $e^{2\pi i\sigma/N}$, $e^{2\pi iv}$ truncated at the exponents $Q^2/2 \leq L$, $NP^2/2 \leq K$ wanted in the tables. The exponent $c = \sum_s e^{2\pi isl/N} c_b^{(r,s)}(4k'l - j^2)$ is an integer for $N \leq 2$, where the root of unity is ± 1; for $N = 3, 5, 7$ it would require cyclotomic arithmetic, and we have not computed those models. The one factor with $k' = l = 0$, $(1 - e^{\mp 2\pi iv})^{-2}$ after the prefactor is taken into account, is the only one whose expansion depends on the chamber: in the chamber $\operatorname{Im} v < 0$ it is expanded in powers of $e^{-2\pi iv}$, in the chamber $\operatorname{Im} v > 0$ in powers of $e^{2\pi iv}$. The two expansions give the two columns of Table 6

Q^2	P^2	$d(Q,P)$ for $Q\cdot P=-2,\,-1,\,0,\,1,\,2,\,3,\,4$						
2	2	-209 304	130 329	50 064	25 353	648	327	0
4	2	-2 023 536	1 598 376	1 127 472	561 576	50 064	8 376	-648
6	2	-15 493 728	16 844 421	16 491 600	8 533 821	1 127 472	130 329	-15 600
4	4	-16 620 544	28 698 000	32 861 184	18 458 000	3 859 456	561 576	12 800
6	4	-53 249 700	474 507 528	632 078 672	392 427 528	110 910 300	18 458 000	1 127 472
6	6	2 857 656 828	11 890 608 225	16 193 130 552	11 232 685 725	4 173 501 828	920 577 636	110 910 300

TABLE 6. The index $d(Q,P)=(-1)^{Q\cdot P+1}c(Q,P)$ of quarter-BPS dyons of the heterotic string on T^6, from $1/\Phi_{10}$, in the chamber $\operatorname{Im} v<0$ (the expansion in powers of $e^{-2\pi i v}$). The table agrees entry by entry with Table 1 of [100]. Entries with $Q\cdot P<0$ are the other chamber's values at $|Q\cdot P|$ by the symmetry $v\mapsto -v$; their negativity for $Q\cdot P\le -1$ is the two-centred contribution of Theorem 4.2.

	$Q\cdot P=0$			$Q\cdot P=1$			$Q\cdot P=2$		
Q^2	$P^2=0$	$P^2=2$	$P^2=4$	$P^2=0$	$P^2=2$	$P^2=4$	$P^2=0$	$P^2=2$	$P^2=4$
0	2	64	816	0	0	-8	0	0	0
1	-12	-224	-1 248	1	96	1 968	0	0	-12
2	48	1 152	18 240	-8	-256	840	0	64	2 592
3	-168	-3 392	-10 320	37	1 376	34 656	-2	-224	2 432
4	528	11 520	200 736	-136	-3 840	16 632	16	1 152	43 392
5	-1 512	-30 336	-55 424	439	13 152	343 152	-72	-3 392	33 720
6	4 032	83 968	1 544 832	-1 288	-33 536	171 152	256	11 520	414 336
7	-10 176	-202 560	-179 022	3 503	92 928	2 476 752	-806	-30 336	302 400
8	24 528	496 512	9 480 000	-8 968	-220 672	1 265 256	2 320	83 968	2 926 080
9	-56 796	-1 118 496	-155 232	21 854	540 416	14 545 584	-6 200	-202 560	2 049 968
10	127 008	2 521 600	49 523 328	-51 080	-1 204 992	7 558 560	15 616	496 512	16 919 712
11	-275 544	-5 374 656	2 684 560	115 154	2 711 616	73 540 080	-37 508	-1 118 496	11 568 000
12	581 952	11 389 440	228 872 064	-251 528	-5 741 824	38 736 600	86 544	2 521 600	84 554 880

TABLE 7. The index $d(Q,P)$ of quarter-BPS dyons in the $\mathbb{Z}_2$ toroidal orbifold of type IIB on T^6, from the product formula for $1/\widetilde{\Phi}_2$ with the twisted elliptic genera (9), in the chamber $\operatorname{Im} v<0$ and with the normalisation of Remark 5.9. The entries agree with Tables 17, 18 and 19 of [16] in their common range $Q^2\le 8$. The column $P^2=0$ lies outside the single-centred cone; its sign is $(-1)^{Q^2}$. The negative entries inside the cone are those of Table 5.

at $\pm Q\cdot P$, and their difference is the wall-crossing jump of Theorem 4.2, which the program checks for every $Q^2,P^2\in\{2,4,6\}$ and $|Q\cdot P|\le 4$.

The index. The coefficient of $e^{2\pi i(\rho Q^2/2+\sigma P^2/2+v\,Q\cdot P)}$ in $1/\widetilde{\Phi}$, multiplied by $(-1)^{Q\cdot P+1}$, is the index $d(Q,P)$ of (2). For the orbifolds the prefactor of the product is $e^{2\pi i(\rho+\sigma/N+v)}$ for the CHL models and $e^{2\pi i(\rho+v)}$ for the toroidal orbifolds, as in [16], and the integral over σ runs over a period of length N (Remark 5.9).

The tables. Table 6 is the heterotic index for $Q^2,P^2\in\{2,4,6\}$ and $-2\le Q\cdot P\le 4$; it reproduces Table 1 of [100] entry by entry. Table 7 is the index of the $\mathbb{Z}_2$ toroidal orbifold for $Q^2\le 12$, $P^2\in\{0,2,4\}$ and $Q\cdot P\in\{0,1,2\}$; it agrees with Tables 17 to 19 of [16] where they overlap and extends them from $Q^2=8$ to $Q^2=12$. Table 5 lists the charges in the cone (8) with $Q^2\le 12$, $P^2\le 8$, $Q\cdot P\le 4$ whose index is negative. The same scan over the cone of the heterotic string and of the $\mathbb{Z}_2$ CHL model, up to $Q^2\le 12$ (heterotic) or $NQ^2/2\le 6$ (CHL), $P^2\le 6$, $Q\cdot P\le 4$, finds no non-positive index, and the CHL values reproduce Table 2 of [100].

Appendix B. The caustic for a periodic perturbation

This appendix works out Theorem 7.4 for the simplest inhomogeneous initial datum in one space dimension, a periodic perturbation of the homogeneous decay,

$$T_0(x) = T_* + a\cos kx, \qquad a > 0,\ k > 0, \tag{19}$$

on the branch $\partial_t T > 0$, and records the quantities that a numerical solution of the full equation (11) can be compared with.

Velocity and its derivative. With $T_0' = -ak\sin kx$ and $W_0 = \sqrt{1 + a^2k^2\sin^2 kx}$,

$$v_0(x) = \frac{ak\sin kx}{\sqrt{1 + a^2k^2\sin^2 kx}}, \qquad v_0'(x) = \frac{ak^2\cos kx}{(1 + a^2k^2\sin^2 kx)^{3/2}}. \tag{20}$$

The dust moves away from the maxima of T_0 (where $\cos kx = 1$ and $v_0' > 0$, a rarefaction) and towards its minima (where $\cos kx = -1$ and $v_0' = -ak^2 < 0$).

The caustic time. The supremum of $-v_0'$ is attained at the minima $x_m = (2m+1)\pi/k$ of T_0, so by Theorem 7.4(iv)

$$\mu_- = \sup_x(-v_0'(x)) = ak^2, \qquad t_c = \frac{1}{ak^2}. \tag{21}$$

The characteristics $X(t;x_0) = x_0 + t\,v_0(x_0)$ from a neighbourhood of x_m converge on x_m, which is a fixed point of the flow, and have a singular differential there at t_c; for $t > t_c$ the map $x_0 \mapsto X(t;x_0)$ folds over near x_m and three characteristics pass through each point of a growing interval around it. The smaller the perturbation, the later the caustic, and no value of $a > 0$ avoids it; the time is finite for every amplitude and every wavelength, and diverges only in the homogeneous limit $a \to 0$ or in the long-wavelength limit $k \to 0$ at fixed a.

The density. Along the characteristic from x_0,

$$\varepsilon\big(t, X(t;x_0)\big) = \frac{\varepsilon_0(x_0)}{1 + t\,v_0'(x_0)}, \qquad \varepsilon(t,x_m) = \frac{\varepsilon_0(x_m)}{1 - t/t_c},$$

and the density at the minimum diverges as $(t_c - t)^{-1}$. Near the caustic, for $x_0 = x_m + \xi$ with ξ small,

$$v_0'(x_0) = -ak^2\Big(1 - \tfrac{k^2\xi^2}{2}(1 + 3a^2k^2) + O(\xi^4)\Big),$$

so at $t = t_c$ the linear term of $X(t_c; x_m + \xi) - x_m$ cancels and

$$X(t_c; x_m + \xi) = x_m + c\,\xi^3 + O(\xi^5), \qquad c = \tfrac{k^2}{6}(1 + 3a^2k^2),$$

while the density there is $\varepsilon_0(x_m)/(\frac{k^2}{2}(1 + 3a^2k^2)\xi^2) \propto |x - x_m|^{-2/3}$: the integrable density singularity at the cubic cusp onset, followed by fold points after the first focusing time, with total energy in any interval finite at t_c and infinite density at one point. For $t > t_c$ the three-valued velocity field carries a finite energy density on each sheet and the dust description of [86] has no single-valued continuation.

Comparison with the full equation. Felder, Kofman and Starobinsky [44] integrated (11) numerically with the exponential potential and inhomogeneous initial data, and found that after a transient the solution is approximated by the free-streaming (Hamilton–Jacobi) solution, which is (12), and that the caustics predicted by it form in the full equation, with second derivatives of T growing without bound at the caustic; Barnaby [4] showed the same analytically for several effective actions. The dust formula gives a comparison time after matching effective initial data. It is not an exact prediction of the full Born–Infeld evolution: the transient and higher-derivative corrections depend on the initial data and the effective action. No amplitude-independent transient time or string-scale conclusion is inferred here.

Appendix C. Independent scope checks and source register

C.1. **The sign of compression.** For the forward branch $T_t = \sqrt{1+|\nabla T|^2}$, the characteristic velocity is minus the normalised gradient. In one dimension let $d = (1+T_0'^2)^{3/2} > 0$. Then

$$v_0' = -T_0''/d < 0 \quad \Longleftrightarrow \quad T_0'' > 0.$$

This sign is independent of the density and reverses if the time branch is reversed. The scalar equivalence and the identity $1+(-1/\lambda)\lambda = 0$ for $\lambda < 0$ are separately checked in the analytic Lean supplement. These identities are not a formalisation of the full characteristic PDE theorem. For the periodic datum in Appendix B, compactness ensures attainment. For a general noncompact datum, replacing a supremum by a maximum is an additional assumption.

C.2. **What an index can and cannot imply.** Write a signed index as $I = B - F$ with nonnegative integer counts. The toy example $B = 3, F = 2$ gives $I = 1 > 0$ although fermionic states remain. Thus positivity cannot alone prove that every contributing state has zero angular momentum. Conversely, if assumptions A_1, A_2, A_3 imply $I > 0$ and a computed index has $I < 0$, one can conclude only $\neg(A_1 \wedge A_2 \wedge A_3)$. Selecting a particular failed assumption requires additional physics. This logical distinction is especially relevant when interpreting a negative full index as a statement about hair in every duality frame.

C.3. **The charge cone and reproducibility.** Equation (3.9) of [100] contains $3(Q \cdot P) \leq 2Q^2 + P^2$. The squared inner product occurs instead in the discriminant constraint. Mixing these two conditions changes the admissible charges. The Python scan uses the linear inequality, integer arithmetic, $Q^2 \leq 12$, $P^2 \leq 8$ and $Q \cdot P \leq 4$ and returns thirty negative triples. The Lean series certificate retains its original table range and verifies the original twenty-one triples with the corrected cone predicate. Its list is a subset, not an exhaustive scan of the larger box. The formal scope report records the exact toolchain, source hashes, successful checks and omissions.

C.4. **The base and the cover.** The local test $\chi = 1$ already separates the two spaces. On the base, $\eta^2 - h = 0$ has derivative $\partial_h = -1$, hence is smooth. On the cover $h = \xi^2$, the equation becomes $(\eta-\xi)(\eta+\xi) = 0$, a singular union at the origin. Blowing up or normalising the cover divisor may be useful for its invariants, but cannot turn this distinction into an error in the original weak-coupling expansion. The discriminant identity of Theorem 9.1 is checked as a polynomial identity in the analytic Lean supplement; the global geometry is not.

C.5. **Primary passages used for scope corrections.** The principal comparisons are reproducible from the following sources, with their original qualifications retained.

- [100], introduction, pp. 3–5 and Eq. (3.9): assumptions about hair and other index contributions, and the linear charge cone.
- [16], Sections 4.1–4.2: known negative full indices in toroidal orbifolds; [17], Sections 4–5: proposed hair factors, finite positive horizon-index tests, and remaining mode-identification questions.
- [86], Eqs. (19)–(25): the pressureless Hamiltonian equations and the static solution with arbitrary density. The homogeneous decay assumption is stated in its abstract.
- [118], abstract, weak-coupling expansion and auxiliary double cover: the reduction is formal and singular auxiliary spaces are explicitly discussed. The D7 cover singularities are developed in [18]; global weak-coupling obstructions and the I_0^* exception are in [131].
- [94], abstract and Gauss–Bonnet calculation: the higher-dimensional coefficient mismatch is reported by Sen, rather than discovered here.
- [113]: domain, ultraviolet and flat-direction issues are raised by Sen himself; they are not new unacknowledged objections.




FIGURE 4. Local distinction at $\chi = 1$. The base divisor $h = \eta^2$ is smooth. Its pullback under $h = \xi^2$ consists of the crossing branches $\eta = \pm\xi$. The diagram displays real slices of complex divisors; it is a schematic, not a rendering of the full compactification.

- [45], Theorem 1.2 and Section 6, state the general coprime case using the announced metric compactification; [60], Theorem A: magnetic charge three is covered under the stated QFB-metric hypothesis. [61] separates proved one-loop universality from conjectural higher quantum extensions. [132] labels its beyond-Newtonian extension a conjecture.

This register is not evidence that every reference or every later citing paper has been independently checked. The bibliography supplies further context; new objections require their own exact source and hypothesis check.

APPENDIX D. TWO LIMITS OF PERTURBATIVE REASONING

D.1. **An exact leading-soft model.** Assume independent soft emissions with intensity $b\,d\omega/\omega$, $b \geq 0$, an infrared cutoff λ, detector resolution E and hard scale Λ, with $0 < \lambda < E < \Lambda$. Assume also the virtual factor $\exp[-b\ln(\Lambda/\lambda)]$. Summing any number of unobserved emissions below E gives

$$\begin{aligned} P_\lambda &= e^{-b\ln(\Lambda/\lambda)} \sum_{n=0}^{\infty} \frac{[b\ln(E/\lambda)]^n}{n!} \\ &= e^{-b\ln(\Lambda/\lambda)+b\ln(E/\lambda)} = e^{-b\ln(\Lambda/E)}. \end{aligned} \tag{22}$$

The series is absolutely convergent and the last expression is independent of λ. This is a self-contained cancellation in the independent-emission model. It assumes soft factorisation and the virtual factor; it does not prove either for every string-field-theory diagram, include subleading emissions, or construct the general four-dimensional observable. The missing physical and analytic steps must not be concealed by the elementary exponential identity. The finite-exponent cancellation is separately checked in Lean.

D.2. **Formal unitarity does not determine a completion.**

Proposition D.1. *There exist two smooth scalar amplitudes with the same Taylor series at zero coupling such that one is unitary and the other is not for every positive coupling.*

Proof. Take $S_0(g) = 1$ and $S_1(g) = 1 + f(g)$, where $f(0) = 0$ and $f(g) = e^{-1/g^2}$ for $g \neq 0$. For every integer $N \geq 0$, choose an integer m with $2m > N$. Since $e^t \geq t^m/m!$ for $t > 0$,

$$0 \leq |g|^{-N} f(g) \leq m!|g|^{2m-N} \longrightarrow 0.$$

Each derivative of f away from zero is a finite linear combination of powers of g^{-1} times $f(g)$. Induction using the displayed bound extends every derivative continuously to zero with value zero. Thus both amplitudes have the same Taylor series, the constant 1. It satisfies the formal scalar unitarity identity to every order, whereas $S_1(g) > 1$ gives $|S_1(g)|^2 > 1$ for $g > 0$. □

This is a counterexample to a general inference from a perturbative series to an exact completion, not a counterexample to Sen's perturbative theorem. The flat-function smoothness argument is proved here in ordinary analysis and is not part of the finite Lean certificate. A nonperturbative completion needs additional input such as a summation prescription, analytic class, boundary conditions or an independently constructed theory.

Appendix E. A finite-energy infrared limit in an explicit soft model

The leading-soft cancellation of Appendix D can be strengthened to a construction allowing infinitely many unresolved quanta and a total-energy observable. The theorem below is a statement about an explicitly defined positive emission model. Its relation to a string amplitude requires additional factorisation estimates, stated afterwards. It is not a claimed solution of the full four-dimensional string-field-theory problem.

Theorem E.1 (finite-energy limit and quantitative cutoff removal)**.** *Fix $b \geq 0$ and $0 < E < \Lambda$. For $0 < \lambda < E$, let N_λ be a Poisson random measure on $[\lambda, E]$ with intensity $b\,d\omega/\omega$, coupled by restriction of one measure on $(0, E]$. Put $S_\lambda = \int_{[\lambda,E]} \omega N(d\omega)$. There is a finite nonnegative random variable S such that $S_\lambda \uparrow S$ almost surely and in L^1, and*

$$\mathbb{E}S = bE, \qquad \mathbb{E}|S - S_\lambda| = b\lambda, \tag{23}$$

$$\mathbb{E}e^{-tS} = \exp\left[b\int_0^E \frac{e^{-t\omega} - 1}{\omega}\, d\omega\right] \quad (t \geq 0). \tag{24}$$

For a bounded Lipschitz function $F : [0, \infty) \to \mathbb{R}$, with Lipschitz constant L_F, define the inclusive model functional

$$\mathcal{I}_\lambda(F) = e^{-b\ln(\Lambda/\lambda)} \sum_{n=0}^{\infty} \frac{b^n}{n!} \int_{[\lambda,E]^n} F\left(\sum_{j=1}^n \omega_j\right) \prod_{j=1}^n \frac{d\omega_j}{\omega_j}. \tag{25}$$

Then

$$\mathcal{I}_0(F) = (E/\Lambda)^b \mathbb{E}F(S), \tag{26}$$

$$|\mathcal{I}_\lambda(F) - \mathcal{I}_0(F)| \leq (E/\Lambda)^b L_F b\lambda. \tag{27}$$

If $b > 0$, the total number of emitted quanta is infinite almost surely, although S is finite, and the law of S has no atoms. Consequently the cutoff also has a limit for each total-energy veto $F(s) = \mathbf{1}_{s \leq \delta}$.

Proof. Partition $(0, E]$ into dyadic bands $J_k = (E2^{-k-1}, E2^{-k}]$, $k \geq 0$. On each band take an independent Poisson number with mean $b \ln 2$, and conditional on that number take independent energies with density $1/(\omega \ln 2)$ on the band. This constructs the measure on the product probability space without assuming a finite total particle number. The expectation of the energy in J_k is $b|J_k|$, so the sum of the expectations is bE. The nonnegative total energy is therefore finite almost surely. Monotone convergence gives the almost-sure limit and

$$\mathbb{E}(S - S_\lambda) = \int_0^\lambda \omega\, b\frac{d\omega}{\omega} = b\lambda.$$

For a finite cutoff, the elementary generating function of a Poisson distribution yields

$$\mathbb{E}e^{-tS_\lambda} = \exp\left[b\int_\lambda^E \frac{e^{-t\omega}-1}{\omega}\,d\omega\right].$$

The bound $|e^{-t\omega}-1| \le t\omega$ makes the exponent integrable at zero. Dominated convergence proves the Laplace formula for S. Writing the Poisson expectation of $F(S_\lambda)$ as its absolutely convergent sum over particle numbers gives exactly

$$\mathcal{I}_\lambda(F) = (E/\Lambda)^b \mathbb{E}F(S_\lambda).$$

The Lipschitz estimate follows from the common coupling and the preceding L^1 identity.

If $b > 0$, the count on $(\lambda, E]$ has Poisson mean $b\ln(E/\lambda) \to \infty$. For every fixed integer K, its probability of being at most K tends to zero. Hence the total count is infinite almost surely. To prove absence of atoms, split at $a \in (0, E)$. The sum of energies in $[a, E]$, conditional on at least one emission, has a continuous distribution, since each energy has a density and convolutions retain that property. It is independent of the energies below a. Thus for every x,

$$\mathbb{P}(S = x) \le \mathbb{P}(N([a, E]) = 0) = (a/E)^b.$$

Let $a \downarrow 0$. Finally $S_\lambda \to S$ almost surely, and the indicator of a fixed total-energy threshold is continuous away from an event of probability zero. Bounded convergence proves the final assertion. For $b = 0$ the measure is concentrated at $S = 0$ and the conclusion is immediate directly. □

This theorem treats total unresolved energy, not just a bound on each individual quantum. It also explains why a finite-energy limit need not be a finite-particle Fock vector. It constructs an energy distribution, not the angular momentum, memory algebra or gauge constraints of a gravitational quantum state.

Proposition E.2 (an explicit transfer criterion)**.** *Let $\Sigma_\lambda(F)$ be a proposed regulated string observable. Suppose, for each observable in a specified class,*

$$\Sigma_\lambda(F) = H\mathcal{I}_\lambda(F) + R_\lambda(F),$$

where H is finite and independent of λ, and $R_\lambda(F) \to R_0(F)$ is separately proved. Then $\Sigma_\lambda(F)$ has a finite limit, with

$$\Sigma_0(F) = H\mathcal{I}_0(F) + R_0(F).$$

For Lipschitz F, its error is at most $|H|(E/\Lambda)^b L_F b\lambda + |R_\lambda(F) - R_0(F)|$.

Proof. Subtract the two expressions and apply Theorem E.1 and the triangle inequality. The convergence assumption on the remainder is essential; the proposition does not establish it. □

The remaining string-theory problem. Weinberg's treatment [134] establishes the familiar gravitational soft cancellations in its stated setting. More recent algebraic scattering constructions [135, 136] explicitly address the state-space problem. They do not justify silently replacing a constructed string-field evolution by an assumed automorphism. For the full string problem one must supply a specified stable compactification and observable, a common regulator compatible with the gauge identities, factorisation over all relevant degeneration regions, the corresponding virtual factors and measure, and control of collinear regions and overlapping limits. In the transfer proposition these obligations reside in the existence of H and the convergence of R_λ. No proof of those obligations for every four-dimensional superstring background is supplied here. The Poisson construction and the scalar exponential identity are not Lean-formalised as probability theory in this release.

Appendix F. A uniform single-soft estimate and the overlapping-limit obstruction

This appendix makes one part of the transfer problem quantitative. The tree soft theorem is an established result of Sen [137]; the estimate below is a consequence on compact sets of massive hard kinematics, not a replacement for that theorem or a new proof of the complete four-dimensional infrared problem. We then give an exact example explaining why the same argument does not establish the required loop remainder.

F.1. **Kinematics and the tree pole decomposition.** Use signature $(+---)$ and write a future null soft momentum as $q = \omega n$, where $n = (1, \widehat{n})$, $|\widehat{n}| = 1$. The hard momenta p_i are future timelike; incoming and outgoing legs have signs $\eta_i = -1$ and $+1$, respectively. At the limiting hard configuration, $\sum_i \eta_i p_i = 0$. In the convention with linear coupling $-(\kappa/2)h_{\mu\nu}T^{\mu\nu}$, the leading coefficient for one physical graviton polarisation is

$$a(n) = \frac{\kappa}{2}\mathcal{M}_N \sum_i \eta_i \frac{\epsilon_{\mu\nu}(n) p_i^\mu p_i^\nu}{p_i \cdot n}. \tag{28}$$

Here $\mathcal{M}_N$ is the stripped hard amplitude. A smooth on-shell recoil prescription is fixed when imposing momentum conservation at nonzero q. Its changes at order ω belong in the remainder. Formula (28) is unchanged by $\epsilon_{\mu\nu} \mapsto \epsilon_{\mu\nu} + n_\mu \xi_\nu + \xi_\mu n_\nu$, since the change is $\kappa \mathcal{M}_N \xi \cdot \sum_i \eta_i p_i = 0$.

Choose a compact set of nonsingular hard configurations, away from additional hard-channel poles and thresholds, and restrict to a recoil prescription defined on a neighbourhood of it. Since each hard leg is massive,

$$p_i \cdot n \geq p_i^0 - |\boldsymbol{p}_i| = \frac{m_i^2}{p_i^0 + |\boldsymbol{p}_i|} > 0. \tag{29}$$

After decreasing the upper soft energy E if necessary, the same lower bound with a smaller positive constant holds uniformly along the recoil family. A physical helicity basis can be chosen in finitely many angular charts, a finite cover of the sphere of the kind whose combinatorics is studied in [5]; the bounds below concern its invariant norms and do not require a globally smooth polarisation vector on the sphere.

In the tree sectors covered by [137], use the usual meromorphic tree amplitude and its external-leg pole factorisation. On the selected compact neighbourhood it has the form

$$\mathcal{M}_{N+1}(\omega, n) = \frac{1}{\omega}\sum_i f_i(\omega, n) + h(\omega, n), \qquad \sum_i f_i(0, n) = a(n), \tag{30}$$

where the f_i are continuously differentiable in ω and h is bounded, uniformly in the hard configurations and the angular charts. These are tree analytic inputs, not assumptions about a loop amplitude. One obtains the f_i by absorbing the nonzero factor $2\eta_i p_i(\omega, n) \cdot n$ into the numerator of each external-leg propagator. Other channels are regular by the choice of neighbourhood. The amplitude may contain infinitely many string poles elsewhere; the statement uses only local regularity away from those poles, rather than a truncation of the string spectrum.

Proposition F.1 (uniform tree remainder)**.** *Let $H = \sup|h|$, $D_i = \sup|\partial_\omega f_i|$, and $C = H + \sum_i D_i$ on the neighbourhood just specified. Then*

$$\mathcal{M}_{N+1}(\omega, n) = \frac{a(n)}{\omega} + r(\omega, n), \qquad |r(\omega, n)| \leq C \quad (0 < \omega \leq E). \tag{31}$$

Proof. Subtract the coefficient at $\omega = 0$ in (30). The fundamental theorem of calculus gives

$$r(\omega, n) = h(\omega, n) + \sum_i \int_0^1 \partial_\omega f_i(t\omega, n)\, dt.$$

The stated bound follows by the triangle inequality. All suprema are finite by the regularity and compactness hypotheses. The lower bound (29) is what makes the estimate uniform in angle for this argument. It supplies no such constant for a massless hard leg. □

F.2. **An explicit bound after the real-emission subtraction.** Let Ω be the angular and physical-polarisation space, with finite measure $\mu(\Omega)$; for both helicities integrated over the full sphere, $\mu(\Omega) = 8\pi$. Put $A = \sup_\Omega |a|$. The one-particle phase-space convention is

$$d\Phi_q = \frac{\omega\, d\omega\, d\mu(n)}{2(2\pi)^3}.$$

Let a real measurement function satisfy $|F(\omega, n)| \le B$ and $|F(\omega, n) - F(0, n)| \le L\omega$. Smooth recoil Jacobians and bounded hard-bin weights may be included in F when these two estimates hold.

Theorem F.2 (integrable subtracted tree emission)**.** *Under Proposition F.1, define*

$$T(\omega, n) = F(\omega, n)|\mathcal{M}_{N+1}(\omega, n)|^2 - F(0, n)\frac{|a(n)|^2}{\omega^2}.$$

The signed integral $\mathcal{R}_\lambda(F) = \int_{\lambda<\omega<E} T\, d\Phi_q$ has a limit as $\lambda \downarrow 0$, and

$$|\mathcal{R}_0(F) - \mathcal{R}_\lambda(F)| \le \frac{\mu(\Omega)}{2(2\pi)^3}\left[(LA^2 + 2BAC)\lambda + \frac{BC^2}{2}\lambda^2\right]. \tag{32}$$

Proof. The complex amplitude identity in (31) gives

$$\left|\left|\frac{a}{\omega} + r\right|^2 - \frac{|a|^2}{\omega^2}\right| \le \frac{2|a||r|}{\omega} + |r|^2.$$

Separate T into $(F - F(0, n))|a|^2/\omega^2$ and F times the difference of the two squared amplitudes. Consequently

$$|T(\omega, n)| \le \frac{LA^2 + 2BAC}{\omega} + BC^2.$$

Multiplication by the phase-space factor ω leaves an integrable bound at zero. Integrating it over $(0, \lambda)$ proves absolute convergence and (32). □

This proves cutoff removal for the *subtracted single-real-emission tree sector*. It neither assigns a finite value to its unsubtracted logarithmic term nor supplies the corresponding virtual contribution. Those are separate ingredients in an inclusive observable.

The lower bound (29) applies to a real on-shell null q, not to an arbitrary virtual loop momentum. For example $p = (m, 0, 0, 0)$ and $k = (0, \boldsymbol{k})$ give $p \cdot k = 0$ even when $\boldsymbol{k} \neq 0$. Hence there is no estimate $|p \cdot k| \ge c\|k\|$ on the complete real loop domain merely because p is massive. A virtual estimate must handle the pole prescription, contour and cancellations; it cannot be obtained by copying the real-emission angular bound.

F.3. **Logarithmic remainders: a sufficient quantitative bound.** A bounded tree remainder must not be assumed unchanged at four-dimensional loop level. Logarithmic soft terms are known [80]. Their presence alone does not prevent integration after the leading subtraction. For a nonnegative integer m, define

$$h_m(\omega) = (1 + \ln(E/\omega))^m, \tag{33}$$

$$P_m(x) = \sum_{j=0}^{m} \binom{m}{j} j! x^{m-j}, \tag{34}$$

$$Q_k(x) = \sum_{j=0}^{k} \binom{k}{j} \frac{j!}{2^j} x^{k-j}. \tag{35}$$

For $0 < \lambda \le E$ and $x = 1 + \ln(E/\lambda)$, direct integration gives

$$\int_0^\lambda h_m(\omega)\, d\omega = \lambda P_m(x), \tag{36}$$

$$\int_0^\lambda \omega h_m(\omega)^2\, d\omega = \frac{\lambda^2}{2} Q_{2m}(x). \tag{37}$$

Indeed put $\omega = \lambda e^{-u}$, expand $(x+u)^m$, and use $\int_0^\infty e^{-cu} u^j du = j!/c^{j+1}$, proved by j integrations by parts. The exponential decay makes each boundary term vanish.

Proposition F.3 (logarithmic extension)**.** *If a specified amplitude has the decomposition* (31) *with the weaker bound* $|r(\omega, n)| \le C h_m(\omega)$*, the same subtraction is absolutely integrable and its tail is at most*

$$\frac{\mu(\Omega)}{2(2\pi)^3}\left[LA^2\lambda + 2BAC\lambda P_m(x) + \frac{BC^2\lambda^2}{2} Q_{2m}(x)\right]. \tag{38}$$

If a regulator-dependent family r_λ *has this bound with the* same C, m *and converges pointwise to* r_0*, its subtracted integrals converge by dominated convergence. If in addition* $|r_\lambda - r_0| \le \varepsilon(\lambda) h_m$ *for* $\omega \ge \lambda$*, their difference is bounded by the tail* (38) *plus*

$$\frac{\mu(\Omega)}{2(2\pi)^3}\varepsilon(\lambda)\left[2BAEP_m(1) + BCE^2 Q_{2m}(1)\right]. \tag{39}$$

Proof. Repeat the proof of Theorem F.2, replacing C by $Ch_m(\omega)$, and then use (36). These moments give a regulator-independent integrable majorant when C, m are uniform. For the quantitative last assertion,

$$\left||a/\omega + r_\lambda|^2 - |a/\omega + r_0|^2\right| \le \frac{2A\varepsilon(\lambda)h_m}{\omega} + 2C\varepsilon(\lambda)h_m^2.$$

Integrate the right side with the measurement bound B and phase space, extending its domain to $(0, E)$. This proves (39). □

The proposition does not assert that the remainders of every string loop amplitude satisfy its uniform hypothesis. Bounds whose constant diverges with the virtual infrared regulator do not meet it.

F.4. **Why separate soft limits do not control a corner.**

Proposition F.4 (an overlapping-limit obstruction)**.** *For every fixed* $x > 0$*, the integral* $\int_0^E (x+y)^{-2}\, dy$ *is finite, and the analogous statement holds with* x *and* y *exchanged. Nevertheless,*

$$\int_\lambda^E \int_\lambda^E \frac{dx\, dy}{(x+y)^2} = \ln \frac{(E+\lambda)^2}{4E\lambda} \longrightarrow +\infty. \tag{40}$$

Proof. The first assertion follows from the antiderivative $-1/(x+y)$. Integrating first in x gives $1/(\lambda+y) - 1/(E+y)$. Integrating in y yields (40). In coordinates $r = x+y$, $t = x/(x+y)$, the Jacobian has absolute value r, so the density becomes $dr\, dt/r$. Resolving the corner by this change of variables exposes a logarithmic radial divergence; it does not remove it. □

This is an analytic counterexample to an inference from separate integrability to joint integrability. It is not asserted to be an actual string diagram or a counterexample to a gravitational Ward identity. It shows why a proof about one soft external momentum cannot replace an estimate at every overlapping degeneration.

For clarity, the useful estimate after such a resolution can also be stated exactly. Suppose one *subtracted* sector of an integral is represented by variables $u_j \in (0, 1]$, an angular variable θ, and density

$$G(u, \theta) \prod_{j=1}^{s} \frac{du_j}{u_j}\, d\nu(\theta).$$

If there are $\delta_j > 0$ and an integrable $C(\theta) \geq 0$ with

$$|G(u,\theta)| \leq C(\theta) \prod_j u_j^{\delta_j}, \tag{41}$$

then its absolute integral is at most $\|C\|_1 \prod_j \delta_j^{-1}$. Removing the region where any $u_j < t$ changes the integral by at most

$$\|C\|_1 \left(\prod_j \delta_j^{-1} \right) \sum_j t^{\delta_j}. \tag{42}$$

This follows from Tonelli's theorem, $\int_0^t u^{\delta-1} du = t^\delta/\delta$, and a union bound on the removed regions. The argument proves a sufficient estimate once the sector representation and its bound are established. It does not establish (41) for the string integrand or identify the physical regulator with this box cutoff.

F.5. **A useful power count for the pure-graviton soft sector.** There is also an all-graph homogeneous estimate worth separating from a full convergence theorem. Consider soft graviton propagators connecting massive eikonal lines. Suppose there are I soft propagators, V internal soft interaction vertices of valences $d_v \geq 3$, and E_a attachments to the eikonal lines. Take each interaction to have at least two soft derivatives. Exclude vacuum components and tadpoles; absorb two-point kinetic insertions into the propagator. Assume the V internal momentum-conservation constraints are independent, so the number of independent soft loop momenta is $\ell = I - V$. Incidence counting gives $E_a = 2I - \sum_v d_v$.

Under a uniform rescaling of all these loop momenta by r, the loop measures, soft propagators, eikonal denominators and vertex numerators have total degree at least

$$\gamma = 4\ell - 2I - E_a + 2V = \sum_v (d_v - 2). \tag{43}$$

Thus $V > 0$ implies $\gamma \geq V > 0$, whereas $V = 0$ gives the marginal degree zero of independent exchanges. The equality follows by substitution of the two counting identities; the inequality uses $d_v \geq 3$. If the remaining angular factors are bounded, a radial integral has density bounded by a constant times $r^{\gamma-1} dr$ and converges for $\gamma > 0$.

This is consistent with the established gravitational infrared analysis [138]. It explains the distinguished role of independent eikonal exchanges. It is not a bound on every partial rescaling: if some momenta stay fixed, a vertex need not supply two powers of the momenta being scaled. Nor does the count bound angular pinch surfaces or other massless species. To use it in (41), every nested and overlapping scaling region must be controlled after the relevant Ward-identity cancellations. A favourable homogeneous degree for the whole graph does not prove this last assertion.

F.6. **What remains for the full string-field-theory observable.** Gravitational soft factorisation and cancellation results extend well beyond a single tree diagram [134, 138]. Those results must retain their scattering assumptions and the cancellations between complete sets of diagrams. They do not identify an arbitrary positive Poisson model with a complete compactified string theory containing its other massless fields.

To apply the transfer proposition of Appendix E, one still needs a specified background, external states and infrared-safe measurement; the common real and virtual regulator; and a subtraction construction compatible with the gauge identities. At each fixed perturbative order that construction must control every simultaneous soft, collinear and degeneration region, the relevant string-state sums, and the remaining hard coefficient. A sufficient route is a uniform integrable bound after all required cancellations, such as (41), on the appropriate contours or cut phase spaces, with control of regulator removal. A rigorously controlled distributional limit could provide another route. Neither route is supplied for the complete observable here. An ordinary nonperturbative S-matrix would require further construction, beyond finiteness at each order.

The tree estimate (32), the logarithmic integration formulas and the corner calculation are established here. A factorisation theorem and convergent remainder for the complete four-dimensional string-field observable have not been proved. The finite algebraic certificates accompanying this appendix do not formalise the soft theorem, the phase-space integrals or the full string amplitude.

Declarations

Funding. No funding was received for this work.

Conflict of interest. The author declares no conflict of interest.

Data availability. The program that computes the Fourier coefficients of Appendix A and every table of this paper, the Lean 4 certificates of its finite statements, the scripts that check the explicit constants of Appendices E and F, and the source of the paper are in release `v2.0.0` of the repository named in the note under the abstract, archived on Zenodo as 10.5281/zenodo.23669248 [10]. The catalogue of Section 2 is kept in the ledger format of the engine Anaxia [11]. The source register in Appendix C records the primary passages used for the scope corrections.

References

[1] D. Astefanesei, K. Goldstein, R. P. Jena, A. Sen and S. P. Trivedi, *Rotating attractors*, JHEP **10** (2006) 058, doi:10.1088/1126-6708/2006/10/058, arXiv:hep-th/0606244.

[2] P. C. Argyres, A. Kapustin and N. Seiberg, *On S-duality for non-simply-laced gauge groups*, JHEP **06** (2006) 043, doi:10.1088/1126-6708/2006/06/043, arXiv:hep-th/0603048.

[3] M. F. Atiyah and N. J. Hitchin, *The geometry and dynamics of magnetic monopoles*, Princeton University Press, Princeton (1988), doi:10.1515/9781400859306.

[4] N. Barnaby, *Caustic formation in tachyon effective field theories*, JHEP **07** (2004) 025, doi:10.1088/1126-6708/2004/07/025, arXiv:hep-th/0406120.

[5] D. Bhattacharjee and U. Bhattacharya, *Iterated nerve complexes for k-fold sensor coverage*, Preprints.org (2026), doi:10.20944/preprints202606.1351.v2.

[6] N. Banerjee, S. Banerjee, R. K. Gupta, I. Mandal and A. Sen, *Supersymmetry, Localization and Quantum Entropy Function*, JHEP **02** (2010) 091, doi:10.1007/JHEP02(2010)091, arXiv:0905.2686.

[7] S. Banerjee, R. K. Gupta, I. Mandal and A. Sen, *Logarithmic Corrections to $\mathcal{N}=4$ and $\mathcal{N}=8$ Black Hole Entropy: A One Loop Test of Quantum Gravity*, JHEP **11** (2011) 143, doi:10.1007/JHEP11(2011)143, arXiv:1106.0080.

[8] D. Bhattacharjee, *Establishing equivariant class $[O]$ for hyperbolic groups*, Asian Res. J. Math. **18** (2022), no. 11, 362–369, doi:10.9734/arjom/2022/v18i11615.

[9] D. Bhattacharjee, *Non-commutativity over canonical suspension η for genus $g \geq 1$ in hypercomplex structures for potential $\rho\phi$*, Asian Res. J. Math. **18** (2022), no. 11, 332–341, doi:10.9734/arjom/2022/v18i11607.

[10] D. Bhattacharjee, *The conjectures of Ashoke Sen: computations, tables and Lean 4 certificate*, Zenodo (2026), doi:10.5281/zenodo.23669248.

[11] D. Bhattacharjee, *BLLM: Beyond the LLM with Deep Human-Mind-AI*, version 3.1.1, Zenodo (2026), doi:10.5281/zenodo.23484395.

[12] N. Banerjee, I. Mandal and A. Sen, *Black Hole Hair Removal*, JHEP **07** (2009) 091, doi:10.1088/1126-6708/2009/07/091, arXiv:0901.0359.

[13] K. Bringmann and S. Murthy, *On the positivity of black hole degeneracies in string theory*, Commun. Num. Theor. Phys. **7** (2013) 15, doi:10.4310/CNTP.2013.v7.n1.a2, arXiv:1208.3476.

[14] B. Balthazar, V. A. Rodriguez and X. Yin, *ZZ instantons and the non-perturbative dual of $c = 1$ string theory*, JHEP **05** (2023) 048, doi:10.1007/JHEP05(2023)048, arXiv:1907.07688.

[15] N. Berkovits, A. Sen and B. Zwiebach, *Tachyon condensation in superstring field theory*, Nucl. Phys. B **587** (2000) 147, doi:10.1016/S0550-3213(00)00501-0, arXiv:hep-th/0002211.

[16] A. Chattopadhyaya and J. R. David, *Properties of dyons in $\mathcal{N}=4$ theories at small charges*, JHEP **05** (2019) 005, doi:10.1007/JHEP05(2019)005, arXiv:1810.12060.

[17] A. Chattopadhyaya and J. R. David, *Horizon states and the sign of their index in $\mathcal{N}=4$ dyons*, JHEP **03** (2021) 106, doi:10.1007/JHEP03(2021)106, arXiv:2010.08967.

[18] A. Collinucci, F. Denef and M. Esole, *D-brane Deconstructions in IIB Orientifolds*, JHEP **02** (2009) 005, doi:10.1088/1126-6708/2009/02/005, arXiv:0805.1573.

[19] G. Lopes Cardoso, B. de Wit, J. Kappeli and T. Mohaupt, *Asymptotic degeneracy of dyonic $\mathcal{N} = 4$ string states and black hole entropy*, JHEP **12** (2004) 075, doi:10.1088/1126-6708/2004/12/075, arXiv:hep-th/0412287.

[20] A. Clingher, R. Donagi and M. Wijnholt, *The Sen Limit*, Adv. Theor. Math. Phys. **18** (2014) 613, doi:10.4310/ATMP.2014.v18.n3.a2, arXiv:1212.4505.

[21] A. Chowdhury, R. S. Garavuso, S. Mondal and A. Sen, *Do All BPS Black Hole Microstates Carry Zero Angular Momentum?*, JHEP **04** (2016) 082, doi:10.1007/JHEP04(2016)082, arXiv:1511.06978.

[22] A. Chowdhury and S. Maji, *Counting $\mathcal{N} = 8$ black holes as algebraic varieties*, JHEP **05** (2024) 091, doi:10.1007/JHEP05(2024)091, arXiv:2311.04786.

[23] A. Chowdhury and S. Maji, *Pure D-brane black holes: BPS counting and non-BPS vacua*, JHEP **08** (2026) 070, doi:10.1007/JHEP08(2026)070, arXiv:2601.07927.

[24] A. Chowdhury, A. Kidambi, S. Murthy, V. Reys and T. Wrase, *Dyonic black hole degeneracies in $\mathcal{N} = 4$ string theory from Dabholkar-Harvey degeneracies*, JHEP **10** (2020) 184, doi:10.1007/JHEP10(2020)184, arXiv:1912.06562.

[25] A. Dabholkar, *Exact counting of black hole microstates*, Phys. Rev. Lett. **94** (2005) 241301, doi:10.1103/PhysRevLett.94.241301, arXiv:hep-th/0409148.

[26] A. Dabholkar and D. Gaiotto, *Spectrum of CHL dyons from genus-two partition function*, JHEP **12** (2007) 087, doi:10.1088/1126-6708/2007/12/087, arXiv:hep-th/0612011.

[27] A. Dabholkar and S. Nampuri, *Spectrum of dyons and black holes in CHL orbifolds using Borcherds lift*, JHEP **11** (2007) 077, doi:10.1088/1126-6708/2007/11/077, arXiv:hep-th/0603066.

[28] J. R. David and A. Sen, *CHL Dyons and Statistical Entropy Function from D1-D5 System*, JHEP **11** (2006) 072, doi:10.1088/1126-6708/2006/11/072, arXiv:hep-th/0605210.

[29] A. Dabholkar, F. Denef, G. W. Moore and B. Pioline, *Precision counting of small black holes*, JHEP **10** (2005) 096, doi:10.1088/1126-6708/2005/10/096, arXiv:hep-th/0507014.

[30] F. Denef and G. W. Moore, *Split states, entropy enigmas, holes and halos*, JHEP **11** (2011) 129, doi:10.1007/JHEP11(2011)129, arXiv:hep-th/0702146.

[31] N. Dorey, C. Fraser, T. J. Hollowood and M. A. Kneipp, *S-duality in $\mathcal{N} = 4$ supersymmetric gauge theories with arbitrary gauge group*, Phys. Lett. B **383** (1996) 422, doi:10.1016/0370-2693(96)00773-3, arXiv:hep-th/9605069.

[32] A. Dabholkar, J. Gomes and S. Murthy, *Quantum black holes, localization and the topological string*, JHEP **06** (2011) 019, doi:10.1007/JHEP06(2011)019, arXiv:1012.0265.

[33] A. Dabholkar, J. Gomes and S. Murthy, *Localization & Exact Holography*, JHEP **04** (2013) 062, doi:10.1007/JHEP04(2013)062, arXiv:1111.1161.

[34] J. R. David, D. P. Jatkar and A. Sen, *Product representation of Dyon partition function in CHL models*, JHEP **06** (2006) 064, doi:10.1088/1126-6708/2006/06/064, arXiv:hep-th/0602254.

[35] J. R. David, D. P. Jatkar and A. Sen, *Dyon spectrum in generic $\mathcal{N} = 4$ supersymmetric $\mathbb{Z}_N$ orbifolds*, JHEP **01** (2007) 016, doi:10.1088/1126-6708/2007/01/016, arXiv:hep-th/0609109.

[36] A. Dabholkar, S. Murthy and D. Zagier, *Quantum black holes, wall crossing, and mock modular forms*, preprint, arXiv:1208.4074.

[37] R. Dijkgraaf, E. P. Verlinde and H. L. Verlinde, *Counting dyons in $\mathcal{N} = 4$ string theory*, Nucl. Phys. B **484** (1997) 543, doi:10.1016/S0550-3213(96)00640-2, arXiv:hep-th/9607026.

[38] D. Bhattacharjee, *Which string vibrations become which generations on the Dwork quintic: code, checks and engine record*, version 1.1.0, Zenodo (2026), doi:10.5281/zenodo.23484427.

[39] I. Ellwood and M. Schnabl, *Proof of vanishing cohomology at the tachyon vacuum*, JHEP **02** (2007) 096, doi:10.1088/1126-6708/2007/02/096, arXiv:hep-th/0606142.

[40] T. Erler, *Analytic solution for tachyon condensation in Berkovits' open superstring field theory*, JHEP **11** (2013) 007, doi:10.1007/JHEP11(2013)007, arXiv:1308.4400.

[41] T. Erler and C. Maccaferri, *String Field Theory Solution for Any Open String Background*, JHEP **10** (2014) 029, doi:10.1007/JHEP10(2014)029, arXiv:1406.3021.

[42] T. Erler and M. Schnabl, *A Simple Analytic Solution for Tachyon Condensation*, JHEP **10** (2009) 066, doi:10.1088/1126-6708/2009/10/066, arXiv:0906.0979.

[43] G. N. Felder and L. Kofman, *Inhomogeneous fragmentation of the rolling tachyon*, Phys. Rev. D **70** (2004) 046004, doi:10.1103/PhysRevD.70.046004, arXiv:hep-th/0403073.

[44] G. N. Felder, L. Kofman and A. Starobinsky, *Caustics in tachyon matter and other Born–Infeld scalars*, JHEP **09** (2002) 026, doi:10.1088/1126-6708/2002/09/026, arXiv:hep-th/0208019.

[45] K. Fritzsch, C. Kottke and M. Singer, *Monopoles and the Sen Conjecture*, preprint, arXiv:1811.00601.

[46] E. Fuchs and M. Kroyter, *On the validity of the solution of string field theory*, JHEP **05** (2006) 006, doi:10.1088/1126-6708/2006/05/006, arXiv:hep-th/0603195.

[47] J. P. Gauntlett and J. A. Harvey, *S-duality and the spectrum of magnetic monopoles in heterotic string theory*, preprint, arXiv:hep-th/9407111.

[48] J. P. Gauntlett and D. A. Lowe, *Dyons and S-duality in $\mathcal{N} = 4$ supersymmetric gauge theory*, Nucl. Phys. B **472** (1996) 194, doi:10.1016/0550-3213(96)00218-0, arXiv:hep-th/9601085.

[49] L. Girardello, A. Giveon, M. Porrati and A. Zaffaroni, *S-duality in $\mathcal{N} = 4$ Yang–Mills theories with general gauge groups*, Nucl. Phys. B **448** (1995) 127, doi:10.1016/0550-3213(95)00177-T, arXiv:hep-th/9502057.

[50] V. A. Gritsenko and V. V. Nikulin, *Siegel automorphic form corrections of some Lorentzian Kac–Moody Lie algebras*, Amer. J. Math. **119** (1997) 181, doi:10.1353/ajm.1997.0002.

[51] P. Heidmann and S. Mondal, *The full space of BPS multicenter states with pure D-brane charges*, JHEP **06** (2019) 011, doi:10.1007/JHEP06(2019)011, arXiv:1810.10019.

[52] T. Hausel, E. Hunsicker and R. Mazzeo, *Hodge cohomology of gravitational instantons*, Duke Math. J. **122** (2004) 485, doi:10.1215/S0012-7094-04-12233-X, arXiv:math/0207169.

[53] N. Hitchin, *L^2 cohomology of hyperKahler quotients*, Commun. Math. Phys. **211** (2000) 153, doi:10.1007/s002200050806, arXiv:math/9909002.

[54] L. V. Iliesiu, S. Murthy and G. J. Turiaci, *Black hole microstate counting from the gravitational path integral*, JHEP **08** (2025) 152, doi:10.1007/JHEP08(2025)152, arXiv:2209.13602.

[55] L. V. Iliesiu, S. Murthy and G. J. Turiaci, *Revisiting the logarithmic corrections to the black hole entropy*, JHEP **07** (2025) 058, doi:10.1007/JHEP07(2025)058, arXiv:2209.13608.

[56] D. P. Jatkar and A. Sen, *Dyon spectrum in CHL models*, JHEP **04** (2006) 018, doi:10.1088/1126-6708/2006/04/018, arXiv:hep-th/0510147.

[57] A. Kapustin and E. Witten, *Electric-Magnetic Duality And The Geometric Langlands Program*, Commun. Num. Theor. Phys. **1** (2007) 1, doi:10.4310/CNTP.2007.v1.n1.a1, arXiv:hep-th/0604151.

[58] D. Karan, B. Khatun, B. Sahoo and A. Sen, *All order classical electromagnetic soft theorems*, JHEP **11** (2025) 025, doi:10.1007/JHEP11(2025)025, arXiv:2501.07328.

[59] C. Keeler, F. Larsen and P. Lisbao, *Logarithmic Corrections to $N \geq 2$ Black Hole Entropy*, Phys. Rev. D **90** (2014) 043011, doi:10.1103/PhysRevD.90.043011, arXiv:1404.1379.

[60] C. Kottke and F. Rochon, *L^2-Cohomology of quasi-fibered boundary metrics*, Invent. Math. **236** (2024) 1083, doi:10.1007/s00222-024-01253-5, arXiv:2103.16655.

[61] H. Krishna and B. Sahoo, *Universality of loop corrected soft theorems in 4d*, JHEP **11** (2023) 233, doi:10.1007/JHEP11(2023)233, arXiv:2308.16807.

[62] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson-Thomas invariants and cluster transformations*, preprint, arXiv:0811.2435.

[63] M. Kudrna and M. Schnabl, *Universal Solutions in Open String Field Theory*, preprint, arXiv:1812.03221.

[64] A. Laddha and A. Sen, *Classical proof of the classical soft graviton theorem in $D > 4$*, Phys. Rev. D **101** (2020) 084011, doi:10.1103/PhysRevD.101.084011, arXiv:1906.08288.

[65] K.-M. Lee, E. J. Weinberg and P. Yi, *Electromagnetic duality and $SU(3)$ monopoles*, Phys. Lett. B **376** (1996) 97, doi:10.1016/0370-2693(96)00286-9, arXiv:hep-th/9601097.

[66] N. D. Lambert, H. Liu and J. M. Maldacena, *Closed strings from decaying D-branes*, JHEP **03** (2007) 014, doi:10.1088/1126-6708/2007/03/014, arXiv:hep-th/0303139.

[67] Y. Matsuo and K. Okuyama, *BPS condition of string junction from M theory*, Phys. Lett. B **426** (1998) 294, doi:10.1016/S0370-2693(98)00288-3, arXiv:hep-th/9712070.

[68] C. Montonen and D. Olive, *Magnetic monopoles as gauge particles?*, Phys. Lett. B **72** (1977) 117, doi:10.1016/0370-2693(77)90076-4.

[69] D. R. Morrison and W. Taylor, *Classifying bases for 6D F-theory models*, Cent. Eur. J. Phys. **10** (2012) 1072, doi:10.2478/s11534-012-0065-4, arXiv:1201.1943.

[70] D. R. Morrison and W. Taylor, *Non-Higgsable clusters for 4D F-theory models*, JHEP **05** (2015) 080, doi:10.1007/JHEP05(2015)080, arXiv:1412.6112.

[71] S. Murthy and V. Reys, *Single-centered black hole microstate degeneracies from instantons in supergravity*, JHEP **04** (2016) 052, doi:10.1007/JHEP04(2016)052, arXiv:1512.01553.

[72] Y. Okawa, *Comments on Schnabl's analytic solution for tachyon condensation in Witten's open string field theory*, JHEP **04** (2006) 055, doi:10.1088/1126-6708/2006/04/055, arXiv:hep-th/0603159.

[73] H. Osborn, *Topological charges for $N = 4$ supersymmetric gauge theories and monopoles of spin 1*, Phys. Lett. B **83** (1979) 321, doi:10.1016/0370-2693(79)91118-3.

[74] R. Pius and A. Sen, *Cutkosky rules for superstring field theory*, JHEP **10** (2016) 024, doi:10.1007/JHEP10(2016)024; erratum JHEP **09** (2018) 122, doi:10.1007/JHEP09(2018)122, arXiv:1604.01783.

[75] M. Porrati, *On the existence of states saturating the Bogomolny bound in $\mathcal{N} = 4$ supersymmetry*, Phys. Lett. B **377** (1996) 67, doi:10.1016/0370-2693(96)00346-2, arXiv:hep-th/9505187.

[76] S.-J. Rey and J.-T. Yee, *BPS dynamics of triple (p, q) string junction*, Nucl. Phys. B **526** (1998) 229, doi:10.1016/S0550-3213(98)00401-5, arXiv:hep-th/9711202.

[77] M. Rosselló, *The Immortal Dyon Index Is Positive*, preprint, arXiv:2403.19630.

[78] L. Rastelli, A. Sen and B. Zwiebach, *String field theory around the tachyon vacuum*, Adv. Theor. Math. Phys. **5** (2002) 353, doi:10.4310/ATMP.2001.v5.n2.a5, arXiv:hep-th/0012251.

[79] L. Rastelli, A. Sen and B. Zwiebach, *Vacuum string field theory*, preprint, arXiv:hep-th/0106010.

[80] B. Sahoo and A. Sen, *Classical and Quantum Results on Logarithmic Terms in the Soft Theorem in Four Dimensions*, JHEP **02** (2019) 086, doi:10.1007/JHEP02(2019)086, arXiv:1808.03288.
[81] M. Schnabl, *Analytic solution for tachyon condensation in open string field theory*, Adv. Theor. Math. Phys. **10** (2006) 433, doi:10.4310/ATMP.2006.v10.n4.a1, arXiv:hep-th/0511286.
[82] J. H. Schwarz and A. Sen, *Duality symmetries of 4-D heterotic strings*, Phys. Lett. B **312** (1993) 105, doi:10.1016/0370-2693(93)90495-4, arXiv:hep-th/9305185.
[83] J. H. Schwarz and A. Sen, *Duality symmetric actions*, Nucl. Phys. B **411** (1994) 35, doi:10.1016/0550-3213(94)90053-1, arXiv:hep-th/9304154.
[84] G. Segal and A. Selby, *The cohomology of the space of magnetic monopoles*, Commun. Math. Phys. **177** (1996) 775, doi:10.1007/BF02099547.
[85] N. Seiberg, *Why is the matrix model correct?*, Phys. Rev. Lett. **79** (1997) 3577, doi:10.1103/PhysRevLett.79.3577, arXiv:hep-th/9710009.
[86] A. Sen, *Field theory of tachyon matter*, Mod. Phys. Lett. A **17** (2002) 1797, doi:10.1142/S0217732320008071, arXiv:hep-th/0204143.
[87] A. Sen, *Tachyon matter*, JHEP **07** (2002) 065, doi:10.1088/1126-6708/2002/07/065, arXiv:hep-th/0203265.
[88] A. Sen, *Rolling tachyon*, JHEP **04** (2002) 048, doi:10.1088/1126-6708/2002/04/048, arXiv:hep-th/0203211.
[89] A. Sen, *Dirac–Born–Infeld action on the tachyon kink and vortex*, Phys. Rev. D **68** (2003) 066008, doi:10.1103/PhysRevD.68.066008, arXiv:hep-th/0303057.
[90] A. Sen, *Open–closed duality at tree level*, Phys. Rev. Lett. **91** (2003) 181601, doi:10.1103/PhysRevLett.91.181601, arXiv:hep-th/0306137.
[91] A. Sen, *Tachyon dynamics in open string theory*, Int. J. Mod. Phys. A **20** (2005) 5513, doi:10.1142/S0217751X0502519X, arXiv:hep-th/0410103.
[92] A. Sen, *How does a fundamental string stretch its horizon?*, JHEP **05** (2005) 059, doi:10.1088/1126-6708/2005/05/059, arXiv:hep-th/0411255.
[93] A. Sen, *Black hole entropy function and the attractor mechanism in higher derivative gravity*, JHEP **09** (2005) 038, doi:10.1088/1126-6708/2005/09/038, arXiv:hep-th/0506177.
[94] A. Sen, *Stretching the horizon of a higher dimensional small black hole*, JHEP **07** (2005) 073, doi:10.1088/1126-6708/2005/07/073, arXiv:hep-th/0505122.
[95] A. Sen, *Three String Junction and $\mathcal{N} = 4$ Dyon Spectrum*, JHEP **12** (2007) 019, doi:10.1088/1126-6708/2007/12/019, arXiv:0708.3715.
[96] A. Sen, *Two centered black holes and $\mathcal{N} = 4$ dyon spectrum*, JHEP **09** (2007) 045, doi:10.1088/1126-6708/2007/09/045, arXiv:0705.3874.
[97] A. Sen, *Walls of Marginal Stability and Dyon Spectrum in $\mathcal{N} = 4$ Supersymmetric String Theories*, JHEP **05** (2007) 039, doi:10.1088/1126-6708/2007/05/039, arXiv:hep-th/0702141.
[98] A. Sen, *Quantum Entropy Function from AdS_2/CFT_1 Correspondence*, Int. J. Mod. Phys. A **24** (2009) 4225, doi:10.1142/S0217751X09045893, arXiv:0809.3304.
[99] A. Sen, *Negative discriminant states in $\mathcal{N} = 4$ supersymmetric string theories*, JHEP **10** (2011) 073, doi:10.1007/JHEP10(2011)073, arXiv:1104.1498.
[100] A. Sen, *How Do Black Holes Predict the Sign of the Fourier Coefficients of Siegel Modular Forms?*, Gen. Rel. Grav. **43** (2011) 2171, doi:10.1007/s10714-011-1175-9, arXiv:1008.4209.
[101] A. Sen, *Logarithmic Corrections to $\mathcal{N} = 2$ Black Hole Entropy: An Infrared Window into the Microstates*, Gen. Rel. Grav. **44** (2012) 1207, doi:10.1007/s10714-012-1336-5, arXiv:1108.3842.
[102] A. Sen, *Logarithmic Corrections to Rotating Extremal Black Hole Entropy in Four and Five Dimensions*, Gen. Rel. Grav. **44** (2012) 1947, doi:10.1007/s10714-012-1373-0, arXiv:1109.3706.
[103] A. Sen, *Logarithmic Corrections to Schwarzschild and Other Non-extremal Black Hole Entropy in Different Dimensions*, JHEP **04** (2013) 156, doi:10.1007/JHEP04(2013)156, arXiv:1205.0971.
[104] A. Sen, *Off-shell Amplitudes in Superstring Theory*, Fortsch. Phys. **63** (2015) 149, doi:10.1002/prop.201500002, arXiv:1408.0571.
[105] A. Sen, *Gauge Invariant 1PI Effective Superstring Field Theory: Inclusion of the Ramond Sector*, JHEP **08** (2015) 025, doi:10.1007/JHEP08(2015)025, arXiv:1501.00988.
[106] A. Sen, *BV Master Action for Heterotic and Type II String Field Theories*, JHEP **02** (2016) 087, doi:10.1007/JHEP02(2016)087, arXiv:1508.05387.
[107] A. Sen, *Unitarity of Superstring Field Theory*, JHEP **12** (2016) 115, doi:10.1007/JHEP12(2016)115, arXiv:1607.08244.
[108] A. Sen, *Subleading Soft Graviton Theorem for Loop Amplitudes*, JHEP **11** (2017) 123, doi:10.1007/JHEP11(2017)123, arXiv:1703.00024.
[109] A. Sen, *Fixing an Ambiguity in Two Dimensional String Theory Using String Field Theory*, JHEP **03** (2020) 005, doi:10.1007/JHEP03(2020)005, arXiv:1908.02782.
[110] A. Sen, *D-instantons, string field theory and two dimensional string theory*, JHEP **11** (2021) 061, doi:10.1007/JHEP11(2021)061, arXiv:2012.11624.

[111] A. Sen, *Normalization of type IIB D-instanton amplitudes*, JHEP **12** (2021) 146, doi:10.1007/JHEP12(2021)146, arXiv:2104.11109.
[112] A. Sen, *Normalization of D-instanton amplitudes*, JHEP **11** (2021) 077, doi:10.1007/JHEP11(2021)077, arXiv:2101.08566.
[113] A. Sen, *Revisiting localization for BPS black hole entropy*, JHEP **08** (2026) 174, doi:10.1007/JHEP08(2026)174, arXiv:2302.13490.
[114] A. Sen, *Rotating charged black hole solution in heterotic string theory*, Phys. Rev. Lett. **69** (1992) 1006, doi:10.1103/PhysRevLett.69.1006, arXiv:hep-th/9204046.
[115] A. Sen, *Strong–weak coupling duality in four-dimensional string theory*, Int. J. Mod. Phys. A **9** (1994) 3707, doi:10.1142/S0217751X94001497, arXiv:hep-th/9402002.
[116] A. Sen, *Dyon–monopole bound states, self-dual harmonic forms on the multi-monopole moduli space, and $SL(2,\mathbb{Z})$ invariance in string theory*, Phys. Lett. B **329** (1994) 217, doi:10.1016/0370-2693(94)90763-3, arXiv:hep-th/9402032.
[117] A. Sen, *F theory and orientifolds*, Nucl. Phys. B **475** (1996) 562, doi:10.1016/0550-3213(96)00347-1, arXiv:hep-th/9605150.
[118] A. Sen, *Orientifold limit of F theory vacua*, Phys. Rev. D **55** (1997) R7345, doi:10.1103/PhysRevD.55.R7345, arXiv:hep-th/9702165.
[119] A. Sen, *Tachyon condensation on the brane anti-brane system*, JHEP **08** (1998) 012, doi:10.1088/1126-6708/1998/08/012, arXiv:hep-th/9805170.
[120] A. Sen, *D0-branes on T^n and matrix theory*, Adv. Theor. Math. Phys. **2** (1998) 51, doi:10.4310/ATMP.1998.v2.n1.a2, arXiv:hep-th/9709220.
[121] A. Sen, *String network*, JHEP **03** (1998) 005, doi:10.1088/1126-6708/1998/03/005, arXiv:hep-th/9711130.
[122] A. Sen, *SO(32) spinors of type I and other solitons on brane–anti-brane pair*, JHEP **09** (1998) 023, doi:10.1088/1126-6708/1998/09/023, arXiv:hep-th/9808141.
[123] A. Sen, *Descent relations among bosonic D-branes*, Int. J. Mod. Phys. A **14** (1999) 4061, doi:10.1142/S0217751X99001901, arXiv:hep-th/9902105.
[124] A. Sen, *Non-BPS states and branes in string theory*, preprint, arXiv:hep-th/9904207.
[125] A. Sen, *Universality of the tachyon potential*, JHEP **12** (1999) 027, doi:10.1088/1126-6708/1999/12/027, arXiv:hep-th/9911116.
[126] A. Sen and B. Zwiebach, *String field theory: a review*, preprint, arXiv:2405.19421.
[127] A. Sen and B. Zwiebach, *Quantum background independence of closed string field theory*, Nucl. Phys. B **423** (1994) 580, doi:10.1016/0550-3213(94)90145-7, arXiv:hep-th/9311009.
[128] C. Vafa and E. Witten, *A Strong coupling test of S-duality*, Nucl. Phys. B **431** (1994) 3, doi:10.1016/0550-3213(94)90097-3, arXiv:hep-th/9408074.
[129] E. Witten, *D-branes and K-theory*, JHEP **12** (1998) 019, doi:10.1088/1126-6708/1998/12/019, arXiv:hep-th/9810188.
[130] D. Bhattacharjee and U. Bhattacharya, *Lattice Yang–Mills theory: spacing-uniform obstructions, the mass gap up to the Dobrushin ceiling and in a forest gauge, and Lean certificate*, version 3.3.1, Zenodo (2026), doi:10.5281/zenodo.23490028.
[131] J. Halverson, C. Long and B. Sung, *On the scarcity of weak coupling in the string landscape*, JHEP **02** (2018) 113, doi:10.1007/JHEP02(2018)113, arXiv:1710.09374.
[132] D. Karan, B. Khatun, R. Mahajan, B. Sahoo and A. Sen, *All-Orders Classical Soft Graviton Theorems in the Newtonian Limit and Beyond*, preprint (2026), arXiv:2610.07134.
[133] A. Sen, *Extended Supergravity Needs String Scale Cut-off*, preprint (2026), arXiv:2606.17149.
[134] S. Weinberg, *Infrared photons and gravitons*, Phys. Rev. **140** (1965) B516, doi:10.1103/PhysRev.140.B516.
[135] K. Prabhu, G. Satishchandran and R. M. Wald, *Infrared finite scattering theory in quantum field theory and quantum gravity*, Phys. Rev. D **106** (2022) 066005, doi:10.1103/PhysRevD.106.066005, arXiv:2203.14334.
[136] K. Prabhu and G. Satishchandran, *Infrared finite scattering theory: Amplitudes and soft theorems*, preprint (2024), arXiv:2402.18637.
[137] A. Sen, *Soft Theorems in Superstring Theory*, JHEP **06** (2017) 113, doi:10.1007/JHEP06(2017)113, arXiv:1702.03934.
[138] R. Akhoury, R. Saotome and G. Sterman, *Collinear and Soft Divergences in Perturbative Quantum Gravity*, Phys. Rev. D **84** (2011) 104040, doi:10.1103/PhysRevD.84.104040, arXiv:1109.0270.

Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL), Bhubaneswar, Odisha 751030, India

Formerly, University of Calcutta, Senate House, 87/1 College Street, Kolkata 700073, West Bengal, India
Email address: d.bhattacharjee@erl-forschung.de
Email address: itsdeep@live.com
Email address: egspl@proton.me