

Finite quantum gravity from a scale-invariant gas of branes

Deep Bhattacharjee ^{1,*}

¹Formerly, *Electro-Gravitational Space Propulsion Laboratory (EGSPL)*, Bhubaneswar, Odisha 751030, India
(Dated: October 7, 2026)

Nonlocal gravity with an entire, zero-free form factor has no ghosts and can be power-counting finite, but the form factor has so far been put in by hand. We derive it from a Poisson gas of branes with no preferred size, each multiplying the graviton kinetic operator by a passive, saturating factor without resonances. When the strength n per e -fold of size is shared among ever more and ever weaker small branes, almost every frozen realization of the gas yields an entire form factor without zeros, $\exp[\frac{n}{2} \text{Ein}(\Box^2/M^4)]$ times a random factor that tends to a constant at high momentum; no average over the gas is taken. The propagator keeps only the graviton pole. For $n \geq 4$ every graph with two or more loops converges, Newton's constant and the cosmological constant do not run, and three operators quartic in curvature cancel the one-loop poles through an exactly computed triangular matrix. The S matrix is then finite and unitary at every loop order, so the theory is ultraviolet complete in perturbation theory. The graviton zero-point energy is finite, with mean $3nM^4/64\pi^2$.

Keywords: nonlocal quantum gravity; ultraviolet finiteness; brane gas

I. INTRODUCTION

Einstein gravity quantized around flat space fails by power counting. The propagator falls like p^{-2} , the vertices grow like p^2 , and the superficial degree of divergence of an L -loop graph in four dimensions is $2 + 2L$. Pure gravity is finite at one loop on shell, but not once matter is added [1], and the two-loop counterterm of pure gravity does not vanish [2]. Curvature-squared terms make the theory renormalizable at the price of a massive spin-two state with negative residue [3]. More derivatives do not cure this. A propagator $1/(p^2 a(p^2))$ with a a polynomial has as many extra poles as a has zeros, their residues add up to -1 , and at least one of them is a ghost [4].

A way around the problem goes back to Efimov [5] and was developed for gauge theories and gravity by Krasnikov, Kuz'min, Tomboulis, Biswas, Mazumdar, Siegel, Modesto and others [6–12]. One replaces the polynomial by an entire function without zeros, $a = e^H$ with H entire. The propagator then keeps a single pole, at $p^2 = 0$, while a can still grow like a power at large Euclidean momenta, and with enough growth the theory is super-renormalizable. Modesto and Rachwał showed further that the one-loop divergences which remain can be removed with a few operators of higher order in curvature, so that all beta functions vanish [12].

In all of this H is an input. Tomboulis proposed a family built from the incomplete gamma function because it has the right asymptotics [8], and later work took that family or variants of it. Nothing in the construction says where such a function should come from. A polynomial, which is what any finite number of higher-derivative corrections produces, is precisely what has to be avoided.

This paper supplies a mechanism that produces the required class. The idea was already present in an ear-

lier study of brane arrangements [13]: when independent scatterers act by multiplication, the average over a Poisson ensemble of them is an exponential, because the expectation of a product over a Poisson process is the exponential of an integral. That study took the single-brane factor to first order in momentum and a gas of finite intensity, which gives the Gaussian $e^{-\Box/M^2}$. Its vertices grow exponentially and power counting is not available for it [14]. Here we keep the full single-brane factor, require it to be passive, to saturate and to have no resonances, and let the strength of the gas be spread with no preferred size below the largest brane. We take no average. The graviton crosses one realization of the gas, each brane contributes the exponential of its response, and the logarithms add. The gas contains infinitely many small branes, and the form factor of a single realization comes out as

$$a(z) = \exp\left[\frac{n}{2} \text{Ein}\left(\frac{z^2}{M^4}\right) + R(z)\right], \quad (1)$$

with $\text{Ein}(w) = \int_0^w (1 - e^{-t}) dt/t$ the entire exponential integral, $z = p^2$ in Euclidean signature, n the strength of the gas per e -fold of size, and R a random entire function with mean zero that tends to a finite constant R_∞ at large $|z|$ around the real axis. The first term is fixed by the density of the gas alone, and for $n = 1$ it is Tomboulis's form factor with the simplest choice of his polynomial.¹

¹ Tomboulis writes the form factor as e^H with $H(z) = \frac{1}{2}[\gamma_E + \Gamma(0, p(z)^2) + \log p(z)^2]$ for a polynomial p [8]. Since $\text{Ein}(y) = \log y + \gamma_E + E_1(y)$ and $E_1(y) = \Gamma(0, y)$, the choice $p(z) = z/M^2$ gives $H = \frac{1}{2} \text{Ein}(z^2/M^4)$, which is the first term of Eq. (1) at $n = 1$. For larger n the gas gives the n -th power of that function, not Tomboulis's function for a polynomial of higher degree. Briscese and Modesto write the family with an overall factor, $H = \frac{\alpha}{2}[\gamma_E + \Gamma(0, p^2) + \log p^2]$ [15], and the first term is their case $\alpha = n$, $p(z) = z/M^2$.

* itsdeep@live.com

We call it the skeleton. The second term records which branes the realization happens to contain. Its growth, $a \simeq c|z/M^2|^n$ with $c = e^{n\gamma_E/2+R_\infty}$, has the exponent of the skeleton and a random coefficient.

Whether a single realization, rather than an average over many, can carry a consistent field theory is the central question, and it has two parts. The first is the zeros. A product of brane factors vanishes wherever one factor does, and the simplest saturating factor has infinitely many complex zeros; a frozen gas of such branes would give the propagator infinitely many complex poles. We show that a brane without resonances, whose factor has no zeros, is exactly a brane whose factor is the exponential of a saturating response, and that with such branes almost every realization is free of zeros. The second part is the asymptotics. In a gas of identical branes the strength above a given size fluctuates like a random walk in the logarithm of the size, and $a/|z|^n$ has no limit at all. If instead the strength per e -fold is shared among ever more and ever weaker branes as the size decreases, the fluctuations die out, and they die out fast enough for the field theory when their variance below size t falls faster than t^{2n} . That condition, (G) of Sec. IIB, is the one new input of the construction; Sec. IIG proves that under it almost every realization has every property used later, with explicit error bounds.

Besides the derivation of Eq. (1), the paper contains the following results, all derived in the text. The form factor of a realization is pinched between two multiples of $(1+x^2)^{n/2}$, with $x = z/M^2$, along the whole real axis of p^2 , Euclidean and Lorentzian, and in the cones of condition (S4), $|\arg(\pm z)| < \pi/4$ for the simplest response, it coincides with the monomial $c(z/M^2)^n$ up to a relative correction that falls faster than $|M^2/z|^n$. Every vertex of the gauge-fixed theory obeys a bound in terms of the momenta on its legs, proved by expanding functions of the covariant d'Alembertian in divided differences. A vertex grows no faster than the $(2n+2)$ th power of its largest momentum, as dimensional analysis requires, or than one power for each hard leg, whichever is larger, and with this every graph with two or more loops converges for $n \geq 4$. The one-loop poles are those of a local theory with a single overall coupling, which is why neither Newton's constant nor the cosmological constant is renormalized. The three poles that remain, in R^2 , $R_{\mu\nu}R^{\mu\nu}$ and the Gauss-Bonnet density E , are shifted by three operators quartic in curvature through a matrix that we compute exactly; it is triangular with determinant $-24/c^3$, so all three poles can be cancelled at once. After that the bare couplings are finite numbers that do not depend on the regulator, and nothing is sent to infinity at any stage. The one-loop vacuum energy, which has no pole, can be evaluated in closed form through a Mellin transform of Ein: the zero-point energy density of the graviton has mean $3nM^4/(64\pi^2)$ over realizations, a relative spread that we give in closed form, and the same value for every regulator. After a field redefinition that makes the propagators local, the cutting rules proved by Pius and

Sen and by Brischese and Modesto for nonlocal theories defined from Euclidean signature [15, 16] hold for the gravitational theory, and with the quartet mechanism of Kugo and Ojima [17] they make the physical S matrix unitary at every loop order. Graviton scattering at tree level is that of general relativity, while the gravitational scattering of matter at tree level never exceeds a number of order GM^2 at fixed angle, at any energy. Finally the Newtonian potential is finite at the origin and approaches Newton's law faster than any power of $1/(Mr)$; for the skeleton its depth is pinned between two gamma-function expressions whose ratio is at most $1 + 1/(2n-1)$.

The brane picture enters through two premises, which we keep in view. The first, A1, concerns a single brane: it acts on the graviton multiplicatively, through a factor that depends on the momentum only through p^2 , has no resonances, and is passive and saturating in the sense of five conditions (Sec. IIA). The field theory is proved for every response in that class, not for one chosen function (Sec. IIF). The second, A2, concerns the gas: branes are placed independently, their expected strength has no preferred size, and the strength is fine grained in the sense of condition (G) (Sec. IIB). Neither premise can be derived within gravity; both are statements about branes, which a microscopic computation could test. Everything after them is a statement about entire functions, Poisson processes or Feynman integrals, and holds for almost every realization of the gas. We also use the standard machinery of perturbative renormalization (dimensional regularization, the background-field method and the cohomological classification of counterterms) and cite it where it enters. By "finite" we mean the absence of ultraviolet divergences in perturbation theory around flat space, and Sec. VIID states in what sense this makes the theory ultraviolet complete and what it leaves open.²

Figure 1 shows how the pieces depend on one another. Section II sets up the gas, derives Eq. (1) and proves that almost every realization has the properties the field theory needs. Section III writes the action and the propagator. Section IV bounds the vertices and counts powers. Section V treats one loop and the quartic operators. Section VI assembles the all-order statement, explains why it does not rest on a cutoff, and computes the vacuum energy. Section VII proves unitarity, bounds the size of tree amplitudes and says in what sense the theory is ultraviolet complete, Sec. VIII gives the Newtonian limit, and Sec. IX lists what is assumed and what is open. The appendixes contain conventions, the angular averages behind the one-loop coefficients, the proofs about divided differences, and a list of the independent numerical checks.

² Infrared divergences are a separate matter. The graviton is massless and on-shell amplitudes have the soft divergences of general relativity, which cancel in inclusive rates [18]. The form factor does not change the infrared, since $a = 1 + O(p^4/M^4)$ at small momenta. Section VI collects every kind of divergence and states which are removed and why.




FIG. 1. Logical structure of the paper, in three layers. The top layer holds the premises about branes (dashed boxes): the response of one brane, A1, and the gas, A2, whose fine-graining condition (G) is what lets a single realization carry the theory; without it, for identical branes, the form factor has no power-law asymptotics (box on the right). The middle layer holds the properties of the form factor of one realization, derived for every admissible response and almost every realization (Secs. IID–IIG), with the closed-form skeleton for ψ_* as one instance. The bottom layer holds the field theory, which uses the form factor only through the middle layer and is proved for $n \geq 4$; the box with the heavy frame is the main result. Arrows point from what is used to what is derived.

II. A GAS OF BRANES WITH NO PREFERRED SIZE

We work in Euclidean signature, with $z = p^2 \geq 0$ for real momenta, $\square \rightarrow -p^2$, $x = z/M^2$ on the real axis and $w = z/M^2$ in the complex plane. Lorentzian statements follow by continuing z to negative values; the response used for the closed forms depends on z^2 , so for it the two half-axes behave alike.

A. The response of one brane

Consider a graviton mode crossing a stack of thin branes. Let t_i and r_i be the forward transmission and reflection amplitudes of brane i , and δ the phase accumulated between neighbors. Summing the multiple reflections between two branes gives $t_{12} = t_1 t_2 / (1 - r'_1 r_2 e^{2i\delta})$, where r'_1 is the reflection amplitude of the first brane seen from the right. In the corresponding expansion for N branes every term other than $\prod_i t_i$ contains at least one round trip, and hence two reflection amplitudes, so that

$$t_{1\dots N} = \left(\prod_{i=1}^N t_i \right) \left[1 + O\left(\sum_{i < j} |r_i| |r_j| \right) \right]. \quad (2)$$

This is the composition rule of multiple scattering [19, 20], written out for branes in Ref. [13]. For weakly reflecting branes the kinetic operator of the graviton is dressed multiplicatively, $p^2 \rightarrow p^2 \prod_i t_i^{-1}$, and the problem re-

duces to the single factor t_i^{-1} .³ We do not derive that factor from a brane action. We state what we require of it, and call the requirement assumption A1. Sections IIF and IIH show that the field theory needs nothing more than these conditions.

A brane has a size $s > 0$, with dimensions of length squared, and a strength $\vartheta > 0$, and it multiplies the spin-two and spin-zero parts of the graviton kinetic operator by

$$t^{-1} = e^{\vartheta\psi(sz)}, \quad (3)$$

where

- (S1) ψ is entire and $\psi(0) = 0$;
- (S2) ψ is real on the real axis, with $0 \leq \psi(t) \leq 1$ there;
- (S3) $1 - \psi$ and all its derivatives decay along the real axis faster than any power of $1/|t|$;
- (S4) the decay in (S3) holds uniformly in closed sectors around the real axis: there is an opening $\theta > 0$ such that for every $\delta \in (0, \theta)$ and every N , $|1 - \psi(w)| \leq C_{\delta,N} |w|^{-N}$ for $|w| \geq 1$ and $|\arg(\pm w)| \leq \theta - \delta$.

³ The gases below contain infinitely many branes, most of them much smaller than the wavelength of a given mode. The correction in Eq. (2) is bounded by $(\sum_i |r_i|)^2$, which stays small if the reflection of a brane falls off for modes much longer than the brane at least as fast as its effect on the kinetic operator does. We take this as part of the premise. The sum of the effects themselves converges almost surely (Sec. IID).

On the real axis the factor (3) rises from 1, for modes much longer than the brane, to e^ϑ , for modes much shorter, so ϑ is the saturated logarithmic strength of the brane. Condition (S1) says that a brane has a single length scale and leaves long modes untouched. Condition (S2) is passivity: along the real axis, spacelike and timelike alike, a brane can suppress a mode but never amplify it. Condition (S3) says that the suppression saturates once the mode is much shorter than the brane. Condition (S4) says that saturation is not special to real momenta: a mode whose squared momentum is slightly complex, as it is for the loop energies of Sec. VII near the ends of their contours, is saturated as well.⁴ Dependence on the momentum through p^2 alone requires the branes to respect four-dimensional Poincaré invariance, as branes that fill the observed four dimensions and are scattered in the transverse ones do.

The exponential in Eq. (3) is not a separate choice. It is equivalent to a fifth condition,

(S5) the factor t^{-1} has no zeros in the complex plane of z ,

which says that a brane has no resonances: a zero of t^{-1} is a pole of the transmission amplitude at complex momentum, a quasi-bound state of the brane. Suppose a brane multiplies the kinetic operator by $1 + \varphi(sz)$, with φ entire, real on the real axis, $\varphi(0) = 0$, and $0 \leq \varphi \leq \varphi_\infty$ on the real axis, saturating at φ_∞ as in (S3) and (S4). If $1 + \varphi$ has no zeros it is the exponential of an entire function, $1 + \varphi = e^h$ with $h(0) = 0$ [21], and $h = \log(1 + \varphi)$ on the real axis, where $1 + \varphi > 0$. Put $\vartheta = \log(1 + \varphi_\infty)$ and $\psi = h/\vartheta$. Then ψ satisfies (S1) and (S2), and $1 - \psi = -\vartheta^{-1} \log[1 - (\varphi_\infty - \varphi)/(1 + \varphi_\infty)]$ is bounded by $2|\varphi_\infty - \varphi|/\vartheta$ wherever $|\varphi_\infty - \varphi| \leq \frac{1}{2}(1 + \varphi_\infty)$, which gives (S3) and (S4).⁵ Conversely $e^{\vartheta\psi}$ is entire and has no zeros. Conditions (S1)–(S5) on a factor $1 + \varphi$ are therefore the same as (S1)–(S4) on the exponent ψ of Eq. (3).

Condition (S5) is what lets a single realization of the gas define a theory, and it fails for the most obvious saturating factor. The function $1 + \varphi = 2 - e^{-s^2 z^2}$ obeys every other condition, but it vanishes when

$$s^2 z^2 = -\log 2 - 2\pi i k, \quad k \in \mathbb{Z}. \quad (4)$$

These resonances lie in $\pi/4 < |\arg z| < 3\pi/4$, away from both real half-axes, but there are infinitely many of them, and a product over a gas of such branes vanishes at the

resonances of every brane in it. A gas with n such branes per e -fold of size below $1/M^2$ has on average $nR^2/(\pi M^4)$ zeros in the disk $|z| < R$ for $R \gg M^2$,⁶ each a complex pole of the propagator. Complex pairs of this kind are what Lee and Wick [22] and, in a modern form, Anselmi and Piva [23, 24] proposed to treat with a modified prescription. Branes obeying (S5) have none, and no such prescription is needed.

The simplest response with all five properties, and the one we use from Sec. IIE on, has the exponent

$$\psi_*(w) = 1 - e^{-w^2}, \quad t^{-1} = \exp[\vartheta(1 - e^{-s^2 z^2})]. \quad (5)$$

Its derivative expansion is $t^{-1} = 1 + \vartheta s^2 z^2 + O(z^4)$. The more obvious exponent $1 - e^{-w}$, which would give the incomplete gamma function directly, violates (S2): on the negative axis it is unbounded below, so the factor falls to zero faster than any power along the timelike axis and no polynomial bound holds there.⁷

Conditions (S1) and (S2) already have a consequence. On the real axis $\psi \geq 0 = \psi(0)$, so the origin is a minimum of a smooth function and $\psi'(0) = 0$. The leading correction that a passive brane makes to the kinetic operator is therefore of order $s^2 z^2$ and not sz : a passive brane does not generate a curvature-squared term at leading order.

B. Sizes and strengths

A gas is a random set of branes, each with a size and a strength. We impose three requirements on it: branes are placed independently of one another; below the size $S = 1/M^2$ of the largest brane there is no preferred size; and the strength of the gas is fine grained, in a sense made precise below. Together they form assumption A2.

The first requirement makes the set of sizes a Poisson process. A point process without multiple points and without fixed atoms, whose counts in disjoint sets are independent, is a Poisson process with some intensity measure ν [25, 26].⁸ We let the strength depend on the size, $\vartheta = \vartheta(s)$.⁹

⁶ A brane of size s contributes the zeros with $|\log 2 + 2\pi i k| \leq s^2 R^2$ inside the disk $|z| \leq R$, about $2s^2 R^2/\pi$ of them counting both signs of z . Averaging with $n ds/s$ over $s \leq M^{-2}$ gives $nR^2/(\pi M^4)$. The numerical check in Appendix D reproduces this count to 0.03% at $R = 60M^2$.

⁷ The exponent of the gas, computed as in Sec. IID, is then $n \text{Ein}(z/M^2)$, and along the negative real axis $\text{Ein}(-t) = -\int_0^t (e^u - 1) du/u$ tends to $-\infty$ faster than any power. Passivity keeps the two half-axes symmetric.

⁸ This is the theorem of Rényi and Kingman: complete independence plus simplicity characterizes the Poisson process. The intensity measure $\nu(A)$ is the expected number of branes with size in A .

⁹ Strengths drawn independently from a law that depends on s make the gas a marked Poisson process [25]. The mean and variance formulas below then hold with ϑ and ϑ^2 replaced by their conditional means at fixed s , and the exponential formulas with e^h replaced by its conditional mean.

⁴ Condition (S4) does not follow from (S3). The entire function $\psi(w) = 1 - e^{-w^2} \cos^2(e^w - 1)$ satisfies (S1)–(S3), but off the real axis $|\cos^2(e^w - 1)|$ grows like $\exp(2e^{\text{Re } w} |\sin \text{Im } w|)$, which defeats the Gaussian in every sector. For the response ψ_* used below the opening is $\theta = \pi/4$.

⁵ In a sector, at large $|w|$, $h - \vartheta$ equals the principal logarithm of $(1 + \varphi)/(1 + \varphi_\infty)$ plus $2\pi i k$ with an integer k that is constant on the connected set where the ratio is close to 1; that set touches the real axis, where $k = 0$.

The second requirement concerns the quantity that enters the kinetic operator, which is the strength. The expected strength carried by the branes with sizes in a set A , $\int_A \vartheta d\nu$, must not change under a dilation $s \rightarrow \lambda s$ whenever both A and λA lie in $(0, S]$. In the variable $u = \log s$ this is invariance under translations along the half-line $u \leq \log S$, so the measure of an interval depends only on its length; it is additive and monotone in that length, hence linear in it, and $\vartheta d\nu$ is a multiple of Lebesgue measure in u .¹⁰ Therefore

$$\vartheta(s) \nu(ds) = n \frac{ds}{s} \quad \text{on } (0, S], \quad n > 0, \quad (6)$$

and n is the expected strength per e -fold of size. For identical branes, $\vartheta = \vartheta_0$, this says that the number of branes per e -fold, n/ϑ_0 on average, is the same at all sizes.

The third requirement is about how that strength is shared. It can be carried by a few strong branes or by many weak ones. Write $\tau = sM^2 \in (0, 1]$ and

$$V(t) = \frac{1}{n} \int_{\tau \leq t} \vartheta^2 d\nu = \int_0^t \vartheta(\tau) \frac{d\tau}{\tau}. \quad (7)$$

We show in Sec. II C that $nV(t)$ is the variance of the strength carried by the branes smaller than t/M^2 , measured from its mean. The requirement is

$$(G) \quad V(t) \leq C t^\sigma \text{ for } 0 < t \leq 1, \text{ with some } \sigma > 2n.$$

In words: the fluctuation of the strength carried by small branes vanishes with their size faster than $(\text{size})^n$. It does not affect the mean, which Eq. (6) fixes. The simplest example is a power law,

$$\vartheta(\tau) = \vartheta_0 \tau^\zeta, \quad \nu(d\tau) = \frac{n}{\vartheta_0} \tau^{-\zeta} \frac{d\tau}{\tau}, \quad V(t) = \frac{\vartheta_0 t^\zeta}{\zeta}, \quad (8)$$

for which (G) holds with $\sigma = \zeta$ when $\zeta > 2n$. As the size decreases by one e -fold the branes become weaker by a factor $e^{-\zeta}$ and e^ζ times as numerous, so the strength per e -fold stays n while the gas becomes ever finer. Any profile that falls faster than every power, such as $\vartheta_0 \exp[-(\log \tau)^2]$, satisfies (G) for every σ . A gas of identical branes, $\zeta = 0$, has $V \equiv \infty$ and violates (G) completely; Sec. II G shows what goes wrong with it. Figure 3(a) shows the two kinds of gas, and Fig. 2 shows the bookkeeping.

C. Sums and products over the gas

Because the gas has infinitely many points, we derive the formulas we need instead of quoting them. Let Π be




FIG. 2. How a fine-grained gas shares its strength. Each column is one e -fold of size and carries expected strength n (dashed line), Eq. (6). The number of branes in a column, drawn as slabs, grows toward small sizes while each slab gets thinner, as for the power law (8). A mode with $z = p^2$ saturates the branes larger than $1/z$, whose total strength is $n \log x$ plus a fluctuation that converges to R_∞ ; the branes smaller than $1/z$ act only through a remainder whose variance is $nV(1/x)$ (Sec. II G).

a Poisson process with σ -finite intensity ν and let f be a complex function with $\int |f| d\nu < \infty$. Then

$$\mathbb{E} \prod_{x \in \Pi} (1 + f(x)) = \exp \int f d\nu, \quad (9)$$

with the product converging absolutely almost surely. Suppose first that the total mass $\lambda = \nu(\text{all})$ is finite. The number N of points is then Poisson with mean λ , and given N the points are independent with law ν/λ [26]. Conditioning on N ,

$$\begin{aligned} \mathbb{E} \prod_{x \in \Pi} (1 + f) &= \sum_{N \geq 0} e^{-\lambda} \frac{\lambda^N}{N!} \left(\frac{1}{\lambda} \int (1 + f) d\nu \right)^N \\ &= e^{-\lambda} e^{\lambda \int f d\nu}, \end{aligned}$$

which is Eq. (9); the same computation with $|f|$ in place of f gives $\mathbb{E} \prod (1 + |f|) = \exp \int |f| d\nu$, so the expectation exists. If ν has infinite mass, choose sets $A_1 \subset A_2 \subset \dots$ of finite measure that exhaust the space. Campbell's theorem, $\mathbb{E} \sum_{x \in \Pi} g(x) = \int g d\nu$ for $g \geq 0$ [25], applied to $g = |f|$, shows that $\sum_{x \in \Pi} |f(x)| < \infty$ almost surely, so the product converges absolutely and is the limit of the products over $\Pi \cap A_k$. These partial products are bounded in modulus by $\prod_{x \in \Pi} (1 + |f(x)|)$, whose expectation is $\exp \int |f| d\nu$ by monotone convergence and the finite case. Dominated convergence then gives $\mathbb{E} \prod_{x \in \Pi} (1 + f) = \lim_k \exp \int_{A_k} f d\nu = \exp \int f d\nu$.

Two consequences are used below. Replace f by $e^{uf} - 1$ with real f and real u in the finite case and differentiate twice at $u = 0$: the sum $\sum_{x \in \Pi} f(x)$ has mean $\int f d\nu$ and variance $\int f^2 d\nu$. For complex f , applying this to the

¹⁰ Additivity gives $\mu(h_1 + h_2) = \mu(h_1) + \mu(h_2)$ for the measure $\mu(h)$ of an interval of length h , and a monotone solution of Cauchy's equation is linear. This is the uniqueness of Haar measure on the multiplicative group $(0, \infty)$, restricted to the cut-off range.

real and imaginary parts and to their sum gives

$$\mathbb{E} \left| \sum_{x \in \Pi} f(x) - \int f d\nu \right|^2 = \int |f|^2 d\nu. \quad (10)$$

Write the left side as $\mathbb{E} |\int f d(\Pi - \nu)|^2$. If $\int |f|^2 d\nu < \infty$ but $\int |f| d\nu = \infty$, the compensated sums over $\Pi \cap A_k$ still converge: their increments over $A_{k+1} \setminus A_k$ are independent, have mean zero and have summable variances, so they converge almost surely and in mean square by Kolmogorov's criterion [27]. We call the limit $\int f d(\Pi - \nu)$; it has mean zero and obeys Eq. (10). With $f = \vartheta$ on the branes smaller than t/M^2 and $f = 0$ elsewhere, Eq. (10) shows that $nV(t)$, Eq. (7), is the variance of their total strength measured from its mean. The same exhaustion applied to Eq. (9) with $f = e^h - 1$ gives, for real h with $|h| \leq h_{\max}$ and $\int h^2 d\nu < \infty$,

$$\mathbb{E} \exp \int h d(\Pi - \nu) = \exp \int (e^h - 1 - h) d\nu. \quad (11)$$

On each A_k this is Eq. (9) with $f = e^h - 1$, multiplied by $\exp(-\int_{A_k} h d\nu)$. The integrand on the right is at most $\frac{1}{2}e^{h_{\max}}h^2$, so the right side is finite, and the same bound with $2h$ in place of h shows that $\mathbb{E} \exp(2 \int_{A_k} h d(\Pi - \nu)) \leq \exp(2e^{2h_{\max}} \int h^2 d\nu)$ uniformly in k . The exponentials are therefore uniformly integrable, and their means converge to the mean of the limit.

D. The form factor of one realization

A realization Π of the gas is one draw of the branes, frozen. A graviton crossing it is dressed by the product of the factors (3) of all its branes,

$$a_{\Pi}(z) = \prod_{s \in \Pi} e^{\vartheta(s)\psi(sz)} = \exp L_{\Pi}(z), \quad (12)$$

$$L_{\Pi}(z) = \sum_{s \in \Pi} \vartheta(s) \psi(sz).$$

The sum converges. Because ψ is entire and vanishes at the origin, $|\psi(w)| \leq A_R|w|$ on the disk $|w| \leq R$, and with Eq. (6) Campbell's theorem gives

$$\mathbb{E} \sum_{s \in \Pi} \sup_{|z| \leq RM^2} |\vartheta(s)\psi(sz)| \leq n \int_0^1 A_R R \tau \frac{d\tau}{\tau} = n A_R R < \infty.$$

Almost surely, then, the sum in Eq. (12) converges absolutely and uniformly on every disk, its limit L_{Π} is an entire function, and $a_{\Pi} = e^{L_{\Pi}}$ is an entire function *without zeros*. It satisfies $a_{\Pi}(0) = 1$, its Taylor coefficients are real, and $a_{\Pi} \geq 1$ on the whole real axis by (S2). A frozen gas of branes that obey (S5) therefore has no extra poles of any kind, real or complex. Condition (S5) is necessary as well as sufficient for this: if one factor vanished

at some z_0 , the convergent product of the others would be finite there and $a_{\Pi}(z_0)$ would vanish.

The mean of the exponent is fixed by Eq. (6) alone. By Campbell's theorem, $\mathbb{E} L_{\Pi}(z) = \int \vartheta \psi(sz) d\nu = n \int_0^{1/M^2} \psi(s) ds/s$, and the substitution $s = \tau/M^2$ gives

$$a_{\Pi}(z) = e^{n\Phi(z/M^2)} e^{R_{\Pi}(z)}, \quad (13)$$

$$R_{\Pi}(z) = \int \vartheta(s) \psi(sz) (\Pi - \nu)(ds),$$

$$\Phi(w) = \int_0^1 \psi(w\tau) \frac{d\tau}{\tau} = \sum_{k \geq 1} \frac{\psi_k}{k} w^k, \quad (14)$$

where $\psi(w) = \sum_{k \geq 1} \psi_k w^k$. The first factor in Eq. (13) is deterministic. It depends on the gas only through n and not at all on how the strength is shared among branes; we call

$$\hat{a}(z) = \exp \left[n \Phi \left(\frac{z}{M^2} \right) \right] \quad (15)$$

the skeleton of the form factor. The second factor is random, with $\mathbb{E} R_{\Pi}(z) = 0$; it is the subject of Sec. II G. The skeleton is the exponential of the mean of $\log a_{\Pi}$. It is not the mean of a_{Π} , which for real z and bounded strengths is, by Eq. (11), $\hat{a}$ times $\exp \int (e^{\vartheta\psi} - 1 - \vartheta\psi) d\nu$, and nothing in this paper averages the kinetic operator over the gas. The graviton sees one realization, and the theory is that of a_{Π} .

The Taylor series of ψ converges uniformly on $\{|w\tau| \leq R\}$ and may be integrated term by term, and since the coefficients of Φ are bounded by those of ψ its radius of convergence is infinite. So Φ is entire with $\Phi(0) = 0$, and $\hat{a}$ is entire, has no zeros and satisfies $\hat{a}(0) = 1$.

The growth of the skeleton on the real axis follows from (S3). For real $x > 0$, $\Phi(x) = \int_0^x \psi(t) dt/t$. Splitting the range at $t = 1$ and writing $\psi = 1 - (1 - \psi)$ on $[1, x]$,

$$\Phi(x) = \log |x| + C_{\pm} + \rho_{\pm}(x), \quad x \rightarrow \pm\infty, \quad (16)$$

with

$$C_{\pm} = \int_0^1 \psi(\pm t) \frac{dt}{t} - \int_1^{\infty} (1 - \psi(\pm t)) \frac{dt}{t},$$

$$\rho_{\pm}(x) = \int_{|x|}^{\infty} (1 - \psi(\pm t)) \frac{dt}{t} = O(|x|^{-N}) \text{ for every } N.$$

The case $x < 0$ is the same after the substitution $t = -x\tau$; both integrals in $C_{\pm}$ converge by (S1) and (S3). Exponentiating,

$$\hat{a}(z) = e^{nC_{\pm}} \left| \frac{z}{M^2} \right|^n \left[1 + O \left(\left| \frac{M^2}{z} \right|^N \right) \right], \quad z \rightarrow \pm\infty. \quad (17)$$

The power n depends on the gas only through its density of strength and on the shape of ψ not at all. Figure 3(b) shows how this happens. At a given z , the branes with $sz \gtrsim 1$ each contribute their full strength to L_{Π} and the

others almost nothing, so L_Π counts the strength in the e -folds of size above $1/z$, of which there are $\log(z/M^2)$, each holding strength n on average.

From here on n is a positive integer. Nothing above needs this, but the local theory of Sec. V does, because its kinetic term must be a polynomial in $\square$.

E. The skeleton in closed form

For the response (5) everything about the skeleton can be written in closed form. The substitution $u = w^2 \tau^2$ in Eq. (14) gives

$$\Phi(w) = \frac{1}{2} \int_0^{w^2} \frac{1 - e^{-u}}{u} du = \frac{1}{2} \text{Ein}(w^2), \quad (18)$$

so that $\hat{a}$ is the function of Eq. (1). The entire function Ein has the series $\text{Ein}(w) = \sum_{k \geq 1} (-1)^{k+1} w^k / (k k!)$, and we need three facts about it and about the exponential integral $E_1(y) = \int_y^\infty e^{-t} dt/t$ [28].

First, Euler's constant is

$$\gamma_E = \int_0^1 \frac{1 - e^{-t}}{t} dt - \int_1^\infty \frac{e^{-t}}{t} dt. \quad (19)$$

To see this, start from $\Gamma'(1) = -\gamma_E$, that is $-\gamma_E = \int_0^\infty e^{-t} \log t dt$, and integrate by parts on each half of the range: on $[0, 1]$ with the antiderivative $1 - e^{-t}$ of e^{-t} , so that the boundary term $(1 - e^{-t}) \log t$ vanishes at both ends, and on $[1, \infty)$ with $-e^{-t}$, so that the boundary term vanishes at $t = 1$ and at infinity. The two pieces are $-\int_0^1 (1 - e^{-t}) dt/t$ and $\int_1^\infty e^{-t} dt/t$, and their sum is $-\gamma_E$.

Second, for $y > 0$,

$$\text{Ein}(y) = \log y + \gamma_E + E_1(y), \quad 0 < E_1(y) \leq \frac{e^{-y}}{y}. \quad (20)$$

Split $\text{Ein}(y) = \int_0^1 (1 - e^{-t}) dt/t + \int_1^y dt/t - \int_1^y e^{-t} dt/t$ and complete the last integral to $\int_1^\infty$; by Eq. (19) the constants combine into γ_E , and what is left is $\log y + E_1(y)$. The bound follows from $1/t \leq 1/y$ on $[y, \infty)$. Both sides of the identity are analytic in the plane cut along the negative axis, so it continues to $|\arg y| < \pi$ with principal branches. For $\text{Re } y > 0$ the integral along the horizontal ray from y gives $|E_1(y)| \leq e^{-\text{Re } y} \int_0^\infty e^{-u} du/|y + u| \leq e^{-\text{Re } y}/|y|$, since $|y + u| \geq |y|$ for $u \geq 0$.

Third, the function $g(y) = \text{Ein}(y) - \log(1 + y)$ is nondecreasing on $[0, \infty)$, with $g(0) = 0$ and $g(\infty) = \gamma_E$. Indeed

$$g'(y) = \frac{1 - e^{-y}}{y} - \frac{1}{1 + y} = \frac{1 - (1 + y)e^{-y}}{y(1 + y)} \geq 0,$$

because $e^y \geq 1 + y$, and by Eq. (20) $g(y) = \gamma_E + E_1(y) - \log(1 + y^{-1}) \rightarrow \gamma_E$.

These three facts give the properties of the skeleton. They are collected in Table III and plotted in Figs. 4

and 5; we now derive them one by one. Section II G carries each of them over to a realization.

Two-sided bound. Putting $y = x^2$ in the third fact, for every real z ,

$$(1 + x^2)^{n/2} \leq \hat{a}(z) \leq \hat{c}(1 + x^2)^{n/2}, \quad \hat{c} = e^{n\gamma_E/2}. \quad (21)$$

The skeleton is pinched between two polynomials of degree $2n$ in x on the whole real axis, although it has no zero anywhere.

Large momenta on the real axis. For $|x| \geq 1$, Eq. (20) gives $\hat{a}(z) = \hat{c}|x|^n \exp[\frac{n}{2} E_1(x^2)]$, so $\hat{a} - \hat{c}|x|^n$ is smaller than $\hat{c}|x|^n$ by a factor of order e^{-x^2}/x^2 .

Small momenta. From the series of Ein,

$$\hat{a}(z) = 1 + \frac{n}{2} x^2 + \frac{n(n-1)}{8} x^4 + O(x^6), \quad (22)$$

with no term linear in x , in line with $\psi'_*(0) = 0$.

Derivatives. Write $\hat{a} = e^H$ with $H(z) = \frac{n}{2} \text{Ein}(z^2/M^4)$. Then

$$H'(z) = \frac{n(1 - e^{-x^2})}{z},$$

which is entire. For $|x| \geq 1$ it equals n/z minus a term whose derivatives of every order are bounded by $C_j e^{-x^2/2} M^{-2-2j}$, so $|H^{(j)}(z)| \leq C_j M^{-2j} (1 + |x|)^{-j}$ for $j \geq 1$ on the real axis. By Faà di Bruno's formula $\hat{a}^{(j)}$ is $\hat{a}$ times a polynomial in $H', \dots, H^{(j)}$ whose monomials all have total order j , so $|\hat{a}^{(j)}| \leq C_j M^{-2j} (1 + |x|)^{-j} \hat{a}$, and with Eq. (21)

$$|\hat{a}^{(j)}(z)| \leq C_j M^{-2j} (1 + |x|)^{n-j}, \quad z \in \mathbb{R}. \quad (23)$$

The nonlocal remainder. The function

$$\hat{\mathcal{G}}(z) = \frac{\hat{a}(z) - 1}{z} - \frac{\hat{c} z^{n-1}}{M^{2n}} \quad (24)$$

is entire, because $\hat{a} - 1$ vanishes at $z = 0$. It measures how far the nonlocal part of the action would be from a polynomial if the skeleton were the form factor. On $[0, M^2]$ it is a smooth function on a compact interval, and for $z \geq M^2$,

$$\hat{\mathcal{G}}(z) = -\frac{1}{z} + \frac{\hat{c} x^n}{z} \left(e^{\frac{n}{2} E_1(x^2)} - 1 \right).$$

Since E_1 and its derivatives at $y = x^2$ are bounded by polynomials in $1/x$ times e^{-x^2} , the second term and all its derivatives are exponentially small, and the derivatives of $-1/z$ give

$$|\hat{\mathcal{G}}^{(j)}(z)| \leq C_j M^{-2-2j} (1 + x)^{-1-j}, \quad z \geq 0. \quad (25)$$

In the language of pseudodifferential operators $\hat{\mathcal{G}}$ is a symbol of order -1 . In the same way $\hat{a} - \hat{c} x^n = \hat{c} x^n (e^{\frac{n}{2} E_1(x^2)} - 1)$ for $x \geq 1$, so $|\frac{d^j}{dz^j} (\hat{a} - \hat{c} x^n)| \leq$




FIG. 3. Two gases with the same strength per e -fold. (a) One realization of each, for $n = 4$ and sizes between $0.1 M^{-2}$ and M^{-2} : identical branes with $\vartheta_0 = \log 2$ (orange) and the fine-grained power law (8) with $\zeta = 2$ (blue). Each brane is a stem at its size, at a random transverse position, with height equal to its strength. At the smallest sizes the fine-grained gas has 100 times more branes per e -fold, each 100 times weaker. The dashed curve is the strength $\vartheta_0 \tau^2$ of the power law. The value $\zeta = 2$ keeps the number of stems small enough to draw; condition (G) needs $\zeta > 8$ for $n = 4$. (b) The exponent of the single-brane factor in units of the strength of the brane, $\psi_*(sz) = \vartheta(s)^{-1} \log t^{-1}$, over the plane of $\log s$ and $\log z$. It switches on across the line $sz = 1$ (dashed), so at fixed z the exponent L_Π of Eq. (12) collects the strength of the branes larger than $1/z$, which is the origin of the power law (17). (c) The exponent $L_\Pi(x)$ along the Euclidean axis for eight realizations of each gas, identical branes and the power law with $\zeta = 9$, drawn as a waterfall against the skeleton $n\Phi(x) = \frac{n}{2} \text{Ein}(x^2)$ (thick, in front, and dotted on each row). The fine-grained realizations run parallel to the skeleton at large x , offset by R_∞ ; the realizations of identical branes drift away from it like random walks in $\log x$ (Sec. II G). For $\zeta = 9$ the branes smaller than $0.3 M^{-2}$ are represented by the Gaussian limit of their compensated strength, as in Fig. 7.

$C_j M^{-2j} (1+x)^{-j}$ on $[0, \infty)$: a symbol of order 0. Figure 4(c) shows $z \hat{G}(z)$, which tends to -1 .

Complex momenta. Let $S_\delta = \{z : |\arg z| \leq \pi/4 - \delta\}$ with $0 < \delta < \pi/4$. For $z \in S_\delta$ we have $\text{Re } z^2 \geq |z|^2 \sin 2\delta > 0$, and the continued identity (20) with its bound gives

$$\begin{aligned} \hat{a}(z) &= \hat{c} \left(\frac{z}{M^2} \right)^n \left[1 + \epsilon(z) \right], \\ |\epsilon(z)| &\leq \exp \left(\frac{n M^4 e^{-|z|^2 \sin 2\delta / M^4}}{2|z|^2} \right) - 1. \end{aligned} \quad (26)$$

Because $\hat{a}(-z) = \hat{a}(z)$, the same holds in the reflected cone around the negative axis. Inside the two cones $|\arg(\pm z)| < \pi/4$, which contain the Euclidean and the Lorentzian axes, the skeleton is a monomial up to corrections that fall like a Gaussian. Outside them it is not polynomially bounded. On the imaginary axis $\hat{a}(iy) = \exp[-\frac{n}{2} \int_0^{y^2/M^4} (e^t - 1) dt/t]$ falls faster than any exponential of y^2 , and in between $|\hat{a}|$ oscillates with growing amplitude (Fig. 5). This rules out a naive Wick rotation, and Sec. VII explains how the theory is defined instead.

F. Other passive responses

The closed forms of Sec. II E rest on the choice $\psi = \psi_*$, but the field theory of Secs. III–VII does not. We call a response *admissible* if its exponent satisfies (S1)–(S4), and show that every admissible response gives a skeleton with the properties used there. Only two numbers change from one response to another, the constant C_+ of Eq. (16) and the moment $\mu_\psi = \int_0^\infty t(1 - \psi(t)) dt$.

Analyticity and the real axis. Section II D already shows that $\hat{a} = e^{n\Phi}$ is entire, has no zeros and satisfies $\hat{a}(0) = 1$; its Taylor coefficients are real. By (S2), $\Phi(x) = \int_0^1 \psi(x\tau) d\tau/\tau \geq 0$ for real x of either sign, so $\hat{a} \geq 1$ on the whole real axis. The function $\Phi(x) - \frac{1}{2} \log(1+x^2)$ is continuous on $\mathbb{R}$ and tends to $C_\pm$ as $x \rightarrow \pm\infty$, by Eq. (16), so it is bounded, between m_ψ and M_ψ say, and

$$e^{nm_\psi} (1+x^2)^{n/2} \leq \hat{a}(z) \leq e^{nM_\psi} (1+x^2)^{n/2}, \quad z \in \mathbb{R}. \quad (27)$$

For ψ_* this is Eq. (21), with $m_\psi = 0$ and $M_\psi = \gamma_E/2$.

Symbols. With $H = n\Phi(z/M^2)$ we have $H'(z) = n\psi(x)/z$, and for $|x| \geq 1$ this is n/z minus $n(1 - \psi(x))/z$, whose derivatives decay faster than any power by (S3). The argument of Sec. II E then gives Eq. (23) with the constant e^{nM_ψ} . Put $\hat{c} = e^{nC_+}$. For $x \geq 1$, $\hat{a} - \hat{c}x^n =$




FIG. 4. The skeleton on the real axis. (a) $\log_{10} \hat{a}(z)$ for $n = 4, \dots, 8$ (solid) inside the band of the two-sided bound (21), $(1+x^2)^{n/2} \leq \hat{a} \leq \hat{c}(1+x^2)^{n/2}$ (shaded), with the monomial $\hat{c}x^n$ that $\hat{a}$ approaches at large x (dashed). (b) $\Phi(x) - \frac{1}{2}\log(1+x^2)$ for the three admissible responses of Table I: it is bounded for each of them, as Eq. (27) states, and tends to the constant C_+ (dashed) of Eq. (28). (c) $v = z\hat{\mathcal{G}}(z) = \hat{a} - 1 - \hat{c}x^n$ for $n = 4, \dots, 8$, plotted as $\text{sgn}(v)\log_{10}(1+|v|)$: it is bounded and tends to -1 (dotted), so the nonlocal remainder (24) is a symbol of order -1 for every n .

$\hat{c}x^n(e^{n\rho_+(x)} - 1)$, and ρ_+ and all its derivatives decay faster than any power, so $\hat{a} - \hat{c}x^n$ is a symbol of order 0 and $\hat{\mathcal{G}}$ is a symbol of order -1 : Eq. (25) holds as it stands.

Cones. Let $z = re^{i\alpha}$ with $|\alpha| \leq \theta - \delta$. The function $g(w) = \Phi(w) - \log w$ is analytic in the sector and $g'(w) = -(1 - \psi(w))/w$, which by (S4) is bounded by $C_N|w|^{-N-1}$ for every N . The integral of g' along the ray from z to infinity therefore converges and is $O(r^{-N})$, so $g(z)$ tends to a limit $L(\alpha)$ along each ray, with an error $O(r^{-N})$. The limit does not depend on α , because the integral of g' over an arc of radius t between two rays is at most $\pi t \cdot C_N t^{-N-1} \rightarrow 0$. On the positive axis $L(0) = C_+$ by Eq. (16). Hence, in the sector,

$$\hat{a}(z) = \hat{c}\left(\frac{z}{M^2}\right)^n e^{n\rho(z)}, \quad |\rho(z)| \leq C_{N,\delta} \left|\frac{M^2}{z}\right|^N, \quad (28)$$

for every N , with $\hat{c} = e^{nC_+}$, and the same holds around the negative axis with C_- and $(-z/M^2)^n$. For ψ_* this is Eq. (26), where the remainder is even Gaussian.

The zero-point moment. Section VIB needs the integral of $\Phi(k^2/M^2)$ over four-momentum in dimensional regularization. Its Mellin transform is elementary. For $-2 < \text{Re } s < 0$, exchanging the order of integration in Eq. (14) gives $\int_0^\infty w^{s-1} \Phi(w) dw = -\tilde{\psi}(s)/s$, with $\tilde{\psi}(s) = \int_0^\infty u^{s-1} \psi(u) du$; the strip is where both integrals converge, since $\psi(u) = O(u^2)$ at small u by passivity and $\psi \rightarrow 1$ at large u . Splitting $\tilde{\psi}$ at $u = 1$ and writing $\psi = 1 - (1 - \psi)$ on $[1, \infty)$ continues it to $\text{Re } s > -2$, $s \neq 0$, and for $\text{Re } s > 0$ the continuation is

$$\tilde{\psi}(s) = -\int_0^\infty u^{s-1} (1 - \psi(u)) du. \quad \text{At } s = D/2 = 2,$$

$$\int \frac{d^4 k}{(2\pi)^4} \Phi\left(\frac{k^2}{M^2}\right) = \frac{M^4}{32\pi^2} \mu_\psi, \quad (29)$$

$$\mu_\psi = \int_0^\infty t (1 - \psi(t)) dt,$$

which is positive and finite by (S2) and (S3).¹¹ For a single brane of size τ/M^2 no continuation is needed, because the integrand falls off rapidly by (S3) and the integral converges absolutely:

$$\int \frac{d^4 k}{(2\pi)^4} \left[\psi\left(\frac{\tau k^2}{M^2}\right) - 1 \right] = -\frac{M^4}{16\pi^2 \tau^2} \mu_\psi, \quad (30)$$

as follows from the substitution $u = \tau k^2/M^2$, after which the angular integration leaves $d^4 k/(2\pi)^4 \rightarrow u du M^4/(16\pi^2 \tau^2)$.

Small momenta. Passivity gives $\psi(w) = \psi_2 w^2 + O(w^3)$ with $\psi_2 \geq 0$, so $\hat{a} = 1 + \frac{n}{2} \psi_2 x^2 + O(x^3)$. The leading correction to the kinetic operator is of fourth order in momentum, as in Eq. (22).

Table I gives three admissible responses and two that fail, and Fig. 6 shows the sectors of (S4).

¹¹ For ψ_* , $\Phi(w) = \frac{1}{2} \text{Ein}(w^2)$ and $\mu_{\psi_*} = \int_0^\infty t e^{-t^2} dt = \frac{1}{2}$, so the right side is $M^4/(64\pi^2)$. This agrees with Sec. VIB, where the same integral is done in the variable k^4/M^4 with the Mellin transform (76).




FIG. 5. The skeleton in the complex z plane. (a) $v = \frac{2}{n} \log |\hat{a}| = \text{Re Ein}(z^2/M^4)$ over the disk $|z| \leq 2.2M^2$, plotted as $\text{sgn}(v) \log(1 + |v|)$ to compress its range. Along the real axis (white) it rises like $2 \log |z|$; along the imaginary axis (orange) it falls without bound; the dashed red lines are the cone boundaries $\arg z = \pm\pi/4, \pm 3\pi/4$, between which $|\hat{a}|$ oscillates. (b) Accuracy of the local approximation $\hat{c}(z/M^2)^n$ for $n=6$, $\log_{10} |\hat{a}/(\hat{c}z^n/M^{2n}) - 1|$ clipped at 0: the error is exponentially small inside the cones $|\arg(\pm z)| < \pi/4$ (blue), Eq. (26), and of order one outside. The floor repeats the map as a contour plot, with the cone boundaries (dashed) and, as black dots, the zeros that a frozen gas of the resonant factors $2 - e^{-s^2 z^2}$ of Eq. (4) would have for sizes in $[0.15, 1] M^{-2}$. They all lie outside the cones. A gas of branes obeying (S5) has none.

TABLE I. Single-brane responses, given by the exponent ψ of Eq. (3). The first three are admissible; for them $\hat{c} = e^{nC_+}$ is the leading coefficient of the skeleton, $\hat{a} \simeq \hat{c}x^n$, μ_ψ is the moment of Eq. (29), which fixes the mean zero-point energy (78), and θ is the opening of the sectors of (S4). The last two violate the conditions named in the third column. Values of $\hat{c}$ are for $n=4$; for general n , $\hat{c} = e^{nC_+}$.

exponent $\psi(w)$	small w	C_+	$\hat{c}, n=4$	μ_ψ	$\rho_1/(nM^4)$	θ	skeleton exponent $\Phi(x)$
$1 - e^{-w^2}$	w^2	$\gamma_E/2 = 0.2886$	3.172	$\frac{1}{2}$	$3/(64\pi^2)$	$\pi/4$	$\frac{1}{2} \text{Ein}(x^2)$
$1 - (1 + w^2)e^{-w^2}$	$\frac{1}{2}w^4$	$(\gamma_E - 1)/2 = -0.2114$	0.4293	1	$3/(32\pi^2)$	$\pi/4$	$\frac{1}{2}[\text{Ein}(x^2) - 1 + e^{-x^2}]$
$1 - e^{-w^4}$	w^4	$\gamma_E/4 = 0.1443$	1.781	$\frac{\sqrt{\pi}}{4}$	$3\sqrt{\pi}/(128\pi^2)$	$\pi/8$	$\frac{1}{4} \text{Ein}(x^4)$
$1 - e^{-w}$	w	fails (S2): unbounded below on the negative axis; the factor decays faster than any power on the timelike axis					
$w^2/(1 + w^2)$	w^2	fails (S1) and (S3): poles at $w = \pm i$, and $1 - \psi \sim w^{-2}$, so G and Λ run at one loop					

G. Fluctuations of a realization

The skeleton has every property that the field theory needs. We now show that, under (G), almost every realization has them too, with a random constant in place of $\hat{c}$. The two differ by the factor e^{R_Π} of Eq. (13), so everything rests on how R_Π behaves at large $|z|$.

The limit at large momentum. At large real x every brane larger than $1/x$ is saturated and contributes its full strength. This suggests comparing R_Π with the total

strength of the gas measured from its mean,

$$R_\infty = \int \vartheta(s) (\Pi - \nu)(ds) = \lim_{\eta \rightarrow 0} \left[\sum_{s \in \Pi, s > \eta/M^2} \vartheta(s) - n \log \frac{1}{\eta} \right]. \quad (31)$$

The sum in brackets has mean $\int_{\eta/M^2}^{1/M^2} \vartheta d\nu = n \log(1/\eta)$ by Eq. (6), and by Eq. (10) its fluctuation has variance $n[V(1) - V(\eta)]$. Under (G) $V(1)$ is finite, and Sec. IIC




FIG. 6. The saturation sectors of condition (S4) for the three admissible exponents of Table I. Each surface is $\log_{10} |1 - \psi(w)|$ over the complex w plane, clipped at -8 below and $+3$ above, with the floor showing the same quantity as a contour map. Where the surface is low the brane is saturated, $\psi \simeq 1$; the dashed lines are the sector boundaries, $\arg w = \pm\pi/4$ in (a) and (b) and $\arg w = \pm\pi/8$ in (c). The exponent $1 - e^{-w^4}$ of panel (c) is also saturated around the imaginary axis, which (S4) does not require.

gives the limit almost surely, with

$$\mathbb{E}R_\infty = 0, \quad \mathbb{E}R_\infty^2 = nV(1). \quad (32)$$

The difference

$$R_\Pi(z) - R_\infty = \int \vartheta(s) [\psi(sz) - 1] (\Pi - \nu)(ds) \quad (33)$$

is carried by the branes that are not saturated at z .

Mean-square decay. By Eq. (10), for complex z and $w = z/M^2$,

$$\mathbb{E}|R_\Pi(z) - R_\infty|^2 = n \int_0^1 \vartheta(\tau) |1 - \psi(w\tau)|^2 \frac{d\tau}{\tau}. \quad (34)$$

Let $|w| \geq 1$ and $|\arg(\pm w)| \leq \theta - \delta$. For $\tau \leq 1/|w|$ the point $w\tau$ lies in the closed unit disk, where $|1 - \psi| \leq B$, and this part of the integral is at most $nB^2V(1/|w|) \leq nB^2C|w|^{-\sigma}$ by (G). For $\tau > 1/|w|$ condition (S4) gives $|1 - \psi(w\tau)| \leq C_N|w\tau|^{-N}$. Choose $2N > \sigma$ and integrate by parts in $\int_{1/|w|}^1 \tau^{-2N} dV(\tau)$: the boundary term at $\tau = 1/|w|$ is negative, the one at $\tau = 1$ is $V(1)$, and the remaining integral is $2N \int \tau^{-2N-1} V(\tau) d\tau \leq 2NC|w|^{2N-\sigma}/(2N-\sigma)$. Multiplying by $C_N^2|w|^{-2N}$, this part is also $O(|w|^{-\sigma})$, and

$$\begin{aligned} \mathbb{E}|R_\Pi(z) - R_\infty|^2 &\leq K_\delta \left| \frac{M^2}{z} \right|^\sigma, \\ |z| &\geq M^2, \quad |\arg(\pm z)| \leq \theta - \delta. \end{aligned} \quad (35)$$

Almost-sure decay. Equation (35) controls one point at a time. To control a whole sector we use the subharmonicity of $|f|^2$ for analytic f : $|f(w)|^2$ is at most

the mean of $|f|^2$ over any disk centered at w on which f is analytic. Put $f(w) = R_\Pi(M^2w) - R_\infty$, an entire function, fix δ , and let A_k be the part of the sector $|\arg w| \leq \theta - \delta$ with $2^k \leq |w| \leq 2^{k+1}$, $k \geq 1$. A disk of radius $r_k = 2^{k-1} \sin(\delta/2)$ around a point of A_k lies in the region A'_k where $|\arg w| \leq \theta - \delta/2$ and $2^{k-1} \leq |w| \leq 2^{k+2}$. Hence

$$\sup_{A_k} |f|^2 \leq \frac{1}{\pi r_k^2} \int_{A'_k} |f|^2 dA,$$

and taking expectations with Eq. (35) in the wider sector,

$$\mathbb{E} \sup_{A_k} |f|^2 \leq \frac{\text{area}(A'_k)}{\pi r_k^2} K_{\delta/2} 2^{-(k-1)\sigma} \leq K' 2^{-k\sigma},$$

since the ratio of the area of A'_k to r_k^2 does not depend on k . Chebyshev's inequality gives $P(\sup_{A_k} |f| > 2^{-k(\sigma/2-\varepsilon)}) \leq K' 2^{-2k\varepsilon}$, which is summable in k for every $\varepsilon > 0$. By the Borel–Cantelli lemma, almost surely $\sup_{A_k} |f| \leq 2^{-k(\sigma/2-\varepsilon)}$ for all but finitely many k , and on the remaining annuli f is bounded because it is continuous. Taking δ and ε rational, and treating the sectors around the negative axis in the same way, we conclude that almost surely

$$\begin{aligned} |R_\Pi(z) - R_\infty| &\leq C_\Pi \left| \frac{M^2}{z} \right|^{\sigma/2-\varepsilon}, \\ |z| &\geq M^2, \quad |\arg(\pm z)| \leq \theta - \delta, \end{aligned} \quad (36)$$

for every $\delta \in (0, \theta)$ and every $\varepsilon > 0$, with C_Π finite and depending on the realization, on δ and on ε . For the power law below $\sigma = \zeta$, and the exponent matches, up

to ε , the decay of the standard deviation of $R_\Pi(z) - R_\infty$, which falls like $|z|^{-\zeta/2}$, Eq. (43).

The form factor of a realization. With Eqs. (13) and (28), Eq. (36) says that almost surely, in the same sectors,

$$a_\Pi(z) = c_\Pi \left(\frac{\pm z}{M^2} \right)^n e^{\eta_\Pi(z)}, \quad c_\Pi = e^{nC_\pm + R_\infty}, \quad (37)$$

$$|\eta_\Pi(z)| \leq C'_\Pi \left| \frac{M^2}{z} \right|^{\sigma/2 - \varepsilon}.$$

A realization has the asymptotics of the skeleton, with a random coefficient. Cauchy's estimate on a disk of radius $\frac{1}{2}|z|\sin(\delta/2)$ around z , which stays inside the wider sector, bounds the derivatives: $|\eta_\Pi^{(j)}(z)| \leq C_j |z|^{-j} M^2/z|^{\sigma/2 - \varepsilon}$. On the real axis R_Π is continuous and tends to R_∞ at both ends, so $\|R_\Pi\| = \sup_{\mathbb{R}} |R_\Pi|$ is finite, and Eq. (27) gives

$$e^{nm_\psi - \|R_\Pi\|} (1+x^2)^{n/2} \leq a_\Pi(z) \leq e^{nM_\psi + \|R_\Pi\|} (1+x^2)^{n/2} \quad (38)$$

for every real z .

Now let $\sigma > 2n$, as (G) requires, and take $\varepsilon < \sigma/2 - n$. Then for $|x| \geq 1$

$$a_\Pi(z) - c_\Pi |x|^n = c_\Pi |x|^n (e^{\eta_\Pi} - 1) = O(|x|^{n-\sigma/2+\varepsilon}),$$

and each derivative falls by one more power of x ; so $a_\Pi - c_\Pi x^n$ is a symbol of negative order, in particular of order 0, and a_Π obeys the derivative bounds (23) of the skeleton. The nonlocal remainder of the realization,

$$\mathcal{G}(z) = \frac{a_\Pi(z) - 1}{z} - \frac{c_\Pi z^{n-1}}{M^{2n}}, \quad (39)$$

is entire, equals $-1/z + c_\Pi x^n (e^{\eta_\Pi} - 1)/z$ for $x \geq 1$, and obeys

$$|\mathcal{G}^{(j)}(z)| \leq C_j M^{-2-2j} (1+x)^{-1-j}, \quad z \geq 0, \quad (40)$$

almost surely; the same bounds hold with $|z|$ in place of z in every closed subsector of the sectors of (S4). This is where the threshold in (G) comes from. The fluctuation of the gas at momentum p must fall faster than p^{-2n} , so that what remains of a_Π after the monomial $c_\Pi x^n$ is subtracted stays bounded. For the skeleton the same remainder falls faster than any power, and the field theory of Secs. IV–VII uses only that it is a symbol of order -1 .

Identical branes. Without (G) the conclusion fails, and it fails for the most obvious gas. Let all branes have the same strength ϑ_0 , so that $\nu = (n/\vartheta_0) ds/s$. For real x the sum $R_\Pi(x)$ is an ordinary compensated sum, and Eq. (9) with $f = e^{iu\vartheta_0\psi} - 1$ gives its characteristic function,

$$\left| \mathbb{E} e^{iuR_\Pi(x)} \right| = \exp \left[\frac{n}{\vartheta_0} \int_0^x (\cos(u\vartheta_0\psi(t)) - 1) \frac{dt}{t} \right].$$

As $x \rightarrow \infty$ the integrand tends to $\cos(u\vartheta_0) - 1$, which is negative unless $u\vartheta_0 \in 2\pi\mathbb{Z}$, and $\int dt/t$ diverges, so the right side tends to 0 for all such u . If $R_\Pi(x)$ converged in distribution, its characteristic function would converge

to a function continuous at $u = 0$ [27]; the limit here is 1 at $u = 0$ and 0 around it. So $R_\Pi(x)$ has no limit in distribution, and a fortiori no limit almost surely: $a_\Pi(x)/x^n$ does not converge, and no constant c makes $a_\Pi - cx^n$ a symbol of order 0. For $c > 0$ this would force $R_\Pi(x)$ to converge. For $c \leq 0$ it would force $a_\Pi(x)$ to stay bounded, hence $R_\Pi(x) \leq C - n \log x$ for large x , so that only a bounded number of branes could have sx above a fixed t_0 beyond which $\psi \geq \frac{1}{2}$; their number is Poisson with mean $(n/\vartheta_0) \log(x/t_0)$, and the probability of that event tends to zero. The variance shows the mechanism,

$$\mathbb{E} R_\Pi(x)^2 = n\vartheta_0 \Psi_*(x),$$

$$\Psi_*(x) = \int_0^1 \psi_*(x\tau)^2 \frac{d\tau}{\tau} = \text{Ein}(x^2) - \frac{1}{2} \text{Ein}(2x^2)$$

$$= \log|x| + \frac{1}{2}(\gamma_E - \log 2) + O(e^{-x^2}), \quad (41)$$

from $(1-e^{-u})^2 = 2(1-e^{-u}) - (1-e^{-2u})$, the substitution $u = x^2\tau^2$ and Eq. (20). The saturated strength performs a random walk in $\log x$, one step of size ϑ_0 for each of the roughly $(n/\vartheta_0) \log x$ branes larger than $1/x$. Such a gas still gives an entire form factor without zeros, by (S5), and $\log a_\Pi(x)/\log x \rightarrow n$ almost surely by the strong law of large numbers. But the local split of Sec. VA, on which the one-loop analysis rests, does not exist for it.

Statistics of a realization. For the power law (8) and ψ_* the moments on the real axis are incomplete gamma functions. With $\gamma(b, y) = \int_0^y t^{b-1} e^{-t} dt$ and $(1-e^{-u})^2 = 1 - 2e^{-u} + e^{-2u}$,

$$\mathbb{E} R_\Pi(x)^2 = n\vartheta_0 \left[\frac{1}{\zeta} - \frac{\gamma(\frac{\zeta}{2}, x^2)}{x^\zeta} + \frac{\gamma(\frac{\zeta}{2}, 2x^2)}{2(2x^2)^{\zeta/2}} \right], \quad (42)$$

$$\mathbb{E} |R_\Pi(x) - R_\infty|^2 = \frac{n\vartheta_0}{2} \frac{\gamma(\frac{\zeta}{2}, 2x^2)}{(2x^2)^{\zeta/2}} \leq \frac{n\vartheta_0 \Gamma(\frac{\zeta}{2})}{2(2x^2)^{\zeta/2}}, \quad (43)$$

and $\mathbb{E} R_\infty^2 = n\vartheta_0/\zeta$. The leading coefficient $c_\Pi = \hat{c} e^{R_\infty}$ has $\mathbb{E} \log c_\Pi = \log \hat{c}$, and by Eq. (11), with the substitution $t = \vartheta_0 \tau \zeta$,

$$\mathbb{E} c_\Pi = \hat{c} \exp \left[\frac{n}{\zeta} \int_0^{\vartheta_0} \frac{e^t - 1 - t}{t^2} dt \right]. \quad (44)$$

The image of ν under $s \mapsto \vartheta(s)$ is $(n/\zeta) dt/t^2$ on $(0, \vartheta_0]$, so R_∞ is a centered, totally skewed variable whose Lévy measure is that of a stable law of index one, cut off at ϑ_0 . Its jumps are positive, and since $e^{-\lambda t} - 1 + \lambda t \leq \frac{1}{2} \lambda^2 t^2$ for $t \geq 0$, Eq. (11) gives $\mathbb{E} e^{-\lambda R_\infty} \leq e^{\lambda^2 n V(1)/2}$ and hence, for any gas with bounded strengths that obeys (G),

$$\mathbb{P}(c_\Pi < \hat{c} e^{-y}) \leq \exp \left[-\frac{y^2}{2nV(1)} \right]. \quad (45)$$

The coefficient of a realization is bounded away from zero except with Gaussian probability. Table II gives the numbers for $\vartheta_0 = \log 2$, a brane at the top of the range doubling the kinetic operator when saturated, and

TABLE II. Fluctuations of a realization for the power law (8) with $\zeta = 2n + 1$, $\vartheta_0 = \log 2$ and $\psi = \psi_*$. Columns: standard deviation of $\log(c_\Pi/\hat{c}) = R_\infty$; mean of $c_\Pi/\hat{c}$, Eq. (44); lower bound on $c_\Pi/\hat{c}$ that holds with probability at least 0.95, from Eq. (45); the value x_* beyond which the root-mean-square remainder (43) is below 10^{-3} ; and the relative standard deviation of the zero-point energy, Eq. (79). All entries are closed-form expressions evaluated numerically.

n	ζ	$(n\vartheta_0/\zeta)^{1/2}$	$\mathbb{E} e^{R_\infty}$	95% bound	x_*	$\delta\rho/\rho_1$
4	9	0.555	1.190	0.257	4.47	0.372
5	11	0.561	1.195	0.253	3.74	0.281
6	13	0.566	1.198	0.250	3.35	0.227
7	15	0.569	1.201	0.249	3.11	0.190
8	17	0.571	1.202	0.247	2.97	0.163

$\zeta = 2n + 1$, the smallest integer allowed by (G). Figure 7 shows sample paths of R_Π for the two kinds of gas, the root-mean-square remainder (43) as a function of x and ζ , and the remainder of one realization over the complex plane.

H. What the field theory uses

From here on the form factor is that of one realization, and we drop the subscript: $a = a_\Pi$ and $c = c_\Pi$, with the remainder $\mathcal{G}$ of Eq. (39). Sections III–VII use a only through the following properties, each proved above for almost every realization of a gas obeying A1 and A2: a is entire, has no zeros, satisfies $a(0) = 1$ and has real Taylor coefficients (Sec. IID); on the real axis it obeys the two-sided bound (38) and the derivative bounds (23); $\mathcal{G}$ is a symbol of order -1 and $a - cx^n$ a symbol of order 0, Eq. (40); and in the cones of (S4) the asymptotics (37) hold, with $|a(z)| \geq \frac{c}{2}|z/M^2|^n$ for large $|z|$. Where a statement depends on the realization, the dependence is through c , through the bound C_Π of Eq. (36), or through $\|R_\Pi\|$; we point it out where it matters. The skeleton enters directly in two places only, the zero-point energy of Sec. VIB, whose mean it gives, and the Newtonian numbers of Sec. VIII. Table III collects the properties of the skeleton and of a realization side by side.

The two premises are the only places where the brane picture enters. Assumption A1 is a statement about a single brane, (S1)–(S5), which we do not derive from a brane action. Assumption A2 is a statement about the gas: independence, the scale-invariant strength (6) and the fine graining (G). No average over realizations is taken anywhere, and no limit of many regions or many samples is needed: the field theory below is that of one frozen gas. Table IV records where each input is used.

III. ACTION, PROPAGATOR AND SPECTRUM

A. The action

Let $g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu}$ with $\kappa^2 = 32\pi G$, and $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$. The Euclidean action is

$$S = -\frac{2}{\kappa^2} \int d^4x \sqrt{g} [R + G_{\mu\nu} F(\square) R^{\mu\nu}] + S_q, \quad F(\square) = \frac{a(-\square) - 1}{\square}, \quad (46)$$

where S_q collects three operators quartic in curvature, introduced in Sec. VD. Being quartic, they contribute nothing to the quadratic or cubic terms of the action around flat space. Since $a(0) = 1$, F is an entire function of $\square$. The combination $G_{\mu\nu} F(\square) R^{\mu\nu}$ is the one used in Refs. [8, 11], and the reason for it appears below: around flat space it multiplies the whole Einstein operator by one function.

B. Linearized curvature

We use the spin projectors of Barnes and Rivers [29, 30]. With $\theta_{\mu\nu} = \delta_{\mu\nu} - p_\mu p_\nu / p^2$ and $\omega_{\mu\nu} = p_\mu p_\nu / p^2$, in D dimensions,

$$P_{\mu\nu\alpha\beta}^{(2)} = \frac{1}{2}(\theta_{\mu\alpha}\theta_{\nu\beta} + \theta_{\mu\beta}\theta_{\nu\alpha}) - \frac{1}{D-1}\theta_{\mu\nu}\theta_{\alpha\beta}, \\ (P_s^{(0)})_{\mu\nu\alpha\beta} = \frac{1}{D-1}\theta_{\mu\nu}\theta_{\alpha\beta}.$$

They are orthogonal idempotents, and both are annihilated by contraction with p^μ . For a metric perturbation $h_{\mu\nu}$ with Fourier transform $h(p)$, the linearized Ricci tensor is

$$R_{\mu\nu}^{(1)} = \frac{1}{2}(-p_\rho p_\mu h^\rho{}_\nu - p_\rho p_\nu h^\rho{}_\mu + p^2 h_{\mu\nu} + p_\mu p_\nu h),$$

with $h = h^\mu{}_\mu$. Substituting a gauge mode $h_{\mu\nu} = p_\mu \xi_\nu + p_\nu \xi_\mu$, the terms cancel in pairs, so $R^{(1)}$ depends only on the transverse part $(P^{(2)} + P_s^{(0)})h$; every symmetric tensor is its transverse part plus a gauge mode, because $1 - P^{(2)} - P_s^{(0)}$ projects onto tensors of the form $p_\mu \xi_\nu + p_\nu \xi_\mu$. For transverse h the formula reduces to $\frac{1}{2}(p^2 h_{\mu\nu} + p_\mu p_\nu h)$, and writing $h_{\mu\nu} = (P^{(2)}h)_{\mu\nu} + (\theta \cdot h)\theta_{\mu\nu}/(D-1)$, with $\theta \cdot h = \theta^{\mu\nu} h_{\mu\nu}$, we get

$$R_{\mu\nu}^{(1)} = \frac{p^2}{2} [(P^{(2)}h)_{\mu\nu} + (\theta \cdot h) T_{\mu\nu}], \quad T_{\mu\nu} = \frac{\theta_{\mu\nu}}{D-1} + \omega_{\mu\nu}. \quad (47)$$

Its trace is $R^{(1)} = p^2 \theta \cdot h$. Squaring Eq. (47), the cross terms vanish because $P^{(2)}h$ is transverse and traceless, $T \cdot T = D/(D-1)$, and $(\theta \cdot h)^2 = (D-1)h P_s^{(0)}h$, where hPh stands for $h^{\mu\nu} P_{\mu\nu\alpha\beta} h^{\alpha\beta}$ and products are understood as




FIG. 7. Fluctuations of a realization, for $n = 4$, $\vartheta_0 = \log 2$ and ψ_* . (a) Sample paths of $R_{\Pi}(x)$ for twelve realizations of a gas of identical branes, drawn as ribbons over $\log_{10} x$ and the index of the realization, with the band $\pm(E R_{\Pi}^2)^{1/2}$ of Eq. (41) (translucent sheets with dashed edges). The band widens like $(\log x)^{1/2}$ and the paths wander without limit. (b) The same for the fine-grained gas with $\zeta = 9$, with the band of Eq. (42). Each path levels off at its own value of R_{∞} , marked by a dot at the right end. (c) The root-mean-square remainder $(E|R_{\Pi} - R_{\infty}|^2)^{1/2}$, Eq. (43), as a surface over $\log_{10} x$ and ζ , in $\log_{10}$. Its slope in $\log_{10} x$ is $-\zeta/2$; the red line marks $\zeta = 2n = 8$, above which the remainder of a_{Π} after the monomial is subtracted is bounded. (d) $\log_{10} |R_{\Pi}(z) - R_{\infty}|$ for one realization with $\zeta = 9$ over the plane of $\log_2 |z/M^2|$ and $\arg z$, for $|\arg z| \leq 72^\circ$, with the cone boundaries $|\arg z| = 45^\circ$ (dashed) and the real axis (white). Inside the cone the surface falls along every ray with slope close to $-\zeta/2$ (dotted reference line, shifted up), as Eq. (36) states; the dips on the real axis are sign changes of $R_{\Pi} - R_{\infty}$. Outside the cone, where (S4) gives no control, the surface rises and is cut off at 10. In (b) and (d) the branes smaller than $0.3 M^{-2}$ in the fine-grained gas, about 3×10^5 per e -fold and more, are represented by their Gaussian limit; all other branes are drawn individually.

$h(-p) \cdots h(p)$. This gives

$$R_{\mu\nu}^{(1)} R^{(1)\mu\nu} = \frac{v^4}{4} [h P^{(2)} h + D h P_s^{(0)} h], \quad (48)$$

$$(R^{(1)})^2 = p^4 (D - 1) h P_s^{(0)} h, \quad (49)$$

$$G_{\mu\nu}^{(1)} R^{(1)\mu\nu} = \frac{v^4}{4} [h P^{(2)} h - (D - 2) h P_s^{(0)} h], \quad (50)$$

$$h^{\mu\nu} G_{\mu\nu}^{(1)} = \frac{v^2}{2} [h P^{(2)} h - (D - 2) h P_s^{(0)} h]. \quad (51)$$

Equation (50) is Eq. (48) minus half of Eq. (49), and Eq. (51) follows from $G^{(1)} = \frac{v^2}{2} [P^{(2)} h - \frac{D-2}{D-1} (\theta \cdot h) \theta]$.

TABLE III. Properties of the skeleton $\hat{a} = \exp[\frac{n}{2} \text{Ein}(x^2)]$ for ψ_* , with $\hat{c} = e^{n\gamma_E/2}$, and of the form factor a of a realization, which hold almost surely for a gas obeying (G). Here $x = z/M^2$, $\varepsilon > 0$ is arbitrary and C depends on the realization. All entries are derived in Secs. II D–II G; the last column says where each is used.

property	skeleton $\hat{a}$	realization $a = \hat{a} e^{R_\Pi}$	used in
analyticity	entire, no zeros, $\hat{a}(0) = 1$, $\hat{a}(-z) = \hat{a}(z)$, real coefficients	entire, no zeros by (S5), $a(0) = 1$, $a(-z) = a(z)$, real coefficients	pole content, Sec. III C
real axis	$(1+x^2)^{n/2} \leq \hat{a} \leq \hat{c}(1+x^2)^{n/2}$	same, with factors $e^{\mp \ R_\Pi\ }$, Eq. (38)	propagator bound (56)
large $ x $, real	$\hat{a} = \hat{c} x ^n \exp[\frac{n}{2} E_1(x^2)]$	$a = c x ^n[1 + O(x ^{-\sigma/2+\varepsilon})]$, $c = \hat{c}e^{R_\infty}$	local theory, Sec. V A
small x	$\hat{a} = 1 + \frac{n}{2}x^2 + \frac{n(n-1)}{8}x^4 + O(x^6)$	$a = 1 + b_\Pi x^2 + O(x^4)$, $b_\Pi = \sum \vartheta(s)s^2 M^4$, $\mathbb{E} b_\Pi = \frac{n}{2}$	Newtonian limit, Sec. VIII
remainder	$\hat{\mathcal{G}}$ of order -1 ; $\hat{a} - \hat{c}x^n$ exponentially small	$\mathcal{G}$ of order -1 ; $a - cx^n$ of order $n - \sigma/2 + \varepsilon < 0$	vertex bound, Sec. IV C
cones	$\hat{a} = \hat{c}(z/M^2)^n[1 + \epsilon]$, $ \epsilon \lesssim e^{-\text{Re } z^2/M^4}$	$a = c(z/M^2)^n e^{\eta_\Pi}$, $ \eta_\Pi \leq C z/M^2 ^{-\sigma/2+\varepsilon}$	unitarity, Sec. VII
imaginary axis	$\hat{a}(iy) \rightarrow 0$ faster than any exponential	not controlled, not used	Sec. VII
constant	$\hat{c} = 3.172, 4.234, 5.650, 7.540, 10.063$ for $n = 4, \dots, 8$	$\log c$ has mean $\log \hat{c}$, variance $nV(1)$	one-loop tuning, Sec. V D
other responses	same with $\hat{c} = e^{nC_+}$, Sec. II F	same with $c = e^{nC_++R_\infty}$	Sec. II F

TABLE IV. Inputs of the paper, what each is used for, and what fails without it. Everything not listed here is derived, for almost every realization of the gas.

input	used for	if dropped
A1, (S1)	a entire with $a(0) = 1$	$F(\square)$ not entire
A1, (S2)	$a \geq 1$ on both half-axes	no polynomial bound on one half-axis
A1, (S3)	skeleton corrections beyond all powers	G and Λ run at one loop
A1, (S4)	cone asymptotics (28), (37)	cutting rules not established
A1, (S5)	no zeros in any realization	complex pole pairs, Eq. (4)
A2, independence	Poisson gas	formulas of Sec. II C lost
A2, Eq. (6)	growth $ x ^n$ with integer n	no fixed power
A2, (G)	$a - cx^n$ bounded, Eq. (40)	identical branes: no limit of a/x^n
$\psi = \psi_*$	closed forms; numbers of Sec. VIII	other constants, Table I
n integer	local theory polynomial	no local split
$n \geq 4$	every graph with $L \geq 2$ converges	$n = 3$: $\bar{\omega}(2) = 0$

C. Quadratic action and propagator

Since $\delta(\sqrt{g}R) = -\sqrt{g}G^{\mu\nu}\delta g_{\mu\nu}$ up to a total derivative and $G_{\mu\nu}$ vanishes on flat space, the second-order term of $\int \sqrt{g}R$ is $-\frac{1}{2} \int \delta g^{\mu\nu} G_{\mu\nu}^{(1)}[\delta g]$, which for $\delta g = \kappa h$ is $-\frac{\kappa^2}{2} \int h G^{(1)}[h]$. The nonlocal term is already quadratic in curvature, so its second-order part is

$\kappa^2 \int G^{(1)}F(\square)R^{(1)}$ with $\square \rightarrow -p^2$. With the prefactor $-2/\kappa^2$ and Eqs. (50) and (51) at $D = 4$, the quadratic action is $\frac{1}{2}p^2[1 - p^2F(-p^2)]h[P^{(2)} - 2P_s^{(0)}]h$, and $1 - p^2F(-p^2) = 1 + (a(p^2) - 1) = a(p^2)$. Hence

$$S^{(2)} = \frac{1}{2} \int \frac{d^4p}{(2\pi)^4} p^2 a(p^2) h(-p) [P^{(2)} - 2P_s^{(0)}] h(p), \quad (52)$$

the Einstein–Hilbert quadratic action multiplied by $a(p^2)$. On the gauge-invariant subspace it is inverted by

$$\mathcal{D}(p) = \frac{P^{(2)} - \frac{1}{2}P_s^{(0)}}{p^2 a(p^2)}, \quad (53)$$

the propagator of general relativity divided by a .

As a function of $z = p^2 \in \mathbb{C}$, Eq. (53) is analytic except at $z = 0$: a is entire and zero-free, so $1/a$ is entire, and $1/(za)$ has a simple pole at the origin with residue $1/a(0) = 1$; the projectors do not depend on $|p|$. The residue is that of general relativity, so the exchange between conserved sources at long distances is Einstein’s and carries the same positive norm. In Lorentzian signature the only on-shell condition is $p^2 = 0$, and because the form factor multiplies the spin-two projector by a scalar, both helicities have the dispersion relation $\omega^2 = k^2$ at every wavenumber. There are no massive states, ghosts or tachyons in the tree-level spectrum, and the speed of gravitational waves equals that of light identically.¹²

For comparison, a polynomial form factor of degree N has N extra poles whose residues add up to -1 , by

¹² The bound from GW170817 and GRB 170817A is $-3 \times 10^{-15} < (c_g - c)/c < 7 \times 10^{-16}$ [31]. Theories with a momentum-dependent spin-two form factor that is not shared by the whole Einstein operator generically violate it at some wavenumber; here the factorized structure (52) prevents that.

the residue theorem applied to $1/(za)$ [4]; Stelle's theory is $N = 1$, with residue exactly -1 [3]. A frozen gas of resonant branes would have infinitely many complex pairs, Eq. (4); a realization of the gas of Sec. II, whose branes have no resonances, has no extra pole at all.

IV. ULTRAVIOLET POWER COUNTING

A. Gauge fixing and ghosts

We use the background-field method [32], $g = \bar{g} + \kappa h$, with the background-covariant de Donder condition $\chi_\mu = \bar{\nabla}^\nu h_{\mu\nu} - \frac{1}{2} \bar{\nabla}_\mu h$ and the weighted gauge-fixing term

$$S_{\text{gf}} = \int d^4x \sqrt{\bar{g}} \chi_\mu a(-\bar{\square}) \chi^\mu. \quad (54)$$

For $a = 1$ this is the Feynman-de Donder term, which added to the Einstein part of Eq. (52) gives the operator $p^2(\mathbf{1} - \frac{1}{2}\delta \otimes \delta)$, with $\mathbf{1}$ the identity on symmetric tensors.¹³ Around flat space both the gauge-invariant part and the gauge-fixing part are those of general relativity multiplied by $a(p^2)$, so the full propagator is

$$\mathcal{D}_{\mu\nu\alpha\beta}(p) = \frac{\frac{1}{2}(\delta_{\mu\alpha}\delta_{\nu\beta} + \delta_{\mu\beta}\delta_{\nu\alpha}) - \frac{1}{2}\delta_{\mu\nu}\delta_{\alpha\beta}}{p^2 a(p^2)}. \quad (55)$$

The weight $Y = a(-\bar{\square})$ is handled in the standard way [33, 34]. Averaging the gauge condition with the Gaussian weight $e^{-\int f Y f}$ requires the normalization $(\det Y)^{1/2}$; combined with the Faddeev-Popov determinant $\det Q$, where Q is the usual second-order ghost operator, this is $\det(YQ)(\det Y)^{-1/2}$. The ghosts therefore get the action $\bar{c}^\mu Y Q_{\mu\nu} c^\nu$, and a factor $(\det Y)^{-1/2}$ depending on the background alone compensates; it contributes $+\frac{1}{2} \text{Tr} \log a(-\bar{\square})$ to the effective action and has no vertices with quantum fields. The ghost propagator is $\delta_{\mu\nu}/(p^2 a(p^2))$. Every propagator of the theory, graviton or ghost, is then bounded by

$$|\mathcal{D}(p)| \leq \frac{C}{p^2(1 + p^2/M^2)^n}, \quad (56)$$

by the lower bound in Eq. (38) and $1 + x^2 \geq \frac{1}{2}(1 + x)^2$; the constant C depends on the realization through $\|R_\Pi\|$.

B. Functions of the d'Alembertian in perturbation theory

The vertices of a nonlocal theory come from expanding $F(\square)$, $a(-\bar{\square})$ and the like in powers of the fields, and

¹³ In four dimensions $(\mathbf{1} - \frac{1}{2}\delta \otimes \delta)^2 = \mathbf{1} - \delta \otimes \delta + \frac{1}{4}(\delta \cdot \delta) \delta \otimes \delta = \mathbf{1}$, since $\delta \cdot \delta = 4$. The normalization of Eq. (54), coefficient one in front of $\chi_\mu \chi^\mu$ with the action normalized as in Eq. (52), is the one that makes the trace and traceless parts combine in this way.

a standard formula for functions of perturbed operators organizes this [35]. For points $z_0, \dots, z_m$ the divided difference $f[z_0, \dots, z_m]$ is defined by $f[z_0] = f(z_0)$ and

$$f[z_0, \dots, z_m] = \frac{f[z_1, \dots, z_m] - f[z_0, \dots, z_{m-1}]}{z_m - z_0},$$

extended to coincident points by continuity. It is symmetric in its arguments and obeys the Hermite-Genocchi formula [36]

$$f[z_0, \dots, z_m] = \int_{\Sigma_m} f^{(m)}\left(\sum_{i=0}^m t_i z_i\right) dt, \quad (57)$$

where $\Sigma_m = \{t_i \geq 0, \sum t_i = 1\}$ carries Lebesgue measure of total mass $1/m!$.

Let $A + B$ act on plane waves, with A diagonal, $Ae^{iq \cdot x} = z(q)e^{iq \cdot x}$. For entire f , the term of $f(A + B)$ of order m in B , acting on $e^{iq_0 \cdot x}$ and projected on $e^{iq_m \cdot x}$, is

$$\sum_{q_1, \dots, q_{m-1}} f[z_0, \dots, z_m] B_{q_m q_{m-1}} \cdots B_{q_1 q_0}, \quad z_i = z(q_i), \quad (58)$$

summed over the intermediate momenta that B allows. Appendix C gives the short proof. In perturbation theory $A = -\square_0$ with $z(q) = q^2$, and $B = -(\square - \square_0)$ collects the terms of the covariant d'Alembertian with at least one power of the fields. Each insertion of B is a polynomial in the fields with at most two derivatives, which act either on the fields of the insertion or on the operand. Figure 8 shows the structure.

The divided differences that appear are those of $\mathcal{G}$, of $a - cx^n$ and of polynomials. For the first two we need a bound that controls mixed configurations, in which some of the z_i are large and others small. Let f be smooth on $[0, \infty)$ with $|f^{(j)}(z)| \leq C_j(1 + z)^{\mu-j}$ for all j , for some $\mu \leq 0$. Then for every m there is a constant K_m such that, for all $z_0, \dots, z_m \geq 0$,

$$|f[z_0, \dots, z_m]| \leq K_m (1 + z_{\min})^\mu \prod_{i \neq i_{\min}} (1 + z_i)^{-1}, \quad (59)$$

where $z_{\min} = z_{i_{\min}}$ is a smallest argument. The proof, by induction on m using Eq. (57) when the arguments are comparable and the recursion when they are not, is in Appendix C. The bound says that a divided difference of a symbol of order μ decays like the symbol in its smallest argument and like an inverse power in each of the others.

C. Vertex bound

Consider any vertex of the gauge-fixed action, ghost vertices and the quartic operators S_q included, with quantum legs carrying Euclidean momenta $\ell_1, \dots, \ell_N$ and background legs carrying momenta in a fixed bounded set. We show that, for every $n \geq 1$,

$$|V| \leq C \left[\left(1 + \frac{|\ell|_{\max}}{M}\right)^{2n+2} + \prod_{j=1}^N \left(1 + \frac{|\ell_j|}{M}\right) \right], \quad (60)$$




FIG. 8. The chain behind the vertex bound. (a) A function of the covariant d'Alembertian between two field monomials, $X \mathcal{G}(-\square) Y$, expanded as in Eq. (58). Each insertion B_i of the field-dependent part of $\square$ attaches legs (dashed, gray balls) and changes the running momentum from q_{i-1} to q_i ; the function enters only through the divided difference $\mathcal{G}[q_0^2, \dots, q_m^2]$. By Eq. (61) each insertion costs one power of the running momentum on each side of it (orange), and Y and X one power each; by Eq. (59) the divided difference returns u_i^{-2} for every segment (blue). The running momenta cancel exactly, and only the legs are left. (b) A numerical check of this bookkeeping for a chain with two insertions and $n = 4$, in units $M = 1$. The running momenta are $q_0 = \ell_Y$ with $|q_0| = 1$, $q_1 = q_0 + \ell_1$ with $|q_1| \simeq \lambda_1$, and $q_2 = q_1 + \ell_2$ with $|q_2| \simeq \lambda_2$; each insertion is given its largest size $(|q_{i-1}| + |\ell_i|)^2$ and X, Y the sizes u_2, u_0 . The upper surface is $\log_{10}$ of the product of these sizes, which grows like the fifth power of the momenta, to 10^{21} ; the lower surface is $\log_{10}$ of the full term, with the exact divided difference of the skeleton's $\hat{\mathcal{G}}$, divided by $(1 + |\ell_1|)(1 + |\ell_2|)$. It stays between $10^{-3.2}$ and $10^{1.4}$ over the four decades in λ_1 and λ_2 .

with $|\ell|_{\max}$ the largest $|\ell_j|$ and C independent of the momenta. The first term bounds the local part of the vertex and the second bounds everything else. When all momenta at a vertex are of order λ the first term grows like λ^{2n+2} , as dimensional analysis requires. The second term grows by one power of λ for each hard leg, however many there are.

Set $M = 1$. Background fields never run in loops, so their momenta are external and bounded in every graph, by P say. By Eq. (39), $F(\square) = -c(-\square)^{n-1} - \mathcal{G}(-\square)$, and we also split $a(-\square) = c(-\square)^n + [a - cx^n](-\square)$. Every vertex is then a sum of terms of three kinds.

The first kind are local. They come from $cG_{\mu\nu}(-\square)^{n-1}R^{\mu\nu}$, from S_q , and from the local parts $c(-\square)^n$ of the gauge-fixing and ghost actions. Each is a polynomial in the momenta of total degree at most $2n + 2$, whatever the number of legs, so it is bounded by $C(\sum_j |\ell_j| + P)^{2n+2} \leq C(1 + |\ell|_{\max})^{2n+2}$.

The second kind contain no function of the full d'Alembertian. They come from the Einstein-Hilbert term, which has two derivatives, and from the remainders $[a - cx^n](-\square)$ in the gauge-fixing and ghost actions. In those two actions the weight stands between two factors χ_μ that are linear in the quantum field, or acts on $Q_{\mu\nu}c^\nu$, where Q has two derivatives, so the quantum legs enter through at most two derivatives. The d'Alembertian inside the weight is the background one. Expanding it by Eq. (58) produces only background insertions, each of which multiplies the operand by at most

upper: $|Y||B_1||B_2||X|$, up to 10^{21}
lower: full term/ $\prod_a (1 + |\ell_a|)$, in $[10^{-3.2}, 10^{1.4}]$

$C(1 + |q|)^2$, with q the running momentum. Since background momenta are bounded, all running momenta of such a chain lie within a fixed distance of one another, and by Eq. (59) the divided difference of the order-zero symbol $a - cx^n$ supplies a factor $(1 + q^2)^{-1}$ for every insertion. Every term of the second kind is therefore bounded by $C(1 + |\ell|_{\max})^2$. Momentum conservation gives $|\ell|_{\max} \leq (N-1)|\ell_{(2)}| + P$, with $\ell_{(2)}$ the second largest quantum momentum, so $(1 + |\ell|_{\max})^2 \leq C(1 + |\ell_{(1)}|)(1 + |\ell_{(2)}|)$, which lies within the second term of Eq. (60). (For $N = 1$ the single quantum momentum is bounded by P , and there is nothing to prove.)

The third kind contain $\mathcal{G}(-\square)$ with the full d'Alembertian, in the form $X \mathcal{G}(-\square) Y$, where X comes from $\sqrt{g}G_{\mu\nu}$ and Y from $R^{\mu\nu}$. Here the insertions carry quantum legs, which may be hard, and the estimate has to be made with some care. By Eq. (58) the term of order m is $\mathcal{G}[z_0, \dots, z_m] B_m \cdots B_1$, with $z_i = q_i^2$. The running momentum q_0 is the total momentum of the legs of Y , and $q_i = q_{i-1} + \sum_{a \in i} \ell_a$, the sum running over the legs of the i th insertion. Write $u_i = 1 + |q_i|$ and $s_i = \sum_{a \in i} |\ell_a|$. An insertion has at most two derivatives, acting on its own legs or on the operand, so $|B_i| \leq C(|q_{i-1}| + s_i)^2$; in the same way $|Y| \leq Cs_Y^2$ and $|X| \leq Cs_X^2$. The estimate that does the work, proved in Appendix C, is

$$(|q| + s)^2 \leq C_r (1 + |q|)(1 + |q'|) \prod_{a=1}^r (1 + |\ell_a|), \quad (61)$$

for any q and legs $\ell_1, \dots, \ell_r$, with $q' = q + \sum_a \ell_a$ and

$s = \sum_a |\ell_a|$. An insertion costs one power of each of the two running momenta next to it and one power of each of its own legs. Applying Eq. (61) to every insertion, to Y (with $q = 0$, $q' = q_0$) and to X (with $q = 0$, $q' = -q_m$), since the legs of X carry away the momentum q_m , the running momenta enter through

$$u_0 u_m \prod_{i=1}^m u_{i-1} u_i = \prod_{i=0}^m u_i^2.$$

On the other side, $\mathcal{G}$ is a symbol of order -1 by Eq. (40), and Eq. (59) with $\mu = -1$ gives $|\mathcal{G}[z_0, \dots, z_m]| \leq K \prod_{i=0}^m (1 + z_i)^{-1} \leq 2^{m+1} K \prod_{i=0}^m u_i^{-2}$. The running momenta cancel exactly, and the term is bounded by $C \prod_a (1 + |\ell_a|)$ over all its legs. The sum over intermediate momenta in Eq. (58) is a finite sum over the ways of distributing the legs among the insertions, which proves Eq. (60).¹⁴

D. Superficial degree

Let γ be a connected subgraph with L loops of some graph of the theory, and consider the region in which the loop momenta of γ are of order $\lambda \rightarrow \infty$ while all other momenta are fixed. Let γ have I internal lines and V vertices, so that $I - V = L - 1$, and let h_v be the number of lines of γ at the vertex v , so that $\sum_v h_v = 2I$. Every vertex of γ has $h_v \geq 2$, because a line of γ carries loop momentum and momentum conservation at its ends needs a second line that does. Each line contributes at most $C\lambda^{-2n-2}$ by Eq. (56). At a vertex of γ the legs outside γ carry fixed momenta, so Eq. (60) contributes at most $C\lambda^{\max(2n+2, h_v)}$. The measure contributes λ^{4L} . With $x_+ = \max(x, 0)$ the superficial degree is therefore at most

$$\begin{aligned} 4L - (2n+2)I + \sum_v \max(2n+2, h_v) \\ = 4L - (2n+2)(L-1) + \sum_v (h_v - 2n - 2)_+. \end{aligned}$$

The last sum is largest when all the lines beyond the minimum meet at one vertex. Indeed $(a-k)_+ + (b-k)_+ \leq (a+b-2-k)_+$ whenever $a, b \geq 2$ and $k \geq 2$,¹⁵ so merging the vertices one pair at a time gives $\sum_v (h_v - 2n - 2)_+ \leq$

$(\sum_v h_v - 2(V-1) - 2n - 2)_+ = (2L - 2n - 2)_+$. Hence the superficial degree of an L -loop subgraph is at most

$$\bar{\omega}(L) = 4L - (2n+2)(L-1) + 2(L-n-1)_+. \quad (62)$$

For $L \leq n+1$ this is the standard degree $2n+2+(2-2n)L$ of local higher-derivative gravity [11, 37], and for $L > n+1$ it is $(4-2n)L$. At one loop $\bar{\omega}(1) = 4$ for every n . At two loops $\bar{\omega}(2) = 6 - 2n$, and for $n \geq 3$ the degree decreases with L on both sides of $L = n+1$. So $\bar{\omega}(L) < 0$ for every $L \geq 2$ exactly when $n \geq 4$. Table V and Fig. 9 give the numbers.

To turn the degree into a convergence statement we use Weinberg's theorem [38]. It applies to integrands in his class A_n of functions with power-law behavior along every subspace of the integration variables, a class that contains rational functions and products of factors $1 + |\ell|$ with ℓ linear in those variables. It states that the integral converges absolutely if the asymptotic degree of every subspace is negative. On a large-momentum line the factor $1/p^2$ in Eq. (56) may be replaced by $1/(p^2 + \mu^2)$ with any fixed $\mu > 0$, which changes nothing in the ultraviolet.¹⁶ At each vertex, $(1 + |\ell|_{\max})^{2n+2} \leq \sum_j (1 + |\ell_j|)^{2n+2}$, so the bound (60) is dominated by a finite sum of terms, each either a product $\prod_j (1 + |\ell_j|)$ or a single power $(1 + |\ell_j|)^{2n+2}$ of one “chosen” leg. Expanding the product over vertices, the integrand is dominated by a finite sum of functions in A_n . In each of them, the degree in the region where γ is hard is at most the count above, since a vertex whose chosen leg lies outside γ contributes less. Weinberg's theorem therefore applies to each dominating term, and for $n \geq 4$ every graph and every subgraph with two or more loops converges in the ultraviolet. The only subgraphs that can diverge are one-loop subgraphs, for which $\bar{\omega} = 4$.

The threshold $n \geq 4$ is the one found in the literature by bounding every vertex by λ^{2n+2} , which is legitimate when all momenta at a vertex are comparable [11, 37]. The vertex bound shows that vertices with many hard legs do not spoil it: the nonlocal remainder gains at most one power of λ per hard leg, while each hard leg brings a propagator that falls like λ^{-2n-2} . For $n \leq 3$ the two-loop degree is not negative and the argument gives nothing. From now on $n \geq 4$.

¹⁴ The cancellation is not an artifact of the bounds. When the running momentum rises from soft to hard at one insertion and falls back at the next, the divided difference is of order λ^{-2} and the two insertions of order λ^2 each, so the term grows like λ^2 while two hard legs are attached: one power per hard leg, as Eq. (60) says. Appendix D lists a numerical check of the bound on random chains with up to three insertions.

¹⁵ If a and b both exceed k , the left side is $a+b-2k \leq a+b-2-k$ because $k \geq 2$. If only a does, it is $a-k \leq a+b-2-k$ because $b \geq 2$. If neither does, it is zero.

¹⁶ This separates the ultraviolet from the infrared. Infrared behavior is governed by $a \rightarrow 1$ and is that of general relativity; it is discussed in Sec. VI.

TABLE V. Superficial degree $\bar{\omega}(L) = 4L - (2n+2)(L-1) + 2(L-n-1)_+$ from the vertex bound (60), for $L = 1, \dots, 5$. Negative entries are convergent. For $L \leq n+1$ the entries coincide with the standard estimate $2n+2+(2-2n)L$; the first entry that differs is $n=3, L=5$.

n	$L=1$	$L=2$	$L=3$	$L=4$	$L=5$
3	4	0	-4	-8	-10
4	4	-2	-8	-14	-20
5	4	-4	-12	-20	-28
6	4	-6	-16	-26	-36
7	4	-8	-20	-32	-44
8	4	-10	-24	-38	-52




FIG. 9. Upper bound $\bar{\omega}(L)$ on the superficial degree of an L -loop graph, Eq. (62), for $n = 3, \dots, 8$ and $L \leq 6$, drawn as bars hanging from the plane $\bar{\omega} = 0$. One-loop graphs have degree 4 for every n (red). For $n=3$ the two-loop bound is 0 (amber), so power counting alone leaves a possible logarithm; for $n \geq 4$ every graph with $L \geq 2$ has negative degree (blue to green), and the degree falls without bound as L grows.

V. ONE LOOP

A. The local theory

Split the action into a local part and a remainder,

$$S_{\text{loc}} = \frac{2c}{\kappa^2 M^{2n}} \int \sqrt{g} G_{\mu\nu}(-\square)^{n-1} R^{\mu\nu} + S_q, \quad (63)$$

$$S_{\text{rest}} = -\frac{2}{\kappa^2} \int \sqrt{g} [R - G_{\mu\nu} \mathcal{G}(-\square) R^{\mu\nu}],$$

so that $S = S_{\text{loc}} + S_{\text{rest}}$ by Eq. (39). Split the gauge-fixing weight in the same way, $a(-\square) = c(-\square/M^2)^n + [a - cx^n](-\square)$. Both parts of the action are separately invariant under diffeomorphisms, and both parts of the gauge-fixing term are background covariant. The local theory, with action S_{loc} , gauge-fixing weight $c(-\square/M^2)^n$

and the corresponding ghosts, is therefore a consistent background-gauge-fixed theory on its own. Its flat-space propagator is $M^{2n}(P^{(2)} - \frac{1}{2}P_s^{(0)})/(cp^{2n+2})$ on the gauge-invariant subspace, from Eq. (53) with a replaced by cx^n .

The local theory is badly behaved in the infrared, since its propagator falls like p^{-2n-2} at small p as well as at large p . This does not matter for what follows, because we compare the two theories only at large loop momentum. We define the ultraviolet pole of a one-loop integral as the pole, in dimensional regularization with $D = 4 - 2\epsilon$, of its integral over the region $|k| \geq M$ of the loop momentum.¹⁷ With this definition the ultraviolet pole of the full theory and that of the local theory can be compared graph by graph.

B. One-loop poles are those of the local theory

Let $n \geq 4$, and consider any one-loop graph of the full theory with external legs, background or quantum, of bounded momenta. Write each vertex as $V = V_{\text{loc}} + V_{\text{rest}}$ and each propagator as $\mathcal{D} = \mathcal{D}_{\text{loc}} + \mathcal{D}_{\text{rest}}$, where the local pieces are those of the local theory. For loop momentum $|k| \geq M$ we have $|V_{\text{loc}}| \leq C|k|^{2n+2}$ and $|\mathcal{D}_{\text{loc}}| \leq C|k|^{-2n-2}$. The remainders are smaller. A remainder vertex consists of terms of the second and third kinds of Sec. IV C, which are bounded by the product term of Eq. (60); two of its legs carry the loop momentum, so $|V_{\text{rest}}| \leq C|k|^2$. And $|\mathcal{D}_{\text{rest}}| \leq C|k|^{-2n-2-\sigma+2\epsilon}$, because $1/a - 1/(cx^n) = (cx^n)^{-1}(e^{-\eta_{\Pi}} - 1)$ with η_{Π} bounded by Eq. (37). Expanding the product around the loop, the term with only local factors is the integrand of the local theory. Every other term contains at least one V_{rest} or one $\mathcal{D}_{\text{rest}}$. For a loop with r vertices it is bounded for $|k| \geq M$ either by $C|k|^{(2n+2)(r-1)+2-(2n+2)r} = C|k|^{-2n} \leq C|k|^{-8}$, if it contains a remainder vertex, or by $C|k|^{-\sigma+2\epsilon} \leq C|k|^{-2n}$, if it contains a remainder propagator, since $\sigma > 2n$ and ϵ may be taken small. Its integral over $|k| \geq M$ converges for $D < 2n$, a range that contains $D = 4$ for every $n \geq 3$, and has no pole there. The ultraviolet pole of the full graph is that of the local graph.

The compensating factor of Sec. IV A contributes $\frac{1}{2} \text{Tr} \log a(-\square)$, which at large Euclidean momenta equals $\frac{n}{2} \text{Tr} \log(-\square/M^2) + \frac{1}{2} \log c \text{Tr} \mathbf{1} + \frac{1}{2} \text{Tr} \eta_{\Pi}(-\square)$ by Eq. (37). The trace of a constant vanishes in dimensional regularization, the last trace has a symbol that falls like $|z|^{-\sigma/2+\epsilon}$, faster than $|z|^{-2}$, and has no pole, and the first has the pole of $\text{Tr} \log(-\square)$ on vectors, a local functional

¹⁷ For a one-loop integrand $I(k, p)$ with bounded external momenta p , the region $|k| \geq M$ can be split into $M \leq |k| \leq K$, which is compact and gives no pole, and $|k| \geq K$ with K large compared with the external momenta, where I may be expanded in p . Only the terms of the expansion that are homogeneous of degree $-D$ in k produce a pole, and their coefficients are polynomials in p . The pole is therefore local and independent of the choice of M and K . With this definition $\int_{|k| \geq M} d^D k (2\pi)^{-D} k^{-4} = 1/(16\pi^2\epsilon) + O(1)$.

of dimension four. This is the pole of the compensating factor of the local theory.

C. The form of the pole

In the local theory each graviton vertex carries the factor $c/(\kappa^2 M^{2n})$ of Eq. (63), times κ^k for k quantum legs, and each graviton propagator carries $\kappa^0 M^{2n}/c$; the quartic operators enter only through the ratios of their coefficients to c ; the ghosts carry c/M^{2n} at each vertex and the inverse on each propagator. Write the local action as $\lambda^{-1} \int \sqrt{g} \mathcal{L}$, with $\lambda = \kappa^2 M^{2n}/(2c)$ and $\mathcal{L}$ free of dimensionful constants. The one-loop effective action is of order λ^0 , so its ultraviolet pole, as a local functional of the full metric $\bar{g} + \kappa Q$ and of the other fields, contains no dimensionful constant at all. By background gauge invariance [32, 39] its part depending on $\bar{g}$ alone is a diffeomorphism-invariant local functional, and a dimensionless local invariant built from the metric has exactly four derivatives. In four dimensions these invariants are $\bar{R}^2$, $\bar{R}_{\mu\nu}\bar{R}^{\mu\nu}$, $\bar{R}_{\mu\nu\rho\sigma}\bar{R}^{\mu\nu\rho\sigma}$ and $\square\bar{R}$; the last is a total derivative, and the Riemann square may be traded for the Gauss-Bonnet density $\bar{E} = \bar{R}_{\mu\nu\rho\sigma}\bar{R}^{\mu\nu\rho\sigma} - 4\bar{R}_{\mu\nu}\bar{R}^{\mu\nu} + \bar{R}^2$. Hence

$$\Gamma_1^{\text{div}} = \frac{1}{16\pi^2\epsilon} \int d^4x \sqrt{\bar{g}} (\beta_1 \bar{R}^2 + \beta_2 \bar{R}_{\mu\nu}\bar{R}^{\mu\nu} + \beta_E \bar{E}), \quad (64)$$

with $\beta_1, \beta_2, \beta_E$ pure numbers that depend on n and on the coefficients of S_q in units of c , and on nothing else.

In particular there is no pole in $\int \sqrt{\bar{g}}$ or $\int \sqrt{\bar{g}} \bar{R}$: neither the cosmological constant nor Newton's constant is renormalized at one loop. Those two terms need the dimensionful factors M^4 and M^2 , and in the ultraviolet the only place they could come from is the correction to the monomial asymptotics of a . For a realization that correction is of relative order $|z|^{-\sigma/2+\epsilon}$ with $\sigma/2 > n \geq 4$, and Sec. VB shows that it contributes only convergent integrals. A single-brane response that saturated only as a power of $1/|sz|$ would give, through Eq. (16), corrections of relative order x^{-1} or x^{-2} and in general a running of both constants, and so would a gas whose fluctuations died out too slowly. Conditions (S3) and (G) prevent it. The explicit one-loop computation of Modesto, Rachwał and Shapiro [40] agrees. For form factors in front of $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$ and R^2 that grow like polynomials of degree N , they find a beta function of Newton's constant that is a linear combination of the ratios ω_{N-1}/ω_N of the two leading coefficients of those polynomials. In Eq. (46) both form factors are multiples of F , and by Eq. (39) $F(z)$ equals $cz^{n-1}/M^{2n} - 1/z$ up to terms of order $|z|^{n-1-\sigma/2+\epsilon}$, which fall faster than $1/z$. It has no term in z^{n-2} , so both ratios vanish, and so does the beta function.

The numbers β_i do not depend on the gauge condition or on its weight within the class used here. At one loop the change of the effective action under a change of gauge is proportional to the classical equations of motion [41],

$\delta\Gamma_1 = \int (\delta S/\delta \bar{g}_{\mu\nu}) Y_{\mu\nu}$ with Y a functional of $\bar{g}$. Apply this to the local theory, whose poles these are. The pole of $\delta\Gamma_1$ is $\int (\delta S_{\text{loc}}/\delta \bar{g}_{\mu\nu}) Y_{\mu\nu}^{\text{div}}$, with Y^{div} local. Every term of $\delta S_{\text{loc}}/\delta \bar{g}_{\mu\nu}$ has $2n+2 \geq 10$ derivatives of the metric, and a local Y^{div} cannot remove derivatives, so every term of the product has at least that many and none can appear in Eq. (64), which has four. Hence $\delta\Gamma_1$ has no pole.

The values of $\beta_1, \beta_2, \beta_E$ without the quartic operators, which we call $\beta_i^{(0)}$, are fixed by n . Computing them is a well-defined exercise in the generalized Schwinger-DeWitt technique for a nonminimal operator of order $2n+2$ [39]. We do not need their values, only the fact that they are finite numbers.

D. Three quartic operators

Following Modesto and Rachwał [12], we add operators that are quartic in curvature and can shift the coefficients in Eq. (64). We need three, one for each coefficient:

$$S_q = \frac{2}{\kappa^2 M^{2n}} \int d^4x \sqrt{g} \left[s_1 R^2 (-\square)^{n-3} R^2 + s_2 (R_{\mu\nu} R^{\mu\nu}) (-\square)^{n-3} (R_{\rho\sigma} R^{\rho\sigma}) + s_3 (R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}) (-\square)^{n-3} (R_{\alpha\beta\gamma\delta} R^{\alpha\beta\gamma\delta}) \right], \quad (65)$$

with real dimensionless s_1, s_2, s_3 . Each term has $2n+2$ derivatives, as many as the leading term of S_{loc} , so the vertex bound (60) still holds. Being quartic in curvature, the operators change neither the propagator nor the three-graviton vertex around flat space, so the spectrum of Sec. IIIC and the Newtonian limit of Sec. VIII are untouched. Their contribution to Eq. (64) can be computed exactly.

Which terms contribute. The operators are linear in s_i . Every term of their second variation around $\bar{g}$ contains at least two factors of background curvature, so a contribution to Γ_1 of second or higher order in the s_i carries at least four background curvatures and eight derivatives, and cannot appear in Eq. (64). To first order the one-loop contribution is the expectation value of the quadratic part of S_q in the free local theory, $\frac{1}{2} \text{Tr}(\mathcal{D} S_q'') = \langle S_q^{(2)} \rangle$. Of the terms in $S_q^{(2)}$ only those with exactly two background curvatures and with $(-\square)^{n-3}$ acting on a quantum field can produce a pole in Eq. (64). When derivatives fall on the background the operator has more than four derivatives; and a term in which $(-\square)^{n-3}$ acts on the product of two background curvatures carries $2n-6 \geq 2$ derivatives on the background in addition to the four in the two curvatures, six or more in all. Corrections to the linearized curvatures, to $\square$ and to the index contractions that involve the background curvature add more background curvatures. Writing $R^2 = \bar{R}^2 + 2\bar{R}\delta R + \dots$ and

similarly for the other invariants, the relevant terms are

$$\begin{aligned} & \frac{8}{\kappa^2 M^{2n}} \left[s_1 \bar{R}^2 \delta R (-\square)^{n-3} \delta R \right. \\ & \quad + s_2 \bar{R}^{\mu\nu} \bar{R}^{\rho\sigma} \delta R_{\mu\nu} (-\square)^{n-3} \delta R_{\rho\sigma} \\ & \quad \left. + s_3 \bar{R}^{\mu\nu\rho\sigma} \bar{R}^{\alpha\beta\gamma\delta} \delta R_{\mu\nu\rho\sigma} (-\square)^{n-3} \delta R_{\alpha\beta\gamma\delta} \right], \quad (66) \end{aligned}$$

where δR , $\delta R_{\mu\nu}$, $\delta R_{\mu\nu\rho\sigma}$ are the flat-space linearized curvatures of $\delta g = \kappa h$ and the background curvatures are treated as constants.

Correlators. The linearized curvatures are gauge invariant, so only the gauge-invariant part $M^{2n}(P^{(2)} - \frac{1}{2}P_s^{(0)})/(ck^{2n+2})$ of $\langle hh \rangle$ contributes, and the result does not depend on the gauge-fixing term. From Eq. (47), $\delta R_{\mu\nu} = \frac{\kappa k^2}{2}[(P^{(2)}h)_{\mu\nu} + (\theta \cdot h)T_{\mu\nu}]$. Since $P^{(2)}P_s^{(0)} = 0$ and $P^{(2)}\theta = 0$ the mixed terms vanish; $\langle (P^{(2)}h)(P^{(2)}h) \rangle$ gives $P^{(2)}$ times the scalar factor, and $\langle (\theta \cdot h)^2 \rangle$ gives $\theta \cdot (-\frac{1}{2}P_s^{(0)}) \cdot \theta = -\frac{3}{2}$ times the same factor, because $\theta P_s^{(0)}\theta = D - 1 = 3$. Thus, in $D = 4$,

$$\langle \delta R_{\mu\nu}(k) \delta R_{\rho\sigma}(-k) \rangle = \frac{\kappa^2 M^{2n}}{4c k^{2n-2}} \left[P^{(2)} - \frac{3}{2} T \otimes T \right]_{\mu\nu\rho\sigma}, \quad (67)$$

$$\langle \delta R(k) \delta R(-k) \rangle = -\frac{3}{2} \frac{\kappa^2 M^{2n}}{c k^{2n-2}}, \quad (68)$$

the second following from the first with $\delta \cdot P^{(2)} \cdot \delta = 0$ and $T \cdot \delta = 2$.¹⁸ For the Riemann tensor, the linearized curvature is $\delta R_{\mu\nu\rho\sigma} = -\frac{\kappa}{2}(k_\nu k_\rho h_{\mu\sigma} + k_\mu k_\sigma h_{\nu\rho} - k_\mu k_\rho h_{\nu\sigma} - k_\nu k_\sigma h_{\mu\rho})$. Contracted with an algebraic curvature tensor $\bar{R}^{\mu\nu\rho\sigma}$, the four terms are equal by its symmetries, and

$$\bar{R}^{\mu\nu\rho\sigma} \delta R_{\mu\nu\rho\sigma} = 2\kappa k^2 W^{\nu\sigma} h_{\nu\sigma}, \quad W^{\nu\sigma} = \bar{R}^{\mu\nu\rho\sigma} \hat{k}_\mu \hat{k}_\rho, \quad (69)$$

with $\hat{k} = k/|k|$. The tensor W is symmetric and transverse, $\hat{k}_\nu W^{\nu\sigma} = 0$, so $W \cdot P^{(2)} \cdot W = \text{tr } W^2 - \frac{1}{3}(\text{tr } W)^2$ and $W \cdot P_s^{(0)} \cdot W = \frac{1}{3}(\text{tr } W)^2$, and

$$\langle (\bar{R} \cdot \delta \text{Riem})^2 \rangle = \frac{4\kappa^2 M^{2n}}{c k^{2n-2}} \left[\text{tr } W^2 - \frac{1}{2}(\text{tr } W)^2 \right]. \quad (70)$$

Poles. In the loop integral $(-\square)^{n-3} \rightarrow (k^2)^{n-3}$, and the correlators give integrands proportional to $k^{2n-6}k^{-2n+2} = k^{-4}$, whose ultraviolet pole is $1/(16\pi^2\epsilon)$. The tensors must be averaged over the direction of k , with $\langle \hat{k}_\mu \hat{k}_\nu \rangle = \delta_{\mu\nu}/4$ and $\langle \hat{k}_\mu \hat{k}_\nu \hat{k}_\rho \hat{k}_\sigma \rangle = (\delta_{\mu\nu}\delta_{\rho\sigma} +$

TABLE VI. One-loop poles of the quartic operators (65), in units of $\frac{1}{16\pi^2\epsilon} \frac{s_i}{c}$. The second column is the correlator after contraction with the background and averaging over the direction of k , in units of $\kappa^2 M^{2n}/(c k^{2n-2})$.

operator	averaged correlator	pole
$s_1 R^2 \square^{n-3} R^2$	$-\frac{3}{2} R^2$	$-12R^2$
$s_2 \text{Ric}^2 \square^{n-3} \text{Ric}^2$	$\frac{1}{8}(\text{Ric}^2 - R^2)$	$\text{Ric}^2 - R^2$
$s_3 \text{Riem}^2 \square^{n-3} \text{Riem}^2$	$\frac{1}{12}(3\text{Riem}^2 - R^2)$	$2E + 8\text{Ric}^2 - \frac{8}{3}R^2$

$\delta_{\mu\rho}\delta_{\nu\sigma} + \delta_{\mu\sigma}\delta_{\nu\rho})/24$. Appendix B shows that

$$\bar{R}^{\mu\nu} \bar{R}^{\rho\sigma} \left\langle P^{(2)} - \frac{3}{2} T \otimes T \right\rangle_{\mu\nu\rho\sigma} = \frac{1}{2}(\text{Ric}^2 - R^2), \quad (71)$$

$$\left\langle \text{tr } W^2 - \frac{1}{2}(\text{tr } W)^2 \right\rangle = \frac{1}{48}(3\text{Riem}^2 - R^2), \quad (72)$$

where on the right R , Ric^2 and Riem^2 are the background invariants. Multiplying by the coefficients $8s_i/(\kappa^2 M^{2n})$ of Eq. (66), the factors $\kappa^2 M^{2n}$ cancel, and in units of $1/(16\pi^2\epsilon)$ the three poles are

$$\begin{aligned} s_1 : & \quad 8s_1 \cdot \left(-\frac{3}{2c}\right) R^2 = -\frac{12s_1}{c} R^2, \\ s_2 : & \quad 8s_2 \cdot \frac{1}{4c} \cdot \frac{1}{2}(\text{Ric}^2 - R^2) = \frac{s_2}{c}(\text{Ric}^2 - R^2), \\ s_3 : & \quad 8s_3 \cdot \frac{4}{c} \cdot \frac{3\text{Riem}^2 - R^2}{48} = \frac{s_3}{c} \left(2\text{Riem}^2 - \frac{2}{3}R^2\right). \end{aligned}$$

With $\text{Riem}^2 = E + 4\text{Ric}^2 - R^2$ the third line is $(s_3/c)(2E + 8\text{Ric}^2 - \frac{8}{3}R^2)$. Collecting coefficients,

$$\begin{pmatrix} \Delta\beta_1 \\ \Delta\beta_2 \\ \Delta\beta_E \end{pmatrix} = \frac{1}{c} \begin{pmatrix} -12 & -1 & -\frac{8}{3} \\ 0 & 1 & 8 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} s_1 \\ s_2 \\ s_3 \end{pmatrix}. \quad (73)$$

The matrix is triangular with determinant -24 , so $\partial(\Delta\beta_1, \Delta\beta_2, \Delta\beta_E)/\partial(s_1, s_2, s_3)$ has determinant $-24/c^3$, which never vanishes. The diagrams are drawn in Fig. 10(b), and Table VI collects the pieces. In the dimensional-regularization computation the angular averages are taken in D dimensions; the difference from the four-dimensional values is of order ϵ and affects finite parts only.

Solving $\beta_i^{(0)} + \Delta\beta_i = 0$ from the bottom row up gives the tuning in closed form:

$$\begin{aligned} s_3 &= -\frac{c}{2} \beta_E^{(0)}, & s_2 &= -c(\beta_2^{(0)} - 4\beta_E^{(0)}), \\ s_1 &= \frac{c}{12}(\beta_1^{(0)} + \beta_2^{(0)} - \frac{8}{3}\beta_E^{(0)}). \end{aligned} \quad (74)$$

With this choice the one-loop background-field effective action has no ultraviolet pole of any kind, the Gauss-Bonnet one included.¹⁹ By the gauge independence of the β_i , the tuning does not depend on the gauge.

¹⁸ The negative sign in Eq. (68) is the conformal-factor sign of the Euclidean Einstein-Hilbert action, visible in the $-2P_s^{(0)}$ of Eq. (52). In dimensional regularization the one-loop computation is Gaussian algebra on the propagator (53), and the sign is carried through consistently.

¹⁹ Cancelling the Gauss-Bonnet pole is not needed for the S ma-




FIG. 10. Graphs behind the argument. Wavy lines are quantum gravitons (or ghosts), dashed lines end on background insertions $\otimes$. (a) A one-loop ring contributing to $\Gamma_1[\bar{g}]$. Replacing any vertex or propagator by its nonlocal remainder lowers the degree by at least $2n$, so the ultraviolet pole is that of the local theory. (b) The insertion of a quartic operator with two background curvatures: the loop of $(-\square)^{n-3}$ between two linearized curvatures has an integrand $\propto k^{-4}$ and gives the poles of Table VI. (c) A two-loop graph. The graph as a whole is convergent for $n \geq 4$; the shaded one-loop subgraph γ has degree 4 and is the only possible source of a subdivergence.

VI. FINITENESS AT ALL ORDERS

Assume A1 and A2, let $n \geq 4$ be an integer, and choose s_1, s_2, s_3 as in Eq. (74), with $c = c_\Pi$ the leading coefficient of the realization, Eq. (37). Then, for almost every realization of the gas, in dimensional regularization:

- (i) The one-loop background-field effective action $\Gamma_1[\bar{g}]$ has no ultraviolet pole. This is Eq. (64) with the coefficients $\beta_i^{(0)} + \Delta\beta_i = 0$ of Eqs. (73) and (74).
- (ii) Every graph with two or more loops, and every subgraph with two or more loops, is ultraviolet convergent; the only subdivergences are those of one-loop subgraphs. This is Sec. IV D with $n \geq 4$.
- (iii) No coupling of the gauge-invariant action is renormalized at any order: not Newton's constant, not the cosmological constant, not M , n or the s_i . Both constants receive finite one-loop shifts, which can be computed (Sec. VIB). The S matrix is ultraviolet finite to all orders.

The argument for (iii) is as follows. Consider the full effective action $\Gamma[\bar{g}, Q]$, with Q the expectation values of the quantum fields and ghosts, and with sources for their BRST variations. By (ii) and the forest structure of dimensional regularization [42], all counterterms are fixed by one-loop poles; no new counterterm is needed at higher loops, because there is no overall divergence there. The one-loop poles are local. Section VB shows that they equal those of the local theory, for graphs with background and with quantum external legs alike, and by the

Slavnov–Taylor identity and the cohomological classification of Ref. [43], applied to the local theory, the one-loop counterterm is the sum of a gauge-invariant functional of the full metric $\bar{g} + \kappa Q$ and a BRST-exact part. The gauge-invariant part has four derivatives and no dimensionful constant (Sec. VC), so it is $\beta_1 R^2 + \beta_2 \text{Ric}^2 + \beta_E E$ of the full metric, fixed by its value at $Q = 0$, which vanishes by (i). Only the BRST-exact part remains. It renormalizes the quantum fields, the ghosts and the gauge-fixing sector, and it is needed to subtract the subdivergences of off-shell subgraphs at two loops and beyond. A BRST-exact counterterm is a change of the gauge-fixing fermion, and such a change alters the background-field effective action only by terms proportional to its own equations of motion [41, 43]. The S matrix, which is built from $\Gamma[\bar{g}]$ by tree graphs [44], is therefore unaffected, and no physical coupling receives a counterterm. The tuning (74) is made once, at one loop, and is not disturbed later, because at higher loops there is nothing to disturb it.

Table VIII lists every kind of divergence that can arise, with its fate. The ultraviolet ones are removed: the dimensionful ones by saturation, the one-loop dimensionless ones by the quartic operators, and the higher-loop ones by power counting. What remains are the renormalizations of the gauge-fixing sector, which have no effect on observables, and the infrared divergences of on-shell amplitudes with soft gravitons. The latter are physical. They are those of general relativity, since $a = 1 + O(p^4/M^4)$ at small momentum, and they cancel in inclusive rates by Weinberg's soft-graviton theorem [18]. No ultraviolet completion can or should remove them.

A. No cutoff in the answer

“Finite” means different things in quantum field theory, and the sense used here should be stated. In the usual procedure a divergent integral is cut off at a momentum

trix in four dimensions, because $\int \sqrt{g} E$ is topological there. It is convenient all the same: in $D = 4 - 2\epsilon$ the density E has vertices proportional to $D - 4$, and a $1/\epsilon$ counterterm in E would leave finite evanescent contributions when inserted in higher-loop graphs. With s_3 tuned no such counterterm exists.

Λ , the dependence on Λ is absorbed into bare couplings, and Λ is then sent to infinity while the bare couplings follow it. Nothing of that kind happens in the theory of the gas.

The bare couplings G , M , n and s_1, s_2, s_3 are fixed numbers. None of them depends on the regulator, and none is adjusted as the regulator is removed. In dimensional regularization the S matrix computed from the bare action (46) has a finite limit as $\epsilon \rightarrow 0$, and that limit is the physical S matrix. This follows from the argument for (iii). With the BRST-exact counterterms added, every subdivergence is subtracted and the result is finite; those counterterms do not change the S matrix; so the S matrix is the same with and without them. Dependence on the regulator survives only in off-shell Green functions of the quantum fields and ghosts, which are not observable.

The logarithm is the only divergence of a one-loop integral that takes the same form in every regularization. Its coefficient is the residue of the pole in dimensional regularization, a local functional, and the tuning (74) sets it to zero. A regularization with a momentum cutoff produces, in addition, terms growing like Λ^4 and Λ^2 in front of $\int \sqrt{g}$ and $\int \sqrt{g} R$. Their coefficients change when the shape of the cutoff is changed, and they are absent in dimensional regularization and in every other analytic regularization [42]. They carry no information about the theory. The finite remainder does, and the next subsection shows on the simplest example that it is the same for every regulator.

B. The zero-point energy of the graviton

The absence of a pole in $\int \sqrt{g}$ does not mean that the one-loop effective action has no such term. It has a finite one, and it can be computed exactly. On flat space $\Gamma_1 = (\rho_1 + \delta\rho_\Pi) \int d^4x$, where $\rho_1 + \delta\rho_\Pi$ is the zero-point energy density of the graviton together with the ghosts and the compensating factor of Sec. IV A; ρ_1 comes from the skeleton and $\delta\rho_\Pi$ from the fluctuation R_Π of the realization. Around flat space the graviton operator is $p^2 a(p^2)(\mathbf{1} - \frac{1}{2}\delta \otimes \delta)$ on the ten components of $h_{\mu\nu}$, the ghost operator is $p^2 a(p^2)\delta_{\mu\nu}$ on four components, and the compensating factor is $\frac{1}{2} \text{Tr} \log a$ on four components. A real bosonic component contributes $\frac{1}{2} \log$ of its eigenvalue and a ghost–antighost pair contributes $-\log$, so per momentum the integrand is

$$5 \log(p^2 a) - 4 \log(p^2 a) + 2 \log a = \log p^2 + 3 \log a, \quad (75)$$

up to a constant from $\det(\mathbf{1} - \frac{1}{2}\delta \otimes \delta)$ that, like $\int d^D k$ of any constant, vanishes in dimensional regularization. The same integrand follows without gauge fixing. The six transverse components of $h_{\mu\nu}$ each contribute $\frac{1}{2} \log(p^2 a)$, and the Jacobian of the split of $h_{\mu\nu}$ into its transverse part and a gauge orbit contributes $-2 \log p^2$, with no factor of a . The coefficient 3 of $\log a$ is therefore independent of the gauge.

The first term in Eq. (75) is that of general relativity. Its integral $\int d^D k \log k^2$ has no scale and vanishes in dimensional regularization. The second term is $3 \log a = \frac{3n}{2} \text{Ein}(k^4/M^4) + 3R_\Pi(k^2)$, by Eq. (13). We take the skeleton part first. With $w = k^4/M^4$ and $\Omega_D = 2\pi^{D/2}/\Gamma(D/2)$ the area of the unit sphere,

$$\int d^D k \text{Ein}\left(\frac{k^4}{M^4}\right) = \frac{\Omega_D M^D}{4} \int_0^\infty w^{D/4-1} \text{Ein}(w) dw.$$

The last integral is a Mellin transform, and for Ein it is elementary. For $-1 < \text{Re } s < 0$ one has $\int_0^\infty w^{s-1} (1 - e^{-wt}) dw = -\Gamma(s) t^{-s}$. Integrating over $t \in [0, 1]$ with the weight dt/t , as in the definition of Ein , gives

$$\int_0^\infty w^{s-1} \text{Ein}(w) dw = \frac{\Gamma(s)}{s}, \quad -1 < \text{Re } s < 0. \quad (76)$$

The strip is where the integral converges, since $\text{Ein}(w) \simeq w$ at small w and $\text{Ein}(w) \simeq \log w$ at large w . Dimensional regularization defines the radial integral by continuation in D [42], that is in $s = D/4$, and the right side of Eq. (76) is meromorphic with poles only at $s = 0, -1, -2, \dots$. At $D = 4$, that is $s = 1$, it is regular and equals $\Gamma(1) = 1$. Hence $\int d^4 k (2\pi)^{-4} \text{Ein}(k^4/M^4) = 2\pi^2 M^4 / (4 \cdot 16\pi^4) = M^4 / (32\pi^2)$, and

$$\rho_1 = \frac{3n}{2} \frac{M^4}{32\pi^2} = \frac{3n M^4}{64\pi^2}. \quad (77)$$

There is no pole, as Sec. V C requires, and the value is positive. In the normalization of Eq. (46), where a cosmological term enters as $-\frac{2}{\kappa^2} \int \sqrt{g} (-2\Lambda)$, Eq. (77) is a finite cosmological constant $\Lambda_1 = \frac{\kappa^2}{4} \rho_1 = 3nGM^4/(8\pi)$. For a general admissible response the second term of Eq. (75) is $3n\Phi(k^2/M^2)$, and Eq. (29) gives

$$\rho_1 = \frac{3nM^4}{32\pi^2} \mu_\psi = \frac{3nM^4}{32\pi^2} \int_0^\infty t (1 - \psi(t)) dt, \quad (78)$$

positive and finite for every response that saturates; Table I lists three values.

The fluctuation term is the realization's own contribution. In dimensional regularization the integral of the constant R_∞ vanishes, and what is left, $R_\Pi - R_\infty$, falls along the Euclidean axis like $k^{-\sigma+2\epsilon}$ by Eq. (36), faster than k^{-4} , so its integral converges absolutely and is the same in every regularization. Inserting Eq. (33) and integrating brane by brane with Eq. (30),²⁰

$$\begin{aligned} \delta\rho_\Pi &= 3 \int \frac{d^4 k}{(2\pi)^4} [R_\Pi(k^2) - R_\infty] \\ &= -\frac{3M^4 \mu_\psi}{16\pi^2} \int \frac{\vartheta(s)}{(sM^2)^2} (\Pi - \nu)(ds). \end{aligned} \quad (79)$$

²⁰ The exchange of the momentum integral and the compensated sum is justified by truncation. For the branes larger than η/M^2 the sum is finite and the exchange is trivial. As $\eta \rightarrow 0$ the right side of Eq. (79) converges in mean square by Eq. (10), and so does the left side, by Minkowski's inequality: the root-mean-square difference at momentum k is at most $[nV(\eta)]^{1/2}$ and, by Eq. (35), at most $C|k|^{-\sigma}$ for large k , which is integrable.

This is a compensated sum over the branes, with mean zero and, by Eq. (10), variance $(3M^4\mu_\psi/16\pi^2)^2 n \int_0^1 \vartheta(\tau)\tau^{-4}d\tau/\tau$. The integral converges when $V(t) = O(t^\sigma)$ with $\sigma > 4$, which (G) guarantees. For the power law (8) the variance is $(3M^4\mu_\psi/16\pi^2)^2 n\vartheta_0/(\zeta - 4)$, and relative to the mean (78) the standard deviation is

$$\frac{(\mathbb{E} \delta\rho_\Pi^2)^{1/2}}{\rho_1} = 2 \left[\frac{\vartheta_0}{n(\zeta - 4)} \right]^{1/2}, \quad (80)$$

independent of the response. Table II gives 0.37 for $n = 4$ and $\zeta = 9$. The zero-point energy of a realization is therefore $\rho_1 + \delta\rho_\Pi$, a finite number that is the same for every regulator, whose mean over realizations is Eq. (78). Each brane lowers it, by an amount proportional to $\vartheta(s)/s^2$; the positive mean is the dimensionally continued sum over the infinitely many small branes, $\int_0^1 \tau^{-2} d\tau/\tau = -\frac{1}{2}$ by continuation, which is how a sum of negative terms acquires the positive value (77).

The number in Eq. (77) is not a property of dimensional regularization. Cut the radial integral off at $w = W$, that is at $k = \Lambda$ with $W = \Lambda^4/M^4$. Integrating by parts with $\text{Ein}'(w) = (1 - e^{-w})/w$ and using Eq. (20),

$$\begin{aligned} \int_0^W \text{Ein}(w) dw &= W \text{Ein}(W) - W + 1 - e^{-W} \\ &= W \log W + (\gamma_E - 1)W + 1 + O(e^{-W}). \end{aligned}$$

With a smooth cutoff the same integral is

$$\begin{aligned} \int_0^\infty \text{Ein}(w) e^{-w/W} dw &= W \log(1 + W) \\ &= W \log W + 1 + O(W^{-1}), \end{aligned}$$

because $\int_0^\infty e^{-w/W} (1 - e^{-wt}) dw = W^2 t/(1 + Wt)$ and $\int_0^1 W^2 dt/(1 + Wt) = W \log(1 + W)$. The terms that grow with W , of order $\Lambda^4 \log(\Lambda/M)$ and Λ^4 , differ between the two cutoffs. The constant is 1 in both, the value of $\Gamma(s)/s$ at $s = 1$, and this is true for any cutoff function.²¹ The power-law terms are the scheme artifacts of Sec. VIA; the finite part is universal. Figure 11 shows both halves of the argument: the Mellin transform with its strip of convergence and its value at $s = 1$, and the remainder after the growing terms are removed, for a one-parameter family of cutoffs that ends in the sharp one.

²¹ Let r be smooth on $[0, \infty)$ and rapidly decreasing, with $r(0) = 1$, and let $\tilde{r}(s) = \int_0^\infty u^{s-1} r(u) du$. By the Mellin-Parseval formula, $\int_0^\infty \text{Ein}(w) r(w/W) dw = \frac{1}{2\pi i} \int_{\text{Re } s=s_0} \frac{\Gamma(s)}{s} \tilde{r}(1-s) W^{1-s} ds$ with $-1 < s_0 < 0$. Move the contour to the right. The double pole of $\Gamma(s)/s$ at $s = 0$ gives the terms $W \log W$ and W , whose coefficients involve $\tilde{r}(1)$ and $\tilde{r}'(1)$ and so depend on the cutoff. The simple pole of $\tilde{r}(1-s)$ at $s = 1$, where $\tilde{r}(1-s) \simeq r(0)/(1-s)$, gives the constant $\Gamma(1)r(0) = 1$ for every r . What remains decays as a negative power of W .

In general relativity the zero-point integrand is $\log k^2$ alone. Its integral is zero in dimensional regularization and quartic with a cutoff, and it has no finite part that is independent of the regulator. In the theory of the gas the form factor turns it into a definite number set by M , n and the realization. Newton's constant receives a finite shift of the same kind, proportional to M^2 , which we have not computed. The cosmological constant is then the sum of a bare term, which is not renormalized, and Λ_1 . Making the total small is still a choice of the bare term, but it is now a choice between two finite numbers rather than the subtraction of a divergence, and the theory does not explain why the total is small. Matter fields, which this paper does not include, would add zero-point energies of their own.

VII. UNITARITY

The spectrum of Sec. IIIC is a statement about the tree-level propagator. Loops raise a separate question, because $1/a$ is unbounded near the imaginary z axis, as Sec. IIE shows for the skeleton, and the loop energies cannot be Wick-rotated in the usual way. A theory with such a form factor has to be defined by a prescription. We use the one that goes back to Efimov [5] and was put in its modern form by Pius and Sen [16]. Loop integrals are defined with Euclidean loop momenta. Lorentzian amplitudes are the analytic continuations of the Euclidean ones in the external energies, and during the continuation the contours of the loop energies are deformed to avoid the poles of the propagators while their end points stay at $\pm i\infty$. Figure 12(a) shows why the end points matter: the propagator is small along the Euclidean contour and along the real axis away from its pole, and in the wedges between them it swings between very large and very small values.

In this prescription the physical S matrix is unitary at every order of perturbation theory. The proof has the two steps familiar from local gauge theories. First, the amplitudes obey the Cutkosky cutting rules with all ten components of the graviton and the ghosts on the cut lines, which is unitarity in the space of all asymptotic states, whose metric is indefinite (Sec. VIIA). Second, BRST invariance removes the states of negative and zero norm by the quartet mechanism of Kugo and Ojima (Sec. VIIB). Table VII lists every input and where it is established.

A. Cutting rules

Briscese and Modesto proved the Cutkosky rules for nonlocal theories whose form factor e^H has the following four properties [15]:

- (U1) H is an entire function;
- (U2) e^H has no zeros and no poles at finite momentum;




FIG. 11. The finite zero-point energy. (a) $\log_{10} |\Gamma(s)/s|$ over the complex s plane, the Mellin transform (76) of Ein. The integral converges in the strip $-1 < \text{Re } s < 0$ (orange band on the floor, which also carries the contours of the surface); its continuation has a double pole at $s = 0$ and simple poles at $s = -1, -2, \dots$ (black curve along the real axis), and is regular at $s = 1$, where it equals 1 (red). That value is the whole answer (77). (b) The integral $\int_0^\infty \text{Ein}(w) r(w/W) dw$ with the cutoffs $r(u) = e^{-u^p}$, after the terms that grow with W are subtracted, $\tilde{r}(1) W \log W + [\tilde{r}'(1) + \gamma_E \tilde{r}(1)] W$, with $\tilde{r}$ the Mellin transform of r . The surface is drawn over $\log_{10} W$ and $1/p \in [0, 1]$. The coefficients of the subtracted terms depend on p , the remainder does not: it tends to 1 (gray plane) for every p , and at $W = 10^3$ it lies within 10^{-3} of 1 across the whole family. The red edge $1/p = 0$ is the sharp cutoff, where the remainder is $1 - e^{-W} + WE_1(W)$.

- (U3) e^H grows without bound at the ends of the loop-energy contours and as the spatial loop momentum grows;
- (U4) H is asymptotically logarithmic in a cone around the real axis of the squared momentum.

Their proof is written out for a scalar field with a local kinetic term, obtained by a field redefinition, and vertices that carry a factor $e^{-H/2}$ for every leg. It follows the proof of Pius and Sen for string field theory [16], and Chin and Tomboulis reached the same cutting rule for general entire vertices by way of the Landau equations [45]. In all three the Feynman rules enter through three properties only:²²

- (P1) the propagators are those of a local theory;
- (P2) the vertices are entire functions of the momenta on their legs, with real Taylor coefficients in Euclidean variables;

- (P3) on the deformed contours the integrand of every graph, together with its counterterms, is absolutely integrable at the ends of the contours, uniformly for external momenta in compact sets.

We verify them for the gravitational theory of the gas.

(P1): *a field redefinition.* In the Feynman–de Donder gauge of Sec. IV A the quadratic operators of $h_{\mu\nu}$ and of the ghosts around flat space are $a(p^2)$ times those of general relativity, Eq. (55). Put

$$h_{\mu\nu} = a(-\square)^{-1/2} \tilde{h}_{\mu\nu}, \quad c^\mu = a(-\square)^{-1/2} \tilde{c}^\mu, \quad \bar{c}_\mu = a(-\square)^{-1/2} \tilde{\bar{c}}_\mu, \quad (81)$$

with $\square$ the flat d'Alembertian. Since $a = e^H$ with H entire, $a^{-1/2} = e^{-H/2}$ is entire as well, and the quadratic action of $\tilde{h}$ and the ghosts is that of general relativity in Feynman–de Donder gauge. That is (P1); the redefined propagators have a single pole, at $p^2 = 0$, with the residue of general relativity. The Jacobian of Eq. (81) does not depend on the fields and drops out. The redefinition does not change the S matrix. It is linear, and on shell it is the identity, since $a(0) = 1$, so the residues that enter the reduction formula are unchanged.²³

²² The Gaussian or exponential form of the vertices in Refs. [15, 16] is used to show that the integrals converge at the ends of the deformed contours, which is (P3). The rest of the argument, the continuation of the Euclidean amplitude into the physical region from the upper half plane of the external energies, the pinching of the contours at the poles of the propagators, and the identification of the anti-Hermitian part with the sum over cuts, uses only the location of the poles and the analyticity of the vertices.

²³ This is the equivalence theorem for field redefinitions in its simplest form: a linear redefinition $h = Z(-\square)\tilde{h}$ multiplies every amputated Green function by $Z(p_j^2)$ on each external leg, and at $p_j^2 = 0$ these factors are $a(0)^{-1/2} = 1$.

(a) $n = 4$, $|\mathbf{k}| = 1.2M$, $z = \mathbf{k}^2 - (k^0)^2$
 white: Euclidean contour; red: real axis;
 gray wedges: $|\hat{a}|^{\pm 1}$ oscillates beyond $10^{\pm 6}$





FIG. 12. Unitarity in the Efimov–Pius–Sen prescription. (a) The size of a propagator, $\log_{10} |z \hat{a}(z)|^{-1}$ for the skeleton, over the complex plane of the loop energy k^0 at fixed spatial momentum $|\mathbf{k}| = 1.2M$, for $n = 4$, with $z = \mathbf{k}^2 - (k^0)^2$. Along the Euclidean contour (white, $\text{Re } k^0 = 0$, drawn on the floor and on the surface) and along the real axis away from the poles at $k^0 = \pm|\mathbf{k}|$ (red, black dots) the propagator is small. In the wedges around $\arg k^0 = \pm\pi/4, \pm 3\pi/4$, where z is close to the imaginary axis, $\log |\hat{a}|$ has an oscillating part of size $\frac{n}{2}|E_1(z^2/M^4)|$, which grows like $e^{|\text{Re } z^2|/M^4}$, so the propagator swings between extremely large and extremely small values; this is why the loop energies cannot be rotated to the real axis. The surface is clipped at ± 6 and left out where that oscillating part exceeds 6 decades (gray on the floor). (b) The cutting rule of Sec. VII A: the anti-Hermitian part of an amplitude is the sum over cuts with all eighteen on-shell modes of the redefined fields. Between physical states, those in $\ker Q_B$, the contributions of the four BRST quartets cancel (Sec. VII B), and only the two helicities of the graviton remain.

(P2): *entire vertices*. After the redefinition every vertex is the vertex of Sec. IV C multiplied by $a^{-1/2}(\ell_j^2)$ for each quantum leg. By Sec. IV B the vertex of Sec. IV C is a finite sum of polynomials in the momenta times divided differences of the entire functions $\mathcal{G}$ and $a - cx^n$, and a divided difference of an entire function is an entire function of all its arguments, by Eq. (57). Products of entire functions are entire, and all Taylor coefficients are real because ψ is real on the real axis and the sizes and strengths of the branes are real. The counterterms of Sec. VI are local, with real coefficients. This is (P2).

(P3): *the ends of the contours*. Take a line that carries the loop momentum k plus a combination q of external momenta. Near the end of a loop-energy contour the Euclidean loop energy is real and large, $k_4 = \kappa \rightarrow \pm\infty$, while the continued external energies are complex but bounded. The squared Euclidean momentum of the line is then $z = (\kappa + q_4)^2 + |\mathbf{k} + \mathbf{q}|^2 = (\kappa^2 + \mathbf{k}^2)[1 + O(|q|/|\kappa|)]$, so $\arg z \rightarrow 0$, and the same holds when $|\mathbf{k}| \rightarrow \infty$ at fixed κ . In the language of Appendix C, all squared momenta of lines and of running momenta in vertex chains lie in a region $\Omega = \text{conv } K + S_\delta$, a compact set plus a cone around the positive axis. On Ω the cone asymptotics (37) give $|a(z)| \geq \frac{\epsilon}{2}|z/M^2|^n$ once $|z|$ is large, so the propagator obeys Eq. (56) with p^2 replaced by $|z|$, and Appendix C shows that the divided-difference bound (59), and with it the vertex bound (60), hold on Ω with $|\ell|$ replaced by $|\text{Re } \ell|$ plus a constant. Every factor of the integrand

therefore obeys on the contour the bound it obeys on the Euclidean section, and the power counting of Sec. IV D, applied to the subtracted integrands,²⁴ gives (P3). Properties (U3) and (U4) are the special case of this statement for a single propagator.

With (P1)–(P3) the argument of Refs. [15, 16] applies as it stands. For real external momenta the amplitude is the boundary value of an analytic function from the upper half plane of the external energies, and its anti-Hermitian part is the sum of the Cutkosky cuts, in which every cut propagator is replaced by $2\pi\delta(p^2)\theta(p^0)$ times its numerator. The numerators are those of general relativity in Feynman–de Donder gauge, so the cut lines carry all ten polarizations of $\tilde{h}_{\mu\nu}$ and the ghosts. Summed over all graphs, this is the statement $S^\dagger S = \mathbf{1}$ in the Fock space $\mathcal{F}$ of the free fields $\tilde{h}_{\mu\nu}$, $\tilde{c}^\mu$, $\tilde{\bar{c}}_\mu$, with its indefinite

²⁴ The integrals that diverge are the one-loop subgraphs, with $\bar{\omega} = 4$. Subtract from each its Taylor expansion to fourth order in its external momenta at a Euclidean point, as in the BPHZ scheme [42]. Derivatives with respect to external momenta lower the degree of the vertex and propagator bounds by one each, because the functions involved are symbols in the sense of Eqs. (40) and (59), so the subtracted subgraphs have negative degree and the subtracted integrand of every graph is absolutely convergent by Weinberg’s theorem. The subtraction differs from the counterterms of Sec. VI by local terms with real coefficients, which are vertices of the kind allowed by (P2).

inner product.

B. Physical states

The space $\mathcal{F}$ contains states of negative and zero norm, and the cuts of Sec. VII A run over all of them. They are removed by BRST symmetry, as in Einstein gravity. Kugo and Ojima proved [17, 46] that the physical S matrix is unitary with a positive inner product if

- (Q1) the gauge-fixed theory has a nilpotent BRST charge Q_B that commutes with the S matrix;
- (Q2) the asymptotic fields are free fields on which Q_B acts by the linearized BRST transformation, so that the unphysical modes form quartets;²⁵
- (Q3) the S matrix is unitary in the full space $\mathcal{F}$.

Under these conditions S maps $\ker Q_B$ into itself, and it is unitary on the quotient $\ker Q_B / \text{im } Q_B$, whose inner product is positive definite. For linearized gravity in de Donder gauge the eight unphysical polarizations of the graviton and the eight ghost modes form four quartets, one for each value of the vector index, and the quotient contains the two helicities of the graviton and nothing else [17].

Condition (Q3) is Sec. VII A. For (Q1), the action (46) is invariant under diffeomorphisms, being built from curvatures and the covariant d'Alembertian, and the gauge-fixing and ghost terms of Sec. IV A are together the BRST variation of a gauge-fixing fermion, $\bar{c}^\mu a(-\square)(\chi_\mu - \frac{1}{2}b_\mu)$ with a Nakanishi–Lautrup field b_μ that is integrated out. The weight depends on the background only and is inert under BRST, so the gauge-fixed action is BRST invariant and $Q_B^2 = 0$. The Slavnov–Taylor identities follow from that invariance and the invariance of the measure, as in local theories, and with them $[Q_B, S] = 0$ [17]; the counterterms of Sec. VI are gauge invariant or BRST exact and preserve this. For (Q2), the redefinition (81) leaves the linear part of the BRST transformation unchanged. Indeed $s h_{\mu\nu} = \partial_\mu c_\nu + \partial_\nu c_\mu + O(\text{fields}^2)$ becomes

$$s \tilde{h}_{\mu\nu} = a^{1/2}(\partial_\mu c_\nu + \partial_\nu c_\mu) + \dots = \partial_\mu \tilde{c}_\nu + \partial_\nu \tilde{c}_\mu + \dots,$$

because $a^{1/2}$ and $a^{-1/2}$ act on the same momentum. The free action is that of general relativity in Feynman–de Donder gauge (P1). So the asymptotic fields and the asymptotic BRST charge are exactly those of linearized

TABLE VII. Inputs of the unitarity argument and where each is established. The last row is the only assumption, and it is the one made for Einstein gravity.

input	content	established in
(P1)	local propagators after $h = a^{-1/2}\tilde{h}$	Eq. (81), a zero-free
(P2)	vertices entire, real coefficients	Secs. IV B, IV C
(P3)	integrable at the ends of the contours	Eq. (37), App. C, Sec. IV D
(Q1)	BRST invariance, $[Q_B, S] = 0$	Sec. IV A, Slavnov–Taylor
(Q2)	free asymptotic fields, four quartets	$a(0) = 1$, Eq. (81)
(Q3)	$S^\dagger S = \mathbf{1}$ on $\mathcal{F}$	cutting rules, Sec. VII A
—	asymptotic states free; soft gravitons treated inclusively	assumed, as for general relativity [17, 18]

general relativity, and the quartets are the ones found by Kugo and Ojima. There are no further asymptotic states, because the redefined propagators have no other poles, which is the content of (U2).²⁶

Therefore the physical S matrix of the gas is unitary at every order of perturbation theory in κ , in the Efimov–Pius–Sen prescription. What is assumed beyond the derivations above is what Kugo and Ojima assume for Einstein gravity: that the asymptotic states are described by the free fields. For massless gravitons this holds only after the soft divergences have been dealt with, by an infrared regulator or by passing to inclusive rates [18], exactly as in general relativity. The prescription has a price that local theories do not pay: it is acausal on scales of order $1/M$, where the time order of production and decay vertices can be reversed [47]. With M near the Planck mass this is far from observable.

C. The size of amplitudes

Unitarity in this sense constrains the cuts of amplitudes, not their size. For actions quadratic in the Ricci tensor and the scalar curvature, with form factors inserted, the on-shell tree amplitudes coincide with those of general relativity for any number of gravitons [48], and $R + G_{\mu\nu}F(\square)R^{\mu\nu}$ is of that type. The quartic operators are different. Those built from R and $R_{\mu\nu}$ add nothing to on-shell graviton scattering, because the linearized

²⁵ A quartet is a pair of BRST doublets: a mode χ with $Q_B\chi = \gamma$ and an antighost mode $\tilde{\gamma}$ with $Q_B\tilde{\gamma} = \beta$, where γ and $\tilde{\gamma}$ have ghost number ± 1 and the only nonzero inner products are $\langle\chi|\beta\rangle$ and $\langle\gamma|\tilde{\gamma}\rangle$. Every state built with at least one quartet mode is either outside $\ker Q_B$ or in $\text{im } Q_B$, or has zero norm, and the quotient $\ker Q_B / \text{im } Q_B$ is spanned by states without quartet modes [46].

²⁶ This is a statement about perturbation theory. Shapiro has pointed out that the resummed propagator $1/(p^2a + \Sigma)$ of a nonlocal theory may acquire complex-conjugate poles from the self-energy Σ [4]. At any finite order of perturbation theory the self-energy enters as insertions on lines with the bare propagator, and no such pole appears; what the resummed theory does is a nonperturbative question.

Ricci tensor of an on-shell graviton vanishes. The Riemann operator does add something: a contact term of order $G|s_3|E^{2n+2}/M^{2n}$ at center-of-mass energy E , which grows faster than any amplitude of general relativity and reaches order one at $E_* = (M^{2n}/G|s_3|)^{1/(2n+2)}$. That scale can be avoided. Set $s_3 = 0$ and solve for s_1 and s_2 alone, with the upper-left 2×2 block of Eq. (73), whose determinant is $-12/c^2$. The Gauss–Bonnet pole then stays, but $\int \sqrt{g} E$ is topological in four dimensions and does not enter the S matrix.

With $s_3 = 0$ every tree amplitude of gravitons is that of general relativity, and at fixed angle it grows like GE^2 . It reaches order one near the Planck energy $E_P = G^{-1/2}$, and there the loop expansion, finite term by term, stops being an expansion in a small number. General relativity becomes strongly coupled at the same scale, and the form factor does not move it.

Matter that couples minimally to the metric behaves differently, because its gravitational scattering at tree level goes through the propagator (53) itself. Between conserved sources the propagator gives $T_1 \cdot T_2 - \frac{1}{2}T_1T_2$ divided by $p^2a(p^2)$, and for two distinct massless scalars $T_1 \cdot T_2 - \frac{1}{2}T_1T_2 = -su$ in terms of the Mandelstam variables. The one-graviton exchange is therefore the amplitude of general relativity divided by the form factor at the momentum transfer,

$$|A(s, t)| = \frac{8\pi G |su|}{|t|a(-t)}, \quad (82)$$

with a evaluated at the Euclidean point $p^2 = -t > 0$. At a fixed scattering angle θ in the center-of-mass frame, $-t = s \sin^2(\theta/2)$ and $-u = s \cos^2(\theta/2)$, and the lower half of the two-sided bound (38) gives

$$|A| \leq 8\pi GM^2 C_\Pi \frac{\cos^2(\theta/2)}{\sin^4(\theta/2)} \frac{x}{(1+x^2)^{n/2}}, \quad x = \frac{-t}{M^2},$$

with $C_\Pi = e^{-nm_\psi + \|R_\Pi\|}$. The last factor is at most $(n-1)^{(n-1)/2}/n^{n/2}$, which is 0.325 for $n = 4$, and it falls like x^{1-n} . At every fixed angle the tree amplitude for the gravitational scattering of matter therefore rises like GE^2 only up to $E \sim M$, then falls like $GM^2(M/E)^{2n-2}$, and never exceeds a number of order $GM^2 = (M/M_P)^2$. For M below the Planck mass, matter does not scatter strongly through one graviton at any energy. This is a tree-level statement about matter added to the theory; the finiteness proof of Sec. VI concerns pure gravity, and loops of matter are not treated here. The forward peak at $|t| \rightarrow 0$ is the Newtonian one of general relativity, an infrared effect untouched by the form factor.

D. In what sense the theory is ultraviolet complete

A quantum field theory is ultraviolet complete in perturbation theory when its loop expansion is defined at every order without a cutoff and without new degrees of freedom: each term is finite with fixed bare couplings, the

S matrix is unitary order by order, and the asymptotic states are the same at every energy. For $n \geq 4$, almost every realization of a gas obeying A1 and A2, and the assumption on asymptotic states that is made for general relativity, Secs. VI and VII establish all three. The bare couplings do not depend on the regulator (Sec. VIA), the physical S matrix is unitary at every loop order, and its only asymptotic states are the two helicities of the graviton. No physical coupling is renormalized, so none of them runs. The perturbative renormalization group of the physical couplings is trivial, there is no Landau pole, and no order of perturbation theory calls for a new coupling, whereas in general relativity the two-loop divergence does [2]. In this sense, and to every order in κ , the theory of the gas is ultraviolet complete.

The statement concerns the loop expansion, and Sec. VII C shows where that expansion stops being useful for gravitons. A definition of the theory that does not rest on the expansion, by a Euclidean functional integral or by a proof that the series is Borel summable, would make it ultraviolet complete without qualification. We do not have one, and two obstacles are concrete. The quadratic action (52) is that of general relativity multiplied by $a(p^2)$, which is positive on the Euclidean axis, so its spin-zero part, with coefficient -2 , is unbounded below exactly as the conformal factor of general relativity is [49]; the gas does not cure this. The resummed propagator may also acquire complex poles from the self-energy [4], which no finite order sees. Neither obstacle is special to the gas. Both are open for every nonlocal theory of this class, and no nontrivial relativistic quantum field theory in four dimensions is yet known to satisfy the Wightman axioms [50]. Classically, the linearized field of any mass below $m_c \sim M_P^2/M$ is weak everywhere and has no horizon (Sec. VIII). Collisions at higher energies, where general relativity forms black holes, are a nonlinear problem that this paper leaves open.

VIII. NEWTONIAN LIMIT

The quartic operators do not touch the propagator, so the static field of a point mass follows from Eq. (53) alone. For a source $T_{\mu\nu} = m \delta^3(\mathbf{x}) u_\mu u_\nu$ at rest, the linearized field is the one of general relativity with $1/\mathbf{k}^2$ replaced by $1/(\mathbf{k}^2 a(\mathbf{k}^2))$, because the form factor multiplies both projectors by the same scalar. In general relativity the computation gives $\tilde{\Phi}_N(\mathbf{k}) = -4\pi Gm/\mathbf{k}^2$, so here

$$\tilde{\Phi}_N(\mathbf{k}) = -\frac{4\pi Gm}{\mathbf{k}^2 a(\mathbf{k}^2)}, \quad (83)$$

and the Fourier transform of a radial function in three dimensions,

$$\int \frac{d^3k}{(2\pi)^3} e^{i\mathbf{k}\cdot\mathbf{x}} f(k) = \frac{1}{2\pi^2 r} \int_0^\infty k \sin(kr) f(k) dk,$$

TABLE VIII. Bookkeeping of divergences. “Removed” means that the corresponding pole or singularity is absent for $n \geq 4$ with the tuning (74); the third column gives the reason.

divergence	fate	reason
one loop, $\int \sqrt{g}$ and $\int \sqrt{g} R$ (running of Λ , G)	absent for $n \geq 4$	one-loop poles are those of a local theory with no dimensionful coupling; saturation (S3) and fine graining (G), Sec. VC; the finite vacuum energy has mean $3nM^4/(64\pi^2)$, Sec. VIB
one loop, R^2 , $R_{\mu\nu}R^{\mu\nu}$, E	removed	three quartic operators, triangular matrix with determinant $-24/c^3$, Eqs. (73), (74)
two or more loops, overall	absent for $n \geq 4$	vertex bound and Weinberg’s theorem, Sec. IVD
one-loop subgraphs, gauge-invariant part	removed	equals the one-loop pole of the full metric, which vanishes after tuning
one-loop subgraphs, BRST-exact part	present off shell	renormalizes the gauge-fixing sector; no effect on the S matrix, Sec. VI
power-law terms (Λ^4 , Λ^2 with a cutoff)	scheme artifacts	absent in any analytic regularization; the finite part does not depend on the regulator, Sec. VIA
soft gravitons on shell (infrared)	physical	as in general relativity; cancel in inclusive rates [18]
static potential at $r \rightarrow 0$ (classical)	removed	$\Phi_N(0)$ finite; Gamma-function bounds for the skeleton, Sec. VIII
extra poles of the propagator	absent	almost every realization entire and zero-free, by (S5); resonant branes would give complex pairs, Eq. (4)

gives

$$\Phi_N(r) = -\frac{2Gm}{\pi r} \int_0^\infty \frac{\sin kr}{k a(k^2)} dk. \quad (84)$$

With $a = 1$ the integral is $\pi/2$ and Eq. (84) is Newton’s potential. With the form factor of the gas, the integrand is bounded by $1/a(k^2)$ uniformly in r , and $1/a$ falls like k^{-2n} . Below we find how deep the well is, how fast the potential approaches $-Gm/r$, and how large a mass the linearized description can carry; all three have closed answers.

A. Depth of the well

At the origin $\sin(kr)/(kr) \rightarrow 1$, and

$$\Phi_N(0) = -\frac{2Gm}{\pi} I_n, \quad I_n = \int_0^\infty \frac{dk}{a(k^2)}. \quad (85)$$

Since $|\sin(kr)/(kr)| \leq 1$ and the integrand of I_n is positive, $|\Phi_N(r)| \leq |\Phi_N(0)|$ for every r : the potential is deepest at the center, and it is finite there. The value of I_n is bracketed from both sides by elementary integrals, and the bracket is narrow. We give the numbers for the skeleton, $a = \hat{a}$. A realization has $a = \hat{a} e^{R_\Pi}$ with R_Π bounded on the real axis, so its I_n lies between $e^{-\sup R_\Pi}$ and $e^{-\inf R_\Pi}$ times the skeleton value, the supremum and infimum being taken over the real axis; the qualitative statements below, a finite well, $2n - 2$ continuous derivatives and an approach to Newton’s law faster than any power, hold for almost every realization as they stand, because they use only Eq. (38), the derivative bounds that follow Eq. (37), which take the place of Eq. (23), and $a - 1 = O(k^4)$; only the constants change from one realization to another.

For an upper bound we use the left inequality of Eq. (21), $a \geq (1 + k^4/M^4)^{n/2}$. With $t = k^4/M^4$,

$$I_n \leq \int_0^\infty \frac{dk}{(1 + k^4/M^4)^{n/2}} = \frac{M}{4} \int_0^\infty \frac{t^{-3/4} dt}{(1 + t)^{n/2}} = J_n, \quad (86)$$

where

$$J_n = \frac{M}{4} B\left(\frac{1}{4}, \frac{n}{2} - \frac{1}{4}\right) = \frac{M}{4} \frac{\Gamma(\frac{1}{4})\Gamma(\frac{n}{2} - \frac{1}{4})}{\Gamma(\frac{n}{2})}. \quad (87)$$

For even $n = 2m$ the gamma functions collapse. From $\Gamma(m - \frac{1}{4}) = \Gamma(\frac{3}{4}) \prod_{j=1}^{m-1} (j - \frac{1}{4})$ and the reflection formula $\Gamma(\frac{1}{4})\Gamma(\frac{3}{4}) = \pi\sqrt{2}$,

$$J_{2m} = \frac{\pi\sqrt{2}}{4} M \prod_{j=1}^{m-1} \left(1 - \frac{1}{4j}\right), \quad (88)$$

so $J_4 = \frac{3\pi\sqrt{2}}{16} M$, $J_6 = \frac{21\pi\sqrt{2}}{128} M$, $J_8 = \frac{77\pi\sqrt{2}}{512} M$, and so on.

For a lower bound the right inequality of Eq. (21) is too weak, since $\hat{c} = e^{n\gamma_E/2}$ grows exponentially with n . A better one comes from the definition of Ein itself. The integrand $(1 - e^{-t})/t$ is at most 1, so $\text{Ein}(w) \leq w$ for $w \geq 0$, and therefore $a(k^2) \leq \exp(\frac{n}{2} k^4/M^4)$. Then

$$I_n \geq \int_0^\infty e^{-nk^4/(2M^4)} dk = \frac{M}{4} \Gamma\left(\frac{1}{4}\right) \left(\frac{2}{n}\right)^{1/4} \equiv K_n. \quad (89)$$

The two bounds are close. Their ratio is $J_n/K_n = (n/2)^{1/4} \Gamma(\frac{n}{2} - \frac{1}{4})/\Gamma(\frac{n}{2})$, and Wendel’s inequality for ratios of gamma functions [51],²⁷ $\Gamma(x + s) \geq x \Gamma(x) (x +$

²⁷ For $x > 0$ and $0 < s < 1$, Hölder’s inequality with exponents $1/s$

$s)^{s-1}$ with $x = \frac{n}{2} - \frac{1}{4}$, $s = \frac{1}{4}$, gives $J_n/K_n \leq 2n/(2n-1)$. Hence

$$\Phi_N(0) = -\frac{\Gamma(\frac{1}{4})}{2\pi} \left(\frac{2}{n}\right)^{1/4} GmM \left[1 + \frac{\theta_n}{2n-1}\right], \quad (90)$$

$$0 \leq \theta_n \leq 1.$$

For $n = 4$, Eqs. (86) and (89) give $0.4852 \leq |\Phi_N(0)|/(GmM) \leq 0.5303$, and the value of the integral itself, obtained by quadrature as a check, is 0.5100; for $n = 6$ the bracket is $[0.4385, 0.4640]$ around 0.4519. For comparison, the Gaussian form factor e^{p^2/M^2} gives $\Phi_N = -Gm \operatorname{erf}(Mr/2)/r$ and $|\Phi_N(0)| = GmM/\sqrt{\pi} \approx 0.564 GmM$ [10, 52]. The well of the gas is about ten percent shallower for $n = 4$, and it becomes slowly shallower as the gas becomes denser, as $n^{-1/4}$.

The smoothness of Φ_N is fixed by the decay of Eq. (83). Since $\tilde{\Phi}_N = O(k^{-2n-2})$, the integral $\int d^3k |\mathbf{k}|^j |\tilde{\Phi}_N|$ converges for $j < 2n - 1$, and Φ_N has $2n - 2$ continuous derivatives everywhere, the origin included. Equivalently, $\nabla^2 \Phi_N = 4\pi G \rho_{\text{eff}}$ with an effective source

$$\rho_{\text{eff}}(\mathbf{x}) = m \int \frac{d^3k}{(2\pi)^3} \frac{e^{i\mathbf{k}\cdot\mathbf{x}}}{a(\mathbf{k}^2)}, \quad \int \rho_{\text{eff}} d^3x = m, \quad (91)$$

the total being m because $a(0) = 1$. The point mass is spread over a region of size $1/M$. The same two estimates of a give

$$\frac{\Gamma(\frac{3}{4})}{8\pi^2} \left(\frac{2}{n}\right)^{3/4} \leq \frac{\rho_{\text{eff}}(0)}{mM^3} \leq \frac{1}{8\pi^2} B\left(\frac{3}{4}, \frac{n}{2} - \frac{3}{4}\right),$$

and the linearized tidal tensor at the center, $\partial_i \partial_j \Phi_N(0) = \frac{4\pi G}{3} \rho_{\text{eff}}(0) \delta_{ij}$, is finite. The curvature singularity of the linearized Schwarzschild field is gone.

B. Approach to Newton's law

Using $\int_0^\infty \sin(kr) dk/k = \pi/2$, write

$$\Phi_N(r) = -\frac{Gm}{r} [1 - \delta(r)], \quad (92)$$

$$\delta(r) = \frac{2}{\pi} \int_0^\infty \sin(kr) \chi(k) dk,$$

with $\chi(k) = [1 - 1/a(k^2)]/k$. Because $a(k^2)$ is an even entire function of k and $1 - 1/a = O(k^4)$ (for the skeleton $\frac{n}{2}k^4/M^4 + O(k^8)$ by Eq. (22)), χ is odd, smooth, and $O(k^3)$ at the origin. At large k , Eq. (23)

and the chain rule give $\partial_k^j [1/a(k^2)] = O(k^{-2n-j})$, hence $\chi^{(j)}(k) = O(k^{-1-j})$. Now integrate by parts twice:

$$\int_0^\infty \sin(kr) \chi(k) dk = -\frac{1}{r^2} \int_0^\infty \sin(kr) \chi''(k) dk.$$

The boundary terms at infinity vanish because χ and χ' do, and those at $k = 0$ vanish because $\chi(0) = 0$ and $\sin 0 = 0$. The function χ'' is again odd and smooth, so the step can be repeated, and after N repetitions

$$|\delta(r)| \leq \frac{2}{\pi} \frac{\|\chi^{(2N)}\|_{L^1}}{r^{2N}} = \frac{C_N}{(Mr)^{2N}}, \quad (93)$$

where C_N depends on n and N only, since $\chi(k) = M^{-1} \hat{\chi}(k/M)$ with $\hat{\chi}$ independent of M ; for a realization C_N depends also on the sizes sM^2 of its branes. As N is arbitrary, the correction to Newton's law falls faster than any power of $1/(Mr)$. Figure 13 shows the potential for $n = 4$ and the deviation for $n = 4, \dots, 8$; the oscillations of $\delta(r)$ come from the oscillating Fourier kernel and do not affect the bound.

This is the classical counterpart of the infrared statement of Sec. VI. Every term of the low-momentum expansion of $1/a$ is a polynomial in k^4 , and its Fourier transform is a contact term supported at $r = 0$, so no inverse power of r can appear at any order. The long-range quantum corrections of the effective field theory of gravity, which come from loops of massless particles and are non-analytic in k [53], are not affected, because the infrared of the theory is that of general relativity. The inverse-square law has been tested down to $52 \mu\text{m}$ [54], and by Eq. (93) these tests require only $M \cdot 52 \mu\text{m} \gg 1$, that is M well above 4×10^{-3} eV.

C. Mass scale of the linear regime

The linearized description is self-consistent at every point when $2|\Phi_N| < 1$ everywhere, which by Eq. (85) and $|\Phi_N(r)| \leq |\Phi_N(0)|$ happens exactly when

$$m < m_c = \frac{\pi}{4G I_n}, \quad \frac{\pi}{4G J_n} \leq m_c \leq \frac{\pi}{4G K_n}. \quad (94)$$

In units with $\hbar = c = 1$ and $G = M_{\text{P}}^{-2}$, $m_c = \mu_n M_{\text{P}}^2/M$ with μ_n of order one; Table IX gives it for several n . Below m_c the weak-field expansion holds down to $r = 0$ and the linearized field has no horizon. Above it, the expansion fails near the center and the question becomes nonlinear, which is beyond this paper. For M near the Planck mass, m_c is itself of order the Planck mass, so this is a statement about the smallest objects, not about astrophysical black holes.

IX. DISCUSSION

Table X sets the result beside the better-known alternatives, for the quantities that decide the ultraviolet

and $1/(1-s)$ applied to $\Gamma(x+s) = \int_0^\infty t^{s+x} t^{(1-s)(x-1)} e^{-t} dt$ gives $\Gamma(x+s) \leq \Gamma(x+1)^s \Gamma(x)^{1-s} = x^s \Gamma(x)$. The same inequality with $x \rightarrow x+s$ and $s \rightarrow 1-s$ gives $\Gamma(x+1) \leq (x+s)^{1-s} \Gamma(x+s)$, that is $\Gamma(x+s) \geq x \Gamma(x)/(x+s)^{1-s}$. Together, $(x/(x+s))^{1-s} \leq \Gamma(x+s)/(x^s \Gamma(x)) \leq 1$. We use the lower half with $x = \frac{n}{2} - \frac{1}{4}$ and $s = \frac{1}{4}$.




FIG. 13. Newtonian limit. (a) The potential (84) of a point mass for the skeleton with $n = 4$, in units of GmM , drawn over a plane through the source with one quarter cut away. The surface and the solid profiles on the cut faces are Φ_N ; the mesh and the dashed profiles are Newton's $-Gm/r$. The red bar on the axis is the bracket $0.4852 \leq |\Phi_N(0)|/GmM \leq 0.5303$ of Eqs. (86) and (89); the floor carries the contours of Φ_N . (b) The relative deviation $|\delta(r)|$ of Eq. (92) for $n = 4, \dots, 8$, on logarithmic axes. The dips are zeros of δ , which oscillates because of the Fourier kernel. The envelope falls faster than the reference lines $(Mr)^{-4}$ (dotted) and $(Mr)^{-8}$ (dash-dot) drawn at the back, as Eq. (93) requires. The curves are numerical evaluations of the closed-form integrals and serve as illustration only.

TABLE IX. Constants of the Newtonian limit, in units $M = 1$ (for J_n , K_n , I_n) and M_P^2/M (for m_c). K_n and J_n are the closed-form bounds (89) and (87); the column I_n is a quadrature of Eq. (85), given as a check. The bracket for m_c is Eq. (94), and $\hat{c} = e^{n\gamma_E/2}$. All entries are for the skeleton $\hat{a}$.

n	$\hat{c}$	K_n	I_n	J_n	$ \Phi_N(0) /GmM$	m_c
4	3.172	0.7622	0.8011	0.8330	0.5100	[0.943, 1.030]
5	4.234	0.7208	0.7485	0.7725	0.4765	[1.017, 1.090]
6	5.650	0.6887	0.7099	0.7289	0.4519	[1.077, 1.140]
7	7.540	0.6627	0.6797	0.6953	0.4327	[1.130, 1.185]
8	10.06	0.6409	0.6550	0.6682	0.4170	[1.175, 1.225]
10	17.92	0.6061	0.6165	0.6264	0.3925	[1.254, 1.296]

behavior: the form factor, the loop orders at which divergences occur, and the extra poles of the propagator. The last column records where the form factor comes from, which is the point of this paper.

The argument has two layers, which could fail independently. The first is about the gas. A product of independent factors, each of the exponential form that (S5) forces, is the exponential of a sum over the branes, and for a Poisson gas that sum splits into a deterministic integral and a compensated fluctuation. If the gas has no preferred scale, the integral grows like a logarithm, so the form factor grows like a power, and the exponent counts the strength per e -fold. The fluctuation is where a single realization could differ from its skeleton. It is a sum of independent terms with zero mean, and at large mo-

mentum it settles to a random constant when the small branes are weak enough, fast enough for the field theory under condition (G); in a gas of identical branes it never settles and performs a random walk in $\log p$. This is how an entire function with polynomial growth on the real axis and no zeros arises without being chosen, and it ties the growth of a to a density rather than to a parameter of the action. Nothing is averaged: the graviton propagates in one frozen gas, and the statements hold for almost every gas. The second layer is field theory. Once the form factor has the properties listed in Sec. IIH, the vertex bound and power counting confine the divergences to one loop, saturation and (G) keep the dimensionful couplings from running, three operators cancel the dimensionless one-loop poles, and the cutting rules with the quartet mechanism make the physical S matrix unitary. None of the field-theory steps uses the gas, and they apply to any mechanism that produces a form factor with those properties.

Under A1 and A2 with $n \geq 4$, and with the tuning (74), almost every realization of the gas gives a theory with no ultraviolet divergence at any order of perturbation theory around flat space. No physical coupling needs a counterterm, the S matrix has no ultraviolet pole, and the one-loop background-field effective action has none either. The bare couplings are finite numbers, the same for every regulator (Sec. VIA), and quantities that have no pole can be computed outright. The zero-point energy of the graviton is $\rho_1 + \delta\rho_\Pi$, whose mean over realizations is $3nM^4/(64\pi^2)$ for ψ_* and Eq. (78) in general, and whose

TABLE X. Ultraviolet behavior and pole content of the graviton propagator $1/(p^2 a)$ in four dimensions. For a polynomial form factor of degree N the residues of the extra poles add up to -1 , so at least one is a ghost [4]. ^aRenormalizable, with divergences at every order. ^bVertices grow exponentially and power counting is not available; see Ref. [14]. ^cAfter adding operators cubic or quartic in curvature [12]. ^dFor almost every realization of a gas obeying (G), with the one-loop tuning (74); no coupling is renormalized at any order, and the physical S matrix is unitary at every loop order in the Euclidean prescription (Sec. VII).

theory	form factor $a(p^2)$	divergent loop orders	extra poles	origin of a
Einstein [1, 2]	1	$L \geq 1$ off shell, $L \geq 2$ on shell	none	—
Stelle [3]	$1 + p^2/M^2$	all ^a	one massive ghost	local action
local, degree $N \geq 2$ [37]	polynomial	$L \leq (N + 1)/(N - 1)$	N	local action
Gaussian [9, 10]	e^{p^2/M^2}	open ^b	none	chosen
Tomboulis, Modesto [8, 11]	$e^{H(p^2)}$, polynomial growth	one loop; none ^c	none	chosen
this work, $n \geq 4$	$\exp[\frac{n}{2} \text{Ein}(p^4/M^4) + R_{\Pi}(p^2)]$	none ^d	none	one frozen Poisson gas of branes, Sec. II

relative fluctuation is Eq. (80). The static potential of a point mass is finite at the origin in almost every realization, and for the skeleton its depth is known in closed form to within $1/(2n - 1)$, Eq. (90). The renormalizations that remain act only on the gauge-fixing sector and have no physical effect. The physical S matrix is unitary at every loop order. Infrared divergences remain, as they must for a massless graviton, and they cancel in inclusive rates exactly as in general relativity [18]. Table VIII is the full list of divergences, and Table XI separates what is proved from what is assumed.

The following points are not settled here.

The two premises. Assumption A1 is a statement about a single brane. The composition law (2) justifies the product structure for weakly reflecting branes, and Sec. IIF shows that the field theory needs only the five conditions (S1)–(S5), but the response itself is not derived from a brane action. Condition (S5) is the one with the sharpest consequence. A brane with a resonance, such as the factor $2 - e^{-s^2 z^2}$, puts complex zeros into almost every realization, Eq. (4), and a theory built on such a gas has complex-conjugate pairs of poles that need a Lee–Wick or fakeon prescription [22–24]; it is not covered by Secs. IIIC–VII. Assumption A2 asks for independence, for a scale-invariant strength, and for (G), which says that the strength of a brane of size s falls faster than $(sM^2)^{2n}$ as $s \rightarrow 0$ in the sense of Eq. (7). We have given no microscopic reason for (G). Its role is clear, however: it is what makes the fluctuation of the gas at momentum p fall faster than p^{-2n} , and without it the local split of Sec. VA does not exist. Deriving the graviton effective action from a microscopic brane theory [55, 56] would test A1 and A2 and would fix ψ , $\vartheta(s)$ and n .

Dependence on the realization. Finiteness, the spectrum and unitarity hold for almost every realization, but several numbers are random: the leading coefficient $c = \hat{c} e^{R_\infty}$, and with it the tuning (74), which is fixed once the gas is given; the zero-point energy; and the depth of the Newtonian well. Their distributions are known in

closed form (Sec. IIG and Table II), and for the power law (8) with $\zeta = 2n + 1$ they are narrow. A universe with a frozen gas is one realization, and its constants are those of that realization.

Unitarity and strong coupling. Section VII proves that the physical S matrix is unitary at every loop order in the Efimov–Pius–Sen prescription, under the assumption about asymptotic states that is made for Einstein gravity. The prescription brings acausality on the scale $1/M$ [47]. Unitarity bounds the cuts of amplitudes, not their size. Graviton scattering is that of general relativity at tree level and becomes strong near the Planck energy, or already at E_* if the Riemann operator is kept, while the scattering of matter through one graviton stays of order GM^2 (Sec. VIIC).

The one-loop coefficients. The tuning (74) exists for every n because the matrix (73) is invertible, and it is given in closed form in terms of $\beta_1^{(0)}$, $\beta_2^{(0)}$ and $\beta_E^{(0)}$. Those three numbers are the one-loop poles of the local theory of Sec. VA, which is a higher-derivative theory of the type studied in Ref. [37]. We have not computed them, and the values of s_1 , s_2 , s_3 are therefore known as functions of them, not as numbers. For super-renormalizable theories of this class these beta functions are also still to be calculated in the literature [40].

The range of n . The threshold $n \geq 4$ is sharp for power counting. At $n = 3$ the two-loop degree $\bar{\omega}(2)$ is zero, and power counting alone does not exclude a logarithmic divergence at two loops; whether it is present would need the two-loop poles themselves.

Beyond perturbation theory. Everything here is perturbative in κ around flat space, and in that sense the theory is ultraviolet complete (Sec. VIID). A nonperturbative definition is not given. The Euclidean conformal mode is unbounded below, as in general relativity, the resummed propagator may acquire complex poles from the self-energy [4], and nothing is claimed about the nonlinear regime, about compact objects heavier than m_c of Eq. (94), or about the value of M .

TABLE XI. Status of the claims. “Proved” means derived in the text from the stated inputs, with the cited theorems; “premise” means a statement about branes that the paper does not derive. Statements about a realization hold for almost every realization of the gas.

statement	status	Sec.
response of one brane, A1	premise; (S1)–(S5), any admissible ψ	II A , II F
statistics of the gas, A2	premise; Poisson, Eq. (6), (G)	II B
form factor of a realization	derived from A1, A2: entire, zero-free, $a \simeq c x^n$	II C – II G
gas of identical branes	fails (G); no limit of a/x^n	II G
one pole, residue of GR	proved	III C
vertex bound	proved, every n	IV C , C
$L \geq 2$ graphs converge	proved, $n \geq 4$; $n = 3$ marginal	IV D
G , Λ do not run	proved	V C
one-loop poles cancel	proved; $\beta_i^{(0)}$ not computed	V D
no physical counterterm	proved, all orders	VI
graviton zero-point energy	computed, regulator independent; mean and variance	VI B
cutting rules	proved from (P1)–(P3)	VII A
unitarity on physical states	proved; asymptotic states as in GR	VII B
Newtonian potential	finite for almost every realization; closed-form bounds for the skeleton	VIII
soft gravitons	physical, as in GR	VI
ultraviolet complete in perturbation theory	proved: finite, unitary, no running, no new states	VII D
matter scattering, tree level	bounded by $O(GM^2)$ at fixed angle	VII C
gravitons above M_P ; nonperturbative definition	open; strong coupling as in GR	VII C , VII D

To sum up: a scale-invariant Poisson gas of branes that have no resonances and whose small members are weak enough, condition (G), gives in almost every frozen realization a form factor that is entire, has no zeros and grows like cp^{2n} with a remainder that falls faster than the field theory needs. That form factor removes every extra pole from the graviton propagator. For $n \geq 4$ the theory built on it has no ultraviolet divergence at any order once three numbers are fixed at one loop, and those numbers are finite. Newton’s constant and the cosmological constant are not renormalized, the zero-point energy of the graviton is a finite number that does not depend on the regulator, the physical S matrix is unitary at every

loop order, and the potential of a point mass is finite at the origin. Within perturbation theory the construction is therefore ultraviolet complete. Whether it stays complete beyond perturbation theory is the main question it leaves open.

DATA AVAILABILITY

All results in this paper are derived analytically in the text, and no argument depends on a numerical computation. Deep Bhattacharjee carried out all computations and checks, in C, Python, Julia and Lean 4, with the assistance of Claude Code. The code consists of three independent programs that share no code, a Python script, `verify.py` (178 checks), a C99 program, `formfactor_check.c` (80 checks), and a Julia program, `verify.jl` (96 checks, in 256-bit arithmetic with its own special functions and quadrature), which re-evaluate the identities, inequalities, distributions and coefficients of Secs. [II](#)–[VIII](#) and of Appendices [B](#) and [C](#) at sample points; a Lean 4 file, `PowerCounting.lean`, in which the integer statements behind the power counting and the one-loop tuning are proved for all values of their arguments and checked by the Lean kernel; and a Python script, `make_figures.py`, which draws the plotted panels of Figs. [3](#), [4](#), [5](#), [6](#), [7](#), [8](#), [9](#), [11](#), [12](#) and [13](#). The schematic diagrams are drawn in the L^AT_EX source with TikZ. Appendix [D](#) lists what is checked. The code accompanies this paper as Supplemental Material and is deposited, together with the code of the earlier study, in the archive of Ref. [\[13\]](#).

USE OF AI TOOLS

AI tools assisted with drafting and editing the text, with algebraic and numerical calculations, and with writing the verification code and the scripts that draw the figures. The author directed this work, checked its results, and takes full responsibility for the content of the paper.

FUNDING

The author received no funding for this work.

COMPETING INTERESTS

The author declares no competing interests.

Appendix A: Conventions and notation

Spacetime has dimension D , set to 4 except in dimensional regularization, where $D = 4 - 2\epsilon$. Momentum-

TABLE XII. Notation.

symbol	meaning
M	inverse of the largest brane size
n	strength per e -fold of size, Eq. (6)
s, τ	brane size (squared length); $\tau = sM^2 \in (0, 1]$
$\vartheta, \vartheta_0, \zeta$	strength of a brane; power law $\vartheta_0 \tau^\zeta$
$V(t), \sigma$	fine graining, Eq. (7) and condition (G)
Π, ν	a realization of the gas (Poisson process) and its intensity
ψ, ψ_*	exponent of one brane; $\psi_*(w) = 1 - e^{-w^2}$
θ	opening of the sectors in (S4)
z, x	$z = p^2$ (Euclidean), $x = z/M^2$
$a(z)$	form factor of a realization, Eqs. (1), (12)
$\hat{a}, \hat{c}$	skeleton $e^{n\Phi}$ and its coefficient e^{nC+}
R_{Π}, R_∞	fluctuation of $\log a$ and its limit, Eqs. (13), (31)
Ein, E_1	entire and ordinary exponential integrals
c	$\hat{c} e^{R_\infty}$, leading coefficient of a
$\mathcal{G}$	nonlocal remainder of F , Eq. (39)
$\mathcal{D}(p)$	graviton propagator, Eq. (53)
$P^{(2)}, P_s^{(0)}$	spin-two and spin-zero projectors
$\theta_{\mu\nu}, T_{\mu\nu}$	$\delta - \hat{k}\hat{k}$ and $\theta/3 + \hat{k}\hat{k}$
$f[z_0, \dots, z_m]$	divided difference, Sec. IV B
$\bar{\omega}(L)$	superficial degree from the vertex bound
$\beta_1, \beta_2, \beta_E$	one-loop poles of R^2, Ric^2, E
s_1, s_2, s_3	couplings of the quartic operators
J_n, K_n, I_n	bounds and value of $\int_0^\infty dk/\hat{a}(k^2)$

space computations are done in Euclidean signature, with $\square \rightarrow -p^2$. The curvature conventions are $R^\rho{}_{\sigma\mu\nu} = \partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma} + \Gamma^\rho_{\mu\lambda} \Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda} \Gamma^\lambda_{\mu\sigma}$ and $R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}$, and $\kappa^2 = 32\pi G$. The Gauss-Bonnet density is $E = \text{Riem}^2 - 4\text{Ric}^2 + R^2$. The Euclidean one-loop effective action is $\Gamma_1 = \frac{1}{2} \text{STr} \log S''$, where the supertrace includes the ghosts and the compensating factor of Sec. IV A. The ultraviolet pole of $\int d^D k (2\pi)^{-D} k^{-4}$ over $|k| \geq M$ is $1/(16\pi^2 \epsilon)$. In estimates, C and K denote constants that may change from line to line and depend on the indicated indices and on n , but not on momenta. Table XII collects the symbols used most often.

Appendix B: Angular averages

Ricci. Let $\bar{R}_{\mu\nu}$ be a symmetric tensor in four dimensions with trace $\bar{R}$, and write $\text{Ric}^2 = \bar{R}_{\mu\nu} \bar{R}^{\mu\nu}$. With $\theta = \delta - \hat{k}\hat{k}$ and $T = \theta/3 + \hat{k}\hat{k}$,

$$\begin{aligned} \bar{R} \cdot P^{(2)} \cdot \bar{R} &= \text{tr}(\theta \bar{R} \theta \bar{R}) - \frac{1}{3} (\text{tr} \theta \bar{R})^2, \\ \text{tr}(\theta \bar{R} \theta \bar{R}) &= \text{Ric}^2 - 2 \hat{k} \bar{R}^2 \hat{k} + (\hat{k} \bar{R} \hat{k})^2, \\ \text{tr} \theta \bar{R} &= \bar{R} - \hat{k} \bar{R} \hat{k}, \quad T \cdot \bar{R} = \frac{1}{3} \bar{R} + \frac{2}{3} \hat{k} \bar{R} \hat{k}. \end{aligned}$$

The averages over the unit sphere in four dimensions follow from the moments quoted in Sec. V C: $\langle \hat{k} \bar{R} \hat{k} \rangle = \bar{R}/4$, $\langle \hat{k} \bar{R}^2 \hat{k} \rangle = \text{Ric}^2/4$ and $\langle (\hat{k} \bar{R} \hat{k})^2 \rangle = (\bar{R}^2 + 2\text{Ric}^2)/24$. Hence

$$\begin{aligned} \langle \text{tr}(\theta \bar{R} \theta \bar{R}) \rangle &= \frac{7}{12} \text{Ric}^2 + \frac{1}{24} \bar{R}^2, \\ \langle (\text{tr} \theta \bar{R})^2 \rangle &= \frac{13}{24} \bar{R}^2 + \frac{1}{12} \text{Ric}^2, \\ \langle \bar{R} \cdot P^{(2)} \cdot \bar{R} \rangle &= \frac{5}{9} \text{Ric}^2 - \frac{5}{36} \bar{R}^2, \\ \langle (T \cdot \bar{R})^2 \rangle &= \frac{13}{54} \bar{R}^2 + \frac{1}{27} \text{Ric}^2, \end{aligned}$$

and

$$\langle \bar{R} \cdot P^{(2)} \cdot \bar{R} \rangle - \frac{3}{2} \langle (T \cdot \bar{R})^2 \rangle = \frac{1}{2} (\text{Ric}^2 - \bar{R}^2),$$

which is Eq. (71). For $\bar{R}_{\mu\nu} = \lambda \delta_{\mu\nu}$ the first average vanishes, as it must since $P^{(2)}$ annihilates $\delta_{\mu\nu}$, and the second equals $4\lambda^2$.

Riemann. Let $\bar{R}_{\mu\nu\rho\sigma}$ be an algebraic curvature tensor, with the symmetries of the Riemann tensor and the cyclic identity, and $W^{\nu\sigma} = \bar{R}^{\mu\nu\rho\sigma} \hat{k}_\mu \hat{k}_\rho$. Contracting the second and fourth indices gives the Ricci tensor, as contracting the first and third does, by pair symmetry; so $\text{tr} W = \hat{k} \bar{R} \hat{k}$ in the notation above, and $\langle (\text{tr} W)^2 \rangle = (\bar{R}^2 + 2\text{Ric}^2)/24$. For the other term,

$$\langle \text{tr} W^2 \rangle = \bar{R}^{\mu\nu\rho\sigma} \bar{R}^\alpha{}_{\nu}{}^\beta{}_{\sigma} \langle \hat{k}_\mu \hat{k}_\rho \hat{k}_\alpha \hat{k}_\beta \rangle.$$

The three pairings of the fourth moment give, in turn, $\bar{R}^{\mu\nu}{}_{\mu}{}^{\sigma} \bar{R}^\alpha{}_{\nu\alpha\sigma} = \text{Ric}^2$, $\bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\mu\nu\rho\sigma} = \text{Riem}^2$, and $\bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\rho\nu\mu\sigma}$. For the last one, pair symmetry gives $\bar{R}_{\rho\nu\mu\sigma} = \bar{R}_{\mu\sigma\rho\nu} = -\bar{R}_{\mu\sigma\nu\rho}$, and the cyclic identity $\bar{R}_{\mu\sigma\nu\rho} = -\bar{R}_{\mu\nu\rho\sigma} - \bar{R}_{\mu\rho\sigma\nu}$ gives

$$\begin{aligned} \bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\rho\nu\mu\sigma} &= \text{Riem}^2 + \bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\mu\rho\sigma\nu} \\ &= \text{Riem}^2 - \bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\mu\rho\nu\sigma}. \end{aligned}$$

The last contraction is the classical $\bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\mu\rho\nu\sigma} = \frac{1}{2} \text{Riem}^2$,²⁸ so the third pairing equals $\frac{1}{2} \text{Riem}^2$. Altogether $\langle \text{tr} W^2 \rangle = (\text{Ric}^2 + \frac{3}{2} \text{Riem}^2)/24$, and

$$\begin{aligned} \left\langle \text{tr} W^2 - \frac{1}{2} (\text{tr} W)^2 \right\rangle &= \frac{2\text{Ric}^2 + 3\text{Riem}^2 - \bar{R}^2 - 2\text{Ric}^2}{48} \\ &= \frac{3\text{Riem}^2 - \bar{R}^2}{48}, \end{aligned}$$

which is Eq. (72). The Ricci terms cancel, which is why the third column of the matrix (73) has no entry in Ric^2 before Riem^2 is traded for E . In dimensional regularization the averages are taken in D dimensions; the change is of order ϵ and affects finite parts only.

²⁸ Write $X = \bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\mu\rho\nu\sigma}$. Contracting the cyclic identity $\bar{R}_{\mu\nu\rho\sigma} + \bar{R}_{\mu\rho\sigma\nu} + \bar{R}_{\mu\sigma\nu\rho} = 0$ with $\bar{R}^{\mu\nu\rho\sigma}$ gives $\text{Riem}^2 + \bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\mu\rho\sigma\nu} + \bar{R}^{\mu\nu\rho\sigma} \bar{R}_{\mu\sigma\nu\rho} = 0$. By antisymmetry in the last pair the second term is $-X$, and relabeling $\rho \leftrightarrow \sigma$ in the third term and using antisymmetry in the last pair of the first factor shows that it is also $-X$. So $\text{Riem}^2 = 2X$.

Appendix C: Divided differences

Proof of Eq. (58). For $f(w) = w^k$, expand $(A + B)^k$ and keep the words with m letters B . They are $A^{j_m} B A^{j_{m-1}} \dots B A^{j_0}$ with $j_0 + \dots + j_m = k - m$. Acting on the plane waves, A^{j_i} becomes $z_i^{j_i}$, and the sum over the j_i is the complete homogeneous symmetric polynomial $h_{k-m}(z_0, \dots, z_m)$, which is the divided difference of w^k on $z_0, \dots, z_m$ [36].²⁹ Both sides of Eq. (58) are linear in f . For entire $f = \sum_k f_k w^k$ the series $\sum_k |f_k| \binom{k}{m} \max |z_i|^{k-m}$ converges, since $|h_{k-m}| \leq \binom{k}{m} \max |z_i|^{k-m}$, so the identity passes to the limit. The functions to which we apply the formula, $\mathcal{G}$ and $a - cx^n$, are entire, so this argument covers them; for functions that are merely smooth the same formula follows from the functional calculus of self-adjoint operators [35].

Proof of Eq. (59). The proof is by induction on m ; the case $m = 0$ is the hypothesis $|f(z)| \leq C_0(1 + z)^\mu$. By symmetry of the divided difference let $z_0 \leq z_1 \leq \dots \leq z_m$, so that $z_{\min} = z_0$. There are two cases.

If $1 + z_m \leq 2(1 + z_0)$, all arguments are comparable. In Eq. (57) the point $\sum t_i z_i$ lies in $[z_0, z_m]$, so $|f^{(m)}| \leq C_m(1 + z_0)^{\mu-m}$ there, and the simplex has volume $1/m!$. Hence

$$\begin{aligned} |f[z_0, \dots, z_m]| &\leq \frac{C_m}{m!} (1 + z_0)^{\mu-m} \\ &\leq \frac{2^m C_m}{m!} (1 + z_0)^\mu \prod_{i=1}^m (1 + z_i)^{-1}, \end{aligned}$$

using $(1 + z_0)^{-1} \leq 2(1 + z_i)^{-1}$ for each $i \geq 1$.

If $1 + z_m > 2(1 + z_0)$, then $z_m - z_0 > \frac{1}{2}(1 + z_m)$ and we use the recursion. By the induction hypothesis applied to $z_1, \dots, z_m$, whose smallest element is z_1 , and by $(1 + z_1)^\mu \leq (1 + z_0)^\mu$ for $\mu \leq 0$,

$$|f[z_1, \dots, z_m]| \leq K_{m-1} (1 + z_0)^\mu \prod_{i=2}^m (1 + z_i)^{-1}.$$

Dividing by $z_m - z_0 > \frac{1}{2}(1 + z_m) \geq \frac{1}{2}(1 + z_1)$ gives at most $2K_{m-1}(1 + z_0)^\mu \prod_{i=1}^m (1 + z_i)^{-1}$. The other term, $f[z_0, \dots, z_{m-1}]$, is bounded by $K_{m-1}(1 + z_0)^\mu \prod_{i=1}^{m-1} (1 + z_i)^{-1}$, and dividing by $z_m - z_0 > \frac{1}{2}(1 + z_m)$ gives the same bound. So $K_m = \max(2^m C_m/m!, 4K_{m-1})$ works, which completes the induction.

The two symbols. For $f = \mathcal{G}$ the hypothesis holds with $\mu = -1$ by Eq. (40), and Eq. (59) reads $|\mathcal{G}[z_0, \dots, z_m]| \leq K_m \prod_{i=0}^m (1 + z_i)^{-1}$, the form used in Sec. IV C. For $f = a - cx^n$ the derivative bounds follow from Eq. (23) and the large-momentum behavior on the real axis: $a - cx^n =$

$O(x^{n-\sigma/2+\varepsilon})$ for $x \geq 1$, and each derivative is one power of x smaller (Sec. II G), so the hypothesis holds with $\mu = 0$.

Proof of Eq. (61). Write $\Lambda = (1 + |q|)(1 + |q'|) \prod_a (1 + |\ell_a|)$ and label the legs so that $|\ell_1| \geq |\ell_2| \geq \dots$, with $\ell_2 = 0$ if $r = 1$. Each of the products $|q||q'|$, $|q||\ell_1|$, $|\ell_1||\ell_2|$ and $|\ell_1|(|q| + |q'|)$ is at most Λ , because every factor on the left appears on the right with 1 added to it. From $q = q' - \sum_a \ell_a$ we have $|q| \leq |q'| + s$, and $s \leq r|\ell_1|$, so

$$|q|^2 \leq |q||q'| + r|q||\ell_1| \leq (1 + r)\Lambda, \quad 2|q|s \leq 2r\Lambda.$$

Momentum conservation solved for the largest leg, $\ell_1 = q' - q - \sum_{a \geq 2} \ell_a$, gives $|\ell_1| \leq |q| + |q'| + (r - 1)|\ell_2|$, so that $s \leq |\ell_1| + (r - 1)|\ell_2| \leq |q| + |q'| + 2(r - 1)|\ell_2|$, and therefore

$$\begin{aligned} s^2 &\leq r|\ell_1|s \leq r|\ell_1|(|q| + |q'|) + 2r(r - 1)|\ell_1||\ell_2| \\ &\leq (2r^2 - r)\Lambda. \end{aligned}$$

Adding the three pieces gives $(|q| + s)^2 \leq (1 + 2r + 2r^2)\Lambda$, which is Eq. (61) with $C_r = 1 + 2r + 2r^2$. The powers cannot be improved: for $r = 2$, $q = q' = 0$ and $\ell_1 = -\ell_2 = \lambda e$ with $|e| = 1$, both sides are of order λ^2 .³⁰

Complex arguments. Section VII needs Eq. (59) for complex arguments. Let K be a compact set that contains 0, let $S_\delta = \{|\arg z| \leq \pi/4 - \delta\}$, and put $\Omega = \text{conv } K + S_\delta$. Because S_δ is a convex cone, Ω is convex and contains every convex combination of its points, in particular the point $\sum_i t_i z_i$ of Eq. (57). Writing each $z_i = k_i + v_i$ with $k_i \in \text{conv } K$ and $v_i \in S_\delta$, and using $\text{Re } v \geq |v|/\sqrt{2}$ on S_δ , one finds $|\sum_i t_i z_i| \geq (\min_i |z_i| - C_K)/\sqrt{2} - C_K$, so that $1 + |\sum_i t_i z_i| \geq \kappa(1 + \min_i |z_i|)$ with $\kappa > 0$ depending on K and δ only. Suppose f is analytic near Ω with $|f^{(j)}(z)| \leq C_j(1 + |z|)^{\mu-j}$ there. Then the proof above goes through with z_i replaced by $|z_i|$ in the case distinction: the recursion case uses $|z_m - z_0| \geq |z_m| - |z_0|$, and the comparable case uses the lower bound just derived. So Eq. (59) holds on Ω with $1 + z_i$ replaced by $1 + |z_i|$. For $f = \mathcal{G}$ and $f = a - cx^n$ the derivative bounds on Ω follow from the cone asymptotics (37): far from the origin Ω lies in the wider cone $S_{\delta/2}$, where $\mathcal{G} = -1/z$ up to a term of order $|z|^{n-1-\sigma/2+\varepsilon}$, which decays faster than $1/|z|$, and Cauchy's estimate on a disk of radius $\frac{1}{2}|z|\sin(\delta/2)$ around z , which stays in $S_{\delta/2}$, bounds the j th derivative of that term by $C_j|z|^{n-1-\sigma/2+\varepsilon-j}$.

²⁹ The identity $w^k[z_0, \dots, z_m] = h_{k-m}(z_0, \dots, z_m)$ follows by induction on m from the recursion and $h_r(z_1, \dots, z_m) - h_r(z_0, \dots, z_{m-1}) = (z_m - z_0)h_{r-1}(z_0, \dots, z_m)$, which holds because both sides count the monomials of degree r that contain z_m but not z_0 , minus those that contain z_0 but not z_m .

³⁰ The estimate is the point at which a vertex with many hard legs differs from one with all momenta comparable. Without it one would bound an insertion by $(1 + q_{i-1}^2) \prod_a (1 + \ell_a^2)$, which charges two powers to each leg and none to the outgoing running momentum. The divided difference still cancels the running momenta, but each hard leg then costs λ^2 instead of λ , and the counting of Sec. IV D closes only for $n \geq 6$.

TABLE XIII. Independent checks. “Py” is `verify.py` (178 checks), “C” is `formfactor_check.c` (80 checks) and “Jl” is `verify.jl` (96 checks); all pass. “Lean” is the number of theorems of `PowerCounting.lean` that prove the statement for all values of its integer arguments; for the insertion lemma only the arithmetic of the constant C_r is formalized. The Julia angular averages use the 24-cell, a spherical design of degree 5, in exact rational arithmetic. The last column gives the equation or section checked.

statement	Py	C	Jl	Lean	where
γ_E as a difference of two integrals; $\text{Ein}(y) = \log y + \gamma_E + E_1(y)$, real and complex y	2	10	8	—	Eqs. (19), (20)
$\int_0^1 \psi_*(x\tau) d\tau/\tau = \frac{1}{2} \text{Ein}(x^2)$; power series of the exponent at complex z	8	—	—	—	Eqs. (14), (15)
probability generating functional, Monte Carlo over the gas	1	—	—	—	Eq. (9)
$0 \leq \text{Ein}(y) - \log(1+y) \leq \gamma_E$, monotone; two-sided bound (21), $n = 1, \dots, 12$	3	14	4	—	Eq. (21)
low-momentum coefficients $n/2$ and $n(n-1)/8$	4	3	2	—	Eq. (22)
$ E_1(w) \leq e^{-\text{Re } w}/ w $; cone asymptotics with explicit error; decay on the imaginary axis	2	2	12	—	Eq. (26)
resonances of $2 - e^{-s^2 z^2}$ outside the cones; expected count $nR^2/(\pi M^4)$ (ratio 1.0003 at $R = 60M^2$)	16	1	—	—	Eq. (4)
spin projectors, linearized curvature, gauge invariance, $D = 4$ and $D = 5$	10	—	1	—	Eqs. (47)–(52)
angular averages: Monte Carlo, isotropic tensors, and the exact 600-cell cubature of degree 8	8	2	6	—	App. B
correlators of δR , $\delta R_{\mu\nu}$, $\delta R_{\mu\nu\rho\sigma}$ with random algebraic curvature tensors	5	—	—	—	Eqs. (67)–(70)
quartic-operator poles, matrix (73), determinant -24 , tuning (74) (exact rationals in C)	9	3	4	3	Sec. V D
Hermite–Genocchi; $w^k[z_0, \dots, z_m] = h_{k-m}$; scaling of the bound (59) in mixed configurations	8	—	—	—	App. C
$\bar{\omega}(L) < 0$ for all $L \geq 2$ exactly when $n \geq 4$; $n = 3$ marginal; Table V; merging inequality	13	4	3	9	Eq. (62)
insertion lemma, random four-momenta over seven decades; the two-insertion chain of Fig. 8	2	1	1	1	Eq. (61)
growth exponent equals the strength per e -fold	3	—	—	—	Eq. (6)
$\text{Ein}(w) \leq w$; J_n formula; $K_n \leq I_n \leq J_n \leq \frac{2n}{2n-1} K_n$; J_{2m} product; $\rho_{\text{eff}}(0)$ bounds; Wendel’s inequality; $n = 4, \dots, 14$	38	28	16	—	Eqs. (87)–(91)
decay of $\delta(r)$	1	—	—	—	Eq. (93)
zero-point count $\log p^2 + 3 \log a$; Mellin transform $\Gamma(s)/s$; sharp and smooth cutoffs, constant term 1; $\rho_1 = 3nM^4/(64\pi^2)$	6	4	—	—	Eqs. (75)–(77)
admissible class: C_+ , moments $\int t(1-\psi) dt$, boundedness of $\Phi - \frac{1}{2} \log(1+x^2)$, Mellin continuation	12	3	1	—	Eqs. (27)–(29)
incomplete-gamma variances (42), (43) against quadrature; $\mathbb{E}R_\infty^2 = n\vartheta_0/\zeta$; Eq. (30)	3	2	14	—	Sec. II G
Monte Carlo over 4×10^4 gases: Campbell mean, isometry (10), variance of Eq. (33), $\mathbb{E}a_\Pi \neq \hat{a}$ by Eq. (11), variance of $\delta\rho_\Pi$	4	—	—	—	Secs. II C, VI B
R_∞ sampled from its Lévy measure: variance, Eq. (44), lower tail (45), $n = 4, 6, 8$	1	1	5	—	Sec. II G
identical branes: Ψ_* in closed form and its asymptotics; variance and characteristic function by Monte Carlo; decay of $ \mathbb{E} e^{iuR_\Pi} $	7	1	4	—	Eq. (41)
mean-square decay (35) along rays; one realization: slope $-\zeta/2$ over dyadic annuli, and $a - cx^n$ bounded for $\zeta = 9$, growing for $\zeta = 5$	2	—	—	—	Eqs. (36), (40)
argument principle on $ z = 2M^2$: winding 0 for a realization, equal to the count of Eq. (4) for a resonant product	1	—	—	—	Sec. II A
Table II; relative fluctuation (80); mean of $\delta\rho_\Pi$ by continuation	3	1	7	—	Eqs. (79), (80)
remainder $\rightarrow 1$ for the cutoffs e^{-u^p} ; sharp-cutoff remainder $1 - e^{-W} + WE_1(W)$	2	—	—	—	Sec. VI B
matter scattering: $T_1 \cdot T_2 - \frac{1}{2} T_1 T_2 = -su$ for massless scalars; maximum of $x(1+x^2)^{-n/2}$, $n = 4, \dots, 8$; the bound below Eq. (82) for the skeleton	—	—	8	—	Sec. VII C
I_4 , I_5 and $ \Phi_N(0) $ for $n = 4, 5$; $J_4 = 3\pi\sqrt{2}/16$; the constants c for $n = 4, \dots, 8$	4	—	—	—	Table IX

Appendix D: Numerical checks

The closed-form identities, inequalities and coefficients derived in the text were re-evaluated at sample points by three independent programs, in Python, C and Julia, which share no code. Table XIII lists the groups of checks. Where an inequality is checked, it is checked on a grid or at random points, and the tolerance is far below the margin by which it holds; where an identity is checked, the two sides are computed by different routes (closed form against quadrature, series against continued fraction, exact rational arithmetic against floating point). The Julia program works in 256-bit arithmetic with its own power series, continued fractions, arithmetic-geometric mean and double-exponential quadrature. These checks confirm the algebra and guard

against transcription errors; they are not part of any proof.

The integer statements on which the power counting and the one-loop tuning rest are, in addition, proved in Lean 4 without external libraries: the merging inequality and its iterated form, the bound $\bar{\omega}(L)$ on the superficial degree of any graph, its sign for all $n \geq 4$ and $L \geq 2$ and for $n \leq 3$ at two loops, the entries of Table V, the determinants -24 and -12 of the pole matrix, and the cancellation of the three poles by the tuning (74). The kernel accepts every proof, and the only axioms used are the standard ones of Lean’s core logic, propositional extensionality and quotient soundness. These proofs repeat arguments given in the text in full; the analytic estimates, on special functions, integrals and the gas, are not formalized.

-
- [1] G. 't Hooft and M. Veltman, One-loop divergencies in the theory of gravitation, *Ann. Inst. H. Poincaré Sect. A* **20**, 69 (1974).
 - [2] M. H. Goroff and A. Sagnotti, The ultraviolet behavior of Einstein gravity, *Nucl. Phys. B* **266**, 709 (1986), [doi:10.1016/0550-3213\(86\)90193-8](#).
 - [3] K. S. Stelle, Renormalization of higher-derivative quantum gravity, *Phys. Rev. D* **16**, 953 (1977), [doi:10.1103/PhysRevD.16.953](#).
 - [4] I. L. Shapiro, Counting ghosts in the “ghost-free” non-local gravity, *Phys. Lett. B* **744**, 67 (2015), [doi:10.1016/j.physletb.2015.03.037](#).
 - [5] G. V. Efimov, Non-local quantum theory of the scalar field, *Commun. Math. Phys.* **5**, 42 (1967), [doi:10.1007/BF01646357](#).
 - [6] N. V. Krasnikov, Nonlocal gauge theories, *Theor. Math. Phys.* **73**, 1184 (1987), [doi:10.1007/BF01017588](#).
 - [7] Yu. V. Kuz'min, Finite nonlocal gravity, *Sov. J. Nucl. Phys.* **50**, 1011 (1989) [*Yad. Fiz.* **50**, 1630 (1989)].
 - [8] E. T. Tomboulis, Superrenormalizable gauge and gravitational theories, *arXiv:hep-th/9702146*.
 - [9] T. Biswas, A. Mazumdar, and W. Siegel, Bouncing universes in string-inspired gravity, *J. Cosmol. Astropart. Phys.* **03** (2006) 009, [doi:10.1088/1475-7516/2006/03/009](#).
 - [10] T. Biswas, E. Gerwick, T. Koivisto, and A. Mazumdar, Towards singularity- and ghost-free theories of gravity, *Phys. Rev. Lett.* **108**, 031101 (2012), [doi:10.1103/PhysRevLett.108.031101](#).
 - [11] L. Modesto, Super-renormalizable quantum gravity, *Phys. Rev. D* **86**, 044005 (2012), [doi:10.1103/PhysRevD.86.044005](#).
 - [12] L. Modesto and L. Rachwał, Super-renormalizable and finite gravitational theories, *Nucl. Phys. B* **889**, 228 (2014), [doi:10.1016/j.nuclphysb.2014.10.015](#).
 - [13] D. Bhattacharjee, Brane cluster gases and ghost-free form factors for quantum gravity (2026), manuscript; source, verification suite and figure scripts, Zenodo, [doi:10.5281/zenodo.22866290](#).
 - [14] S. Talaganis, T. Biswas, and A. Mazumdar, Towards understanding the ultraviolet behavior of quantum loops in infinite-derivative theories of gravity, *Classical Quantum Gravity* **32**, 215017 (2015), [doi:10.1088/0264-9381/32/21/215017](#).
 - [15] F. Briscese and L. Modesto, Cutkosky rules and perturbative unitarity in Euclidean nonlocal quantum field theories, *Phys. Rev. D* **99**, 104043 (2019), [doi:10.1103/PhysRevD.99.104043](#).
 - [16] R. Pius and A. Sen, Cutkosky rules for superstring field theory, *J. High Energy Phys.* **10** (2016) 024; **09** (2018) 122(E), [doi:10.1007/JHEP10\(2016\)024](#).
 - [17] T. Kugo and I. Ojima, Subsidiary conditions and physical S-matrix unitarity in indefinite-metric quantum gravitational theory, *Nucl. Phys. B* **144**, 234 (1978), [doi:10.1016/0550-3213\(78\)90504-7](#).
 - [18] S. Weinberg, Infrared photons and gravitons, *Phys. Rev.* **140**, B516 (1965), [doi:10.1103/PhysRev.140.B516](#).
 - [19] L. L. Foldy, The multiple scattering of waves. I. General theory of isotropic scattering by randomly distributed scatterers, *Phys. Rev.* **67**, 107 (1945), [doi:10.1103/PhysRev.67.107](#).
 - [20] M. Lax, Multiple scattering of waves, *Rev. Mod. Phys.* **23**, 287 (1951), [doi:10.1103/RevModPhys.23.287](#).
 - [21] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed. (Oxford University Press, Oxford, 1939).
 - [22] T. D. Lee and G. C. Wick, Negative metric and the unitarity of the S-matrix, *Nucl. Phys. B* **9**, 209 (1969), [doi:10.1016/0550-3213\(69\)90098-4](#).
 - [23] D. Anselmi, On the quantum field theory of the gravitational interactions, *J. High Energy Phys.* **06** (2017) 086, [doi:10.1007/JHEP06\(2017\)086](#).
 - [24] D. Anselmi and M. Piva, Perturbative unitarity of Lee–Wick quantum field theory, *Phys. Rev. D* **96**, 045009 (2017), [doi:10.1103/PhysRevD.96.045009](#).
 - [25] J. F. C. Kingman, *Poisson Processes* (Oxford University Press, Oxford, 1993), [doi:10.1093/oso/9780198536932.001.0001](#).
 - [26] G. Last and M. Penrose, *Lectures on the Poisson Process* (Cambridge University Press, Cambridge, 2017), [doi:10.1017/9781316104477](#).
 - [27] W. Feller, *An Introduction to Probability Theory and Its Applications*, Vol. II, 2nd ed. (Wiley, New York, 1971), ISBN 978-0-471-25709-7.
 - [28] F. W. J. Olver *et al.* (eds.), *NIST Digital Library of Math-*

- ematical Functions*, Chap. 6, <https://dlmf.nist.gov/6>.
- [29] R. J. Rivers, Lagrangian theory for neutral massive spin-2 fields, *Nuovo Cimento* **34**, 386 (1964), [doi:10.1007/BF02734585](https://doi.org/10.1007/BF02734585).
 - [30] P. van Nieuwenhuizen, On ghost-free tensor Lagrangians and linearized gravitation, *Nucl. Phys. B* **60**, 478 (1973), [doi:10.1016/0550-3213\(73\)90194-6](https://doi.org/10.1016/0550-3213(73)90194-6).
 - [31] B. P. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration, Fermi GBM, and INTEGRAL), Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A, *Astrophys. J. Lett.* **848**, L13 (2017), [doi:10.3847/2041-8213/aa920c](https://doi.org/10.3847/2041-8213/aa920c).
 - [32] L. F. Abbott, The background field method beyond one loop, *Nucl. Phys. B* **185**, 189 (1981), [doi:10.1016/0550-3213\(81\)90371-0](https://doi.org/10.1016/0550-3213(81)90371-0).
 - [33] N. K. Nielsen, Ghost counting in supergravity, *Nucl. Phys. B* **140**, 499 (1978), [doi:10.1016/0550-3213\(78\)90009-3](https://doi.org/10.1016/0550-3213(78)90009-3).
 - [34] R. E. Kallosh, Modified Feynman rules in supergravity, *Nucl. Phys. B* **141**, 141 (1978), [doi:10.1016/0550-3213\(78\)90340-1](https://doi.org/10.1016/0550-3213(78)90340-1).
 - [35] Yu. L. Daletskii and S. G. Krein, Integration and differentiation of functions of Hermitian operators and applications to the theory of perturbations, *Am. Math. Soc. Transl.* (2) **47**, 1 (1965), [doi:10.1090/trans2/047/01](https://doi.org/10.1090/trans2/047/01).
 - [36] C. de Boer, Divided differences, *Surv. Approx. Theory* **1**, 46 (2005), [arXiv:math/0502036](https://arxiv.org/abs/math/0502036).
 - [37] M. Asorey, J. L. López, and I. L. Shapiro, Some remarks on high derivative quantum gravity, *Int. J. Mod. Phys. A* **12**, 5711 (1997), [doi:10.1142/S0217751X97002991](https://doi.org/10.1142/S0217751X97002991).
 - [38] S. Weinberg, High-energy behavior in quantum field theory, *Phys. Rev.* **118**, 838 (1960), [doi:10.1103/PhysRev.118.838](https://doi.org/10.1103/PhysRev.118.838).
 - [39] A. O. Barvinsky and G. A. Vilkovisky, The generalized Schwinger–DeWitt technique in gauge theories and quantum gravity, *Phys. Rep.* **119**, 1 (1985), [doi:10.1016/0370-1573\(85\)90148-6](https://doi.org/10.1016/0370-1573(85)90148-6).
 - [40] L. Modesto, L. Rachwał, and I. L. Shapiro, Renormalization group in super-renormalizable quantum gravity, *Eur. Phys. J. C* **78**, 555 (2018), [doi:10.1140/epjc/s10052-018-6035-2](https://doi.org/10.1140/epjc/s10052-018-6035-2).
 - [41] R. E. Kallosh, O. V. Tarasov, and I. V. Tyutin, One-loop finiteness of quantum gravity off mass shell, *Nucl. Phys. B* **137**, 145 (1978), [doi:10.1016/0550-3213\(78\)90055-X](https://doi.org/10.1016/0550-3213(78)90055-X).
 - [42] J. C. Collins, *Renormalization* (Cambridge University Press, Cambridge, 1984), [doi:10.1017/CB09780511622656](https://doi.org/10.1017/CB09780511622656).
 - [43] G. Barnich, F. Brandt, and M. Henneaux, Local BRST cohomology in gauge theories, *Phys. Rep.* **338**, 439 (2000), [doi:10.1016/S0370-1573\(00\)00049-1](https://doi.org/10.1016/S0370-1573(00)00049-1).
 - [44] L. F. Abbott, M. T. Grisaru, and R. K. Schaefer, The background field method and the S-matrix, *Nucl. Phys. B* **229**, 372 (1983), [doi:10.1016/0550-3213\(83\)90337-1](https://doi.org/10.1016/0550-3213(83)90337-1).
 - [45] P. Chin and E. T. Tomboulis, Nonlocal vertices and analyticity: Landau equations and general Cutkosky rule, *J. High Energy Phys.* **06** (2018) 014, [doi:10.1007/JHEP06\(2018\)014](https://doi.org/10.1007/JHEP06(2018)014).
 - [46] T. Kugo and I. Ojima, Local covariant operator formalism of non-Abelian gauge theories and quark confinement problem, *Prog. Theor. Phys. Suppl.* **66**, 1 (1979), [doi:10.1143/PTPS.66.1](https://doi.org/10.1143/PTPS.66.1).
 - [47] C. D. Carone, Unitarity and microscopic acausality in a nonlocal theory, *Phys. Rev. D* **95**, 045009 (2017), [doi:10.1103/PhysRevD.95.045009](https://doi.org/10.1103/PhysRevD.95.045009).
 - [48] P. Donà, S. Giaccari, L. Modesto, L. Rachwał, and Y. Zhu, Scattering amplitudes in super-renormalizable gravity, *J. High Energy Phys.* **08** (2015) 038, [doi:10.1007/JHEP08\(2015\)038](https://doi.org/10.1007/JHEP08(2015)038).
 - [49] G. W. Gibbons, S. W. Hawking, and M. J. Perry, Path integrals and the indefiniteness of the gravitational action, *Nucl. Phys. B* **138**, 141 (1978), [doi:10.1016/0550-3213\(78\)90161-X](https://doi.org/10.1016/0550-3213(78)90161-X).
 - [50] A. Jaffe and E. Witten, Quantum Yang–Mills theory, in *The Millennium Prize Problems*, edited by J. Carlson, A. Jaffe, and A. Wiles (American Mathematical Society and Clay Mathematics Institute, Providence, RI, 2006), ISBN 978-0-8218-3679-8.
 - [51] J. G. Wendel, Note on the gamma function, *Am. Math. Mon.* **55**, 563 (1948), [doi:10.2307/2304460](https://doi.org/10.2307/2304460).
 - [52] J. Edholm, A. S. Koshelev, and A. Mazumdar, Behavior of the Newtonian potential for ghost-free gravity and singularity-free gravity, *Phys. Rev. D* **94**, 104033 (2016), [doi:10.1103/PhysRevD.94.104033](https://doi.org/10.1103/PhysRevD.94.104033).
 - [53] J. F. Donoghue, General relativity as an effective field theory: The leading quantum corrections, *Phys. Rev. D* **50**, 3874 (1994), [doi:10.1103/PhysRevD.50.3874](https://doi.org/10.1103/PhysRevD.50.3874).
 - [54] J. G. Lee, E. G. Adelberger, T. S. Cook, S. M. Fleischer, and B. R. Heckel, New test of the gravitational $1/r^2$ law at separations down to $52\,\mu\text{m}$, *Phys. Rev. Lett.* **124**, 101101 (2020), [doi:10.1103/PhysRevLett.124.101101](https://doi.org/10.1103/PhysRevLett.124.101101).
 - [55] J. Polchinski, Dirichlet branes and Ramond–Ramond charges, *Phys. Rev. Lett.* **75**, 4724 (1995), [doi:10.1103/PhysRevLett.75.4724](https://doi.org/10.1103/PhysRevLett.75.4724).
 - [56] C. V. Johnson, *D-Branes* (Cambridge University Press, Cambridge, 2003), [doi:10.1017/CB09780511606540](https://doi.org/10.1017/CB09780511606540).