

A COUNTEREXAMPLE TO A CONJECTURE OF GRAFFITI.PC ON THE FOREST NUMBER

DEEP BHATTACHARJEE

ABSTRACT. Conjecture 66 of Graffiti.pc bounds the order of a largest induced forest of a connected graph from below by the even mode of the degrees of its complement divided by its average degree. We show that it is false, under every reading of its definitions and by an unbounded margin.

1. INTRODUCTION

Graffiti.pc is a program written by DeLaViña [3] after Fajtlowicz’s Graffiti [5]; it proposes inequalities between graph invariants that hold on its database of graphs. DeLaViña’s list *Written on the Wall II* [4] records its conjectures with their status. Conjecture 66, generated on 25 March 2004, is still marked as open (in the version of 26 July 2026). This note shows that it is false. The proofs are by hand; as in [2], every finite statement is also re-checked by computer, and the counterexamples are also confirmed in the Lean proof assistant (Section 3 and Appendix A).

All graphs are finite and simple. For a graph G with n vertices and m edges, $\deg_{\text{avg}}(G) = 2m/n$ is the average degree and $\overline{G}$ the complement, in which a vertex of degree d in G has degree $n - 1 - d$. The *forest number* $f(G)$ is the largest number of vertices of an induced subgraph of G that is a forest. The list defines the *even mode* of a graph as “the most frequently occurring degree that is an even integer”, with $\text{even_mode}_{\min}$ the smallest one when several are equally frequent. We write $\text{em}(H)$ for this number: among the even values in the degree sequence of H , the most frequent one, and the smallest of these in case of a tie. Conjecture 66 reads as follows in [4].

Conjecture 66. If G is a simple connected graph, then

$$f(G) \geq 2 \left\lfloor \frac{\text{em}(\overline{G})}{\deg_{\text{avg}}(G)} \right\rfloor.$$

The definition of the even mode also admits a second reading: the smallest even value among the most frequent degrees, defined only when some most frequent degree is even. And while the statement uses $\lfloor \cdot \rfloor$, the definitions that the list attaches to it are those of f , the even mode, the complement, the average degree and the floor function $\lfloor \cdot \rfloor$. The list also records that in 2008 all 11,716,571 connected graphs with 10 vertices were added to the database of Graffiti.pc (see its notes to Conjectures 42 and 187). We show in Section 3 that the printed conjecture fails on seven of these graphs, while the version with $\lfloor \cdot \rfloor$ holds on all of them; so the floor may well be what the program meant. We therefore treat all four readings.

2020 *Mathematics Subject Classification*. Primary 05C69; Secondary 05C05, 05C07.

Key words and phrases. induced forest, forest number, degree sequence, complement, Graffiti.pc, counterexample.

Corresponding author: d.bhattacharjee@erl-forschung.de; ORCID [0000-0003-0466-750X](https://orcid.org/0000-0003-0466-750X).




FIGURE 1. The graph H_c : copies $Q_1, \dots, Q_c$ of K_4 , each drawn as a tetrahedron, joined by the edges $x_{i,1}x_{i+1,0}$. The shaded vertices $x_{i,2}, x_{i,3}$ induce a forest of order $2c$.

For $c \geq 1$ let H_c be the graph made of c disjoint copies $Q_1, \dots, Q_c$ of the complete graph K_4 , with vertices $x_{i,0}, x_{i,1}, x_{i,2}, x_{i,3}$ in Q_i , together with the $c - 1$ edges $x_{i,1}x_{i+1,0}$ for $1 \leq i < c$ (Figure 1).

Theorem 1.1. *For every $c \geq 1$, $f(H_c) = 2c$, and the right side of Conjecture 66 for H_c equals $2\lceil q_c \rceil$ with*

$$q_c = \frac{8c(c-1)}{7c-1},$$

under both readings of the even mode. Consequently Conjecture 66 fails for H_c if and only if $c \geq 8$, and the conjecture with $\lfloor \cdot \rfloor$ in place of $\lceil \cdot \rceil$ fails for H_c if and only if $c \geq 14$. In both cases the right side exceeds $f(H_c)$ by at least $2\lfloor (c-7)/7 \rfloor$, which tends to infinity.

Theorem 1.2. *No connected graph with 2 to 9 vertices violates any of the four readings. Among the 11,716,571 connected graphs with 10 vertices, exactly 7 violate Conjecture 66 as printed, 5 violate it when the even mode is read in the second way, and none violates either reading with $\lfloor \cdot \rfloor$.*

The smallest member of the family that refutes all four readings, H_{14} , has 56 vertices. We did not determine the smallest counterexample to the readings with $\lfloor \cdot \rfloor$; the smallest one we found has 37 vertices (Remark 2.5).

2. THE CHAINS OF COMPLETE GRAPHS

Lemma 2.1. $f(H_c) = 2c$.

Proof. Three vertices of one Q_i induce a triangle, so an induced forest contains at most two vertices of each Q_i ; since the Q_i partition the vertex set, it has at most $2c$ vertices. The $2c$ vertices $x_{i,2}, x_{i,3}$ induce c disjoint edges, which form a forest. $\square$

Lemma 2.2. H_c is connected, has $7c - 1$ edges, and $\deg_{\text{avg}}(H_c) = (7c - 1)/(2c)$. It has $2c + 2$ vertices of degree 3 and $2c - 2$ vertices of degree 4. In both readings the even mode of $\overline{H_c}$ equals $4c - 4$.

Proof. The copies Q_i are connected and consecutive ones are joined by an edge, so H_c is connected; it has $6c$ edges inside the copies and $c - 1$ between them. Every vertex has degree 3 inside its copy, and the $2(c - 1)$ endpoints $x_{i,1}$ ($i < c$) and $x_{i,0}$ ($i > 1$) of the joining edges, which are distinct, have one more neighbour. In $\overline{H_c}$, which has $n = 4c$ vertices, the degrees are therefore $4c - 4$, which is even, taken $2c + 2$ times, and $4c - 5$, which is odd, taken $2c - 2$ times. So $4c - 4$ is the only even degree of $\overline{H_c}$, and it is also its most frequent degree, because $2c + 2 > 2c - 2$. $\square$

Lemma 2.3. *Let $c \geq 1$ and $q_c = 8c(c-1)/(7c-1)$. Then $\lceil q_c \rceil \geq c+1$ if and only if $c \geq 8$, and $\lfloor q_c \rfloor \geq c+1$ if and only if $c \geq 14$. For $c \geq 7$,*

$$\lfloor q_c \rfloor - c \geq \left\lfloor \frac{c-7}{7} \right\rfloor.$$

Proof. We have

$$q_c - c = \frac{8c^2 - 8c - 7c^2 + c}{7c-1} = \frac{c(c-7)}{7c-1}.$$

Since c is an integer, $\lceil q_c \rceil \geq c+1$ if and only if $q_c > c$, that is, $c > 7$. Next, $q_c \geq c+1$ if and only if $c(c-7) \geq 7c-1$, that is, $c^2 - 14c + 1 \geq 0$. The roots of $t^2 - 14t + 1$ are $7 \pm \sqrt{48}$, which lie in $(0, 1)$ and $(13, 14)$, so for positive integers c this holds exactly when $c \geq 14$. Finally, for $c \geq 7$ we have $c/(7c-1) \geq 1/7$ and $c-7 \geq 0$, so $q_c - c \geq (c-7)/7$, and taking integer parts gives the last claim. $\square$

Proof of Theorem 1.1. By Lemma 2.2, under either reading of the even mode,

$$\frac{\text{em}(\overline{H_c})}{\deg_{\text{avg}}(H_c)} = \frac{(4c-4) \cdot 2c}{7c-1} = q_c.$$

By Lemma 2.1, Conjecture 66 fails for H_c exactly when $2\lceil q_c \rceil > 2c$, and the version with the floor fails exactly when $2\lfloor q_c \rfloor > 2c$; Lemma 2.3 identifies these cases. The difference between the right side and $f(H_c)$ is at least $2\lfloor q_c \rfloor - 2c \geq 2\lfloor (c-7)/7 \rfloor$. $\square$

For example, H_8 has 32 vertices, $f(H_8) = 16$ and $q_8 = 448/55 \approx 8.15$, so the printed right side is 18. For H_{14} , with 56 vertices, $f(H_{14}) = 28$ and $q_{14} = 1456/97 \approx 15.01$, so the right side is 32 with $\lceil \cdot \rceil$ and 30 with $\lfloor \cdot \rfloor$.

Remark 2.4. The family is close to extremal in the following sense. Alon, Kahn and Seymour [1] proved that $f(G) \geq \sum_v \min\{1, 2/(d(v)+1)\}$, with equality for disjoint unions of complete graphs. For a connected graph whose degrees are all close to d , the right side of Conjecture 66 is at most about $2n/d$ while $f(G)$ is at least about $2n/(d+1)$; a counterexample therefore needs the near-equality of disjoint cliques, a large even mode of the complement, and few extra edges to make the graph connected. The chains H_c have all three.

Remark 2.5. Let T be the graph made of 12 disjoint triangles $a_i b_i c_i$, the 11 edges $b_i a_{i+1}$, and one pendant vertex attached to a_1 . It has 37 vertices and 48 edges, $f(T) = 25$ (at most two vertices of each triangle, and the pendant vertex), and the only even degree of $\overline{T}$ is 34, taken by the 13 vertices of degree 2 in T . With the first reading of the even mode, $34/\deg_{\text{avg}}(T) = 34 \cdot 37/96 \approx 13.10$, so the right side is 26 with $\lfloor \cdot \rfloor$ and 28 with $\lceil \cdot \rceil$. The most frequent degree of $\overline{T}$ is the odd number 33, so T says nothing about the second reading.

3. SMALL GRAPHS

Let G_{10} be the graph obtained from a triangle $t_1 t_2 t_3$ and a complete graph on $\{q_1, q_2, q_3, q_4\}$ by adding the edge $t_1 q_1$ and three pendant vertices ℓ_1, ℓ_2, ℓ_3 , with ℓ_i adjacent to q_i (Figure 2).

Proposition 3.1. *G_{10} is a counterexample to Conjecture 66 as printed, under both readings of the even mode: $f(G_{10}) = 7$, while the right side is 8.*

Proof. An induced forest contains at most two vertices of the triangle, at most two of the K_4 , and the three pendant vertices, so $f(G_{10}) \leq 7$; the seven shaded vertices in Figure 2 induce the forest $t_2 t_3, \ell_2 q_2 q_3 \ell_3, \ell_1$. The degrees of G_{10} are 5 (q_1), 4 (q_2, q_3), 3 (t_1, q_4), 2 (t_2, t_3) and 1 (ℓ_1, ℓ_2, ℓ_3); so $m = 13$, $\deg_{\text{avg}}(G_{10}) = 13/5$, and the degrees of $\overline{G}_{10}$ are 4, 5, 5, 6, 6, 7, 7, 8, 8.




FIGURE 2. The graph G_{10} . The shaded vertices induce a forest of order $7 = f(G_{10})$.

TABLE 1. Counterexamples among the connected graphs with n vertices. A dash means that no graph of that order has an even mode under the reading, or none violates it; the readings are the even mode as defined (first) or as the smallest even most frequent degree (second), each with $\lceil \cdot \rceil$ or $\lfloor \cdot \rfloor$.

n	graphs	first, $\lceil \cdot \rceil$	second, $\lceil \cdot \rceil$	first, $\lfloor \cdot \rfloor$	second, $\lfloor \cdot \rfloor$
2–8	12,112	0	0	0	0
9	261,080	0	0	0	0
10	11,716,571	7	5	0	0

The value 8 occurs three times and every other value at most twice, so it is the even mode under both readings, and the right side is $2\lceil 8 \cdot 5/13 \rceil = 2\lceil 40/13 \rceil = 8$. $\square$

With $\lfloor \cdot \rfloor$ the right side for G_{10} is $2\lfloor 40/13 \rfloor = 6$, and the inequality holds.

Theorem 1.2 comes from an exhaustive search. The connected graphs with at most 10 vertices were generated with **geng** from **nauty** [7], and for each of them the four right sides were computed in integer arithmetic (the quotient $x/\deg_{\text{avg}}(G)$ equals $xn/(2m)$) and compared with the forest number, computed by branch and bound over vertex subsets. Table 1 gives the results. The seven counterexamples with 10 vertices all have 13 edges, three pendant vertices and forest number 7; their graph6 codes are I?ACJ@PqW (this is G_{10}), I?AACJaf0, I?AACIdu_, I?AADJaf0, I?AADHQbW, I?ABAeK0w and I?`CR@PoW, and the last two violate only the first reading of the even mode.

APPENDIX A. VERIFICATION

The proofs above use no computation. The computer checks the finite claims and the search of Section 3, independently in C, C++, Bash, Python, Julia and Lean; all files are in the repository listed under Data and code availability, where **scripts/run_all.sh** runs them.

- (1) **C. wowii66.c** performs the search of Section 3 on graph6 input, and for $c \leq 300$ builds H_c and checks Lemmas 2.1–2.3: the degrees, the even modes and both right sides, the partition into cliques and the explicit forest of order $2c$, and $f(H_c) = 2c$ by branch and bound for $c \leq 6$.
- (2) **C++. wowii66.cpp** repeats both computations with no shared code and a different method for f (a greedy lower bound, and an exact value over all vertex subsets only

- when that bound is below the right side). Its output for all connected graphs with at most 10 vertices is identical, line for line, to that of the C program.
- (3) **Bash** (shell integer arithmetic only): G_{10} , with $f(G_{10}) = 7$ over all 1024 vertex subsets; H_c for $c \leq 40$; and the inequalities of Lemma 2.3 for $c \leq 20000$.
 - (4) **Python** (exact rational arithmetic): G_{10} and the seven graphs of Table 1, with f over all vertex subsets and a colour-refinement check that the seven are pairwise non-isomorphic; every labelled connected graph with at most 6 vertices; H_c for $c \leq 60$, with f over all subsets for $c \leq 4$; and the inequalities of Lemma 2.3 for $c \leq 10^5$.
 - (5) **Julia** (exact rational arithmetic): the same graphs, H_c for $c \leq 300$, and $f(H_c) = 2c$ over all 2^{4c} vertex subsets for $c \leq 5$.
 - (6) **Lean 4** [8]: a self-contained file without Mathlib computes, in the kernel, every quantity of Proposition 3.1 from the edge list of G_{10} , including $f(G_{10}) = 7$ over all 1024 vertex subsets, and proves that Conjecture 66 fails for G_{10} under both readings; the proof depends only on the axiom `propext`.
 - (7) **Lean 4 with Mathlib** [6]: a second file defines the forest number, both readings of the even mode and the average degree for every finite simple graph, and proves for every c that H_c is connected, that $f(H_c) \leq 2c$, the degrees of $\overline{H_c}$, both even modes and both right sides. It concludes that H_c violates Conjecture 66 under both readings for every $c \geq 8$, and the version with $\lfloor \cdot \rfloor$ for every $c \geq 14$. The proof uses only the standard axioms `propext`, `Classical.choice` and `Quot.sound`.

STATEMENTS AND DECLARATIONS

Funding. No funding was received for conducting this study.

Competing interests. The author has no competing interests to declare that are relevant to the content of this article.

Data and code availability. The programs, their outputs and the Lean files are available at <https://github.com/creelie/eConjectureWOWII66>.

Use of generative AI. The author used Claude (Anthropic) for programming, computation and drafting; the author checked the mathematics and takes full responsibility for the content.

REFERENCES

- [1] N. Alon, J. Kahn and P. D. Seymour, *Large induced degenerate subgraphs*, Graphs Combin. **3** (1987), 203–211. <https://doi.org/10.1007/BF01788542>
- [2] D. Bhattacharjee, P. Mandal and U. Bhattacharya, *Resolving Erdős–Ulam monochromatic union-closed family conjectures*, preprint, 2026. [arXiv:2610.02833](https://arxiv.org/abs/2610.02833)
- [3] E. DeLaViña, *Graffiti.pc: a variant of Graffiti*, in: Graphs and Discovery, DIMACS Ser. Discrete Math. Theoret. Comput. Sci. 69, Amer. Math. Soc., Providence, RI, 2005, 71–79. <https://doi.org/10.1090/dimacs/069/05>
- [4] E. DeLaViña, *Written on the Wall II (conjectures of Graffiti.pc)*, <http://cms.uhd.edu/faculty/delavinae/research/wowII/>; version of 26 July 2026 archived at <https://web.archive.org/web/20260726061534/http://cms.uhd.edu/faculty/delavinae/research/wowII/all.html>
- [5] S. Fajtlowicz, *On conjectures of Graffiti*, Discrete Math. **72** (1988), 113–118. [https://doi.org/10.1016/0012-365X\(88\)90199-9](https://doi.org/10.1016/0012-365X(88)90199-9)
- [6] The mathlib Community, *The Lean mathematical library*, in: Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), ACM, New York, 2020, 367–381. <https://doi.org/10.1145/3372885.3373824>
- [7] B. D. McKay and A. Piperno, *Practical graph isomorphism, II*, J. Symbolic Comput. **60** (2014), 94–112. <https://doi.org/10.1016/j.jsc.2013.09.003>

- [8] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in: Automated Deduction – CADE 28, Lecture Notes in Comput. Sci. 12699, Springer, Cham, 2021, 625–635.
https://doi.org/10.1007/978-3-030-79876-5_37

FORMERLY, ELECTRO-GRAVITATIONAL SPACE PROPULSION LABORATORY (EGSPL),
BHUBANESWAR, ODISHA 751030, INDIA

Email address: `d.bhattacharjee@erl-forschung.de`

Email address: `itsdeep@live.com`

URL: <https://orcid.org/0000-0003-0466-750X>