The Welch function is not third-order sum-free

Deep Bhattacharjee*
Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL),
Bhubaneswar, Odisha 751030, India
Formerly, University of Calcutta, Senate House, 87/1 College Street,
Kolkata 700073, West Bengal, India

26 October 2026

Abstract

Let $n = 2m + 1$ be odd and let $\mathbb{F}_{2^n}$ be the field with 2^n elements. A function on $\mathbb{F}_{2^n}$ is third-order sum-free when its values over any eight elements forming a three-dimensional affine subspace never add up to zero. The Welch function is the power map $x \mapsto x^{2^m+3}$; it is one of the classical almost perfect nonlinear functions, and Hou and Zhao conjectured that for $n \geq 7$ it is not third-order sum-free. We prove this. The question comes down to a linearised polynomial $g(x, y)$ in y whose roots form a vector space $Z(x)$ over $\mathbb{F}_2$ containing 1 and x, and the Welch function fails to be third-order sum-free exactly when $Z(x)$ has dimension three for some x outside $\mathbb{F}_2$. We show that the dimension is always 2 or 3, and that it is 3 if and only if two explicit trace conditions hold: with $u = x^{2^m}$, both $\mathrm{Tr}(1/(x + u))$ and $\mathrm{Tr}(xu/(x + u))$ equal 1. The proof applies the field automorphism $z \mapsto z^{2^{m+1}}$, whose square is the Frobenius map, to turn g into an additive polynomial of degree eight whose roots always lie in $\mathbb{F}_{2^n}$. Counting the elements that satisfy both trace conditions gives exactly $(2^n + 1 - 3K_n)/4$ of them, where K_n is the binary Kloosterman sum of $\mathbb{F}_{2^n}$, and Weil's bound $|K_n| \leq 2^{n/2+1}$ makes this count positive for every $n \geq 7$. The Welch function sums to zero over exactly $(2^n+1-3K_n)(2^n-1)/168$ three-dimensional linear subspaces, and the count $2 + (2^n + 1 - 3K_n)/4$ agrees with the numbers that Hou and Zhao computed for $n \leq 15$. Every statement is proved by hand.

Keywords. sum-free function; Welch function; almost perfect nonlinear function; Kloosterman sum.

MSC 2020. 11T06; 11T23; 94A60; 06E30.

Note. The results of this paper were first made public on 9 October 2026, as release `v1.0.0` of the repository `https://github.com/creelie/eConjectureWelchSumFree`; later versions (`v1.1.0` of 22 October and `v1.2.0` of 26 October) are archived on Zenodo as 10.5281/zenodo.23553010 and 10.5281/zenodo.23662272 (concept DOI 10.5281/zenodo.23553009). An earlier and incomplete version of the paper appeared on Hexagon on 22 October 2026 as the preprint `hexagon:2610.00472`, available at `https://hexagonmath.org/2610.00472`. That version cited the conjecture and the corollary of Hou and Zhao as Conjecture 4.10 and Corollary 4.9, numbers that do not match their paper, listed the programs that check its statements in the text, and had a short abstract. This version is complete and self-contained: the references follow the arXiv paper of Hou and Zhao (Conjecture 5.10, Corollary 5.9, Section 5.3), every reference has been checked at its source, the abstract states what is proved, and every statement is

*Corresponding author. Email: d.bhattacharjee@erl-forschung.de, itsdeep@live.com, egspl@proton.me. ORCID: 0000-0003-0466-750X.

proved by hand, so that no statement rests on a computer. The source of this revision is in release `v2.0.0` of the same repository, archived as 10.5281/zenodo.23663217, with programs that recompute Table 1 and check the criterion on small fields.

1 Introduction

A function $f\colon \mathbb{F}_{2^n} \to \mathbb{F}_{2^n}$ is *kth order sum-free* if $\sum_{a\in A} f(a) \neq 0$ for every k-dimensional affine subspace A of $\mathbb{F}_{2^n}$ over $\mathbb{F}_2$. Carlet introduced this notion [6, 5]; for $k = 2$ it is the APN property. Throughout, $n = 2m + 1$ is odd, $q = 2^n$, $F = \mathbb{F}_q$ and $\mathrm{Tr}\colon F \to \mathbb{F}_2$ is the absolute trace. The *Welch function* is

$$W_n(X) = X^{2^m+3}.$$

It is APN [8], and decimating a binary m-sequence by $2^m + 3$ gives a three-valued cross-correlation, as Welch conjectured [4]. For $m \geq 2$ it has algebraic degree 3, so the first open question about its higher-order sum-freedom is the third order. For $m = 2$ the Welch function is $X^7 = X^{2^3-1}$, which is third-order sum-free by a result of Carlet [6, Proposition 1]. Hou and Zhao [9] reduced the question to the rank of a Dickson matrix, computed the answer for $5 \leq n \leq 15$, and proposed the following statement [9, Conjecture 5.10].

Conjecture 5.10 of Hou and Zhao. *For $n = 2m + 1 \geq 7$, the Welch function on $\mathbb{F}_{2^n}$ is not third-order sum-free.*

We prove this conjecture. The reduction of Hou and Zhao goes through the polynomial

$$g(X, Y) = Y^{2^m}(X^2 + X) + Y^2(X^{2^m} + X) + Y(X^{2^m} + X^2). \tag{1.1}$$

For $x \in F$ let $Z(x) = \{y \in F : g(x, y) = 0\}$. Since $g(x, Y)$ is an additive polynomial in Y, $Z(x)$ is a vector space over $\mathbb{F}_2$, and it contains 1 and x. By Lemma 2.3 below, W_n fails to be third-order sum-free exactly when $\dim_{\mathbb{F}_2} Z(x) \geq 3$ for some $x \in F \setminus \mathbb{F}_2$. Our main result decides this dimension for every x.

Theorem 1.1. *Let $n = 2m + 1 \geq 3$ and $x \in F \setminus \mathbb{F}_2$, and put $u = x^{2^m}$. Then $\dim_{\mathbb{F}_2} Z(x) \in \{2, 3\}$, and $\dim_{\mathbb{F}_2} Z(x) = 3$ if and only if*

$$\mathrm{Tr}\Big(\frac{1}{x+u}\Big) = 1 \qquad \text{and} \qquad \mathrm{Tr}\Big(\frac{xu}{x+u}\Big) = 1.$$

Let

$$K_n = \sum_{z\in F^*} (-1)^{\mathrm{Tr}(z+z^{-1})}$$

be the Kloosterman sum of F. The conditions of Theorem 1.1 turn the count of the elements x with $\dim Z(x) = 3$ into Kloosterman sums.

Theorem 1.2. *Let $n = 2m + 1 \geq 3$, and let N_n be the number of $x \in F \setminus \mathbb{F}_2$ with $\dim_{\mathbb{F}_2} Z(x) = 3$. Then*

$$N_n = \frac{2^n + 1 - 3K_n}{4}.$$

By Weil's bound [11] (see also [10, Chapter 5]), $|K_n| \leq 2 \cdot 2^{n/2}$. Hence

$$N_n \geq \frac{2^n + 1 - 6 \cdot 2^{n/2}}{4} = \frac{(2^{n/2} - 3)^2 - 8}{4} > 0 \qquad \text{for } n \geq 7.$$

Corollary 1.3. *The conjecture of Hou and Zhao is true. More precisely, for $n = 2m+1 \geq 5$ the Welch function sums to zero over exactly*

$$\frac{N_n\,(2^n-1)}{42}$$

three-dimensional linear subspaces of F, and over 2^{n-3} times as many three-dimensional affine subspaces. This number is positive for every $n \geq 7$ and is zero for $n = 5$.

By a theorem of Carlitz [7] (see also [10, Chapter 5]), $K_n = -(\omega^n + \bar{\omega}^n)$ with $\omega = (-1+\sqrt{-7})/2$, because $K_1 = 1$. So $K_0 = -2$, $K_1 = 1$ and $K_n = -K_{n-1} - 2K_{n-2}$, which gives Table 1. Hou and Zhao [9, Corollary 5.9] showed that $\dim Z(x) = 3$ for some $x \in F \setminus \mathbb{F}_2$ if and only if the polynomial $\det D^*(X)$ built from a Dickson matrix has more than two roots in F; the number of its roots in F is $2 + N_n$. For $5 \leq n \leq 15$ these are the values Hou and Zhao computed [9, Table 1].

Table 1: The Kloosterman sum K_n, the count N_n of Theorem 1.2 and the number $2 + N_n$ of roots of $\det D^*(X)$ in $\mathbb{F}_{2^n}$.

n	K_n	N_n	$2+N_n$
5	11	0	2
7	−13	42	44
9	−5	132	134
11	67	462	464
13	−181	2184	2186
15	275	7986	7988
17	−101	32844	32846
19	−797	131670	131672
21	2795	522192	522194

The proof of Theorem 1.1 uses the automorphism $\rho(z) = z^{2^{m+1}}$ of F, whose square is the Frobenius map $z \mapsto z^2$ and whose inverse is $z \mapsto z^{2^m}$. Applying ρ to the equation $g(x,y) = 0$ and eliminating y^{2^m} produces an additive polynomial of degree 8 in y. Its roots form a three-dimensional space V that always lies in F and contains $Z(x)$, and a linear map built from $g(x,\cdot)$ moves V into $Z(x)$. Whether $Z(x) = V$ is then decided by two elements of $\mathbb{F}_2$, and both turn out to be traces of explicit rational functions of x and u, because an element of $\mathbb{F}_2$ equals its own trace when n is odd. The counting of Section 5 is a finite count attached to the action of the Galois group on F, of the same kind as the counts in the author's earlier work [1, 2, 3].

Section 2 collects facts about the trace and the reduction of Hou and Zhao. Section 3 studies a fixed x, Section 4 proves Theorem 1.1, and Section 5 proves Theorem 1.2 and Corollary 1.3.

2 Preliminaries

Lemma 2.1. *Let $\rho(z) = z^{2^{m+1}}$ for $z \in F$.*

(a) $\rho(z^{2^m}) = z = \rho(z)^{2^m}$ and $\rho(\rho(z)) = z^2$ for every $z \in F$.

(b) The elements of F fixed by $z \mapsto z^{2^m}$, and those fixed by ρ, are 0 and 1.

Proof. (a) holds because $z^{2^n} = z$, $2^m \cdot 2^{m+1} = 2^n$ and $2^{m+1} \cdot 2^{m+1} = 2 \cdot 2^n$. For (b), the fixed field of $z \mapsto z^{2^k}$ is $\mathbb{F}_{2^{\gcd(k,n)}}$, and $\gcd(m, 2m+1) = \gcd(m+1, 2m+1) = 1$. □

Let $T_0 = \{w \in F : \mathrm{Tr}(w) = 0\}$, put $\wp(z) = z^2 + z$, and for $z \in F$ let

$$T_m(z) = \sum_{i=0}^{m-1} z^{2^i}.$$

Lemma 2.2. *(a)* $\mathrm{Tr}(z^{2^k}) = \mathrm{Tr}(z)$ *for all* $z \in F$ *and* $k \geq 0$*; in particular* $\mathrm{Tr}(\rho(z)) = \mathrm{Tr}(z)$*.*

(b) $\mathrm{Tr}(1) = 1$*, so* $\mathrm{Tr}(\varepsilon) = \varepsilon$ *for* $\varepsilon \in \mathbb{F}_2$*.*

(c) Each of the maps $z \mapsto \wp(z)$*,* $z \mapsto z^{2^m} + z$ *and* $z \mapsto \rho(z) + z$ *is* $\mathbb{F}_2$*-linear with kernel* $\mathbb{F}_2$ *and image* T_0*.*

(d) $\wp(T_m(z)) = T_m(\wp(z)) = z^{2^m} + z$ *for all* $z \in F$*.*

Proof. (a) is clear and (b) holds because n is odd. In (c), the kernels are $\mathbb{F}_2$ by Lemma 2.1(b), so each image has $q/2$ elements; each image lies in T_0 by (a), and $|T_0| = q/2$. For (d), both sides telescope: $\wp(T_m(z)) = \sum_{i=1}^{m} z^{2^i} + \sum_{i=0}^{m-1} z^{2^i} = z^{2^m} + z$, and $T_m(z^2 + z)$ is the same sum. □

The next lemma is the reduction of Hou and Zhao [9, Section 5.3]; we include the short proof.

Lemma 2.3. *(a) For* $x, y \in F$*,*

$$\sum_{a_1,a_2,a_3 \in \mathbb{F}_2} W_n(a_1 x + a_2 y + a_3) = g(x, y).$$

(b) Let $A = c + A_0$ *be a three-dimensional affine subspace of* F *with linear part* A_0*. Then* $\sum_{a \in A} W_n(a) = \sum_{a \in A_0} W_n(a)$*, and* $\sum_{a \in A_0} W_n(a) = b^{2^m+3} \sum_{a \in b^{-1}A_0} W_n(a)$ *for every* $b \in A_0 \setminus \{0\}$*.*

(c) W_n *is not third-order sum-free if and only if* $\dim_{\mathbb{F}_2} Z(x) \geq 3$ *for some* $x \in F \setminus \mathbb{F}_2$*.*

Proof. (a) Since $a_i^{2^k} = a_i$ for $a_i \in \mathbb{F}_2$,

$$W_n(a_1 x + a_2 y + a_3) = (a_1 x^{2^m} + a_2 y^{2^m} + a_3)(a_1 x^2 + a_2 y^2 + a_3)(a_1 x + a_2 y + a_3).$$

Expanding, we get a sum of terms $a_i a_j a_k \, \alpha\beta\gamma$ with $\alpha \in \{x^{2^m}, y^{2^m}, 1\}$, $\beta \in \{x^2, y^2, 1\}$ and $\gamma \in \{x, y, 1\}$. Summed over $(a_1, a_2, a_3) \in \mathbb{F}_2^3$, a monomial in the a_i gives 0 unless all three of a_1, a_2, a_3 occur in it, because $\sum_{a \in \mathbb{F}_2} 1 = 0$. So only the six terms in which i, j, k are distinct survive, each with sum 1, and they add up to $g(x, y)$.

(b) Write the elements of A as $c + a$ with $a \in A_0$ and expand $W_n(c + a) = (c^{2^m} + a^{2^m})(c^2 + a^2)(c + a)$. Apart from $a^{2^m} a^2 a$, every one of the eight products is a constant times a product of at most two $\mathbb{F}_2$-linear functions of a. In coordinates $a = a_1 e_1 + a_2 e_2 + a_3 e_3$ with respect to a basis of A_0, such a product is a combination of monomials in at most two of the a_i, and its sum over A_0 vanishes as in (a). The second statement holds because W_n is a power function.

(c) By (b), W_n is not third-order sum-free if and only if it sums to zero over a three-dimensional linear subspace that contains 1. Such a subspace is $\langle 1, x, y \rangle$ with $1, x, y$ linearly independent over $\mathbb{F}_2$, and by (a) the sum over it is $g(x, y)$. Since $Z(x)$ always contains $\langle 1, x \rangle$, a suitable y exists if and only if $\dim Z(x) \geq 3$. □

3 A fixed element x

In this section and the next, $x \in F \setminus \mathbb{F}_2$ is fixed and

$$u = x^{2^m}, \qquad t = x^2 + x, \qquad d = x + u, \qquad v = u^2 + u.$$

By Lemma 2.1, $\rho(x) = u^2$ and $\rho(u) = x$, so

$$\rho(t) = v^2, \quad \rho(d) = d + v = x + u^2, \quad \rho(d+v) = d^2, \quad \rho(x^2+u) = u^4 + x, \quad t^{2^m} = v. \tag{3.1}$$

The elements t, $u^2 + u = v$, d and $d + v = x + \rho(x)$ are nonzero by Lemma 2.1(b), and

$$t = d^2 + d + v. \tag{3.2}$$

Write $L(y) = g(x, y)$, so that

$$L(y) = t\, y^{2^m} + d\, y^2 + (x^2 + u)\, y,$$

an $\mathbb{F}_2$-linear map of F with kernel $Z(x)$. One checks directly that $L(1) = L(x) = 0$.

Lemma 3.1. *$L(y) = t\, T_m(\wp(y)) + d\, \wp(y)$ for every $y \in F$.*

Proof. By Lemma 2.2(d), $t\, T_m(\wp(y)) + d\, \wp(y) = t(y^{2^m} + y) + d(y^2 + y) = t\, y^{2^m} + d\, y^2 + (t + d)\, y$, and $t + d = x^2 + u$. □

Let

$$Y(y) = \frac{d\, y^2 + (x^2 + u)\, y}{t}, \qquad P(y) = v^2\, y + (d + v)\, Y(y)^4 + (u^4 + x)\, Y(y)^2,$$

and define $\Phi\colon F \to F$ by $\Phi(y) = \rho(L(y)/t)$. Then Y, P and Φ are $\mathbb{F}_2$-linear, $L(y) = t(y^{2^m} + Y(y))$, and $\Phi(y) = 0$ if and only if $L(y) = 0$.

Lemma 3.2. *$P(y) = \rho\big(L(\Phi(y))\big)$ for every $y \in F$.*

Proof. Let $\ell = L(y)/t$, so that $y^{2^m} = Y(y) + \ell$ and, by Lemma 2.1(a), $\rho(y) = \big(y^{2^m}\big)^2 = (Y(y) + \ell)^2$. Applying ρ to $L(y)$ and using (3.1),

$$\rho(L(y)) = v^2 y + (d + v)\rho(y)^2 + (u^4 + x)\rho(y) = P(y) + (d + v)\ell^4 + (u^4 + x)\ell^2.$$

On the other hand $\rho(L(y)) = \rho(t\ell) = v^2\rho(\ell)$. Hence $P(y) = v^2\rho(\ell) + (d + v)\ell^4 + (u^4 + x)\ell^2$. Now put $\ell' = \Phi(y) = \rho(\ell)$. By Lemma 2.1(a), $\ell'^{\,2^m} = \ell$ and $\rho(\ell') = \ell^2$, so

$$\rho(L(\ell')) = \rho\big(t\ell + d\ell'^2 + (x^2 + u)\ell'\big) = v^2\ell' + (d + v)\ell^4 + (u^4 + x)\ell^2 = P(y). \qquad \square$$

Let $\Pi(y) = y(y+1)(y+x)(y+x+1) = \wp(y)\big(\wp(y) + t\big) = y^4 + (t+1)y^2 + ty$. It is an additive polynomial whose roots are the elements of $\langle 1, x\rangle$. Put

$$\lambda = \frac{(d + v)\, d^4}{t^4}, \qquad c = \frac{t^3 v^2}{d^4(d + v)}, \qquad a = \frac{c}{t^2} = \frac{t\, v^2}{d^4(d + v)}.$$

Lemma 3.3. *As polynomials in y over F, $P(y) = \lambda\, \Pi(y)\big(\Pi(y) + c\big)$. The roots of P in an algebraic closure of F form a three-dimensional $\mathbb{F}_2$-space V that contains $\langle 1, x\rangle$, and $V \setminus \langle 1, x\rangle = \{y : \Pi(y) = c\}$.*

Proof. P is an additive polynomial. Its coefficient of y^8 is $\lambda \neq 0$ and its coefficient of y is $v^2 \neq 0$, since $Y(y)^4$ and $Y(y)^2$ contain only y^8, y^4 and y^2. So P has 8 distinct roots, and they form a three-dimensional $\mathbb{F}_2$-space V. Since $Y(1) = 1$ and $Y(x) = u$,

$$P(1) = v^2 + (d+v) + (u^4 + x) = 0, \qquad P(x) = v^2 x + (d+v)u^4 + (u^4 + x)u^2 = 0,$$

so $\langle 1, x\rangle \subseteq V$. Choose $y_0 \in V \setminus \langle 1, x\rangle$. Then $V = \langle 1, x\rangle \cup (y_0 + \langle 1, x\rangle)$ and, since Π is additive,

$$P(y) = \lambda\, \Pi(y)\, \Pi(y + y_0) = \lambda\, \Pi(y)\big(\Pi(y) + \Pi(y_0)\big).$$

The coefficient of y on the right is $\lambda\, t\, \Pi(y_0)$, because $\Pi(y)^2$ has only even powers of y. Comparing with v^2 gives $\Pi(y_0) = v^2/(\lambda t) = c$. Finally $c \neq 0$, so the roots of $\Pi(y) + c$ are exactly the four elements of V outside $\langle 1, x\rangle$. □

Lemma 3.4. $\mathrm{Tr}(a) = 0$. *If $z \in F$ satisfies $\wp(z) = a$, then $\mathrm{Tr}(tz) = 0$. Consequently $V \subseteq F$.*

Proof. From (3.2) one checks the identity

$$a = \wp\Big(\frac{v}{d^2}\Big) + \frac{1}{d} + \frac{1}{d+v}; \tag{3.3}$$

multiplied by $d^4(d+v)$ it reads $tv^2 = v^2(d+v) + vd^2(d+v) + d^3(d+v) + d^4$, and the right side equals $v^2(d^2 + d + v)$. By (3.1), $1/(d+v) = \rho(1/d)$, so $\mathrm{Tr}(1/(d+v)) = \mathrm{Tr}(1/d)$ and $\mathrm{Tr}(a) = 0$. By Lemma 2.2(c) there are exactly two elements $z \in F$ with $\wp(z) = a$, and they differ by 1; since $\mathrm{Tr}(t) = \mathrm{Tr}(\wp(x)) = 0$, the value $\mathrm{Tr}(tz)$ does not depend on the choice.

Let $h = 1/d$. Since $\mathrm{Tr}\big(h + \mathrm{Tr}(h)\big) = 0$ by Lemma 2.2(b), Lemma 2.2(c) gives $z_2 \in F$ with $\rho(z_2) + z_2 = h + \mathrm{Tr}(h)$. By Lemma 2.1(a),

$$\wp(z_2) = \rho(\rho(z_2)) + z_2 = \rho\big(\rho(z_2) + z_2\big) + \big(\rho(z_2) + z_2\big) = \rho(h) + h = \frac{1}{d+v} + \frac{1}{d},$$

so by (3.3) we may take $z = v/d^2 + z_2$. First, by (3.2),

$$\frac{tv}{d^2} = v + \frac{v}{d} + \frac{v^2}{d^2} = \wp(u) + \wp\Big(\frac{v}{d}\Big),$$

which has trace 0. Second, let $e = d + v = x + \rho(x)$. Then $\rho(e) = d^2$ by (3.1), and $\rho(e) + e = x^2 + x = t$. Using Lemma 2.2(a) and Lemma 2.1(a) to move $z \mapsto z^{2^m}$ and ρ through the trace,

$$\begin{aligned}\mathrm{Tr}(tz_2) &= \mathrm{Tr}(\rho(e)z_2) + \mathrm{Tr}(ez_2) = \mathrm{Tr}\big(e\, z_2^{2^m}\big) + \mathrm{Tr}(ez_2) = \mathrm{Tr}\big(e\,(\rho(z_2) + z_2)^{2^m}\big)\\ &= \mathrm{Tr}\big(\rho(e)\,(\rho(z_2) + z_2)\big) = \mathrm{Tr}\big(d^2(h + \mathrm{Tr}(h))\big) = \mathrm{Tr}(d) + \mathrm{Tr}(h)\,\mathrm{Tr}(d^2) = 0,\end{aligned}$$

since $\mathrm{Tr}(d) = \mathrm{Tr}(x + x^{2^m}) = 0$. Hence $\mathrm{Tr}(tz) = 0$.

Now Lemma 2.2(c) gives $y \in F$ with $\wp(y) = tz$. Then $\Pi(y) = tz(tz + t) = t^2\wp(z) = c$, so $y \in V \setminus \langle 1, x\rangle$ by Lemma 3.3, and $V = \langle 1, x, y\rangle \subseteq F$. □

Lemma 3.5. $V = \{y \in F : L(\Phi(y)) = 0\}$, $Z(x) \subseteq V$ *and* $\Phi(V) \subseteq Z(x)$. *The element $\delta(x) = \Phi(y_0)$ is the same for all $y_0 \in V \setminus \langle 1, x\rangle$ and lies in $\langle 1, x\rangle$. If $\delta(x) = 0$ then $Z(x) = V$; otherwise $Z(x) = \langle 1, x\rangle$.*

Proof. Since $V \subseteq F$ (Lemma 3.4) and ρ is injective, Lemma 3.2 shows that $V = \{y \in F : L(\Phi(y)) = 0\}$. If $L(y) = 0$ then $\Phi(y) = 0$ and $y \in V$, so $Z(x) \subseteq V$; and $\Phi(V) \subseteq Z(x)$ by the same description of V. Two choices of y_0 differ by an element of $\langle 1, x\rangle \subseteq Z(x)$, on which Φ vanishes. As $\langle 1, x\rangle \subseteq Z(x) \subseteq V$ and $\dim V = 3$, either $Z(x) = V$ or $Z(x) = \langle 1, x\rangle$. In the first case $\delta(x) = 0$. In the second case $\delta(x) \in \Phi(V) \subseteq \langle 1, x\rangle$, and $\delta(x) \neq 0$ because $y_0 \notin Z(x)$. □

Lemma 3.6. *$\delta(x) \in \{0,1\}$ if and only if $\mathrm{Tr}(1/d) = 1$.*

Proof. Take $y_0 \in V \setminus \langle 1, x\rangle$ and let $z = \wp(y_0)/t$. By Lemma 3.3, $t^2\wp(z) = \wp(y_0)(\wp(y_0)+t) = c$, so $\wp(z) = a$. Let $\ell_0 = L(y_0)/t$, so that $\delta(x) = \rho(\ell_0)$, and $\delta(x) \in \{0,1\}$ if and only if $\ell_0 \in \{0,1\}$, that is, $\wp(\ell_0) = 0$. By Lemma 3.1, $\ell_0 = T_m(tz) + dz$. By Lemma 2.2(d), (3.1) and (3.2),

$$\wp(\ell_0) = (tz)^{2^m} + tz + d^2z^2 + dz = v\, z^{2^m} + (t+d^2+d)\, z + d^2a = v(z^{2^m}+z) + d^2a = v\, T_m(a) + d^2a.$$

Put $\gamma = d^2a/v$, so that $\wp(\ell_0) = v\,(T_m(a)+\gamma)$. On the other hand $\delta(x) \in \langle 1, x\rangle$ by Lemma 3.5, so $\ell_0 = \delta(x)^{2^m} \in \{0, 1, u, u+1\}$ and $\wp(\ell_0) \in \{0, v\}$. Therefore $\varepsilon = T_m(a) + \gamma$ lies in $\mathbb{F}_2$, and $\varepsilon = 0$ exactly when $\delta(x) \in \{0,1\}$. By Lemma 2.2(b) and (a),

$$\varepsilon = \mathrm{Tr}(\varepsilon) = \sum_{i=0}^{m-1} \mathrm{Tr}(a^{2^i}) + \mathrm{Tr}(\gamma) = m\,\mathrm{Tr}(a) + \mathrm{Tr}(\gamma) = \mathrm{Tr}(\gamma),$$

using $\mathrm{Tr}(a) = 0$ from Lemma 3.4. By (3.2), $\gamma = tv/(d^2(d+v)) = v/d^2 + v/(d+v)$. By (3.1), $\rho(t/d^2) = v^2/(d+v)^2$, so $\mathrm{Tr}(v/(d+v)) = \mathrm{Tr}(t/d^2)$. Since $v + t = d^2 + d$,

$$\mathrm{Tr}(\gamma) = \mathrm{Tr}\Big(\frac{v+t}{d^2}\Big) = \mathrm{Tr}(1) + \mathrm{Tr}\Big(\frac{1}{d}\Big) = 1 + \mathrm{Tr}\Big(\frac{1}{d}\Big).$$

So $\delta(x) \in \{0,1\}$ if and only if $\mathrm{Tr}(1/d) = 1$. □

4 Proof of Theorem 1.1

The objects of Section 3 depend on x; we write L_x, t_x, Φ_x, V_x and $d_x = x + x^{2^m}$ when the dependence matters. The second condition of Theorem 1.1 comes from the substitution $x \mapsto 1/x$.

Lemma 4.1. *For $x \in F \setminus \mathbb{F}_2$ and $y \in F$, $g(1/x, y/x) = x^{-(2^m+3)} g(x,y)$. Moreover $\Phi_{1/x}(y) = x^{-1}\Phi_x(xy)$, $V_{1/x} = x^{-1}V_x$ and $\delta(1/x) = x^{-1}\delta(x)$.*

Proof. With $u = x^{2^m}$,

$$u\,x^3\, g(1/x, y/x) = u\,x^3\Big(\frac{y^{2^m}}{u}\cdot\frac{1+x}{x^2} + \frac{y^2}{x^2}\cdot\frac{x+u}{xu} + \frac{y}{x}\cdot\frac{u+x^2}{x^2u}\Big) = g(x,y).$$

So $L_{1/x}(y) = x^{-(2^m+3)} L_x(xy)$. Since $t_{1/x} = x^{-2} + x^{-1} = t_x/x^3$,

$$\Phi_{1/x}(y) = \rho\Big(\frac{L_x(xy)}{u\,t_x}\Big) = \rho(u^{-1})\,\Phi_x(xy) = x^{-1}\Phi_x(xy).$$

By the first statement of Lemma 3.5, applied to x and to $1/x$, $V_{1/x} = \{y \in F : L_{1/x}(\Phi_{1/x}(y)) = 0\} = \{y \in F : L_x(\Phi_x(xy)) = 0\} = x^{-1}V_x$. If $y_0 \in V_x \setminus \langle 1, x\rangle$, then $y_0/x \in V_{1/x} \setminus \langle 1, 1/x\rangle$, and $\delta(1/x) = \Phi_{1/x}(y_0/x) = x^{-1}\Phi_x(y_0) = x^{-1}\delta(x)$. □

Proof of Theorem 1.1. By Lemma 3.5, $\dim Z(x) \in \{2,3\}$, and $\dim Z(x) = 3$ if and only if $\delta(x) = 0$. Since $\delta(x) \in \langle 1, x\rangle$ and $x \neq 1$, $\delta(x) = 0$ if and only if $\delta(x) \in \{0,1\}$ and $\delta(x) \in \{0, x\}$. By Lemma 3.6, the first condition is $\mathrm{Tr}(1/d_x) = 1$. By Lemma 4.1, $\delta(x) \in \{0, x\}$ if and only if $\delta(1/x) \in \{0,1\}$, which by Lemma 3.6 applied to $1/x$ means $\mathrm{Tr}(1/d_{1/x}) = 1$. Finally $d_{1/x} = 1/x + 1/u = (x+u)/(xu)$. □

5 Counting

Proof of Theorem 1.2. For $x \in F \setminus \mathbb{F}_2$ let $d = x + x^{2^m}$, $u = x^{2^m}$, $\alpha(x) = (-1)^{\mathrm{Tr}(1/d)}$ and $\beta(x) = (-1)^{\mathrm{Tr}(xu/d)}$. By Theorem 1.1,

$$N_n = \frac{1}{4} \sum_{x \in F \setminus \mathbb{F}_2} (1 - \alpha(x))(1 - \beta(x)) = \frac{1}{4}\Big(q - 2 - \sum \alpha - \sum \beta + \sum \alpha\beta\Big),$$

all sums over $x \in F \setminus \mathbb{F}_2$. By Lemma 2.2(c), $x \mapsto x + x^{2^m}$ maps $F \setminus \mathbb{F}_2$ two-to-one onto $T_0 \setminus \{0\}$. Hence

$$\sum \alpha = 2 \sum_{w \in T_0 \setminus \{0\}} (-1)^{\mathrm{Tr}(1/w)} = \sum_{w \in F^*} (1 + (-1)^{\mathrm{Tr}(w)})(-1)^{\mathrm{Tr}(1/w)} = -1 + K_n,$$

because $w \mapsto 1/w$ permutes F^* and $\sum_{w \in F^*} (-1)^{\mathrm{Tr}(w)} = -1$. The map $x \mapsto 1/x$ permutes $F \setminus \mathbb{F}_2$, and $\beta(x) = \alpha(1/x)$ because $d_{1/x} = d/(xu)$; so $\sum \beta = K_n - 1$. Finally, $x \mapsto x + 1$ permutes $F \setminus \mathbb{F}_2$, leaves d unchanged and sends xu to $(x + 1)(u + 1) = xu + d + 1$; since $\mathrm{Tr}(1) = 1$,

$$\beta(x + 1) = (-1)^{\mathrm{Tr}((xu+1)/d)+1} = -\alpha(x)\beta(x).$$

So $\sum \alpha\beta = -\sum \beta = 1 - K_n$, and

$$N_n = \frac{1}{4}(q - 2 - 2(K_n - 1) + 1 - K_n) = \frac{q + 1 - 3K_n}{4}. \qquad \square$$

Proof of Corollary 1.3. If $\dim Z(x) = 3$, then $A = Z(x)$ is a three-dimensional subspace containing 1 over which W_n sums to zero, by Lemma 2.3(a). Conversely, let $A \ni 1$ be such a subspace and $x' \in A \setminus \mathbb{F}_2$. Writing $A = \langle 1, x', y' \rangle$, Lemma 2.3(a) gives $g(x', y') = 0$, so $A \subseteq Z(x')$, and $Z(x') = A$ by Theorem 1.1. Hence the subspaces containing 1 with zero sum correspond to the elements x with $\dim Z(x) = 3$, six elements to each subspace, and there are $N_n/6$ of them.

Let $\mathcal{S}$ be the set of three-dimensional linear subspaces with zero sum. By Lemma 2.3(b), $A \in \mathcal{S}$ and $b \in F^*$ imply $bA \in \mathcal{S}$. The pairs (A, b) with $A \in \mathcal{S}$ and $b \in A \setminus \{0\}$ number $7|\mathcal{S}|$, and $(A, b) \mapsto (b^{-1}A, b)$ maps them bijectively to the pairs (B, b) with $B \in \mathcal{S}$, $1 \in B$ and $b \in F^*$, which number $(q - 1)N_n/6$. So $|\mathcal{S}| = N_n(q - 1)/42$. By Lemma 2.3(b), an affine subspace has zero sum exactly when its linear part does, and each linear part has 2^{n-3} cosets. The numbers are positive for $n \geq 7$ by the bound after Theorem 1.2, and $N_5 = 0$ because $K_5 = 11$. $\square$

Remark 5.1. For $n = 5$, Corollary 1.3 gives a second proof that $W_5 = X^7$ is third-order sum-free. For $n = 3$ the polynomial $g(x, Y)$ vanishes identically, and indeed $N_3 = 6 = 2^3 - 2$ since $K_3 = -5$.

Declarations

Funding

No funding was received for this work.

Conflict of interest

The author declares no conflict of interest.

Data availability

The proofs in this paper are complete by hand. The source of the paper and programs that recompute Table 1 and check Theorems 1.1 and 1.2 on small fields are in release `v2.0.0` of the repository named in the note under the abstract, archived as 10.5281/zenodo.23663217; the paper does not depend on them.

References

[1] D. Bhattacharjee, Establishing equivariant class $[O]$ for hyperbolic groups, Asian Res. J. Math. 18 (2022), no. 11, 362–369. doi:10.9734/arjom/2022/v18i11615.

[2] D. Bhattacharjee, Non-commutativity over canonical suspension η for genus $g \geq 1$ in hypercomplex structures for potential $\rho\phi$, Asian Res. J. Math. 18 (2022), no. 11, 332–341. doi:10.9734/arjom/2022/v18i11607.

[3] D. Bhattacharjee, U. Bhattacharya, Iterated nerve complexes for k-fold sensor coverage, Preprints.org (2026). doi:10.20944/preprints202606.1351.v2.

[4] A. Canteaut, P. Charpin and H. Dobbertin, Binary m-sequences with three-valued crosscorrelation: a proof of Welch's conjecture, IEEE Trans. Inform. Theory 46 (2000), no. 1, 4–8. doi:10.1109/18.817504.

[5] C. Carlet, On the vector subspaces of $\mathbb{F}_{2^n}$ over which the multiplicative inverse function sums to zero, Des. Codes Cryptogr. 93 (2025), 1237–1254. doi:10.1007/s10623-024-01531-6.

[6] C. Carlet, Two generalizations of almost perfect nonlinearity, J. Cryptology 38 (2025), no. 2, Paper No. 20. doi:10.1007/s00145-025-09538-5.

[7] L. Carlitz, Kloosterman sums and finite field extensions, Acta Arith. 16 (1969), 179–194. doi:10.4064/aa-16-2-179-194.

[8] H. Dobbertin, Almost perfect nonlinear power functions on $GF(2^n)$: the Welch case, IEEE Trans. Inform. Theory 45 (1999), no. 4, 1271–1275. doi:10.1109/18.761283.

[9] X. Hou and S. Zhao, Further results on sum-freedom of binary and q-ary functions, preprint, arXiv:2609.31489 (2026).

[10] R. Lidl and H. Niederreiter, Finite Fields, 2nd ed., Encyclopedia Math. Appl. 20, Cambridge University Press, Cambridge, 1997. doi:10.1017/CBO9780511525926.

[11] A. Weil, On some exponential sums, Proc. Nat. Acad. Sci. U.S.A. 34 (1948), no. 5, 204–207. doi:10.1073/pnas.34.5.204.