A SIXTEEN-PRIME-FACTOR BOUND FOR LEHMER'S TOTIENT PROBLEM AND EXACT PRIME-COMPLETION CRITERIA

DEEP BHATTACHARJEE

ABSTRACT. We study composite integers n for which Euler's totient divides $n-1$. The finite-search results give at least sixteen distinct prime factors, at least 18001 when $(n-1)/\varphi(n) \geq 3$, and more than 10^9 when $3 \mid n$. A solution with k prime factors satisfies $n < 2^{2^{k-7}}$, and every solution exceeds 10^{35}. For the companion divisibility $\varphi(n) \mid n+1$, the nine known solutions exhaust the cases with at most eight prime factors. For a fixed prime prefix $A = \prod p$, $B = \prod(p-1)$, the quotient is forced to be $M = \lceil A/B \rceil$ and a possible largest prime is uniquely $q = 1 + (A-1)/(MB-A)$. We derive its exact divisibility, coprimality and valuation consequences, and prove that deleting the largest prime of a hypothetical counterexample cannot yield a smaller Lehmer counterexample. A two-prime factorisation identifies the additional integrality and primality conditions required by a replacement construction. A complete finite construction is proved for every prescribed factor count. Excluding its output at every factor count, or excluding the final prime for every prefix, remains unproved; the full Lehmer conjecture is not resolved.

1. INTRODUCTION

1.1. **Two questions of Lehmer.** Every prime p satisfies $\varphi(p) = p-1$, and so $\varphi(p)$ divides $p-1$. In 1932 Lehmer [19] asked whether any composite number n has the same property,

$$\varphi(n) \mid n-1. \tag{1.1}$$

Nobody has found one, and nobody has shown that none exists. Lehmer proved that a composite solution of (1.1) must be odd, squarefree and divisible by at least seven distinct primes. The question is Problem B37 in Guy's collection [16]. Lehmer's paper also lists solutions of the companion equation[1]

$$\varphi(n) \mid n+1. \tag{1.2}$$

The two equations differ only in a sign. Equation (1.1) has no known composite solution, whereas (1.2) has the solutions

$$1,\ 2,\ 3,\ 15,\ 255,\ 65535,\ 83623935,\ 4294967295,\ 6992962672132095, \tag{1.3}$$

and no others are known. Except for 2, each of them satisfies $n+1 = 2\varphi(n)$. Five of them are products of the first Fermat primes[2], namely 3, $15 = 3\cdot 5$, $255 = 3\cdot 5\cdot 17$, $65535 = 3\cdot 5\cdot 17\cdot 257$ and

Date: 27 October 2026.

2020 *Mathematics Subject Classification.* Primary 11A25; Secondary 11A41, 11A51, 11D68, 11N36, 11Y11, 11Y16, 11Y70.

Key words and phrases. Lehmer's totient problem, Euler's totient function, exact completion, prime factors, exhaustive search, primality certificates.

Research revision prepared 27 October 2026. Earlier confirmed archive: doi:10.5281/zenodo.23638948. Original preprint: hexagon:2610.00454v1. The full Lehmer conjecture remains unproved.

[1]The comments to sequence A050474 of the OEIS [30] record this, and a misprint in Lehmer's summary where $3\cdot 5\cdot 353\cdot 929$ stands for $83623935 = 3\cdot 5\cdot 17\cdot 353\cdot 929$.

[2]The Fermat numbers $F_i = 2^{2^i}+1$ are prime for $0 \leq i \leq 4$, and $F_0F_1\cdots F_{r-1} = 2^{2^r}-1$. For $1 \leq r \leq 5$ the number $n = 2^{2^r}-1$ is therefore a product of primes F_i, and $\varphi(n) = 2^{1+2+\cdots+2^{r-1}} = 2^{2^r-1} = (n+1)/2$. The chain breaks at $r=6$ because $F_5 = 641\cdot 6700417$, as Euler found.

$4294967295 = 3{\cdot}5{\cdot}17{\cdot}257{\cdot}65537$. Apart from 1 and 2, the other two are $83623935 = 3{\cdot}5{\cdot}17{\cdot}353{\cdot}929$ and $6992962672132095 = 83623935 \cdot 83623937$. These numbers form sequence A203966 of the OEIS [30].

Much of the literature on (1.1) concerns the number $\omega(n)$ of distinct prime factors of a composite solution. Pomerance [33] proved that a composite solution with k prime factors satisfies $n < k^{2^k}$, so that for each k the question is a finite one, and Burek and Żmija [7] sharpened this to $n \le 2^{2^k} - 2^{2^{k-1}}$, using a lemma that goes back to Heath-Brown's work on odd perfect numbers [18] and was sharpened by Cook [10] and Nielsen [28].[3] In binary, their bound allows n one digit for each subset of the set of its prime factors. Cohen and Hagis [8] settled all $k \le 13$ and obtained $\omega(n) \ge 14$. When $3 \mid n$ much more is true: Hagis [17] proved $\omega(n) \ge 298848$, and Burcsi, Czirbusz and Farkas [6] raised this to $\omega(n) \ge 40000000$. In the same paper they report that a computation of Renze [35], made available as a Mathematica notebook, gives $\omega(n) \ge 15$ in general, and Pinch [32] outlined a proof of the same bound in a poster presented at ANTS VII, which also reports that a composite solution exceeds 10^{30}. Neither account has been refereed. Since $p - 1 \mid n - 1$ for every prime p dividing a solution, a composite solution of (1.1) would be a Carmichael number;[4] Banks, Nevans and Pomerance [3] discuss this and the connection with Giuga's conjecture.[5]

More is known when the quotient $M = (n-1)/\varphi(n)$ exceeds 2. Hagis [17] proved $\omega(n) \ge 1991$ for $M = 3$, and Grytczuk and Wójtowicz [15] gave lower bounds for $\omega(n)$ that grow exponentially with $M \ge 4$ (see [39] for both). The prime factors of a solution form what Golomb [14] called a G-sequence: none of them divides another minus one. Meijer [23], who points out the connection with (1.1), proved with the Brun–Selberg sieve that the product of $p/(p-1)$ over the elements of a G-sequence up to x is at most a constant times $\log\log x$. Yamada [39] made this explicit with the large sieve and deduced that $M < 16.03235 \log\log \omega(n)$ for every composite solution of $\varphi(n) \mid n \pm 1$ with $\omega(n) \ge 4$, so that $\omega(n)$ is at least doubly exponential in M.

Norman [29] claimed that $M \ge 3$ for every composite solution of (1.1), and hence $\omega(n) \ge 1991$. The proof rests on [29, Lemma 2.1], which asserts that $\psi(G)/|G|^2 > \varphi(m)/(2m)$ for every noncyclic nilpotent group G of order $4m$ with m odd and squarefree, where $\psi(G)$ is the sum of the orders of the elements of G. The lemma is false. For $G = C_2 \times C_2 \times C_m$ with m prime to 3 we have $\psi(G) = 7\prod_{p\mid m}(p^2 - p + 1)$, since ψ is multiplicative over factors of coprime order, and the ratio of $\psi(G)/|G|^2$ to $\varphi(m)/(2m)$ is $\frac{7}{8}\prod_{p\mid m}(1 + \frac{1}{p(p-1)})$. The product over all primes $p \ge 5$ is $\frac{4}{7}\zeta(2)\zeta(3)/\zeta(6) = 1.1106\ldots < \frac{8}{7}$, so the ratio is less than 1; for $m = 5$ the two sides are $147/400$ and $2/5$. The proof of the lemma uses $\psi(V)/16 \ge \frac{1}{2}$ for the Sylow 2-subgroup V, whereas $\psi(C_2 \times C_2) = 7$. The claim $M \ge 3$ is therefore unproved; if it holds, Theorem 1.1 below gives $\omega(n) \ge 18001$ for every composite solution.

Less is known about (1.2). The comments to sequences A050474 and A203966 of the OEIS [30] report that the M.Sc. thesis of Wong [38] shows that a further solution of $n + 1 = 2\varphi(n)$ would need at least eight prime factors, that there are no further solutions of $n + 1 = 2\varphi(n)$ below 10^{31} and none of (1.2) below $1.6 \cdot 10^{30}$ (Alekseyev), and that $(n + 1)/\varphi(n) = 3$ would

[3] Heath-Brown proved that an odd perfect number with k distinct prime factors is less than 4^{4^k}, and Nielsen lowered this to 2^{4^k}. The lemma of Cook and Nielsen concerns integers $x_1 \le \cdots \le x_m$ for which $\prod(1 - 1/x_i)$ lies just below a given fraction; we state and sharpen it in Section 6.1.

[4] By Korselt's criterion a composite n satisfies $a^{n-1} \equiv 1 \pmod n$ for every a prime to n exactly when n is squarefree and $p - 1 \mid n - 1$ for every prime $p \mid n$. A solution of (1.1) is squarefree by Lemma 2.1, and $p - 1 \mid \varphi(n) \mid n - 1$.

[5] Giuga conjectured in 1950 that an integer $n > 1$ with $1^{n-1} + 2^{n-1} + \cdots + (n-1)^{n-1} \equiv -1 \pmod n$ is prime. He showed that a composite n satisfies this congruence exactly when $p \mid n/p - 1$ and $p - 1 \mid n/p - 1$ for every prime $p \mid n$; the second condition makes n a Carmichael number.

force at least 33 prime factors. None of this has appeared in a refereed paper, and we know of no refereed bound on $\omega(n)$ for (1.2).

1.2. **Results.** We write $\omega(n)$ for the number of distinct prime factors of n. Our first theorem goes one prime further than the bound announced by Renze and by Pinch, and its proof includes a complete proof of that bound.

Theorem 1.1. *Let n be composite with $\varphi(n) \mid n-1$. Then $\omega(n) \geq 16$. If $(n-1)/\varphi(n) \geq 3$, then $\omega(n) \geq 18001$, and if $3 \mid n$, then $\omega(n) > 10^9$.*

Apart from Norman's claim, we know of no earlier claim beyond fifteen prime factors in general. The second assertion raises the bound 1991 of Hagis for the quotient 3 to 18001, and for $3 \nmid n$ it also improves the bounds of Grytczuk and Wójtowicz for the quotients from 4 to 7. For $3 \mid n$ the third assertion raises the bound of Burcsi, Czirbusz and Farkas from 40000000 to more than 10^9. Our second theorem bounds the size of a solution in terms of the number of its prime factors, and the third bounds it from below.

Theorem 1.2. *Let n be composite with $\varphi(n) \mid n-1$, let $k = \omega(n)$, and put $M = (n-1)/\varphi(n)$.*

(i) $n < 2^{2^{k-7}}$.

(ii) *If $M \geq 3$, then* $n < 2^{2^{k-17981}}$.

(iii) *If $3 \mid n$, then* $n < 2^{2^{k-10^9}}$.

Theorem 1.3. *Let n be composite with $\varphi(n) \mid n-1$. Then $n > 10^{35}$.*

Part (i) of Theorem 1.2 divides the number of binary digits allowed by the bound of Burek and Żmija by $2^7 = 128$. Together with Theorems 1.1 and 1.3 it shows that a solution with sixteen prime factors, the first case not yet excluded, lies between 10^{35} and $2^{512} < 1.35 \cdot 10^{154}$. Theorem 1.3 raises the bound 10^{30} reported in Pinch's poster; it is proved in Section 5.3 by the search of Section 3 with one additional rule, which controls the size of n at every node.

The other results concern (1.2).

Theorem 1.4. *The solutions of $\varphi(n) \mid n+1$ with $\omega(n) \leq 8$ are exactly the nine numbers (1.3).*

Theorem 1.5. *Every solution of $\varphi(n) \mid n+1$ with $n > 3$ and $3 \nmid n$ has $\omega(n) \geq 16$.*

Theorem 1.6. *Let $n > 3$ satisfy $\varphi(n) \mid n+1$ with $(n+1)/\varphi(n) \geq 3$. Then $\omega(n) \geq 18001$, and $\omega(n) > 10^9$ if $3 \mid n$.*

The next theorem explains the list (1.3). Say that n is a *Fermat-type* solution if $n+1 = 2\varphi(n)$; by convention $n = 1$ is one.

Theorem 1.7. *Let n_0 be a Fermat-type solution.*

(i) *For a prime $q \nmid n_0$, the number $n_0 q$ is a Fermat-type solution if and only if $q = n_0 + 2$.*

(ii) *For primes $p < q$ not dividing n_0, the number $n_0 pq$ is a Fermat-type solution if and only if*

$$(p - n_0 - 1)(q - n_0 - 1) = n_0^2 + n_0 + 1.$$

(iii) *The Fermat-type solutions in (1.3), which are all of them except 2, are exactly the numbers obtained from 1 by repeated use of* (i) *and* (ii)*: every such extension of one of them is again in (1.3).*

Theorem 1.4 contains the bound reported from Wong's thesis and goes one prime factor further: a solution of (1.2) outside (1.3) has at least nine prime factors, whatever the quotient $(n+1)/\varphi(n)$. Theorem 1.6 raises the 33 prime factors of the observation on the quotient 3 recorded in the OEIS to 18001. Part (iii) of Theorem 1.7 places 83623935 in the same family as

the products of Fermat primes: it is the two-prime extension of 255 coming from the factorisation $255^2 + 255 + 1 = 65281 = 97 \cdot 673$, just as 4294967295 comes from the trivial factorisation $65281 = 1 \cdot 65281$ (Figure 15).

The last result of this section is not new. It follows from the bound of Meijer [23], and Yamada [39] proved it with $c = 1/16.03235$ when $\omega(n) \geq 4$. We give a short proof by the large sieve in Section 2.2, because the estimate behind it explains the size of the bounds in Theorems 1.1 and 1.6.

Theorem 1.8 (Meijer, Yamada)**.** *There is an absolute and effective constant $c > 0$ with the following property. Let n be composite with $\varphi(n) \mid n-1$, or let $n > 3$ with $\varphi(n) \mid n+1$, and let M be the quotient $(n-1)/\varphi(n)$ or $(n+1)/\varphi(n)$. Then*

$$\omega(n) \geq \exp(\exp(cM)) - 2.$$

The first primes alone give only $\omega(n) \geq \exp(c'M)$, since the product of $p/(p-1)$ over the first K primes grows like $\log K$. Theorem 1.8 uses the fact that no prime factor of a solution divides another prime factor minus one, and it is proved by the large sieve (Section 2.2). It explains why the bounds 18001 and 10^9 in Theorems 1.1 and 1.6 lie so far above the bounds 33 and 1540 that the first primes give.

1.3. **Theory and computation.** Every search in this paper rests on a theorem, proved by hand, that reduces the statement to finitely many cases and shows that the search covers all of them. The results divide as follows.

Proved entirely by hand. Theorem 1.8 and Meijer's bound (Theorem 2.7), by the large sieve; the sharp product lemma (Theorem 6.2) and the size bound $n < 2^{2^{k-2}}$ (Proposition 6.11); the reductions of Section 2, up to the evaluation of the explicit finite products in Lemmas 2.2 and 2.3; parts (i) and (ii) of Theorem 1.7; and the equivalence of Proposition 9.1.

Proved by hand up to a finite computation.

- $\omega(n) \geq 16$ in Theorem 1.1, and Theorem 1.5: the interval bounds, the divisibility condition for the last two primes and the completeness of the search (Propositions 3.2 and 3.5, Theorem 3.6, Corollary 3.3), and the reduction of the last three primes to divisors in a residue class (Proposition 4.1, Theorem 4.4, Proposition 4.5), leave a search over at most fifteen primes, which ends in 33865004 two-prime problems for fifteen (Section 5).
- The bounds 18001 and 10^9 in Theorems 1.1 and 1.6: Corollary 2.6 reduces them to the branch-and-bound of Proposition 2.5.
- Theorem 1.3: the size rule of Proposition 5.1 leaves the trees of Table 4, with 1 381 783 982 two-prime problems in all (Section 5.3).
- Theorem 1.2: Corollary 6.10 reduces part (i) to the tree $\mathcal{T}_7$ of 33678 nodes (Theorem 6.15), and parts (ii) and (iii) to the evaluation of two explicit rational numbers.
- Theorem 1.4: the same reductions leave 228260561 two-prime problems for eight primes (Section 7.4).
- Theorem 1.7(iii): the factorisation of eight integers.

Every search was carried out by at least two programs written independently, and every count reported here is the same for all of them (Section 8). A search that finds nothing proves something only if it would have found what it looks for, and the programs were therefore also run on equations that have solutions.

1.4. **Scope.** Theorems 1.1–1.8 bound the number of prime factors and the size of a new solution from both sides and describe the known ones; they do not decide whether (1.1) has a composite solution or whether (1.2) has solutions beyond (1.3). Section 9 estimates the cost of the next cases and states the assertion, equivalent to Lehmer's conjecture, that this paper does not prove.

1.5. **Method.** The bound $\omega(n) \geq 16$ of Theorem 1.1 and Theorems 1.4 and 1.5 rest on three reductions proved in Sections 2 to 4. The first (Section 2) shows that one may assume $n + \varepsilon = 2\varphi(n)$, with $\varepsilon = -1$ for Lehmer's equation and $\varepsilon = +1$ for the companion, and, except in Theorem 1.4, that $3 \nmid n$. The second (Section 3) shows that once the first j prime factors are fixed, the next one lies in an interval whose ends are explicit functions of them (Proposition 3.2), that for each k only finitely many sequences of primes satisfy these conditions (Corollary 3.3), and that the last two prime factors are determined by a divisibility condition (Proposition 3.5). Each value of k is therefore a finite problem, and the traversal of an explicit finite tree finds every solution (Theorem 3.6). The bounds for the two signs of ε differ only by terms of size $1/\varphi(n)$; for $p_1 \geq 5$ they give identical trees for every $k \leq 14$, and for $k = 15$ the same nodes down to depth 13 (Figure 8). The third reduction (Section 4) treats the last three primes together: once p_{k-2} is fixed, the last two correspond to a divisor of an explicit integer N in a known residue class modulo an integer c (Proposition 4.1). Such divisors are the lattice points of a coset near a hyperbola (Lemmas 4.2 and 4.3). Since the two complementary divisors lie in the same class, their sum lies in a single class modulo c, which reduces the problem to one square test for each value of the sum (Section 4.4), and when $c^3 \geq N + 2c^2$ there are at most four such divisors (Proposition 4.5). With these reductions the case $k \leq 14$ becomes a tree of at most 31962 nodes and the case $k = 15$ a set of 33865004 problems of this kind, which we solved by computer. One of the routes never factors an integer, so that for every $k \leq 15$ the conclusion depends on no primality proof.

The companion equation with eight prime factors and $3 \mid n$ behaves differently. After the Fermat primes $3, 5, 17, 257, 65537$ the product $\prod p/(p-1)$ falls short of 2 by only 2^{-31}, the interval for the sixth prime has length 2^{33}, and each of the $1.3 \cdot 10^8$ admissible choices of that prime leaves a two-prime problem whose modulus is far below the cube root of the integer to be split. We treat these problems by the sum route of Section 4.4, combined with trial division for the small divisors and sieved by congruences that primes satisfy, and we settle the 1967265 cases in which this would be slow by factoring the integer completely, with every prime factor proved prime (Section 7.4). Theorem 1.7 rests on a short identity together with the factorisation of eight integers.

The bounds for a quotient of at least 3 and for $3 \mid n$, in Theorems 1.1 and 1.6, come from independence. The prime factors of a solution form a set in which no prime divides another minus one, and over such a set the product of $p/(p-1)$ grows only like the double logarithm of its size (Theorem 2.7). A branch-and-bound over these sets (Proposition 2.5) shows that such a set must be very large before the product reaches 3, as it must when the quotient is at least 3 and $3 \nmid n$, or reaches $\frac{8}{3}$ with all its primes congruent to 2 modulo 3, as it must for the prime factors other than 3 when $3 \mid n$.

Theorem 1.2 is proved in Section 6. Its first ingredient is a sharp form of the lemma of Cook and Nielsen (Theorem 6.2), in which the bound for the product of the remaining prime factors depends on how far the product $\prod p/(p-1)$ over the prime factors already chosen falls short of M. This lemma alone gives $n < 2^{2^{k-2}}$ (Proposition 6.11), and more generally $n < 2^{2^{k-s}}$ for every solution whose first prime factors are small in a precise sense. In every other solution the prime factors grow rapidly, and for $s = 7$ an exhaustive search over their possible initial segments, which visits 33678 of them, shows that there are none. Parts (ii) and (iii) combine the lemma of Cook and Nielsen with Proposition 2.5.

The paper is organised as follows. Section 2 proves the reductions, the bounds for large quotients and for $3 \mid n$, Theorem 1.6, and, by the large sieve, Theorem 1.8. Section 3 proves the interval bounds and the completeness of the search, and Section 4 the reduction of the last three primes to divisors in a residue class. Section 5 completes the proof of Theorem 1.1, Section 6

proves Theorem 1.2, and Section 7 proves Theorems 1.4, 1.5 and 1.7. Section 8 summarises the checks on the computations, and Section 9 states the assertion that remains open.

Notation. Throughout, $n = p_1 p_2 \cdots p_k$ with primes $p_1 < \cdots < p_k$, and $\varepsilon \in \{-1, +1\}$ selects the equation $\varphi(n) \mid n + \varepsilon$; we put $M = (n+\varepsilon)/\varphi(n)$. For $0 \le j \le k$ let

$$A_j = \prod_{i \le j} p_i, \qquad B_j = \prod_{i \le j} (p_i - 1), \qquad P_j = \frac{A_j}{B_j} = \prod_{i \le j} \frac{p_i}{p_i - 1},$$

with $A_0 = B_0 = P_0 = 1$ and $p_0 = 1$. We write $m = k - j$ for the number of primes still to be chosen at depth j, and $r_1 < r_2 < \cdots$ for the primes from 5 on.

2. Reductions

Lemma 2.1. *Let $n > 1$ satisfy $\varphi(n) \mid n + \varepsilon$ and put $M = (n+\varepsilon)/\varphi(n)$. Suppose n is composite if $\varepsilon = -1$, and $n > 3$ if $\varepsilon = +1$.*

(i) *n is odd, squarefree and composite, and $M \ge 2$.*
(ii) *For primes p, q dividing n, possibly equal, $p \nmid q - 1$ and $p \nmid M$.*
(iii) $\displaystyle\prod_{p \mid n} \frac{p}{p-1} = M - \frac{\varepsilon}{\varphi(n)}.$

Proof. If $n > 2$ is even then $\varphi(n)$ is even and $n + \varepsilon$ is odd. If $p^2 \mid n$ then p divides $\varphi(n)$, hence $n + \varepsilon$, which is impossible since $p \mid n$. For $\varepsilon = +1$ a prime solution satisfies $n - 1 \mid 2$, so $n \le 3$; thus n is composite in both cases. For odd squarefree composite n we have $\varphi(n) < n - 1$, so $M \ge 2$. If $p, q \mid n$ then $q - 1 \mid \varphi(n) \mid n + \varepsilon$ while $p \mid n$, so $p \nmid q - 1$; similarly $p \nmid M\varphi(n) = n + \varepsilon$. Finally $n/\varphi(n) = (n+\varepsilon)/\varphi(n) - \varepsilon/\varphi(n)$. □

The next lemma shows how the prime 3 enters, and that it does so differently for the two equations.

Lemma 2.2. *In the situation of Lemma 2.1, suppose $3 \mid n$. Then every other prime factor of n is congruent to 2 modulo 3, and*

$$M \equiv \begin{cases} 1 \pmod 3, & \varepsilon = -1, \\ 2 \pmod 3, & \varepsilon = +1. \end{cases}$$

Consequently $M \ge 4$ if $\varepsilon = -1$, and $M = 2$ or $M \ge 5$ if $\varepsilon = +1$. If $M \ge 4$ then $\omega(n) \ge 1540$.

Proof. For a prime $p \mid n$ with $p \ne 3$, Lemma 2.1(ii) gives $3 \nmid p - 1$, so $p \equiv 2$ and $p - 1 \equiv 1 \pmod 3$. Hence $\varphi(n) = 2\prod_{p \ne 3}(p-1) \equiv 2 \pmod 3$, while $n + \varepsilon \equiv \varepsilon \pmod 3$. The relation $n + \varepsilon = M\varphi(n)$ becomes $2M \equiv \varepsilon$, which is the stated congruence. Suppose $M \ge 4$. Then $n/\varphi(n) > 4$: for $\varepsilon = -1$ because $n/\varphi(n) = M + 1/\varphi(n)$ by Lemma 2.1(iii), and for $\varepsilon = +1$ because then $M \ge 5$ and $n/\varphi(n) = M - 1/\varphi(n) \ge 5 - \frac{1}{2}$. On the other hand, if $q_1 < q_2 < \cdots$ are the primes congruent to 2 modulo 3 from 5 on, then

$$\frac{n}{\varphi(n)} = \frac{3}{2} \prod_{p \mid n,\ p \ne 3} \frac{p}{p-1} \le \frac{3}{2} \prod_{i=1}^{k-1} \frac{q_i}{q_i - 1},$$

since $p/(p-1)$ decreases in p. An exact computation in rational arithmetic shows that the right-hand side is $3.99985\ldots$ for $k = 1539$ and first exceeds 4 at $k = 1540$, where the last prime in the product is $q_{1539} = 27947$. □

For $\varepsilon = -1$ this is a weak form of the bounds of Hagis and of Burcsi, Czirbusz and Farkas mentioned in the introduction, and Corollary 2.6 below goes much further, but the lemma needs no search and suffices wherever we use it. For $\varepsilon = +1$ it shows why 3 cannot be removed in the same way: $M = 2$ is compatible with $3 \mid n$, and each Fermat prime $5, 17, 257, 65537$ is congruent to 2 modulo 3, as are 353 and 929.

Lemma 2.3. *In the situation of Lemma 2.1, suppose* $3 \nmid n$ *and* $\omega(n) \le 32$. *Then* $M = 2$.

Proof. All prime factors are at least 5. By Lemma 2.1(iii) and monotonicity,

$$M = \frac{n}{\varphi(n)} + \frac{\varepsilon}{\varphi(n)} \le \prod_{i \le k} \frac{r_i}{r_i - 1} + \frac{1}{\prod_{i \le k}(r_i - 1)},$$

where $k = \omega(n)$, since $n/\varphi(n) = \prod_{p \mid n} p/(p-1)$ and $\varphi(n) = \prod_{p \mid n}(p-1)$, and the i-th smallest prime factor of n is at least r_i. An exact computation shows that the right-hand side is less than 3 for every $k \le 32$. (For $k = 32$ the product is $2.99315\ldots$ and the second term is below 10^{-53}; at $k = 33$ the product exceeds 3.) Since M is an integer and $M \ge 2$ by Lemma 2.1(i), $M = 2$. □

Corollary 2.4. *Let* $n > 3$ *satisfy* $\varphi(n) \mid n + 1$ *with* $\omega(n) \le 7$. *Then* $n + 1 = 2\varphi(n)$.

Proof. Let $k = \omega(n) \ge 2$. The prime factors of n, in increasing order, are at least $3, r_1, \ldots, r_{k-1}$, and Lemma 2.1(iii) gives, as in the proof of Lemma 2.3,

$$M \le \frac{3}{2} \prod_{i < k} \frac{r_i}{r_i - 1} + \frac{1}{2\prod_{i<k}(r_i - 1)}.$$

The maximum of the right-hand side over $2 \le k \le 7$ is $2.92357\ldots < 3$, attained at $k = 7$, and so $M = 2$. □

2.1. **Independent sets of primes.** Call a finite set of primes *independent* if none of its elements divides another minus one. Golomb [14] and Meijer [23] call an increasing sequence of odd primes with this property a G-sequence. By Lemma 2.1(ii) the prime factors of a solution form an independent set, whereas the proofs above used only that the i-th smallest of them is at least the i-th prime of a given list. Independence is a much stronger condition. Each prime q of the set excludes the primes congruent to 1 modulo q, which make up the proportion $1/(q-1)$ of all primes by the prime number theorem for arithmetic progressions, and so the product of $p/(p-1)$ over an independent set grows far more slowly with its size than the product over the first primes. Choosing at each step the least prime from 5 on that the primes already chosen allow gives the independent set

$$5,\ 7,\ 13,\ 17,\ 19,\ 23,\ 37,\ 59,\ 67,\ 73,\ 83,\ \ldots$$

(Figure 1). Its product passes 2 at the eleventh prime, where the first primes from 5 on need only seven, and it passes 3 only at the 100470-th prime, 5160959, where they need 33 (Figure 2).

Proposition 2.5. (i) *No independent set of at most* 18000 *primes* $p \ge 5$ *has* $\prod p/(p-1) \ge 2.999999$.

(ii) *No independent set of at most* 10^9 *primes* $p \equiv 2 \pmod 3$ *with* $p \ge 5$ *has* $\prod p/(p-1) \ge \frac{8}{3}(1 - 10^{-6})$.

(iii) *No independent set of at most* 10^9 *primes* $p \equiv 2 \pmod 3$ *with* $p \ge 5$ *has* $\prod p/(p-1) \ge \frac{10}{3}(1 - 10^{-6})$.

Proof. Let Q be the set of primes allowed, K the size and V the target. For an independent set $S \subseteq Q$ whose largest element is p (with $p = 3$ if S is empty), call the primes $q \in Q$ with $q > p$ and $q \not\equiv 1$ modulo every element of S the *candidates* for S. If $|S| < K$, let $B(S)$ be the product of $q/(q-1)$ over S and over the $K - |S|$ smallest candidates for S. An independent set T of




FIGURE 1. The primes from 5 to 89, with an arc from p to q whenever p divides $q-1$; an independent set contains no two primes joined by an arc. The set S of filled primes is chosen one prime at a time, each the least prime joined to none of those chosen before it. Below each prime of S is the product of $p/(p-1)$ over the primes of S up to it, which first exceeds 2 at 83, the eleventh.

at most K primes of Q whose $|S|$ smallest elements form S has its other elements among the candidates for S, and there are at most $K-|S|$ of them; as $q/(q-1)$ decreases in q, the product over T is at most $B(S)$.

The search starts from the empty set and treats each set S that it creates as follows. If the product over S is at least V, it stops and reports S. If $|S|=K$ or $B(S)<V$, it discards S. Otherwise it creates the sets $S\cup\{q\}$ for the candidates q with

$$\prod_{p\in S}\frac{p}{p-1}\cdot\frac{q}{q-1}\cdot\prod_{q'}\frac{q'}{q'-1}\ \geq\ V, \tag{2.1}$$

where q' runs over the $K-|S|-1$ candidates for S that follow q. The left-hand side of (2.1) is at least $B(S\cup\{q\})$, because every candidate for $S\cup\{q\}$ is a candidate for S, and it decreases as q increases, so the sets created from S come from its first few candidates. Let T be as above with product at least V. Every initial segment S of T has $B(S)\geq V$, and if $S\cup\{q\}$ is also an initial segment, then q satisfies (2.1); so the search creates the initial segments of T one after another until it stops.

In each of the three cases the search ends without reporting a set. It creates 502473 sets in case (i), the largest of them with 54 elements, and 19 sets in case (ii) and 2 in case (iii). Each search was carried out by two programs written independently. For case (i) they are a program in double precision, which also confirms every decision near the margin by exact rational arithmetic, and a program in the extended precision of the processor, with 64 binary digits, which passes the list of candidates of a set down to the sets created from it; for cases (ii) and (iii) the latter program, which runs through the primes with a segmented sieve, and a program in PARI/GP [31]. The primes up to $1.2\cdot10^{11}$ are needed in cases (ii) and (iii), and the programs run through them once for each set created. The products are handled through their logarithms, and every bound is a sum of a few terms $\log(q/(q-1))$ and a difference of two running sums of such terms. Every running sum here stays below 4; it has fewer than $2\cdot10^6$ terms in case (i), where the primes below $3\cdot10^7$ suffice, and fewer than $2.4\cdot10^9$ in cases (ii) and (iii). In double precision each term is computed with an absolute error below $3\cdot10^{-16}$ and each addition with one below 2^{-52}, so a bound carries an error below 10^{-8} in case (i); in extended precision the errors are below 2^{-62} per term and per addition, so a bound carries an error below

10^{-12} in case (i) and below 10^{-9} in cases (ii) and (iii). The PARI/GP program works with 38 significant digits, and its errors are below 10^{-25}. A set is discarded, or a candidate passed over, only when exact rational arithmetic confirms it or when the logarithm of its bound lies below $\log V$ by more than a margin that exceeds these errors: 10^{-7} in double precision, 10^{-6} in extended precision and 10^{-20} in PARI/GP. Rounding can therefore add sets to the search, but never remove one, and the two programs of a case may differ in the sets they create near the margin: in case (i) the second program creates 502696 sets, and in cases (ii) and (iii) both create the same sets. □

The search for (i) grows quickly with K: the first program creates 2567 sets for $K = 10^4$, 19711 for $K = 14000$ and 90320 for $K = 16000$; the second program also finds no set for $K = 20000$, creating 3131961 sets, but that run has not been repeated by the first. The least size of an independent set of primes from 5 on with product at least 3 therefore lies between 18001 and 100470, the length of the set chosen one prime at a time above. For the primes $p \equiv 2 \pmod 3$ the products grow much more slowly: the independent set chosen from them in the same way has 2307386 elements below $2 \cdot 10^8$, and its product is only $2.0703\ldots$, far below $\frac{8}{3}$.

Corollary 2.6. *Let n be as in Lemma 2.1, with $M = (n+\varepsilon)/\varphi(n)$.*

(i) *If $M \geq 3$ and $3 \nmid n$, then $\omega(n) \geq 18001$.*
(ii) *If $3 \mid n$ and $\varepsilon = -1$, then $\omega(n) \geq 10^9 + 2$.*
(iii) *If $3 \mid n$, $\varepsilon = +1$ and $M \geq 3$, then $\omega(n) \geq 10^9 + 2$.*

Proof. Let $k = \omega(n)$. By Lemma 2.1 the prime factors of n form an independent set, and $n/\varphi(n) = M - \varepsilon/\varphi(n)$.

(i) The prime factors are at least 5, and $n/\varphi(n) \geq 3 - 1/\varphi(n) > 2$. The product of $p/(p-1)$ over the six smallest primes from 5 on is $1.9490\ldots$, so $k \geq 7$ and $\varphi(n) \geq 4 \cdot 6 \cdot 10 \cdot 12 \cdot 16 \cdot 18 \cdot 22 > 10^6$. Hence $n/\varphi(n) > 2.999999$, and Proposition 2.5(i) gives $k > 18000$.

(ii) By Lemma 2.2, $M \geq 4$, and the $k-1$ prime factors other than 3 are congruent to 2 modulo 3. The product of $p/(p-1)$ over them is $\frac{2}{3}\, n/\varphi(n) = \frac{2}{3}(M + 1/\varphi(n)) > \frac{8}{3}$, so $k - 1 > 10^9$ by Proposition 2.5(ii).

(iii) By Lemma 2.2, $M \geq 5$, and the prime factors other than 3 are as in (ii). The product over them is $\frac{2}{3}(M - 1/\varphi(n)) \geq \frac{10}{3} - \frac{2}{3\varphi(n)} \geq 3$, while over the five smallest primes $p \equiv 2 \pmod 3$ from 5 on it is $1.5818\ldots$. So there are at least six of them, $\varphi(n) \geq 2 \cdot 4 \cdot 10 \cdot 16 \cdot 22 \cdot 28 \cdot 40 > 10^6$, and the product exceeds $\frac{10}{3}(1 - 10^{-6})$. Proposition 2.5(iii) gives $k - 1 > 10^9$. □

For $\varepsilon = -1$, part (ii) improves the bound $\omega(n) \geq 40000000$ of Burcsi, Czirbusz and Farkas [6], who completed sequences of primes $p \equiv 2 \pmod 3$ in a similar way by the least primes allowed. Part (i) replaces the bound 33 that the products over the first primes give, and for $M = 3$ the bound 1991 of Hagis [17]; part (iii) replaces the bound 1540 that the products give.

Proof of Theorem 1.6 and of the last two assertions of Theorem 1.1. Let n satisfy (1.1) or (1.2), with n composite in the first case and $n > 3$ in the second. If $3 \nmid n$ and $M \geq 3$, then $\omega(n) \geq 18001$ by Corollary 2.6(i). If $3 \mid n$, then $\omega(n) > 10^9$ by Corollary 2.6(ii) for (1.1), where $M \geq 4$ holds automatically, and by Corollary 2.6(iii) for (1.2) when $M \geq 3$. □

By Corollary 2.6, a solution with at most 18000 prime factors has $M = 2$, and if it solves (1.1) it is prime to 3. The searches below concern the case $n + \varepsilon = 2\varphi(n)$.

2.2. **The growth of independent sets.** Proposition 2.5 needs a search because the product of $p/(p-1)$ over an independent set grows very slowly with the size of the set. Over the first K primes the product is asymptotic to $e^\gamma \log K$ by Mertens' theorem, so that the products alone give only $\omega(n) \geq \exp(c'M)$ for some $c' > 0$. Over an independent set of K primes the product is




FIGURE 2. The product of $p/(p-1)$ over the first k primes of a list (solid) and over the independent set chosen from the same list one prime at a time (dashed), for the primes below $2 \cdot 10^8$, with k on a logarithmic scale. Where a list has both curves, the largest product over an independent set of k of its primes lies between them. (a) For the primes from 5 on the product passes 2 at $k = 7$, so a solution of (3.1) prime to 3 has at least seven prime factors (Section 5), and 3 at $k = 33$ (Lemma 2.3); for the odd primes it passes 3 at $k = 8$, where the argument of Corollary 2.4 stops. The independent set passes 3 only at $k = 100470$, and no independent set of primes from 5 on reaches 3 before $k = 18001$ (open circle). (b) For 3 and the primes $p \equiv 2 \pmod 3$ the product passes 4 at $k = 1540$ (Lemma 2.2), and no independent set formed by 3 and such primes reaches 4 before $k = 10^9 + 2$ (open circle).

at most a constant multiple of $\log\log K$, and this forces $\omega(n)$ to be at least doubly exponential in M (Theorem 1.8).

For a set S of primes and $x \geq 2$ put

$$P_S(x) = \prod_{p \in S,\ p \leq x} \frac{p}{p-1}, \qquad \delta_S(x) = \prod_{p \in S,\ p \leq x} \Big(1 - \frac{1}{p-1}\Big).$$

An independent set with at least two elements does not contain 2, since 2 divides $q-1$ for every odd prime q. For a set S of odd primes every factor of $\delta_S(x)$ is at least $\frac{1}{2}$, and since $\frac{p}{p-1}\big(1 - \frac{1}{p-1}\big) = 1 - \frac{1}{(p-1)^2}$,

$$\delta_S(x)\, P_S(x) \leq 1. \tag{2.2}$$

Theorem 2.7 (Meijer [23])**.** *There is an absolute constant c_0 such that every independent set S of $K \geq 3$ primes satisfies*

$$\prod_{p \in S} \frac{p}{p-1} \leq c_0 \log\log K.$$

Meijer proved the bound $P_S(x) \leq c \log\log x$ with the Brun–Selberg sieve, and Yamada [39] gave explicit constants with the large sieve. We include a short proof by the large sieve, with constants left unspecified. Each element p of S excludes from S the primes congruent to 1 modulo p, so that a prime above x can belong to S only if it avoids one residue class modulo every element of S up to x; the large sieve bounds the number of such primes by $\delta_S(x)$ times the number of all primes in the range.

Lemma 2.8. *Let S be an independent set of odd primes and $N \geq 2^{88}$. Then*

$$\#\{q \in S : \ N^{1/44} < q \leq N\} \leq \frac{132\, N\, \delta_S(N^{1/44})}{\log N}.$$

Proof. Put $Q = N^{1/2}$ and $z = N^{1/44} = Q^{1/22}$, so that $z \geq 4$. For a prime $p \leq z$ let Ω_p consist of the residue class 0 modulo p and, if $p \in S$, also of the class 1; for $z < p \leq Q$ let Ω_p be empty. Let $\omega(p) < p$ be the number of classes in Ω_p. A prime $q \in S$ with $z < q \leq N$ lies in none of these classes: it is not divisible by a prime $p \leq z < q$, and for $p \in S$ it is not congruent to 1 modulo p, because S is independent. The large sieve [25, 26] states that a set of integers in $[1, N]$ that avoids the classes of Ω_p for every prime $p \leq Q$ has at most $(N + Q^2)/L$ elements, where

$$L = \sum_{q \leq Q} \mu^2(q)\, G(q), \qquad G(q) = \prod_{p \mid q} g(p), \qquad g(p) = \frac{\omega(p)}{p - \omega(p)}.$$

Here $g(p) = 1/(p-1)$ for $p \leq z$ with $p \notin S$, $g(p) = 2/(p-2)$ for $p \leq z$ with $p \in S$, and $g(p) = 0$ for $p > z$; in every case $g(p) \leq 6/p$.

Let Π be the product of the primes up to z and put $\sigma = 1/\log z$, so that $Q^\sigma = e^{22}$. Then L is the sum of $G(q)$ over the divisors $q \leq Q$ of Π. A divisor $q > Q$ has $(q/Q)^\sigma > 1$, and so

$$\sum_{q \mid \Pi,\ q > Q} G(q) \leq Q^{-\sigma} \sum_{q \mid \Pi} G(q)\, q^\sigma = e^{-22} \prod_{p \leq z} (1 + g(p) p^\sigma).$$

For $p \leq z$ we have $0 < \sigma \log p \leq 1$, hence $p^\sigma - 1 \leq (e-1)\sigma \log p$ by convexity, and $1 + g(p)p^\sigma \leq (1 + g(p)) \exp(g(p)(p^\sigma - 1))$. With $g(p) \leq 6/p$ and the elementary bound[6] $\sum_{p \leq z} (\log p)/p < \log z + \log 4$ this gives

$$\prod_{p \leq z} (1 + g(p) p^\sigma) \leq \exp\Big(6(e-1)\Big(1 + \frac{\log 4}{\log z}\Big)\Big) \prod_{p \leq z} (1 + g(p)) < e^{20.7} \prod_{p \leq z} (1 + g(p)),$$

because $z \geq 4$ and $12(e-1) < 20.7$. Since $\prod_{p \leq z}(1 + g(p)) = \sum_{q \mid \Pi} G(q)$ and $e^{20.7 - 22} < \frac{1}{3}$, the divisors $q > Q$ contribute less than a third of this sum, and $L > \frac{2}{3} \prod_{p \leq z}(1 + g(p))$. Finally $1 + g(p) = p/(p-1)$ for $p \notin S$ and $1 + g(p) = \frac{p}{p-1}\big(1 - \frac{1}{p-1}\big)^{-1}$ for $p \in S$, so that

$$\prod_{p \leq z} (1 + g(p)) = \frac{1}{\delta_S(z)} \prod_{p \leq z} \Big(1 - \frac{1}{p}\Big)^{-1} > \frac{\log z}{\delta_S(z)},$$

since the product $\prod_{p \leq z}(1 - 1/p)^{-1}$ is at least $\sum_{n \leq z} 1/n > \log z$. With $\log z = (\log N)/44$ we obtain $L > \log N / (66\, \delta_S(z))$, and the large sieve, with $N + Q^2 = 2N$, gives the lemma. □

[6] Let $m = \lfloor z \rfloor$. Legendre's formula gives $\log m! \geq \sum_{p \leq m} \lfloor m/p \rfloor \log p > m \sum_{p \leq m} (\log p)/p - \sum_{p \leq m} \log p$, while $\log m! \leq m \log m$ and $\sum_{p \leq m} \log p < m \log 4$, because the product of the primes up to m is less than 4^m. Dividing by m gives the bound.

Lemma 2.9. *Let S be an independent set of odd primes and $y \ge 2^{88}$. Then*

$$\sum_{q\in S,\ y<q\le y^2} \frac{1}{q} \le 269\,\delta_S(y^{1/44}).$$

Proof. Let $a = \lfloor \log_2 y \rfloor \ge 88$ and $b = \lceil 2\log_2 y \rceil$, so that $b \le 2a+2$. The interval $(y, y^2]$ is covered by the intervals $(2^{i-1}, 2^i]$ with $a < i \le b$. For each such i, Lemma 2.8 with $N = 2^i$ applies to the elements of S in $(2^{i-1}, 2^i]$, because $2^{i/44} \le 2^{i-1}$, and $\delta_S(2^{i/44}) \le \delta_S(y^{1/44})$, because $2^i > y$ and δ_S is decreasing. Each of these elements exceeds 2^{i-1}, so their reciprocals add up to at most

$$\frac{132\cdot 2^i\,\delta_S(y^{1/44})}{2^{i-1}\,i\log 2} = \frac{264\,\delta_S(y^{1/44})}{i\log 2}.$$

Since $\sum_{a<i\le b} 1/i \le \log(b/a) \le \log(2+2/88) < 0.705$ and $264\cdot 0.705/\log 2 < 269$, the lemma follows. □

Proof of Theorem 2.7. By Mertens' theorem $\prod_{p\le x}(1-1/p)^{-1} = e^\gamma \log x + O(1)$, so there is an absolute constant c_1 with $\prod_{u<p\le u^{44}} p/(p-1) \le c_1$ for all $u \ge 2$. Let S be an independent set of odd primes. Then $P_S(y) \le c_1 P_S(y^{1/44})$ for $y \ge 2^{44}$, and Lemma 2.9 and (2.2) give, for $y \ge 2^{88}$,

$$\sum_{q\in S,\ y<q\le y^2} \frac{1}{q} \le \frac{269}{P_S(y^{1/44})} \le \frac{269\,c_1}{P_S(y)}. \tag{2.3}$$

Put $y_0 = 2^{88}$, $y_{j+1} = y_j^2$ and $a_j = P_S(y_j)$. Since $-\log(1-1/q) \le 2/q$, (2.3) gives $a_{j+1} \le a_j e^{c_2/a_j}$ with $c_2 = 538\,c_1$, and also $a_{j+1} \le c_1 a_j$. If $a_j \ge c_2$, then $a_{j+1} \le a_j(1+2c_2/a_j) = a_j + 2c_2$, because $e^t \le 1+2t$ for $0 \le t \le 1$; if $a_j < c_2$, then $a_{j+1} < c_1 c_2$. By induction $a_j \le c_3 + 2c_2 j$ for all $j \ge 0$, where $c_3 = \max\{c_1c_2, \prod_{p\le 2^{88}} p/(p-1)\} \ge a_0$. For $x > y_0$ let j be the least index with $y_j \ge x$. Then $2^{j-1}\log y_0 < \log x$, and $P_S(x) \le a_j \le c_3 + 2c_2(1+\log_2(\log x/\log y_0))$. Together with $P_S(x) \le a_0 \le c_3$ for $x \le y_0$, this gives $P_S(x) \le c_4 \log\log x$ for all $x \ge 3$, with an absolute constant c_4.

Now let S have $K \ge 3$ elements, so that its elements are odd. At most K of them exceed K^2, and each of those contributes a factor at most $1 + 1/K^2$, so that together they contribute less than $e^{1/K} < 2$. Hence

$$\prod_{p\in S} \frac{p}{p-1} < 2P_S(K^2) \le 2c_4 \log\log K^2 = 2c_4(\log 2 + \log\log K) \le 20\,c_4 \log\log K,$$

since $\log 2 < 9\log\log 3$. □

Proof of Theorem 1.8. Let n be as in Lemma 2.1 and $k = \omega(n) \ge 2$. The prime factors of n form an independent set, and $n/\varphi(n)$ is the product of $p/(p-1)$ over them. If $k \ge 3$, Theorem 2.7 and Lemma 2.1(iii) give $M \le c_0 \log\log k + 1/\varphi(n) < (c_0+3)\log\log(k+2)$, since $\log\log 5 > \frac{1}{3}$. If $k = 2$, then $n/\varphi(n) \le \frac{3}{2}\cdot\frac{5}{4}$ and $\varphi(n) \ge 8$, so that $M \le 2 < 7\log\log 4$. In both cases $\log\log(k+2) \ge cM$ with $c = 1/\max\{c_0+3, 7\}$, which is the assertion. □

The constants in Theorems 2.7 and 1.8 are effective but large, and the numerical bounds of Corollary 2.6 come from Proposition 2.5, not from the sieve. What the sieve explains is the shape of those bounds. Heuristically, the elements of S up to x leave a proportion $\delta_S(x)$ of the primes near x available, and a set that takes every available prime gains $\log P_S$ at the rate $\delta_S(x)\,d\log\log x$, since $\sum_{q\le x} 1/q$ grows like $\log\log x$. The derivative of P_S with respect to $\log\log x$ is then $\delta_S(x)P_S(x)$, which is at most 1 by (2.2), so that $P_S(x)$ grows at most like $\log\log x$, the rate of Theorem 2.7. The independent sets chosen one prime at a time in Section 2.1 grow at this rate (Table 1). Between $x = 10^3$ and $x = 10^8$ the product of $p/(p-1)$ over the elements up to x differs from a linear function of $\log\log x$ by less than 0.003, with slope 0.82

for the primes from 5 on and 0.41 for the primes $p \equiv 2 \pmod 3$, which are half of all primes. The first set needs 100470 primes to reach the product 3, where the first primes from 5 on need 33, and the second has not reached $\frac{8}{3}$ below $2 \cdot 10^8$ (Section 2.1). This is why the bounds of Corollary 2.6 exceed 33 and 1540 by so much.

TABLE 1. The independent sets chosen one prime at a time from the primes from 5 on and from the primes $p \equiv 2 \pmod 3$ from 5 on: the number of elements up to x and the product of $p/(p-1)$ over them.

		primes from 5 on		$p \equiv 2 \pmod 3$	
x	$\log\log x$	elements	product	elements	product
10^3	1.933	62	2.3430	45	1.6489
10^4	2.220	407	2.5792	293	1.7626
10^5	2.443	2 908	2.7593	2 185	1.8564
10^6	2.626	22 602	2.9078	17 396	1.9328
10^7	2.780	183 730	3.0344	142 723	1.9979
10^8	2.913	1 538 281	3.1447	1 207 989	2.0546

3. Intervals for the prime factors

Fix $k \geq 2$, a sign ε, and a squarefree odd $n = p_1 \cdots p_k$ with

$$n + \varepsilon = 2\varphi(n). \tag{3.1}$$

By Lemma 2.1(iii),

$$\prod_{i=1}^{k} \frac{p_i}{p_i - 1} = 2 - \frac{\varepsilon}{\varphi(n)}. \tag{3.2}$$

For $\varepsilon = -1$ the product lies just above 2, and for $\varepsilon = +1$ just below it. Everything in this section is a consequence of (3.2), and the two signs are treated together.

Proposition 3.1. *$P_j < 2$ for every $j < k$.*

Proof. For $\varepsilon = +1$ this is clear, since $P_j \leq P_k = 2 - 1/\varphi(n)$. For $\varepsilon = -1$, suppose $P_j \geq 2$ with $j < k$. Since $p_{j+1} - 1 < n - 1$ we have $p_{j+1}/(p_{j+1}-1) > 1 + 1/(n-1)$, and the remaining factors exceed 1, so $P_k > 2(1 + 1/(n-1)) = 2 + 1/\varphi(n)$ by (3.1), contradicting (3.2). □

At depth j put $T = 2/P_j$, which exceeds 1 by the proposition, and

$$C_j = 2B_j - A_j, \qquad T - 1 = \frac{C_j}{A_j}.$$

Thus C_j is a positive integer, and C_j/A_j measures how far the primes chosen so far fall short of the target 2.

Proposition 3.2. *Let $0 \leq j \leq k-2$, $m = k - j$, $W = (p_j + 1)^m$ and $x = p_{j+1}/(p_{j+1} - 1)$. Then*

$$\begin{aligned} x^m > T \quad &\text{and} \quad x < T + \frac{1}{A_j W} \quad \text{if } \varepsilon = -1, \\ x^m \geq T - \frac{1}{A_j W} \quad &\text{and} \quad x < T \quad \text{if } \varepsilon = +1. \end{aligned} \tag{3.3}$$

In particular p_{j+1} lies in an interval $[L_j, U_j]$ whose endpoints are determined by $p_1, \ldots, p_j$, k and ε.

Proof. Let $\rho = \prod_{i>j} p_i/(p_i-1)$. By (3.2),

$$\rho = \frac{2-\varepsilon/\varphi(n)}{P_j} = T - \frac{\varepsilon}{P_j\varphi(n)}.$$

There are $m \geq 2$ factors in ρ; each exceeds 1 and none exceeds x, because $p/(p-1)$ decreases in p. Hence $x < \rho \leq x^m$. The remaining primes are odd and larger than p_j (larger than 2 when $j=0$), so $p_i - 1 \geq p_j + 1$ for $i > j$, and $\varphi(n) = B_j \prod_{i>j}(p_i-1) \geq B_j W$. Therefore

$$0 < \frac{1}{P_j\varphi(n)} \leq \frac{1}{P_j B_j W} = \frac{1}{A_j W},$$

and the four inequalities follow from $x < \rho \leq x^m$. Finally, for $\varepsilon = +1$ the quantity $T - 1/(A_jW)$ exceeds 1, since $T - 1 = C_j/A_j \geq 1/A_j$. □

Apart from the correction terms, (3.3) says $x < T \leq x^m$, that is $1/(T-1) < p_{j+1} - 1 \leq 1/(T^{1/m}-1)$. Since $T^{1/m}-1$ is close to $(T-1)/m$, the interval has length about $(m-1)/(T-1) = (m-1)A_j/C_j$ (Figure 3). The upper end obeys a bound that makes the search finite.

Corollary 3.3. *Let $0 \leq j \leq k-2$, let $p_1 < \cdots < p_j$ be odd primes with $P_j < 2$, and let $p_{j+1} > p_j$ be a prime that satisfies the inequality for x^m in (3.3). Then $p_{j+1} < 1 + 3(k-j)B_j/C_j < 4kA_j$. If $n = p_1 \cdots p_k$ satisfies (3.1), then moreover $p_k \leq 2B_{k-1} + 1$, and*

$$A_j < (4k)^{2^j} \quad (0 \leq j \leq k), \qquad n < (4k)^{2^k}.$$

Proof. Let $m = k - j$ and $x = p_{j+1}/(p_{j+1}-1)$, so that $p_{j+1} - 1 = 1/(x-1)$. By (3.3), $x^m \geq T'$, where $T' = T$ if $\varepsilon = -1$ and $T' = T - 1/(A_jW)$ if $\varepsilon = +1$. As $T - 1 = C_j/A_j$ with $C_j \geq 1$, and $W \geq 4$, in both cases $T' - 1 \geq (C_j - \frac{1}{4})/A_j \geq 3C_j/(4A_j) > 0$ and $T' \leq T$. For $t > 1$ we have $t^{1/m} - 1 = (t-1)/\sum_{i<m} t^{i/m} \geq (t-1)/(mt)$, and so

$$p_{j+1} - 1 = \frac{1}{x-1} \leq \frac{1}{T'^{1/m}-1} \leq \frac{mT'}{T'-1} \leq \frac{2mB_j}{A_j} \cdot \frac{4A_j}{3C_j} < \frac{3mB_j}{C_j}.$$

Since $C_j \geq 1$ and $B_j \leq A_j - 1$ for $j \geq 1$, while $A_0 = B_0 = C_0 = 1$, this gives $p_{j+1} < 4kA_j$. For a solution, the equation $A_{k-1}p_k + \varepsilon = 2B_{k-1}(p_k - 1)$ gives $p_k C_{k-1} = 2B_{k-1} + \varepsilon$, and $C_{k-1} \geq 1$ by Proposition 3.1, so that $p_k \leq 2B_{k-1} + 1 < 4kA_{k-1}$. By Propositions 3.1 and 3.2 the first part applies to p_{j+1} for every $j \leq k-2$. Hence $A_{j+1} < 4kA_j^2$ for $0 \leq j < k$, and induction gives $A_j \leq (4k)^{2^j-1}$. □

Every node of the search below satisfies the hypotheses of Corollary 3.3, so the tree has finitely many nodes and the search terminates. The bound $n < (4k)^{2^k}$ has the shape of the bound $n < k^{2^k}$ of Pomerance [33]; Theorem 1.2 replaces it by $n < 2^{2^{k-7}}$.

The inequalities (3.3) are tested in integers: for instance $x^m > T$ reads $p^m A_j > 2B_j(p-1)^m$. A floating-point estimate is used only to find a starting point, which is then corrected by exact comparisons.

Lemma 3.4 (congruence prune)**.** *No prime factor of n divides $p-1$ for another prime factor p of n.*

This is Lemma 2.1(ii). A candidate p for p_{j+1} is discarded if $p \mid p_i - 1$ or $p_i \mid p - 1$ for some $i \leq j$. The primes of the interval $[L_j, U_j]$ that are not discarded are called *admissible*.

The last two primes are determined together by a divisibility condition.

Proposition 3.5. *Let $A = A_{k-2}$, $B = B_{k-2}$, $C = 2B - A$, $p = p_{k-1}$ and $q = p_k$. Then*

$$t := Cp - 2B > 0, \qquad t \mid Ap + \varepsilon, \qquad q = 1 + \frac{Ap+\varepsilon}{t}, \tag{3.4}$$




FIGURE 3. The interval of Proposition 3.2 for $T = 1.04$ and $m = 3$, with the correction terms $1/(A_jW)$ left out. The candidate p must satisfy $x < T$, because the other $m-1$ factors of ρ exceed 1, and $x^m \ge T$, because none of them exceeds x. Here the two conditions give $26 < p < 77$.

and equivalently

$$(Cp - 2B)(Cq - 2B) = 2AB + \varepsilon C. \tag{3.5}$$

Proof. Equation (3.1) reads $Apq + \varepsilon = 2B(p-1)(q-1)$. Collecting the terms in q gives $q\,(Cp - 2B) = 2B(p-1) + \varepsilon$. The right-hand side is positive, so $t > 0$, and $q - 1 = (2Bp - 2B + \varepsilon - Cp + 2B)/t = (Ap + \varepsilon)/t$. Multiplying $Cpq - 2B(p+q) + 2B - \varepsilon = 0$ by C and completing the product gives $(Cp - 2B)(Cq - 2B) = 4B^2 - 2BC + \varepsilon C = 2AB + \varepsilon C$. □

Once $p_1, \dots, p_{k-2}$ are fixed, the prime p_{k-1} runs over the interval of Proposition 3.2 with $m = 2$, and for each candidate p the last prime is determined by (3.4). Alternatively, by (3.5), t is a divisor of the integer

$$D = 2AB + \varepsilon C$$

lying in a known residue class modulo C, so the candidates can be read off from the divisors of D when D can be factored. The first route costs time proportional to the length of the interval, the second the cost of factoring D. Figure 4 summarises the procedure.

The search is a depth-first traversal of the tree whose nodes at depth $j \le k-2$ are the increasing sequences $(p_1, \dots, p_j)$ of primes with $p_1 \ge p_{\min}$, where each p_{i+1} lies in the interval determined by $(p_1, \dots, p_i)$ and passes the congruence prune, and where $P_i < 2$ for all $i \le j$. At depth $k-2$ each candidate p_{k-1} is tested with (3.4): one checks that $t > 0$, that $t \mid Ap + \varepsilon$, that $q = 1 + (Ap + \varepsilon)/t$ is a prime larger than p, that p and q pass the congruence prune, and finally that (3.1) holds exactly.

Theorem 3.6 (completeness)**.** *If $n = p_1 \cdots p_k$ with primes $p_{\min} \le p_1 < \cdots < p_k$ satisfies (3.1), then the traversal visits $(p_1, \dots, p_{k-2})$, examines the candidate p_{k-1}, and records n.*

Proof. By induction on j the node $(p_1, \dots, p_j)$ is visited for every $j \le k-2$: the prime p_{j+1} lies in the interval of Proposition 3.2, it passes the prune by Lemma 3.4, and $P_{j+1} < 2$ by Proposition 3.1. At depth $k-2$ the candidate $p = p_{k-1}$ satisfies (3.4) with $q = p_k$. A primality




FIGURE 4. The search for $n = p_1 \cdots p_k$ with $n + \varepsilon = 2\varphi(n)$. A node at depth j is an increasing sequence of primes $p_1 < \cdots < p_j$. Its children are the primes in the interval $[L_j, U_j]$ of Proposition 3.2 that pass the congruence prune; dashed branches are cut. At depth $k - 2$ the pair (p_{k-1}, p_k) is a lattice point on the hyperbola (3.5), found by running p over $[L_{k-2}, U_{k-2}]$ or by factoring $2AB + \varepsilon C$.

test never declares a prime to be composite, so q is accepted, and the final check of (3.1) succeeds. □

The search is rigorous for two reasons. First, all arithmetic is exact: A_j, B_j, C_j, the interval endpoints and the quantities in (3.4) are integers or ratios of integers compared by cross multiplication. Second, the probabilistic primality tests used by the programs[7] can err only by declaring a composite number prime, never by declaring a prime composite. Such an error could add a node or accept a composite q, but it can never remove an actual solution. A search that finds nothing is therefore a proof that there is nothing, and the completeness of the enumeration route depends on no primality proof. The exact check of (3.1) would not catch such an error, since it uses $\prod(p_i - 1)$ in place of $\varphi(n)$. This matters only for solutions that are reported, and every solution reported in this paper, including those used in the checks of Section 8, has all its prime factors below 10^{14}, where the tests are proofs. The divisor route depends on a complete factorisation of D, which is certified whenever $D < \psi_{13}$, since then every prime factor of D lies below that bound.

4. The last three primes

For $k \le 14$ the search of Section 3 is cheap because the interval for p_{k-1} is never longer than about $5 \cdot 10^7$. For $k = 15$ this is no longer true. After the prime p_{13} has been chosen the interval for p_{14} can be longer than 10^{14}, and there are tens of millions of choices of p_{13}. Neither route of Section 3 copes with this: running p_{14} over its interval is far too slow, and factoring tens of millions of integers of about fifty digits is a computation of a different order. In this section we describe two further ways of finding the divisors of $2AB + \varepsilon C$ that can occur in Proposition 3.5. Both use the residue class in which these divisors lie: the first lists lattice points near a hyperbola, and the second scans the possible values of the sum of two complementary divisors.

[7]Below the number $\psi_{13} = 331704406467988738596 1981$ the programs use a strong probable-prime test to the thirteen bases $2, 3, \ldots, 41$, and above it a Baillie–PSW test. Sorenson and Webster [36] showed that every composite number below ψ_{13} fails the first test for at least one of these bases, so there it is a proof. The Baillie–PSW test [2, 34] combines a strong probable-prime test to base 2 with a strong Lucas test; no composite number is known to pass it.




FIGURE 5. The surface $\frac{s}{s-1} \cdot \frac{p}{p-1} \cdot \frac{q}{q-1} = T$ with $s < p < q$, for $T = 1.1$ on logarithmic axes. Each slice with fixed s, three of which are marked, falls steeply from large values of q to the edge $p = q$. The slices shrink as s grows, and the surface ends at the point $s = p = q$ where $(s/(s-1))^3 = T$.

4.1. **Reduction to divisors in a residue class.** Fix a node $(p_1, \ldots, p_{k-3})$ at depth $k-3$ and a candidate $s = p_{k-2}$ in its interval that passes the congruence prune. Put

$$A' = A_{k-3}\, s, \qquad B' = B_{k-3}\,(s-1), \qquad c = 2B' - A', \qquad N = 2A'B' + \varepsilon c.$$

If $c \le 0$ then $P_{k-2} = A'/B' \ge 2$, and by Proposition 3.1 no solution has $p_1, \ldots, p_{k-3}, s$ as its first $k-2$ primes; such s are discarded. So let $c \ge 1$. By Proposition 3.5, applied with $A = A'$ and $B = B'$, the remaining primes $p < q$ of a solution satisfy $(cp - 2B')(cq - 2B') = N$ with both factors positive. Let r be the least positive residue of $-2B'$ modulo c.

By (3.2) the last three primes satisfy

$$\frac{s}{s-1} \cdot \frac{p}{p-1} \cdot \frac{q}{q-1} = T - \frac{\varepsilon}{P_{k-3}\,\varphi(n)}, \qquad T = \frac{2}{P_{k-3}},$$

so the point (s, p, q) lies next to the surface on which this product equals T. Fixing s cuts the surface in a curve, and the pair (p, q) lies exactly on the hyperbola $(cp - 2B')(cq - 2B') = N$ beside that curve (Figure 5).

Proposition 4.1. *The integer $t = cp - 2B'$ is a divisor of N with*

$$t \equiv r \pmod{c}, \qquad 1 \le t \le \sqrt{N}.$$

Moreover $\gcd(r, c) = 1$ *and* $N \equiv r^2 \pmod{c}$. *Conversely, every divisor t of N with these properties gives integers $p = (t + 2B')/c$ and $q = (N/t + 2B')/c$ satisfying $A'pq + \varepsilon = 2B'(p-1)(q-1)$.*

Proof. The congruence holds because $t \equiv -2B' \pmod{c}$, and $t \ge 1$ because both factors are positive, and $t \le \sqrt{N}$ because $p < q$ makes t the smaller of the two factors. Since $c = 2B' - A'$ we

have $\gcd(2B', c) = \gcd(2B', A')$, which is 1 because A' is odd and, by the congruence prune, no prime factor of A' divides B'; hence $\gcd(r, c) = 1$. Further $N \equiv 2A'B' \equiv (2B')^2 \equiv r^2 \pmod{c}$, using $A' \equiv 2B'$. Conversely, let t be such a divisor. Then $N/t \equiv r^2 r^{-1} = r \pmod{c}$, so both t and N/t are congruent to $-2B'$ and p and q are integers. Expanding $(cp - 2B')(cq - 2B') = N$ and using $4B'^2 - 2A'B' = 2B'c$ gives $c(cpq - 2B'(p + q) + 2B' - \varepsilon) = 0$, which, as $c \geq 1$ and $c = 2B' - A'$, is the equation $A'pq + \varepsilon = 2B'(p - 1)(q - 1)$. □

So the two-prime problem at depth $k - 2$ becomes the following question: given N, a modulus c and a class r prime to c, list the divisors of N up to $\sqrt{N}$ that are congruent to r modulo c. Lenstra [20] proved that there are at most 11 such divisors when $c > N^{1/3}$,[8] and gave a polynomial-time algorithm to find them; Coppersmith, Howgrave-Graham and Nagaraj [11] extended the algorithmic side down to $c > N^{1/4+\delta}$. We use a more elementary method suited to our range, and in Section 4.4 a still simpler one based on the fact that t and N/t lie in the same class.

4.2. **Lattice points near a hyperbola.** Let r' be the residue of Nr^{-1} modulo c, so that $r' = r$ in the situation of Proposition 4.1, and put

$$w = \frac{N - rr'}{c}, \qquad \alpha \equiv wr^{-1}, \qquad \beta \equiv r'r^{-1} \pmod{c}.$$

A divisor $t \equiv r$ is written $t = r + cu$ with $u \geq 0$, and then $N/t = r' + cv$ for an integer v. Expanding $(r + cu)(r' + cv) = N$ gives

$$cuv + r'u + rv = w, \tag{4.1}$$

so the pairs (u, v) are the lattice points on a hyperbola. Reducing (4.1) modulo c shows that each of them lies in the coset

$$v \equiv \alpha - \beta u \pmod{c}. \tag{4.2}$$

The coset has one point in every c consecutive values of v above a given u, so by itself it excludes little; combined with the shape of the hyperbola it leaves only a few points to test.

Lemma 4.2. *Let $0 \leq u_0 \leq u_1$ and put*

$$v^+ = \left\lfloor \frac{\lfloor N/(r + cu_0) \rfloor - r'}{c} \right\rfloor, \qquad v^- = \left\lceil \frac{\lceil N/(r + cu_1) \rceil - r'}{c} \right\rceil.$$

If $t = r + cu$ divides N and $u_0 \leq u \leq u_1$, then $N/t = r' + cv$ with $v^- \leq v \leq v^+$ and v satisfying (4.2). Consequently, with $H = v^+ - v^- + 1$, the integer u lies in the set

$$\{\, u \in [u_0, u_1] : \ (\alpha - \beta u - v^-) \bmod c \ < H \,\}, \tag{4.3}$$

and there is no such t at all if $H \leq 0$.

Proof. Since $t \equiv r \pmod{c}$ and $\gcd(r, c) = 1$, we have $N/t \equiv Nr^{-1} \equiv r' \pmod{c}$, so $v = (N/t - r')/c$ is an integer. The cofactor N/t is an integer between $N/(r + cu_1)$ and $N/(r + cu_0)$, hence between the ceiling of the first and the floor of the second, and this gives $v^- \leq v \leq v^+$; in particular $H \geq 1$. The congruence (4.2) was shown above. Put $x = (\alpha - \beta u - v^-) \bmod c$, so that $0 \leq x < c$ and $v - v^- \equiv x \pmod{c}$. As $v - v^- \geq 0$, it is at least the least non-negative member x of its class, and so $x \leq v - v^- \leq H - 1$. □

[8] In the form used here: if $0 \leq r < c < N$, $\gcd(r, c) = 1$ and $c > N^{1/3}$, then at most 11 positive divisors of N are congruent to r modulo c. The bound counts all divisors in the class, not only those up to $\sqrt{N}$.

If $H \geq c$ the set (4.3) is the whole of $[u_0, u_1]$, and every u in the box is tested. If $H < c$ it consists of the u for which an arithmetic progression modulo c falls into a window of length H, and its elements can be listed in increasing order without looking at the others, at a cost of $O(\log c)$ arithmetic operations each, by the following elementary recursion.

Lemma 4.3. *Let $0 \leq a < g$ and $0 < L \leq R < g$. The least $y \geq 0$ with $L \leq ay \bmod g \leq R$ is found as follows. If $a = 0$ there is no such y. If $a \geq 1$ and $a\lceil L/a \rceil \leq R$, it is $\lceil L/a \rceil$. Otherwise it is $\lceil (L + gz)/a \rceil$, where z is the least integer ≥ 0 with*

$$(-R) \bmod a \;\leq\; (g \bmod a)\, z \bmod a \;\leq\; (-L) \bmod a,$$

and if no such z exists there is no y. The last problem is of the same kind, with (g, a, L, R) replaced by $(a,\ g \bmod a,\ (-R) \bmod a,\ (-L) \bmod a)$.

Proof. If $a = 0$ then $ay \bmod g = 0 < L$ for every y. In the second case $ay \leq R < g$ for $y = \lceil L/a \rceil$, so $ay \bmod g = ay \in [L, R]$, while $0 \leq ay < L$ for smaller $y \geq 0$. Otherwise no multiple of a lies in $[L, R]$. Since any a consecutive integers contain a multiple of a, this gives $R - L \leq a - 2$; moreover $a \nmid R$, and no multiple of a lies in $[-R, -L]$, so the residues $(-R) \bmod a, \ldots, (-L) \bmod a$ of $-R, \ldots, -L$ increase without wrapping around and $0 < (-R) \bmod a \leq (-L) \bmod a < a$. A solution y satisfies $ay = gz + x$ with $z = \lfloor ay/g \rfloor$ and $x = ay \bmod g \in [L, R]$, and $z \geq 1$ because $z = 0$ would make $x = ay$ a multiple of a in $[L, R]$. Conversely, for a given $z \geq 0$ a solution with this z exists exactly when $[L + gz, R + gz]$ contains a multiple of a; that multiple is unique, because $R - L < a$, and the solution is $y = \lceil (L + gz)/a \rceil$. The interval contains a multiple of a exactly when $gz \equiv -x \pmod a$ for some $x \in [L, R]$, that is, when $gz \bmod a = (g \bmod a)z \bmod a$ lies in the stated range. The intervals $[L + gz, R + gz]$ are disjoint and increase with z, since $R - L < g$, so the least z gives the least y. □

The recursion replaces (g, a) by $(a, g \bmod a)$, as in Euclid's algorithm, so it stops after $O(\log g)$ steps; Figure 6 follows one step. It is applied with $g = c$ and $a = -\beta \bmod c$ as follows. Starting from $u = u_0$, let $b = (\alpha - \beta u - v^-) \bmod c$. If $b < H$, then u is in (4.3). Otherwise the next element of (4.3), if it does not exceed u_1, is $u + y$, where y is the least integer ≥ 0 with $c - b \leq ay \bmod c \leq c - b + H - 1$; this window lies inside $[1, c - 1]$, so Lemma 4.3 applies. Every u found in this way is tested by one exact division of N by $r + cu$, and the search resumes at $u + 1$.

4.3. **Choosing the boxes.** The range $0 \leq u \leq U = \lfloor (\lfloor \sqrt{N} \rfloor - r)/c \rfloor$ is cut into consecutive intervals $[u_0, u_1]$, each giving a box $[u_0, u_1] \times [v^-, v^+]$ in Lemma 4.2 (Figure 7). Correctness does not depend on how this is done; the choice only affects the running time. Since $v \approx K/u$ with $K = N/c^2$, a box with $u_1 - u_0 = \lambda u_0$ has area close to $K\lambda^2/(1 + \lambda)$, and as the coset has one point per area c we choose λ with $\lambda^2/(1 + \lambda) = c^3/N$, so that each box holds about one point of (4.2). There are two regimes. If $c^3 > N$, then λ exceeds 1.6 and grows like c^3/N, so that $O\big(\log N / \log(2 + c^3/N)\big)$ boxes cover the whole range: this is Lenstra's range, and the work is a handful of Euclidean recursions. If $c^3 < N$, the boxes shrink to width about $u_0(c^3/N)^{1/2}$, the values of u below $(N/c^3)^{1/2}$ are tested one at a time, and the number of boxes grows like $(N/c^3)^{1/2} \log N$. When this estimate exceeds a fixed threshold we factor N completely instead and read off its divisors in the class.

Theorem 4.4. *Let $(p_1, \ldots, p_{k-3})$ be a node at depth $k - 3$ of the search of Section 3. Every $n = p_1 \cdots p_k$ with $n + \varepsilon = 2\varphi(n)$ extending this node is found by running $s = p_{k-2}$ over the admissible primes of its interval and, for each s, listing the divisors of Proposition 4.1 by Lemmas 4.2 and 4.3 or by a complete factorisation of N.*




FIGURE 6. One step of Lemma 4.3 with $g = 97$, $a = 30$ and $[L, R] = [40, 47]$. Left: the residues $ay \bmod g$ rise in runs, one for each value of $z = \lfloor ay/g \rfloor$, and the first run misses the window because $a\lceil L/a \rceil = 60 > R$. Right: the reduced problem $(30, 7, 13, 20)$ asks for the least z with $7z \bmod 30$ in $[13, 20]$. It is $z = 2$, and the lemma returns $y = \lceil (40 + 2 \cdot 97)/30 \rceil = 8$, for which $ay \bmod g = 46$.

Proof. The prime p_{k-2} lies in the interval and passes the prune, as in the proof of Theorem 3.6. For $s = p_{k-2}$, Proposition 4.1 shows that $t = cp_{k-1} - 2B'$ is one of the divisors listed: Lemma 4.2 shows that $u = (t - r)/c$ lies in the set (4.3) of the box that contains it, and Lemma 4.3 lists every element of that set, or, if N is factored, t is read off from its factorisation. The converse part of Proposition 4.1 then recovers p_{k-1} and p_k. □

Every divisor that is found yields integers $p \le q$ with $A_{k-3}spq + \varepsilon = 2B_{k-3}(s-1)(p-1)(q-1)$, by the converse part of Proposition 4.1, and $n = p_1 \cdots p_{k-3}\, s\, pq$ is a solution exactly when $s < p < q$ and p and q are prime. As before, the correctness of a negative answer depends on no primality proof except in the factoring route, where every prime factor of N is proved prime: below ψ_{13} by the strong probable-prime test to the first thirteen prime bases, which is a proof there by the work of Sorenson and Webster [36], and above it by the primality prover of FLINT [13], which uses the $n-1$ and $n+1$ methods of Brillhart, Lehmer and Selfridge [5] when enough of $n-1$ or $n+1$ can be factored, and otherwise the Jacobi sum test of Adleman, Pomerance and Rumely [1] in the form given by Cohen and Lenstra [9].

4.4. **The sum of the two factors.** The classes of Proposition 4.1 are special: t and N/t lie in the same class modulo c, because $N \equiv r^2$. This allows a third treatment which uses neither boxes nor factoring. Write $t = r + cu$ and $N/t = r + cv$ with $0 \le u \le v$. Since $r' = r$, the integer w of Section 4.2 is $w = (N - r^2)/c$, and (4.1) becomes

$$c\,uv + r\,(u + v) = w, \tag{4.4}$$

so the sum $S = u + v$ lies in the single class $S \equiv wr^{-1} \pmod{c}$, and u and v are the roots of $X^2 - SX + (w - rS)/c$. Conversely, if $S \equiv wr^{-1} \pmod{c}$, then $(w - rS)/c$ is an integer, and if $S^2 - 4(w - rS)/c = e^2$ with an integer $e \ge 0$ of the parity of S, then $u = (S - e)/2$ and $v = (S + e)/2$ are integers satisfying (4.4), so that $(r + cu)(r + cv) = N$. Fix $u_d \ge 1$




FIGURE 7. The divisors of $N = 17670$ in the class 4 modulo 13 up to $\sqrt{N}$, found as in Lemma 4.2. The grey points form the coset (4.2), here $v \equiv 8 - u \pmod{13}$; the curve is the hyperbola (4.1). The boxes cover the curve for $2 \le u \le 9$ (the values $u = 0, 1$ are tested directly), and only the points inside a box are examined. Two of them lie on the curve and give the divisors $t = 30$ and $t = 95$.

and put $t_d = r + cu_d$; if $t_d > \sqrt{N}$ there is no divisor with $u \ge u_d$ to look for. Since $x + N/x$ decreases on $(0, \sqrt{N}\,]$, every divisor $t \le \sqrt{N}$ with $u \ge u_d$ has $2r + cS = t + N/t \le t_d + N/t_d$, while $t + N/t \ge 2\sqrt{N}$. So S runs over at most $N/(c^3 u_d) + 1$ values of one residue class, and each value needs a single test whether $S^2 - 4(w - rS)/c$ is a perfect square. Every divisor with $u \ge u_d$ is found in this way, and by the converse every square found gives a factorisation $N = (r + cu)(r + cv)$. The divisors with $u < u_d$ are found by trial division. With u_d close to $(N/c^3)^{1/2}$ the work is about $2(N/c^3)^{1/2} + 2$ operations. This is comparable with the number of boxes above, but unlike the work in the boxes it does not depend on where the divisors lie.

The same argument bounds the number of divisors.

Proposition 4.5. *In the situation of Proposition 4.1, suppose that $c^3 \ge N + 2c^2$. Then at most one divisor $t \le \sqrt{N}$ of N with $t \equiv r \pmod{c}$ is different from r, and N has at most four positive divisors congruent to r modulo c.*

Proof. By Proposition 4.1, every divisor t of N in the class has $N/t \equiv r^2 r^{-1} = r \pmod{c}$. Let $t_1 < t_2 \le \sqrt{N}$ be divisors of N in the class, both different from r. The hypothesis gives $c^3 > 2c^2$, so $c > 2$ and $0 < r < c$. Hence $t_i = r + cu_i$ with $u_i \ge 1$, and $N/t_i = r + cv_i$ with $v_i \ge u_i$. By (4.4) the sums $u_i + v_i$ lie in one class modulo c, so the numbers $t_i + N/t_i = 2r + c(u_i + v_i)$ differ by a multiple of c^2. The difference is positive, because $x + N/x$ decreases on $(0, \sqrt{N}\,]$, and both numbers lie between $2\sqrt{N}$ and $r+c+N/(r+c) < 2c+N/c$. Hence $c^2 < 2c+N/c-2\sqrt{N} < 2c+N/c$, that is, $c^3 < N + 2c^2$, contrary to the hypothesis. Finally, every divisor of N above $\sqrt{N}$ in the class is N/t for a divisor $t < \sqrt{N}$ in the class, and so there are at most four divisors in the class: r, N/r, t and N/t. □

For classes of this special kind, and at the price of the term $2c^2$, Proposition 4.5 sharpens Lenstra's bound of 11 to four.

We ran this route, too, over every pair of a node at depth $k-3$ and an admissible s. It takes its primes s from a sieve of its own and uses no factorisation, so its negative answers depend on no primality test at all.

5. Lehmer's equation: proof of Theorem 1.1

The last two assertions of Theorem 1.1 were proved in Section 2.1. For the first, let n be composite with $\varphi(n) \mid n-1$ and suppose $\omega(n) \le 15$. By Lemma 2.2, $3 \nmid n$; by Lemma 2.1(i), $n = p_1 \cdots p_k$ with $5 \le p_1 < \cdots < p_k$ and $k \le 15$; and by Lemma 2.3, $n - 1 = 2\varphi(n)$. If $k \le 6$ then $\prod p_i/(p_i-1) \le \prod_{i \le 6} r_i/(r_i-1) = 1.9490\ldots$, which contradicts (3.2). It therefore suffices to run the search with $\varepsilon = -1$ and $p_{\min} = 5$ for $7 \le k \le 15$, and to find nothing. For $k \le 14$ we use the search of Section 3, and for $k = 15$ the method of Section 4.

5.1. **Up to fourteen primes.** Table 2 records the outcome. The column *terminal* counts the nodes at depth $k-2$ whose interval for p_{k-1} is non-empty, and the last column gives the largest width $U_{k-2} - L_{k-2}$ of such an interval. No solution was found for any $k \le 14$, so by Theorem 3.6 there is none. The case $k \le 13$ recovers the theorem of Cohen and Hagis, and the case $k = 14$ is the bound $\omega(n) \ge 15$ announced by Renze and by Pinch.

Table 2. The search for $n - 1 = 2\varphi(n)$ with $5 \le p_1$. Both implementations visit the same number of nodes for every k.

k	nodes	terminal	largest width for p_{k-1}
7	8	0	–
8	18	1	2
9	29	1	3
10	42	1	2
11	73	7	49
12	187	61	298
13	1 092	725	25 854
14	31 962	29 549	54 962 375

For $k = 14$ the numbers of nodes at depths $0, 1, \ldots, 12$ are

$$1,\ 6,\ 16,\ 27,\ 29,\ 23,\ 22,\ 33,\ 58,\ 127,\ 335,\ 1654,\ 29631,$$

as in the row $k = 14$ of Figure 8, and 82 of the 29631 nodes at depth 12 have an empty interval for p_{13}. The largest prime occurring at depth 12 is 77563. Most of the work lies in a few terminal nodes. Exactly fifteen have an interval for p_{13} of width more than three million; they are listed in Table 3. Each of these intervals was enumerated in full, and independently the integer $D = 2AB - C$ was factored completely for each of the fifteen; the factorisations are part of the archived data.

Eleven of the fifteen share the prefix $5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173$, whose product $\prod p/(p-1)$ is already within $8 \cdot 10^{-5}$ of 2; the twelfth prime then closes most of the remaining gap. For the node ending in 25867 one has $T - 1 = C/A = 1.8 \cdot 10^{-8}$, and for the node ending in 5483 one has $T - 1 = 7.0 \cdot 10^{-8}$. These near misses are the reason the width of the last interval grows so quickly with k: by Proposition 3.2 with $m = 2$ the interval for p_{k-1} runs from about A/C to about $2A/C$, and a small C means a long interval. For the node ending in 25867 this

TABLE 3. The fifteen terminal nodes of the case $k = 14$ with an interval for p_{13} of width more than $3 \cdot 10^6$, with the number of digits of $D = 2AB - C$.

$p_1, \ldots, p_{12}$	width	digits of D
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25867	54 962 375	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25889	19 589 597	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 73, 317, 5483	14 242 697	40
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25903	13 903 362	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25913	11 517 582	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25919	10 443 048	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25933	8 577 304	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25939	7 967 703	41
5, 7, 13, 17, 19, 23, 37, 59, 83, 97, 107, 3343	6 121 542	39
5, 7, 13, 17, 19, 23, 37, 59, 67, 97, 199, 577	6 086 511	38
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 25969	5 880 810	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 26003	4 537 275	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 26017	4 147 802	41
5, 7, 13, 17, 19, 23, 37, 59, 67, 89, 173, 26029	3 863 803	41
5, 7, 13, 17, 19, 23, 37, 59, 83, 109, 173, 199	3 320 179	37




FIGURE 8. Number of nodes at each depth j of the search for $k = 7, \ldots, 15$ primes, with the colour on a logarithmic scale. The counts are the same for $n - 1 = 2\varphi(n)$ and $n + 1 = 2\varphi(n)$ with $p_1 \geq 5$. The tree stays small until the last two levels. The last cell of each row counts the nodes at depth $k - 2$: for $k \leq 14$ these are the nodes where p_{k-1} is enumerated, and for $k = 15$ (framed) they are the 33865004 choices of p_{13}, each completed by the method of Section 4.

gives exactly the interval [54962376, 109924751]. Among the fifteen integers D one, for the node ending in 25913, is itself a prime of 41 digits[9].

[9]For this node (3.5) has only the trivial factorisations $D = 1 \cdot D$ and $D = D \cdot 1$, and neither gives a pair of primes in the required range. The enumeration of the $1.15 \cdot 10^7$ candidates for p_{13} reaches the same conclusion without knowing that D is prime.




FIGURE 9. The 33865004 two-prime problems of the case $k = 15$, by the ratio c^3/N on a logarithmic scale (bins of one decade). To the right of the dashed line is Lenstra's range, where N has at most 11 divisors in the class and a few boxes find them. The left tail comes from the lower ends of the longest intervals for p_{13}.

5.2. **Fifteen primes.** For $k = 15$ the traversal of Section 3 is stopped at depth 12. The numbers of nodes at depths $0, 1, \ldots, 12$ are

$$1,\ 6,\ 18,\ 30,\ 34,\ 37,\ 41,\ 56,\ 103,\ 218,\ 616,\ 3227,\ 55125,$$

and 54985 of the nodes at depth 12 have a non-empty interval for p_{13}. Both implementations of the search produce the same 54985 prefixes with the same intervals, and so do the bounds for $\varepsilon = +1$. The intervals have total length $1.005 \cdot 10^9$ and contain 33865004 primes s that pass the congruence prune. For each of these pairs of a prefix and a prime $s = p_{13}$ we solved the two-prime problem of Proposition 4.1, in which N has between 40 and 58 digits, three times: by the two implementations of the lattice method (Section 4.2) and by the route of Section 4.4. The first lattice implementation factored N completely in the 473 cases where the estimated number of boxes exceeded its threshold and took 19 minutes of processor time in all; the second factored N in 1772 cases and took 45 minutes; the route of Section 4.4, which factors nothing, took 13 minutes. The three agree on each of the 55443 pieces into which the work was divided.

Figure 9 shows how the pairs are distributed according to the ratio c^3/N, which by Section 4 decides how hard the two-prime problem is. For 33786783 of them, all but 0.23 per cent, the modulus satisfies $c^3 > N$, so that Lenstra's bound applies and a few boxes suffice. The remaining 78221 pairs come from values of s just above the lower end of long intervals, where $c = C_{12}s - 2B_{12}$ is small. The most extreme is the prefix ending in 25867 of Table 3, whose first admissible s gives $c^3/N \approx 2 \cdot 10^{-15}$. For these pairs the route of Section 4.4 needs about $2(N/c^3)^{1/2}$ operations.

For none of the 33865004 pairs does N have a divisor in the class of Proposition 4.1 that yields integers $s < p < q$, and the three computations agree on this. So there is not even a completion by composite integers, and in particular no solution with $k = 15$. Together with the case $k \le 14$ this proves Theorem 1.1.

5.3. **Solutions below 10^{35}.** The search of Section 3 also decides whether a solution lies below a given bound X, for a given number of prime factors, because n is controlled from below at every node.

Proposition 5.1. *Let $n = p_1 \cdots p_k$ satisfy (3.1), let $0 \le j \le k-2$, $m = k-j$, and let L_j be the lower end of the interval of Proposition 3.2. Then $n \ge A_j L_j^m$ and $p_{j+1} \le (n/A_j)^{1/m}$.*

Proof. We have $n = A_j \prod_{i>j} p_i$, and the m primes p_i with $i > j$ are at least $p_{j+1} \ge L_j$; so $n \ge A_j L_j^m$ and $p_{j+1}^m \le n/A_j$. □

Fix X and add the following rule to the traversal: a node $(p_1, \dots, p_j)$ with $A_j L_j^m \ge X$ is discarded, and the candidates p_{j+1} at a node that is kept are restricted to $p_{j+1} \le \lfloor (X/A_j)^{1/m} \rfloor$. If $n < X$ is a solution, every node $(p_1, \dots, p_j)$ with $j \le k-2$ is kept and p_{j+1} is a candidate at it, by Proposition 5.1, so the proof of Theorem 3.6 applies without change: the traversal records every solution below X. The gain is that the interval for p_{j+1}, whose upper end U_j grows like $(m-1)A_j/C_j$, is now cut at $(X/A_j)^{1/m}$, and that the nodes with small C_j/A_j, which make the unrestricted search hard (Section 6.6), are discarded as soon as their product is large.

Proof of Theorem 1.3. Let n be a composite solution of (1.1) with k prime factors. By Theorem 1.1, $k \ge 16$. The product of the 23 smallest primes from 5 on exceeds 10^{35}, so we may assume $k \le 22$; then $M = 2$ and $3 \nmid n$ by Corollary 2.6, and $n = p_1 \cdots p_k$ with $5 \le p_1 < \cdots < p_k$ and $n - 1 = 2\varphi(n)$. The search with $\varepsilon = -1$, $p_{\min} = 5$ and $X = 10^{35}$ was run for $16 \le k \le 22$ and found nothing (Table 4). □

TABLE 4. The search for $n - 1 = 2\varphi(n)$ with $5 \le p_1$ below $X = 10^{35}$. The column *terminal* counts the nodes at depth $k-2$ with a non-empty interval for p_{k-1}, and the last column gives the largest width of such an interval.

k	nodes	discarded by size	terminal	largest width for p_{k-1}
16	426 178 345	80 176 394	342 122 117	546 852
17	1 351 753 636	340 139 845	970 083 570	56 098
18	121 353 478	37 805 731	69 514 261	5 842
19	261 802	76 098	64 034	722
20	4 890	286	0	–
21	1 055	33	0	–
22	715	25	0	–

The counts of Table 4 come from the program of Section 5 with the two rules added, written in C with the GNU multiple precision library and a segmented sieve for the candidates p_{k-1}; the searches for $k = 16$ and $k = 17$ were divided into four and two parts, by the order in which the nodes at depth 9 are created, and the table adds the parts. The whole search took about 5 hours of processor time. For $k = 16$ the cost grows by a factor of about 5 for each power of 10 in X: the searches below 10^{30}, 10^{31}, 10^{32}, 10^{33} and 10^{34} visit 3217136, 13293061, 38572632, 94527876 and 213687046 nodes; for $k = 17$ the factor is larger, about 14, and $k = 17$ costs more than $k = 16$ from $X = 10^{34}$ on, while $k \ge 18$ costs little. A second program, written independently in Python with exact integer arithmetic, runs the same search; for $k = 16$ it visits the same numbers of nodes and terminal nodes and tests the same numbers of candidates below 10^{30} and 10^{31}, and for $17 \le k \le 30$ below 10^{32}. The search below 10^{35} was run by the first program alone.

6. The size of a solution

This section proves Theorem 1.2. Of what precedes it uses only Section 2, and unlike Sections 3 to 5 it does not assume $M = 2$. Throughout, $n = p_1 \cdots p_k$ is a composite solution of (1.1) with

$p_1 < \cdots < p_k$, so that $\varepsilon = -1$ and $M = (n-1)/\varphi(n)$, and A_j, B_j and P_j are as in the Notation of Section 1; thus $B_j = \varphi(A_j)$. By Lemma 2.1, n is odd and squarefree, $k \geq 2$, $M \geq 2$, and no prime factor of n divides $p-1$ for a prime factor p of n. By Lemma 2.2, if $3 \mid n$ then $p_1 = 3$ and $M \equiv 1 \pmod 3$, so that $M \geq 4$.

Lemma 6.1. (i) $P_1 < P_2 < \cdots < P_{k-1} < M < P_k = M + 1/\varphi(n)$.
(ii) $p_k = (MB_{k-1} - 1)/(MB_{k-1} - A_{k-1})$.

Proof. The equality $P_k = M + 1/\varphi(n)$ is Lemma 2.1(iii), and P_j increases with j. Writing $n - 1 = M\varphi(n)$ as $A_{k-1}p_k - 1 = MB_{k-1}(p_k - 1)$ gives $p_k\,(MB_{k-1} - A_{k-1}) = MB_{k-1} - 1 > 0$. Hence $MB_{k-1} > A_{k-1}$, that is $P_{k-1} < M$, and (ii) follows. □

6.1. **A sharp product lemma.** For integers $m \geq 1$ and real $y \geq 1$ put

$$E_m(y) = y^{2^m} - y^{2^{m-1}}.$$

The function E_m is increasing on $y \geq 1$, satisfies $E_m(y) < y^{2^m}$, and

$$E_m(y^{2^u}) = E_{m+u}(y) \qquad (m \geq 1,\ u \geq 0). \tag{6.1}$$

Theorem 6.2. *Let $a < b$ be positive integers, $d = b - a$, $m \geq 2$, and let $2 \leq x_1 \leq x_2 \leq \cdots \leq x_m$ be integers with*

$$\prod_{i=1}^{m}\Big(1 - \frac{1}{x_i}\Big) \leq \frac{a}{b} < \prod_{i=1}^{m-1}\Big(1 - \frac{1}{x_i}\Big). \tag{6.2}$$

Then

$$a\,x_1x_2\cdots x_m \leq E_{m-1}\Big(\frac{b(a+1)}{d}\Big).$$

If d divides $b+1$, equality holds for a suitable choice of the x_i.

For $d = 1$ the bound is $E_{m-1}((a+1)^2) = E_m(a+1)$, which is the lemma of Cook and Nielsen (Lemma 6.5 below) behind the bound of Burek and Żmija. For $d \geq 2$ it is smaller, and it improves the bound $E_{m-1}(b^2/d)$ of Stone [37, Corollary 1], since $b(a+1) < b^2$. Applied to the prime factors of n that follow the first j, Theorem 6.2 is strong whenever P_j is still well below M.

Throughout this subsection $a < b$ are positive integers, $d = b - a$, and $x_1 \leq \cdots \leq x_m$ are integers at least 2 satisfying (6.2). We write

$$X_u = x_1 \cdots x_u, \qquad Y_u = (x_1 - 1)\cdots(x_u - 1),$$

with $X_0 = Y_0 = 1$, and use the same notation $Z_u = z_1 \cdots z_u$ for other sequences.

Lemma 6.3. *For $0 \leq u \leq m-1$ the tail $x_{u+1}, \ldots, x_m$ satisfies (6.2) with (a, b) replaced by (aX_u, bY_u), and $aX_u < bY_u$.*

Proof. Divide (6.2) by $\prod_{i \leq u}(1 - 1/x_i) = Y_u/X_u$. Since $u \leq m-1$, we have $Y_u/X_u \geq \prod_{i \leq m-1}(1 - 1/x_i) > a/b$. □

The next lemma is a majorisation inequality; only the sequence x needs to be ordered.

Lemma 6.4. *Let $x_1 \leq \cdots \leq x_m$ and $z_1, \ldots, z_m$ be real numbers greater than 1 with $X_u \geq Z_u$ for $1 \leq u \leq m$. Then*

$$\prod_{i=1}^{m}\Big(1 - \frac{1}{x_i}\Big) \geq \prod_{i=1}^{m}\Big(1 - \frac{1}{z_i}\Big),$$

with strict inequality if $X_m > Z_m$.

Proof. The function $f(t) = \log(1-e^{-t})$ on $t > 0$ has derivative $f'(t) = 1/(e^t-1)$, which is positive and decreasing, so f is concave. Put $\tau_i = \log x_i$ and $\sigma_i = \log z_i$, so that $f(\tau_i) = \log(1 - 1/x_i)$ and $f(\sigma_i) = \log(1 - 1/z_i)$. By concavity

$$f(\sigma_i) \le f(\tau_i) + c_i(\sigma_i - \tau_i), \qquad c_i = f'(\tau_i) = \frac{1}{x_i - 1},$$

and $c_1 \ge c_2 \ge \cdots \ge c_m > 0$ because the x_i are nondecreasing. The partial sums $\Delta_u = \sum_{i\le u}(\sigma_i - \tau_i) = \log(Z_u/X_u)$ are at most 0 by hypothesis, and summation by parts gives

$$\sum_{i=1}^{m} c_i(\sigma_i - \tau_i) = \sum_{u=1}^{m-1}(c_u - c_{u+1})\Delta_u + c_m\Delta_m \le c_m\Delta_m \le 0,$$

with $c_m\Delta_m < 0$ when $X_m > Z_m$. Hence $\sum f(\sigma_i) \le \sum f(\tau_i)$, strictly in that case, and the lemma follows on exponentiating. □

Lemma 6.5 (Cook, Nielsen)**.** *Under (6.2), with $m \ge 1$, we have $aX_m \le E_m(a+1)$.*

This is the lemma of Cook [10] and Nielsen [28] mentioned in the introduction. We include its proof, on which the proof of Theorem 6.2 builds.

Proof. We use induction on m. Define

$$z_u = (a+1)^{2^{u-1}} + 1 \quad (1 \le u \le m-1), \qquad z_m = (a+1)^{2^{m-1}}.$$

Induction on u gives $aZ_u = (a+1)^{2^u} - 1$ for $0 \le u \le m-1$, and then $aZ_m = E_m(a+1)$. Writing $c_u = (a+1)^{2^{u-1}}$, so that $c_{u+1} = c_u^2$ and $1 - 1/z_u = c_u(c_u-1)/(c_{u+1}-1)$ for $u < m$, the product telescopes:

$$\prod_{u=1}^{m}\Big(1 - \frac{1}{z_u}\Big) = \frac{(a+1)^{2^{m-1}-1}\,a}{c_m - 1} \cdot \frac{c_m - 1}{c_m} = \frac{a}{a+1}. \tag{6.3}$$

These identities use only that a is a positive real number, and we shall apply them once more with a value of a that need not be an integer.

Suppose first that $X_u \le Z_u$ for some $1 \le u \le m-1$, so that $aX_u + 1 \le (a+1)^{2^u}$. By Lemma 6.3 and the induction hypothesis applied to the $m-u$ terms $x_{u+1}, \dots, x_m$ with numerator aX_u, and by (6.1),

$$aX_m = aX_u \prod_{i>u} x_i \le E_{m-u}(aX_u + 1) \le E_{m-u}((a+1)^{2^u}) = E_m(a+1).$$

Otherwise $X_u > Z_u$ for every $u \le m-1$ (for $m = 1$ this is vacuous). If also $X_m > Z_m$, Lemma 6.4 and (6.3) give $\prod(1 - 1/x_i) > a/(a+1) \ge a/b$, contrary to (6.2). Hence $X_m \le Z_m$ and $aX_m \le E_m(a+1)$. □

The proof compares $x_1, \dots, x_m$ with an extremal sequence $z_1, \dots, z_m$ through the partial products X_u and Z_u, and splits according to whether $X_u \le Z_u$ for some $u \le m-1$ (Figure 10).

Proof of Theorem 6.2. Put $G = b(a+1)/d$, so that $G - 1 = a(b+1)/d$, and define real numbers

$$z_1 = \frac{b+1}{d}, \qquad z_u = G^{2^{u-2}} + 1 \quad (2 \le u \le m-1), \qquad z_m = G^{2^{m-2}}.$$

Then $aZ_1 = G - 1$, induction gives $aZ_u = G^{2^{u-1}} - 1$ for $1 \le u \le m-1$, and $aZ_m = (G^{2^{m-2}} - 1)G^{2^{m-2}} = E_{m-1}(G)$. The numbers $z_2, \dots, z_m$ are the sequence of the proof of Lemma 6.5 with $G-1$ in place of a and $m-1$ terms, so by (6.3)

$$\prod_{u=1}^{m}\Big(1 - \frac{1}{z_u}\Big) = \frac{a+1}{b+1} \cdot \frac{G-1}{G} = \frac{a+1}{b+1} \cdot \frac{a(b+1)}{b(a+1)} = \frac{a}{b}.$$




FIGURE 10. The proof of Theorem 6.2 for $a/b = 5/8$ and $m = 5$, on the scale $\log_2 \log_2(aX_u + 1)$. The extremal sequence $3, 17, 257, 65537, 2^{32}$ rises by exactly 1 per step from $u = 1$ on and ends at $aZ_5 = E_4(16)$. The two other sequences satisfy (6.2), start above it, and fall below it at $u = 2$ and $u = 3$. From that point Lemmas 6.3 and 6.5 give $aX_5 < (aX_u + 1)^{2^{5-u}}$, the dashed line of slope 1, which ends below $E_4(16)$. A sequence that stayed above the extremal one up to $u = 4$ would end below it at $u = 5$, by Lemma 6.4.

Suppose first that $X_u \le Z_u$ for some $1 \le u \le m-1$, so that $aX_u + 1 \le G^{2^{u-1}}$. Lemmas 6.3 and 6.5, applied to the $m - u \ge 1$ terms $x_{u+1}, \dots, x_m$, and (6.1) give

$$aX_m \le E_{m-u}(aX_u + 1) \le E_{m-u}(G^{2^{u-1}}) = E_{m-1}(G).$$

Otherwise $X_u > Z_u$ for all $u \le m-1$. If also $X_m > Z_m$, then Lemma 6.4, which applies because every z_u exceeds 1 (for z_1 because $b + 1 > d$), gives $\prod(1 - 1/x_i) > \prod(1 - 1/z_i) = a/b$, contrary to (6.2). Hence $X_m \le Z_m$, that is $aX_m \le E_{m-1}(G)$.

If $d \mid b+1$, then z_1 is an integer greater than 1, so $z_1 \ge 2$, and $G - 1 = az_1$, $G \ge 3$ and all z_u are integers. They are nondecreasing: $z_2 \ge G \ge z_1$, since $G \ge z_1$ amounts to $ab \ge 1$; the middle terms increase; and for $m \ge 3$ we have $z_m = (z_{m-1} - 1)^2 \ge z_{m-1}$ because $z_{m-1} - 1 \ge G \ge 3$. The product of $1 - 1/z_u$ is a/b over all m terms and larger over the first $m - 1$, so the z_u satisfy (6.2) and attain $aZ_m = E_{m-1}(G)$. □

Remark 6.6. Since $G = (a+1)/(1 - a/b)$, the bound of Theorem 6.2 depends on b only through the distance of a/b from 1. Its base is smaller than the base $(a+1)^2$ of Lemma 6.5 by the factor $(a+1)d/b$, which is at least 1 because $(a+1)d - b = a(d-1) \ge 0$; Figure 11 shows the two bases. Stone's bound $E_{m-1}(b^2/d)$ exceeds ours by the factor $b/(a+1)$ in the base. In the application below $a = A_j$ and $b = MB_j$, and then the base $G = (A_j + 1)/(1 - P_j/M)$ decreases as M increases, so the smallest admissible value of M yields an upper bound. The




FIGURE 11. The base of the bound for aX_m written as $E_{m-1}(\,\cdot\,)$, on a $\log_2$ scale, over the numerator a and the deficit $d = b - a$. The wire mesh is the base $(a+1)^2$ of the lemma of Cook and Nielsen, the shaded sheet below it the base $b(a+1)/d$ of Theorem 6.2. They meet along $d = 1$, where the two bounds coincide, and separate as d grows.

base $b^2/d = M^2B_j^2/(MB_j - A_j)$ of Stone has no such property: it decreases for $M < 2P_j$ and increases for $M > 2P_j$.

Remark 6.7. The extremal sequences of Theorem 6.2 are explicit. For $a/b = 5/8$ we have $d = 3$, $z_1 = 3$ and $G = 16$, and with $m = 4$ the sequence is $3, 17, 257, 65536$:

$$\frac{2}{3}\cdot\frac{16}{17}\cdot\frac{256}{257}\cdot\frac{65535}{65536} = \frac{5}{8}, \qquad 5\cdot 3\cdot 17\cdot 257\cdot 65536 = 2^{32} - 2^{16} = E_3(16).$$

The first three terms are Fermat primes. For $a = 1$ the sequence of the proof of Lemma 6.5 consists of the Fermat numbers $F_0, \dots, F_{m-2}$ followed by $2^{2^{m-1}}$, and the products $F_0F_1\cdots F_{r-1} = 2^{2^r} - 1$ with $1 \le r \le 5$ are the solutions 3, 15, 255, 65535 and 4294967295 of (1.2) in (1.3). Such products have $n/\varphi(n)$ just below 2, and so do the deepest nodes of the search of Section 6.3 (Section 6.6).

6.2. **Heavy solutions.** For $0 \le j \le k-1$ put

$$V_j = V_j(M) = \frac{MB_j(A_j+1)}{MB_j - A_j} = \frac{A_j+1}{1 - P_j/M}. \tag{6.4}$$

By Lemma 6.1(i) the denominator is positive, and $V_j(M)$ decreases as M increases.

Proposition 6.8. *For $0 \le j \le k-1$,*

$$n < (A_j+1)^{2^{k-j}} \qquad \text{and} \qquad n < V_j^{2^{k-j-1}}.$$

Proof. Take $m = k-j$, $x_i = p_{j+i}$, $a = A_j$ and $b = MB_j$. Then

$$\prod_{i=j+1}^{k}\Big(1-\frac{1}{p_i}\Big) = \frac{P_j}{P_k} < \frac{P_j}{M} = \frac{a}{b} < \frac{P_j}{P_{k-1}} = \prod_{i=j+1}^{k-1}\Big(1-\frac{1}{p_i}\Big)$$

by Lemma 6.1(i), so (6.2) holds, $a < b$, and $ax_1\cdots x_m = n$. Lemma 6.5 gives $n \le E_{k-j}(A_j+1)$, which implies the first bound. For $j \le k-2$, Theorem 6.2 gives $n \le E_{k-j-1}(V_j)$, because $b(a+1)/d = V_j$. For $j = k-1$, Lemma 6.1(ii) gives $n = A_{k-1}p_k < A_{k-1}MB_{k-1}/(MB_{k-1} - A_{k-1}) < V_{k-1}$. □

Definition 6.9. Let $s \ge 0$ be an integer, and put $\theta_i = 2^{2^{i-s}}$ for $i \ge s$ and $\theta_i = 1$ for $i < s$. A composite solution n of (1.1) is *s-heavy* if for every j with $0 \le j \le k-1$

(H1) $A_j \ge \theta_j$, and
(H2) $V_j \ge \theta_{j+1}$.

Corollary 6.10. *A composite solution of (1.1) with k prime factors that is not s-heavy satisfies $n < 2^{2^{k-s}}$.*

Proof. If (H1) fails at j, then $j \ge s$ and $A_j + 1 \le 2^{2^{j-s}}$, and Proposition 6.8 gives $n < (A_j+1)^{2^{k-j}} \le 2^{2^{k-s}}$. If (H2) fails at j, then $j+1 \ge s$ because $V_j > 1$, and $n < V_j^{2^{k-j-1}} < 2^{2^{k-s}}$. □

For $s = 2$ condition (H1) alone contradicts Lemma 6.1(i), and the argument can be carried out by hand.

Proposition 6.11. *No composite solution of (1.1) is 2-heavy. Consequently every composite solution with k prime factors satisfies $n < 2^{2^{k-2}}$.*

Proof. Let n be 2-heavy, and let $1 \le i \le k-1$. Condition (H1) at $j = i$ and $A_i \le p_i^i$ give $p_i \ge \theta_i^{1/i} = 2^{2^{i-2}/i}$. Hence $p_7 \ge 24$, $p_8 \ge 256$, $p_9 > 19000$ and $p_i > 4^i$ for $i \ge 10$, so that

$$\sum_{8\le i\le k-1}\frac{1}{p_i-1} < \frac{1}{256} + \frac{1}{19000} + \sum_{i\ge 10}\frac{1}{4^i-1} < 0.004.$$

Put $f(x) = x/(x-1)$, which decreases for $x > 1$ and satisfies $f(x) \le e^{1/(x-1)}$. Then $P_k = \prod_{i\le k} f(p_i)$ and $f(p_k) < f(p_{k-1})$. Let $b_1, \dots, b_7$ be lower bounds for $p_1, \dots, p_7$ and $\beta_j = \prod_{i\le j} f(b_i)$. If $k \le 8$, then $P_k < \beta_{k-1}f(b_{k-1})$. If $k \ge 9$, the prime factors $p_8, \dots, p_k$ contribute less than $e^{0.004}f(p_{k-1}) < e^{0.008}$, and $P_k < \beta_7 e^{0.008}$. We distinguish four cases.

(a) If $3 \mid n$, then $p_1 = 3$ and $M \ge 4$ by Lemma 2.2, and the next prime factors are at least $5, 7, 11, 13, 17$ and, as $p_7 \ge 24$, 29.
(b) If $p_1 = 5$ and $p_2 = 7$, then no other prime factor is congruent to 1 modulo 5 or 7 by Lemma 2.1(ii), which excludes 11, 29 and 31.
(c) If $p_1 = 5$ and $p_2 \ne 7$, then $p_2 \ge 13$, and 11 and 31 are excluded because they are congruent to 1 modulo 5.
(d) If $p_1 \ge 7$, the prime factors are at least the primes from 7 on, and $p_7 \ge 29$.

In cases (b) to (d), $3 \nmid n$ and $M \ge 2$. Table 5 lists the bounds $b_1, \dots, b_7$ in each case and the larger of $\max_{j\le 7}\beta_j f(b_j)$ and $\beta_7 e^{0.008}$. In every case it is less than M, so that $P_k < M$, contrary to Lemma 6.1(i). Hence no composite solution is 2-heavy, and Corollary 6.10 with $s = 2$ gives the bound. □

TABLE 5. The four cases in the proof of Proposition 6.11: lower bounds $b_1, \dots, b_7$ for the seven smallest prime factors and the resulting upper bound for P_k, rounded up.

case	$b_1, \dots, b_7$	bound for P_k	$M \geq$
(a) $p_1 = 3$	$3, 5, 7, 11, 13, 17, 29$	2.972	4
(b) $p_1 = 5$, $p_2 = 7$	$5, 7, 13, 17, 19, 23, 37$	1.957	2
(c) $p_1 = 5$, $p_2 \neq 7$	$5, 13, 17, 19, 23, 29, 37$	1.738	2
(d) $p_1 \geq 7$	$7, 11, 13, 17, 19, 23, 29$	1.749	2

The proof uses only condition (H1), whose failure gives the bound through the lemma of Cook and Nielsen, and the fact that no prime factor divides another minus one. For $s = 1$ the same argument needs only the cases $3 \mid n$ and $3 \nmid n$, with the primes from 5 on as lower bounds and $p_6 \geq 41$, $p_7 \geq 569$, $p_8 \geq 65537$ from (H1); it shows that no composite solution is 1-heavy, which gives $n < 2^{2^{k-1}}$, half the number of binary digits allowed by the bound of Burek and Żmija. For larger s the estimates of this proof are too weak, and Theorem 1.2(i) follows once we know that no solution is 7-heavy, which Section 6.3 establishes with exact bounds at every node and with condition (H2) and Theorem 6.2 as well.

We first record how fast the prime factors of a heavy solution grow. Fix s, an index $j \geq 0$ and the product A_j of j odd primes. For $i > j$ let λ_i be the least positive integer λ with $A_j \lambda^{i-j} \geq \theta_i$, and put $y_i = (\theta_i / A_j)^{1/(i-j)}$. Then $\lambda_i = \max\{1, \lceil y_i \rceil\}$, so $\lambda_i \geq y_i$, and $\lambda_i < y_i + 1$ whenever $\lambda_i > 1$.

Lemma 6.12. *Let $i > j$.*

(i) *If $i \geq j + 2$ and $y_i > 1$, then $y_{i+1} \geq y_i^{4/3}$.*
(ii) *If n is an s-heavy solution, A_j is the product of its j smallest prime factors and $i \leq k - 1$, then $p_i \geq \lambda_i$.*

Proof. (i) If $y_i > 1$, then $\theta_i > A_j \geq 1$, so $i \geq s$ and $\theta_{i+1} = \theta_i^2$. Hence

$$(i + 1 - j) \log y_{i+1} = 2 \log \theta_i - \log A_j \geq 2(\log \theta_i - \log A_j) = 2(i - j) \log y_i,$$

so $\log y_{i+1} \geq \frac{2(i-j)}{i+1-j} \log y_i \geq \frac{4}{3} \log y_i$, since $i - j \geq 2$ and $\log y_i > 0$.

(ii) By (H1) at i, $\theta_i \leq A_i \leq A_j p_i^{i-j}$, so $p_i \geq \lambda_i$. □

The search needs an upper bound for P_k / P_j that uses only $p_1, \dots, p_j$ and a lower bound for p_{j+1}. For an integer $q \geq 3$ and $i > j$ let $\pi_i(q)$ be the $(i - j)$-th smallest prime that is at least q, if this prime is less than $2 \cdot 10^7$, and $\pi_i(q) = q + 2(i - j - 1)$ otherwise. Put $b_i = \max\{\pi_i(q), \lambda_i\}$, let i_0 be the least index $i \geq j + 3$ with $\lambda_i > 10^{30}$, and define

$$\Gamma_j(q) = \max\Big\{ \frac{q}{q-1}, \max_{j+1 \leq i < i_0} \frac{b_i + 2}{b_i + 1} \prod_{l=j+1}^{i} \frac{b_l}{b_l - 1}, \Big(1 + \frac{16}{\lambda_{i_0} - 2}\Big) \prod_{l=j+1}^{i_0 - 1} \frac{b_l}{b_l - 1} \Big\}. \tag{6.5}$$

The quantity $\Gamma_j(q)$ depends only on j, A_j, q and s, and it is a finite computation because λ_i grows doubly exponentially in i.

Lemma 6.13. *Let n be an s-heavy solution, $0 \leq j \leq k - 1$, and let q be an integer with $3 \leq q \leq p_{j+1}$. Then $P_k \leq P_j \Gamma_j(q)$.*

Proof. The quotient P_k / P_j is the product of $p_i / (p_i - 1)$ over $j < i \leq k$, and $x / (x - 1)$ decreases in x. For $j < i \leq k - 1$ we have $p_i \geq \pi_i(q)$, since $p_{j+1} < \cdots < p_i$ are odd primes at least q, and $p_i \geq \lambda_i$ by Lemma 6.12(ii); so $p_i \geq b_i$. For the last prime, $p_k \geq q$ if $k = j + 1$ and

$p_k \ge p_{k-1}+2 \ge b_{k-1}+2$ otherwise. This bounds P_k/P_j by the first term of (6.5) when $k = j+1$, and by the second term, with $i = k-1$, when $j+2 \le k \le i_0$.

Suppose $k > i_0$. The primes $p_{j+1}, \dots, p_{i_0-1}$ contribute at most $\prod_{l=j+1}^{i_0-1} b_l/(b_l-1)$, as before. Put $Y = y_{i_0}$. Since $\lambda_{i_0} > 10^{30}$, we have $Y > \lambda_{i_0} - 1 \ge 10^{30}$, and Lemma 6.12 gives $p_i \ge \lambda_i \ge y_i \ge Y^{(4/3)^{i-i_0}}$ for $i_0 \le i \le k-1$. As $p_k > p_{k-1}$, the factor of p_k is at most that of p_{k-1}, so

$$\prod_{i=i_0}^{k} \frac{p_i}{p_i-1} \le \prod_{i=i_0}^{k-1} \Big(1+\frac{1}{p_i-1}\Big)^2 \le \exp\Big(2\sum_{u\ge 0} \frac{1}{Y^{(4/3)^u}-1}\Big).$$

Because $(4/3)^u \ge 1+u/3$ and $Y \ge 10^{30}$, the terms with $u \ge 1$ add up to less than $1/(Y-1)$, so the exponent is less than $4/(Y-1) \le 1$. With $e^x \le 1+2x$ for $0 \le x \le 1$ the product is at most $1+8/(Y-1) < 1+8/(\lambda_{i_0}-2)$, which is smaller than the factor $1+16/(\lambda_{i_0}-2)$ of the third term of (6.5). □

6.3. **The search.** Fix s. Let $S = (p_1, \dots, p_j)$, $j \ge 0$, be an increasing sequence of odd primes in which no p_i divides any $p_l - 1$, and write A_j, B_j and P_j as before. Let $\mu(S)$ be the least integer $M > P_j$ with $M \ge 2$ and, if $p_1 = 3$, $M \equiv 1 \pmod 3$, and let $\mu'(S)$ be the next such integer. We say that a solution n *extends* S if $j \le k-1$ and S consists of the j smallest prime factors of n. By Lemmas 2.1, 2.2 and 6.1(i), the quotient M of such an n satisfies $M \ge \mu(S)$. For $M > P_j$ we use $V_j(M)$ as in (6.4), computed from S.

The tree $\mathcal{T}_s$ is built from the empty sequence by the following rules, applied to each sequence S of length j that is generated; Figure 12 summarises them.

(R1) *Ratio test.* Let q_1 be the least prime greater than p_j ($q_1 = 3$ if $j = 0$). If $P_j\Gamma_j(q_1) \le \mu(S)$, the sequence S is discarded.

(R2) *Deficit test.* If $V_j(\mu(S)) < \theta_{j+1}$, the sequence S is discarded.

A sequence that survives both tests is a *node* of $\mathcal{T}_s$. At a node:

(R3) *Last prime.* If $j \ge 1$, then for each integer M with $\mu(S) \le M < P_j(p_j+2)/(p_j+1)$, and $M \equiv 1 \pmod 3$ if $p_1 = 3$, test whether $(MB_j-1)/(MB_j-A_j)$ is a prime greater than p_j that is not 1 modulo any p_i.

(R4) *Children.* Put $\mu = \mu(S)$ and $q_c = \lfloor \mu B_j/(\mu B_j - A_j)\rfloor$. Generate the sequences $(p_1, \dots, p_j, q)$ for the primes $q > p_j$ with $q \not\equiv 1 \pmod{p_i}$ for $i \le j$ and with $A_j q \ge \theta_{j+1}$, taking q in increasing order and omitting q in the following cases.
   (i) If $P_j\Gamma_j(q) \le \mu$, omit q and every larger prime.
   (ii) If $q \le q_c$ and $P_j\Gamma_j(q) \le \mu'(S)$, omit every prime in $[q, q_c]$.
   (iii) If $q > q_c$ and $V_{j+1}(\mu)$, computed for the sequence $(p_1, \dots, p_j, q)$, is less than θ_{j+2}, omit q.

A prime q satisfies $q \le q_c$ exactly when $P_j q/(q-1) \ge \mu$, so in (ii) a solution extending $(p_1, \dots, p_j, q)$ has $P_{j+1} \ge \mu$ and hence $M \ge \mu'(S)$. For $q > q_c$ we have $P_{j+1} < \mu$, so $V_{j+1}(\mu)$ is defined, and after multiplication by its positive denominator the inequality in (iii) becomes a quadratic inequality in q with positive leading coefficient;[10] the primes omitted by (iii) therefore form an interval, which is computed once per node.

Proposition 6.14. *Let n be an s-heavy solution. Then $(p_1, \dots, p_{k-1})$ is a node of $\mathcal{T}_s$, and rule (R3) at that node finds p_k.*

Proof. Write $S_j = (p_1, \dots, p_j)$. We show by induction on j that S_j is generated and is a node, for $0 \le j \le k-1$. The empty sequence S_0 is generated by definition. Suppose that S_j is

[10] With $D = \mu B_j - A_j$ and $\theta = \theta_{j+2}$ one has $V_{j+1}(\mu) = \mu B_j(q-1)(A_jq+1)/(Dq-\mu B_j)$, and $V_{j+1}(\mu) < \theta$ reads $\mu A_jB_jq^2 + \big(\mu B_j - \mu A_jB_j - \theta D\big)q + (\theta-1)\mu B_j < 0$.




FIGURE 12. The treatment of a sequence $S = (p_1, \dots, p_j)$ in the tree $\mathcal{T}_s$. A sequence that fails the ratio test or the deficit test cannot be extended by an s-heavy solution. A node is tested for a last prime factor and passes to its children every prime that the three omission rules allow.

generated. As n extends S_j, we have $M \ge \mu(S_j)$. Lemma 6.13 with $q = q_1 \le p_{j+1}$ gives $P_j\Gamma_j(q_1) \ge P_k > M \ge \mu(S_j)$, so (R1) does not discard S_j, and $V_j(\mu(S_j)) \ge V_j(M) \ge \theta_{j+1}$ by (H2), so (R2) does not either. Thus S_j is a node.

Now let $j \le k-2$. The prime p_{j+1} is a candidate of (R4) at S_j, by Lemma 2.1(ii) and (H1) at $j+1$. Lemma 6.13 gives $P_j\Gamma_j(q) \ge P_k > M \ge \mu(S_j)$ for every integer q with $3 \le q \le p_{j+1}$, so rule (i) applies at no prime $q \le p_{j+1}$. If $p_{j+1} \le q_c$, then $P_{j+1} \ge \mu(S_j)$, and $M > P_{j+1}$ forces $M \ge \mu'(S_j)$; hence $P_j\Gamma_j(q) > \mu'(S_j)$ for $q \le p_{j+1}$, rule (ii) applies at no prime $q \le p_{j+1}$, and an application at a larger prime does not reach p_{j+1}. If $p_{j+1} > q_c$, rule (ii) omits only primes up to q_c, and $V_{j+1}(\mu(S_j)) \ge V_{j+1}(M) \ge \theta_{j+2}$ by (H2) at $j+1$, so rule (iii) does not omit p_{j+1}. Hence S_{j+1} is generated.

At the node S_{k-1} we have $\mu(S_{k-1}) \le M < P_k$ and $P_k = P_{k-1}p_k/(p_k-1) \le P_{k-1}(p_{k-1}+2)/(p_{k-1}+1)$, so M is one of the values tested in (R3), and p_k is the quotient given by Lemma 6.1(ii). □

6.4. **The computation.** All quantities in the rules are rational numbers, and the search evaluates them exactly. The primes $\pi_i(q)$ below $2 \cdot 10^7$ are taken from a sieve. Primes above that are reached by stepping to the next probable prime; such a step never skips a prime, so a composite number may occasionally be treated as a prime, which can only enlarge the tree. It is equally harmless to replace q_1 in (R1) by the smaller number $p_j + 2$, as one of the two programs does beyond the sieve, because the proof of Proposition 6.14 uses only Lemma 6.13, which holds for every integer q with $3 \le q \le p_{j+1}$. In rule (R3) the quotient was never an integer, so no primality proof is needed.

Theorem 6.15. *The tree $\mathcal{T}_7$ has* 33678 *nodes and depth* 13. *Rule* (R3) *tests* 14814 *values of M, and for none of them is the quotient $(MB_j - 1)/(MB_j - A_j)$ an integer. Consequently, by Proposition 6.14, no composite solution of* (1.1) *is* 7*-heavy.*

TABLE 6. The tree $\mathcal{T}_7$ by depth j: the number of nodes and the number of values of M tested by rule (R3).

j	0	1	2	3	4	5	6	7	8	9	10	11	12	13
nodes	1	1	1	3	6	7	7	13	31	79	266	1 501	29 073	2 689
tests	0	0	0	0	0	0	0	0	0	0	16	215	11 894	2 689




FIGURE 13. The nodes of $\mathcal{T}_7$ by depth j. The bars span the range of $\log_2 \log_2 V_j(\mu)$ (right of each depth) and of $\log_2 \log_2 A_j$ (left of each depth) over the nodes of depth j, and the short marks below them are the thresholds $j+1-7$ of (H2) and $j-7$ of (H1). The deficit condition (H2) is the one that binds: at depth 13 the smallest value of $\log_2 \log_2 V_{13}$ is $7.0000075\ldots$

Table 6 lists the nodes by depth. Every node has $\mu(S) = 2$, so every node has $p_1 \neq 3$ and $P_j < 2$; in particular the sequence (3), for which $\mu = 4$, is discarded at once by the ratio test. The first branch of the tree is $5, 7, 13, 17, 19, 23, 37, 59, 67, 73$, the beginning of the independent set of Section 2.1 chosen one prime at a time, and the deepest nodes have thirteen entries.

Proof of Theorem 1.2(i). By Theorem 6.15 no composite solution of (1.1) is 7-heavy, and Corollary 6.10 with $s = 7$ gives $n < 2^{2^{k-7}}$. □

For $s = 1, \ldots, 7$ the trees $\mathcal{T}_s$ have 1, 3, 7, 12, 24, 118 and 33678 nodes. The deficit test does most of the work: without rules (R2) and (iii), the tree $\mathcal{T}_6$ already has 4469 nodes and depth 12.

6.5. **Larger quotients.** When $M \geq 3$, condition (H1) for a suitable s is incompatible with Lemma 6.1(i) and the bounds of Section 2.1, and no further search is needed. Let $\ell_1 < \ell_2 < \cdots$ be integers with $p_i \geq \ell_i$ for $i \leq k$, let $J \geq 1$, and let V^* be a number with $P_{\min\{k,J\}} < V^*$. If (H1) holds for all $j \leq k-1$, then $A_i \leq p_i^i$ gives $p_i \geq \rho_i$ for $i \leq k-1$, where ρ_i is the least positive

integer x with $x^i \ge \theta_i$. With $w_i = \max\{\ell_i, \rho_i\}$ we obtain, as in the proof of Lemma 6.13,

$$P_k < R_s = V^* \max\Big\{1, \quad \max_{J \le i < i_0} \frac{w_i' + 2}{w_i' + 1} \prod_{l=J+1}^{i} \frac{w_l}{w_l - 1}, \\ \Big(1 + \frac{16}{\rho_{i_0} - 2}\Big) \prod_{l=J+1}^{i_0 - 1} \frac{w_l}{w_l - 1}\Big\}. \tag{6.6}$$

where $w_i' = \max\{w_i, \ell_{i+1} - 2\}$ accounts for $p_{i+1} \ge \max\{\ell_{i+1}, w_i + 2\}$, and $i_0 \ge \max\{J+1, s+4\}$ is an index with $\rho_{i_0} > 10^{30}$. The three terms cover $k \le J$, then $J < k \le i_0$ with $k = i+1$, and $k > i_0$. The tail estimate is that of Lemma 6.13, applied to $y_i = \theta_i^{1/i}$, for which $p_i \ge \rho_i \ge y_i > \rho_i - 1$ when $\rho_i > 1$, and $\log y_{i+1} = \frac{2i}{i+1} \log y_i \ge \frac{4}{3} \log y_i$ when $i \ge \max\{s, 2\}$. We evaluate R_s in exact rational arithmetic, with integer lower bounds in place of the ρ_i, which can only enlarge it.

Proof of Theorem 1.2(ii) *and* (iii). Suppose $3 \mid n$. Then $M \ge 4$, every prime factor other than 3 is congruent to 2 modulo 3, by Lemma 2.2, and $k \ge 10^9 + 2$ by Corollary 2.6(ii). Take $J = 10^9 + 1$, $\ell_i = 2i + 1$, which is at most the i-th odd prime, and $V^* = 4(1 - 10^{-6})$. The J smallest prime factors are 3 and an independent set of 10^9 primes $p \equiv 2 \pmod 3$, so $P_J < \frac{3}{2} \cdot \frac{8}{3}(1 - 10^{-6}) = V^*$ by Proposition 2.5(ii). The computation gives $R_{10^9} = 3.9999960\ldots < 4$. If (H1) held for all $j \le k - 1$ with $s = 10^9$, then (6.6) would give $P_k < 4 \le M$, contrary to Lemma 6.1(i). So n is not 10^9-heavy, and Corollary 6.10 gives (iii).

Suppose $M \ge 3$ and $3 \nmid n$. Then $k \ge 18001$ by Corollary 2.6(i). Take $J = 18000$, $\ell_i = r_i$, the primes from 5 on, and $V^* = 2.999999$; the J smallest prime factors form an independent set, so $P_J < V^*$ by Proposition 2.5(i). Then $R_{17981} = 2.99999900511\ldots < 3$, and the same argument gives $n < 2^{2^{k-17981}}$. If $M \ge 3$ and $3 \mid n$, part (iii) gives a stronger bound. □

The exponent 17981 is the largest that this argument gives with $J = 18000$, since $R_{17982} = 3.000014\ldots$. Everything is decided by the prime factors around the 18000-th. For $s = 17981$, (H1) at $j = 18000$ gives $p_{18000} \ge \rho_{18000} > 5 \cdot 10^8$, so that the remaining prime factors raise $P_{18000} < 2.999999$ by a factor less than $1 + 2 \cdot 10^{-9}$. For $s = 17982$ the corresponding bound $\rho_{18000} < 2.5 \cdot 10^4$ lies below $r_{18000} = 200201$, and the next prime factor alone may raise the product by the factor $1 + 1/200226$.

6.6. **Where the search is hard.** The nodes that survive longest are prefixes with P_j just below 2, the same kind of prefix that makes the next cases expensive (Section 9). Every node at depth 13 has $2 - P_{13} < 10^{-10}$: there A_{13} has between 27 and 29 digits, while $C_{13} = 2B_{13} - A_{13}$ has between 14 and 18. In this respect the deep nodes behave like the products of Fermat primes of Remark 6.7, for which $2\varphi(n) - n = 1$. For such a prefix $1 - P_j/2$ is tiny, so V_j is large and the deficit test cannot discard it. Its children are few for the same reason: a prime $q \le q_c = \lfloor 1/(1 - P_j/2) \rfloor$, which exceeds $3 \cdot 10^{10}$ at depth 13, would raise P_{j+1} to 2 or more and force $M \ge 3$, and rule (ii) removes these primes. Figure 13 shows how close to the threshold of (H2) the deepest nodes come.

The tree grows quickly with s (Figure 14), and for $s = 7$ more than nine tenths of its nodes lie at depths 12 and 13. Each further unit of s halves the number of binary digits allowed by Theorem 1.2(i). The method proves $n < 2^{2^{k-s}}$ for every s for which the tree $\mathcal{T}_s$ can be searched, and the cost of the search is dominated by the deep prefixes with P_j close to 2. We have not completed the search for $s = 8$: at the node $5, 7, 13, 17, 19, 23, 37, 59, 67, 73, 317, 5483$ of depth 12, where $2/P_{12} - 1 = 7.0 \cdot 10^{-8}$, rule (R4) already generates more than 10^6 sequences.




FIGURE 14. The number of nodes of $\mathcal{T}_s$ at each depth, for $s = 4, 5, 6, 7$, on a logarithmic scale.

6.7. **Comparison with other bounds.** Stone [37, (52), (53)] treated tuples of primes with $\prod(1 - 1/p_i) \le \frac{1}{2} < \prod_{i<k}(1 - 1/p_i)$, which include the prime factors of solutions with $M = 2$. For $p_1 = 5$ he obtained $\prod p_i \le E_k(\beta)$ with $\beta \approx 1.66127$, and with $\beta \approx 1.50188$ when also $p_2 = 7$. Theorem 1.2(i) gives the base $2^{1/128} = 1.00543\ldots$ for solutions of (1.1). Table 7 compares, along the first branch of $\mathcal{T}_7$, the exponents e for which the three bases give $n < 2^{2^{k-e}}$. The deficit gains about three quarters of a unit of e over Lemma 6.5, and Theorem 6.2 adds a further gain over Stone's form that is largest for short prefixes.

TABLE 7. Along the branch $5, 7, 13, \ldots$ of $\mathcal{T}_7$ with $M = 2$: the exponent e such that the bound for n obtained at depth j is $2^{2^{k-e}}$, for the base $(A_j + 1)^2$ of Lemma 6.5, the base b^2/d of Stone and the base V_j of Theorem 6.2.

j	p_j	P_j	Cook–Nielsen	Stone	Theorem 6.2
1	5	1.25000	−0.370	−0.142	0.000
2	7	1.45833	−0.370	0.099	0.181
3	13	1.57986	−0.143	0.486	0.530
4	17	1.67860	0.309	1.017	1.041
5	19	1.77186	0.899	1.644	1.657
6	23	1.85240	1.561	2.324	2.330
7	37	1.90385	2.251	3.030	3.033
8	59	1.93668	2.965	3.759	3.761
9	67	1.96602	3.720	4.516	4.517
10	73	1.99333	4.507	5.265	5.265

7. The companion equation

7.1. **Up to seven prime factors.** Let n satisfy (1.2) with $\omega(n) \le 7$. If $n \le 3$ there is nothing to prove. Otherwise n is odd, squarefree and composite by Lemma 2.1, and $n+1 = 2\varphi(n)$ by Corollary 2.4. Since 3 may divide n, the search now starts from $p_{\min} = 3$, and we run it with $\varepsilon = +1$ for $2 \le k \le 7$. The outcome is in Table 8.

Table 8. The search for $n+1 = 2\varphi(n)$ with $3 \le p_1$ and $k \le 7$.

k	nodes	terminal	largest width for p_{k-1}	solutions found
2	1	1	1	15
3	3	2	2	255
4	5	2	15	65535
5	11	4	256	83623935, 4294967295
6	60	44	65 536	6992962672132095
7	6 914	6 826	2^{32}	none

For $k \le 6$ the search returns exactly the known solutions, each at its own depth, and this is also a test of the programs. For $k = 7$ it returns nothing. Here the enumeration route is too slow, because one terminal node, the prefix $3, 5, 17, 257, 65537$, has an interval for p_6 of width 2^{32}; for this prefix $C = 1$ and $D = 2AB + C = 2^{64} - 2^{32} + 1$, which is prime[11]. We therefore use the divisor route throughout, which is rigorous in this range: the largest integer D factored in the case $k \le 7$ has 21 digits, so every factorisation is certified. For $k \le 6$ the enumeration route was also run and agrees node for node. This proves Theorem 1.4 for solutions with at most seven prime factors; the case of eight is treated in Section 7.4.

7.2. **Solutions prime to three.** Let $n > 3$ satisfy (1.2) with $3 \nmid n$ and $\omega(n) \le 15$. Then $n = p_1 \cdots p_k$ with $5 \le p_1$, and $n+1 = 2\varphi(n)$ by Lemma 2.3. For $k \le 6$ the product $\prod p_i/(p_i - 1)$ is at most $1.9490\ldots$, whereas (3.2) requires it to be $2 - 1/\varphi(n)$. For $k \ge 2$ this is at least $2 - \frac{1}{24} > 1.95$, as $\varphi(n) \ge 4 \cdot 6$, and for $k = 1$ it is at least $\frac{7}{4}$, whereas $p_1/(p_1 - 1) \le \frac{5}{4}$. For $7 \le k \le 14$ we ran both implementations with $\varepsilon = +1$ and $p_{\min} = 5$. Neither found a solution, and the node counts, the numbers of terminal nodes and the largest widths coincide with those of Table 2 for every k. For $k = 15$ the nodes at depth 12 and their intervals are again those of Section 5.2, and the method of Section 4, run with $\varepsilon = +1$ by both implementations and by the route of Section 4.4 over the same 33865004 pairs, finds no completion in integers. This proves Theorem 1.5.

The coincidence with Table 2 has a simple cause. The two sets of bounds in (3.3) differ only by the terms $1/(A_j W)$, which in this range are far smaller than the gaps between consecutive admissible values of $p/(p-1)$ near the endpoints, and the computation confirms that no endpoint moves. The tests (3.4) at the last level do differ, but by then the tree is fixed. For $3 \nmid n$ and $k \le 15$ the two equations therefore lead to the same search.

7.3. **Extensions of Fermat-type solutions.**

Proof of Theorem 1.7. Let $2\varphi(n_0) = n_0 + 1$ and let $Q > 1$ be squarefree and coprime to n_0. Then $n = n_0 Q$ satisfies $2\varphi(n) = n + 1$ if and only if $2\varphi(n_0)\varphi(Q) = n_0 Q + 1$, that is, $(n_0 + 1)\varphi(Q) = n_0 Q + 1$, or

$$n_0(Q - \varphi(Q)) = \varphi(Q) - 1. \tag{7.1}$$

[11] Indeed $A = 2^{32} - 1$, $B = 2^{31}$ and $C = 2B - A = 1$, so $D = 2^{32}(2^{32} - 1) + 1 = \Phi_6(2^{32})$, the sixth cyclotomic polynomial at 2^{32}. Its primality is well known and is confirmed by a deterministic test, since $D < 3.3 \cdot 10^{24}$.

(i) For $Q = q$ prime, (7.1) reads $n_0 = q - 2$.

(ii) For $Q = pq$ with primes $p < q$ we have $Q - \varphi(Q) = p + q - 1$ and $\varphi(Q) - 1 = pq - p - q$, so (7.1) becomes $pq - (n_0 + 1)(p + q) + n_0 = 0$, and adding $(n_0 + 1)^2 - n_0$ to both sides gives

$$(p - n_0 - 1)(q - n_0 - 1) = n_0^2 + n_0 + 1.$$

The two factors on the left have the same sign, and they are positive: if both were negative we would have $2 \le p < q \le n_0$, and the left-hand side would be at most $(n_0 - 1)^2 < n_0^2 + n_0 + 1$. Conversely, a factorisation $n_0^2 + n_0 + 1 = de$ with $d < e$ gives $p = n_0 + 1 + d$ and $q = n_0 + 1 + e$, and if both are prime then n_0pq is a Fermat-type solution; the conditions $p, q \nmid n_0$ hold because $p, q > n_0$. The divisor $d = 1$ gives $p = n_0 + 2$ and $q = (n_0 + 1)^2 + 1 = n_0(n_0 + 2) + 2$, which is the composition of two one-prime extensions.

(iii) Factoring $n_0^2 + n_0 + 1$ and testing $n_0 + 2$ for each of the eight numbers gives Table 9. The one-prime extensions are $1 \to 3 \to 15 \to 255 \to 65535 \to 4294967295$ and $83623935 \to 6992962672132095$. The two-prime extensions are those listed in the table, and the only one that is not a composition of one-prime extensions is $255 \to 83623935$. Every extension of each of the eight numbers is therefore again one of the eight, and each of them is reached from 1, so the eight numbers are exactly the closure of $\{1\}$. The rows for 65535, 83623935, 4294967295 and 6992962672132095 show where the chains stop. Every factor in Table 9 is below 2^{64}, where the primality tests we use are proofs, and a test that declares a candidate p or q composite is never wrong. □

TABLE 9. One- and two-prime extensions of the Fermat-type solutions.

n_0	$n_0 + 2$ prime	factorisation of $n_0^2 + n_0 + 1$; two-prime extensions (p, q)
1	yes	3; $(3, 5)$
3	yes	13; $(5, 17)$
15	yes	241; $(17, 257)$
255	yes	$97 \cdot 673$; $(257, 65537)$ and $(353, 929)$
65535	yes	$193 \cdot 22253377$; none
83623935	yes	$67 \cdot 180097 \cdot 579535339$; none
4294967295	no	$2^{64} - 2^{32} + 1$, prime; none
6992962672132095	no	$13 \cdot 37 \cdot 109 \cdot 30802180789 \cdot 30280941915943441$; none

The chains stop for the following reasons. For $n_0 = 65535$ the trivial divisor gives $p = 65537$ and $q = 2^{32} + 1 = 641 \cdot 6700417$, which is Euler's factorisation of F_5, and the divisor 193 gives $p = 65729$, a prime, but $q = 22318913 = 3037 \cdot 7349$. For $n_0 = 4294967295$ the one-prime extension fails because $n_0 + 2 = 2^{32} + 1$ is composite once more. Figure 16 shows the hyperbolas for $n_0 = 255$ and $n_0 = 65535$ with their lattice points.

Remark 7.1. For three new primes, $Q = spq$ with $s < p < q$, write $x = s - 1$, $y = p - 1$, $z = q - 1$. Equation (7.1) becomes

$$xyz = n_0(xy + yz + zx) + n_0(x + y + z) + n_0 + 1,$$

and dividing by n_0xyz shows that $1/x + 1/y + 1/z$ falls short of $1/n_0$ by $(n_0(x + y + z) + n_0 + 1)/(n_0xyz)$, which is less than $4/x^2$ because $x < y < z$. Hence x, the smallest of the three, satisfies $3/x > 1/n_0 - 4/x^2$ and $1/x < 1/n_0$, that is, $n_0 < x < 3n_0 + 2$. For fixed x the equation is bilinear in y and z:

$$\big((x - n_0)y - n_0(x + 1)\big)\big((x - n_0)z - n_0(x + 1)\big) = (x - n_0)(n_0x + n_0 + 1) + n_0^2(x + 1)^2,$$




FIGURE 15. The Fermat-type solutions of $n + 1 = 2\varphi(n)$ as the closure of 1 under extensions. Horizontal arrows are one-prime extensions $n_0 \mapsto n_0(n_0 + 2)$, labelled by the prime $n_0 + 2$. The bent arrow is the two-prime extension of 255 by $(353, 929)$, coming from the factorisation $65281 = 97 \cdot 673$. Dashed arrows end the two chains: 4294967297 and 6992962672132097 are composite, and neither $n_0^2 + n_0 + 1$ has a factorisation that yields two primes.




FIGURE 16. The two-prime extensions of n_0 are the lattice points $(d, e) = (p - n_0 - 1, q - n_0 - 1)$ on $de = n_0^2 + n_0 + 1$ with both p and q prime, shown on logarithmic axes. For $n_0 = 255$ both lattice points with $d \le e$ give primes (filled). For $n_0 = 65535$ both lattice points give a prime p and a composite q (open).

so the pair (y, z) comes from the divisors of an integer of size about $n_0^2 x^2$. For $n_0 = 4294967295$ there are about $3.8 \cdot 10^8$ primes s in the range, and $1.3 \cdot 10^8$ of them pass the congruence prune: $s \equiv 2 \pmod 3$ as Lemma 2.2 requires, and $s \not\equiv 1$ modulo 5, 17, 257 and 65537. Each of them leaves an integer of about forty digits to factor. This is the prefix $3, 5, 17, 257, 65537$ that dominates the case $k = 8$ in Section 7.4.

7.4. **Eight prime factors.** Let n satisfy (1.2) with $\omega(n) = 8$. Then $n > 3$, and n is odd and squarefree by Lemma 2.1. If $3 \nmid n$, then $\omega(n) \ge 16$ by Theorem 1.5; hence $3 \mid n$. By Lemma 2.2 the quotient M is either 2 or at least 5, and in the second case $\omega(n) \ge 1540$. Hence $n+1 = 2\varphi(n)$

and $p_1 = 3$, and it suffices to run the search with $\varepsilon = +1$ and $k = 8$ over the prefixes that begin with 3.

The traversal down to depth 5 takes a few seconds. It yields 10459 prefixes $(p_1, \dots, p_5)$ with a non-empty interval for p_6, and 10458 of them begin with 3. Their intervals have total length $1.3045 \cdot 10^{10}$, and most of it belongs to a few prefixes (Figure 18, left): $8.59 \cdot 10^9$ to $3, 5, 17, 257, 65537$, that is, to the solution 4294967295 itself, where $C_5 = 1$; $1.23 \cdot 10^9$ to $3, 5, 17, 257, 65543$; and $1.67 \cdot 10^8$ to $3, 5, 17, 353, 929$, the solution 83623935, where again $C_5 = 1$. Thirty-four prefixes have an interval longer than 10^7, and 326 one longer than 10^6. The intervals contain 228260561 primes s that pass the congruence prune, 131788942 of them for the prefix $3, 5, 17, 257, 65537$. Each pair of a prefix and such a prime $s = p_6$ leaves the two-prime problem of Proposition 4.1, in which N has between 15 and 40 digits.

These problems lie far to the left of those of Figure 9. For the prefix $3, 5, 17, 257, 65537$ and $p_6 = s$ one has $A' = (2^{32} - 1)s$ and $B' = 2^{31}(s - 1)$, so that the modulus is $c = s - 2^{32}$, and $N = 2A'B' + c$ is close to $2^{64}s^2$. As s runs over $[2^{32} + 1, 3 \cdot 2^{32} - 1]$, the ratio c^3/N stays below $2.1 \cdot 10^{-10}$, and the number $(N/c^3)^{1/2}$ that governs the methods of Section 4 grows from $7 \cdot 10^4$ at the upper end to $4 \cdot 10^{16}$ at the first admissible prime, $s = 4294967357$, where $c = 61$. We therefore combine the sum route with trial division and with a complete factorisation, as follows.

Fix a prefix and an admissible prime s, define A', B', c, N as in Section 4, and let $t = cp - 2B'$ be the divisor of Proposition 4.1. Since $p \geq s + 2$, it satisfies $t \geq t_0 = c(s + 2) - 2B'$, and we choose a split point t_d with $t_0 \leq t_d$.

(i) The divisors with $t < t_d$ are found by running p over the integers with $t_0 \leq cp - 2B' < t_d$ and testing whether t divides $A'p + \varepsilon$, as in Proposition 3.5. Since $c(A'p + \varepsilon) = A't + N$ and $\gcd(t, c) = 1$ by Proposition 4.1, this is the same as testing whether t divides N, and the program does that when $A'p$ is too large for 128-bit arithmetic.

(ii) For the divisors with $t_d \leq t \leq \sqrt{N}$ we use the sum $\sigma = p + q$ and the product $\pi = pq$. By (3.5),

$$c\,\pi = 2B'\sigma - 2B' + \varepsilon, \qquad (q - p)^2 = \sigma^2 - 4\pi, \tag{7.2}$$

so σ lies in the class of $(2B' - \varepsilon)(2B')^{-1}$ modulo c, and $t + N/t = c\sigma - 4B'$. As $x + N/x$ decreases on $(0, \sqrt{N}\,]$, the sum runs over its class in the range $2\sqrt{N} + 4B' \leq c\sigma \leq t_d + N/t_d + 4B'$, and each value is tested by asking whether $\sigma^2 - 4\pi$ is a perfect square. This is the route of Section 4.4, written in terms of p and q instead of u and v: the two sums differ by the constant $(2r + 4B')/c$.

(iii) Both loops are sieved by congruences. Let $p < q$ be integers with $A'pq + 1 = 2B'(p-1)(q-1)$. No p_i divides $p - 1$ or $q - 1$, since it would then divide 1. If moreover p and q have no prime factor up to 61, which holds for primes larger than s because every admissible s exceeds 61, then modulo every prime $\ell \leq 61$ they avoid the residue 0, and also the residue 1 when ℓ is one of the p_i; in particular they are odd and congruent to 2 modulo 3. Call these residues excluded. Since $X^2 - \sigma X + \pi = (X - p)(X - q)$, the roots of this polynomial modulo a prime ℓ are p and q modulo ℓ. The sieve discards only values of σ that are odd, or not congruent to 1 modulo 3, or for which $\sigma^2 - 4\pi$ is not a square modulo one of 2^8, 3^2, 5^2 and 7^2, or for which, for some prime $\ell \leq 43$, the polynomial has no root modulo ℓ or a root at an excluded residue; and in the trial division it discards only values of p at an excluded residue modulo some prime $\ell \leq 61$. Every other value is tested exactly. So the sieve keeps every value that comes from integers p and q without prime factors up to 61, and in particular every value that comes from primes.

(iv) If the estimated work of (i) and (ii) exceeds four milliseconds, the time we allow for factoring N, then N is factored completely with PARI/GP [31], every prime factor is

proved prime,[12] and all divisors t of N with $t_0 \le t \le \sqrt{N}$ in the class of Proposition 4.1 are listed.

Since c is odd and prime to 3, the admissible values of σ form one class modulo $6c$. The work of (i) and (ii) is therefore about $(t_d - t_0)/c$ values of p and $(t_d + N/t_d - 2\sqrt{N})/(6c^2)$ values of σ, and t_d is chosen to minimise it (Figure 17), with the two kinds of steps weighted by their measured costs. With equal weights the minimum lies near $t_d = (N/6c)^{1/2}$ and amounts to about $2(N/6c^3)^{1/2}$ steps, as in Section 4.4. By the argument of that section, (i) and (ii) together find every divisor $t \ge t_0$ of N in the class up to $\sqrt{N}$, and so does (iv); the sieve (iii) loses none that comes from integers p and q without prime factors up to 61. Where σ and π are too large for 128-bit arithmetic, (ii) is carried out in multiprecision arithmetic. Now let n be a solution with $p_1 = 3$ and $k = 8$. By Theorem 3.6 its first five primes form one of the prefixes and p_6 is one of the admissible primes s of its interval, and by Proposition 4.1 its last two primes give a divisor $t \ge t_0$ in the class. Hence n is found. Every completion $s < p < q$ in integers that is found is checked against $A'pq + 1 = 2B'(p-1)(q-1)$ exactly, and p and q are tested for primality.

The computation examined all 228260561 pairs; in 1967265 of them N was factored, with up to 39 digits. The computation was cut into 10897 pieces, each a range of s under one prefix, and run on four processors; it made $3.4 \cdot 10^{11}$ trial divisions, the sums ranged over $6.7 \cdot 10^{13}$ values of σ, the sieve (iii) passed $2.1 \cdot 10^8$ of them on to the exact test, and the whole run took 6.9 hours of processor time, 3.2 of them in PARI/GP. It found 21 completions $s < p < q$ in integers and no solution. Sixteen of the completions lie under the prefix $3, 5, 17, 353, 929$ with $s = 83623937$, where $3 \cdot 5 \cdot 17 \cdot 353 \cdot 929 \cdot s$ is the solution $n_0 = 6992962672132095$. They are the pairs $p = n_0 + 1 + d$, $q = n_0 + 1 + e$ of Theorem 1.7(ii), one for each factorisation $n_0^2 + n_0 + 1 = de$ with $d < e$, and by Table 9 none of them consists of two primes. In each of the other five, listed in Table 10, p or q is divisible by one of $p_1, \dots, p_5$. So (1.2) has no solution with eight prime factors, and together with Section 7.1 this proves Theorem 1.4.

TABLE 10. The completions $s < p < q$ in integers of the case $k = 8$ other than the sixteen two-prime extensions of 6992962672132095, with the factorisations of p and q.

$s = p_6$	p	q
prefix $3, 5, 17, 257, 65537$		
4294967459	$17 \cdot 43 \cdot 673 \cdot 230521189693$	$5 \cdot 409642859 \cdot 26363459459$
4294967969	27419312914203719	$5 \cdot 19 \cdot 5011 \cdot 7699 \cdot 21393991009$
4596195557	65533391633	$17 \cdot 2092323256997814577105 2877$
prefix $3, 5, 17, 353, 929$		
83625827	$3^2 \cdot 5^3 \cdot 13 \cdot 252862031$	$3 \cdot 5 \cdot 7 \cdot 13 \cdot 113 \cdot 794656817253868193$
83760029	$3 \cdot 5 \cdot 3431155289$	$3 \cdot 87389293 \cdot 45294677621$

8. Checks on the computations

A search that finds nothing proves something only if it would have found what it looks for. Every search of this paper was therefore carried out by at least two programs written

[12] With the option `factor_proven` set, PARI/GP certifies every prime factor it returns. A factor below 2^{64} is certified by its Baillie–PSW test, which no composite number below 2^{64} passes; the PARI/GP documentation records that this was checked against the table of base-2 pseudoprimes below 2^{64} of Feitsma and Galway [12]. A larger factor is certified by the $n - 1$ method [5] when enough of $n - 1$ is known, and otherwise by the test of Adleman, Pomerance and Rumely [1] in the form of Cohen and Lenstra [9].




FIGURE 17. Steps (i), (ii) and (iv) for four admissible values $s_1 < s_2 < s_3 < s_4$ of p_6, not to scale. The divisors $t = cp - 2B'$ of N with $t_0 \le t \le \sqrt{N}$ lie on the branch $t \mapsto N/t$. Trial division covers $t < t_d$ and the sums cover $t_d \le t \le \sqrt{N}$. The split point t_d is chosen for each s to minimise the work, and where both routes would be slow, as for s_1, the integer N is factored completely.




FIGURE 18. The case $k = 8$ of the companion equation. Left: the 10458 prefixes $(3, p_2, \ldots, p_5)$ by the width of the interval for p_6, on logarithmic axes; the two in the last bar end in 65537 and 65543. Right: processor time per admissible p_6 under the prefix $3, 5, 17, 257, 65537$, one point for each piece of the run, against the modulus $c = p_6 - 2^{32}$. For c below about 10^8 the integer N is factored, and each p_6 costs about six milliseconds in all; above that the sums and the trial division are faster, and their cost falls as c grows. Beyond $c = 2^{32}$, that is for $p_6 > 2^{33}$, the condition $p > p_6$ leaves only a short range of divisors.

independently, and every count reported here is the same for all of them, with the two exceptions stated in Sections 2.1 and 5.3: the two programs of Proposition 2.5(i) differ in the sets they create

near the margin, and the search below 10^{35} of Theorem 1.3 was run by one program, after the two programs had agreed below 10^{31}. For Theorems 1.1 and 1.5 one of the two programs never factors an integer, so that its negative answers depend on no primality proof. The programs were also run on equations that do have solutions: on the equation $2^k(n-1) = (2^k+h)\varphi(n)$, whose 56 solutions with $k = 4$ and $k = 5$ in the ranges of Appendix A they all return, and on (1.2), where they return every known solution with at most six prime factors at the right depth and nothing else. The divisor routines of Section 4 were compared with complete factorisations and tested on planted divisors, and the program of Section 7.4 was tested in the same ways and partly run twice. The programs, their recorded output and the details of these tests are in the archive [4].

9. The assertion that remains open

The methods of Section 4 settle the case $k = 15$ of Lehmer's equation in a quarter of an hour on a single processor and the case of eight primes of the companion equation in 6.9 hours. They do not reach the next cases: for (1.1) with sixteen prime factors the number of two-prime problems grows from $3.4 \cdot 10^7$ to about $9 \cdot 10^{13}$, and for (1.2) with nine prime factors and $3 \mid n$ the Fermat primes alone leave some 10^{17} of them. In both cases most of the cost comes from a few prefixes whose product $\prod p/(p-1)$ differs from 2 by less than 10^{-7}, the prefixes that Section 6.6 describes. The search with the size rule of Section 5.3 is cheaper by far, but its cost grows by a factor between 5 and 15 for each further power of 10 in the bound, and it cannot reach the bound 2^{512} of Theorem 1.2. We end with the statement, equivalent to Lehmer's conjecture, that a proof for all k would have to establish.

Proposition 9.1. *Equation (1.1) has no composite solution if and only if there are no finite set S of odd primes and no integer $M \geq 2$ such that, with $A = \prod_{p\in S} p$ and $B = \prod_{p\in S}(p-1)$, the number $D = MB - A$ is positive, divides $A-1$, and makes $(MB-1)/D$ a prime larger than every element of S.*

Proof. Let n be a composite solution of (1.1), let q be its largest prime factor, S the set of its other prime factors and $M = (n-1)/\varphi(n)$; by Lemma 2.1 the elements of S are odd. Then $n = Aq$, $\varphi(n) = B(q-1)$, and $n - 1 = M\varphi(n)$ reads $qD = MB - 1$; so $D \geq 1$, D divides $MB-1 = D+(A-1)$ and hence $A-1$, and $q = (MB-1)/D$ is a prime larger than every element of S. Conversely, for S and M as in the statement the number $n = Aq$ with $q = (MB-1)/D$ is composite and satisfies $n - 1 = M\varphi(n)$, since $qD = MB - 1$ is the same equation. □

By Theorem 1.1 a set S with these properties has at least 15 elements, at least 18000 if $M \geq 3$, and at least 10^9 if $3 \in S$. The assertion asks that one number attached to each set S never be a prime above the elements of S. For the companion equation the same construction with $D = 1$ gives the Fermat numbers: the Fermat primes $S = \{3, 5, 17, 257, 65537\}$ give $2^{32} + 1$. For Lehmer's equation $D = 1$ would give $(MB-1)/D = A$, which is not a prime larger than the elements of S; so $D > 1$, and the assertion is not a question about Fermat numbers. Two further constraints on D follow from the proof. Since $qD = D + A - 1$, the divisor is $D = (A-1)/(q-1)$, so $q - 1$ divides $A - 1$ as well and the assertion asks that $1 + (A-1)/(MB-A)$ never be a prime larger than the elements of S; and since $2^{|S|}$ divides B while A is odd, D is odd and $D \equiv -A \pmod{2^{|S|}}$, so $D \geq 3$.

9.1. **One candidate for each prefix.** The quotient in Proposition 9.1 need not be searched independently. The following elementary observation applies to both signs and to every number of prime factors.

Proposition 9.2. *Let S be a nonempty finite set of distinct odd primes, and put*

$$A = \prod_{p \in S} p, \quad B = \prod_{p \in S} (p-1), \quad r = \max S.$$

Suppose that $q > r$ is an odd integer, $\varepsilon \in \{-1, 1\}$, M is an integer, and

$$Aq + \varepsilon = MB(q-1).$$

Then

$$M = \left\lceil \frac{A}{B} \right\rceil, \qquad 0 < M - \frac{A}{B} \leq \frac{1}{2}.$$

The last inequality is strict when $\varepsilon = -1$.

Proof. Since A is odd and B is even, A/B is not an integer. Every factor $u/(u-1)$ with $u \geq 3$ exceeds 1, and S is a subset of the integers from 3 to r. Therefore

$$\frac{A}{B} \leq \prod_{u=3}^{r} \frac{u}{u-1} = \frac{r}{2}.$$

Rearranging the assumed equation gives the exact identity

$$M - \frac{A}{B} = \frac{A + \varepsilon}{B(q-1)} > 0.$$

Both q and r are odd, so $q - 1 \geq r + 1$. Also $B \geq 2$. For $\varepsilon = 1$ these facts give

$$\frac{A+1}{B(q-1)} \leq \frac{A/B + 1/B}{r+1} \leq \frac{r/2 + 1/2}{r+1} = \frac{1}{2}.$$

For $\varepsilon = -1$ we have instead

$$\frac{A-1}{B(q-1)} < \frac{A/B}{r+1} \leq \frac{r}{2(r+1)} < \frac{1}{2}.$$

Thus M is an integer strictly above A/B and less than $A/B + 1$, which proves the ceiling formula. □

Corollary 9.3. *For S, A, B, r as in Proposition 9.2, define*

$$m = \left\lceil \frac{A}{B} \right\rceil, \qquad d = mB - A.$$

Then $0 < d < B$. For each $\varepsilon \in \{-1, 1\}$ there is a prime $q > r$ for which $\varphi(Aq) \mid Aq + \varepsilon$ if and only if

$$d \mid A + \varepsilon \quad \text{and} \quad Q = 1 + \frac{A + \varepsilon}{d} \text{ is a prime greater than } r.$$

If it exists, the prime is unique and equals Q, and $Aq + \varepsilon = m\varphi(Aq)$.

Proof. The inequality $0 < d < B$ follows from $m - 1 < A/B < m$. If q is a completion, it is odd, and $\varphi(Aq) = B(q-1)$, since q is a prime outside S. Proposition 9.2 forces its quotient to be m. Rearranging $Aq + \varepsilon = mB(q-1)$ now gives $d(q-1) = A + \varepsilon$, so $q = Q$ and $d \mid A + \varepsilon$. Conversely, for the prime Q in the statement,

$$mB(Q-1) = (A+d)(Q-1) = AQ - A + d(Q-1) = AQ + \varepsilon,$$

which is the required equation, and the same identity proves uniqueness. □

Corollary 9.4. *Under the hypotheses of Corollary 9.3, suppose that $\varepsilon = -1$ and Q is a prime greater than r. Then d is odd,*

$$3 \le d < \frac{B}{2}, \qquad \gcd(d, mAB) = 1, \qquad A \equiv mB \equiv 1 \pmod{d}.$$

If a prime ℓ divides d, then $\ell \nmid m$ and every $p \in S$ is neither 0 nor 1 modulo ℓ. If $3 \mid d$, then $m \equiv 1 \pmod 3$, every element of S is 2 modulo 3, and $|S|$ is even. In particular $m = 2$ forces $d \ge 5$.

Proof. The integer $d = mB - A$ is odd, and Proposition 9.2 gives $d/B < 1/2$. If $d = 1$, then $Q = A$; this cannot be a prime greater than every element of the nonempty set S. Thus $d \ge 3$. Since $d \mid A - 1$, we have $A \equiv 1 \pmod d$; the identity $mB = A + d$ gives $mB \equiv 1 \pmod d$. Hence d is prime to A, m and B, proving the coprimality assertion and the stated restrictions modulo each prime factor ℓ of d. If $3 \mid d$, no element of S is 0 or 1 modulo 3, so each is 2 modulo 3. Thus $B \equiv 1 \pmod 3$, $mB \equiv 1$ gives $m \equiv 1 \pmod 3$, and $A \equiv 1$ forces $|S|$ even. For $m = 2$ this excludes $3 \mid d$, so the positive odd integer $d \ge 3$ is at least 5. □

Corollary 9.3 removes the unbounded search over M for a fixed prefix. More explicitly, in Proposition 9.1 one may require S to be nonempty and replace its existential integer M by $\lceil A/B \rceil$. It does not remove the unbounded quantifier over S. An exclusion of the candidate Q for every such set S, by showing that the divisibility fails or that Q is not a prime above r, would prove Lehmer's conjecture. No such exclusion is proved here. The elementary reductions in this subsection are not claimed to improve the numerical bounds in Theorem 1.1. Molnar and Singh [24] give an algorithm enumerating spoof Lehmer factorizations for each fixed number of positive bases. Their fixed-length enumeration and the reduction here leave distinct unbounded assertions: neither supplies a bound on the number of prime factors of a hypothetical Lehmer number.

The need to keep the primality test is visible in the companion equation. For $S = \{3, 5, 17, 257, 65537\}$ we have $A = 2^{32} - 1$, $B = 2^{31}$, $m = 2$ and $d = 1$. The candidate is

$$Q = 2^{32} + 1 = 641 \cdot 6700417.$$

It satisfies $AQ + 1 = 2B(Q - 1)$ exactly, but it is composite. Thus an exact integer completion and the ceiling identity do not by themselves produce a prime solution.

Appendix A. Solutions of the test equation

The following 56 integers $n = p_1 \cdots p_k$ satisfy $2^k(n-1) = (2^k + h)\varphi(n)$ for the indicated h. They are the complete output of both implementations for $k = 4$, $1 \le h \le 40$, and $k = 5$, $1 \le h \le 85$, with $p_1 \ge 3$, listed in increasing order of n down the columns. Each was verified by direct arithmetic.

Appendix B. Reproduction checks for this revision

On 24 October 2026 the sum-of-factors implementation `lastthree_c.py` was rerun from the independently recomputed frontier for $k = 15$, $\varepsilon = -1$, using the source at commit `fd398ef410faeca431c3c5bab142f3443e857cdf` of `creelie/eLhmr`. All 55443 tasks completed, covering 33865004 admissible values of p_{13}. There were no integer candidate completions and no prime solutions. This is a fresh reproduction of the fifteen-prime computation, rather than a check only of previously saved output.

The same implementation was rerun for $7 \le k \le 14$ for both signs, agreed with the recorded counts and recovered all 61 known positive-control completions. Its divisor routine was compared with independent factor enumeration on 3100 random or planted instances, with no mismatch;

TABLE 11. Solutions with $k = 4$, $1 \le h \le 40$.

$p_1, \ldots, p_4$	h	$p_1, \ldots, p_4$	h
3, 5, 47, 89	15	13, 29, 433, 2969	2
5, 19, 29, 163	6	7, 97, 139, 8581	3
7, 17, 229, 251	4	13, 29, 383, 28879	2
3, 29, 263, 521	9	11, 47, 1109, 15497	2
13, 37, 151, 271	2	29, 41, 1291, 15199	1
3, 29, 197, 1559	9	11, 47, 1049, 77477	2
17, 23, 83, 1709	2	7, 61, 859, 183397	3
29, 71, 163, 193	1	31, 37, 2377, 66499	1
41, 43, 97, 491	1	19, 157, 9697, 15511	1
37, 41, 137, 829	1	19, 157, 8353, 20887	1

TABLE 12. Solutions with $k = 5$, $1 \le h \le 85$.

$p_1, \ldots, p_5$	h	$p_1, \ldots, p_5$	h
5, 7, 13, 139, 653	19	5, 47, 317, 40253, 499027	9
11, 13, 17, 109, 379	9	19, 269, 443, 1747, 1852699	2
3, 17, 29, 317, 4643	21	23, 71, 983, 1031, 103437589	2
11, 17, 107, 173, 1373	6	19, 199, 757, 5059, 23698783	2
5, 19, 37, 73, 19739	12	31, 43, 233, 318629, 12310321	2
19, 73, 97, 103, 613	3	19, 313, 389, 1277, 590838019	2
11, 17, 127, 131, 2879	6	11, 47, 1051, 67967, 102302009	4
5, 7, 269, 317, 4283	15	7, 59, 1721, 41813, 958695209	6
5, 53, 233, 857, 5689	9	41, 167, 14411, 322633, 1891541	1
31, 53, 149, 569, 3407	2	31, 37, 2411, 47681, 5732949913	2
13, 17, 271, 1567, 5573	5	31, 37, 2297, 2639701, 1379629439	2
7, 13, 17, 1553, 480499	11	37, 463, 827, 1667389, 4325665973	1
17, 23, 131, 199, 167099	4	37, 353, 1867, 25109743, 844718249	1
11, 83, 163, 239, 210461	4	43, 139, 61729, 212167, 32068788511	1
13, 167, 349, 1747, 8237	3	37, 367, 1553, 21098893, 41563512479	1
19, 181, 1439, 3767, 17683	2	43, 139, 47857, 57253657, 72147837793	1
3, 11, 269, 17837, 3861929	21	37, 353, 1867, 24384917, 20504256743083	1
43, 317, 331, 953, 166357	1	43, 139, 47819, 1145259251, 653813324977	1

1610 of the instances had a planted divisor. The exact threshold products, extension calculations, fifteen stored $k = 14$ factorisation certificates and the hand-bound calculations were also rerun. Comparison of the six archived $k = 15$ runs checked agreement of the task sets and their results; that comparison is distinct from the fresh negative-sign run. The positive-sign $k = 15$ sum-route search was also freshly completed: its independently regenerated frontier gave the same 55443 tasks and 33865004 admissible values, with no integer candidates or prime solutions.

The 41 distinct factors stored for the fifteen long $k = 14$ nodes were also proved prime by complete Lucas certificates, with 259 recursive prime nodes. The independent verifier uses only integer multiplication, modular powers and greatest common divisors, and checks that its roots are exactly the stored factors and that those factorisations multiply to the recomputed integers. FLINT is used to discover the certificates, not in their verification. The criterion is elementary: if the prime factorisation of $v - 1$ is known and

$$a^{v-1} \equiv 1 \pmod{v}, \qquad \gcd\bigl(a^{(v-1)/\ell} - 1, v\bigr) = 1 \quad \text{for every prime } \ell \mid v - 1,$$

then the order of a modulo any prime divisor r of v divides $v-1$ and cannot divide $(v-1)/\ell$ for any such ℓ. Its valuation at each ℓ is therefore the full valuation of $v-1$, so the order equals $v-1$. Since it also divides $r-1$, we have $r \geq v$ and hence $r = v$. Thus v is prime. The certificates apply this argument recursively down to 2.

For the new reduction in Section 9.1, an independent integer implementation compared the one-candidate test with direct evaluation at 537786 pairs of a prime prefix and a prime tail, covering 6475 prefixes and both signs. It also recovered six known composite companion solutions and rejected the composite tail $2^{32}+1$ above. These finite checks supplement the proofs of Proposition 9.2 and its corollaries; they do not prove an assertion for all prefixes.

The independent-set computations of Proposition 2.5 were rerun for this revision at the larger sizes $K = 18000$ and $K = 10^9$, each by two programs (Section 2.1), and the search below 10^{35} of Theorem 1.3 was carried out (Section 5.3); their programs and output are supplied with this revision. The eight-prime companion computation was not freshly rerun; its records and implementation remain in the cited archive. These statements describe the 24 October reproduction. Subsequently, the version 1.8.0 archive recorded a local Lean 4.34.1 check of ten statements in `lean/UniformFinite.lean`, including the fixed-count reduction and final-entry identities. That limited check does not formalise the exhaustive searches or the universal exclusion. The present revision does not claim a new complete Lean build.

Appendix C. A uniform finite construction for every factor count

This appendix addresses the quantifier over the number of prime factors. It gives one terminating construction for each prescribed integer k, with a proof valid for every k. It does not prove that the output is empty for every k. Finite algorithms at a prescribed factor count are already part of the literature, including [24]; no priority claim for that general principle is made here. The bounds below and the supplied implementation are written out to make this particular construction self-contained. They use no unproved input from a research preprint.

Fix $\varepsilon \in \{-1, 1\}$ and consider composite solutions of

$$n + \varepsilon = M\varphi(n), \qquad M \in \mathbb{Z}_{\geq 2}. \tag{C.1}$$

Such a solution is odd and squarefree: evenness contradicts the parity of the equation, and a squared prime divisor of n would divide both $\varphi(n)$ and $n+\varepsilon$. Also $\gcd(n, M\varphi(n)) = 1$. Write the prime factors in increasing order. For a proper prefix put

$$A = \prod_{i=1}^{j} p_i, \qquad B = \prod_{i=1}^{j} (p_i - 1), \qquad t = k - j, \qquad D = MB - A.$$

The empty prefix has $A = B = 1$ and $t = k$. If c is the product of the remaining primes, then

$$D\varphi(c) = A(c - \varphi(c)) + \varepsilon > 0. \tag{C.2}$$

For a nonempty prefix, $A \geq 3$ and $c - \varphi(c) \geq 1$; for the empty prefix, $D = M - 1 \geq 1$. Thus every proper prefix of a solution has $D > 0$.

Lemma C.1 (Uniform bound for the next prime)**.** *Let p be the next prime of a solution extending the prefix, with t primes remaining. Then*

$$p \leq 1 + \left\lfloor \frac{tA - 1}{D} \right\rfloor \qquad (\varepsilon = -1), \tag{C.3}$$

$$p \leq 1 + \left\lfloor \frac{tA + 1}{D} \right\rfloor \qquad (\varepsilon = 1). \tag{C.4}$$

For $\varepsilon = -1$ one also has

$$p \le 2 + \left\lfloor \frac{2tA^2 - 1}{D(MB + A)} \right\rfloor. \tag{C.5}$$

Proof. Let $r_1 = p < r_2 < \cdots < r_t$ be the remaining primes. The function $x/(x-1)$ decreases for $x > 1$. Since $r_i \ge p + i - 1$,

$$\prod_{i=1}^{t} \frac{r_i}{r_i - 1} \le \prod_{i=1}^{t} \frac{p + i - 1}{p + i - 2} = 1 + \frac{t}{p - 1}.$$

For the minus sign, $MB < A \prod_i r_i/(r_i - 1)$, so $D(p-1) < tA$. All quantities are integers, giving (C.3). For the plus sign,

$$MB = A \prod_i \frac{r_i}{r_i - 1} + \frac{1}{\prod_i (r_i - 1)} \le A + \frac{tA}{p - 1} + \frac{1}{(p-1)^t}.$$

It follows that $D(p-1) \le tA + (p-1)^{1-t} \le tA + 1$, which gives (C.4).

For the sharper minus-sign estimate, oddness gives $r_i \ge p + 2(i-1)$. For $x > 2$,

$$\left(\frac{x}{x-1}\right)^2 < \frac{x}{x-2},$$

because $x(x-2) < (x-1)^2$. Multiplying the resulting inequalities yields

$$\left(\prod_i \frac{r_i}{r_i - 1}\right)^2 < \prod_{i=1}^{t} \frac{p + 2i - 2}{p + 2i - 4} = \frac{p + 2t - 2}{p - 2}.$$

Consequently $(MB)^2(p-2) < A^2(p + 2t - 2)$, or $D(MB + A)(p-2) < 2tA^2$. The integer form of this strict inequality is (C.5). □

Corollary C.2 (Finite quotient range)**.** *For $k \ge 2$, the quotient of a composite solution satisfies*

$$2 \le M \le \begin{cases} \min\{\lfloor \sqrt{2k} \rfloor, \lfloor (k+1)/2 \rfloor\}, & \varepsilon = -1, \\ \lfloor (k+3)/2 \rfloor, & \varepsilon = 1. \end{cases}$$

Proof. At the empty prefix $D = M - 1$ and the first prime is at least 3. The linear inequalities give the displayed linear bounds. For the minus sign, the squared estimate in the proof of Lemma C.1 gives $M^2 < 2k + 1$. Since M^2 is an integer, $M^2 \le 2k$. □

Theorem C.3 (Complete finite tree for each k)**.** *For every $k \ge 2$ and $\varepsilon \in \{-1, 1\}$, the following finite construction outputs exactly the composite integers with k distinct prime factors satisfying (C.1), each once.*

(i) *For each integer M in Corollary C.2, start with $(A, B, t) = (1, 1, k)$ and no chosen primes.*

(ii) *At a node with $t \ge 2$, let r be the last chosen prime, or 1 at the empty prefix. Enumerate the primes p such that*

$$p > r, \qquad p > \frac{MB}{D},$$

and p is at most the bound of Lemma C.1 (the minimum of its two minus-sign bounds when applicable). Keep only choices with $\gcd(A, p-1) = 1$ and $p \nmid M$. Replace the state by

$$(A, B, t) \longmapsto (Ap, B(p-1), t-1).$$

(iii) *At $t = 1$, calculate the single rational number*

$$q = \frac{MB + \varepsilon}{D} = 1 + \frac{A + \varepsilon}{D}. \tag{C.6}$$

Output Aq exactly when q is an integer prime greater than r.

One may also discard any node containing 3 unless $M \equiv -\varepsilon \pmod 3$.

Proof. Every node has $D > 0$ and therefore a finite upper bound for its next prime. The condition $p > MB/D$ makes the next proper-prefix defect $pD - MB$ positive. The depth decreases by one, so induction on t proves termination. Primality itself can be decided by trial division through the integer square root; no probable-prime assumption is needed.

Every genuine solution has a quotient in the stated finite range and a unique increasing list of primes. Its proper prefixes satisfy (C.2) and Lemma C.1. The gcd tests follow from $\gcd(n, M\varphi(n)) = 1$, so they cannot remove that list. Its last prime is forced by (C.6). This proves completeness.

Conversely, an accepted tail is a prime different from all prefix primes. Thus $\varphi(Aq) = B(q-1)$, and the identity $Dq = MB + \varepsilon$, with $D = MB - A$, gives $Aq + \varepsilon = MB(q-1)$. Every output therefore satisfies the target equation. Its increasing factorization and its quotient are unique.

Finally, if $3 \mid n$, no other prime factor can be 1 modulo 3, by the same gcd condition. Hence $\varphi(n) \equiv 2 \pmod 3$ and $M\varphi(n) \equiv \varepsilon \pmod 3$, giving $M \equiv -\varepsilon \pmod 3$. This proves the optional prune. □

Implementation and exact scope. The implementation `uniform_tree.py` uses integer arithmetic and trial division, and provides resource limits separately from the mathematical construction. Reaching a resource limit returns an explicit incomplete result, never an empty-search certificate. Fresh runs finish the minus-sign counts $2 \le k \le 13$, all empty, and reproduce the six known composite companion solutions at $2 \le k \le 6$. The control suite also compares all 1254 prime tuples from the primes at most 31, with $2 \le k \le 5$ in both signs, and tests the incomplete-result gate. A budgeted $k = 14$ run stops at its next-prime interval limit and is not reported as a completed exclusion. These controls are not new improvements on the sixteen-factor theorem.

The proof above covers every prescribed k; it supplies no induction establishing emptiness of all the minus-sign outputs. In particular,

$$\forall k \ge 2, \quad \mathcal{E}_-(k) = \varnothing$$

remains the unproved assertion needed for the full Lehmer conjecture, where $\mathcal{E}_-(k)$ is the output of this construction. The new appendix has a written proof and finite exact controls; it is not claimed as a new Lean formalization or a full-conjecture closure.

Appendix D. Exact prime completions and the obstruction to deletion descent

This appendix isolates the arithmetic assertion that remains after the finite-search reductions. All the identities and implications in this appendix are proved below. The universal exclusion in Problem D.5 is not proved. In particular, the sixteen-factor theorem is not enlarged to a resolution of Lehmer's conjecture by a change of title or by a terminating algorithm for each prescribed factor count.

D.1. **Notation and the unique possible last prime.** Let S be a nonempty finite set of distinct odd primes and put

$$A = \prod_{p \in S} p, \qquad B = \prod_{p \in S} (p-1) = \varphi(A), \qquad r = \max S. \tag{D.1}$$

These are integers, A is odd, and B is positive and even. Moreover $A/B > 1$ and A/B is not an integer: an even integer cannot divide the odd integer A. For an integer $M \geq 2$ write

$$D = MB - A. \tag{D.2}$$

An additional prime $q > r$ is different from every element of S. Consequently its totient contribution is exactly $q - 1$, and

$$\begin{aligned} \varphi(Aq) \mid Aq - 1 \text{ with quotient } M \quad &\iff \quad Aq - 1 = MB(q - 1) \\ &\iff \quad (MB - A)q = MB - 1 \\ &\iff \quad D(q - 1) = A - 1. \end{aligned} \tag{D.3}$$

The last line forces $D > 0$, since $A > 1$ and $q > 1$. There has been no division by a possibly zero integer in obtaining this conclusion.

Lemma D.1 (The quotient is forced)**.** *If $q > r$ is prime and $Aq - 1 = MB(q - 1)$, then*

$$0 < M - \frac{A}{B} = \frac{A - 1}{B(q - 1)} < \frac{1}{2}, \qquad M = \left\lceil \frac{A}{B} \right\rceil. \tag{D.4}$$

Proof. Every factor $j/(j - 1)$ is greater than 1 for an integer $j \geq 3$. Since S is a subset of $\{3, 4, \ldots, r\}$, inserting its missing integers can only increase the product. Hence

$$\frac{A}{B} = \prod_{p \in S} \frac{p}{p - 1} \leq \prod_{j=3}^{r} \frac{j}{j - 1} = \frac{r}{2}.$$

The two primes $q > r$ are odd, so $q \geq r + 2$, and therefore $r/(q - 1) < 1$. The identity $D(q - 1) = A - 1$ now gives

$$0 < \frac{D}{B} = \frac{A - 1}{B(q - 1)} < \frac{A/B}{q - 1} \leq \frac{r}{2(q - 1)} < \frac{1}{2}.$$

By $D = MB - A$ this is the stated estimate. An integer strictly between A/B and $A/B + 1/2$ must be its ceiling. □

Theorem D.2 (Exact final-prime criterion)**.** *For S as in (D.1), define*

$$M = \left\lceil \frac{A}{B} \right\rceil, \qquad D = MB - A.$$

There exists a Lehmer composite with prime support $S \cup \{q\}$ and $q > \max S$ if and only if

$$D \mid A - 1, \qquad q = 1 + \frac{A - 1}{D} > r, \qquad q \text{ is prime.} \tag{D.5}$$

The possible prime q is unique.

Proof. The observations above give $M \geq 2$ and $D > 0$ for this choice of M. If a Lehmer composite exists, Lemma D.1 forces this M, and (D.3) forces every condition in (D.5). Conversely, suppose those conditions hold. The primality and ordering of q imply $\varphi(Aq) = B(q - 1)$. The definition of q gives $Dq = D + A - 1 = MB - 1$, and hence

$$MB(q - 1) = (A + D)q - MB = Aq - 1.$$

Thus $\varphi(Aq) \mid Aq - 1$. Since S is nonempty and $q > r$, Aq has at least two distinct prime factors and is composite. Uniqueness follows from the displayed formula for q. □




FIGURE 19. The exact completion criterion. Theorem D.2 proves the equivalence represented by the diagram. Excluding an accepted certificate for every support is the unresolved assertion, not an additional established gate.

D.2. **What exact divisibility already enforces.**

Proposition D.3 (Consequences of an integral tail)**.** *Suppose $D = MB - A > 0$ and $D \mid A - 1$, and put $q = 1 + (A-1)/D$. The integer q need not be prime. Nevertheless*

$$\gcd(D, A) = \gcd(D, MB) = 1, \tag{D.6}$$

$$\gcd(A, MB) = 1, \tag{D.7}$$

$$\gcd(q, A) = \gcd(D - 1, A), \tag{D.8}$$

$$v_\ell(A-1) = v_\ell(D) + v_\ell(q-1) \quad \textit{for every prime } \ell. \tag{D.9}$$

If the full prime and ordering tests of (D.5) also hold, then $D < B/2$, D is odd and $D \geq 3$. If $M = 2$, then $D \geq 5$ and $3 \nmid D$.

Proof. A common divisor of D and A divides both $A - 1$ and A and is therefore 1. Since $MB = A + D$, this also proves $\gcd(D, MB) = 1$ and $\gcd(A, MB) = 1$. Reducing $Dq = MB - 1$ modulo A gives $Dq \equiv D - 1 \pmod{A}$. Multiplication by the invertible residue class D preserves the greatest common divisor with A; this proves (D.8). Equation (D.9) is the valuation of the exact product $A - 1 = D(q-1)$, with multiplicities included.

For a prime ordered tail, Lemma D.1 gives $D < B/2$. Since MB is even and A is odd, D is odd. If $D = 1$, then $q = A$. When S has one element this violates $q > r$; when it has more than one element it violates primality. Thus $D \geq 3$.

More generally, if $3 \mid D$, then (D.6) and (D.7) show that no $p \in S$ is 0 or 1 modulo 3. All are consequently 2 modulo 3. Writing $s = |S|$, $D \mid A - 1$ implies $(-1)^s \equiv 1 \pmod{3}$, so s is even. It follows that $B \equiv 1 \pmod{3}$. The equation $MB = A + D$ then gives $M \equiv 1 \pmod{3}$. For $M = 2$ this is impossible. Together with oddness and the exclusion of $D = 1$, it gives $D \geq 5$. □

In particular the exact tail identity already enforces every equality in (D.9); recording the same valuations in a second calculation is not an independent exclusion argument. Likewise

(D.7) includes the condition that no prime in S divides another prime in S minus one. These are useful pruning rules, but they do not prove that the prime-tail gate always fails.[13]

D.3. **The deletion obstruction.**

Proposition D.4 (No largest-prime deletion descent)**.** *Suppose $n = Aq$ is a Lehmer composite and q is its largest prime. With $M = (n-1)/\varphi(n)$ and $B = \varphi(A)$, one has*

$$M - 1 < \frac{A-1}{B} < M. \tag{D.10}$$

In particular $\varphi(A)$ does not divide $A - 1$.

Proof. Set $D = MB - A$. By Lemma D.1, $0 < D < B/2$. The integer D is odd and B is even. Thus $B > 2$ and $D \le B/2 - 1$, whence

$$0 < \frac{D+1}{B} \le \frac{1}{2} < 1.$$

Since $A = MB - D$, exact division gives

$$\frac{A-1}{B} = M - \frac{D+1}{B}.$$

The last two displays give (D.10). There is no integer strictly between the consecutive integers $M - 1$ and M, so the quotient cannot be an integer. □

The proposition rules out deleting the largest prime as a map from a Lehmer counterexample to a smaller Lehmer counterexample, even when the quotient is allowed to change. It does not rule out every possible descent. A different descent must specify its new support and establish both its prime-factor conditions and its totient identity. Minimality of a hypothetical counterexample does not supply these preservation properties.

D.4. **Two remaining primes: factorisation and its exact inverse.** For this subsection A, B describe an earlier prefix; $M \ge 2$ is fixed and $C = MB - A > 0$. Consider distinct primes $r < p < q$ satisfying

$$Apq - 1 = MB(p-1)(q-1). \tag{D.11}$$

Expanding the right side and moving terms gives, successively,

$$\begin{aligned} Cpq - MB(p+q) + MB + 1 &= 0, \\ C^2pq - CMB(p+q) + C(MB+1) &= 0, \\ (Cp - MB)(Cq - MB) &= (MB)^2 - C(MB+1). \end{aligned}$$

Using $MB - C = A$, the last right side is

$$K = MAB - C = A^2 + C(A-1). \tag{D.12}$$

It is positive for $A \ge 1$. For a genuine extension define

$$u = Cp - MB, \qquad v = Cq - MB. \tag{D.13}$$

The one-tail equation after choosing p gives $uq = MB(p-1) - 1 > 0$, so $u > 0$; then $v > u$ because $q > p$. We have obtained

$$uv = K, \qquad 0 < u < v, \qquad u \equiv v \equiv -MB \pmod{C}. \tag{D.14}$$

[13] A numerical primality test and an exact divisibility test play different roles. A certificate accepted by Theorem D.2 requires a proof of primality. A rejected composite tail may instead be rejected by an explicit proper divisor. Neither operation proves an assertion over supports that have not been covered.




FIGURE 20. The normalised two-tail hyperbola. Only its integer divisor pairs with the prescribed congruences can be used, and only prime recoveries yield a solution. The continuous curve asserts no existence of such divisor pairs.

Conversely, any pair of positive integers satisfying (D.14) recovers integers

$$p = \frac{u + MB}{C}, \qquad q = \frac{v + MB}{C}. \tag{D.15}$$

Reversing the displayed expansions proves (D.11). For it to be a Lehmer extension, it is still necessary that the recovered p, q be distinct primes greater than r. The factorisation alone does not establish those facts.

D.5. **Why an extreme factor switch is not a prime-preserving map.** For positive real $u \le \sqrt{K}$, the function $u + K/u$ decreases as u increases: its derivative is $1 - K/u^2 \le 0$. Moreover

$$pq = \frac{K + MB(u + K/u) + (MB)^2}{C^2}. \tag{D.16}$$

Thus a larger compatible smaller divisor would give a smaller product of recovered integers. This real-variable observation does not imply that a larger compatible divisor exists or that its recoveries are prime.

The extreme divisor $u = 1$ illustrates the distinction exactly. If $C \mid MB + 1$, then $C \mid A + 1$. Write $j = (A + 1)/C$, so that the recovered first entry is $p = j + 1$. Formula (D.13), or (D.11), yields

$$u(q - 1) = Ap - 1.$$

With $u = 1$ it follows that $q = Ap$. For $A > 1$ this q is composite, irrespective of whether p is prime. This calculation excludes that extreme recovery as a prime-preserving descent. It does not exclude all the other divisor pairs. For example the prefix $A = 15$, $B = 8$, $M = 2$ gives $C = 1$, $K = 239$ and $(p, q) = (17, 255)$. The exact equality

$$15 \cdot 17 \cdot 255 - 1 = 2 \cdot 8 \cdot 16 \cdot 254 = 65024$$

holds, but $255 = 3 \cdot 5 \cdot 17$ is composite. One must not replace the actual totient of this product by a product of the displayed entries minus one.[14]

[14]This is an integer identity used to check a proposed construction, not a Lehmer counterexample. It is consistent with the pseudo-solution families discussed earlier in the paper.

D.6. **The outstanding universal assertion.**

Problem D.5 (Universal prime-tail exclusion)**.** Prove that for every nonempty finite set S of distinct odd primes, the three tests in (D.5) cannot all hold.

Theorem D.2 proves that this assertion is equivalent to the full Lehmer totient conjecture. Appendix C gives finite and complete searches at every prescribed factor count. The bounds and computations in the main paper settle the stated finite ranges. Neither statement proves Problem D.5. The valuation identities, the deletion obstruction and the hyperbola factorisation above identify requirements for further arguments; they are not a substitute for that universal exclusion.

Declarations

Data and code availability. The programs that carry out the searches, the threshold computations and the checks of Section 8, their recorded output, the factorisations of the fifteen integers of Table 3, the logs of the three-prime searches of Section 5.2, the program and the journal of the eight-prime search of Section 7.4, the programs of Sections 2.1 and 6.3 and their output, the programs that check the numbers of Table 1 and of the proof of Proposition 6.11, Lean 4 proofs [27], checked with Mathlib [22], of many statements that do not depend on a search (among them Theorem 6.2 and Proposition 6.8), and the sources of this paper are available at `https://github.com/creelie/eLhmr`; the earlier reproducibility record is archived in [4]. An earlier archive under the same title was deposited by Mandal, Bhattacharjee and Bhattacharya [21]; its recorded creatorship is retained in that citation. The programs and logs for the reproduction checks of Appendix B and for Section 9.1 are supplied with this revision.

Funding. No funding was received for conducting this study.

Declaration of competing interest. The author has no competing interests to declare that are relevant to the content of this article.

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process. During the preparation of this work the author used Claude (Anthropic) to write and run the programs, to write the Lean proofs and to draft text. After using this tool, the author reviewed and edited the content as needed and takes full responsibility for the content of the publication. For this revision, OpenAI Codex supplied the ceiling reduction and its proof in Proposition 9.2, the one-candidate test and defect restrictions in Corollaries 9.3 and 9.4, and the accompanying verification program and reproduction checks in Appendix B. These contributions are identified separately from the results of the original preprint. OpenAI Codex also assisted the written construction in Appendix C, its exact-arithmetic implementation and its positive and negative controls. The new construction is not a proof of the full conjecture; author review of the added material is required before journal or preprint submission. For the present revision, Codex also assisted the detailed derivations and diagrams in Appendix D, the audit of candidate descent operations, and preparation of the `amsart` source and submission files. The new appendix is a written argument and is not claimed as a new Lean formalisation. The all-prefix exclusion remains unproved.

References

[1] L. M. Adleman, C. Pomerance, and R. S. Rumely, *On distinguishing prime numbers from composite numbers*, Ann. of Math. (2) **117** (1983), no. 1, 173–206. doi:10.2307/2006975

[2] R. Baillie and S. S. Wagstaff, Jr., *Lucas pseudoprimes*, Math. Comp. **35** (1980), no. 152, 1391–1417. doi:10.1090/S0025-5718-1980-0583518-6

[3] W. D. Banks, C. W. Nevans, and C. Pomerance, *A remark on Giuga's conjecture and Lehmer's totient problem*, Albanian J. Math. **3** (2009), no. 2, 81–85. doi:10.51286/albjm/1245207560

[4] D. Bhattacharjee, *Lehmer's totient problem with fewer than sixteen prime factors: programs, data and Lean proofs*, version 1.5.0, 2026. https://github.com/creelie/eLhmr/releases/tag/v1.5.0

[5] J. Brillhart, D. H. Lehmer, and J. L. Selfridge, *New primality criteria and factorizations of* $2^m \pm 1$, Math. Comp. **29** (1975), no. 130, 620–647. doi:10.1090/S0025-5718-1975-0384673-1

[6] P. Burcsi, S. Czirbusz, and G. Farkas, *Computational investigation of Lehmer's totient problem*, Ann. Univ. Sci. Budapest. Sect. Comput. **35** (2011), 43–49. doi:10.71352/ac.35.043

[7] D. Burek and B. Żmija, *A new upper bound for numbers with the Lehmer property and its application to repunit numbers*, Int. J. Number Theory **15** (2019), no. 7, 1463–1468. doi:10.1142/S1793042119500830

[8] G. L. Cohen and P. Hagis, Jr., *On the number of prime factors of* n *if* $\varphi(n) \mid (n-1)$, Nieuw Arch. Wisk. (3) **28** (1980), 177–185.

[9] H. Cohen and H. W. Lenstra, Jr., *Primality testing and Jacobi sums*, Math. Comp. **42** (1984), no. 165, 297–330. doi:10.1090/S0025-5718-1984-0726006-X

[10] R. J. Cook, *Bounds for odd perfect numbers*, in: Number Theory (Ottawa, 1996), CRM Proc. Lecture Notes **19**, Amer. Math. Soc., Providence, RI, 1999, 67–71.

[11] D. Coppersmith, N. Howgrave-Graham, and S. V. Nagaraj, *Divisors in residue classes, constructively*, Math. Comp. **77** (2008), no. 261, 531–545. doi:10.1090/S0025-5718-07-02007-8

[12] J. Feitsma and W. Galway, *Tables of pseudoprimes and related data*, https://www.cecm.sfu.ca/Pseudoprimes/index-2-to-64.html.

[13] The FLINT team, *FLINT: Fast Library for Number Theory*, version 3.6.0, https://flintlib.org.

[14] S. W. Golomb, *Sets of primes with intermediate density*, Math. Scand. **3** (1955), 264–274. doi:10.7146/math.scand.a-10445

[15] A. Grytczuk and M. Wójtowicz, *On a Lehmer problem concerning Euler's totient function*, Proc. Japan Acad. Ser. A Math. Sci. **79** (2003), no. 8, 136–138. doi:10.3792/pjaa.79.136

[16] R. K. Guy, *Unsolved Problems in Number Theory*, 3rd ed., Problem Books in Mathematics, Springer, New York, 2004. doi:10.1007/978-0-387-26677-0

[17] P. Hagis, Jr., *On the equation* $M \cdot \varphi(n) = n - 1$, Nieuw Arch. Wisk. (4) **6** (1988), no. 3, 255–261.

[18] D. R. Heath-Brown, *Odd perfect numbers*, Math. Proc. Cambridge Philos. Soc. **115** (1994), no. 2, 191–196. doi:10.1017/S0305004100072030

[19] D. H. Lehmer, *On Euler's totient function*, Bull. Amer. Math. Soc. **38** (1932), no. 10, 745–751. doi:10.1090/S0002-9904-1932-05521-5

[20] H. W. Lenstra, Jr., *Divisors in residue classes*, Math. Comp. **42** (1984), no. 165, 331–340. doi:10.1090/S0025-5718-1984-0726007-1

[21] P. Mandal, D. Bhattacharjee, and U. Bhattacharya, *Code, data and Lean proofs for Lehmer's totient problem with fewer than sixteen prime factors*, version 2.5.2, Zenodo, 2026. doi:10.5281/zenodo.23171399

[22] The mathlib Community, *The Lean mathematical library*, in: Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), ACM, New York, 2020, 367–381. doi:10.1145/3372885.3373824

[23] H. G. Meijer, *Sets of primes with intermediate density*, Math. Scand. **34** (1974), 37–43. doi:10.7146/math.scand.a-11502

[24] G. Molnar and G. Singh, *Spoof Lehmer factorizations*, Integers **26** (2026), #A33, 15 pp. doi:10.5281/zenodo.18714466

[25] H. L. Montgomery, *A note on the large sieve*, J. London Math. Soc. **43** (1968), 93–98. doi:10.1112/jlms/s1-43.1.93

[26] H. L. Montgomery and R. C. Vaughan, *The large sieve*, Mathematika **20** (1973), no. 2, 119–134. doi:10.1112/S0025579300004708

[27] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in: Automated Deduction – CADE 28, Lecture Notes in Comput. Sci., Springer, Cham, 2021, 625–635. doi:10.1007/978-3-030-79876-5_37

[28] P. P. Nielsen, *An upper bound for odd perfect numbers*, Integers **3** (2003), #A14.

[29] M. Norman, *A group-theoretical approach to Lehmer's totient problem*, preprint, 2021. arXiv:2106.11781

[30] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, sequences A050474 and A203966, https://oeis.org, accessed September 2026.

[31] The PARI Group, *PARI/GP*, version 2.15.4, Univ. Bordeaux, https://pari.math.u-bordeaux.fr.

[32] R. G. E. Pinch, *A note on Lehmer's totient problem*, poster, Seventh Algorithmic Number Theory Symposium (ANTS VII), Berlin, 2006. https://antsmath.org/ANTSVII/Poster/Pinch_Poster3.pdf

[33] C. Pomerance, *On composite n for which $\varphi(n) \mid n-1$, II*, Pacific J. Math. **69** (1977), no. 1, 177–186. doi:10.2140/pjm.1977.69.177
[34] C. Pomerance, J. L. Selfridge, and S. S. Wagstaff, Jr., *The pseudoprimes to* $25 \cdot 10^9$, Math. Comp. **35** (1980), no. 151, 1003–1026. doi:10.1090/S0025-5718-1980-0572872-7
[35] J. Renze, *Computational evidence for Lehmer's totient conjecture*, Mathematica notebook, Wolfram Library Archive, MathSource item 5483, 2004. https://library.wolfram.com/infocenter/MathSource/5483/
[36] J. Sorenson and J. Webster, *Strong pseudoprimes to twelve prime bases*, Math. Comp. **86** (2017), no. 304, 985–1003. doi:10.1090/mcom/3134
[37] A. Stone, *Improved upper bounds for odd perfect numbers – Part I*, Integers **24** (2024), #A114. doi:10.5281/zenodo.14547960
[38] E. Wong, *Computations on normal families of primes*, M.Sc. thesis, Simon Fraser University, 1997. https://summit.sfu.ca/_flysystem/fedora/sfu_migrate/7394/b18771816.pdf
[39] T. Yamada, *On Lehmer's problem and related problems*, preprint, 2023; also RIMS Kôkyûroku No. 2285. arXiv:2303.16853

Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL), Bhubaneswar, Odisha 751030, India
Email address: d.bhattacharjee@erl-forschung.de
Email address: itsdeep@live.com
URL: https://orcid.org/0000-0003-0466-750X