ISOMORPHIC INVARIANT ORDERS WITHOUT HERMITE EQUIVALENCE

DEEP BHATTACHARJEE

Abstract. Bhargava, Evertse, Győry, Remete and Swaminathan asked whether, in every degree $n \geq 4$, there are integral polynomials with isomorphic invariant orders that are not Hermite equivalent; they answered the question for $n = 4$. We answer it for every even n and for every n with $n-2$ divisible by 3, 5 or 7, with explicit pairs. In degree six one polynomial of the pair can be chosen monic.

Keywords: Hermite equivalence; invariant order; binary form.

MSC 2020: 11R04 (primary); 11C08, 11E16, 11R29.

1. Introduction

In 1857 Hermite introduced an equivalence relation on integral polynomials of a given degree n that is coarser than the classical $\mathrm{GL}_2(\mathbb{Z})$-equivalence, and proved that there are only finitely many classes of a given nonzero discriminant. Bhargava, Evertse, Győry, Remete and Swaminathan revived the notion and gave it an algebraic meaning [1]. To a polynomial

$$f(X) = f_0X^n + f_1X^{n-1} + \cdots + f_n \in \mathbb{Z}[X], \qquad f_0 \neq 0, \quad D(f) \neq 0,$$

one attaches its *invariant order* R_f and its *invariant ideal* I_f, a fractional ideal of R_f; both live in the étale algebra $K_f = \mathbb{Q}[X]/(f)$ and go back to Birch–Merriman, Nakagawa, Simon and Wood [5, 9, 10, 11]. For primitive f and g of nonzero discriminant, f and g are Hermite equivalent if and only if there is a ring isomorphism $R_f \cong R_g$ under which the ideal classes of I_f and I_g correspond [1, Corollary 3.12]. In degree three the condition on ideals is automatic, by the Levi–Delone–Faddeev correspondence, and for monic polynomials it is empty, since then $I_f = R_f$. This led the authors to the following question.[1]

Question (Bhargava–Evertse–Győry–Remete–Swaminathan)**.** Let $n \geq 4$. Do there exist (reducible or irreducible) polynomials $f, g \in \mathbb{Z}[X]$ of degree n and nonzero discriminant that have isomorphic invariant orders but are not Hermite equivalent, and if so, can such polynomials be exhibited?

In [1, Theorem 3.20] they exhibited two quartic polynomials with isomorphic invariant orders, one with principal and one with non-principal invariant ideal, which settles $n = 4$. For $n \geq 5$ the question has remained open. Our main result settles it in most degrees.

Theorem A. *Let $n \geq 5$ and suppose that n is even or that $n-2$ is divisible by 3, 5 or 7. Then there exist primitive polynomials $P, P' \in \mathbb{Z}[X]$ of degree n and the same nonzero discriminant whose invariant orders are isomorphic and which are not Hermite equivalent. For $n \geq 6$ there are infinitely many such pairs with pairwise distinct discriminants.*

Together with the quartic case, the question is answered for every $n \geq 4$ except the odd n for which $n-2$ is prime to 105; the first of these are $13, 15, 19, 21, 25$ (Figure 1). In the two smallest degrees the pairs are short enough to write down.

[1] Question 3.18 of [1]. We quote it in full.

Theorem B. *The sextic polynomials*

$$P(X) = X^6 - 18X^4 - 6X^3 + 79X^2 + 60X + 10,$$
$$P'(X) = 3X^6 + 8X^5 - 22X^4 - 82X^3 - 67X^2 - 20X - 2$$

have the same discriminant and isomorphic invariant orders, and they are not Hermite equivalent. The same holds for the quintic polynomials

$$Q(X) = 14X^5 - 19X^4 - 1676X^3 + 3714X^2 + 50843X - 152646,$$
$$Q'(X) = 42X^5 - 169X^4 - 2608X^3 + 12820X^2 + 975X - 45526.$$

The sextic P is monic. By [1, Corollary 3.16] two monic polynomials with isomorphic invariant orders are always Hermite equivalent, so Theorem B shows that this statement fails as soon as only one of the two polynomials is monic.

The idea behind both theorems is simple. Take two binary quadratic forms q_0 and q_1 of the same discriminant d, so that their invariant orders coincide, with I_{q_0} principal and I_{q_1} not. A single binary form H that is coprime to both q_0 and q_1 at every prime, in the sense that $\mathrm{Res}(q_0, H) = \mathrm{Res}(q_1, H) = \pm 1$, splits off a common factor: the invariant order of $q_i H$ is the product of the invariant orders of q_i and H, and the invariant ideal splits in the same way. The two products then have the same ring but invariant ideals that differ in exactly one factor, and a count of the principal factors shows that no isomorphism can identify them. The whole difficulty is to find such an H in the required degree. Section 5 shows that the obstruction lives in the Picard group of the quartic ring attached to $q_0 q_1$, and that one form of degree k produces forms in every degree divisible by k. Section 6 gives the four quadratic fields and the five forms that prove Theorem A.




FIGURE 1. Degrees $4 \le n \le 120$ in which Question 3.18 of [1] is answered. Grey: the quartic example of [1]. Coloured: Theorem A, by the first divisibility condition it satisfies. White: degrees left open.

2. Invariant orders, invariant ideals and resultants

We identify a polynomial $f \in \mathbb{Z}[X]$ of degree n with the binary form $f(x, y) = y^n f(x/y)$ and use the same letter for both. Throughout, $f_0 = f(1, 0) \neq 0$ and $D(f) \neq 0$. Let α be the image of X in $K_f = \mathbb{Q}[X]/(f)$. The $\mathbb{Z}$-module $I_f(k) \subset K_f$, for $0 \le k \le n-1$, has basis

$$1,\ \alpha,\ \ldots,\ \alpha^k,\ \zeta_{k+1},\ \ldots,\ \zeta_{n-1}, \qquad \zeta_i = f_0\alpha^i + f_1\alpha^{i-1} + \cdots + f_{i-1}\alpha,$$

and $R_f = I_f(0)$, $I_f = I_f(1)$ [1, §3.1]. We use four facts from [1, Theorem 3.3], due to Birch–Merriman, Nakagawa, Simon and Wood:

(F1) R_f is an order in K_f and $D(R_f) = D(f)$.
(F2) $I_f(k)$ is a fractional ideal of R_f; it is invertible if and only if f is primitive, and then $I_f(k) = I_f^k$.
(F3) Let $S_f \subset \mathbb{P}^1_{\mathbb{Z}}$ be the closed subscheme cut out by f. Then R_f is the ring of global functions on S_f and $I_f(k)$ is the module of global sections of $\mathcal{O}(k)|_{S_f}$. Both are viewed inside K_f through the generic fibre $S_f \times \mathbb{Q} = \operatorname{Spec} K_f$, sections of $\mathcal{O}(k)$ being divided by y^k.
(F4) Primitive f, g of degree n and nonzero discriminant are Hermite equivalent if and only if there is a ring isomorphism $\varphi\colon R_f \to R_g$ with $\varphi(I_f)$ in the ideal class of I_g [1, Corollary 3.12].

Statement (F3) is Wood's geometric description [11, §2 and Appendix A].[2] The identification inside K_f uses that $f_0 \neq 0$, so the generic fibre of S_f lies in the chart $y \neq 0$, where it is $\operatorname{Spec} \mathbb{Q}[t]/(f(t,1))$; the section $x^j y^{k-j}$ goes to α^j.

For a binary quadratic form $q = ax^2 + bxy + cy^2$ of non-square discriminant $d = b^2 - 4ac$ with $a \neq 0$, let $\theta = (-b + \sqrt{d})/(2a)$. Then $K_q = \mathbb{Q}(\sqrt{d})$, $R_q = \mathbb{Z} + \mathbb{Z}a\theta = \mathbb{Z}[(d + \sqrt{d})/2]$ is the quadratic order $\mathcal{O}_d$ of discriminant d, and $I_q = \mathbb{Z} + \mathbb{Z}\theta$. In particular two quadratic forms of the same discriminant have literally the same invariant order. If q is primitive, then $aI_q = \mathbb{Z}a + \mathbb{Z}\frac{-b+\sqrt{d}}{2}$ is the invertible ideal of norm $|a|$ classically attached to q. If $a = 1$ then $I_q = R_q$ is principal.

For binary forms F, G of degrees $a, b \geq 1$ we write $\operatorname{Res}(F, G)$ for their resultant. We need the following standard properties [8, Chapter IV, §8 and Chapter IX, §3].

Lemma 2.1. *Let F, G, G_1, G_2, M be integral binary forms with $\deg F = a$, $\deg G = b$ and $\deg M = b - a \geq 0$.*

(1) $\operatorname{Res}(F, G) = \pm 1$ *if and only if F and G have no common zero in $\mathbb{P}^1(\overline{\mathbb{F}}_p)$ for any prime p.*
(2) $\operatorname{Res}(F, G_1G_2) = \operatorname{Res}(F, G_1)\operatorname{Res}(F, G_2)$ *and* $\operatorname{Res}(F, cG) = c^a \operatorname{Res}(F, G)$ *for $c \in \mathbb{Z}$.*
(3) $\operatorname{Res}(F, G + FM) = \operatorname{Res}(F, G)$.
(4) $D(FG) = D(F)D(G)\operatorname{Res}(F, G)^2$.

Proof. The resultant of binary forms is a universal integral polynomial in the coefficients, so it commutes with reduction modulo p, and over an algebraically closed field it vanishes exactly when the two forms have a common zero in $\mathbb{P}^1$. Since $\operatorname{Res}(F, G)$ is an integer, it equals ± 1 if and only if it is not divisible by any prime, which is (1). If $F = f_0 \prod_{i=1}^{a}(x - \alpha_i y)$ with $f_0 \neq 0$, then $\operatorname{Res}(F, G) = f_0^b \prod_i G(\alpha_i, 1)$ up to a sign depending only on a and b; this gives (2) and (3), because FM vanishes at every $(\alpha_i, 1)$. The general case of (2) and (3) follows by continuity in the coefficients, or by a change of variables in $\mathrm{GL}_2(\mathbb{Z})$ that makes $F(1, 0) \neq 0$. Part (4) is the usual product formula for discriminants. □

3. Splitting off a common factor

Lemma 3.1. *Let F and G be primitive binary forms of degrees $a, b \geq 2$ with nonzero leading coefficients $F(1, 0)$, $G(1, 0)$, nonzero discriminants, and $\operatorname{Res}(F, G) = \pm 1$. Then the isomorphism of $\mathbb{Q}$-algebras*

$$K_{FG} \longrightarrow K_F \times K_G, \qquad X \longmapsto (\alpha_F, \alpha_G),$$

[2] Wood works over an arbitrary base scheme and also treats forms of discriminant zero; we only need the case of integral forms with $f_0 D(f) \neq 0$, where S_f is finite and flat over $\operatorname{Spec} \mathbb{Z}$.

given by the Chinese remainder theorem maps R_{FG} onto $R_F \times R_G$ and I_{FG} onto $I_F \times I_G$.

Proof. On the affine chart $y \neq 0$ of $\mathbb{P}^1_{\mathbb{Z}}$ the schemes S_F, S_G and S_{FG} are cut out by $F(t,1)$, $G(t,1)$ and their product in $\mathbb{Z}[t]$. By Lemma 2.1(1) the scheme $\operatorname{Spec}\mathbb{Z}[t]/(F(t,1),G(t,1))$ has no points, so the two ideals are comaximal. For comaximal ideals the intersection equals the product, and the Chinese remainder theorem gives

$$\mathbb{Z}[t]/(F(t,1)G(t,1)) \cong \mathbb{Z}[t]/(F(t,1)) \times \mathbb{Z}[t]/(G(t,1)).$$

The same holds on the chart $x \neq 0$. Hence S_{FG} is the disjoint union of its closed subschemes S_F and S_G, and global sections of $\mathcal{O}(k)$ on S_{FG} are pairs of global sections on S_F and on S_G. On the generic fibre this decomposition is the map in the statement, since $\mathbb{Q}[t]/(F(t,1)G(t,1)) \cong \mathbb{Q}[t]/(F(t,1)) \times \mathbb{Q}[t]/(G(t,1))$ sends t to (α_F, α_G), and dividing by y^k commutes with it. Apply (F3) with $k = 0$ and $k = 1$; the latter is allowed because $a, b \geq 2$. □




FIGURE 2. A schematic picture of $\mathbb{P}^1_{\mathbb{Z}} \to \operatorname{Spec}\mathbb{Z}$. The subschemes S_{q_0} and S_{q_1} of two quadratic forms of the same discriminant always meet, here over a single prime, in a finite scheme T. Theorem 4.1 needs a form H whose subscheme S_H avoids both of them in every fibre.

4. The twisting principle

Theorem 4.1. *Let q_0 and q_1 be primitive binary quadratic forms of the same non-square discriminant d with $q_0(1,0)q_1(1,0) \neq 0$, such that I_{q_0} is a principal ideal of $\mathcal{O}_d$ and I_{q_1} is not. Let H be a binary form of degree $k \geq 2$ with $H(1,0)D(H) \neq 0$ and*

$$\operatorname{Res}(q_0, H) = \pm 1, \qquad \operatorname{Res}(q_1, H) = \pm 1.$$

Then $P = q_0 H$ and $P' = q_1 H$ are primitive of degree $k+2$, $D(P) = D(P') = d\,D(H) \neq 0$, $R_P \cong R_{P'}$, and P and P' are not Hermite equivalent.

Proof. If a prime ℓ divided every coefficient of H, then ℓ^2 would divide $\operatorname{Res}(q_0, H)$ by Lemma 2.1(2). So H is primitive, and P, P' are primitive by Gauss's lemma. Their leading coefficients $q_i(1,0)H(1,0)$ are nonzero, and Lemma 2.1(4) gives $D(P) = D(P') = d\,D(H)$, which is nonzero.

By Lemma 3.1, applied to (q_0, H) and to (q_1, H),

$$R_P \cong \mathcal{O}_d \times R_H \cong R_{P'}, \qquad I_P \leftrightarrow I_{q_0} \times I_H, \qquad I_{P'} \leftrightarrow I_{q_1} \times I_H.$$

Suppose that P and P' were Hermite equivalent. By (F4) there is a ring isomorphism $\varphi\colon R_P \to R_{P'}$ and $\kappa \in K_{P'}^{\times}$ with $\varphi(I_P) = \kappa I_{P'}$.

The order R_H lies in the reduced algebra K_H, so it has finitely many idempotents, and it is the product of the rings eR_H over its primitive idempotents e; each eR_H has no idempotents other than 0 and e. The primitive idempotents of $R_P = \mathcal{O}_d \times R_H$ are $(1,0)$ and the $(0,e)$, because $\mathcal{O}_d$ is a domain. A ring isomorphism sends primitive idempotents bijectively to primitive idempotents, and for each primitive idempotent ε of R_P it restricts to an isomorphism $\varepsilon R_P \to \varphi(\varepsilon)R_{P'}$ carrying εI_P onto $\varphi(\varepsilon)\kappa I_{P'}$, a principal multiple of $\varphi(\varepsilon)I_{P'}$.

For a ring A with an ideal J write $N(A, J)$ for the number of primitive idempotents ε of A such that εA is isomorphic to $\mathcal{O}_d$ and εJ is a principal ideal of εA. The previous paragraph shows $N(R_P, I_P) = N(R_{P'}, I_{P'})$. On the other hand the component $(1,0)$ contributes 1 to $N(R_P, I_P)$, because I_{q_0} is principal, and 0 to $N(R_{P'}, I_{P'})$, because I_{q_1} is not; whether an ideal of $\mathcal{O}_d$ is principal does not depend on the choice of isomorphism, since $\mathcal{O}_d$ has only the identity and conjugation as automorphisms. The components $(0,e)$ contribute the same amount to both counts, because they carry the same pair (eR_H, eI_H). Hence $N(R_P, I_P) = N(R_{P'}, I_{P'}) + 1$, a contradiction. □

Remark 4.2. The proof uses only that I_{q_0} and I_{q_1} lie in different orbits of $\operatorname{Pic}(\mathcal{O}_d)$ under conjugation. The principal case is the most convenient, because then q_0 can be taken monic.

5. Finding the common factor

Fix q_0, q_1 as in Theorem 4.1 and call a binary form H *admissible* if $\operatorname{Res}(q_0, H) = \pm 1$ and $\operatorname{Res}(q_1, H) = \pm 1$. Admissibility is stable under products and under adding multiples of $q_0 q_1$.

Proposition 5.1. *Let $H_1, \dots, H_s$ be admissible forms of degrees $k_1, \dots, k_s$, let $m_1, \dots, m_s \geq 0$, and suppose $k = \sum m_j k_j \geq 4$. Put $M = \prod_{i=1}^{k-4}(x - iy)$ and*

$$H_t = \prod_{j=1}^{s} H_j^{m_j} + t\, q_0 q_1 M \qquad (t \in \mathbb{Z}).$$

Then every H_t is admissible of degree k, and $H_t(1,0)D(H_t) \neq 0$ for all but finitely many t. The values $D(H_t)$ are not all equal.

Proof. Admissibility of H_t follows from Lemma 2.1(2) and (3), since q_0q_1M is divisible by q_0 and by q_1. The leading coefficient $H_t(1,0)$ is a polynomial of degree one in t with slope $q_0(1,0)q_1(1,0) \neq 0$. The discriminant is homogeneous of degree $2k-2$ in the coefficients, so $D(H_t)$ is a polynomial in t of degree at most $2k-2$ whose coefficient of t^{2k-2} is

$$D(q_0q_1M) = D(q_0)\,D(q_1)\,D(M)\,\operatorname{Res}(q_0,q_1)^2\,\operatorname{Res}(q_0,M)^2\,\operatorname{Res}(q_1,M)^2.$$

This is nonzero: q_0 and q_1 are irreducible over $\mathbb{Q}$ and not proportional, because their invariant ideals are not in the same class; M splits into distinct rational linear factors. Hence $D(H_t)$ is a nonconstant polynomial in t, which vanishes for finitely many t only and takes infinitely many values. □

The next statement explains when admissible forms exist. It is not needed for the proof of Theorem A, but it is how the forms in Section 6 were found, and it shows what goes wrong in the remaining degrees.

Proposition 5.2. *Let $F = q_0q_1$ and $k \geq 3$. An admissible form of degree k exists if and only if I_F^k is a principal ideal of the quartic order R_F.*

Proof. By Lemma 2.1(1),(2) a form H of degree k is admissible if and only if H has no zero on S_F, that is, if and only if the section $H|_{S_F}$ of the invertible sheaf $\mathcal{O}(k)|_{S_F}$ vanishes nowhere. Since S_F is affine, this holds if and only if $H|_{S_F}$ generates the module of global sections, which is I_F^k by (F2) and (F3); for $k \geq 4$ one uses that $\mathcal{O}(k)|_{S_F}$ is the k-th tensor power of the invertible sheaf $\mathcal{O}(1)|_{S_F}$, whose module of sections I_F is invertible, so that the sections form $I_F^{\otimes k} = I_F^k$. So an admissible H makes I_F^k principal. Conversely, the sequence $0 \to \mathcal{O}(k-4) \xrightarrow{F} \mathcal{O}(k) \to \mathcal{O}(k)|_{S_F} \to 0$ on $\mathbb{P}^1_{\mathbb{Z}}$ and the vanishing of $H^1(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{O}(k-4))$ for $k \geq 3$ [7, Theorem III.5.1] show that every global section of $\mathcal{O}(k)|_{S_F}$ is the restriction of a form of degree k. A generator of I_F^k is therefore the restriction of an admissible form. □

The class of I_F maps to $(1, [I_{q_1}])$ under $\operatorname{Pic}(R_F) \to \operatorname{Pic}(\mathcal{O}_d)^2$, so I_F^k can be principal only if $[I_{q_1}]^k = 1$. The kernel of that map is $\mathcal{O}_T^\times$ modulo the images of the two unit groups, where $\mathcal{O}_T$ denotes the ring of functions on the finite scheme $T = S_{q_0} \cap S_{q_1}$ (Figure 2). The order of I_F can therefore be larger than that of $[I_{q_1}]$, and it depends on the chosen forms within their classes; a large unit group helps, which is why all our examples come from real quadratic fields. Since an admissible form of degree $n-2$ forces $[I_{q_1}]^{n-2} = 1$ while $[I_{q_1}] \neq 1$, an odd degree n always needs a quadratic order whose class group has an element of odd order dividing $n-2$.

6. Four quadratic fields

Table 1 lists four pairs (q_0, q_1). In each case q_0 is monic, so $I_{q_0} = \mathcal{O}_d$ is principal. The discriminant d is fundamental, $\mathcal{O}_d$ is the full ring of integers of $\mathbb{Q}(\sqrt{d})$, and the class number h is given for orientation only.

d	h	q_0	q_1	admissible base forms
40	2	$x^2 - 10y^2$	$2x^2 + 4xy - 3y^2$	H_4, H_6
229	3	$x^2 + xy - 57y^2$	$3x^2 - 5xy - 17y^2$	H_3
401	5	$x^2 + xy - 100y^2$	$2x^2 + 23xy + 16y^2$	H_5
577	7	$x^2 + xy - 144y^2$	$2x^2 - 11xy - 57y^2$	H_7

Table 1. The quadratic data. The base forms are listed in the appendix.

Lemma 6.1. *In each row of Table 1 the ideal I_{q_1} is not principal.*

Proof. Write $q_1 = ax^2 + bxy + cy^2$. By Section 2, $aI_{q_1} = \mathbb{Z}a + \mathbb{Z}\frac{-b+\sqrt{d}}{2}$ is an ideal $\mathfrak{p}$ of norm a, which is the prime 2 or 3. If $\mathfrak{p}$ were principal, $\pm a$ would be the norm of an element of $\mathcal{O}_d$ not divisible by any rational integer, so $\pm a$ would be primitively represented by the principal form q_0.

For $d = 40$ this is impossible modulo 5: $x^2 - 10y^2 \equiv x^2 \pmod 5$, and ± 2 are not squares modulo 5.

For the other three rows we use reduction theory of indefinite forms [6, Chapter 3]. A form (A, B, C) of discriminant $d > 0$ is reduced if $0 < B < \sqrt{d}$ and $\sqrt{d} - B < 2|A| < \sqrt{d} + B$; two reduced forms are equivalent if and only if they lie on the same cycle. If m is primitively represented by q_0 and $|m| < \sqrt{d}/2$, then q_0 is equivalent to a form (m, B, C), and replacing B by its representative modulo $2|m|$ in the interval $(\sqrt{d} - 2|m|, \sqrt{d})$ makes it reduced. So $\pm a$ would appear as a first coefficient on the cycle of q_0. Here $a < \sqrt{d}/2$ in all three cases ($\sqrt{d}/2 > 7.5$), and the cycles of the principal forms are

$$d = 229: \quad (1, 15, -1),\ (-1, 15, 1);$$

$$d = 401: \quad (1, 19, -10),\ (-10, 1, 10),\ (10, 19, -1),\ (-1, 19, 10),\ (10, 1, -10),\ (-10, 19, 1);$$

$$d = 577: \quad (1, 23, -12),\ (-12, 1, 12),\ (12, 23, -1),\ (-1, 23, 12),\ (12, 1, -12),\ (-12, 23, 1).$$

Each step $(A, B, C) \mapsto (C, B', C')$ chooses $B' \equiv -B \pmod{2|C|}$ in $(\sqrt{d} - 2|C|, \sqrt{d})$, and one checks $B'^2 - 4CC' = d$ directly. The first coefficients are ± 1 for $d = 229$ and $\pm 1, \pm 10$, resp. $\pm 1, \pm 12$, for $d = 401$, 577. Neither ± 3 nor ± 2 occurs. □

Lemma 6.2. *The forms H_4, H_6 (for $d = 40$), H_3 (for $d = 229$), H_5 (for $d = 401$) and H_7 (for $d = 577$) listed in the appendix are admissible for the corresponding pair (q_0, q_1), have nonzero leading coefficient and nonzero discriminant, and are irreducible over $\mathbb{Q}$.*

Proof. For each base form H and each i the number $\mathrm{Res}(q_i, H)$ is the determinant of an integer Sylvester matrix of size $\deg H + 2$; equivalently, with $q_i = a_i x^2 + b_i xy + c_i y^2$ and $\theta_i = (-b_i + \sqrt{d})/(2a_i)$, it equals $a_i^{\deg H} N_{\mathbb{Q}(\sqrt{d})/\mathbb{Q}}(H(\theta_i, 1))$. These finite computations give the values in the appendix, all equal to ± 1. Irreducibility and $D(H) \neq 0$ are again finite checks; irreducibility is not used below. □

Proof of Theorem A. Let $n \geq 5$ and $k = n - 2 \geq 3$. If n is even, write $k = 4a + 6b$ with $a, b \geq 0$ and use the row $d = 40$ with H_4, H_6. If $3 \mid k$, $5 \mid k$ or $7 \mid k$, use the row $d = 229$, 401 or 577 with the single base form H_3, H_5 or H_7 and exponent $k/3$, $k/5$ or $k/7$. If k equals the degree of the base form, take H to be that form; it is admissible with $H(1, 0)D(H) \neq 0$ by Lemma 6.2. Otherwise $k \geq 4$, and Proposition 5.1 gives admissible H_t of degree k with $H_t(1, 0)D(H_t) \neq 0$ for all but finitely many t, and with infinitely many distinct discriminants. By Lemma 6.1, Theorem 4.1 applies to $P = q_0 H$ and $P' = q_1 H$. Their common discriminant $d\, D(H_t)$ takes infinitely many values as t varies, which proves the last assertion; $k \geq 4$ holds for every $n \geq 6$. □

Proof of Theorem B. For the sextic pair take $d = 40$, $q_0 = x^2 - 10y^2$, the form $q_1^* = 3x^2 + 8xy + 2y^2$ of discriminant 40, and $H = x^4 - 8x^2y^2 - 6xy^3 - y^4$. Then $P = q_0 H$ and $P' = q_1^* H$, and one computes $\mathrm{Res}(q_0, H) = \mathrm{Res}(q_1^*, H) = 1$ and $D(H) \neq 0$. The ideal $3I_{q_1^*} = \mathbb{Z}3 + \mathbb{Z}(-4 + \sqrt{10})$ has norm 3, and $x^2 - 10y^2 = \pm 3$ has no solution modulo 5, so $I_{q_1^*}$ is not principal. Theorem 4.1 applies. For the quintic pair take the row $d = 229$ of Table 1 and $H = H_3$; then $Q = q_0 H_3$ and $Q' = q_1 H_3$. □

7. Concluding remarks

Remark 7.1. The forms found here come with a great deal of freedom. Proposition 5.1 produces, in each admissible degree $k \geq 4$, an arithmetic progression of pairs, and different choices of q_1 in its $\mathrm{SL}_2(\mathbb{Z})$-orbit, or of the auxiliary form M, give further families. By Proposition 5.2, the existence of an admissible form of degree k depends only on the class of $I_{q_0 q_1}^{k}$, so the search for base forms reduces to a computation in $\mathcal{O}_T^{\times}$ modulo units.

Remark 7.2. The remaining odd degrees, those with $\gcd(n-2, 105) = 1$, need for some prime $\ell \mid n-2$ a quadratic order with an ideal class of order ℓ together with a pair (q_0, q_1) for which $I_{q_0 q_1}^{\ell}$ is principal. Real quadratic fields with class number divisible by ℓ exist for every ℓ, and for each fixed ℓ the condition is a finite computation, so every further prime ℓ adds the residue class $n \equiv 2 \pmod{\ell}$. A proof covering all odd n at once would need control over the unit group of $\mathcal{O}_d$ modulo the intersection scheme T in a family of fields, which we do not have.

Acknowledgements

The author thanks the authors of [1] for formulating the question so precisely, and acknowledges that the framework used here, the invariant orders and ideals and their geometric description, is due to Birch and Merriman, Nakagawa, Simon and Wood [5, 9, 10, 11, 12]. Related earlier work of the author on hyperbolic groups, hypercomplex structures and toric geometry appears in [2, 3, 4].

Declarations

Data availability. All data needed to verify the results are contained in the paper and its appendix. The LaTeX source, the figure script and a script that recomputes the resultants of the appendix are archived at doi:10.5281/zenodo.24126388.

Funding. The author received no funding for this work.

Conflict of interest. The author declares no conflict of interest.

Ethics statement. This work involves no human or animal subjects.

Appendix A. Base forms

A form $h_0 x^k + h_1 x^{k-1} y + \cdots + h_k y^k$ is written as its coefficient vector $[h_0, \ldots, h_k]$. The last two columns give the resultants with q_0 and q_1 of Table 1.

d	form	coefficients	$\mathrm{Res}(q_0, \cdot)$	$\mathrm{Res}(q_1, \cdot)$
40	H_4	$[-213, 254, 1679, -1135, 67]$	-1	1
40	H_6	$[-1, -2, 11, 19, -9, 9, -7]$	-1	1
229	H_3	$[14, -33, -845, 2678]$	1	-1
401	H_5	$[526505, 10598024, 47546124, -52212324,$ $69923173, -93789359]$	-1	1

For $d = 577$ the form $H_7 = [h_0, \ldots, h_7]$ has $\mathrm{Res}(q_0, H_7) = -1$ and $\mathrm{Res}(q_1, H_7) = 1$, with

$$\begin{aligned}
h_0 &= 923365420800961134849136086966499 9,\\
h_1 &= -158785974126310386817980985635657596,\\
h_2 &= 351336977243533453505499571533318677,\\
h_3 &= 295684110654525425618811641878008 2341,\\
h_4 &= -535022898010655232998946727047265289,\\
h_5 &= 228203582350855930536981792812157305,\\
h_6 &= -625031829845431344394388625520027 51,\\
h_7 &= 202185608478454766021956515566341 69.
\end{aligned}$$

The polynomials of Theorem B are $P = (x^2 - 10y^2)(x^4 - 8x^2y^2 - 6xy^3 - y^4)$, $P' = (3x^2 + 8xy + 2y^2)(x^4 - 8x^2y^2 - 6xy^3 - y^4)$, $Q = (x^2 + xy - 57y^2)H_3$ and $Q' = (3x^2 - 5xy - 17y^2)H_3$, written as polynomials in $X = x/y$.

References

[1] M. Bhargava, J.-H. Evertse, K. Győry, L. Remete and A. A. Swaminathan, Hermite equivalence of polynomials, *Acta Arith.* **209** (2023), 17–58. doi:10.4064/aa211113-12-11.

[2] D. Bhattacharjee, Non-commutativity over canonical suspension η for genus $g \geq 1$ in hypercomplex structures for potential $\rho\phi$, *Asian Res. J. Math.* **18** (2022), no. 11, 332–341. doi:10.9734/arjom/2022/v18i11607.

[3] D. Bhattacharjee, Establishing equivariant class $[O]$ for hyperbolic groups, *Asian Res. J. Math.* **18** (2022), no. 11, 362–369. doi:10.9734/arjom/2022/v18i11615.

[4] D. Bhattacharjee, S. Singha Roy and P. Nandi, Hodge–toric geometry beyond CY20, *Int. J. Prof. Stud.* **21** (2026), no. 1. doi:10.37648/ijps.v21i01.019.

[5] B. J. Birch and J. R. Merriman, Finiteness theorems for binary forms with given discriminant, *Proc. London Math. Soc. (3)* **24** (1972), 385–394. doi:10.1112/plms/s3-24.3.385.

[6] D. A. Buell, *Binary Quadratic Forms: Classical Theory and Modern Computations*, Springer, New York, 1989. doi:10.1007/978-1-4612-4542-1.

[7] R. Hartshorne, *Algebraic Geometry*, Grad. Texts in Math. **52**, Springer, New York, 1977. doi:10.1007/978-1-4757-3849-0.

[8] S. Lang, *Algebra*, revised 3rd ed., Grad. Texts in Math. **211**, Springer, New York, 2002. doi:10.1007/978-1-4613-0041-0.

[9] J. Nakagawa, Binary forms and orders of algebraic number fields, *Invent. Math.* **97** (1989), no. 2, 219–235. doi:10.1007/BF01389040.

[10] D. Simon, The index of nonmonic polynomials, *Indag. Math. (N.S.)* **12** (2001), no. 4, 505–517. doi:10.1016/S0019-3577(01)80039-2.

[11] M. M. Wood, Rings and ideals parameterized by binary n-ic forms, *J. London Math. Soc. (2)* **83** (2011), no. 1, 208–231. doi:10.1112/jlms/jdq074.

[12] M. M. Wood, Parametrization of ideal classes in rings associated to binary forms, *J. Reine Angew. Math.* **689** (2014), 169–199. doi:10.1515/crelle-2012-0058.

Formerly, Electro-Gravitational Space Propulsion Laboratory (EGSPL), Bhubaneswar, Odisha 751030, India

Formerly, University of Calcutta (Heramba Chandra College), 23/49 Gariahat Road, Golpark, Kolkata 700029, West Bengal, India

Email address: d.bhattacharjee@erl-forschung.de

Email address: itsdeep@live.com

Email address: egspl@proton.me (GitHub-related contact; GitHub account creelie)

URL: https://orcid.org/0000-0003-0466-750X