

Modules over short graded Gorenstein rings

Courtney Gibbons
University of Nebraska–Lincoln

AMS Contributed Paper Session
January 11, 2013

Background

Let \mathbb{k} be an algebraically closed field. Every short standard graded Gorenstein \mathbb{k} -algebra R with multiplicity e is isomorphic to

$$\mathbb{k}[x_1, \dots, x_e] / \left(x_1^2 - x_\ell^2, x_i x_j \mid \begin{array}{l} 1 \leq i \leq j \leq e \\ 1 < \ell \leq e \end{array} \right)$$

where $\deg x_\ell = 1$.

Background

Let \mathbb{k} be an algebraically closed field. Every short standard graded Gorenstein \mathbb{k} -algebra R with multiplicity e is isomorphic to

$$\mathbb{k}[x_1, \dots, x_e] / \left(x_1^2 - x_\ell^2, x_i x_j \mid \begin{matrix} 1 \leq i \leq j \leq e \\ 1 < \ell \leq e \end{matrix} \right)$$

where $\deg x_\ell = 1$.

$$\begin{array}{ll} \text{Degree 0:} & \mathbb{k} \\ & \oplus \\ \text{Degree 1:} & \mathbb{k}x_1 \oplus \cdots \oplus \mathbb{k}x_e \\ & \oplus \\ \text{Degree 2:} & \mathbb{k}x_e^2 \end{array}$$

Modules over R

We consider only finitely generated, graded R -modules $M = \bigoplus_{j \in \mathbb{Z}} M_j$.

The Hilbert series of M is defined to be

$$\mathcal{H}_M(s) = \sum \dim_{\mathbb{k}} M_j s^j.$$

Example

For example, $\mathcal{H}_R(s) = 1 + es + s^2$.

$$\begin{array}{lcl} \text{Degree 0:} & & \mathbb{k} \\ & & \oplus \\ \text{Degree 1:} & \mathbb{k}x_1 \oplus \cdots \oplus & \mathbb{k}x_e \\ & & \oplus \\ \text{Degree 2:} & & \mathbb{k}x_e^2 \end{array}$$

Modules over R

We consider only finitely generated, graded R -modules $M = \bigoplus_{j \in \mathbb{Z}} M_j$.

The Hilbert series of M is defined to be

$$\mathcal{H}_M(s) = \sum \dim_{\mathbb{k}} M_j s^j.$$

Example

For example, $\mathcal{H}_R(s) = 1 + es + s^2$.

Degree 0:

\mathbb{k}

\oplus

Degree 1: $\mathbb{k}x_1 \oplus \cdots \oplus \mathbb{k}x_e$

\oplus

Degree 2:

$\mathbb{k}x_e^2$

Modules over R

We consider only finitely generated, graded R -modules $M = \bigoplus_{j \in \mathbb{Z}} M_j$.

The Hilbert series of M is defined to be

$$\mathcal{H}_M(s) = \sum \dim_{\mathbb{k}} M_j s^j.$$

Example

For example, $\mathcal{H}_R(s) = 1 + \textcolor{red}{es} + s^2$.

Degree 0: \mathbb{k}

\oplus

Degree 1: $\textcolor{red}{\mathbb{k}x_1} \oplus \cdots \oplus \textcolor{red}{\mathbb{k}x_e}$

\oplus

Degree 2: $\mathbb{k}x_e^2$

Modules over R

We consider only finitely generated, graded R -modules $M = \bigoplus_{j \in \mathbb{Z}} M_j$.

The Hilbert series of M is defined to be

$$\mathcal{H}_M(s) = \sum \dim_{\mathbb{k}} M_j s^j.$$

Example

For example, $\mathcal{H}_R(s) = 1 + es + s^2$.

$$\begin{array}{lcl} \text{Degree 0:} & & \mathbb{k} \\ & & \oplus \\ \text{Degree 1:} & \mathbb{k}x_1 \oplus \cdots \oplus & \mathbb{k}x_e \\ & & \oplus \\ \text{Degree 2:} & & \mathbb{k}x_e^2 \end{array}$$

Koszul R -modules

An R -module K is called Koszul provided there exist free modules F_i and matrices ∂_i of linear forms (and zeroes) such that the following sequence is exact:

$$0 \leftarrow K \leftarrow F_0 \xleftarrow{\partial_1} F_1 \xleftarrow{\partial_2} F_2 \xleftarrow{\partial_3} \dots$$

Koszul R -modules

An R -module K is called Koszul provided there exist free modules F_i and matrices ∂_i of linear forms (and zeroes) such that the following sequence is exact:

$$0 \leftarrow K \leftarrow F_0 \xleftarrow{\partial_1} F_1 \xleftarrow{\partial_2} F_2 \xleftarrow{\partial_3} \dots$$

Example

The R -module $\mathbb{k} \cong R/(x_1, \dots, x_e)$ is Koszul:

$$0 \leftarrow \mathbb{k} \leftarrow R \xleftarrow{(x_1, \dots, x_e)} R(-1)^e \xleftarrow{\begin{pmatrix} 0 & 0 & z & 0 & 0 & y & 0 & x \\ 0 & -z & -z & -z & y & -z & x & z \\ z & y & 0 & x & 0 & 0 & 0 & 0 \end{pmatrix}} \dots$$

What Hilbert series are possible for Koszul R -modules?

Theorem (G—, 2012)

Given a short graded Gorenstein ring R with multiplicity e , there exists a Koszul module with Hilbert series $p + qs$ if and only if the integers p and q satisfy

$$1 \leq p \quad \text{and} \quad 0 \leq \frac{q}{p} \leq \frac{e + \sqrt{e^2 - 4}}{2}.$$

What Hilbert series are possible for Koszul R -modules?

Theorem (G—, 2012)

Given a short graded Gorenstein ring R with multiplicity e , there exists a Koszul module with Hilbert series $p + qs$ if and only if the integers p and q satisfy

$$1 \leq p \quad \text{and} \quad 0 \leq \frac{q}{p} \leq \frac{e + \sqrt{e^2 - 4}}{2}.$$

Proof idea.

We use previously known results due to L. Avramov, S. Iyengar, and L. Şega for $0 \leq \frac{q}{p} \leq e - 1$.

What Hilbert series are possible for Koszul R -modules?

Theorem (G—, 2012)

Given a short graded Gorenstein ring R with multiplicity e , there exists a Koszul module with Hilbert series $p + qs$ if and only if the integers p and q satisfy

$$1 \leq p \quad \text{and} \quad 0 \leq \frac{q}{p} \leq \frac{e + \sqrt{e^2 - 4}}{2}.$$

Proof idea.

We use previously known results due to L. Avramov, S. Iyengar, and L. Şega for $0 \leq \frac{q}{p} \leq e - 1$. To do this, we construct special Koszul modules.



Special Koszul Modules

Definition

Given a free resolution of \mathbb{k} ,

$$0 \leftarrow \mathbb{k} \leftarrow R(0)^1 \leftarrow R(-1)^e \leftarrow R(-2)^{e^2-1} \leftarrow \dots]$$

Define the sequence $b_n = (b_0 = 1, b_1 = e, b_2 = e^2 - 1, \dots)$.

Special Koszul Modules

Definition

Given a free resolution of \mathbb{k} ,

$$0 \leftarrow \mathbb{k} \leftarrow R(0)^1 \leftarrow R(-1)^e \leftarrow R(-2)^{e^2-1} \leftarrow \dots]$$

Define the sequence $b_n = (b_0 = 1, b_1 = e, b_2 = e^2 - 1, \dots)$.

Special Koszul Modules

Definition

Given a free resolution of \mathbb{k} ,

$$0 \leftarrow \mathbb{k} \leftarrow R(0)^1 \leftarrow R(-1)^e \leftarrow R(-2)^{e^2-1} \leftarrow \cdots]$$

Define the sequence $b_n = (b_0 = 1, b_1 = e, b_2 = e^2 - 1, \dots)$.

Special Koszul Modules

Theorem (G—, 2012)

Let M be an indecomposable module. Fix a positive integer $m \leq e - 1$. The following statements are equivalent:

- 1 $M \cong \text{Syz}_{-n-1}(I)$ where I is an ideal minimally generated by m linear forms.
- 2 $\mathcal{H}_M(s) = (b_n - mb_{n-1}) + (b_{n+1} - mb_n)s$.

If one of the above holds, then M is Koszul.

Next steps

- Macaulay2 code

Next steps

- Macaulay2 code
- Betti diagrams of Koszul modules form a subcone of the cone of Betti diagrams; describe this cone
- Equivalently: describe the cone of Hilbert series of Koszul modules

Next steps

- Macaulay2 code
- Betti diagrams of Koszul modules form a subcone of the cone of Betti diagrams; describe this cone
- Equivalently: describe the cone of Hilbert series of Koszul modules
- Larger picture: cone of Betti diagrams for modules over short Gorenstein rings in general

Next steps

- Macaulay2 code
- Betti diagrams of Koszul modules form a subcone of the cone of Betti diagrams; describe this cone
- Equivalently: describe the cone of Hilbert series of Koszul modules
- Larger picture: cone of Betti diagrams for modules over short Gorenstein rings in general

Acknowledgments & References

This research has been supported by GAANN, NebraskaMATH, NSF grant #DMS-1103176, and the University of Nebraska–Lincoln. Thanks to my advisors Luchezar Avramov and Roger Wiegand for their guidance.

Selected References:

- [AIŞ08] L. L. Avramov, S. B. Iyengar, and L. M. Şega, *Free resolutions over short local rings*, J. Lond. Math. Soc. (2) **78** (2008), no. 2, 459–476, DOI 10.1112/jlms/jdn027. MR2439635 (2009h:13011)
- [AIŞ10] ———, *Short Koszul modules*, J. Commut. Algebra **2** (2010), no. 3, 249–279, DOI 10.1216/JCA-2010-2-3-249. MR2728144 (2012a:13025)
- [Sjö79] G. Sjödin, *The Poincaré series of modules over a local Gorenstein ring with $\mathfrak{m}^3 = 0$ (preprint)*, Matematiska institutionen **2** (1979).