

THE SEARCH FOR INDECOMPOSABLE MODULES

Courtney Gibbons

September 11, 2015

Hamilton College

DEFINITIONS AND STUFF

Let $R = \mathbb{k}[x, y, z]/(x^2 - y^2, x^2 - z^2, xy, xz, yz)$.

Let $R = \mathbb{k}[x, y, z]/(x^2 - y^2, x^2 - z^2, xy, xz, yz)$.

As a \mathbb{k} -vector space,

$$R \cong \langle \bar{x} \rangle \oplus \langle \bar{y} \rangle \oplus \langle \bar{z} \rangle \oplus \langle \overline{x^2} \rangle \oplus \langle \overline{1} \rangle$$

degree 2, “ R_2 ”
 degree 1, “ R_1 ”
 degree 0, “ R_0 ”

Let $R = \mathbb{k}[x, y, z]/(x^2 - y^2, x^2 - z^2, xy, xz, yz)$.

As a \mathbb{k} -vector space,

$$R \cong \begin{array}{c} \langle \overline{x^2} \rangle \\ \oplus \\ \langle \overline{x} \rangle \oplus \langle \overline{y} \rangle \oplus \langle \overline{z} \rangle \\ \oplus \\ \langle \overline{1} \rangle \end{array} \begin{array}{l} \text{degree 2, "R}_2\text{"} \\ \text{degree 1, "R}_1\text{"} \\ \text{degree 0, "R}_0\text{"} \end{array}$$

MODULES AND HILBERT FUNCTIONS

Let $R = \mathbb{k}[x, y, z]/(x^2 - y^2, x^2 - z^2, xy, xz, yz)$.

As a \mathbb{k} -vector space,

$$R \cong \begin{array}{c} \langle \overline{x^2} \rangle \\ \oplus \\ \langle \overline{x} \rangle \oplus \langle \overline{y} \rangle \oplus \langle \overline{z} \rangle \\ \oplus \\ \langle \overline{1} \rangle \end{array} \begin{array}{l} \text{degree 2, "R}_2\text{"} \\ \text{degree 1, "R}_1\text{"} \\ \text{degree 0, "R}_0\text{"} \end{array}$$

Let $R = \mathbb{k}[x, y, z]/(x^2 - y^2, x^2 - z^2, xy, xz, yz)$.

As a \mathbb{k} -vector space,

$$\begin{array}{rclcl}
 & & \langle \overline{x^2} \rangle & & \text{degree 2, "R}_2\text{"} \\
 & & \oplus & & \\
 R & \cong & \langle \overline{x} \rangle \oplus \langle \overline{y} \rangle \oplus \langle \overline{z} \rangle & & \text{degree 1, "R}_1\text{"} \\
 & & \oplus & & \\
 & & \langle \overline{1} \rangle & & \text{degree 0, "R}_0\text{"}
 \end{array}$$

The (finite) sequence $H_R = (1, 3, 1)$ is called the **Hilbert function** of R .

The R -module $M = R/(\bar{z})$ has Hilbert function $H_M = (1, 2, 0) = (1, 2)$:

$$M_2 = \langle \bar{x^2} \rangle = \langle \bar{z^2} \rangle = 0$$

$$M_1 = \langle \bar{x} \rangle \oplus \langle \bar{y} \rangle,$$

$$M_0 = \langle \bar{1} \rangle.$$

Fact 1. Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be a (graded) short exact sequence of R -modules. Then $H_N + H_L = H_M$.

Fact 1. Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be a (graded) short exact sequence of R -modules. Then $H_N + H_L = H_M$.

Consider

$$0 \rightarrow (\bar{z}) \rightarrow R \rightarrow R/(\bar{z}) \rightarrow 0.$$

We can calculate $H_{(\bar{z})}$ as follows:

$$\begin{aligned} H_{(\bar{z})} &= H_R - H_{R/(\bar{z})} \\ &= (1, 3, 1) - (1, 2, 0) \\ &= (0, 1, 1). \end{aligned}$$

I want to say $(0, 1, 1) = (1, 1)$.

Fact 1. Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be a (graded) short exact sequence of R -modules. Then $H_N + H_L = H_M$.

Consider

$$0 \rightarrow (\bar{z}) \rightarrow R \rightarrow R/(\bar{z}) \rightarrow 0.$$

We can calculate $H_{(\bar{z})}$ as follows:

$$\begin{aligned} H_{(\bar{z})} &= H_R - H_{R/(\bar{z})} \\ &= (1, 3, 1) - (1, 2, 0) \\ &= (0, 1, 1). \end{aligned}$$

I want to say $(0, 1, 1) = (1, 1)$.

Note that $R \oplus (\bar{z})$ has Hilbert function $(1, 4, 2)$.

As an R -module, R itself is **indecomposable**.

That is, whenever $R = M \oplus N$, it follows that implies $M = R$, $N = 0$.

As an R -module, R itself is **indecomposable**.

That is, whenever $R = M \oplus N$, it follows that implies $M = R$, $N = 0$.

Let I be an ideal generated by homogeneous polynomials. The R -modules $M = R/I$ and I are also indecomposable.

As an R -module, R itself is **indecomposable**.

That is, whenever $R = M \oplus N$, it follows that implies $M = R$, $N = 0$.

Let I be an ideal generated by homogeneous polynomials. The R -modules $M = R/I$ and I are also indecomposable.

Fact 2. Suppose

$$0 \rightarrow I \rightarrow R \rightarrow F^1 \rightarrow F^2 \rightarrow \cdots \rightarrow F^n \rightarrow M \rightarrow 0$$

is an exact sequence of R -modules, where F^i is a free R -module for each i . Then M is called the n -th **cosyzygy** of I , $\text{Cosyz}^n(I)$, and M is indecomposable.

BACKGROUND

Recall, $R = \mathbb{k}[x, y, z]/(x^2 - y^2, x^2 - z^2, xy, xz, yz)$.

More generally, $R = \mathbb{k}[x_1, \dots, x_e]/(x_i^2 - x_j^2, x_i x_j \mid 1 \leq i < j \leq e)$ is called a **Short Gorenstein Ring of embedding dimension e** .

SETTING UP THE QUESTION

More generally, $R = \mathbb{k}[x_1, \dots, x_e]/(x_i^2 - x_j^2, x_i x_j \mid 1 \leq i < j \leq e)$ is called a **Short Gorenstein Ring of embedding dimension e** .

Prop (Avramov-Iyengar-Şega). Let R be a S.G.R. with $e \geq 3$. If M is a Koszul R -module, then the Hilbert function $H_M = (p, q)$ where

$$1 \leq p \quad \text{and} \quad 0 \leq q < p \frac{e + \sqrt{e^2 - 4}}{2}.$$

SETTING UP THE QUESTION

More generally, $R = \mathbb{k}[x_1, \dots, x_e]/(x_i^2 - x_j^2, x_i x_j \mid 1 \leq i < j \leq e)$ is called a **Short Gorenstein Ring of embedding dimension e** .

Prop (Avramov-Iyengar-Şega). Let R be a S.G.R. with $e \geq 3$. If M is a Koszul R -module, then the Hilbert function $H_M = (p, q)$ where


$$1 \leq p \quad \text{and} \quad 0 \leq q < p \frac{e + \sqrt{e^2 - 4}}{2}.$$

Furthermore, given

$$1 \leq p \quad \text{and} \quad 0 \leq q \leq p(e - 1),$$

there exists a Koszul R -module M where $H_M = (p, q)$.

BUT HERE'S HOW TO REALLY THINK ABOUT IT



When $e = 3$,

$$\begin{aligned}\varphi &= \frac{e + \sqrt{e^2 - 4}}{2} \\ &= \frac{3 + \sqrt{5}}{2} \\ &= 1 + \frac{1 + \sqrt{5}}{2}\end{aligned}$$

When $e = 3$,

$$\begin{aligned}\varphi &= \frac{e + \sqrt{e^2 - 4}}{2} \\&= \frac{3 + \sqrt{5}}{2} \\&= 1 + \frac{1 + \sqrt{5}}{2} \\&= 1 + 1 + \frac{1}{1 + \frac{1}{1 + \dots}}.\end{aligned}$$

When $e = 3$,

$$\begin{aligned}\varphi &= \frac{e + \sqrt{e^2 - 4}}{2} \\ &= \frac{3 + \sqrt{5}}{2} \\ &= 1 + \frac{1 + \sqrt{5}}{2} \\ &= 1 + 1 + \frac{1}{1 + \frac{1}{1 + \dots}}.\end{aligned}$$

Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, 21, 34, ...

Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, 21, 34, ...

Using the theory of continued fractions and convergents, we find that the sequence $(\frac{2}{1}, \frac{5}{2}, \frac{13}{5}, \frac{34}{13}, \dots)$ converges **very quickly** to φ from below.

CONTINUED FRACTIONS AND CONVERGENCE


Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, 21, 34, ...

Using the theory of continued fractions and convergents, we find that the sequence $(\frac{2}{1}, \frac{5}{2}, \frac{13}{5}, \frac{34}{13}, \dots)$ converges **very quickly** to φ from below.


Geometrically, that means it's enough to find modules with the Hilbert functions

$$(1,2), (2,5), (5,13), (13,34), \dots$$


ONCE MORE IN PICTURES




ONCE MORE IN PICTURES




ONCE MORE IN PICTURES




ONCE MORE IN PICTURES




ONCE MORE IN PICTURES




ONCE MORE IN PICTURES



ONCE MORE IN PICTURES



ONCE MORE IN PICTURES



Recap:

Fact 1. Hilbert function - additive on S.E.S.

Fact 2. There's an S.E.S.

$$0 \rightarrow \text{Cosyz}_{n-1}(I) \rightarrow F^n \rightarrow \text{Cosyz}_n(I) \rightarrow 0$$

(and $\text{Cosyz}_n(I)$ is indecomposable).

Consider $I = (\bar{z})$. We know $H_{\text{Cosyz}_1(I)} = H_{R/I} = (1, 2)$.

Then the Hilbert function of $\text{Cosyz}_2(I)$ can be found from the exact sequence

$$0 \rightarrow \text{Cosyz}_1(I) \rightarrow R^2 \rightarrow \text{Cosyz}_2(I) \rightarrow 0.$$

Consider $I = (\bar{z})$. We know $H_{\text{Cosyz}_1(I)} = H_{R/I} = (1, 2)$.

Then the Hilbert function of $\text{Cosyz}_2(I)$ can be found from the exact sequence

$$0 \rightarrow \text{Cosyz}_1(I) \rightarrow R^2 \rightarrow \text{Cosyz}_2(I) \rightarrow 0.$$

Now,

$$H_{\text{Cosyz}_2(I)} = (2, 6, 2) - (0, 1, 2) = (2, 5, 0) = (2, 5).$$

Consider $I = (\bar{z})$. We know $H_{\text{Cosyz}_1(I)} = H_{R/I} = (1, 2)$.

Then the Hilbert function of $\text{Cosyz}_2(I)$ can be found from the exact sequence

$$0 \rightarrow \text{Cosyz}_1(I) \rightarrow R^2 \rightarrow \text{Cosyz}_2(I) \rightarrow 0.$$

Now,

$$H_{\text{Cosyz}_2(I)} = (2, 6, 2) - (0, 1, 2) = (2, 5, 0) = (2, 5).$$

Repeating this trick,

$$H_{\text{Cosyz}_3(I)} = (5, 15, 5) - (0, 2, 5) = (5, 13).$$

THEOREM (AVRAMOV-GIBBONS-WIEGAND)

Let R be a S.G.R. with $e \geq 3$. Then there exists a Koszul R -module M with Hilbert function (p, q) if and only if

$$1 \leq p \quad \text{and} \quad 0 \leq q \leq p \frac{e + \sqrt{e^2 - 4}}{2}.$$

QUESTIONS?

Definition. An R -module M is said to be **Koszul** provided M is generated in degree 0, has no nonzero free summand, and has a linear free resolution.

In our setting, if $H_M = (p, q)$, then:

$$\beta(M) = \begin{bmatrix} & & \vdots & & \\ 0 & 0 & \cdots & 0 & \cdots \\ p & ep - q & \cdots & (e\beta_{n-1} - \beta_{n-2}) & \cdots \\ 0 & 0 & \cdots & 0 & \cdots \\ & & \vdots & & \end{bmatrix}.$$

THOSE OTHER ORDERED PAIRS? $(1,3)$, $(3,8)$, ...

The only indecomposable non-Koszul R -modules have the form $\text{Cosyz}_n(\overline{x_1}, \dots, \overline{x_e})$, and their Hilbert functions use the odd Fibonacci numbers.

- [1] Luchezar L. Avramov, Srikanth B. Iyengar, and Liana M. Şega, Free resolutions over short local rings, *J. Lond. Math. Soc.* (2) **78** (2008), no. 2, 459–476, DOI 10.1112/jlms/jdn027. MR2439635 (2009h:13011)
- [2] ———, Short Koszul modules, *J. Commut. Algebra* **2** (2010), no. 3, 249–279, DOI 10.1216/JCA-2010-2-3-249. MR2728144 (2012a:13025)
- [3] Courtney Gibbons, *Decompositions of Betti diagrams*, ProQuest LLC, Ann Arbor, MI, 2013. Thesis (Ph.D.)—The University of Nebraska - Lincoln. MR3153511
- [4] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 6th ed., Oxford University Press, Oxford, 2008. Revised by D. R. Heath-Brown and J. H. Silverman; With a foreword by Andrew Wiles. MR2445243 (2009i:11001)