

APRIORA Plugin: From Flow Estimation to Pharmaceutical Risk Assessment in QGIS

Cristiano Guidi¹, Jens Tränckner², and Philip Marzahn¹

¹ Geodesy and Geoinformatics, University of Rostock, Justus-von-Liebig-Weg 6, 18059 Rostock, Germany ² Water Management, University of Rostock, Satower Straße 48, 18059 Rostock, Germany

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

APRIORA is an open-source QGIS plugin developed for spatial modelling and risk assessment of active pharmaceutical ingredients (APIs) in river basins. Designed to support regional authorities under the updated EU Urban Wastewater Treatment Directive (UWWTD), APRIORA offers a user-friendly, accessible workflow within the QGIS environment. The plugin is organized into two primary modules containing a total of five tools. The Hydro-Module pre-processes river networks and estimates key hydrological data in data-scarce catchments, while the API Emission module combines pharmaceutical statistics, connected inhabitants at the treatment plant and river hydrology to calculate predicted environmental concentrations (PECs) and risk quotients (RQs) for three different types of risk: environmental, antimicrobial resistance and human health. APRIORA allows users to customize emission parameters, plot comprehensive risk maps and simulate and compare the effectiveness of mitigation scenarios (e.g., treatment upgrades or discharge relocation). Tested in collaboration with regional stakeholders and applied across five Baltic Sea country pilots, APRIORA provides an accessible decision-support tool for water resource management.

Statement of need

Pharmaceuticals are essential in modern healthcare. However, they can harm environmental components as they enter surface waters primarily through wastewater treatment plant (WWTP) effluents (Kosek et al., 2020). Evidence of the detrimental effects of pharmaceuticals on non-target organisms at low concentrations (Daughton & Ternes, 1999; Hoeger et al., 2005; Williams et al., 2010), as well as their persistent presence in aquatic systems (Tränckner & Koegst, 2010), has led to an intensified focus on API mitigation in European water management policy. Particularly, the updated EU Urban Wastewater Treatment Directive (UWWTD) mandates quaternary treatment upgrades for large WWTPs (≥ 150.000 population equivalents (PE)) and requires risk-based prioritization for upgrading mid-sized facilities (10.000 to 150.000 PE) by 2030 based on discharge dilution and receiving water sensitivity (European Parliament and Council of the European Union, 2024).

While the UWWTD focuses primarily on medium and large infrastructure, small-sized WWTPs (< 10.000 PE) are frequently overlooked. However, small facilities discharging into tributaries or low-flow streams often create pollution hotspots due to critical mixing ratios between the emission load and receiving flow volume (Büttner et al., 2022; Di Sabatino et al., 2024). At a catchment level, these distributed upstream discharges create localized ecotoxicological risks along individual river sections and substantial cumulative pollution loads reaching sensitive marine basins downstream, such as the Baltic Sea, which is one of the most polluted marine water bodies in the world (Reusch et al., 2018). Risk characterization usually involves calculating risk quotients (RQs) by comparing predicted environmental concentrations (PECs) with predicted no-effect concentrations (PNECs) (Kisielius et al., 2023) or legally more binding

Environmental Quality Standards (EQSs), as updated under Directive 2026/80 (European Parliament and Council of the European Union, 2026).

To deal with the task of an informed and transparent risk assessment, environmental protection agencies (EPAs), regional and local authorities and WWTP operators require a unified, accessible methodology to model surface water API concentrations, assess spatially distributed risk and prioritize mitigation strategies within river catchments.

Although there are established exposure and emission models, such as stand-alone packages (MoRE (Fuchs et al., 2017), ePiE (Oldenkamp et al., 2018)), ArcGIS extensions (GREAT-ER (Kehrein et al., 2015)) and complex research frameworks (STREAM-EU (Lindim et al., 2016)), their practical adoption by regional decision-makers is affected by commercial software restrictions, high technical barriers to entry, unavailable input data, non-adaptable spatial domains and a lack of direct scenario testing capabilities. APRIORA fills this critical gap as a free and open-source QGIS plugin with low input data demands, allowing end-users to independently map RQs and evaluate interactive mitigation scenarios such as WWTP upgrades, effluent redirection, or discharge relocation.

State of the field

To evaluate the water quality of river networks or catchments, several tools already exist, such as those cited previously: ePiE, STREAM-EU, GREAT-ER and MoRE offer distinct modelling capabilities, yet each presents operational challenges for regional applications. ePiE simulates first-order degradation kinetics and phase transport across European basins, but it relies on EU-wide reporting databases, automatically excluding small WWTPs (<2000 PE), leading to systematic exposure underestimation in upstream tributaries. Furthermore, the model is distributed as an R package, compromising its user-friendliness for authorities in charge who might not be skilled in R or basic coding. STREAM-EU provides dynamic, fugacity-based multi-media fate simulations, yet requires extensive physico-chemical and environmental parameterization that turns routine regional screening into a computationally heavy task. GREAT-ER computes reach-specific probabilistic concentration distributions via Monte Carlo simulations; however, it requires a commercial license for ArcGIS software and laborious data preparation for the catchment under investigation. Meanwhile, MoRE provides comprehensive multi-tier regionalized emission calculations, but it is a closed-distribution platform that requires direct coordination with its maintainers. Additionally, it aggregates pollutant loads over large spatial analytical units (>100 km²) rather than providing continuous, segment-by-segment river risk profiles.

A fundamental requirement across all these tools is accurate hydrological flow data to calculate in-river substance dilution and concentrations. Existing frameworks depend entirely on pre-defined external hydrological models or global databases (such as FLO1K for ePiE, LARSIM for MoRE or SWAT for GREAT-ER) limiting their adaptability in data-scarce areas, particularly in small, ungauged catchments where smaller WWTPs frequently discharge. Refactoring one of these existing models into a free and open-source QGIS plugin was technically unfeasible, making a standalone build the best option.

APRIORA provides its own hydrological module to estimate annual mean flow (MQ_a) and annual mean low flow (MLQ_a), while still keeping the possibility for users to import external hydrological datasets. Additionally, it provides an environment where users can freely customize input parameters including local API consumption, WWTP removal efficiencies, PNEC thresholds and monitoring data. Beyond standard environmental risk assessment (ERA), its framework also supports human health (HHRA) and antimicrobial resistance (AMR-RA) evaluations. It can also simulate environmental propagation for other micropollutants such as PFAS or microplastics whenever emission or consumption data are available. Finally, different mitigation scenario can be tested to support authorities in the decision-making process.

Software design

The APRIORA plugin is an open-source QGIS extension designed to make pharmaceutical risk modelling easy to use for regional decision-makers and environmental agencies working under the UWWTD. Its main design goal is to provide a low technical barrier to entry by using mostly publicly available data and fitting directly into standard QGIS workflows. The plugin is organized into two main modules containing five total tools:

- Hydro-Module: preprocesses stream vector lines (1 - *Fix River Network*) and estimates MQ_a and MLQ_a in data-scarce catchments (2 - *Flow Estimation*)
- API Emission: calculates pollutant loads from WWTP (3 - *Emission Loads*), routes these loads downstream (4 - *Accumulation*) and perform risk assessment (5 - *Risk Assessment*)

A scheme of the plugin is illustrated in Figure 1.




Figure 1: APRIORA plugin scheme.

To make the tool useful across different regions, several practical trade-offs were built into the software design. First, the Hydro-Module works with standard European databases like Copernicus DEM, ERA5 precipitation and CORINE Land Cover, while still allowing users to use local data. Because precipitation data formats can differ between sources (such as ERA5 using single multi-year NetCDF files and local agencies like the German Weather Service using annual rasters), the interface includes a flexible toggle switch so users do not have to manually reformat their files. Additionally, spatial data structures can vary country by country. While downstream routing usually relies on line networks, some databases (like Finland's Vemala model (Huttunen et al., 2016)) store hydrological data as polygons. Because of this reason, 4 - *Accumulation* and 5 - *Risk Assessment* were designed to accept both linear river networks and polygon layers, letting users choose whichever spatial format fits best.

Finally, the plugin is characterised by high user customization. APRIORA comes pre-loaded with consumption data, removal rates and PNEC thresholds for ten common APIs across five Baltic Sea region countries. Users can easily update these parameters or even adjust them for the individual WWTPs within the study catchment. To simplify the collaboration between different

118 users, these customized setups can be exported as simple tables and shared with colleagues,
119 guaranteeing that everyone can reproduce the exact same model run without typing values in
120 by hand.

121 Research impact statement

122 Developed as part of the EU Interreg-sponsored [APRIORA project](#), the plugin was tested and
123 refined based on direct feedback from project partners and environmental authorities. The
124 tool was first introduced to the public in November 2025 at the EMPEREST final conference
125 in Berlin, where around 20 regional stakeholders took part in a workshop on using the API
126 Emission set of tools ([article](#)). A similar workshop was held in June 2026 in Rostock with
127 about 15 regional stakeholders from the Mecklenburg-Western Pomerania region. In July 2026,
128 the plugin was taught during the APRIORA One Health Summer School in Gdansk, where 50
129 participants were trained on using the tool ([article](#)).

130 The research impact of APRIORA is shown through its application across five river catchments
131 in Germany, Sweden, Finland, Latvia and Poland. Consortium partners used the plugin to
132 model pharmaceutical risks and test local mitigation options under different conditions. These
133 case studies formed the basis for pilot reports which can be found in the APRIORA project
134 website ([Latvia](#), [Sweden](#), [Finland](#), [Poland](#) and [Germany](#)). Additionally, a scientific paper about
135 the development and application of the plugin is under writing.

136 AI usage disclosure

137 Generative AI tools were used during the software development and paper writing process.
138 AI was used to help write parts of the Python code, troubleshoot errors, and proofread the
139 manuscript text. All AI-generated code was manually reviewed, tested, and verified by the
140 developer through catchment modeling runs before the software was published. All the text
141 was reviewed and edited by the authors to make sure it was factually correct.

142 Acknowledgments

143 The project leading to this application is funded by the Interreg Baltic Sea Region Programme.
144 Co-funded by the European Union (ERDF), this #MadeWithInterreg project helps to remove
145 pollutants from our waters. The developer would also like to thank all the project partners,
146 regional stakeholders and users who contributed to the development of the plugin by providing
147 continuous feedback and ideas on how to improve it.

148 References

- 149 Büttner, O., Jawitz, J. W., Birk, S., & Borchardt, D. (2022). Why wastewater treatment
150 fails to protect stream ecosystems in Europe. *Water Research*, 217, 118382. <https://doi.org/10.1016/j.watres.2022.118382>
151
- 152 Daughton, C. G., & Ternes, T. A. (1999). Pharmaceuticals and personal care products in the
153 environment: Agents of subtle change? *Environmental Health Perspectives*, 107, 907–938.
154 <https://doi.org/10.1289/ehp.99107s6907>
- 155 Di Sabatino, A., Damiani, G., Ercolino, G., Rossi, F., & Ruggieri, L. (2024). Responses
156 of freshwater invertebrates to Imhoff tank sewage effluents: A preliminary study in four
157 watercourses with different ecological status (Abruzzo, Central Italy). *Sustainability*, 16(6),
158 2452. <https://doi.org/10.3390/su16062452>

- European Parliament and Council of the European Union. (2024). Directive (EU) 2024/3019 concerning urban wastewater treatment. In *Official Journal of the European Union* (L 2024/3019). <https://eur-lex.europa.eu/eli/dir/2024/3019/oj>
- European Parliament and Council of the European Union. (2026). Directive (EU) 2026/805 of the European Parliament and of the Council of 30 March 2026 amending Directive 2000/60/EC, Directive 2006/118/EC and Directive 2008/105/EC as regards environmental quality standards in the field of water policy. In *Official Journal of the European Union* (L 2026/805). <http://data.europa.eu/eli/dir/2026/805/oj>
- Fuchs, S., Kaiser, M., Kiemle, L., Kittlaus, S., Rothvoß, S., Toshovski, S., Wagner, A., Wander, R., Weber, T., & Ziegler, S. (2017). Modeling of regionalized emissions (MoRE) into water bodies: An open-source river basin management system. *Water*, 9(4), 239. <https://doi.org/10.3390/w9040239>
- Hoeger, B., Köllner, B., Dietrich, D. R., & Hitzfeld, B. (2005). Water-borne diclofenac affects kidney and gill integrity and selected immune parameters in brown trout (*salmo trutta f. Fario*). *Aquatic Toxicology*, 75(1), 53–64. <https://doi.org/10.1016/j.aquatox.2005.07.006>
- Huttunen, I., Huttunen, M., Piirainen, V., Korppoo, M., Lepistö, A., Räike, A., Tattari, S., & Vehviläinen, B. (2016). A national-scale nutrient loading model for Finnish watersheds—VEMALA. *Environmental Modeling & Assessment*, 21(1), 83–109. <https://doi.org/10.1007/s10666-015-9470-6>
- Kehrein, N., Berlekamp, J., & Klasmeier, J. (2015). Modeling the fate of down-the-drain chemicals in whole watersheds: New version of the GREAT-ER software. *Environmental Modelling & Software*, 64, 1–8. <https://doi.org/10.1016/j.envsoft.2014.10.018>
- Kisieliu, V., Kharel, S., Skaarup, J., Lauritzen, B. S., Lukas, M., Bogusz, A., Szumska, M., & Bester, K. (2023). Process design for removal of pharmaceuticals in wastewater treatment plants based on predicted no effect concentration (PNEC). *Chemical Engineering Journal*, 476, 146644. <https://doi.org/10.1016/j.cej.2023.146644>
- Kosek, K., Luczkiewicz, A., Fudala-Książek, S., Jankowska, K., Szopińska, M., Svahn, O., Tränckner, J., Kaiser, A., Langas, V., & Björklund, E. (2020). Implementation of advanced micropollutants removal technologies in wastewater treatment plants (WWTPs) - Examples and challenges based on selected EU countries. *Environmental Science & Policy*, 112, 213–226. <https://doi.org/10.1016/j.envsci.2020.06.011>
- Lindim, C., Van Gils, J., & Cousins, I. T. (2016). A large-scale model for simulating the fate & transport of organic contaminants in river basins. *Chemosphere*, 144, 803–810. <https://doi.org/10.1016/j.chemosphere.2015.09.051>
- Oldenkamp, R., Hoeks, S., Čengić, M., Barbarossa, V., Burns, E. E., Boxall, A. B. A., & Ragas, A. M. J. (2018). A high-resolution spatial model to predict exposure to pharmaceuticals in European surface waters: ePiE. *Environmental Science & Technology*, 52(21), 12494–12503. <https://doi.org/10.1021/acs.est.8b03862>
- Reusch, T. B. H., Dierking, J., Andersson, H. C., Bonsdorff, E., Carstensen, J., Casini, M., Czajkowski, M., Hasler, B., Hinsby, K., Hyytiäinen, K., Johannesson, K., Jomaa, S., Jormalainen, V., Kuosa, H., Kurland, S., Laikre, L., MacKenzie, B. R., Margonski, P., Melzner, F., ... Zandersen, M. (2018). The Baltic Sea as a time machine for the future coastal ocean. *Science Advances*, 4(5), eaar8195. <https://doi.org/10.1126/sciadv.aar8195>
- Tränckner, J., & Koegst, T. (2010). Demographic effects on domestic pharmaceutical emissions in Germany. *Novatech 2010 - 7th International Conference on Sustainable Techniques and Strategies for Urban Water Management*, 1–8. <https://hal.science/hal-03296311>
- Williams, R. J., Keller, V. D. J., Johnson, A. C., Young, A. R., Holmes, M. G. R., Wells, C., Gross-Sorokin, M., & Benstead, R. (2010). A national risk assessment for intersex in fish arising from steroid estrogens. *Environmental Toxicology and Chemistry*, 28(1), 220–230.

DRAFT