

Processing of recombinant spider silk proteins into tailor-made materials for biomaterials applications

Kristin Schacht¹ and Thomas Scheibel^{1,2,3,4,5}

Spider silk has extraordinary mechanical properties, is biocompatible and biodegradable, and therefore an ideal material for biomedical applications. However, a drawback for any application is the inhomogeneity of spider silk, as seen for other natural materials, as well as the low availability due to the cannibalism of most spiders. Recently, developed recombinant spider silk proteins ensure constant material properties, as well as scalable production, and further the processing into morphologies other than fibres. Biotechnology enables genetic modification, broadening the range of applications, such as implant coatings, scaffolds for tissue engineering, wound dressing devices as well as drug delivery systems.

Addresses

- ¹ Lehrstuhl Biomaterialien, Fakultät für Ingenieurwissenschaften, Universität Bayreuth, Universitätsstraße 30, Bayreuth 95440, Germany
² Bayreuther Zentrum für Kolloide und Grenzflächen (BZKG), Universität Bayreuth, Universitätsstraße 30, Bayreuth 95440, Germany
³ Institut für Bio-Makromoleküle (bio-mac), Universität Bayreuth, Universitätsstraße 30, Bayreuth 95440, Germany
⁴ Bayreuther Zentrum für Molekulare Biowissenschaften (BZMB), Universität Bayreuth, Universitätsstraße 30, Bayreuth 95440, Germany
⁵ Bayreuther Materialzentrum (BayMAT), Universität Bayreuth, Universitätsstraße 30, Bayreuth 95440, Germany

Corresponding author: Scheibel, Thomas (Thomas.Scheibel@uni-bayreuth.de)

Current Opinion in Biotechnology 2014, 29:62–69

This review comes from a themed issue on **Cell and pathway engineering**

Edited by **Tina Lütke-Eversloh** and **Keith EJ Tyo**

Available online XXX

0958-1669/\$ – see front matter, © 2014 Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.copbio.2014.02.015>

Introduction

Spider silk fibres fascinate scientists especially due to their extraordinary mechanical properties [1]. The combination of strength and elasticity provides a toughness no other natural or synthetic fibre can achieve [2]. Additionally, spider silk is biocompatible, biodegradable and shows hypoallergenic properties suitable for biomedical applications [3,4].


Importantly, spider silk reflects an entire class of materials with different properties, since spiders can produce

several types of silk (for an overview see Heidebrecht and Scheibel [5]). The best characterised spider silk is the *Major Ampullate* (MA)/Dragline silk, constituting the outer frame of orb webs, serving also as a lifeline for the spider and which will be exclusively discussed herein [6]. Two classes of *Major Ampullate* spidroins have been identified in dragline fibres, called MaSp1 and MaSp2, which differ in proline content and hydrophobicity [7]. All *Major Ampullate* spidroins consist of a highly repetitive core domain, flanked by non-repetitive termini [8]. In the core domain distinct amino acid motifs (glycine-rich repeats and polyalanine blocks) enable secondary structures (random coil/helical and β -sheet) accounting for the mechanical properties of the fibre [8–11]. The terminal domains play an important role during storage of spidroins as a spinning dope in the gland and during the initiation of fibre assembly in the spinning duct [12–15].

For centuries, spider's webs have been successfully used to stop bleeding and to promote wound healing [16]. Recently, spider silk has been used as an artificial support for nerve regeneration [17,18]. Defects of peripheral nerves can be repaired by a composite nerve graft made of acellularized veins, spider silk fibres and Schwann cells (SC) mixed with matrigel (a solubilized tissue basement membrane matrix rich in extracellular matrix proteins). In adult sheep, spider silk enhanced Schwann cell migration, axonal-regrowth and remyelination including electrophysiological recovery in a 6.0 cm tibial nerve defect [19]. Further, native spider silk fibres were tested as a braided microsurgical suture to substitute conventional materials in microsurgery and neurosurgery [20,21]. It was shown that the mechanical properties of braided spider silk sutures were superior to those of nylon, the current clinical gold-standard [21]. However, one major drawback of natural spider silk sutures is the inhomogeneity of the fibres, as seen with other natural materials, since differences in silk properties occur between individual spiders and even within single individuals upon environmental changes. Another drawback is the low availability of natural material due to problems in farming based on the cannibalistic behaviour of spiders [22,23].

Biotechnological production of spider silk proteins, as well as the development of silk processing techniques enabled the supply of engineered silk materials for biomedical applications, such as implant coatings, drug delivery systems or scaffolds for tissue engineering, which are reviewed herein.

Figure 1



Genetic engineering to achieve functional spider silk proteins.

Recombinant production of engineered spider silk proteins

In the last decades, several prokaryotic and eukaryotic hosts have been tested concerning recombinant production of spider silk proteins, as recently summarized in Heidebrecht and Scheibel [5[•]].

The benefits of recombinant spider silk proteins (RSSP) are the homogeneity of the starting material as well as the controllable processability into different morphologies, like films, hydrogels, particles or non-woven meshes for various applications [24–29]. Further, biotechnology enables genetic engineering to directly incorporate functional groups into the RSSPs (Figure 1) [24,25,30[•]].

The simplest genetic modification is the incorporation of individual amino acid residues with chemically specific side chains, like cysteine residues comprising thiol groups. A cysteine variant of the RSSP eADF4(C16) (based on the dragline silk protein ADF4 of *A. diadematus*) allowed the covalent coupling of peptides, enzymes or particles before and after silk processing into materials, demonstrating its potential for a broad range of applications [25,30[•]].

Engineered spider silk proteins comprising functional peptide sequences

For biomaterials applications, specific interactions between cells and the surface of a material are essential. Spider silk proteins can be exemplarily modified with cell adhesive peptides to improve cell binding, such as the integrin-binding motif RGD (Arg-Gly-Asp) (Figure 1) [28,30[•],31,32]. Films made of eADF4(C16)-RGD showed a significantly improved attachment and proliferation of fibroblasts (BALB/3T3) in comparison to unmodified eADF4(C16) films [30[•]]. Another RSSP, 4RepCT, genetically functionalized with RGD or the cell binding peptides IKVAV, naturally found in the laminin α 1 chain, or YIGSR, present in the β 1 chain, were processed into fibres, foams and films [33,34]. The adhesion of all tested cell types (fibroblasts, keratinocytes, endothelial and Schwann cells) was significantly improved on RGD-modified in comparison to unmodified 4RepCT films. While only Schwann cells adhered better on matrices comprising the IKVAV-motif, no clear effect of YIGSR could be detected on any of the selected cell types [35].

Functionalizing spider silk proteins or silk hybrids with antimicrobial peptides could be a new approach to

achieve multifunctional biomaterials in order to suppress infections in combination with for example, supporting cell growth. Recently, the human antimicrobial peptides human neutrophil defensin 2 (HNP-2), human neutrophil defensin 4 (HNP-4) and hepcidin were fused to an RSSP 6mer, based on the sequence of MaSp1 from *Nephila clavipes* (*N. clavipes*) (Figure 1). The silk hybrids were processed into films showing antimicrobial activity against Gram negative *E. coli* and Gram positive *S. aureus*. In addition, *in vitro* cell culture studies (cytotoxicity/proliferation) with a human osteosarcoma cell line (SaOs-2) demonstrated the compatibility of these films with mammalian cells [36]. *In vivo* studies in mice showed that the silk-hepcidin protein was highly biocompatible, causing a mild to low inflammatory reaction [37]. In a similar approach, a hybrid between spider silk and a silver binding peptide was produced, which could be processed into films. These films nucleated silver ions from a solution of silver nitrate and inhibited afterwards microbial growth of Gram positive as well as Gram negative bacteria *in vitro* [38].

Silica binding peptides were fused to a consensus sequence of *N. clavipes* MaSp1 introducing silicifying properties [39]. Films thereof promoted osteoblast development with the upregulation of key markers associated with bone formation [40,41].

Other peptides, such as cell penetrating and cell membrane destabilizing peptides (CPPs) are useful candidates, for example, for promoting gene transfer. In this context, recombinant spider silk–polylysine fusions with a ppTG1 peptide, a lysine-rich cell membrane destabilizing peptide to bind plasmid DNA (pDNA), were designed as a highly efficient gene carrier. The newly generated fusion proteins showed useful transfection efficiency, comparable to the transfection reagent Lipofectamine 2000, and at the same time the possibility to control gene release by controlling enzymatic degradation rates of the complexes [42].

Engineering of chimeric spider silk proteins

Chimeric proteins consisting of the RSSP 6mer and the bone sialoprotein (BSP) were processed into films as a support for bone regeneration. Compared to the control (6mer silk films), the Young's modulus of films made of the chimeric silk protein was significantly increased. Furthermore, the chimeric protein retained the ability to form supramolecular aggregates, and in the presence of Ca^{2+} ions these aggregates generated networks [43–45]. *In vitro* studies have demonstrated that human mesenchymal stem cells proliferated and differentiated into the osteogenic lineage on 6mer + BSP films. The presence of cell binding domains in BSP (such as RGD) may be responsible for the cell response when compared to silk alone. For the plain 6mer films, osteoblast-like morphology was not as evident,

and a reduction on viability/proliferation was observed after 3, 7 and 14 days in osteogenic medium [46,47].


Chimeric proteins based on *N. clavipes* MaSp1, and the dentin matrix protein 1 (CDMP1), which provides controlled nucleation and crystallization of hydroxyapatite, targeted self-assembled silk structures with controlled hydroxyapatite (HA) mineralization. Upon processing into films, mineralization was initiated using simulated body fluids (SBF) [48]. Mineralized silk films mediated and promoted bone regeneration around wounds and promoted osteoblastic differentiation in a three dimensional scaffold [49–52].

Processing of engineered spider silk proteins for biomedical applications

Spider silk proteins can be processed into coatings which can be used to improve the biocompatibility and the surface properties of biomaterials, such as medical grade silicone implants. Silicones are highly resistant against hydrolytic and enzymatic degradation, otherwise they are considerably hydrophobic. Thus, adhesion of unspecific proteins and cells as well as proliferation of inflammatory and pro-fibrotic cells is promoted, all of which trigger foreign body-associated fibrosis [53,54]. The most frequently identified complication is capsular fibrosis, which occurs in up to 27% of the patients in the first year after surgery, and no modification of the silicone surface tested showed beneficial effects so far [55–57]. Coatings of eADF4(C16) yielded reduced fibrosis and contraction upon implantation in rats, since the silk coating inhibited fibroblast adhesion, proliferation and collagen I synthesis. Additionally, significant reduction in capsule thickness, post-operative inflammation and re-modeling of extracellular matrix were observed for silk coated implants in comparison to uncoated ones [58•].


Non-woven meshes made of natural (e.g. collagen, fibroin) or synthetic (e.g. PLGA, PCL) polymers have a great potential as an artificial extracellular matrix (ECM) useful for wound healing and tissue engineering [59]. Electrospinning of eADF4(C16) yielded a mesh enabling the adhesion of fibroblasts (BALB/3T3) as well as their proliferation [28]. In contrast, fibroblasts did not adhere and proliferate on flat films indicating that the topography of a silk scaffold influences cell-matrix interactions [32]. Therefore, structured films were analysed to learn more about the influence of topographies on cell attachment [32,60]. Patterned silk films were made of two different silk proteins (RSSP and recombinant lacewing silk protein) using a Polydimethylsiloxan (PDMS) stamp. Both, fibroblasts (BALB/3T3) and myoblasts (C2C12), preferably adhered and aligned on the ground layer (RSSP) and not on the ridges (recombinant lacewing silk protein) [60]. In general, it can be stated that in the absence of specific cell adhesion domains (such as RGD), charge, wettability and topography of a surface

Figure 2




Overview of unspecific cell-matrix interactions determined by wettability and topography, as well as specific cell-matrix interactions mediated by motifs for cell adhesion.

Figure 3



Processing of recombinant spider silk proteins into colloidal systems for various biomaterials applications, such as drug, gene or enzyme delivery or cell and protein encapsulation for diagnostics.

Figure 4



Possible applications of spider silk materials in biomedicine. Recombinant spider silk proteins can be processed into morphologies other than fibres, broadening the spectrum of possible applications.

play an important role for cell attachment (Figure 2) [28,32]. Interestingly, in case of the patterned silk films the presence of the RGD-cell binding motif did not significantly improve cell binding, indicating the importance of topography [60].

In vitro studies demonstrated that three-dimensional porous scaffolds made of the RSSP rS1/9 allowed mouse fibroblast adhesion and proliferation. Within one week, cells migrated into the deeper layers of the porous scaffolds [61]. Next, porous scaffolds made of rS1/9 and natural silkworm fibroin were compared concerning microstructure and physiological behaviour. There was a detectable difference in the microstructure with the walls of the rS1/9 scaffolds being thicker and containing specific micropores in contrast to fibroin-based ones [62]. The vascularization and intergrowth of the connective tissue, penetrated with nerve fibres, was more prominent in Balb/c mice 8 weeks after subcutaneous implantation using the rS1/9 scaffolds. Further, after

implantation into bone defects of Wistar rats, the regeneration, accounting the number of macrophages and multinuclear giant cells, was better in case of the rS1/9 scaffolds.

Foams and fibre-based matrices made of 4RepCT supported growth, attachment and collagen type I production of human primary fibroblast. Further, macroscopic 4RepCT fibres were well tolerated when implanted subcutaneously in rats. In contrast to the control using silkworm silk (Mersilk™) fibres, 4RepCT-fibres supported ingrowth of fibroblasts and formation of capillaries in the centre of 4RepCT fibre bundles, indicating that 4RepCT is superior in supporting the physiological migration of fibroblasts and angioblasts [63,64]. Yang et al., 2010 used supersaturated simulated body fluids to deposit calcium phosphate coatings on 4RepCT spider silk fibre bundles, yielding a homogeneous and thick crystalline calcium phosphate (CaP) layer, which was a perfect template for bone marrow-derived hMSCs [65].

RSSP can further be used to encapsulate active ingredients including drugs, proteins, genes and cells for delivery or diagnostics (Figure 3) [66–68]. Low molecular weight drugs as well as proteins can be loaded into eADF4(C16) particles, which makes these slowly biodegradable particles a promising drug carrier system, with uptake and release of water-soluble substances being controlled by processing conditions and crosslinking [66–68].

Spider silk particles can also be used as gene carriers for tumour cell-specific delivery. Importantly, pDNA complexes of RSSP containing poly(L-lysine) and the tumour-homing peptides (THPs) F3 and CGKRK showed significantly higher target specificity to tumour cells and transfection efficiency in comparison to pDNA complexes of recombinant spider silk proteins without THP [69,70].

Microcapsules made of eADF4(C16) resemble an enclosed reaction chamber with a semipermeable membrane, in which reactions can be initiated from outside. Further, the capsules can protect the enzyme (β -galactosidase was used as a model) against proteolysis. Recently established eADF4(C16) capsules are therefore promising tunable as well as protective enzyme reaction containers for technical and medical applications [71].

Conclusion

In the past five years, the establishment of engineered recombinant spider silk proteins has led to a steadily increasing number of putative applications of spider silk materials (Figure 4).

The benefits of recombinantly produced spider silk proteins include the high quality and homogeneity of the raw material. Biotechnological production is further easily scalable, and processing as well as functionalization through genetic engineering are controllable. By incorporation of individual amino acids, functional peptide sequences or even proteins, novel spider silk hybrid proteins can be designed with a combination of natural and non-natural silk features, opening the road towards novel multifunctional (bio-) materials.

Acknowledgements

This work was supported by DFG SCHE 603/4.

We would like to thank Dr. Martin Humenik and Martina Elsner for discussions and technical support in creating the artwork and Aniela Heidebrecht, Stefanie Wohlrab and Christian Borkner for providing images.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Gerritsen VB: **The tiptoe of an airbus**. *Protein Spotlight Swiss Prot* 2002, **24**:1-2.
2. Gosline JM, Guerette PA, Ortlepp CS, Savage KN: **The mechanical design of spider silks: from fibroin sequence to mechanical function**. *J Exp Biol* 1999, **202**:3295-3303.
3. Altman GH, Diaz F, Jakuba C, Calabro T, Horan RL, Chen JS, Lu H, Richmond J, Kaplan DL: **Silk-based biomaterials**. *Biomaterials* 2003, **24**:401-416.
4. Vollrath F, Barth P, Basedow A, Engström W, List H: **Local tolerance to spider silks and protein polymers in vivo**. *In Vivo* 2002, **16**:229-234.
5. Heidebrecht A, Scheibel T: **Recombinant production of spider silk proteins**. *Adv Appl Microbiol* 2013, **82**:115-153.
- The review summarizes on one hand different types of spider silk and their properties in a web as well as the natural spinning process, and on the other hand the recombinant production of spider silk proteins by different host organisms, such as bacteria, yeast, plants, insect and mammalian cells.
6. Aprhisiart A, Vollrath F: **Design-features of the orb web of the spider, *Araneus-diadematus***. *Behav Ecol* 1994, **5**:280-287.
7. Hinman MB, Lewis RV: **Isolation of a clone encoding a 2nd dragline silk fibroin – *Nephila-clavipes* dragline silk is a 2-protein fiber**. *J Biol Chem* 1992, **267**:19320-19324.
8. Ayoub NA, Garb JE, Tinghitella RM, Collin MA, Hayashi CY: **Blueprint for a high-performance biomaterial: full-length spider dragline silk genes**. *PLoS ONE* 2007, **2**:e514.
9. Andersen SO: **Amino acid composition of spider silks**. *Compar Biochem Physiol* 1970, **35**:705-711.
10. Kummerlen J, van Beek JD, Vollrath F, Meier BH: **Local structure in spider dragline silk investigated by two-dimensional spin-diffusion nuclear magnetic resonance**. *Macromolecules* 1996, **29**:2920-2928.
11. Hayashi CY, Shipley NH, Lewis RV: **Hypotheses that correlate the sequence, structure, and mechanical properties of spider silk proteins**. *Int J Biol Macromol* 1999, **24**:271-275.
12. Askarieh G, Hedhammar M, Nordling K, Saenz A, Casals C, Rising A, Johansson J, Knight SD: **Self-assembly of spider silk proteins is controlled by a pH-sensitive relay**. *Nature* 2010, **465**:236-239.
13. Eisoldt L, Smith A, Scheibel T: **Decoding the secrets of spider silk**. *Mater Today* 2011, **14**:80-86.
14. Hagn F, Eisoldt L, Hardy J, Vendrely C, Coles M, Scheibel T, Kessler H: **A highly conserved spider silk domain acts as a molecular switch that controls fibre assembly**. *Nature* 2010, **465**:239-242.
15. Hagn F, Thamm C, Scheibel T, Kessler H: **pH-dependent dimerization and salt-dependent stabilization of the N-terminal domain of spider dragline silk-implications for fiber formation**. *Angew Chem Int Ed* 2011, **50**:310-313.
16. Bon M: **A discourse upon usefulness of the silk of spiders**. *Philos Trans* 1710, **27**:2-16.
17. Allmeling C, Jokuszies A, Reimers K, Kall S, Choi CY, Brandes G, Kasper C, Scheper T, Guggenheim M, Vogt PM: **Spider silk fibres in artificial nerve constructs promote peripheral nerve regeneration**. *Cell Prolif* 2008, **41**:408-420.
18. Allmeling C, Jokuszies A, Reimers K, Kall S, Vogt PM: **Use of spider silk fibres as an innovative material in a biocompatible artificial nerve conduit**. *J Cell Mol Med* 2006, **10**:770-777.
19. Radtke C, Allmeling C, Waldmann KH, Reimers K, Thies K, Schenk HC, Hillmer A, Guggenheim M, Brandes G, Vogt PM: **Spider silk constructs enhance axonal regeneration and remyelination in long nerve defects in sheep**. *PLoS ONE* 2011, **6**:e16990.
- Spider silk fibres together with an acellularized vein and Schwann cells embedded in a matrigel were used as guidance material for nerve regeneration. It was demonstrated that spider silk enhances Schwann cell migration, axonal regrowth and remyelination including electrophysiological recovery in a long-distance peripheral nerve gap in rats.
20. Hennecke K, Redeker J, Kuhbier JW, Strauss S, Allmeling C, Kasper C, Reimers K, Vogt PM: **Bundles of spider silk, braided into sutures, resist basic cyclic tests: potential use for flexor tendon repair**. *PLoS ONE* 2013, **8**:e61100.

21. Kuhbier JW, Reimers K, Kasper C, Allmeling C, Hillmer A, Menger B, Vogt PM, Radtke C: **First investigation of spider silk as a braided microsurgical suture.** *J Biomed Mater Res B Appl Biomater* 2011, **97**:381-387.
 22. Madsen B, Shao ZZ, Vollrath F: **Variability in the mechanical properties of spider silks on three levels: interspecific, intraspecific and intraindividual.** *Int J Biol Macromol* 1999, **24**:301-306.
 23. Fox LR: **Cannibalism in natural-populations.** *Annu Rev Ecol System* 1975, **6**:87-106.
 24. Hardy JG, Romer LM, Scheibel TR: **Polymeric materials based on silk proteins.** *Polymer* 2008, **49**:4309-4327.
 25. Spiess K, Wohlrab S, Scheibel T: **Structural characterization and functionalization of engineered spider silk films.** *Soft Matter* 2010, **6**:4168-4174.
 26. Schacht K, Scheibel T: **Controlled hydrogel formation of a recombinant spider silk protein.** *Biomacromolecules* 2011, **12**:2488-2495.
 27. Slotta UK, Rammensee S, Gorb S, Scheibel T: **An engineered spider silk protein forms microspheres.** *Angew Chem Int Ed* 2008, **47**:4592-4594.
 28. Leal-Egana A, Lang G, Mauere C, Wickinghoff J, Weber M, Geimer S, Scheibel T: **Interactions of fibroblasts with different morphologies made of an engineered spider silk protein.** *Adv Eng Mater* 2012, **14**:B67-B75.
 29. Hermanson KD, Huemmerich D, Scheibel T, Bausch AR: **Engineered microcapsules fabricated from reconstituted spider silk.** *Adv Mater* 2007, **19**:1810-1815.
 30. Wohlrab S, Muller S, Schmidt A, Neubauer S, Kessler H, Leal-Egana A, Scheibel T: **Cell adhesion and proliferation on RGD-modified recombinant spider silk proteins.** *Biomaterials* 2012, **33**:6650-6659.
- The engineered spider silk protein eADF4(C16) was modified with the integrin recognition sequence RGD both by a genetic and a chemical approach. After film processing it could be demonstrated that adhesion and proliferation of fibroblasts were significantly improved on films made of both RGD-modified proteins.
31. Bini E, Foo CWP, Huang J, Karageorgiou V, Kitchel B, Kaplan DL: **RGD-functionalized bioengineered spider dragline silk biomaterial.** *Biomacromolecules* 2006, **7**:3139-3145.
 32. Leal-Egana A, Scheibel T: **Interactions of cells with silk surfaces.** *J Mater Chem* 2012, **22**:14330-14336.
 33. Grant DS, Tashiro K-I, Segui-Real B, Yamada Y, Martin G, Kleinman H: **Two different laminin domains mediate the differentiation of human endothelial cells into capillary-like structures in vitro.** *Cell* 1989, **58**:933-943.
 34. Graf J, Iwamoto Y, Sasaki M, Martin G, Kleinman H, Robey F, Yamada Y: **Identification of an amino acid sequence in laminin mediating cell attachment, chemotaxis, and receptor binding.** *Cell* 1987, **48**:989-996.
 35. Widhe M, Johansson U, Hillerdahl CO, Hedhammar M: **Recombinant spider silk with cell binding motifs for specific adherence of cells.** *Biomaterials* 2013, **34**:8223-8234.
 36. Gomes SC, Leonor IB, Mano JF, Reis RL, Kaplan DL: **Antimicrobial functionalized genetically engineered spider silk.** *Biomaterials* 2011, **32**:4255-4266.
 37. Gomes S, Gallego-Llamas J, Leonor IB, Mano JF, Reis RL, Kaplan DL: **Biological responses to spider silk-antibiotic fusion protein.** *J Tissue Eng Regen Med* 2011, **6**:356-368.
 38. Currie HA, Deschaume O, Naik RR, Perry CC, Kaplan DL: **Genetically engineered chimeric silk-silver binding proteins.** *Adv Funct Mater* 2011, **21**:2889-2895.
 39. Foo CWP, Patwardhan SV, Belton DJ, Kitchel B, Anastasiades D, Huang J, Naik RR, Perry CC, Kaplan DL: **Novel nanocomposites from spider silk-silica fusion (chimeric) proteins.** *Proc Natl Acad Sci U S A* 2006, **103**:9428-9433.
 40. Mieszawska AJ, Fourligas N, Georgakoudi I, Ouhib NM, Belton DJ, Perry CC, Kaplan DL: **Osteoinductive silk-silica composite biomaterials for bone regeneration.** *Biomaterials* 2010, **31**:8902-8910.
 41. Belton DJ, Mieszawska AJ, Currie HA, Kaplan DL, Perry CC: **Silk-silica composites from genetically engineered chimeric proteins: materials properties correlate with silica condensation rate and colloidal stability of the proteins in aqueous solution.** *Langmuir* 2012, **28**:4373-4381.
 42. Numata K, Kaplan DL: **Silk-based gene carriers with cell membrane destabilizing peptides.** *Biomacromolecules* 2010, **11**:3189-3195.
 43. Gomes S, Numata K, Leonor IB, Mano JF, Reis RL, Kapan DL: **AFM study of morphology and mechanical properties of a chimeric spider silk and bone sialoprotein protein for bone regeneration.** *Biomacromolecules* 2011, **12**:1675-1685.
 44. Mizuno M, Imai T, Fujisawa R, Tani H, Kuboki Y: **Bone sialoprotein (BSP) is a crucial factor for the expression of osteoblastic phenotypes of bone marrow cells cultured on type I collagen matrix.** *Calcif Tissue Int* 2000, **66**:388-396.
 45. Valverde P, Zhang J, Fix A, Zhu J, Ma WL, Tu QS, Chen J: **Overexpression of bone sialoprotein leads to an uncoupling of bone formation and bone resorption in mice.** *J Bone Miner Res* 2008, **23**:1775-1788.
 46. Gomes S, Leonor IB, Mano JF, Reis RL, Kaplan DL: **Spider silk-bone sialoprotein fusion proteins for bone tissue engineering.** *Soft Matter* 2011, **7**:4964-4973.
 47. Gomes S, Gallego-Llamas J, Leonor IB, Mano JF, Reis RL, Kaplan DL: **In vivo biological responses to silk proteins functionalized with bone sialoprotein.** *Macromol Biosci* 2013, **13**:444-454.
 48. Huang J, Wong C, George A, Kaplan DL: **The effect of genetically engineered spider silk-dentin matrix protein 1 chimeric protein on hydroxyapatite nucleation.** *Biomaterials* 2007, **28**:2358-2367.
 49. Yan WQ, Nakamura T, Kawanabe K, Nishigochi S, Oka M, Kokubo T: **Apatite layer-coated titanium for use as bone bonding implants.** *Biomaterials* 1997, **18**:1185-1190.
 50. Li PJ: **Biomimetic nano-apatite coating capable of promoting bone ingrowth.** *J Biomed Mater Res A* 2003, **66A**:79-85.
 51. Barrere F, van der Valk CM, Meijer G, Dalmeijer RAJ, de Groot K, Layrolle P: **Osteointegration of biomimetic apatite coating applied onto dense and porous metal implants in femurs of goats.** *J Biomed Mater Res B Appl Biomater* 2003, **67**:655-665.
 52. Chou YF, Dunn JC, Wu BM: **In vitro response of MC3T3-E1 preosteoblasts within three-dimensional apatite-coated PLGA scaffolds.** *J Biomed Mater Res B Appl Biomater* 2005, **75**:81-90.
 53. Pajkos A, Deva AK, Vickery K, Cope C, Chang L, Cossart YE: **Detection of subclinical infection in significant breast implant capsules.** *Plast Reconstruct Surg* 2003, **111**:1605-1611.
 54. Hezi-Yamit A, Sullivan C, Wong J, David L, Chen MF, Cheng PW, Shumaker D, Wilcox JN, Udipti K: **Impact of polymer hydrophilicity on biocompatibility: implication for DES polymer design.** *J Biomed Mater Res A* 2009, **90**:133-141.
 55. Henriksen TF, Fryzek JP, Holmich LR, McLaughlin JK, Kjoller K, Hoyer AP, Olsen DH, Friis S: **Surgical intervention and capsular contracture after breast augmentation – a prospective study of risk factors.** *Ann Plast Surg* 2005, **54**:343-351.
 56. Dancey A, Nassimzadeh A, Levick P: **Capsular contracture – what are the risk factors? A 14 year series of 1400 consecutive augmentations.** *J Plast Reconstruct Aesth Surg* 2012, **65**:213-218.
 57. Zeplin PH, Larena-Avellaneda A, Jordan M, Laske M, Schmidt K: **Phosphorylcholine-coated silicone implants effect on inflammatory response and fibrous capsule formation.** *Ann Plast Surg* 2010, **65**:560-564.
 58. Zeplin PH, Maksimovikj NC, Jordan MC, Nickel J, Lang G, Leimer AH, Römer L, Scheibel T: **Spider silk coatings as a bioshield to reduce periprosthetic fibrous capsule formation.** *Adv Funct Mater* 2014 <http://dx.doi.org/10.1002/adfm.20130281>.

Silk coating of medical grade silicone was shown to yield reduced major post-operative complications, such as capsule formation and contraction.

59. Sell SA, Wolfe PS, Garg K, McCool JM, Rodriguez IA, Bowlin GL: **The use of natural polymers in tissue engineering: a focus on electrospun extracellular matrix analogues.** *Polymers* 2010, **2**:522-553.
 60. Bauer F, Wohlrab S, Scheibel T: **Controllable cell adhesion, growth and orientation on layered silk protein films.** *Biomater Sci* 2013, **1**:1244-1249.
 61. Agapov II, Pustovalova OL, Moisenovich MM, Bogush VG, Sokolova OS, Sevastyanov VI, Debabov VG, Kirpichnikov MP: **Three-dimensional scaffold made from recombinant spider silk protein for tissue engineering.** *Doklady Biochem Biophys* 2009, **426**:127-130.
 62. Moisenovich MM, Pustovalova O, Shackelford J, Vasiljeva TV, Druzhinina TV, Kamenchuk YA, Guzev VV, Sokolova OS, Bogush VG, Debabov VG *et al.*: **Tissue regeneration in vivo within recombinant spidroin 1 scaffolds.** *Biomaterials* 2012, **33**:3887-3898.
- Two porous scaffolds made from different silk proteins (fibroin of *B. mori* and the recombinant spider silk protein rS1/9) by salt leaching were compared concerning microstructure and biocompatibility. Especially the matrices derived from the recombinant rS1/9 protein were favourable for tissue engineering due to their internal microstructures.
63. Widhe M, Bysell H, Nystedt S, Schenning I, Malmsten M, Johansson J, Rising A, Hedhammar M: **Recombinant spider silk as matrices for cell culture.** *Biomaterials* 2010, **31**:9575-9585.
 64. Fredriksson C, Hedhammar M, Feinstein R, Nordling K, Kratz G, Johansson J, Huss F, Rising A: **Tissue response to subcutaneously implanted recombinant spider silk: an in vivo study.** *Materials* 2009, **2**:1908-1922.
 65. Yang LA, Hedhammar M, Blom T, Leifer K, Johansson J, Habibovic P, van Blitterswijk CA: **Biomimetic calcium phosphate coatings on recombinant spider silk fibres.** *Biomed Mater* 2010, **5**:1-10.
 66. Lammel A, Schwab M, Hofer M, Winter G, Scheibel T: **Recombinant spider silk particles as drug delivery vehicles.** *Biomaterials* 2011, **32**:2233-2240.
 67. Hofer M, Winter G, Myschik J: **Recombinant spider silk particles for controlled delivery of protein drugs.** *Biomaterials* 2012, **33**:1554-1562.
 68. Blüm C, Scheibel T: **Control of drug loading and release properties of spider silk sub-microparticles.** *BioNanoSci* 2012, **2**:67-74.
 69. Numata K, Mieszawska-Czajkowska AJ, Kvenvold LA, Kaplan DL: **Silk-based nanocomplexes with tumor-homing peptides for tumor-specific gene delivery.** *Macromol Biosci* 2012, **12**: 75-82.
- The authors designed nanoscale complexes of recombinant silk molecules containing tumour-homing peptides with DNA as less cytotoxic and highly target-specific gene carriers. The silk-based block copolymers showed relatively high target specificity and transfection efficiency.
70. Numata K, Reagan MR, Goldstein RH, Rosenblatt M, Kaplan DL: **Spider silk-based gene carriers for tumor cell-specific delivery.** *Bioconjug Chem* 2011, **22**:1605-1610.
 71. Blüm C, Nichtl A, Scheibel T: **Spider silk capsules as protective reaction containers for enzymes.** *Adv Funct Mater* 2014, **24**:763-768.
- Microcapsules made of the recombinant spider silk protein eADF4(C16) can be used as a container for high molecular weight compounds such as enzymes. The capsules constitute small enclosed reaction chambers with a semi-permeable membrane.