

1

University of Warsaw

2

Faculty of Mathematics, Informatics and Mechanics

3

Katarzyna Kowalska

Student no. 371053

4

5

# Approximation and Parametrized Algorithms for Segment Set Cover

6

Master's thesis

7

in COMPUTER SCIENCE

8

Supervisor:

**dr Michał Pilipczuk**

Instytut Informatyki

9

June 2020

10 **Supervisor's statement**

11 Hereby I confirm that the presented thesis was prepared under my supervision and  
12 that it fulfils the requirements for the degree of Master of Computer Science.

13 Date

Supervisor's signature

14 **Author's statement**

15 Hereby I declare that the presented thesis was prepared by me and none of its contents  
16 was obtained by means that are against the law.

17 The thesis has never before been a subject of any procedure of obtaining an academic  
18 degree.

19 Moreover, I declare that the present version of the thesis is identical to the attached  
20 electronic version.

21 Date

Author's signature

22

## **Abstract**

23 The work presents a study of different geometric set cover problems. It mostly focuses on  
24 segment set cover and its connection to the polygon set cover.

25

## **Keywords**

26 set cover, geometric set cover, FPT,  $W[1]$ -completeness, APX-completeness, PCP theorem,  
27 NP-completeness

28

## **Thesis domain (Socrates-Erasmus subject area codes)**

29 11.3 Informatyka

30

31

## **Subject classification**

32 D. Software

33 D.127. Blabalgorithms

34 D.127.6. Numerical blabalysis

35

## **Tytuł pracy w języku polskim**

36 Algorytmy parametryzowania i trudność aproksymacji problemu pokrywania zbiorów  
37 odcinkami na płaszczyźnie



# Contents

39	<b>1. Introduction</b>	5
40	<b>2. Definitions</b>	7
41	2.1. Geometric Set Cover	7
42	2.2. Approximation	7
43	2.3. $\delta$ -extensions	7
44	<b>3. Geometric Set Cover with segments</b>	9
45	3.1. FPT for segments	9
46	3.1.1. Axis-parallel segments	9
47	3.1.2. Segments in arbitrary directions	9
48	3.2. APX-completeness for segments parallel to axes	11
49	3.2.1. MAX-(3,3)-SAT and statement of reduction	11
50	3.2.2. Reduction	13
51	3.2.2.1. VARIABLE-gadget	13
52	3.2.2.2. OR-gadget	14
53	3.2.2.3. CLAUSE-gadget	16
54	3.2.2.4. Summary	18
55	3.2.2.5. Summary of construction	18
56	3.2.3. Construction lemmas and proof of Lemma 3	20
57	3.3. FPT for weighted segments with $\delta$ -extensions	22
58	3.4. W[1]-completeness for weighted segments in 3 directions	25
59	3.5. What is missing	27
60	<b>4. Geometric Set Cover with lines</b>	29
61	4.1. Lines parallel to one of the axis	29
62	4.2. FPT for arbitrary lines	29
63	4.3. APX-completeness for arbitrary lines	29
64	4.4. 2-approximation for arbitrary lines	30
65	4.5. Connection with general set cover	30
66	<b>5. Geometric Set Cover with polygons</b>	31
67	5.1. State of the art	31
68	<b>6. Conclusions</b>	33



# Chapter 1

## Introduction

The Set Cover problem is one of the most common NP-complete problems. [tutaj referencja]  
We are given a family of sets and have to choose the smallest subfamily of these sets that cover  
all their elements. This problem naturally extends to settings where we put different weights  
on the sets and look for the subfamily of the minimal weight. This problem is NP-complete  
even without weights and if we put restrictions on what the sets can be. One of such variants  
is Vertex Cover problem, where sets have size 2 (they are edges in a graph).

In this work we focus on another such variant where the sets correspond to some geometric  
shapes and only some points of the plane have to be covered. When these shapes are rectangles  
with edges parallel to the axis, the problem can be proven to be W[1]-complete (solution of  
size  $k$  cannot be found in  $n^o(k)$  time), APX-complete (for sufficiently small  $\epsilon > 0$ , the problem  
does not admit  $1 + \epsilon$ -approximation scheme) [referencje].

Some of these settings are very easy. Set cover with lines parallel to one of the axis can  
be solved in polynomial time.

There is a notion of  $\delta$ -expansions, which loosen the restrictions on geometric set cover. We  
allow the objects to cover the points after  $\delta$ -expansion and compare the result to the original  
setting. This way we can produce both FPT and EPTAS for the rectangle set cover with  
 $\delta$ -extensions [referencje].

**Our contribution.** In this work, we prove that unweighted geometric set cover with seg-  
ments is fixed parameter tractable (FPT).

Moreover, we show that geometric set cover with segments is APX-complete for unweighted  
axis-parallel segments, even with  $1/2$ -extensions. So the problem for very thin rectangles  
also can't admit PTAS. Therefore, in the efficient polynomial-time approximation scheme  
(EPTAS) for *fat polygons* by [Har-Peled and Lee, 2009], the assumption about polygons  
being fat is necessary.

Finally, we show that geometric set cover with weighted segments in 3 directions is  
W[1]-complete. However, geometric set cover with weighted segments is FPT if we allow  
 $\delta$ -extension.

This result is especially interesting, since it's counter-intuitive that the unweighted setting  
is FPT and the weighted setting is W[1]-complete. Most of such problems (like vertex cover  
or [wiecej przykladow]) are equally hard in both weighted and unweighted settings.





## 101 Chapter 2

## 102 Definitions

### 103 2.1. Geometric Set Cover

104 In the geometric set cover problem we are given  $\mathcal{P}$  – a set of objects, which are connected  
105 subsets of the plane,  $\mathcal{C}$  – a set of points in the plane. The task is to choose  $\mathcal{R} \subseteq \mathcal{P}$  such that  
106 every point in  $\mathcal{C}$  is inside some element from  $\mathcal{R}$  and  $|\mathcal{R}|$  is minimized.

107 In the parametrized setting for a given  $k$ , we only look for a solution  $\mathcal{R}$  such that  $|\mathcal{R}| \leq k$ .

108 In the weighted setting, there is some given weight function  $f : \mathcal{P} \rightarrow \mathbb{R}^+$ , and we would  
109 like to find a solution  $\mathcal{R}$  that minimizes  $\sum_{R \in \mathcal{R}} f(R)$ .

### 110 2.2. Approximation

111 Let us recall some definitions related to optimization problems that are used in the following  
112 sections.

113 **Definition 1.** A **polynomial-time approximation scheme (PTAS)** for a minimization  
114 problem  $\Pi$  is a family of algorithms  $\mathcal{A}_\epsilon$  for every  $\epsilon > 0$  such that  $\mathcal{A}_\epsilon$  takes an instance  $I$  of  $\Pi$   
115 and in polynomial time finds a solution that is within a factor  $(1 + \epsilon)$  of being optimal. That  
116 means the reported solution has weight at most  $(1 + \epsilon)\text{opt}(I)$ , where  $\text{opt}(I)$  is the weight of  
117 an optimal solution for  $I$ .

118 **Definition 2.** A problem  $\Pi$  is **APX-hard** if assuming  $P \neq NP$ , there exists  $\epsilon > 0$  such that  
119 there is no polynomial-time  $(1 + \epsilon)$ -approximation algorithm for  $\Pi$ .

### 120 2.3. $\delta$ -extensions

121 TODO PLACEHOLDER for introductory text

122  $\delta$ -extensions is one of the modifications to a problem, that makes geometric set cover  
123 problem easier, it has been already used in literature (place some refrence here).

124 **Definition 3** ( $\delta$ -extensions for center-symmetric objects). For any  $\delta > 0$  and a center-  
125 symmetric object  $L$  with centre of symmetry  $S = (x_s, y_s)$ , the  **$\delta$ -extension** of  $L$  is the  
126 object  $L^{+\delta} = \{(1 + \delta) \cdot (x - x_s, y - y_s) + (x_s, y_s) : (x, y) \in L\}$ , that is,  $L^{+\delta}$  is the image of  $L$   
127 under homothety centered at  $S$  with scale  $(1 + \delta)$

128 The geometric set cover problem with  $\delta$ -extensions is a modified version of geometric set  
129 cover where:

- We need to cover all the points in  $\mathcal{C}$  with objects from  $\{P^{+\delta} : P \in \mathcal{P}\}$  (which always include no fewer points than the objects before  $\delta$ -extensions);

- We look for a solution that is no larger than the optimum solution for the original problem. Note that it does not need to be an optimal solution in the modified problem.

Formally, we have the following.

**Definition 4** (Geometric set cover problem with  $\delta$ -extensions). The geometric set cover problem with  $\delta$ -extensions is the problem where for an input instance  $I = (\mathcal{P}, \mathcal{C})$ , the task is to output a solution  $\mathcal{R} \subseteq \mathcal{P}$  such that the  $\delta$ -extended set  $\{R^{+\delta} : R \in \mathcal{R}\}$  covers  $\mathcal{C}$  and is no larger than the optimal solution for the problem without extensions, i.e.  $|\mathcal{R}| \leq |\text{opt}(I)|$ .

TODO: Some text

**Definition 5** (Geometric set cover PTAS with  $\delta$ -extensions). We define a PTAS for geometric set cover with  $\delta$ -extensions as a family of algorithms  $\{\mathcal{A}_{\delta, \epsilon}\}_{\delta, \epsilon > 0}$  that each takes as an input instance  $I = (\mathcal{P}, \mathcal{C})$ , and in polynomial-time outputs a solution  $\mathcal{R} \subseteq \mathcal{P}$  such that the  $\delta$ -extended set  $\{R^{+\delta} : R \in \mathcal{R}\}$  covers  $\mathcal{C}$  and is within a  $(1 + \epsilon)$  factor of the optimal solution for this problem without extensions, i.e.  $(1 + \epsilon)|\mathcal{R}| \leq |\text{opt}(I)|$ .

## Chapter 3

# Geometric Set Cover with segments

### 3.1. FPT for segments

In this section we consider the fixed-parameter tractable algorithms for unweighted geometric set cover with segments. Setting where segments are limited to be axis-parallel (or limited to constant number of directions) has an FPT algorithm already present in literature. We present an FPT algorithm for unweighted geometric set cover with segments, where segments are in arbitrary directions.

#### 3.1.1. Axis-parallel segments

You can find this in Platypus book. (TODO add referece)

We show an  $\mathcal{O}(2^k)$ -time branching algorithm. In each step, the algorithm selects a point  $a$  which is not yet covered, branches to choose one of the two directions, and greedily chooses a segment in that direction to cover  $a$ . This proceeds until either all points are covered or  $k$  segments are chosen.

Let us take the point  $a = (x_a, y_a)$  which is the smallest among points that are not yet covered in the lexicographic ordering of points in  $\mathbb{R}^2$ . We need to cover  $a$  with some of the remaining segments.

Branch over the choice of one of the coordinates ( $x$  or  $y$ ); without loss of generality, let us assume we chose  $x$ . Among the segments lying on line  $x = x_a$ , we greedily add to the solution the one that covers the most points. As  $a$  was the smallest in the lexicographical order, then all points on line  $x = x_a$  have the  $y$ -coordinate larger than  $y_a$ . Therefore, if we denote the greedily chosen segment as  $s$ , then any other segment on  $x = x_a$  that covers  $a$  can only cover a (possibly improper) subset of points covered by  $s$ . Thus, greedily choosing  $s$  is optimal.

In each step of the algorithm we add one segment to the solution, thus each branch can stop at depth  $k$ . If no branch finds a solution, then that means a solution of size at most  $k$  does not exist.

TODO: Maybe split it into theorem + algorithm + explanation like in section 3.1.2

**Remark 1.** *The same algorithm can be used for segments in  $d$  directions, where we branch over  $d$  directions and it runs in complexity  $\mathcal{O}(d^k)$ .*

#### 3.1.2. Segments in arbitrary directions

In this section we consider setting where segments are not constrained to only  $d$  directions. We present a fixed-parameter tractable algorithm, where parameter is the size of the solution.

**Theorem 1. (FPT for segment cover).** *There exists an algorithm that given a family  $\mathcal{P}$  of  $n$  segments (in any direction), a set of  $m$  points  $\mathcal{C}$  and a parameter  $k$ , runs in time  $k^{O(k)} \cdot (nm)^2$ , and outputs a subfamily  $\mathcal{R} \subseteq \mathcal{P}$  such that  $|\mathcal{R}| \leq k$  and  $\mathcal{R}$  covers all points in  $\mathcal{C}$ , or determines that such a set  $\mathcal{R}$  does not exist.*

We will need the following lemmas.

**Lemma 1.** *Given an instance  $(\mathcal{P}, \mathcal{C})$  of the segment cover problem, without a loss of generality we can assume that no segment covers a superset of what another segment covers. That is, for any distinct  $A, B \in \mathcal{P}$ , we have  $A \cap \mathcal{C} \not\subseteq B \cap \mathcal{C}$  and  $A \cap \mathcal{C} \not\supseteq B \cap \mathcal{C}$ .*

*Proof.* Trivial. □

**Lemma 2.** *Given an instance  $(\mathcal{P}, \mathcal{C})$  of the segment cover problem, if there exists a line  $L$  with at least  $k + 1$  points on it, then there exists a subset  $\mathcal{A} \subseteq \mathcal{P}$ ,  $|\mathcal{A}| \leq k$ , such that every solution  $\mathcal{R}$  with  $|\mathcal{R}| \leq k$  satisfies  $|\mathcal{A} \cap \mathcal{R}| \geq 1$ . Moreover, such a subset can be found in polynomial time.*

*Proof.* First we use Lemma 1.

Let us enumerate the points from  $\mathcal{C}$  that lie on  $L$  as  $x_1, x_2, \dots, x_t$  in the order in which they appear on  $L$ . Every segment that is not collinear with  $L$  can cover at most one of these points. Therefore, in any solution of size not larger than  $k$ , among any  $k$  of these points at least one must be covered with segment collinear with  $L$ .

Therefore, every solution needs to take one of the segments collinear with  $L$  that covers any of the points  $x_1, x_2, \dots, x_k$ . After using reduction from Lemma 1, there are at most  $k$  such segments that are distinct. □

We are ready to prove Theorem 1.

*Proof of Theorem 1.*

We will prove this theorem by presenting a branching algorithm that works in desired complexity. It branches over the choice of segments to cover lines with a lot of points, then finally solving the small instance, where every line has at most  $k$  points by checking all possible solutions.

**Algorithm.** First we use Lemma 1.

Next, we present a recursive algorithm. Given an instance of the problem:

- (1) If there exist a line with at least  $k + 1$  points from  $\mathcal{C}$ , we branch over adding to the solution one of the at most  $k$  possible segments provided by Lemma 2; name this segment  $S$ . Then we find a solution  $\mathcal{R}$  for the problem for points  $\mathcal{C} - S$ , segments  $\mathcal{P} - \{S\}$ , and parameter  $k - 1$ . We return  $\mathcal{R} \cup \{S\}$ .
- (2) If every line has at most  $k$  points on it and  $|\mathcal{C}| > k^2$ , then answer NO.
- (3) If  $|\mathcal{C}| \leq k^2$ , solve the problem by brute force: check all subsets of  $\mathcal{P}$  of size at most  $k$ .

**Correctness.** Lemma 2 proves that at least one segment that we branch over in (1) must be present in every solution  $\mathcal{R}$  with  $|\mathcal{R}| \leq k$ . Therefore, the recursive call can find a solution, provided there exists one.

In (2) the answer is no, because every line covers no more than  $k$  points from  $\mathcal{C}$ , which implies the same about every segment from  $\mathcal{P}$ . Under this assumption we can cover only  $k^2$  points with a solution of size  $k$ , which is less than  $|\mathcal{C}|$ .

Checking all possible solutions in (3) is trivially correct.

219 **Complexity.** In the leaves of recursion we have  $|\mathcal{C}| \leq k^2$ , so  $|\mathcal{P}| \leq k^4$ , because every  
220 segments can be uniquely identified by the two extreme points it covers (by Lemma 1).  
221 Therefore, there are  $\binom{k^4}{k}$  possible solutions to check, each can be checked in time  $O(k|\mathcal{C}|)$ .  
222 Therefore, (3) takes time  $k^{O(k)}$ .

223 In this branching algorithm our parameter  $k$  is decreased with every recursive call, so we  
224 have at most  $k$  levels of recursion with branching over  $k$  possibilities. Candidates to branch  
225 over can be found on each level in time  $O((nm)^2)$ .

226 Reduction from Lemma 1 can be implemented in time  $O(n^2m)$ .

227 It follows that the overall complexity is  $O((nm)^2 \cdot k^{O(k)})$  □

## 228 3.2. APX-completeness for segments parallel to axes

229 In this section we analyze whether there exists PTAS for geometric set cover for rectangles.  
230 We show that we can restrict this problem to a very simple setting: segments parallel to axes  
231 and allow  $(1/2)$ -extension, and the problem is still APX-hard. Note that segments are just  
232 degenerated rectangles with one side being very narrow.

233 Our results can be summarized in the following theorem and this section aims to prove it.

234 **Theorem 2.** *(axis-parallel segment set cover with  $1/2$ -extension is APX-hard).*  
235 *Unweighted geometric set cover with axis-parallel segments in 2D (even with  $1/2$ -extension)*  
236 *is APX-hard. That is, assuming  $P \neq NP$ , there does not exist a PTAS for this problem.*

237 Theorem 2 implies the following.

238 **Corollary 1.** *(rectangle set cover is APX-hard).* *Unweighted geometric set cover with*  
239 *rectangles (even with  $1/2$ -extension) is APX-hard.*

240 We prove Theorem 2 by taking a problem that is APX-hard and showing a reduction. For  
241 this problem we choose MAX-(3,3)-SAT which we define below.

### 242 3.2.1. MAX-(3,3)-SAT and statement of reduction

243 **Definition 6.** MAX-3SAT is the following maximization problem. We are given a 3-CNF  
244 formula, and need to find an assignment of variables that satisfies the most clauses.

245 **Definition 7.** MAX-(3,3)-SAT is a variant of MAX-3SAT with an additional restriction  
246 that every variable appears in exactly 3 clauses. Note that thus, the number of clauses is  
247 equal to the number of variables.

248 In our proof of Theorem 2 we use hardness of approximation of MAX-(3,3)-SAT proved  
249 in [Håstad, 2001] and described in Theorem 3 below.

250 **Definition 8** ( $\alpha$ -satisfiable MAX-3SAT formula). MAX-3SAT formula of size  $n$  is at most  
251  $\alpha$ -satisfiable, if every assignment of variables satisfies no more than  $\alpha n$  clauses.

252 **Theorem 3.** [Håstad, 2001]

253 *For any  $\epsilon > 0$ , it is NP-hard to distinguish satisfiable (3,3)-SAT formulas from at most*  
254  *$(7/8 + \epsilon)$ -satisfiable (3,3)-SAT formulas.*

Given an instance  $I$  of MAX-(3,3)-SAT, we construct an instance  $J$  of axis-parallel segment set cover problem, such that for a sufficiently small  $\epsilon > 0$ , a polynomial time  $(1 + \epsilon)$ -approximation algorithm for  $J$  would be able to distinguish whether an instance  $I$  of MAX-(3,3)-SAT is fully satisfiable or is at most  $(7/8 + \epsilon)$ -satisfiable. However, according to (Theorem 3) the latter problem is NP-hard. This would imply  $P = NP$ , contradicting the assumption.

The following lemma encapsulates the properties of the reduction described in this section, and it allows us to prove Theorem 2.

**Lemma 3.** *Given an instance  $S$  of MAX-(3,3)-SAT with  $n$  variables and optimum value  $opt(S)$ , we can construct an instance  $I$  of geometric set cover with axis-parallel segments in 2D, such that:*

(1) *For every solution  $X$  of instance  $I$ , there exists a solution of  $S$  that satisfies at least  $15n - |X|$  clauses.*

(2) *For every solution of instance  $S$  that satisfies  $w$  clauses, there exists a solution of  $I$  of size  $15n - w$ .*

(3) *Every solution with 1/2-extensions of  $I$  is also a solution to the original instance  $I$ .*

*Therefore, the optimum size of a solution of  $I$  is  $opt(I) = 15n - opt(S)$ .*

We prove Lemma 3 in subsequent sections, but meanwhile let us prove Theorem 2 using Lemma 3 and Theorem 3.

TODO: This below can't use current template

*Proof of Theorem 2.*

Consider any  $0 < \epsilon < 1/(15 \cdot 8)$ .

Let us assume that there exists a polynomial-time  $(1 + \epsilon)$ -approximation algorithm for unweighted geometric set cover with axis-parallel segments in 2D with (1/2)-extensions. We construct an algorithm that solves the problem stated in Theorem 3, thereby proving that  $P = NP$ .

Take an instance  $S$  of MAX-(3,3)-SAT to be distinguished and construct an instance of geometric set cover  $I$  using Lemma 3. We now use the  $(1 + \epsilon)$ -approximation algorithm for geometric set cover on  $I$ . Denote the size of the solution returned by this algorithm as  $approx(I)$ . We prove that if in  $S$  one can satisfy at most  $(\frac{7}{8} + \epsilon)n$  clauses, then  $approx(I) \geq 15n - (\frac{7}{8} + \epsilon)n$  and if  $S$  is satisfiable, then  $approx(I) < 15n - (\frac{7}{8} + \epsilon)n$ .

**Assume  $S$  satisfiable.** From the definition of  $S$  being satisfiable, we have:

$$opt(S) = n.$$

From Lemma 3 we have:

$$opt(I) = 14n.$$

Therefore,

$$\begin{aligned} approx(I) &\leq (1 + \epsilon)opt(I) = 14n(1 + \epsilon) = 14n + 14\epsilon \cdot n = \\ &= 14n + (15\epsilon - \epsilon)n < 14n + \left(\frac{1}{8} - \epsilon\right)n = 15n - \left(\frac{7}{8} + \epsilon\right)n \end{aligned}$$

**Assume  $S$  is at most  $(\frac{7}{8} + \epsilon)$  satisfiable.** From the definition of  $S$  being at most  $(\frac{7}{8} + \epsilon)n$  satisfiable, we have:

$$opt(S) \leq \left(\frac{7}{8} + \epsilon\right)n$$

From Lemma 3 we have:

$$opt(I) \geq 15n - \left(\frac{7}{8} + \epsilon\right)n$$

Since a solution to  $I$  with  $\frac{1}{2}$ -extensions is also a solution without extensions, by Lemma 3 (3.), we have:

$$approx(I) \geq opt(I) = 15n - \left(\frac{7}{8} + \epsilon\right)n$$

Therefore, by using the assumed  $(1 + \epsilon)$ -approximation algorithm, it is possible to distinguish the case when  $S$  is satisfiable from the case when it is at most  $(\frac{7}{8} + \epsilon)n$  satisfiable, it suffices to compute  $approx(I)$  with  $15n - (\frac{7}{8} + \epsilon)n$ . Hence, the assumed approximation algorithm cannot exist, unless  $P = NP$ .  $\square$

### 3.2.2. Reduction

We proceed to the proof of Lemma 3. That is, we show a reduction from MAX-(3,3)-SAT problem to geometric set cover with segments parallel to axis. Moreover, the obtained instance of geometric set cover will be robust to 1/2-extensions (have the same optimal solution after 1/2-extension).

The construction will be composed of 2 types of gadgets: **VARIABLE-gadgets** and **CLAUSE-gadgets**. **CLAUSE-gadgets** would be constructed using two **OR-gadgets** connected together. Every gadget consists of a point set and a segment set.

#### 3.2.2.1. VARIABLE-gadget

**VARIABLE-gadget** is responsible for choosing the value of a variable in a CNF formula. It allows two minimum solutions of size 3 each. These two choices correspond to the two Boolean values of the variable.

**Points.** Define points  $a, b, c, d, e, f, g, h$  as follows, where  $L = 12n$ :




Figure 3.1: **VARIABLE-gadget**. We denote the set of points marked with black circles as  $\text{pointsVariable}_i$ , and they need to be covered (are part of the set  $\mathcal{C}$ ). Note that some of the points are not marked as black dots and exists only to name segments for further reference. We denote the set of red segments as  $\text{chooseVariable}_i^{\text{false}}$  and the set of blue segments as  $\text{chooseVariable}_i^{\text{true}}$ .

$$\begin{array}{llll}
a = (-L, 0) & b = (-\frac{2}{3}L, 0) & c = (-\frac{1}{3}L, 0) & d = (-L, 1) \\
e = (-\frac{2}{3}L, 1) & f = (-\frac{2}{3}L, 2) & g = (L, 0) & h = (L, 2)
\end{array}$$

Let us define:

$$\text{pointsVariable} = \{a, b, c, d, e, f\}$$

and

$$\text{pointsVariable}_i = \text{pointsVariable} + (0, 4i)$$

We denote  $a_i = a + (0, 4i)$  etc.

**Segments.** Let us define:

$$\text{chooseVariable}_i^{\text{true}} = \{(a_i, d_i), (b_i, f_i), (c_i, g_i)\}$$

$$\text{chooseVariable}_i^{\text{false}} = \{(a_i, c_i), (d_i, e_i), (f_i, h_i)\}$$

$$\text{segmentsVariable}_i = \text{chooseVariable}_i^{\text{true}} \cup \text{chooseVariable}_i^{\text{false}}$$

**Lemma 4.** For any  $1 \leq i \leq n$ , points in  $\text{pointsVariable}_i$  can be covered using 3 segments from  $\text{segmentsVariable}_i$ .

*Proof.* We can use either set  $\text{chooseVariable}_i^{\text{true}}$  or  $\text{chooseVariable}_i^{\text{false}}$ .  $\square$

**Lemma 5.** For any  $1 \leq i \leq n$ , points in  $\text{pointsVariable}_i$  can not be covered with fewer than 3 segments from  $\text{segmentsVariable}_i$ .

*Proof.* No segment of  $\text{segmentsVariable}_i$  covers more than one point from  $\{d_i, f_i, c_i\}$ , therefore  $\text{pointsVariable}_i$  can not be covered with fewer than 3 segments.  $\square$

**Lemma 6.** For every set  $A \subseteq \text{segmentsVariable}_i$  such that  $A$  covers  $\text{pointsVariable}_i$  and  $(c_i, g_i), (f_i, h_i) \in A$ , it holds that  $|A| \geq 4$ .

*Proof.* No segment from  $\text{segmentsVariable}_i$  covers more than one point from  $\{a_i, e_i\}$ , therefore  $\text{pointsVariable}_i - \{c_i, f_i, g_i, h_i\}$  can not be covered with fewer than 2 segments.  $\square$

### 3.2.2.2. OR-gadget

TODO: expand the figure to include x, y and maybe name (t,v) as result

OR-gadget has 3 important segments –  $x, y, \text{result}$ , where  $x$  and  $y$  are inputs and part of different gadgets.

Gadget has a minimal solution of weight  $w$  and  $\text{result}$  can be part of minimal solution only if  $x$  or  $y$  are also chosen for the solution. If none of them are chosen, then solution choosing  $\text{result}$  segment has weight at least  $w + 1$ . Therefore the following formula holds for a solution  $R$  assuming that  $R$  uses only  $w$  from this OR-gadget:

$$(x \in R) \vee (y \in R) \iff \text{result} \in R$$

Only 3 points that belong to this segment  $(l_{i,j}, p_{i,j}, v_{i,j})$  can be covered by seg





Figure 3.2: **OR-gadget**. Figure presenting OR-gadget: segments from  $\text{chooseOr}_{i,j}^{false}$  are red, segments from  $\text{chooseOr}_{i,j}^{true}$  are blue, segments from  $\text{orMoveVariable}_{i,j}$  are yellow and green.

328 **Points.**

$$\begin{array}{llll}
 l_0 = (0, 0) & m_0 = (0, 1) & n_0 = (0, 2) & o_0 = (0, 3) \\
 p_0 = (0, 4) & q_0 = (1, 1) & r_0 = (1, 3) & s_0 = (2, 1) \\
 t_0 = (2, 2) & u_0 = (2, 3) & v_0 = (3, 2) & 
 \end{array}$$

$$vec_{i,j} = (10i + 3 + 3j, 4n + 2j)$$

330 Define  $\{l_{i,j}, m_{i,j} \dots v_{i,j}\}$  as  $\{l_0, m_0 \dots v_0\}$  shifted by  $vec_{i,j}$

331 Note that  $v_{i,0} = l_{i,1}$  (see Figure 3.3)

$$\text{pointsOr}_{i,j} = \{l_{i,j}, m_{i,j}, n_{i,j}, o_{i,j}, p_{i,j}, q_{i,j}, r_{i,j}, s_{i,j}, t_{i,j}, u_{i,j}\}$$

332 Note that  $\text{pointsOr}_{i,j}$  does not include point  $v_{i,j}$

333 **Segments.** We define names subsets of segments, to refer to them in lemmas.

$$\text{chooseOr}_{i,j}^{false} = \{(q_{i,j}, r_{i,j}), (s_{i,j}, u_{i,j})\}$$

$$\text{chooseOr}_{i,j}^{true} = \{(m_{i,j}, s_{i,j}), (o_{i,j}, u_{i,j}), (t_{i,j}, v_{i,j})\}$$

$$\text{orMoveVariable}_{i,j} = \{(l_{i,j}, n_{i,j}), (n_{i,j}, p_{i,j})\}$$

334 Segments in OR-gadget:

$$\text{segmentsOr}_{i,j} = \text{chooseOr}_{i,j}^{false} \cup \text{chooseOr}_{i,j}^{true} \cup \text{orMoveVariable}_{i,j}$$

335 **Lemma 7.** For any  $1 \leq i \leq n, j \in \{0, 1\}$  and  $x \in \{l_{i,j}, p_{i,j}\}$ , points in  $\text{pointsOr}_{i,j} - \{x\} \cup \{v_{i,j}\}$   
 336 can be covered with 4 segments from  $\text{segmentsOr}_{i,j}$ .

337 *Proof.* We can do that using one segment from  $\text{orMoveVariable}_{i,j}$ , the one that does not cover  
 338  $x$ , and all segments from  $\text{chooseOr}_{i,j}^{\text{true}}$ .  $\square$

339 **Lemma 8.** For any  $1 \leq i \leq n, j \in \{0, 1\}$ , points in  $\text{pointsOr}_{i,j}$  can be covered with 4 segments  
 340 from  $\text{segmentsOr}_{i,j}$ .

341 *Proof.* We can do that using segments from  $\text{orMoveVariable}_{i,j}$  and  $\text{chooseOr}_{i,j}^{\text{false}}$ .  $\square$

342 TODO: I don't think we need these, we can just show independent set on CLAUSE-gadget  
 343 and it's less confusing

344 **Lemma 9.** For any  $1 \leq i \leq n, j \in \{0, 1\}$  and  $x \in \{l_{i,j}, p_{i,j}\}$  points in  $\text{pointsOr}_{i,j} - \{x\} \cup \{v_{i,j}\}$   
 345 can not be covered with with less than 4 segments from  $\text{segmentsOr}_{i,j}$ .

346 *Proof.* TODO  $\square$

347 **Lemma 10.** For any  $1 \leq i \leq n, j \in \{0, 1\}$ , points in  $\text{pointsOr}_{i,j}$  can not be covered with less  
 348 than 4 segments from  $\text{segmentsOr}_{i,j}$ .

349 *Proof.* TODO  $\square$

### 350 3.2.2.3. CLAUSE-gadget

351 CLAUSE-gadget is responsible for calculating if variables values assigned in variable gadgets  
 352 satisfies the respective clause in CNF. It has minimal solution of weight  $w$  the clause is  
 353 satisfied, ie. at least one of the respective variables is assigned a correct value. Otherwise it  
 354 has minimal solution  $w + 1$ . This way by the minimal solution for the whole problem, we can  
 355 tell how many clauses were possible to satisfy in the optimum solution of CNF.

356 The CLAUSE-gadgets consist of two OR-gadgets. We do not want the CLAUSE-gadgets to  
 357 be crammed somewhere between the very long variable segments. That is why we have a sim-  
 358 ple gadget to *transfer* whether the segment is in a solution, ie. segments  $(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1})$ .  
 359 These consist of two segments that share point  $x_{i,1}$  and these are the only segments that can  
 360 cover  $x_{i,1}$ . If  $x_{i,0}$  is already covered by some other gadget, we can cover  $x_{i,1}$  by the other  
 361 segment covering another point from the gadget, say  $a$ . If  $x_{i,0}$  is not covered, then the only  
 362 way to cover  $x_{i,0}$  is to use segment  $(x_{i,0}, x_{i,1})$ . This emulates the situation as if  $a$  was placed  
 363 on the segment that  $x_{i,0}$  lays on, but uses one extra segment compared to as  $a = x_{i,0}$  and  $x_{i,1}$   
 364 did not existed.

365 **Points.** TODO: maybe add segments choosing value of  $x, y, z$  to the figure with less opacity

366 TODO: Rephrase it

367 Assuming clause  $C_i = x_i \vee y_i \vee z_i$ , function  $\text{idx}(w)$  is returning index of the variable  $w$ ,  
 368 function  $\text{neg}(w)$  is returning whether variable  $w$  is negated in a clause.

$$\begin{aligned} x_{i,0} &= (10i + 1, 4 \cdot \text{idx}(x_i) + 2 \cdot \text{neg}(x_i)) & x_{i,1} &= (10i + 1, 4n) \\ y_{i,0} &= (10i + 2, 4 \cdot \text{idx}(y_i) + 2 \cdot \text{neg}(y_i)) & y_{i,1} &= (10i + 2, 4n + 4) \\ z_{i,0} &= (10i + 3, 4 \cdot \text{idx}(z_i) + 2 \cdot \text{neg}(z_i)) & z_{i,1} &= (10i + 3, 4n + 6) \end{aligned}$$

$$\text{moveVariable}_i = \{x_{i,j} : j \in \{0, 1\}\} \cup \{y_{i,j} : j \in \{0, 1\}\} \cup \{z_{i,j} : j \in \{0, 1\}\}$$

$$\text{pointsClause}_i = \text{moveVariable}_i \cup \text{pointsOr}_{i,0} \cup \text{pointsOr}_{i,1} \cup \{v_{i,1}\}$$




Figure 3.3: **CLAUSE-gadget**. This figure presents CLAUSE-gadget. Every green rectangle is an OR-gadget.  $y$ -coordinates of  $x_{i,0}$ ,  $y_{i,0}$  and  $z_{i,0}$  depend on the values of variables in the  $i$ -th clause.

### Segments.

$$\begin{aligned} \text{segmentsClause}_i &= \{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1}), (x_{i,1}, l_{i,0}), (y_{i,1}, p_{i,0}), (z_{i,1}, p_{i,1}), \} \cup \\ &\cup \text{segmentsOr}_{i,0} \cup \text{segmentsOr}_{i,1} \end{aligned}$$

370 **Lemma 11.** For any  $1 \leq i \leq n$  and  $a \in \{x_{i,0}, y_{i,0}, z_{i,0}\}$ , there is  $\text{solClause}_i^{\text{true},a} \subset \text{segmentsClause}_i$   
 371 with  $|\text{solClause}_i^{\text{true},a}| = 11$  that covers points in  $\text{pointsClause}_i - \{a\}$ .

372 *Proof.* For  $a = x_{i,0}$  (analogous proof for  $y_{i,0}$ ): First we use Lemma 7 twice with excluded  
 373  $x = l_{i,0}$  and  $x = l_{i,1} = v_{i,0}$ , resulting with 8 segments  $\text{chooseOr}_{i,0}^{\text{true}} \cup \text{chooseOr}_{i,1}^{\text{true}}$  which cover  
 374 all required points apart from  $x_{i,1}, y_{i,0}, y_{i,1}, z_{i,0}, z_{i,1}, l_{i,0}$ . We cover those using additional 3  
 375 segments:  $\{(x_{i,1}, l_{i,0}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1})\}$

376 For  $a = z_{i,0}$ : Using Lemma 8 and Lemma 7 with  $x = p_{i,1}$ , resulting with 8 segments  
 377  $\text{chooseOr}_{i,0}^{\text{false}} \cup \text{chooseOr}_{i,1}^{\text{true}}$  which cover all required points apart from  $x_{i,0}, x_{i,1}, y_{i,0}, y_{i,1}, z_{i,1}, p_{i,1}$ .  
 378 We cover those using additional 3 segments:  $\{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,1}, p_{i,1})\}$ .  $\square$

379 **Lemma 12.** For any  $1 \leq i \leq n$  there is  $\text{solClause}_i^{\text{false}} \subset \text{segmentsClause}_i$  with  $|\text{solClause}_i^{\text{false}}| =$   
 380 12 that covers points in  $\text{pointsClause}_i$ .

381 *Proof.* Using Lemma 8 twice we can cover  $\text{pointsOr}_{i,0}$  and  $\text{pointsOr}_{i,1}$  with 8 segments.

382 To cover the remaining points we additionally use:  $\{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1}), (t_{i,1}, v_{i,1})\}$   
 383  $\square$

384 **Lemma 13.** For any  $1 \leq i \leq n$ :

385 (1) points in  $\text{pointsClause}_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$  can not be covered using any subset of segments  
 386 from  $\text{segmentsClause}_i$  of size smaller than 11;

387 (2) points in  $\text{pointsClause}_i$  can not be covered using any subset of segments from  $\text{segmentsClause}_i$   
 388 of size smaller than 12.

389 *Proof of no cover with fewer than 12 segments.* There is independent set of 12 points in  $\text{pointsClause}_i \supseteq$   
 390  $\{x_{i,0}, y_{i,0}, z_{i,0}, l_{i,0}, p_{i,0}, q_{i,0}, u_{i,0}, v_{i,0} = l_{i,1}, p_{i,1}, q_{i,1}, u_{i,1}, v_{i,1}\}$ .  $\square$

391 *Proof of no cover with fewer than 11 segments.* We can choose disjoint sets  $X, Y, Z$  such that  
 392  $X \cup Y \cup Z \subseteq \text{pointsClause}_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$  and there are no segments covering points from  
 393 different sets. And we prove lower bounds for each of these sets.

$$X = \{x_{i,1}, y_{i,1}, z_{i,1}\}$$

394 Set  $X$  is an indendent set, so it must be covered with 3 segments.

$$Y = \text{pointsOr}_{i,0} - \{l_{i,0}, p_{i,0}\}$$

$$Z = \text{pointsOr}_{i,1} - \{l_{i,1}, p_{i,1}\}$$

395 For both  $Y$  and  $Z$  we can check all of the subsets of 3 segments with brutforce that none  
 396 of them cover, so they have to be covered with 4 segments.

397 TODO: Funny fact, neither  $Y$  nor  $Z$  doesn't have independent set of size 4.

398 Therefore  $\text{pointsClause}_i$  must be covered with at least  $3 + 4 + 4 = 11$  segments.  $\square$

#### 399 3.2.2.4. Summary

400 Add some smart lemmas that sets will be exclusive to each other.

401 **Lemma 14. Robustness to 1/2-extensions.** For every segment  $s \in \mathcal{P}$ ,  $s$  and  $s^{+1/2}$  cover  
 402 the same points from  $\mathcal{C}$ .

403 *Proof.* We can just check every segment. Most of the segments  $s$  are collinear only with points  
 404 that lay on  $s$ , so trivially  $s^{+1/2}$  cannot cover more points than  $s$  does.

405 TODO: list problematic segments here  $\square$

#### 406 3.2.2.5. Summary of construction

We define:

$$\mathcal{C} := \bigcup_{1 \leq i \leq n} \text{pointsVariable}_i \cup \text{pointsClause}_i$$

$$\mathcal{P} := \bigcup_{1 \leq i \leq n} \text{segmentsVariable}_i \cup \text{segmentsClause}_i$$

407 The subsequent sections define these sets.

408 We prove some properties of different gadgets. Every segment for a gadget will only cover  
 409 points in this gadget (won't interact with any diferent gadget), so we can prove lemmas *locally*.

410 TODO:  $y$  axis is increasing values downward on figures (not upwards like in normal).




Figure 3.4: **General schema.**

General layout of VARIABLE-gadget and CLAUSE-gadget and how they interact with each other.

TODO: Rename Choose X to VARIABLE-gadget and Clause C to CLAUSE-gadget.

### 3.2.3. Construction lemmas and proof of Lemma 3

In order to prove Lemma 3 we introduce several auxiliary lemmas proving properties of the construction described in the previous section.

Consider an instance  $S$  of MAX-(3,3)-SAT of size  $n$  with optimum solution satisfying  $k$  clauses. Let us construct an instance  $(\mathcal{C}, \mathcal{P})$  of geometric set cover as described in Section 3.2.2 for instance  $S$  of MAX-(3,3)-SAT.

**Lemma 15.** *Instance  $(\mathcal{C}, \mathcal{P})$  of geometric set cover admits a solution of size  $15n - k$ .*

*Proof.* Let the clauses in  $S$  be  $c_1, c_2 \dots c_n$  and the variables be  $x_1, x_2 \dots x_n$ . Let the assignment of the variables in the optimum solution to  $S$  be  $\phi : \{x_1, x_2 \dots x_n\} \rightarrow \{\mathbf{true}, \mathbf{false}\}$ .

We cover every VARIABLE-gadget with solution described in Lemma 4, in the  $i$ -th gadget choosing the set of segments corresponding to the value of  $\phi(x_i)$ .

For every clause that is satisfied, say  $c_i$ , let us name the variable that is **true** in it as  $x_i$  and point corresponding to  $x_i$  in **pointsClause<sub>i</sub>** as  $a$ . Points in **pointsClause<sub>i</sub>** are covered with set **solClause<sub>i</sub><sup>true,a</sup>** described in Lemma 11. For every clause that is not satisfied, say  $c_j$ , points in **pointsClause<sub>j</sub>** are covered with set **solClause<sub>j</sub><sup>false</sup>** described in Lemma 12.

Formally we define sets responsible for choosing variable and satisfying the variable,  $R_i$  and  $C_i$  respectively, as following:

$$\begin{aligned} R_i &= \begin{cases} \mathbf{chooseVariable}_i^{\mathbf{true}} & \text{if } \phi(x_i) = \mathbf{true} \\ \mathbf{chooseVariable}_i^{\mathbf{false}} & \text{if } \phi(x_i) = \mathbf{false} \end{cases} \\ C_i &= \begin{cases} \mathbf{solClause}_i^{\mathbf{true},a} & \text{if } c_i \text{ satisfied} \\ \mathbf{solClause}_i^{\mathbf{false}} & \text{if } c_i \text{ not satisfied} \end{cases} \\ \mathcal{R} &= \bigcup_{i=1}^n \{R_i \cup C_i : 1 \leq i \leq n\}. \end{aligned}$$

This set covers all the points from  $\mathcal{C}$ , because the sets  $R_i, C_i$  individually cover their corresponding gadgets, as proved in the respective lemmas.

All of these sets are disjoint, so the size of the obtained solution is:

$$|\mathcal{R}| = \sum_{i=1}^n R_i + \sum_{i=1}^n C_i = 3n + 11k + 12(n - k) = 15n - k.$$

□

**Lemma 16.** *Suppose we have a solution  $\mathcal{R}$  of the instance  $(\mathcal{C}, \mathcal{P})$  of geometric set cover that is of size  $w$ . Then there exists a solution of  $S$  that satisfies at least  $15n - w$  clauses.*

*Proof.* Let the clauses in  $S$  be  $c_1, c_2 \dots c_n$  and the variables be  $x_1, x_2 \dots x_n$ . Given a solution  $\mathcal{R}$  of the instance  $(\mathcal{C}, \mathcal{P})$  of geometric set cover, we construct a solution of  $S$  by constructing an assignment of variables  $\phi : \{x_1, x_2 \dots x_n\} \rightarrow \{\mathbf{true}, \mathbf{false}\}$  that satisfies at least  $15n - w$  clauses in  $S$ .

**Variables** The solution  $\mathcal{R}$  needs to use at least 3 segments to cover **pointsVariable<sub>i</sub>** for each VARIABLE-gadget (Lemma 5). If  $\mathcal{R}$  contains both segments  $(c_i, g_i)$  and  $(f_i, h_i)$ , then it has used at least 4 segments (Lemma 6).

$$\begin{cases} |\text{pointsVariable}_i \cap \mathcal{R}| \geq 4 & \text{if } (c_i, g_i) \in \mathcal{R} \wedge (f_i, h_i) \in \mathcal{R} \\ |\text{pointsVariable}_i \cap \mathcal{R}| \geq 3 & \text{otherwise} \end{cases}$$

441 We will now create a modified solution  $\mathcal{R}'$  such that  $|\mathcal{R}'| = |\mathcal{R}|$  and  $\mathcal{R}'$  is still a correct  
442 solution for  $(\mathcal{C}, \mathcal{P})$ .

443 If  $\mathcal{R}$  contains exactly one of the segments  $(c_i, g_i)$  or  $(f_i, h_i)$ , choose the corresponding  
444 variable value to the constructed assignment  $\phi$ . If  $\mathcal{R}$  contains none of these segments, set the  
445 value to **false**. If  $\mathcal{R}$  contains both segments, assign the value  $\phi(x_i)$  that satisfies most clauses  
446 in which  $x_i$  occurs. Every variable is used by exactly 3 clauses, so one assignment satisfies  
447 at least 2 of them. In this case we remove one of the segments  $(c_i, g_i)$  or  $(f_i, h_i)$ , depending  
448 on the value of  $\phi(x_i)$ , from  $\mathcal{R}'$  and replace it with a segment that covers the point  $v_{j,1}$  of a  
449 CLAUSE-gadget for a clause  $c_j$  that uses  $x_i$  with value  $\neg\phi(x_i)$

Formally, we define the value  $\phi(x_i)$  for the variable  $x_i$  as follows:

$$\begin{cases} \phi(x_i) = \text{majority}(X_i) & \text{if } (c_i, g_i) \in \mathcal{R} \wedge (f_i, h_i) \in \mathcal{R} \\ \phi(x_i) = \text{true} & \text{if } (c_i, g_i) \in \mathcal{R} \\ \phi(x_i) = \text{false} & \text{if } (f_i, h_i) \in \mathcal{R} \\ \phi(x_i) = \text{false} & \text{otherwise} \end{cases} \quad (3.1)$$

450 and  $\mathcal{R}'$  as:

451 TODO: This is not a correct mathematical notation, but I am not sure how to define it  
452 clearly

$$\mathcal{R}' = \mathcal{R} \begin{cases} -\{(f_i, h_i)\} \cup \{(t_{j,1}, v_{j,1})\} & \phi(x_i) = \text{true} \wedge \text{if } (c_i, g_i) \in \mathcal{R} \wedge (f_i, h_i) \in \mathcal{R} \wedge \\ & \text{clause } c_j \text{ is satisfied by } \phi(x_i) = \text{false} \\ -\{(c_i, g_i)\} \cup \{(t_{j,1}, v_{j,1})\} & \phi(x_i) = \text{false} \wedge \text{if } (c_i, g_i) \in \mathcal{R} \wedge (f_i, h_i) \in \mathcal{R} \wedge \\ & \text{clause } c_j \text{ is satisfied by } \phi(x_i) = \text{true}, \end{cases}$$

453 therefore  $|\text{pointsVariable}_i \cap \mathcal{R}'| \geq 3$  for every  $i$ .

454 **Clauses** For a clause  $C_i = x \vee y \vee z$ ,  $\mathcal{R}'$  needs to use at least 11 segments to cover  
455  $\text{pointsClause}_i - \{x, y, z\}$  in CLAUSE-gadget (Lemma 13).

456 TODO: maybe put something with cases and names of sets as above

457 Moreover, if all of the points  $\{x_{i,0}, y_{i,0}, z_{i,0}\}$  are not covered by the segments from  $\mathcal{R}' \cap \text{pointsVariable}_i$ ,  
458 then  $\mathcal{R}'$  needs to cover  $\text{pointsClause}_i$  with at least 12 segments by Lemma 13.

TODO: Maybe remove section below, because we do this calculation at the end anyway  
We covered CLAUSE-gadget with at least 11 or at least 12 segments:

$$|\bigcup_{i=1}^n \text{segmentsClause}_i \cap \mathcal{R}'| \geq 11n + a$$

459 where  $a$  is the number of clauses where none of the points  $x_{i,0}, y_{i,0}, z_{i,0}$  were covered by  
460  $\mathcal{R}' \cap \text{segmentsVariable}_j$  for their respective variable  $x_j$ .

**Satisfied clauses with chosen variable assignment.** Consider a clause, say  $c_i$ . If none of the points  $x_{i,0}, y_{i,0}, z_{i,0}$  in  $\text{pointsClause}_i$  were covered by segments from  $\mathcal{R}' \cap \text{segmentsVariable}_j$ , this clause is not satisfied by assignment  $\phi$ .

If one of these points is covered by segments from VARIBALE-gadget (TODO better this or  $\mathcal{R}' \cap \text{segmentsVariable}_j$ ), then denote this point as  $t$  and say it corresponds to variable  $x_j$ . Consider the cases of choosing value of  $\phi(x_j)$  in equation (3.1).

If  $\mathcal{R}'$  contains only one of the segments  $(c_j, g_j)$  and  $(f_j, h_j)$ , then the value  $\phi(x_j)$  satisfies  $c_i$ .

If  $\mathcal{R}'$  contains neither  $(c_j, g_j)$  nor  $(f_j, h_j)$ , then it is impossible that  $t$  is covered by segments in  $\mathcal{R} \cap \text{segmentsVariable}_j$ .

$\mathcal{R}'$  was chosen in such a way, that it is impossible that it contains both  $(c_i, g_i)$  and  $(f_i, h_i)$  for some variable  $i$ .

This means that  $\phi$  satisfies all but at most  $a$  clauses in  $S$ .

To conclude, we proved that given a solution of  $(\mathcal{C}, \mathcal{P})$  of size  $w$ , we have constructed a variables assignment  $\phi$  that satisfies at least  $n - a$  clauses of  $S$ . Finally, note that

$$w \geq 3n + 11(n - a) + 12a = 3n + 11n + a = 14n + a,$$

hence

$$15n - w \leq 15n - 14n - a = n - a.$$

So  $\phi$  satisfies at least  $15n - w$  clauses of  $S$ . □

We are ready to conclude the proof of Lemma 3.

*Proof of Lemma 3.* By Lemma 15, we know that there exists a solution to  $(\mathcal{C}, \mathcal{P})$  of size  $15n - k$ , so:

$$\text{opt}((\mathcal{C}, \mathcal{P})) \leq 15n - k.$$

Since the optimum solution of  $S$  satisfies  $k$  clauses, then according to Lemma 16:

$$\text{opt}((\mathcal{C}, \mathcal{P})) \geq 15n - k.$$

Therefore, the solution given by Lemma 15 of size  $15n - k$  is an optimum solution to the instance  $(\mathcal{C}, \mathcal{P})$ . □

### 3.3. FPT for weighted segments with $\delta$ -extensions

TODO: Some intro

**Theorem 4** (FPT for weighted segment cover with  $\delta$ -extensions). *There exists an algorithm that given a family  $\mathcal{P}$  of  $n$  weighted segments (in any direction), a set of  $m$  points  $\mathcal{C}$ , and parameters  $k$  and  $\delta > 0$ , runs in time  $f(k, \delta) \cdot (nm)^c$  for some computable function  $f$  and a constant  $c$ , and outputs a set  $\mathcal{R} \subseteq \mathcal{P}$  such that  $|\mathcal{R}| \leq k$  and  $\mathcal{R}^{+\delta}$  covers all points in  $\mathcal{C}$ , or determines that such a set  $\mathcal{R}$  does not exist.*

To solve this problem we will introduce a lemma about choosing a *good* subset of points.

TODO: Some intuition

**Definition 9.** For a set of collinear points  $C$ , a subset  $A \subseteq C$  is  $(k, \delta)$ -**good** if for any set of segments  $R$  that covers  $A$  and such that  $|R| \leq k$ , it holds that  $R^{+\delta}$  covers  $C$ .



**Lemma 17.** *There exists an algorithm that for any set of collinear points  $C$ ,  $\delta > 0$  and  $k \geq 1$ , outputs a  $(k, \delta)$ -good set  $A \subseteq C$  of size at most  $f(k, \delta)$  for some computable function  $f$ . This algorithm runs in time  $O(|C| \cdot f(k, \delta))$ .*

*Proof.* We prove this for a fixed  $\delta$  by induction over  $k$ .

**Inductive hypothesis.** For any set of collinear points  $C$ , there exists an algorithm that runs in time  $O(|C|k(1 + \frac{1}{\delta}))$  and finds a set  $A$  such that:

- $A$  is  $(\ell, \delta)$ -good for every  $1 \leq \ell \leq k$ ,
- $A$  has size  $|A| < f(\delta, k)$  for some computable function  $f$ ,
- extreme points from  $C$  are in  $A$ .

**Base case for  $k = 1$ .** It is sufficient that  $A$  consists of 2 points: extreme points from  $C$  or a single point if  $|C| = 1$ .

If they are covered with one segment, it must be a segment that includes the extreme points from  $C$ , so it covers the whole set  $C$ .

**Inductive step.** Assuming inductive hypothesis for any set of collinear points  $C$  and for parameter  $k$ , we will prove hypothesis for  $k + 1$ .

Let be  $s$  the minimal segment that includes all points from  $C$ . That is, the extreme points of  $C$  are endpoints of  $s$ .

We define  $M = \lceil 1 + \frac{2}{\delta} \rceil$  subsegments of  $s$  in the following way. We split  $s$  into  $M$  parts  $v_i$  of equal length, that is  $|v_i| = \frac{|s|}{M}$  for each  $1 \leq i \leq M$ .

Let  $C_i$  be the subset of  $C$  consisting of points laying on  $v_i$ .

Let  $t_i$  be the segment with endpoints being the extreme points of  $C_i$  (it might be degenerated segment if  $|C_i| = 1$  or it might be empty if  $C_i$  is empty).

TODO: Add a picture with  $v_i$  and  $t_i$  here

We use the inductive hypothesis to choose  $(k, \delta)$ -good sets  $A_i$  for sets  $C_i$ . Note that if  $|C_i| \leq 1$ , then  $A_i = C_i$  and it's still a  $(k, \delta)$ -good set for  $C_i$ .

Then we define  $A = \bigcup_{i=1}^M A_i$ . Thus  $A$  includes the extreme points of  $C$ , because they are included in the sets  $A_1$  and  $A_M$ .

**Proof that  $A$  is  $(k, \delta)$ -good for  $C$ .** Let us take any cover of  $A$  with  $k + 1$  segments and call it  $\mathcal{R}$ .

For every segment  $t_i$ , if there exists a segment  $x$  in  $\mathcal{R}$  that is disjoint with  $t_i$ , then we have a cover of  $A_i$  with at most  $k$  segments using  $\mathcal{R} - \{x\}$ . Since  $A_i$  is  $(k, \delta)$ -good for  $t_i$  and  $C_i$ , then  $(\mathcal{R} - \{x\})^{+\delta}$  covers  $C_i$ . So  $\mathcal{R}^{+\delta}$  covers  $C_i$  as well.

If there exists a segment  $t_i$  for which a segment  $x$  as defined above does not exist, then all  $k + 1$  segments that cover  $A_i$  intersect with  $t_i$ . (Note: There may exist only one such segment  $t_i$ ). From the inductive hypothesis end points of  $s$  are in  $A_1$  and  $A_M$  respectively, so  $\mathcal{R}$  must cover them. Hence there must exist segments starting in the ends of  $s$  and ending somewhere in  $t_i$ . Let us call these two segments  $y$  and  $z$ . It follows that:  $|y| + |z| + |t_i| \geq |s|$ . Since  $|t_i| \leq |v_i| = \frac{|s|}{M} \leq \frac{|s|}{1 + \frac{2}{\delta}} = \frac{|s|\delta}{\delta + 2}$ , we have  $\max(|y|, |z|) \geq |s|(1 - \frac{\delta}{\delta + 2})/2 = \frac{|s|}{\delta + 2}$ .

TODO: Add a picture with such segments here

After  $\delta$ -extension, the longer of these segments will lengthen both ways by at least:

$$\frac{|s|\delta}{\delta+2} = \frac{|s|}{1+\frac{2}{\delta}} > \frac{|s|}{M} = v_i > t_i.$$

Therefore the longer of segments  $y$  and  $z$  will cover the segment  $t_i$  after  $\delta$ -extension, therefore  $\mathcal{R}^{+\delta}$  covers  $C_i$ .

Since  $C = \bigcup_{i=1}^M C_i$ , then  $\mathcal{R}^{+\delta}$  covers  $C$ .

**Complexity** We use the recursive algorithm for subsets  $C_i$ . Every point from  $C$  belongs to at most 2 sets  $C_i$ .

Apart from recursive algorithm we perform operations linear in size of  $|C|+M$  to calculate the sets  $C_i$ .

Therefore it has complexity:

$$O(|C|+M) + \sum_i^M O(|C_i|k(1+\frac{1}{\delta})) = O(|C|+(1+\frac{1}{\delta})) + O((\sum_i^M |C_i|)k(1+\frac{1}{\delta})) \leq O(|C|k(1+\frac{1}{\delta})).$$

□

*Proof of Theorem 4.* To construct an algorithm for this problem let us formulate some claims about the problem first.

**Definition 10.** Line is **long** if there are at least  $k+1$  points from  $\mathcal{C}$  on it.

**Claim 1.** *If there are more than  $k$  long lines, then  $\mathcal{C}$  can not be covered with  $k$  segments.*

**Claim 2.** *If there is more than  $k^2$  points from  $\mathcal{C}$  that do not lie on any long line, then  $\mathcal{C}$  can not be covered with  $k$  segments.*

Applying the above claims, if we have more than  $k$  long lines or more than  $k^2$  points from  $\mathcal{C}$  that do not lie on any long line, then we answer that there is no solution of size at most  $k$ .

Otherwise, we can split  $\mathcal{C}$  into at most  $k+1$  sets:  $D$ , at most  $k^2$  points that do not lie on any long line and  $C_i$  – points that lay on  $i$ -th long line. Sets  $C_i$  do not need to be disjoint.

Then for every set  $C_i$ , we can use Lemma 17 to get  $(k, \delta)$ -good set  $A_i$  for  $C_i$ .

Then we have set  $D \cup \bigcup A_i$  of size at most  $f(k, \delta)$  for some computable function  $f$ , that if we have a solution  $\mathcal{R}$  of size at most  $k$  that covers  $D \cup \bigcup A_i$ , then  $\mathcal{R}^{+\delta}$  covers  $\mathcal{C}$ . This is because  $\mathcal{R}$  already covers points  $D$ , they cover  $C_i$ , because they cover  $(k, \delta)$ -good set  $A_i$  with at most  $k$  segments, so  $\mathcal{R}^{+\delta}$  covers  $C_i$ .

After that we shrunk down size of  $\mathcal{C}$  to size of  $f(k, \delta)$  for some computable function  $f$ . Then we would like to shrink down size of  $\mathcal{P}$ . For every collinear subset of  $D$ , we can choose one segment from  $\mathcal{P}$  that covers these points and have the lowest weight or decide there is no segment that cover them. There are at most  $|D|^2$  different segments, because we can distinguish these collinear sets by their extreme points.

This has complexity  $O(|D|^2|\mathcal{P}|)$  and produce shrunk down set  $\mathcal{P}$  of size  $f(k, \delta)$  for some computable function  $f$ .

Then we can iterate over all subsets of shrunk down set  $\mathcal{P}$  and choose the set with the lowest sum of weights that cover  $D$ . This solution would have weight not larger than optimal solution for the problem without extension, because we iterate over all possibilities of covering the subset of  $\mathcal{C}$ .

□

### 565 3.4. $W[1]$ -completeness for weighted segments in 3 directions

566 **Theorem 5.**  *$W[1]$ -completeness for weighted segments in 3 directions.* Consider the  
 567 problem of covering a set  $\mathcal{C}$  of points by selecting  $k$  axis-parallel or right-diagonal weighted  
 568 segments with weights from a set  $\mathcal{P}$  with minimal weight. Assuming ETH, there is no algorithm  
 569 for this problem with running time  $f(k) \cdot (|\mathcal{C}| + |\mathcal{P}|)^{o(\sqrt{k})}$  for any computable function  $f$ .

570 We will show reduction from grid tiling problem.

571 Let's have an instance of grid tiling problem – size of the grid  $k$ , number of elements  
 572 available  $n$  and  $k^2$  sets of available pairs in every tile  $S_{i,j} \subseteq \{1, n\} \times \{1, n\}$ .

573 **Construction.** We construct a set  $\mathcal{P}$  of segments and a set  $\mathcal{C}$  of points.

574 First let's choose any ordering of  $n^2$  elements  $\{1, n\} \times \{1, n\}$  and name this sequence  
 575  $a_1 \dots a_{n^2}$ .

$$match_v(i, j) \iff a_i = \{x_i, y_i\} \wedge a_j = \{x_j, y_j\} \wedge x_i = x_j$$

$$match_h(i, j) \iff a_i = \{x_i, y_i\} \wedge a_j = \{x_j, y_j\} \wedge y_i = y_j$$

**Points.** Define points:

$$h_{i,j,t} = (j \cdot (n^2 + 1) + t, (n^2 + 1) \cdot i)$$

$$v_{i,j,t} = ((n^2 + 1) \cdot i, j \cdot (n^2 + 1) + t)$$

Let's define sets  $H$  and  $V$  as:

$$H = \{h_{i,j,t} : 1 \leq i, j, \leq k, 1 \leq t \leq n^2\}$$

$$V = \{v_{i,j,t} : 1 \leq i, j, \leq k, 1 \leq t \leq n^2\}$$

576 Let's define  $\epsilon = 0.1$ . For a point  $\{x, y\} = p$  we define points  $p^L = \{x - \epsilon, y\}$ ,  $p^R = \{x + \epsilon, y\}$ ,  
 577  $p^U = \{x, y - \epsilon\}$ , and  $p^D = \{x, y + \epsilon\}$ .

Then we define:

$$\mathcal{C} := H \cup \{p^L : p \in H\} \cup \{p^R : p \in H\} \cup V \cup \{p^U : p \in V\} \cup \{p^D : p \in V\}$$

578 **Segments.** Define horizontal segments.

$$hor_{i,j,t_1,t_2} = (h_{i,j,t_1}^R, h_{i,j+1,t_2}^L)$$

$$ver_{i,j,t_1,t_2} = (v_{i,j,t_1}^D, v_{i,j+1,t_2}^U)$$

$$hor_{beg_{i,t}} = (h_{i,1,1}^L, h_{i,1,t}^L)$$

$$hor_{end_{i,t}} = (h_{i,n,t}^R, h_{i,n,n^2}^R)$$

$$ver_{beg_{i,t}} = (v_{i,1,1}^U, v_{i,1,t}^U)$$

$$ver_{end_{i,t}} = (v_{i,n,t}^D, v_{i,n,n^2}^D)$$

$$\begin{aligned}
HOR &= \{hor_{i,j,t_1,t_2} : 1 \leq i \leq k, 1 \leq j < k, 1 \leq t_1, t_2 \leq n^2, match_h(t_1, t_2)\} \\
&\cup \{horbeg_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\} \\
&\cup \{horend_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\}
\end{aligned}$$

$$\begin{aligned}
VER &= \{ver_{i,j,t_1,t_2} : 1 \leq i \leq k, 1 \leq j < k, 1 \leq t_1, t_2 \leq n^2, match_v(t_1, t_2)\} \\
&\cup \{verbeg_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\} \\
&\cup \{verend_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\}
\end{aligned}$$

$$DIAG := \{(h_{i,j,t}, v_{j,i,t}) : 1 \leq i, j \leq k, 1 \leq t \leq n^2, a_t \in S_{i,j}\}$$

579 TODO: explain that these segments are in fact diagonal

$$\mathcal{P} := HOR \cup VER \cup DIAG$$

580 **Lemma 18.** *If there exists solution for grid tiling, then there exists solution for our construc-*  
581 *tion using  $2(k+1)k + k^2$  segments with weight exactly  $2k \cdot (k(n^2 + 1) - 2 - 2\epsilon(k-1))$ .*

**Claim 3.** *If there exists a solution to the grid tiling  $c_1 \dots c_k$  and  $r_1 \dots r_k$ , then there exists a solution covering all points*

$$\{h_{i,j,t} : 1 \leq i, j \leq k, t = (c_i, r_j)\} \cup \{v_{i,j,t} : 1 \leq i, j \leq k, t = (c_j, r_i)\}$$

582 *with segments in DIAG and the rest in VER or HOR and has weight  $2k \cdot (k(n^2 + 1) -$*   
583  *$2 - 2\epsilon(k-1))$ .*

584 **Proof.** TODO: jakiś prosty z definicji

585 **Lemma 19.** *If there exists solution for our construction using  $2(k+1)k + k^2$  segments with*  
586 *weight exactly  $2k \cdot (k(n^2 + 1) - 2 - 2\epsilon(k-1))$ , then there exists a solution for grid tiling*

587 **Proof.** This follows from Lemma 20, because we just take which points are covered with  
588 *DIAG.*

589 **Claim 4.** *Points  $p^L, p^R, p^U, p^D$  cannot be covered with DIAG.*

590 **Claim 5.** *Points in  $H \cup \{p^L : p \in H\} \cup \{p^R : p \in H\}$  cannot be covered with VER.*

591 *Points in  $V \cup \{p^U : p \in V\} \cup \{p^D : p \in V\}$  cannot be covered with HOR.*

592 **Claim 6.** *For given  $i, j$  if none of the points  $h_{i,j,t}$  ( $v_{i,j,t}$ ) for  $1 \leq t \leq n^2$  are covered with*  
593 *DIAG, then some spaces between neighbouring points were covered twice.*

594 **Claim 7.** *For given  $i, j$  two points  $h_{i,j,t_1}, h_{i,j,t_2}$  ( $v_{i,j,t_1}, v_{i,j,t_2}$ ) for  $1 \leq t_1 < t_2 \leq n^2$  are covered*  
595 *with DIAG, then one of them had to be also covered with a segment from HOR (VER).*

596 **Proof.** Point  $v_{i,j,t_2}^L$  had to be covered with VER from Claims 4 and 5. And every segment  
597 *in VER covering  $v_{i,j,t_2}^L$ , covers also  $v_{i,j,t_1}^L$ .*

598 **Lemma 20.** *If there exists solution for our construction with weight at most (exactly)  $2k \cdot$*   
599  *$(k(n^2 + 1) - 2 - 2\epsilon(k-1))$ , then for every  $i, j$  there must be exactly one  $t$  such that  $h_{i,j,t}$  ( $v_{i,j,t}$ )*  
600 *is covered with DIAG and moreover if  $h_{i,j,t_1}$  and  $h_{i,j+1,t_2}$  are uncovered, then  $match_h(t_1, t_2)$ .*  
601 *Analogically for  $v$ .*

602 **Proof.** Only  $k^2$  points can be covered only in *DIAG*, the rest has to be covered with  
603 *VER*  $\cup$  *HOR*. Therefore every result must be at least *ALL\_LINES* -  $2k^2\epsilon$ , because only  
604  $2k^2$  spaces of length  $\epsilon$  can be uncovered in this axis.

605 Of course if  $h_{i,j,t_1}$  and  $h_{i,j+1,t_2}$  are uncovered, then there must exist a segment in *HOR*  
606 between  $h_{i,j,t_1}^R$  and  $h_{i,j+1,t_2}^L$ , so  $math_h(t_1, t_2)$  must be true.

### 607 3.5. What is missing

608 We don't know FPT for axis-parallel segments without  $\delta$ -extensions.



## Chapter 4

# Geometric Set Cover with lines

### 4.1. Lines parallel to one of the axis

When  $\mathcal{R}$  consists only of lines parallel to one of the axis, the problem can be solved in polynomial time.

We create bipartial graph  $G$  with node for every line on the input split into sets:  $H$  – horizontal lines and  $V$  – vertical lines. If any two lines cover the same point from  $\mathcal{C}$ , then we add edge between them.

Of course there will be no edges between nodes inside  $H$ , because all of them are pararell and if they share one point, they are the same lines. Similar argument for  $V$ . So the graph is bipartial.

Now Geometric Set Cover can be solved with Vertex Cover on graph  $G$ . Since Vertex Cover (even in weighted setting) on bipartial graphs can be solved in polynomial time.

Short note for myself just to remember how to this in polynomial time:

Non-weighted setting - Konig theorem + max matching

Weighted setting - Min cut in graph of  $\neg A$  or  $\neg B$  (edges directed from  $V$  to  $H$ )

### 4.2. FPT for arbitrary lines

You can find this is Platypus book. We will show FPT kernel of size at most  $k^2$ .

(Maybe we need to reduce lines with one point/points with one line).

For every line if there is more than  $k$  points on it, you have to take it. At the end, if there is more than  $k^2$  points, return NO. Otherwise there is no more than  $k^4$  lines.

In weighted settings among the same lines with different weights you leave the cheapest one and use the same algorithm.

### 4.3. APX-completeness for arbitrary lines

We will show a reduction from Vertex Cover problem. Let's take an instance of the Vertex Cover problem for graph  $G$ . We will create a set of  $|V(G)|$  pairwise non-pararell lines, such that no three of them share a common point.

Then for every edge in  $(v, w) \in E(G)$  we put a point on crossing of lines for vertices  $v$  and  $w$ . They are not pararell, so there exists exactly one such point and any other line don't cover this point (any three of them don't cross in the same point).

Solution of Geometric Set Cover for this instance would yield a sound solution of Vertex Cover for graph  $G$ . For every point (edge) we need to choose at least one of lines (vertices)  $v$  or  $w$  to cover this point.

Vertex Cover for arbitrary graph is APX-complete, so this problem is also APX-complete.

#### 4.4. 2-approximation for arbitrary lines

Vertex Cover has an easy 2-approximation algorithm, but here very many lines can cross through the same point, so we can do  $d$ -approximation, where  $d$  is the biggest number of lines crossing through the same point. So for set where any 3 lines don't cross in the same point it yields 2-approximation.

The problematic cases are where through all points cross at least  $k$  points and all lines have at least  $k$  points on them. It can be created by casting  $k$ -grid in  $k$ -D space on 2D space.

Greedy algorithm yields  $\log |\mathcal{R}|$ -approximation, but I have example for this for bipartial graph and reduction with taking all lines crossing through some point (if there are no more than  $k$ ) would solve this case. So maybe it works.

Unfortunately I haven't done this :(

I can link some papers telling it's hard to do.

#### 4.5. Connection with general set cover

Problem with finite set of lines with more dimensions is equivalent to problem in 2D, because we can project lines on the plane which is not perpendicular to any plane created by pairs of (point from  $\mathcal{C}$ , line from  $\mathcal{P}$ ).

Of course every two lines have at most one common point, so is every family of sets that have at most one point in common equivalent to some geometric set cover with lines?

No, because of Desargues's theorem. Have to write down exactly what configuration is banned.



## Chapter 5

# Geometric Set Cover with polygons

### 5.1. State of the art

Covering points with weighted discs admits PTAS [Li and Jin, 2015] and with fat polygons with  $\delta$ -extensions with unit weights admits EPTAS [Har-Peled and Lee, 2009].

Although with thin objects, even if we allow  $\delta$ -expansion, the Set Cover with rectangles is APX-complete (for  $\delta = 1/2$ ), it follows from APX-completeness for segments with  $\delta$ -expansion in Section 3.2.

Covering points with squares is W[1]-hard [Marx, 2005]. It can be proven that assuming *SETH*, there is no  $f(k) \cdot (|\mathcal{C}| + |\mathcal{P}|)^{k-\epsilon}$  time algorithm for any computable function  $f$  and  $\epsilon > 0$  that decides if there are  $k$  polygons in  $\mathcal{P}$  that together cover  $\mathcal{C}$ , *Theorem 1.9* in [Marx and Pilipczuk, 2015].



<sup>675</sup> Chapter 6

<sup>676</sup> Conclusions



## 677 Bibliography

- 678 [Har-Peled and Lee, 2009] Har-Peled, S. and Lee, M. (2009). Weighted geometric set cover  
679 problems revisited. *Journal of Computational Geometry*, 3.
- 680 [Håstad, 2001] Håstad, J. (2001). Some optimal inapproximability results. *J. ACM*,  
681 48(4):798–859.
- 682 [Li and Jin, 2015] Li, J. and Jin, Y. (2015). A PTAS for the weighted unit disk cover problem.  
683 *CoRR*, abs/1502.04918.
- 684 [Marx, 2005] Marx, D. (2005). Efficient approximation schemes for geometric problems? In  
685 Brodal, G. S. and Leonardi, S., editors, *Algorithms – ESA 2005*, pages 448–459, Berlin,  
686 Heidelberg. Springer Berlin Heidelberg.
- 687 [Marx and Pilipczuk, 2015] Marx, D. and Pilipczuk, M. (2015). Optimal parameterized algo-  
688 rithms for planar facility location problems using voronoi diagrams. *CoRR*, abs/1504.05476.