

1

University of Warsaw

2

Faculty of Mathematics, Informatics and Mechanics

3

Katarzyna Kowalska

Student no. 371053

4

5

Approximation and Parametrized Algorithms for Segment Set Cover

6

Master's thesis

7

in COMPUTER SCIENCE

8

Supervisor:

dr Michał Pilipczuk

Instytut Informatyki

9

June 2020

10 **Supervisor's statement**

11 Hereby I confirm that the presented thesis was prepared under my supervision and
12 that it fulfils the requirements for the degree of Master of Computer Science.

13 Date

Supervisor's signature

14 **Author's statement**

15 Hereby I declare that the presented thesis was prepared by me and none of its contents
16 was obtained by means that are against the law.

17 The thesis has never before been a subject of any procedure of obtaining an academic
18 degree.

19 Moreover, I declare that the present version of the thesis is identical to the attached
20 electronic version.

21 Date

Author's signature

22

Abstract

23 The work presents a study of different geometric set cover problems. It mostly focuses on
24 segment set cover and its connection to the polygon set cover.

25

Keywords

26 set cover, geometric set cover, FPT, $W[1]$ -completeness, APX-completeness, PCP theorem,
27 NP-completeness

28

Thesis domain (Socrates-Erasmus subject area codes)

29 11.3 Informatyka

30

31

Subject classification

32 D. Software

33 D.127. Blabalgorithms

34 D.127.6. Numerical blabalysis

35

Tytuł pracy w języku polskim

36 Algorytmy parametryzowania i trudność aproksymacji problemu pokrywania zbiorów
37 odcinkami na płaszczyźnie

Contents

39	1. Introduction	5
40	2. Definitions	7
41	2.1. Geometric Set Cover	7
42	2.2. Approximation	7
43	2.3. δ -extensions	7
44	3. Geometric Set Cover with segments	9
45	3.1. FPT for segments	9
46	3.1.1. Axis-parallel segments	9
47	3.1.2. Segments in d directions	9
48	3.1.3. Segments in arbitrary direction	9
49	3.2. APX-completeness for segments parallel to axes	11
50	3.2.1. MAX-(3,3)-SAT problem	11
51	3.2.2. Reduction construction	13
52	3.2.2.1. VARIABLE-gadget	13
53	3.2.2.2. OR-gadget	14
54	3.2.2.3. CLAUSE-gadget	15
55	3.2.2.4. Summary	17
56	3.2.3. Summary of construction	17
57	3.2.4. Proofs of construction Lemma 3.2.1	19
58	3.3. Weighted segments	20
59	3.3.1. FPT for weighted segments with δ -extensions	20
60	3.3.2. W[1]-completeness for weighted segments in 3 directions	21
61	3.3.3. What is missing	23
62	4. Geometric Set Cover with lines	25
63	4.1. Lines parallel to one of the axis	25
64	4.2. FPT for arbitrary lines	25
65	4.3. APX-completeness for arbitrary lines	25
66	4.4. 2-approximation for arbitrary lines	26
67	4.5. Connection with general set cover	26
68	5. Geometric Set Cover with polygons	27
69	5.1. State of the art	27
70	6. Conclusions	29

Chapter 1

Introduction

The Set Cover problem is one of the most common NP-complete problems. [tutaj referencja]
We are given a family of sets and have to choose the smallest subfamily of these sets that cover
all their elements. This problem naturally extends to settings where we put different weights
on the sets and look for the subfamily of the minimal weight. This problem is NP-complete
even without weights and if we put restrictions on what the sets can be. One of such variants
is Vertex Cover problem, where sets have size 2 (they are edges in a graph).

In this work we focus on another such variant where the sets correspond to some geometric
shapes and only some points of the plane have to be covered. When these shapes are rectangles
with edges parallel to the axis, the problem can be proven to be W[1]-complete (solution of
size k cannot be found in $n^o(k)$ time), APX-complete (for sufficiently small $\epsilon > 0$, the problem
does not admit $1 + \epsilon$ -approximation scheme) [referencje].

Some of these settings are very easy. Set cover with lines parallel to one of the axis can
be solved in polynomial time.

There is a notion of δ -expansions, which loosen the restrictions on geometric set cover. We
allow the objects to cover the points after δ -expansion and compare the result to the original
setting. This way we can produce both FPT and EPTAS for the rectangle set cover with
 δ -extensions [referencje].

Our contribution. In this work, we prove that unweighted geometric set cover with seg-
ments is fixed parameter tractable (FPT).

Moreover, we show that geometric set cover with segments is APX-complete for unweighted
axis-parallel segments, even with $1/2$ -extensions. So the problem for very thin rectangles
also can't admit PTAS. Therefore, in the efficient polynomial-time approximation scheme
(EPTAS) for *fat polygons* by [Har-Peled and Lee, 2009], the assumption about polygons
being fat is necessary.

Finally, we show that geometric set cover with weighted segments in 3 directions is
W[1]-complete. However, geometric set cover with weighted segments is FPT if we allow
 δ -extension.

This result is especially interesting, since it's counter-intuitive that the unweighted setting
is FPT and the weighted setting is W[1]-complete. Most of such problems (like vertex cover
or [wiecej przykladow]) are equally hard in both weighted and unweighted settings.

103 Chapter 2

104 Definitions

105 2.1. Geometric Set Cover

106 In the geometric set cover problem we are given \mathcal{P} – a set of objects, which are connected
107 subsets of the plane, \mathcal{C} – a set of points in the plane. The task is to choose $\mathcal{R} \subseteq \mathcal{P}$ such that
108 every point in \mathcal{C} is inside some element from \mathcal{R} and $|\mathcal{R}|$ is minimized.

109 In the parametrized setting for a given k , we only look for a solution \mathcal{R} such that $|\mathcal{R}| \leq k$.

110 In the weighted setting, there is some given weight function $f : \mathcal{P} \rightarrow \mathbb{R}^+$, and we would
111 like to find a solution \mathcal{R} that minimizes $\sum_{R \in \mathcal{R}} f(R)$.

112 2.2. Approximation

113 Let us recall some definitions related to optimization problems that are used in the following
114 sections.

115 **Definition 2.2.1.** A **polynomial-time approximation scheme (PTAS)** for a minimiza-
116 tion problem Π is a family of algorithms \mathcal{A}_ϵ for every $\epsilon > 0$ such that \mathcal{A}_ϵ takes an instance I
117 of Π and in polynomial time finds a solution that is within a factor $(1 + \epsilon)$ of being optimal.
118 That means the reported solution has weight at most $(1 + \epsilon)opt(I)$, where $opt(I)$ is the weight
119 of an optimal solution for I .

120 **Definition 2.2.2.** A problem Π is **APX-hard** if assuming $P \neq NP$, there exists $\epsilon > 0$ such
121 that there is no polynomial-time $(1 + \epsilon)$ -approximation algorithm for Π .

122 2.3. δ -extensions

123 TODO PLACEHOLDER for introductory text

124 δ -extensions is one of the modifications to a problem, that makes geometric set cover
125 problem easier, it has been already used in literature (place some refrence here).

126 **Definition 2.3.1** (δ -extensions for center-symmetric objects). For any $\delta > 0$ and a center-
127 symmetric object L with centre of symmetry $S = (x_s, y_s)$, the **δ -extension** of L is the object
128 $L^{+\delta} = \{(1 + \delta) \cdot (x - x_s, y - y_s) + (x_s, y_s) : (x, y) \in L\}$, that is, $L^{+\delta}$ is the image of L under
129 homothety centered at S which scale $(1 + \delta)$

130 The geometric set cover problem with δ -extensions is a modified version of geometric set
131 cover where:

- We need to cover all the points in \mathcal{C} with objects from $\{P^{+\delta} : P \in \mathcal{P}\}$ (which always include no fewer points than the objects before δ -extensions);

- We look for a solution that is no larger than the optimum solution for the original problem. Note that it does not need to be an optimal solution in the modified problem.

Formally we have the following.

Definition 2.3.2 (Geometric set cover problem with δ -extensions). The geometric set cover problem with δ -extensions is the problem where for an input instance $I = (\mathcal{P}, \mathcal{C})$, the task is to output a solution $\mathcal{R} \subseteq \mathcal{P}$, such that the δ -extended set $\{R^{+\delta} : R \in \mathcal{R}\}$ covers \mathcal{C} and is no larger than the optimal solution for the problem without extensions, i.e. $|\mathcal{R}| \leq |\text{opt}(I)|$.

TODO: Some text

Definition 2.3.3 (Geometric set cover PTAS with δ -extensions). We define a PTAS for geometric set cover with δ -extensions as a family of algorithms $\{\mathcal{A}_{\delta, \epsilon}\}_{\delta, \epsilon > 0}$ that each take as an input instance $I = (\mathcal{P}, \mathcal{C})$, and in polynomial time outputs a solution $\mathcal{R} \subseteq \mathcal{P}$ such that the δ -extended set $\{R^{+\delta} : R \in \mathcal{R}\}$ covers \mathcal{C} and is within a $(1 + \epsilon)$ factor of the optimal solution for this problem without extensions, i.e. $(1 + \epsilon)|\mathcal{R}| \leq |\text{opt}(I)|$.

Chapter 3

Geometric Set Cover with segments

3.1. FPT for segments

3.1.1. Axis-parallel segments

You can find this in Platypus book. (TODO add referece)

We show $\mathcal{O}(2^k)$ branching algorithm. Let us take the point K which is the smallest under a lexicographic ordering on coordinates among points that are not covered yet. We need to cover K with some of the remaining segments.

We branch over choice of direction among the 2 axis-parallel directions. In this direction we greedily take the segment that covers the most points. As K was the smallest in lexicographical order, all points in \mathcal{C} colinear with K in both axis-parallel directions are only on one side of K , because their coordinates are larger. Therefore segments covering K in this direction create monotone sequence of sets and we can greedily take one segment that covers superset of all of these segments.

TODO: Maybe split it into theorem + algorithm + explanation like in section 3.1.3

3.1.2. Segments in d directions

The same algorithm as described in the previous section, but we branch over d directions and it runs in complexity $\mathcal{O}(d^k)$.

3.1.3. Segments in arbitrary direction

Theorem 3.1.1. (FPT for segment cover). *There exists an algorithm that given a family \mathcal{P} of n segments (in any direction), a set of m points \mathcal{C} and a parameter k , runs in time $f(k) \cdot (nm)^c$ for some computable function f and constant c , and outputs a subfamily $\mathcal{R} \subseteq \mathcal{P}$ such that $|\mathcal{R}| \leq k$ and \mathcal{R} covers all points in \mathcal{C} or determines that the solution of size at most k doesn't exist.*

This theorem is proved by following lemmas.

Lemma 3.1.1. (Reduction). *Given a family \mathcal{P} of n segments (in any direction) and a set of m points \mathcal{C} for segment cover problem, without a loss of generality we can assume that no segment covers a superset of what another segment covers. That is, for any $A, B \in \mathcal{P}$, we have $A \neq B \Rightarrow A \not\subseteq B$.*

Proof. Trivial. □

Lemma 3.1.2. *Given an instance of a problem, if there exists a line L with at least $k + 1$ points on it, there exists a subset $\mathcal{A} \subseteq \mathcal{P}$, $|\mathcal{A}| \leq k$, such that every solution \mathcal{R} with $|\mathcal{R}| \leq k$ satisfies $|\mathcal{A} \cap \mathcal{R}| \geq 1$.*

Proof. First we use Lemma 3.1.1.

Let us name points from \mathcal{C} that lay on L , x_1, x_2, \dots, x_t in the order they appear on the line.

Every segment that is not colinear with L can cover at most one of these points. Therefore in any solution of size not larger than k , among any k of these points at least one must be covered with segment colinear with L .

Therefore we need to take one of the segments colinear with L that covers any of the points x_1, x_2, \dots, x_k . After using reduction from Lemma 3.1.1, there are at most k such segments that are distinct. □

Proof of theorem 3.1.1.

Algorithm. First we use Lemma 3.1.1.

We present a recursive algorithm. Given an instance of the problem:

- (1) If there exist a line with at least $k + 1$ points, we branch over adding to the solution one of at most k possible segments from Lemma 3.1.2, name this segment S . Then we find a solution \mathcal{R} for problem for points $\mathcal{C} - S$, segments $\mathcal{P} - \{S\}$ and parameter $k - 1$ and return $\mathcal{R} \cup \{S\}$.
- (2) If every line has at most k points on it and $|\mathcal{C}| > k^2$, then answer NO.
- (3) If $|\mathcal{C}| \leq k^2$, solve the problem by brute force algorithm.

Correctness. Lemma 3.1.2 proves that at least one segment that we branch over in (1) must be present in every solution \mathcal{R} with $|\mathcal{R}| \leq k$, therefore the recursive call can find the optimal solution.

In (2) the answer is no, because every line covers no more than k points from \mathcal{C} , which implies that every segment from \mathcal{P} covers at most k . Under this assumption we can cover only k^2 points with a solution of size k , which is less than $|\mathcal{C}|$.

Checking all possible solutions in (3) is trivially correct.

Complexity. In leaves of branching (3) $|\mathcal{C}| < k^2$, so $|\mathcal{P}| < k^4$, because every segments can be uniquely identified by 2 extreme points it covers (by Lemma 3.1.1). Therefore there are $\binom{k^4}{k}$ possible solutions to check, each can be checked in time $O(k|\mathcal{C}|)$. Therefore (3) takes time $O(f(k))$.

In this branching algorithm our parameter k is decreased with every recursive call, so we have at most k levels of recursion with branching over k possibilities. Candidates to branch over can be found on each level in time $O(nm \log(nm))$.

Reduction from Lemma 3.1.1 can be implemented in $O(n^2m)$.

Overall complexity is $O(n^2m + nm \log(nm) \cdot f(k))$ □

216 3.2. APX-completeness for segments parallel to axes

217 In this section we analyze whether there exists an $(1 + \epsilon)$ -approximation scheme for geometric
 218 set cover for rectangles. We show that we can restrict this problem to a very simple setting:
 219 segments parallel to axes and allow $(1/2)$ -extension, and the problem is still APX-hard. Note
 220 that segments are just degenerated rectangles with one side being very narrow.

221 Our results can be summarized in the following theorem and this section aims to prove it.

222 **Theorem 3.2.1.** *(axis-parallel segment set cover with $1/2$ -extension is APX-hard).*
 223 *Unweighted geometric set cover with axis-parallel segments in 2D (even with $1/2$ -extension)*
 224 *is APX-hard. That is, assuming $P \neq NP$, there does not exist a PTAS for this problem.*

225 Theorem 3.2.1 implies the following.

226 **Corollary 3.2.1.** *(rectangle set cover is APX-hard).* *Unweighted geometric set cover*
 227 *with rectangles (even with $1/2$ -extension) is APX-hard.*

228 We prove Theorem 3.2.1 by taking a problem that is APX-hard and showing a reduction.
 229 For this problem we choose MAX-(3,3)-SAT which we define below.

230 Given an instance I of MAX-(3,3)-SAT, we construct an instance J of axis-parallel segment
 231 set cover problem, such that for a sufficiently small $\epsilon > 0$, a polynomial $(1 + \epsilon)$ -approximation
 232 algorithm for J would be able to distinguish whether an instance I of MAX-(3,3)-SAT is fully
 233 satisfiable or $(7/8 + \epsilon)$ -satisfiable. However, according to (Theorem 3.2.2) that problem is
 234 NP-hard. That would imply $P = NP$, contradicting the assumption.

235 3.2.1. MAX-(3,3)-SAT problem

236 **Definition 3.2.1.** MAX-3SAT is a maximization problem. We are given a 3-CNF formula,
 237 and need to find an assignment of variables that satisfies the most clauses.

238 **Definition 3.2.2.** MAX-(3,3)-SAT is MAX-3SAT with an additional restriction that every
 239 variable appears in exactly 3 clauses. Note that thus, the number of clauses is equal to number
 240 of variables.

241 In the lemmas above we use a property of MAX-(3,3)-SAT proved in [Håstad, 2001] and
 242 described in Theorem 3.2.2.

243 **Theorem 3.2.2.** [Håstad, 2001]

244 *For any $\epsilon > 0$, it is NP-hard to distinguish satisfiable (3,3)-SAT formulas from at most*
 245 *$(7/8 + \epsilon)$ -satisfiable (3,3)-SAT formulas.*

246 The following lemma encapsulates the properties of the reduction described in this section,
 247 and it allows us to prove Theorem 3.2.1.

248 **Lemma 3.2.1.** *Given an instance S of MAX-(3,3)-SAT with n variables and optimum value*
 249 *$OPT(S)$, we can construct an instance I of geometric set cover with axis-parallel segments in*
 250 *2D with $1/2$ -extensions, such that:*

251 (1) *For every solution X of instance I , there exists a solution of S of size at least $15n - |X|$.*

252 (2) *For every solution X of instance S , there exists a solution of I of size $15n - |X|$.*

253 (3) *Every solution with $1/2$ -extensions for I is also a solution to the original instance I .*

254 *Therefore, the optimal solution of I is $OPT(I) = 15n - OPT(S)$.*

We prove Lemma 3.2.1 in subsequent sections, but meanwhile let us prove Theorem 3.2.1 using Lemma 3.2.1 and Theorem 3.2.2.

TODO: This below can't use current template

Proof of Theorem 3.2.1. . Consider any $0 < \epsilon < 1/(15 \cdot 8)$.

Let us assume that there exists a polynomial-time $(1 + \epsilon)$ -approximation algorithm for unweighted geometric set cover with axis-parallel segments in 2D with $(1/2)$ -extensions. We construct an algorithm that solves the problem stated in Theorem 3.2.2, thereby proving that $P = NP$.

Take an instance S of MAX-(3,3)-SAT to be distinguished and construct an instance of geometric set cover I using Lemma 3.2.1.

We now use the $(1 + \epsilon)$ -approximation algorithm for geometric set cover on I , denote the cost of the result of applying this algorithm as $approx(I)$.

We prove that if in S one can satisfy at most $(\frac{7}{8} + \epsilon)n$ clauses, then $approx(I) \geq 15n - (\frac{7}{8} + \epsilon)n$ and if S is satisfiable, then $approx(I) < 15n - (\frac{7}{8} + \epsilon)n$.

Assume S satisfiable. From the definition of S being satisfiable, we have:

$$OPT(S) = n.$$

From Lemma 3.2.1 we have:

$$OPT(I) = 14n.$$

Therefore,

$$\begin{aligned} approx(I) &\leq (1 + \epsilon)OPT(I) = 14n(1 + \epsilon) = 14n + 14\epsilon \cdot n = \\ &= 14n + (15\epsilon - \epsilon)n < 14n + \left(\frac{1}{8} - \epsilon\right)n = 15n - \left(\frac{7}{8} + \epsilon\right)n \end{aligned}$$

Assume S is at most $(\frac{7}{8} + \epsilon)$ satisfiable . From the definition of S being at most $(\frac{7}{8} + \epsilon)n$ satisfiable, we have:

$$OPT(S) \leq \left(\frac{7}{8} + \epsilon\right)n$$

From Lemma 3.2.1 we have:

$$OPT(I) \geq 15n - \left(\frac{7}{8} + \epsilon\right)n$$

Since a solution to I with extensions is also a solution without extensions, by Lemma 3.2.1 (3.), we have:

$$approx(I) \geq OPT(I) = 15n - \left(\frac{7}{8} + \epsilon\right)n$$

Therefore, by using the assumed $(1 + \epsilon)$ -approximation algorithm, it is possible to distinguish the case when S is satisfiable from the case when it is at most $(\frac{7}{8} + \epsilon)n$ satisfiable, since there exists a threshold on the approximation result in segment set cover that distinguishes these two cases. Hence, the assumed approximation algorithm cannot exist, unless $P = NP$.

□

277 3.2.2. Reduction construction

278 We show reduction from MAX-(3,3)-SAT problem to geometric set cover with segments parallel to axis. Moreover the instance of geometric set cover will be robust to $1/2$ -extensions (have the same optimal solution after $1/2$ -extension).

281 The construction will be composed of 2 types of gadgets: **VARIABLE-gadgets** and **CLAUSE-gadgets**. **CLAUSE-gadgets** would be constructed using two **OR-gadgets** connected together.

284 3.2.2.1. VARIABLE-gadget

285 VARIABLE-gadget is responsible for choosing the value of a variable in a CNF formula. It allows two minimal solutions and every minimal solution must use exactly one of the (c_i, g_i) and (f_i, h_i) segments. These two choices correspond to the two Boolean values of the variable.

Points. Define points:




Figure 3.1: **VARIABLE-gadget**. We denote the set of points marked with black circles as C_{var}^i , and they need to be covered (are part of the set \mathcal{C}). Note that some of the points are not marked as black dots and exists only to name segments for further reference. We denote the set of red segments as X_{false}^i and the set of green segments as X_{true}^i .

288

289 With $L = 12n$:

$$290 \quad \begin{array}{llll} a = (-L, 0) & b = (-\frac{2}{3}L, 0) & c = (-\frac{1}{3}L, 0) & d = (-L, 1) \\ e = (-\frac{2}{3}L, 1) & f = (-\frac{1}{3}L, 2) & g = (L, 0) & h = (L, 2) \end{array}$$

Let us define:

$$C_{var} = \{a, b, c, d, e, f\}$$

and

$$C_{var}^i = C_{var} + (0, 4i)$$

291 **Segments.** Let us define:

$$X_{true}^i = \{(a_i, d_i), (b_i, f_i), (c_i, g_i)\}$$

$$X_{false}^i = \{(a_i, c_i), (d_i, e_i), (f_i, h_i)\}$$

$$P_{var}^i = X_{true}^i \cup X_{false}^i$$

292 **Lemma 3.2.2.** For any $1 \leq i \leq n$, points C_{var}^i can be covered using 3 segments from P_{var}^i .

293 *Proof.* We can use either set X_{true}^i or X_{false}^i . □

294 **Lemma 3.2.3.** For any $1 \leq i \leq n$, points C_{var}^i can not be covered with less than 3 segments
 295 from P_{var}^i .

296 *Proof.* No segment of P_{var}^i covers more than one point from $\{d_i, f_i, c_i\}$, therefore C_{var}^i can not
 297 be covered with less than 3 segments. \square

298 **Lemma 3.2.4.** If both segments (c_i, g_i) and (f_i, h_i) are chosen, then the covering the remain-
 299 ing points from C_{var}^i requires at least 2 different segments from P_{var}^i .

300 *Proof.* No segment of P_{var}^i covers more than one point from $\{a_i, e_i\}$, therefore $C_{var}^i - \{c_i, f_i, g_i, h_i\}$
 301 can not be covered with less than 2 segments. \square

302 3.2.2.2. OR-gadget

303 OR-gadget has 3 important segments – $x, y, result$. x and y don't count to the weight of
 304 solution of OR-gadget (they are part of different gadgets). It has a minimal solution of weight
 305 w and $result$ can be chosen only if x or y are also chosen for the solution. If none of them
 306 are chosen, then solution choosing $result$ segment has weight at least $w + 1$. Therefore the
 307 following formula holds for a solution R assuming that R uses only w from this OR-gadget:

$$(x \in R) \vee (y \in R) \iff result \in R$$



Figure 3.2: **OR-gadget.** We denote these point as $or_gadget_{i,j}$. We denote set of red segments as $or_{i,j}^{false}$, set of blue segments as $or_{i,j}^{true}$, green and yellow segments as $or_move_variable_{i,j}$.

308 **Points.**

$$\begin{array}{llll}
l_0 = (0, 0) & m_0 = (0, 1) & n_0 = (0, 2) & o_0 = (0, 3) \\
p_0 = (0, 4) & q_0 = (1, 1) & r_0 = (1, 3) & s_0 = (2, 1) \\
t_0 = (2, 2) & u_0 = (2, 3) & v_0 = (3, 2) &
\end{array}$$

$$vec_{i,j} = (10i + 3 + 3j, 4n + 2j)$$

Define $\{l_{i,j}, m_{i,j} \dots v_{i,j}\}$ as $\{l_0, m_0 \dots v_0\}$ shifted by $vec_{i,j}$

Note that $v_{i,0} = l_{i,1}$ (see Figure 3.3)

$$C_or_gadget_{i,j} = \{l_{i,j}, m_{i,j}, n_{i,j}, o_{i,j}, p_{i,j}, q_{i,j}, r_{i,j}, s_{i,j}, t_{i,j}, u_{i,j}\}$$

Segments. We define names subsets of segments, to refer to them in lemmas.

$$or_{i,j}^{false} = \{(q_{i,j}, r_{i,j}), (s_{i,j}, u_{i,j})\}$$

$$or_{i,j}^{true} = \{(m_{i,j}, s_{i,j}), (o_{i,j}, u_{i,j}), (t_{i,j}, v_{i,j})\}$$

$$or_move_variable_{i,j} = \{(l_{i,j}, n_{i,j}), (n_{i,j}, p_{i,j})\}$$

Segments in OR-gadget:

$$P_or_gadget_{i,j} = or_{i,j}^{false} \cup or_{i,j}^{true} \cup or_move_variable_{i,j}$$

Lemma 3.2.5. For any $1 \leq i \leq n, j \in \{0, 1\}$ and $x \in \{l_{i,j}, p_{i,j}\}$ we can cover points in $C_or_gadget_{i,j} - \{x\} \cup \{v_{i,j}\}$ with 4 segments.

Proof. We can do that using one segment from $or_move_variable_{i,j}$ (chosen depending on the value of x) and all segments from $or_{i,j}^{true}$. \square

Lemma 3.2.6. For any $1 \leq i \leq n, j \in \{0, 1\}$, we can cover points in $C_or_gadget_{i,j}$ with 4 segments from $P_or_gadget_{i,j}$.

Proof. We can do that using $or_move_variable_{i,j}$ and $or_{i,j}^{false}$. \square

3.2.2.3. CLAUSE-gadget

CLAUSE-gadget is responsible for calculating if choice of the variable values meets the clause in formula. It has minimal solution of weight w if at least one variable in the clause has a correct value. Otherwise it has minimal solution $w + 1$. This way by the minimal solution for the whole problem, we can tell how many clauses were satisfiable.

The CLAUSE-gadgets consist of two OR-gadgets. We don't want the CLAUSE-gadgets to be crammed somewhere between the very long variable segments. That's why we have a simple gadget to *pass* the value of the segment, ie. segments $(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1})$. Two segments and one of them is chosen if x was chosen in the solution and the other one if x wasn't.



Figure 3.3: **CLAUSE-gadget**. We denote set of these points as C_clause_i . Every green rectangle is an OR-gadget. y -coordinates of $x_{i,0}$, $y_{i,0}$ and $z_{i,0}$ depend on the values of variables in the i -th clause.

331 **Points.** TODO: Rephrase it

332 Assuming clause $C_i = x_i \vee y_i \vee z_i$, function $idx(w)$ is returning index of the variable w ,
 333 function $neg(w)$ is returning whether variable w is negated in a clause.

$$\begin{aligned}
 x_{i,0} &= (10i + 1, 4 \cdot idx(x_i) + 2 \cdot neg(x_i)) & x_{i,1} &= (10i + 1, 4n) \\
 y_{i,0} &= (10i + 2, 4 \cdot idx(y_i) + 2 \cdot neg(y_i)) & y_{i,1} &= (10i + 2, 4n + 4) \\
 z_{i,0} &= (10i + 3, 4 \cdot idx(z_i) + 2 \cdot neg(z_i)) & z_{i,1} &= (10i + 3, 4n + 6)
 \end{aligned}$$

$$move_variable_i = \{x_{i,j} : j \in \{0, 1\}\} \cup \{y_{i,j} : j \in \{0, 1\}\} \cup \{z_{i,j} : j \in \{0, 1\}\}$$

$$C_clause_i = move_variable_i \cup C_or_gadget_{i,0} \cup C_or_gadget_{i,1} \cup \{v_{i,1}\}$$

Segments.

$$\begin{aligned}
 P_clause_i &= \{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1}), (x_{i,1}, l_{i,0}), (y_{i,1}, p_{i,0}), (z_{i,1}, p_{i,1}), \} \cup \\
 &\cup P_or_gadget_{i,0} \cup P_or_gadget_{i,1}
 \end{aligned}$$

335 **Lemma 3.2.7.** For any $1 \leq i \leq n$ and $a \in \{x_{i,0}, y_{i,0}, z_{i,0}\}$, points $C_clause_i - \{a\}$ can be
 336 covered using 11 segments from P_clause_i .

337 *Proof.* For $a = x_{i,0}$ (analogous proof for $y_{i,0}$): First we use Lemma 3.2.5 twice with excluded
 338 $x = l_{i,0}$ and $x = l_{i,1} = v_{i,0}$, resulting with 8 segments $or_{i,0}^{true} \cup or_{i,1}^{true}$ which cover all required

points apart from $x_{i,1}, y_{i,0}, y_{i,1}, z_{i,0}, z_{i,1}, l_{i,0}$. We cover those using additional 3 segments:
 $\{(x_{i,1}, l_{i,0}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1})\}$

For $a = z_{0,i}$: Using Lemma 3.2.6 and Lemma 3.2.5 with $x = p_{i,1}$, resulting with 8 segments
 $or_{i,0}^{false} \cup or_{i,1}^{true}$ which cover all required points apart from $x_{i,0}, x_{i,1}, y_{i,0}, y_{i,1}, z_{i,1}, p_{i,1}$. We cover
those using additional 3 segments: $\{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,1}, p_{i,1})\}$. \square

Lemma 3.2.8. *Points C_clause_i can be covered with 12 segments from P_clause_i .*

Proof. Using Lemma 3.2.6 twice we can cover $or_gadget_{i,0}$ and $or_gadget_{i,1}$ with 8 segments.

To cover the remaining points we additionally use: $\{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1}), (t_{i,1}, v_{i,1})\}$
 \square

Lemma 3.2.9. *For any $1 \leq i \leq n$, points $C_clause_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$ can not be covered using less than 11 segments from P_clause_i .*

All points C_clause_i can not be covered with less than 12 segments from P_clause_i .

Proof of no cover with less than 12 segments. There is independent set of 12 points in $C_clause_i \supseteq$
 $\{x_{i,0}, y_{i,0}, z_{i,0}, l_{i,0}, p_{i,0}, q_{i,0}, u_{i,0}, v_{i,0} = l_{i,1}, p_{i,1}, q_{i,1}, u_{i,1}, v_{i,1}\}$. \square

Proof of no cover with less than 11 segments. We can choose disjoint sets X, Y, Z such that
 $X \cup Y \cup Z \subseteq C_clause_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$ and there are no segments covering points from
different sets. And we prove lower bounds for each of these sets.

$$X = \{x_{i,1}, y_{i,1}, z_{i,1}\}$$

Set X is an indendent set, so it must be covered with 3 segments.

$$Y = or_gadget_{i,0} - \{l_{i,0}, p_{i,0}\}$$

$$Z = or_gadget_{i,1} - \{l_{i,1}, p_{i,1}\}$$

For both Y and Z we can check all of the subsets of 3 segments with brutforce that none
of them cover, so they have to be covered with 4 segments.

TODO: Funny fact, neither Y nor Z doesn't have independent set of size 4.

Therefore C_clause_i must be covered with at least $3 + 4 + 4 = 11$ segments. \square

3.2.2.4. Summary

Add some smart lemmas that sets will be exclusive to each other.

Lemma 3.2.10. Robustness to 1/2-extensions. *For every segment $s \in \mathcal{P}$, s and $s^{+1/2}$ cover the same points from \mathcal{C} .*

3.2.3. Summary of contruction

We define:

$$\mathcal{C} := \bigcup_{1 \leq i \leq n} C_variable_i \cup C_clause_i$$

$$\mathcal{P} := \bigcup_{1 \leq i \leq n} P_variable_i \cup P_clause_i$$

The subsequent sections define these sets.

We prove some properties of different gadgets. Every segment for a gadget will only cover
points in this gadget (won't interact with any diferent gadget), so we can prove lemmas *locally*.

TODO: y axis is increasing values downward on figures (not upwards like in normal).




Figure 3.4: **General schema.**

General layout of VARIABLE-gadget and CLAUSE-gadget and how they interact with each other.

TODO: Rename Choose X to VARIABLE-gadget and Clause C to CLAUSE-gadget.

370 3.2.4. Proofs of construction Lemma 3.2.1

371 **Lemma 3.2.11.** *Given an instance of MAX-(3,3)-SAT of size n with optimal solution k . For*
 372 *instance of geometric cover, constructed according to Lemma 3.2.1, there exists a solution of*
 373 *weight $15n - k$.*

374 *Proof.* Let's name the assignments of the variables in MAX-(3,3)-SAT instance, that achieve
 375 the optimal solution, $y_1, y_2 \dots y_n$, Let's cover every VARIABLE-gadget with solution de-
 376 scribed in Lemma 3.2.2, in the i -th gadget choosing the set of segments responsible for the
 377 value of y_i (true - x_i^{true} or false - x_i^{false}).

378 Cover every satisfied CLAUSE-gadget with solution described in Lemma 3.2.7 and unsat-
 379 isfied CLAUSE-gadget with solution from Lemma 3.2.8.

380 This solution uses $3n + (11m + (m - k)) = 15n - k$ segments.

381 □

382 **Lemma 3.2.12.** *Given an instance of MAX-(3,3)-SAT of size n , and solution of size w to*
 383 *the instance of geometric cover, constructed according to Lemma 3.2.1, there exists a solution*
 384 *to MAX-(3,3)-SAT of size at least $15n - w$.*

385 *Proof.* Among $x_i^{true} \cup x_i^{false}$, we need to use at least 3 segments (Lemma 3.2.3). If we have
 386 chosen both segments (c_i, g_i) and (f_i, h_i) , then we have used at least 4 segments (Lemma 3.2.4).

387 If we chose at most one of the segments (c_i, g_i) and (f_i, h_i) , choose the corresponding
 388 variable value to the solution. If we chose both segments, choose the value that appears in
 389 most (at least 2) clauses. If we have chosen none of the segments, choose any value.

390 To cover $\bigcup_{1 \leq i \leq n} C_variable_i$ we have used at least $3n + a$ segments, where a is the number
 391 of i such that we have chosen both values (c_i, g_i) and (f_i, h_i) .

392 Among the segments responsible for the clause $C_i = x \vee y \vee z$ we need to use at least 11
 393 segments (Lemma 3.2.9) and if we can cover it with 11 segments, then we have earlier chosen
 394 segment responsible for the value of variable x, y or z that satisfies C_i .

395 So we have at least 11 segments for satisfied clauses and at least 12 segments for unsatisfied
 396 clauses, so we cover it with at least $11n + b$ segments, where b is number of clauses where
 397 none of the variables x, y, z were chosen. If the segment responsible for value of x was taken,
 398 but this variable is set to have different value, then we have chosen segments for both x and
 399 $\neg x$ for this variable, so "we cheated" and this maybe clause is not met, but we assigned the
 400 value for this x_i that meets the most clauses, so for each of such "cheated" variables, at most
 401 one of the clauses isn't met.

402 So there are at most $a + b$ unsatisfied clauses in this instance, so we have shown the
 403 assignment with at least $n - (a + b)$ satisfied clauses.

$$w \geq 3n + a + 11n + b = 14n + a + b$$

$$15n - w \leq 15n - 14n - a - b = n - (a + b)$$

404 □

405 *Proof of Lemma 3.2.1.* Given an instance of MAX-(3,3)-SAT of size n with optimal result k .
 406 Let's construct an instance of geometric cover, constructed in aforementioned manner.

407 Given the Lemma 3.2.11, we know the optimal solution for the constructed geometric cover
 408 is at most $15n - k$ and since the k is optimal solution for MAX-(3,3)-SAT, then according to
 409 Lemma 3.2.12 there doesn't exist a solution with cost less than $15n - k$. □

3.3. Weighted segments

3.3.1. FPT for weighted segments with δ -extensions

Theorem 3.3.1. (*FPT for weighted segment cover with δ -extensions*). *There exists an algorithm that given a family \mathcal{P} of n weighted segments (in any direction), a set of m points \mathcal{C} and a parameter k , runs in time $f(k) \cdot (nm)^c$ for some computable function f and constant c , and outputs a subfamily $\mathcal{R} \subseteq \mathcal{P}$ such that $|\mathcal{R}| \leq k$ and $\mathcal{R}^{+\delta}$ covers all points in \mathcal{C} .*

To solve this problem we will introduce kernel for slightly different problem: Weighted segment cover of points and segments. In shortcut: WSCPS.

Lemma 3.3.1. (*Algorithm for kernel of WSCPS*). *There exists an algorithm that given a family \mathcal{P} of n weighted segments (in any direction), a set of m_1 points \mathcal{C}_1 and m_2 segments \mathcal{C}_2 and a parameter k , runs in time $f(k) \cdot g(m_1, m_2) \cdot n^c$ for some computable functions f, g and constant c , and outputs a subfamily $\mathcal{sol} \subseteq \mathcal{P}$ such that $|\mathcal{sol}| \leq k$ and \mathcal{sol} covers all points in \mathcal{C}_1 and all segments in \mathcal{C}_2 .*

Proof Only sketch for now.

We can compute dynamic programming $dp(A, B, z)$ – the best cost to cover at least whole segment A, B using at most z segments. A, B are all interesting points – ends of any segment given on the input or points given on the input. We can compute it in polynomial time.

Then we can create a new double weighted set (original weight, number of used segments from \mathcal{P}) – \mathcal{P}_2 that has only segments which never cover partially any segment from \mathcal{C}_2 (covers the whole segment or doesn't cover at all). In such \mathcal{P}_2 we can find solution \mathcal{R} where any 2 segments have empty intersection (don't cover each other and don't meet at the ends). Because if we had such solution, we can merge these two segments and such segment there's also in \mathcal{P}_2 .

In that case we can find kernel of \mathcal{P}_2 of size $k \cdot (m_1 + 2m_2)^2$, because we only need to take the best weight covering some subset of $\mathcal{C}_1 \cup \mathcal{C}_2$.

Lemma 3.3.2. *Kernel in WSCPS.* *TODO: formulate it properly*

For segment cover, there is a kernel of size $f(k)$ in WSCPS.

Claim 3.3.1. *If there are more than k lines with at least $k+1$ points on them, then they can't be covered with k segments.*

Claim 3.3.2. *If there is more than k^2 points that don't lie on any line with more than k points on it, then they can't be covered with k segments.*

Claim 3.3.3. *For every long line L (with more than k points on them) we can choose $f(k)$ points on them, that if we cover all of these points with at most k segments, then the rest of the points with δ -extensions will be covered by segments in the direction of line L .*

Proof of Lemma 3.3.2. After applying the previous lemmas, we have at most $k^2 + k \cdot f(k)$ points that can be covered in any direction and for the rest of the points we can draw at most $k \cdot f(k)$ segments along their respective long lines that have to be covered by segments after δ -extensions.

Then we extend every available segment by δ -extension and we achieve the kernel in WSCPS for this instance of problem.

Lemma 3.3.3. *If all the points are covered with k segments and the biggest $2(1 + 1/\delta)^{k+1}$ spaces between points are filled, the whole segment is filled after δ -extensions of these segments.*

452 **Proof.** Let's name the $2(1+1/\delta)^{k+1}$ -st biggest space between points as y . We have guarantee
453 that all segments of length $x > y$ are covered without δ -extensions.

454 Let's take one space between points that is not covered before δ -extension and we will
455 prove it will be covered after δ -extensions. Let's assume it isn't.

456 This space has length x . Since it's uncovered, $x \leq y$.

457 Let's take side where the sum of lengths of segments covering the points is greater (left or
458 right). Without loss of generality, let us assume it's right.

459 There are at most k segments to the right of this space between points. Name their lengths
460 $l_1, l_2 \dots l_k$. If the point is covered in the other direction, the segment is degenerated to the
461 point and $l_i = 0$. Name the space between endpoints of l_i and $l_{i+1} - x_i$. Of course, x_i is
462 uncovered space between two points, therefore $x_i \leq y$.

463 TUTAJ BEDZIE PEWNIE RYSUNEK Z TYMI SUPER RZECZAMI DO PRZERW

464 Let's write equations meaning that i -th segment doesn't cover space x after δ -expansion.

$$l_1\delta < x \leq y \Rightarrow l_1 < y/\delta$$

$$l_2\delta < x + l_1 + x_1 < 2y + y/\delta \Rightarrow l_2 < 2y/\delta + y/\delta^2$$

$$l_3\delta < x + l_1 + x_1 + l_2 + x_2 < 3y + 3y/\delta + y/\delta^2 \Rightarrow l_3 < 3y/\delta + 3y/\delta^2 + y/\delta^3$$

465 From this we can "guess" induction $l_i < y((1 + 1/\delta)^i - 1)$

466 Trivially for $l_1 < y/\delta$.

Assume that for all $j < i$:

$$l_j < y((1 + 1/\delta)^j - 1)$$

467 .

$$\begin{aligned} 468 \quad l_i\delta &< x + \sum_{j=1}^{i-1} (l_j + x_j) < iy \sum_{j=1}^{i-1} l_j < iy + \sum_{j=1}^{i-1} j = 1^{i-1}y((1 + 1/\delta)^j - 1) = iy - (i - \\ 469 \quad 1)y + \sum_{j=1}^{i-1} j &= 1^{i-1}y(1 + 1/\delta)^j = y(1 + \sum_{j=1}^{i-1} (1 + 1/\delta)^j) = y(2 + \sum_{j=1}^{i-1} (1 + 1/\delta)^j - 1) = \\ 470 \quad y(\sum_{j=0}^{i-1} (1 + 1/\delta)^j - 1) &= y((1 + 1/\delta)^i / (1 - (1 + 1/\delta)) - 1) = y((1 + 1/\delta)^i \delta - 1) < y((1 + 1/\delta)^i \delta - \delta) \end{aligned}$$

Of course we also know that (since we have chosen the side with greater sum of the width of segments):

$$\sum_{i=1}^k l_i \geq 1/2 \cdot y \cdot 2(1 + 1/\delta)^{k+1} = y \cdot (1 + 1/\delta)^{k+1}$$

471 But $\sum_{i=1}^k l_i < \sum_{i=1}^k y((1 + 1/\delta)^i - 1) = y((1 + 1/\delta)^{k+1} / (1 - (1 + 1/\delta)) - k) = y((1 +$
472 $1/\delta)^{k+1} \delta - k) < y(1 + 1/\delta)^{k+1}$

473 Therefore the space must have been covered after δ -expansions.

474 3.3.2. W[1]-completeness for weighted segments in 3 directions

475 **Theorem 3.3.2.** *W[1]-completeness for weighted segments in 3 directions.* Consider
476 the problem of covering a set \mathcal{C} of points by selecting k axis-pararell or right-diagonal weighted
477 segments with weights from a set \mathcal{P} with minimal weight. Assuming ETH, there is no algorithm
478 for this problem with running time $f(k) \cdot (|\mathcal{C}| + |\mathcal{P}|)^{o(\sqrt{k})}$ for any computable function f .

479 We will show reduction from grid tiling problem.

480 Let's have an instance of grid tiling problem – size of the grid k , number of elements
481 available n and k^2 sets of available pairs in every tile $S_{i,j} \subseteq \{1, n\} \times \{1, n\}$.

482 **Construction.** We construct a set \mathcal{P} of segments and a set \mathcal{C} of points.

483 First let's choose any ordering of n^2 elements $\{1, n\} \times \{1, n\}$ and name this sequence

484 $a_1 \dots a_{n^2}$.

$$match_v(i, j) \iff a_i = \{x_i, y_i\} \wedge a_j = \{x_j, y_j\} \wedge x_i = x_j$$

$$match_h(i, j) \iff a_i = \{x_i, y_i\} \wedge a_j = \{x_j, y_j\} \wedge y_i = y_j$$

Points. Define points:

$$h_{i,j,t} = (j \cdot (n^2 + 1) + t, (n^2 + 1) \cdot i)$$

$$v_{i,j,t} = ((n^2 + 1) \cdot i, j \cdot (n^2 + 1) + t)$$

Let's define sets H and V as:

$$H = \{h_{i,j,t} : 1 \leq i, j \leq k, 1 \leq t \leq n^2\}$$

$$V = \{v_{i,j,t} : 1 \leq i, j \leq k, 1 \leq t \leq n^2\}$$

485 Let's define $\epsilon = 0.1$. For a point $\{x, y\} = p$ we define points $p^L = \{x - \epsilon, y\}$, $p^R = \{x + \epsilon, y\}$,

486 $p^U = \{x, y - \epsilon\}$, and $p^D = \{x, y + \epsilon\}$.

Then we define:

$$\mathcal{C} := H \cup \{p^L : p \in H\} \cup \{p^R : p \in H\} \cup V \cup \{p^U : p \in V\} \cup \{p^D : p \in V\}$$

487 **Segments.** Define horizontal segments.

$$hor_{i,j,t_1,t_2} = (h_{i,j,t_1}^R, h_{i,j+1,t_2}^L)$$

$$ver_{i,j,t_1,t_2} = (v_{i,j,t_1}^D, v_{i,j+1,t_2}^U)$$

$$horbeg_{i,t} = (h_{i,1,1}^L, h_{i,1,t}^L)$$

$$horend_{i,t} = (h_{i,n,t}^R, h_{i,n,n^2}^R)$$

$$verbeg_{i,t} = (v_{i,1,1}^U, v_{i,1,t}^U)$$

$$verend_{i,t} = (v_{i,n,t}^D, v_{i,n,n^2}^D)$$

$$\begin{aligned} HOR &= \{hor_{i,j,t_1,t_2} : 1 \leq i \leq k, 1 \leq j < k, 1 \leq t_1, t_2 \leq n^2, match_h(t_1, t_2)\} \\ &\cup \{horbeg_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\} \\ &\cup \{horend_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\} \end{aligned}$$

$$\begin{aligned} VER &= \{ver_{i,j,t_1,t_2} : 1 \leq i \leq k, 1 \leq j < k, 1 \leq t_1, t_2 \leq n^2, match_v(t_1, t_2)\} \\ &\cup \{verbeg_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\} \\ &\cup \{verend_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2\} \end{aligned}$$

$$DIAG := \{(h_{i,j,t}, v_{j,i,t}) : 1 \leq i, j \leq k, 1 \leq t \leq n^2, a_t \in S_{i,j}\}$$

488 TODO: explain that these segments are in fact diagonal

$$\mathcal{P} := HOR \cup VER \cup DIAG$$

489 **Lemma 3.3.4.** *If there exists solution for grid tiling, then there exists solution for our construction using $2(k+1)k + k^2$ segments with weight exactly $2k \cdot (k(n^2 + 1) - 2 - 2\epsilon(k-1))$.*

Claim 3.3.4. *If there exists a solution to the grid tiling $c_1 \dots c_k$ and $r_1 \dots r_k$, then there exists a solution covering all points*

$$\{h_{i,j,t} : 1 \leq i, j \leq k, t = (c_i, r_j)\} \cup \{v_{i,j,t} : 1 \leq i, j \leq k, t = (c_j, r_i)\}$$

491 *with segments in DIAG and the rest in VER or HOR and has weight $2k \cdot (k(n^2 + 1) - 2 - 2\epsilon(k-1))$.*

493 **Proof.** TODO: jakiś prosty z definicji

494 **Lemma 3.3.5.** *If there exists solution for our construction using $2(k+1)k + k^2$ segments with weight exactly $2k \cdot (k(n^2 + 1) - 2 - 2\epsilon(k-1))$, then there exists a solution for grid tiling*

496 **Proof.** This follows from Lemma 3.3.6, because we just take which points are covered with
497 *DIAG.*

498 **Claim 3.3.5.** *Points p^L, p^R, p^U, p^D cannot be covered with DIAG.*

499 **Claim 3.3.6.** *Points in $H \cup \{p^L : p \in H\} \cup \{p^R : p \in H\}$ cannot be covered with VER.*

500 *Points in $V \cup \{p^U : p \in V\} \cup \{p^D : p \in V\}$ cannot be covered with HOR.*

501 **Claim 3.3.7.** *For given i, j if none of the points $h_{i,j,t}$ ($v_{i,j,t}$) for $1 \leq t \leq n^2$ are covered with
502 *DIAG, then some spaces between neighbouring points were covered twice.**

503 **Claim 3.3.8.** *For given i, j two points h_{i,j,t_1}, h_{i,j,t_2} (v_{i,j,t_1}, v_{i,j,t_2}) for $1 \leq t_1 < t_2 \leq n^2$ are
504 *covered with DIAG, then one of them had to be also covered with a segment from HOR*
505 *(VER).**

506 **Proof.** Point v_{i,j,t_2}^L had to be covered with VER from Claims 3.3.5 and 3.3.6. And every
507 segment in VER covering v_{i,j,t_2}^L , covers also v_{i,j,t_1}^L .

508 **Lemma 3.3.6.** *If there exists solution for our construction with weight at most (exactly)
509 $2k \cdot (k(n^2 + 1) - 2 - 2\epsilon(k-1))$, then for every i, j there must be exactly one t such that
510 $h_{i,j,t}$ ($v_{i,j,t}$) is covered with DIAG and moreover if h_{i,j,t_1} and $h_{i,j+1,t_2}$ are uncovered, then
511 $\text{math}_h(t_1, t_2)$. Analogically for v .*

512 **Proof.** Only k^2 points can be covered only in DIAG, the rest has to be covered with
513 *VER \cup HOR.* Therefore every result must be at least *ALL_LINES* - $2k^2\epsilon$, because only
514 $2k^2$ spaces of length ϵ can be uncovered in this axis.

515 Of course if h_{i,j,t_1} and $h_{i,j+1,t_2}$ are uncovered, then there must exist a segment in HOR
516 between h_{i,j,t_1}^R and $h_{i,j+1,t_2}^L$, so $\text{math}_h(t_1, t_2)$ must be true.

517 3.3.3. What is missing

518 We don't know FPT for axis-pararell segments without δ -extensions.

519 Chapter 4

520 Geometric Set Cover with lines

521 4.1. Lines parallel to one of the axis

522 When \mathcal{R} consists only of lines parallel to one of the axis, the problem can be solved in
523 polynomial time.

524 We create bipartial graph G with node for every line on the input split into sets: H –
525 horizontal lines and V – vertical lines. If any two lines cover the same point from \mathcal{C} , then we
526 add edge between them.

527 Of course there will be no edges between nodes inside H , because all of them are pararell
528 and if they share one point, they are the same lines. Similar argument for V . So the graph is
529 bipartial.

530 Now Geometric Set Cover can be solved with Vertex Cover on graph G . Since Vertex
531 Cover (even in weighted setting) on bipartial graphs can be solved in polynomial time.

532 Short note for myself just to remember how to this in polynomial time:

533 Non-weighted setting - Konig theorem + max matching

534 Weighted setting - Min cut in graph of $\neg A$ or $\neg B$ (edges directed from V to H)

535 4.2. FPT for arbitrary lines

536 You can find this is Platypus book. We will show FPT kernel of size at most k^2 .

537 (Maybe we need to reduce lines with one point/points with one line).

538 For every line if there is more than k points on it, you have to take it. At the end, if there
539 is more than k^2 points, return NO. Otherwise there is no more than k^4 lines.

540 In weighted settings among the same lines with different weights you leave the cheapest
541 one and use the same algorithm.

542 4.3. APX-completeness for arbitrary lines

543 We will show a reduction from Vertex Cover problem. Let's take an instance of the Vertex
544 Cover problem for graph G . We will create a set of $|V(G)|$ pairwise non-pararell lines, such
545 that no three of them share a common point.

546 Then for every edge in $(v, w) \in E(G)$ we put a point on crossing of lines for vertices v
547 and w . They are not pararell, so there exists exactly one such point and any other line don't
548 cover this point (any three of them don't cross in the same point).

Solution of Geometric Set Cover for this instance would yield a sound solution of Vertex Cover for graph G . For every point (edge) we need to choose at least one of lines (vertices) v or w to cover this point.

Vertex Cover for arbitrary graph is APX-complete, so this problem is also APX-complete.

4.4. 2-approximation for arbitrary lines

Vertex Cover has an easy 2-approximation algorithm, but here very many lines can cross through the same point, so we can do d -approximation, where d is the biggest number of lines crossing through the same point. So for set where any 3 lines don't cross in the same point it yields 2-approximation.

The problematic cases are where through all points cross at least k points and all lines have at least k points on them. It can be created by casting k -grid in k -D space on 2D space.

Greedy algorithm yields $\log |\mathcal{R}|$ -approximation, but I have example for this for bipartial graph and reduction with taking all lines crossing through some point (if there are no more than k) would solve this case. So maybe it works.

Unfortunately I haven't done this :(

I can link some papers telling it's hard to do.

4.5. Connection with general set cover

Problem with finite set of lines with more dimensions is equivalent to problem in 2D, because we can project lines on the plane which is not perpendicular to any plane created by pairs of (point from \mathcal{C} , line from \mathcal{P}).

Of course every two lines have at most one common point, so is every family of sets that have at most one point in common equivalent to some geometric set cover with lines?

No, because of Desargues's theorem. Have to write down exactly what configuration is banned.

573 Chapter 5

574 Geometric Set Cover with polygons

575 5.1. State of the art

576 Covering points with weighted discs admits PTAS [Li and Jin, 2015] and with fat polygons
577 with δ -extensions with unit weights admits EPTAS [Har-Peled and Lee, 2009].

578 Although with thin objects, even if we allow δ -expansion, the Set Cover with rectangles is
579 APX-complete (for $\delta = 1/2$), it follows from APX-completeness for segments with δ -expansion
580 in Section 3.2.

581 Covering points with squares is W[1]-hard [Marx, 2005]. It can be proven that assuming
582 *SETH*, there is no $f(k) \cdot (|\mathcal{C}| + |\mathcal{P}|)^{k-\epsilon}$ time algorithm for any computable function f and
583 $\epsilon > 0$ that decides if there are k polygons in \mathcal{P} that together cover \mathcal{C} , *Theorem 1.9* in [Marx
584 and Pilipczuk, 2015].

585 Chapter 6

586 Conclusions

587 Bibliography

- 588 [Har-Peled and Lee, 2009] Har-Peled, S. and Lee, M. (2009). Weighted geometric set cover
589 problems revisited. *Journal of Computational Geometry*, 3.
- 590 [Håstad, 2001] Håstad, J. (2001). Some optimal inapproximability results. *J. ACM*,
591 48(4):798–859.
- 592 [Li and Jin, 2015] Li, J. and Jin, Y. (2015). A PTAS for the weighted unit disk cover problem.
593 *CoRR*, abs/1502.04918.
- 594 [Marx, 2005] Marx, D. (2005). Efficient approximation schemes for geometric problems? In
595 Brodal, G. S. and Leonardi, S., editors, *Algorithms – ESA 2005*, pages 448–459, Berlin,
596 Heidelberg. Springer Berlin Heidelberg.
- 597 [Marx and Pilipczuk, 2015] Marx, D. and Pilipczuk, M. (2015). Optimal parameterized algo-
598 rithms for planar facility location problems using voronoi diagrams. *CoRR*, abs/1504.05476.