

CLINICAL REVIEW: Rare Causes of Hypercalcemia

Thomas P. Jacobs and John P. Bilezikian

Columbia University College of Physicians and Surgeons, New York, New York 10032

Context: Although hypercalcemia is usually caused by primary hyperparathyroidism or malignancy, a number of other conditions can be important to consider. This review considers unusual causes of hypercalcemia that are generally not found in reviews on this subject.

Evidence Acquisition: Articles describing rarely reported associations between hypercalcemia and unusual causes were identified through a computer search of the terms hypercalcemia/etiology and through the references listed in those articles. We grouped the 58 different reports into categories defined by a presumed etiology: increased levels of 1,25-dihydroxyvitamin D or PTHrP, occult milk-alkali syndrome, and undefined mechanisms. Reports in infants and children are listed separately, as are reports of pseudohypercalcemia, situations that are not truly hypercalcemic because the ionized calcium is normal.

Evidence Synthesis: In some situations, as this review points out, a number of unusual causes of hypercalcemia are important to consider. The search for an elusive cause of hypercalcemia is best accomplished by the most likely potential mechanism. An orderly search in this manner is likely to reveal the underlying cause.

Conclusions: That so many patients have been described with rare and usually poorly understood causes of hypercalcemia highlights our incomplete understanding of calcium metabolism in humans and suggests additional areas in which directed clinical investigation might improve our knowledge of the normal metabolism of calcium. (*J Clin Endocrinol Metab* 90: 6316–6322, 2005)

HYPERCALCEMIA IS OFTEN a clue to the presence of unsuspected illness. Major textbooks and reviews correctly point out that the great majority of patients with elevated serum calcium will be found to have either primary hyperparathyroidism or malignancy, although the differential diagnosis is much longer. These other causes of hypercalcemia, including vitamin D intoxication, sarcoidosis, tuberculosis, some fungal infections, thyrotoxicosis, Addison's disease, milk-alkali syndrome related to the prescription of absorbable alkali and calcium, vitamin A intoxication, therapy with thiazide diuretics or lithium carbonate, familial hypocalciuric hypercalcemia, prolonged immobilization in patients with high skeletal turnover, and the recovery phase of rhabdomyolysis-associated acute renal failure, all amount to fewer than about 10% of all causes of hypercalcemia. Nevertheless, they are important to consider in certain clinical situations when the underlying cause of hypercalcemia cannot be attributed to primary hyperparathyroidism or overt malignancy. Despite the rather inclusive nature of this list of potential causes of hypercalcemia, there are still other more unusual etiologies reported in single patients or in small groups of patients. In the typical practice of many endocrinologists, patients are seen occasionally for whom the etiology of the hypercalcemia is not attributable to the illnesses listed above and never becomes well defined. In this review, we summarize published reports of patients with hypercalcemia whose etiology is so unusual that it is not listed in many reviews of hypercalcemia and is even more rarely considered in patients with hypercalcemia of obscure etiology. In some reports, a presumed etiology is offered (e.g.

elevated levels of PTHrP). In others, no presumed mechanism has been defined. Some reports were published before the availability of current methodological tools for investigating hypercalcemia, but have been included because of the unusual clinical circumstances. This review adds to what has previously been reported in the literature for adults and children (1–6).

Methods

The reports included in this review were identified through a comprehensive computer-based search for hypercalcemia/etiology and a search of references in the identified reports. We excluded all forms of hypercalcemia of malignancy and hyperparathyroidism. We also excluded reports of hypercalcemia that occurred in patients with unusual medical problems in which the hypercalcemia seemed to be caused by a commonly encountered mechanism [e.g. tertiary hyperparathyroidism in Low's syndrome (7)]. We have grouped these reports into categories defined by a presumed etiology, although it is clear in some conditions associated with hypercalcemia that the mechanism underlying the hypercalcemia may differ among patients. For example, hypercalcemia in patients with tuberculosis has been reported to occur in patients with elevated (8), normal (9), or low (10) levels of serum 1,25-dihydroxyvitamin D.

Reports in Adults

Hypercalcemia mediated by elevated levels of calcitriol (Table 1)

In addition to sarcoidosis, tuberculosis, and some systemic fungal infections, other illnesses in which the pathology is characterized by granuloma formation have been associated with hypercalcemia in association with elevated levels of 1,25-dihydroxyvitamin D. It is presumed that the mechanism is similar to the previously described etiology: namely, in-

First Published Online August 30, 2005

Abbreviation: SLE, Systemic lupus erythematosus.

JCEM is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.

TABLE 1. Rare causes of hypercalcemia associated with elevated 1,25-dihydroxyvitamin D

1. Wegener's granulomatosis (11–13)
2. Cat scratch fever (14)
3. Crohn's disease (15)
4. Acute granulomatous pneumonia (16)
5. Hepatic granulomatosis in chronic dialysis (17)
6. Talc granulomatosis (18)
7. Silicone granulomatosis (19)
8. BCG therapy (20)
9. 8-Cl-cAMP therapy (21)
10. Lipoid pneumonia (22)
11. Subcutaneous fat necrosis of the newborn (86)

BCG, Calmette-Guérin bacillus.

creased conversion of 25-hydroxyvitamin D to 1,25-dihydroxyvitamin D. The activated macrophages within the granulomas, confirmed only in sarcoidosis-associated granulomas, are believed to contain 1 α -hydroxylase activity that is responsible for the enzymatic activation of 25-hydroxyvitamin D.

In Wegener's granulomatosis, hypercalcemia associated with elevated levels of 1,25-dihydroxyvitamin D has been reported on at least three occasions (11, 12). In one patient, serum calcium and 1,25-dihydroxyvitamin D levels changed in parallel with cyclophosphamide and glucocorticoid therapy (13). Identical 18-yr-old male twins with cat scratch fever developed hypercalcemia and hypercalciuria with elevated 1,25-dihydroxyvitamin D levels. In each patient, serum and urinary calcium concentrations returned to normal when the bacterial infection was successfully treated (14). Hypercalcemia has also been described in two patients with Crohn's disease who had typical granulomas on bowel biopsy. With glucocorticoid therapy, the hypercalcemia and elevated 1,25-dihydroxyvitamin D levels both remitted (15). Acute granulomatous pneumonitis, a rare complication of methotrexate therapy, has been reported in association with hypercalcemia and inappropriately elevated levels of 1,25-dihydroxyvitamin D. Glucocorticoid therapy was effective in controlling the hypercalcemia and reducing the 1,25-dihydroxyvitamin D levels (16). Four patients receiving chronic hemodialysis developed hypercalcemia in association with hepatic granulomatosis of obscure etiology. The inappropriately normal PTH levels could well have been due to the accumulation of inactive PTH fragments, but the high-normal 1,25-dihydroxyvitamin D levels argue for a nonrenal source of enzymatic activation. In each instance, hypercalcemia responded promptly to glucocorticoid therapy (17).

Foreign material introduced into the human body often elicits a granulomatous inflammatory reaction. Hypercalcemia associated with elevated levels of 1,25-dihydroxyvitamin D has been reported in patients with granulomatous pneumoconiosis pneumonia caused by talc (18), a granulomatous reaction to silicone injections around the hips (19), and after a systemic reaction to Calmette-Guérin bacillus therapy for bladder cancer complicated by granulomatous hepatitis and marked hypercalcemia (20).

In some situations, the increase in 1,25-dihydroxyvitamin D has not been clearly associated with a granulomatous reaction. For example, an experimental analog of cAMP (8-Cl-cAMP) given to patients with advanced solid malignancies regularly caused hypercalcemia associated with high

levels of 1,25-dihydroxyvitamin D (21). Lipoid pneumonia in a man with mental retardation was complicated by recurrent hypercalcemia and elevated levels of calcitriol (22).

Hypercalcemia caused by occult milk-alkali syndrome (Table 2)

Hypercalcemia with hypercalciuria and renal dysfunction is usually easily identified as milk-alkali syndrome in patients who ingest large amounts of calcium and absorbable alkalis (23). Rarely, the diagnosis may be undetected in patients who are either not aware that they are ingesting these substances or who conceal their use. Three patients have been reported from Taiwan who chewed betel nuts along with oyster shell powder, the latter to neutralize the bitter taste of the betel nuts. All patients had findings typical of milk-alkali syndrome (24, 25). Two patients who attempted suicide with buffered aspirin tablets each presented with hypercalcemia reasonably attributed to the associated ingestion of calcium carbonate (26). A woman with a severe eating disorder was hospitalized on three occasions with hypercalcemia caused by massive cheese ingestion in the setting of metabolic alkalosis and dehydration caused by recurrent vomiting and use of a thiazide diuretic (27). Similarly, a nurse with an extensive medical history developed extreme hypercalcemia caused by surreptitious ingestion of large amounts of calcium carbonate with a thiazide diuretic (28).

Hypercalcemia caused by PTHrP (Table 3)

Humoral hypercalcemia of malignancy is thought in many instances to be caused by secretion of PTHrP by the tumor (29–31). Although this presentation is now well recognized, hypercalcemia due to high serum levels of PTHrP in the setting of benign disease is very uncommon. A patient with systemic lupus erythematosus (SLE) with renal, central nervous system, and pleural involvement with diffuse lymphadenopathy presented with hypercalcemia accompanied by high levels of PTHrP. Large lymphocytes examined from an excised lymph node contained immunochemically identifiable PTHrP (32). The same group reported a 63-yr-old man with HIV-associated lymphadenopathy and recurrent hypercalcemia of unclear cause, whose cervical lymph node contained large lymphoid cells that stained positively for both PTHrP and its mRNA (33). Another young woman with a known diagnosis of SLE presented with bilateral chylous pleural effusions, diffuse pulmonary calcification, and lymphedematous bilateral breast enlargement. Elevated levels of PTHrP accompanied severe hypercalcemia, and calcium and PTHrP fluctuated in parallel in response to glucocorticoid therapy. Breast tissue did not stain for PTHrP, and no malignancy could be detected over 15 months of observation (34). Massive mammary hyperplasia during pregnancy in a 25-yr-old woman was accompanied by serious hypercalcemia, which responded promptly to bilateral

TABLE 2. Rare presentations of the milk-alkali syndrome

1. Oyster shell calcium in betel nut chewing (24, 25)
2. Overdose with buffered aspirin (26)
3. Massive cheese ingestion (27)
4. Munchausen's syndrome (28)

TABLE 3. Unusual PTHrP-mediated hypercalcemic conditions

1. SLE (32)
2. HIV-associated lymphadenopathy (33)
3. Lymphedema of chest and pleural cavities (34)
4. Massive mammary hyperplasia during pregnancy (35, 36)
5. During late pregnancy and lactation in hypoparathyroidism (37–39, 41)
6. With benign tumors of ovary and kidney, and in benign pheochromocytoma (43–46)

mastectomy (35). Subsequent immunostaining of the excised breast tissue demonstrated PTHrP reactivity in myoepithelial cells, but not in the normal breast tissue (36).

The development of hypercalcemia occurring just before and continuing for weeks after delivery was described in 1969 in two patients with hypoparathyroidism who had previously been well controlled with supplements of ergocalciferol. Neither woman breast fed, and hypercalcemia resolved quickly in one patient after a reduction in the dose of ergocalciferol (37). Hypercalcemia occurred during two successive pregnancies in an otherwise healthy woman with low PTH levels. During the second pregnancy, serum levels of PTHrP were found to be elevated, with persistent elevation of both serum calcium and PTHrP for several months after delivery (38). Another patient with surgical hypoparathyroidism became hypercalcemic while lactating after her first pregnancy, requiring withdrawal of calcium and calcitriol supplements. After a second pregnancy, her serum calcium remained normal off calcitriol and calcium, apparently supported by PTHrP, the serum concentration of which rose immediately after delivery to high levels and remained elevated during lactation for 72 wk (39). A number of groups have demonstrated that serum PTHrP levels are elevated during lactation and probably contribute substantially to the movement of calcium from the maternal skeleton to the mammary glands, causing some suppression of PTH (40, 41). Patients with hypoparathyroidism who are taking pharmacological amounts of vitamin D seem clearly at higher risk for hypercalcemia during the puerperium and lactation and are advised to reduce their intake of vitamin D until they are no longer breast feeding.

It is now clear that PTHrP has other important normal physiological functions (42). It is not surprising therefore to find a few reports of patients whose benign tumor was associated with PTHrP production and hypercalcemia. Benign ovarian tumor (43), renal adenoma (44), and pheochromocytomas (45, 46) have all been reported to be associated with hypercalcemia and elevated levels of PTHrP. When the benign tumor was removed, the hypercalcemia remitted. This syndrome has been dubbed by some humoral hypercalcemia of benignancy (44).

Hypercalcemia caused by medications (Table 4)

Several medications and chemicals other than calcium, vitamins A and D, lithium, and thiazide diuretics have been rarely associated with hypercalcemia. A 31-yr-old woman developed hypercalcemia in the setting of postpregnancy acute interstitial granulomatous nephritis when given omeprazole (47). Theophylline toxicity in 60 patients was complicated in 11 by hypercalcemia with normal levels of PTH,

TABLE 4. Medications that have been unusually associated with hypercalcemia

1. Omeprazole in acute interstitial nephritis (47)
2. Theophylline toxicity (48)
3. GH in intensive care unit patients (49)
4. Parenteral nutrition (50)
5. Fosarnet (52)
6. Hepatitis B vaccination (51)
7. 8-Cl-cAMP chemotherapy (21)
8. Manganese toxicity (53)

suggesting decreased sensitivity of the parathyroids to ionized calcium (48). High doses of human GH given to severely ill patients in a surgical intensive care unit appeared to cause hypercalcemia, which in some patients was severe, although it is not clear from the report whether parenteral nutrition and immobilization may have contributed to the hypercalcemia in some of the patients (49). Modest hypercalcemia in association with osteomalacia, bone pain, and fractures has been observed in several patients being treated with long-term parenteral nutrition, and in one small study was improved by removing vitamin D from the parenteral nutrition solutions (50). A 44-yr-old man developed polyarthritis and hypercalcemia shortly after his third vaccination for hepatitis B, eventually showing evidence of increased bone turnover and lytic bone lesions. A bone biopsy showed increased osteoclastic bone resorption, and his illness responded well to calcitonin, prednisone, furosemide, and clodronate (51). Although it is more often reported to cause a reduction in ionized calcium related to a chelating effect, fosarnet was believed to be the cause of hypercalcemia in two patients with AIDS complicated by cytomegalic infection (52). Manganese toxicity appears to cause hypercalcemia in addition to a characteristic neurological syndrome (53).

Hypercalcemia of unknown mechanism (Table 5)

Hypercalcemia has been reported in several medical settings in which the mechanisms of the hypercalcemia are unknown, either because they were reported before reliable measurements of PTH, PTHrP, or vitamin D metabolites were available or because despite measurements of these

TABLE 5. Rare causes of hypercalcemia in which the mechanism is not known

1. Eosinophilic granuloma (54)
2. Leprosy in rheumatoid arthritis (55)
3. Mycobacterium avium complicating AIDS (56)
4. Cytomegalic virus infection in AIDS (57)
5. Chronic berylliosis (58)
6. Nocardia asteroides pericarditis (59)
7. Diffuse osteoclastosis (70)
8. Paraffin granulomatosis (60)
9. Brucellosis (61)
10. Isolated ACTH deficiency (62)
11. Glucocorticoid withdrawal (63)
12. Hypocaloric diet in hypoparathyroidism (64)
13. Advanced chronic liver disease (65)
14. Type I Gaucher's disease with acute pneumonia (66)
15. Lymphedema in SLE (67)
16. Juvenile rheumatoid arthritis (68)
17. Lymphadenopathy with high IL-6 (69)

substances, a mechanism of hypercalcemia could not be defined.

Several conditions characterized by granulomatous pathology have been reported in patients with hypercalcemia in whom 1,25-dihydroxyvitamin D levels were normal or low. Eosinophilic granuloma has been occasionally complicated by hypercalcemia, which has responded to treatment with glucocorticoids (54). A patient with leprosy complicating rheumatoid arthritis developed hypercalcemia in association with low levels of 1,25-dihydroxyvitamin D and responded to dapsone and prednisone (55). In addition to the occasional association of hypercalcemia with Mycobacterium tuberculosis, hypercalcemia was also observed in a patient with Mycobacterium avium complicating AIDS (56). In other cases of AIDS, hypercalcemia has been described in patients with diffuse cytomegalic virus infection (57). The granulomatous pneumonitis of chronic berylliosis in two patients was found to be associated with hypercalcemia, which resolved spontaneously and without explanation after several months (58). Hypercalcemia complicated pericarditis caused by *Nocardia asteroides* in a patient with hypoparathyroidism (59). It complicated the management of a 63-yr-old male with multiorgan failure histologically associated with the presence of widespread granulomatous lesions associated with paraffin 28 yr after a paraffin mammoplasty (60). Finally, serum calcium levels adjusted for serum albumin were found to be slightly higher in a large group of patients with brucellosis than in a control group, although the differences were small, and additional investigation of mechanisms were not performed (61).

Classic endocrine disorders, such as Addison's disease and thyrotoxicosis, are known to be complicated at times by hypercalcemia. In addition, isolated ACTH deficiency can result in hypercalcemia (62), and even glucocorticoid withdrawal after surgical cure of Cushing's syndrome has been associated with persistently elevated serum calcium levels (63). In a patient with hypoparathyroidism and a stable low serum calcium level while taking large doses of vitamin D, a hypocaloric diet was complicated by hypercalcemia sufficiently severe to require treatment with glucocorticoids and iv fluids (64). Advanced chronic liver disease was complicated by hypercalcemia in 10 patients without concomitant malignancy. Most had mild azotemia, and PTH levels were suppressed or in the low-normal range, as were levels of 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D (65).

A patient with type I Gaucher's disease presented with acute pneumonia and hypercalcemia postulated to be caused by pathological activation of osteoclasts via undegraded glucocerebroside activation of proinflammatory cytokines (66). The lymphadema syndrome described above in association with SLE has been reported to occur in the presence of normal levels of PTHrP as well (67). A boy with juvenile rheumatoid arthritis and persistent hypercalcemia had a serum activity that stimulated bone resorption in vitro and was neutralized by antibody to IL-1 β (68). A 24-yr-old man presented with widespread musculoskeletal pain, lymphadenopathy, hypercalcemia, and hyperphosphatemia. Histological and chemical indices of bone turnover were high, and serum levels of IL-6 were 100-fold higher than normal. The patient improved after administration of iv pamidronate

(69). Finally, a 31-yr-old African woman presented with multiple painful, lytic bony lesions associated with systemic inflammatory symptoms responding to glucocorticoids, but eventually died of severe hypercalcemia. Histological examination of three of the lesions showed only intense osteoclastic bone resorption by normal-appearing osteoclasts, with some foci of osteoblastic new bone formation (70).

Diseases Occurring in Childhood (Table 6)

Children present with hypercalcemia probably less frequently than adults, but the causes that we recognize to be common in adults are also common in children. Young children and infants, however, have been reported with hypercalcemia in association with a number of conditions seen almost exclusively in that population.

Heritable diseases of childhood

A number of monogenic heritable disorders are complicated by hypercalcemia. Jansen's disease, a heritable disorder of dwarfism, is characterized by short stature, scoliosis, short limbs, cranial abnormalities, and hypercalcemia in some cases. It is associated with constitutive activation of the PTH/PTHrP receptor (71) and, somewhat surprisingly, is not associated with osteitis fibrosis cystica even after many years (72). Hypophosphatasia is caused by loss of function mutations in the gene for tissue-nonspecific alkaline phosphatase. Phenotypic presentation varies widely, but in the more severe infantile form, extensive demineralization caused by failure of osteoblastic new bone formation leads to multiple fractures and early mortality. Hypercalcemia probably results from failure of bone matrix mineralization in the face of unimpeded osteoclastic bone resorption (73). Williams-Beuren syndrome, an autosomal dominant heritable disorder, causes elfin facies, supravalvular aortic stenosis, stenosis of pulmonary arteries, short stature, and developmental delay and often presents with hypercalcemia during infancy. Deletions in maternal or paternal chromosome 7q11.23 have been identified in most patients (74). Although evidence supporting a mechanism of increased intestinal calcium absorption or reduced calcitonin excretion has been offered, there is at present no defined mechanism causing the hypercalcemia that has gained consensus (75). In most children, the hypercalcemia resolves by 4 yr of age. A child with primary oxalosis who had required both renal and liver transplants developed recurrent hypercalcemia with lytic

TABLE 6. Rare causes of hypercalcemia in infants and young children

Associated with heritable diseases of childhood

1. Jansen's osteodystrophy (71, 72)
2. Hypophosphatasia (73)
3. Williams-Beuren syndrome (75)
4. Primary oxalosis (76)
5. Congenital lactase deficiency (77)
6. Down's syndrome (78, 79)

Associated with acquired diseases of childhood

1. Renal tubular acidosis (80–83)
2. Phosphate depletion in severe prematurity (84)
3. Subcutaneous fat necrosis of the newborn (86)
4. Infantile hypothyroidism (87)

bone lesions. Bone biopsy demonstrated numerous oxalate crystals associated with macrophages and increased areas of bone erosion (76). Infants with congenital lactase deficiency frequently present with hypercalcemia, hypercalciuria, and nephrocalcinosis. The mechanism of the hypercalcemia is unclear, and it quickly resolves on a lactose-free diet (77). Down's syndrome complicated by hypercalcemia has been reported in four patients, probably, but not clearly, on the basis of excessive calcium supplementation (78, 79).

Acquired diseases of childhood

In infants, hypercalcemia has been reported to complicate renal tubular acidosis. So-called Lightwood's syndrome is rarely seen now (80), but at times was complicated by hypercalcemia in children not exposed to supplemental dietary calcium (81, 82). Severe hypercalcemia caused hospitalization of a 21-d-old male subsequently shown to have distal renal tubular acidosis, which had not resolved by age 4 yr (83). Very premature infants with sustained hypercalcemia and hypophosphatemia corrected their metabolic abnormalities when phosphate supplements were added to expressed breast milk (84) or total parenteral nutrition supplementation. The authors of this report ascribed the hypercalcemia to severe phosphate depletion syndrome, which has been observed in laboratory rodents, but not in adult humans (85).

Subcutaneous fat necrosis of the newborn, an unusual form of self-limited localized panniculitis, often presents shortly after birth in infants who have sustained prenatal or perinatal complications, especially birth asphyxia and meconium aspiration. The etiology of this condition is unknown, but it is often accompanied by hypercalcemia associated with low levels of PTH and is mediated by elevated levels of 1,25-dihydroxyvitamin D, presumably secreted by macrophages participating in the granulomatous inflammatory process. The presence of hypercalcemia carries with it a mortality as high as 15%, although serum calcium usually is normal by 4 months of age (86).

Hypothyroidism in infants, especially when congenital, is associated with mild hypercalcemia. Increased intestinal calcium absorption secondary to increased sensitivity to vitamin D has been proposed as a mechanism, although levels of vitamin D metabolites are not usually high (87).

Pseudohypercalcemia (Table 7)

Persistently elevated total serum calcium in the presence of a normal ionized serum calcium level is rare and has been termed pseudohypercalcemia. Under normal circumstances about half the serum calcium concentration is bound to albumin. Therefore, in patients with elevations of serum albumin, mild hypercalcemia may be observed in the presence of normal free calcium (88). Two patients with essential thrombocythemia were found to have hypercalcemia that fluctuated directly with changes in platelet counts. The researchers believed that the best explanation for the hyper-

TABLE 7. Pseudohypercalcemia

1. High serum albumin (88)
2. Thrombocythemia (89)
3. Calcium binding to M-proteins (90, 93)

calcemia was the in vitro release of calcium from within activated platelets similar to the phenomenon of in vitro hyperkalemia that occurs in such patients (89). Pseudohypercalcemia in the setting of M-protein disorder has been ascribed to abnormal calcium binding by the IgM- κ paraprotein of a patient with Waldenström's macroglobulinemia (90) and by the IgG κ (91) and IgG λ (92) paraproteins as well. A well-studied case demonstrated that an IgG λ M protein in a patient with myeloma quantitatively bound 4 mol calcium/mol IgG (93). Two patients with IgM paraproteinemia were found to have pseudohypercalcemia caused by interference by the paraprotein with calcium measurements by an autoanalyzer, probably caused by hyperviscosity.

Conclusion

Successful management of hypercalcemia usually depends upon determining its etiology. In most patients the cause is obvious from the clinical setting and the results of serum assays of PTH, PTHrP, and vitamin D metabolites. The practicing clinician faced with hypercalcemia not attributable to the more common etiologies listed in most reviews of the subject may benefit from considering some of the rarer entities listed above. As presented in this review, the unusual associations between hypercalcemia and a number of diverse disorders demonstrates that our knowledge of the mechanisms of hypercalcemia is far from complete and points, in some cases, to undescribed humoral and cellular processes that may be important in the pathogenesis of hypercalcemia. As we learn more about the mechanisms underlying these associations, it is likely that we will gain new knowledge about the normal physiological processes that control calcium and skeletal metabolism.

Acknowledgments

Received March 28, 2005. Accepted August 15, 2005.

Address all correspondence and requests for reprints to: Dr. Thomas P. Jacobs, Columbia University Medical Center, 161 Fort Washington Avenue, New York, New York 10032. E-mail: tpj1@columbia.edu.

This work was supported by National Institutes of Health Grant DK-32333.

References

1. Root A 2003 Disorders of calcium and phosphorus metabolism. In: Rudolph CD, Rudolph AM, Hostetter MK, Lister G, Siegel NJ, eds. Rudolph's pediatrics. 21st ed. New York: McGraw-Hill; 2142–2162
2. Bringham F, Demay M, Kronenberg H 2003 Hormones and disorders of mineral metabolism. In: Larsen PR, Kronenberg HM, Melmed S, Polonsky KS, eds. Williams textbook of endocrinology. 10th ed. Philadelphia: Saunders; 1303–1371
3. Potts J 2005 Diseases of the parathyroid gland and other hyper- and hypocalcemic disorders. In: Kasper DL, Braunwald E, Fauci AS, Hauser SL, Longo DL, Jameson JL, eds. Harrison's principles of internal medicine. 16th ed. New York: McGraw-Hill; 2249–2268
4. Pont A 1989 Unusual causes of hypercalcemia. Endocrinol Metab Clin North Am 18:753
5. Wysolmerski JJ 2003 Miscellaneous causes of hypercalcemia. In: Favus MJ, ed. Primer on the metabolic bone diseases and disorders of mineral metabolism. 5th ed. Washington DC: American Society for Bone and Mineral Research; 260–266
6. Rodd C, Goodyer P 1999 Hypercalcemia of the newborn: etiology, evaluation, and management. Pediatr Nephrol 13:542–547
7. Hofbauer L, Kuhne C, Nies C, Heufelder A 2001 Identical twins with hypercalcemia due to Lowe's syndrome. Rheumatology 40:107–115
8. Bell N, Shary J, Shaw S, Turner R 1985 Hypercalcemia associated with

- increased circulating 1,25-dihydroxyvitamin D in a patient with pulmonary tuberculosis. *Calcif Tissue Int* 37:588–591
9. Sullivan J, Salmon W 1987 Hypercalcemia in active pulmonary tuberculosis. *South Med J* 80:572–576
10. Meuthen I, Kirsch L, Saborowski F 1991 Hypercalcemia in active tuberculosis of the lungs and cervical lymph nodes. *Deutsche Med Wochenschr* 116:899–902
11. Edelson G, Talpos G, Bone H 1993 Hypercalcemia associated with Wegener's granulomatosis and hyperparathyroidism: etiology and management. *Am J Nephrol* 13:275–277
12. Shaker J, Redlin K, Warren G, Findling J 1994 Case report: hypercalcemia with inappropriate 1,25-dihydroxyvitamin D in Wegener's granulomatosis. *Am J Med Sci* 308:115–118
13. Bosch X, Lopez-Soto A, Morello A, Olmo A, Urbano-Marquez A 1997 Vitamin D metabolite-mediated hypercalcemia in Wegener's granulomatosis. *Mayo Clin Proc* 72:440–444
14. Bosch X 1998 Hypercalcemia due to endogenous overproduction of active vitamin D in identical twins with cat-scratch disease. *JAMA* 279:532–534
15. Bosch X 1998 Hypercalcemia due to endogenous overproduction of 1,25-dihydroxyvitamin D in Crohn's disease. *Gastroenterology* 114:1061–1065
16. Cook N, Lake F 1996 Hypercalcaemia with methotrexate pneumonitis, possible association with pulmonary granulomata. *Austr NZ J Med* 26:715
17. Hardy P, Moriniere P, Tributou B, Hamdini N, Marie A, Bouffandeau B, Pruna A, Fournier A 1999 Liver granulomatosis is not an exceptional cause of hypercalcemia with hypoparathyroidism in dialysis patients. *J Nephrol* 12:398–403
18. Woywodt A, Schneider W, Goebel U, Luft F 2000 Hypercalcemia due to talc granulomatosis. *Chest* 117:1195–1196
19. Kozeny G, Barbato A, Bansal V, Vertuno L, Hano J 1984 Hypercalcemia associated with silicone-induced granulomas. *N Engl J Med* 311:1103–1105
20. Schattner A, Gilad A, Cohen J 2002 Systemic granulomatosis and hypercalcemia following intravesical bacillus Calmette-Guerin immunotherapy. *J Intern Med* 251:272–277
21. Saunders MP, Salisburn AJ, O'Byrne KJ, Long L, Whitehouse RM, Talbot DC, Mawer EB, Harris AL 1997 A novel cyclic adenosine monophosphate analog induces hypercalcemia via production of 1,25-dihydroxyvitamin D in patients with solid tumors. *J Clin Endocrinol Metab* 82:4004–4048
22. Greenaway T, Caterson I 1989 Hypercalcemia and lipid pneumonia. *Austr NZ J Med* 19:713–715
23. Orwoll E 1982 The milk-alkali syndrome: current concepts. *Ann Intern Med* 97:242–248
24. Wu K-D, Chuang R-B, Wu F, Hsu W-A, Jan I-S, Tsai K-S 1996 The milk-alkali syndrome caused by betelnuts in oyster shell paste. *Clin Toxicol* 34:741–745
25. Lin S, Lin Y, Cheema-Dhadli S, Davids M, Halperin M 2002 Hypercalcemia and metabolic alkalosis with betel nut chewing: emphasis on its integrative pathophysiology. *Nephrol Dialysis Transplant* 17:708–714
26. Reid I 1985 Transient hypercalcemia following overdoses of soluble aspirin tablets. *Austr NZ J Med* 15:364
27. Kallner G, Karlsson H 1987 Recurrent factitious hypercalcemia. *Am J Med* 82:536–538
28. Frame B, Jackson G, Kleerekoper M, Rao D, DeLorenzo A, Garcia M 1981 Acute severe hypercalcemia a la Munchausen. *Am J Med* 70:316–319
29. Grill V, Ho P, Body J, Johanson N, Lee SC, Kukreja SC, Moseley JM, Martin TJ 1991 Parathyroid hormone-related protein: elevated levels in both humoral hypercalcemia of malignancy and hypercalcemia complicating metastatic breast cancer. *J Clin Endocrinol Metab* 73:1309–1315
30. Burtis W, Brady T, Orloff J, Ersbak JB, Warrell Jr RP, Olson BR, Wu TL, Mitnick ME, Broadus AE, Stewart AF 1990 Immunochemical characterization of circulating parathyroid hormone-related protein in patients with humoral hypercalcemia of cancer. *N Engl J Med* 322:1106–1112
31. Stewart AF 2005 Hypercalcemia associated with cancer. *N Engl J Med* 352:373–379
32. Deftos L, Burton D, Baird S, Terkeltaub R 1996 Hypercalcemia and systemic lupus erythematosus. *Arthritis Rheum* 39:2066–2069
33. Burton D, Cheung T, Fierer J, Muchmore EA, Haghighi P, Frankel W, Deftos 2nd LJ 1995 PTHrP production by HIV-associated lymphadenopathy with hypercalcemia. *J Invest Med* 43:226A (Abstract)
34. Braude S, Graham A, Mitchell D 1991 Lymphoedema/hypercalcemia syndrome mediated by parathyroid-hormone-related protein. *Lancet* 337:140–141
35. Van Heerden J, Gharib H, Jackson I 1988 Pseudohyperparathyroidism secondary to gigantic mammary hypertrophy. *Arch Surg* 123:80–82
36. Khosla S, Van Heerden J, Gharib H, Jackson IT, Danks J, Hayman JA, Martin TJ 1990 Parathyroid hormone-related protein and hypercalcemia secondary to massive mammary hyperplasia. *N Engl J Med* 322:1157
37. Wright A, Joplin G, Dixon H 1969 Post-partum hypercalcemia in treated hypoparathyroidism. *Br Med J* 1:23–25
38. Lepre F, Grill V, Martin T 1993 Hypercalcemia in pregnancy and lactation associated with parathyroid hormone-related protein. *N Engl J Med* 328:666–667
39. Mather K, Chik C, Corenblum B 1999 Maintenance of serum calcium by parathyroid hormone-related peptide during lactation in a hypoparathyroid patient. *J Clin Endocrinol Metab* 84:424–427
40. Lippuner K, Zehnder HJ, Casez JP, Takkinen R, Jaeger P 1996 PTH-related protein is released into the mother's bloodstream during lactation: evidence for beneficial effects on maternal calcium-phosphate metabolism. *J Bone Miner Res* 11:1394–1399
41. Dobnig H, Kainer F, Stepan V, Winter R, Lipp R, Schaffer M, Kahr A, Nocnik S, Patterer G, Leb G 1995 Elevated parathyroid hormone-related peptide levels after human gestation: relationship to changes in bone and mineral metabolism. *J Clin Endocrinol Metab* 80:3699–3707
42. Strewler GJ 2000 The physiology of parathyroid-hormone related protein. *N Engl J Med* 342:177–185
43. Knecht T, Behling C, Burton D, Glass C, Deftos L 1996 The humoral hypercalcemia of benignancy. A newly appreciated syndrome. *Am J Clin Pathol* 105:487–492
44. Mahoney C, Cassady C, Weinberger E, Winters W, Benjamin D 1997 Humoral hypercalcemia due to an occult renal adenoma. *Pediatr Nephrol* 11:339–342
45. Kimura S, Nishimura Y, Yamaguchi K, Nagasaki K, Shimada K, Uchida H 1990 A case of pheochromocytoma producing parathyroid hormone-related protein and presenting with hypercalcemia. *J Clin Endocrinol Metab* 70:1559–1573
46. Mune T, Katakami H, Kato Y, Yasuda K, Matsukura S MK 1993 Production and secretion of parathyroid hormone-related protein in pheochromocytoma: a participation of an α -adrenergic mechanism. *J Clin Endocrinol Metab* 76:757–762
47. Wall C, Gaffney E, Mellotte G 2000 Hypercalcemia and acute interstitial nephritis associated with omeprazole therapy. *Nephrol Dial Transplant* 15:1450–1452
48. McPherson M, Prince S, Atamer E, Maxwell D, Ross-Clunis H, Estep H 1986 Theophylline-induced hypercalcemia. *Ann Intern Med* 105:52–54
49. Knox J, Demling R, Wilmore D, Sarraf P, Santos A 1995 Hypercalcemia associated with the use of human growth hormone in an adult surgical intensive care unit. *Arch Surg* 130:442–445
50. Shike M, Sturtridge WC, Tam CS, Harrison JE, Jones G, Murray TM, Husdan H, Whitwell J, Wilson DR, Jeejeebhoy KN 1981 A possible role of vitamin D in the genesis of parenteral-nutrition-induced metabolic bone disease. *Ann Intern Med* 95:560–568
51. Cathebras P, Cartry O, Lafate-Proust M, Lauwers A, Acquart S, Thomas T, Rousset H 1996 Arthritis, hypercalcemia, and lytic bone lesions after hepatitis B vaccination. *J Rheumatol* 23:558–560
52. Gayet S, Ville E, Durand J, Mars ME, Morange S, Kaplanski G, Gallais H, Soubeyrand J 1997 Foscarnet-induced hypercalcemia in AIDS. *AIDS* 11:1068–1070
53. Chandra S, Seth P, Mankeshwar J 1974 Manganese poisoning: Clinical and biochemical observations. *Environ Res* 7:374–380
54. Jurney TH 1984 Hypercalcemia in a patient with eosinophilic granuloma. *Am J Med* 76:527–528
55. Ryzen E, Singer F 1985 Hypercalcemia in leprosy. *Arch Intern Med* 145:1305–1306
56. Aly E-S, Baig M, Khanna D, Baumann M 1999 Hypercalcemia: a clue to mycobacterium avium intracellulare infection in a patient with AIDS. *International Journal of Clinical Practice* 53:227–228
57. Zaloga G, Chernow B, Eil C 1985 Hypercalcemia and disseminated cytomegalovirus infection in the acquired immunodeficiency syndrome. *Ann Intern Med* 102:331–333
58. Stoekle J, Hardy H, Weber A 1969 Chronic beryllium disease. Long term follow-up of sixty cases and selective review of the literature. *Am J Med* 46:545–561
59. Dockbell D, Poland G 1997 Hypercalcemia in a patient with hypoparathyroidism and Nocardia asteroides infection: a novel observation. *Mayo Clin Proc* 72:757–760
60. Albitar S, Genin R, Feu-Chong M, Schohn D, Riviere JP, Serveaux MO, Chuet C, Bourgeon B 1974 Multisystem granulomatous injuries 28 years after paraffin injection. *Nephrol Dial Transplant* 12:1974
61. Malik G 1998 High serum calcium in human brucellosis: a case-control study. *Am J Trop Med Hygiene* 59:397–398
62. Wong R, Gregory R, Lo T 2000 A case of isolated ACTH deficiency presenting with hypercalcemia. *Int J Clin Pract* 54:623–624
63. Suzuki K, Nonaka K, Ichihara K, Fukumoto Y, Miyazaki A, Yamada Y, Itoh Y, Seino Y, Moriawaki K, Tarui S 1986 Hypercalcemia in glucocorticoid withdrawal. *Endocrinol Jpn* 33:203–209
64. Connors M, Sheikholeslam B, Irias J 1976 Vitamin D toxicity after dieting in hypoparathyroidism. *Pediatrics* 57:794–797
65. Gerhardt A, Greenberg A, Reilly J, Van Thiel D 1987 Hypercalcemia, a complication of advanced chronic liver disease. *Arch Intern Med* 147:274–277
66. Byrne C, Bermann L, Cox T 1997 Pathological bone fractures preceded by sustained hypercalcemia in type 1 Gaucher disease. *J Inherited Metab Dis* 20:709–710
67. Berar-Yanay N, Weiner P, Magadie R 2001 Hypercalcemia in systemic lupus erythematosus. *Clin Rheumatol* 20:147–149
68. Schurman S, Bergstrom W, Root A, Sonid A, Hannah W 1998 Interleukin 1 β -mediated calcitropic activity in serum of children with juvenile rheumatoid arthritis. *J Rheumatol* 25:161–165

69. Greenwald R, Stein B, Miller F 1998 Rapid skeletal turnover and hypercalcemia associated with markedly elevated interleukin-6 levels in a young black man. *Bone* 22:285–288
70. Roux S, de Vernejoul M-C, Gossin M, DeBandt M, Palazzo E, N'Zue MK, Meyer O, Kahn M-F 1997 An unusual case of diffuse focal osteolysis with disseminated osteoclastosis. *J Clin Rheumatol* 3:58–61
71. Schipani E, Kruse K, Juppner H 1995 A constitutively active mutant PTH-PTHrP receptor in Jansen-type metaphyseal chondrodysplasia. *Science* 268:98–100
72. Parfitt A, Schipani E, Sudhaker Rao D, Kupin W, Han Z-H, Juppner H 1996 Hypercalcemia due to constitutive activity of the parathyroid hormone (PTH)/PTH-related peptide receptor: comparison with primary hyperparathyroidism. *J Clin Endocrinol Metab* 81:3584–3588
73. Barcia J, Strife C, Langman C 1997 Infantile hypophosphatasia: treatment options to control hypercalcemia, hypercalciuria, and chronic bone demineralization. *J Pediatr* 130:825–828
74. Nickerson E, Greenberg F, Keating M, McCaskill C, Shaffer L 1995 Deletions of the elastin gene at 7q11.23 occur in approximately 90% of patients with Williams syndrome. *Am J Hum Genet* 56:1156–1161
75. Kruse K, Pankau R, Gosch A, Wohlfahrt K 1992 Calcium metabolism in Williams-Beuren syndrome. *J Pediatr* 121:902–907
76. Yamaguchi K, Grant J, Noble-Jamieson G, Jamieson N, Barnes N, Compston J 1995 Hypercalcemia in primary oxalosis: role of increased bone resorption and effects of treatment with pamidronate. *Bone* 16:61–67
77. Saarela T, Simila S, Kolvisto M 1995 Hypercalcemia and nephrocalcinosis in patients with congenital lactase deficiency. *J Pediatr* 127:920–923
78. Andreoli S, Revkees S, Bull M 1995 Hypercalcemia, hypercalciuria, medullary nephrocalcinosis, and renal insufficiency in a toddler with Down syndrome. *Pediatr Nephrol* 9:673
79. Ramage I, Durkan A, Walker K, Beattie T 2002 Hypercalcemia in association with trisomy 21 (Down's syndrome). *J Clin Pathol* 55:543–544
80. Lightwood R, Butler N 1963 Decline in primary infantile renal acidosis: aetiological implications. *Br Med J* 1:885–887
81. Leumann E, Steinmann B 1975 Persistent and transient distal renal tubular acidosis with bicarbonate wasting. *Pediatr Res* 9:767–773
82. Igarashi T, Sekine Y, Kawato H, Kamoshita S, Saigusa Y 1992 Transient neonatal distal renal tubular acidosis with secondary hyperparathyroidism. *Pediatr Nephrol* 6:267–269
83. Rodriguez-Soriano J, Garcia-Fuentes M, Vallo A, Alvarez-Granda J 2000 Hypercalcemia in neonatal distal renal tubular acidosis. *Pediatr Nephrol* 14:354–355
84. Lyon A, McIntosh N, Wheeler K, Brook O 1984 Hypercalcaemia in extremely low birthweight infants. *Arch Dis Child* 59:1141–1144
85. Baylink D, Wergedal J, Stauffer M 1971 Formation, mineralization, and resorption of bone in hypophosphatemic rats. *J Clin Invest* 50:2519–2530
86. Balfour E, Antaya R, Lazova R 2002 Subcutaneous fat necrosis of the newborn presenting as a large plaque with lobulated cystic areas. *Cutis* 70:169–173
87. Tan C, Garabedian M, Farriaux J, Czernichow P, Porriarede R, Balsan S 1986 Hypercalcemia in infants with congenital hypothyroidism and its relation to vitamin D and thyroid hormones. *J Pediatr* 109:808–814
88. Ladenson J, Lewis J, McDonald J, Slatopolsky E, Boyd J 1978 Relationship of free and total calcium in hypercalcemic conditions. *J Clin Endocrinol Metab* 48:393–397
89. Howard M, Ashwell S, Bond L, Holbrook I 2000 Artefactual serum hyperkalcaemia and hypercalcaemia in essential thrombocythaemia. *J Clin Pathol* 53:105–109
90. Side L, Fahie-Wilson M, Mills M 1995 Hypercalcaemia due to calcium binding IgM paraprotein in Waldenstrom's macroglobulinemia. *J Clin Pathol* 48:961–962
91. Jaffe J, Mosher D 1979 Calcium binding by a myeloma protein. *Am J Med* 67:343–346
92. Annesley T, Burritt M, Kyle R 1982 Artifactual hypercalcemia in multiple myeloma. *Mayo Clin Proc* 57:572–575
93. Merlini G, Fitzpatrick L, Siris E, Bilezikian JP, Birken S, Beychok S, Osserman EF 1984 A human myeloma immunoglobulin G binding four molecules of calcium associated with asymptomatic hypercalcemia. *J Clin Immunol* 4:185–196

JCEM is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.