

Screening for Endocrine Hypertension: An Endocrine Society Scientific Statement

William F. Young Jr.,¹ David A. Calhoun,³ Jacques W.M. Lenders,^{4,5} Michael Stowasser,^{6,7,8} and Stephen C. Textor²

¹Division of Endocrinology, Diabetes, Metabolism, Nutrition, and Internal Medicine, and ²Division of Nephrology and Hypertension, Mayo Clinic, Rochester, Minnesota 55905; ³Vascular Biology and Hypertension Program, Division of Cardiovascular Disease, Department of Medicine, University of Alabama at Birmingham, Birmingham, Alabama 35294; ⁴Department of Internal Medicine, Radboud University Medical Centre, Nijmegen, The Netherlands 6525GA; ⁵Department of Internal Medicine III, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany; and ⁶Endocrine Hypertension Research Centre, University of Queensland School of Medicine, ⁷Greenslopes Private Hospital, and ⁸Princess Alexandra Hospital, Queensland 4102, Australia

Hypertension may be the initial clinical presentation for at least 15 endocrine disorders. An accurate diagnosis of endocrine hypertension provides clinicians with the opportunity to render a surgical cure or to achieve an optimal clinical response with specific pharmacologic therapy. It is challenging for the clinician to know when and how to perform case-detection testing for all the endocrine disorders in which hypertension may be the presenting symptom. Herein, we review the different forms of endocrine hypertension, with a focus on prevalence, clinical presentation, guidance on when to perform case detection testing, and currently available case-detection tests. (*Endocrine Reviews* 38: 103–122, 2017)

- | | |
|--|---|
| <ul style="list-style-type: none"> I. Introduction II. Pheochromocytoma and Paraganglioma <ul style="list-style-type: none"> A. Introduction and prevalence B. Clinical presentation C. Who should be screened? D. Case-detection tests <ul style="list-style-type: none"> Preanalytical conditions Reference values Interpretation of test results III. Primary Aldosteronism <ul style="list-style-type: none"> A. Prevalence B. Clinical presentation C. Who should be screened? D. Case-detection tests <ul style="list-style-type: none"> Plasma potassium The aldosterone/renin ratio IV. Other Forms of Mineralocorticoid Excess or Effect <ul style="list-style-type: none"> A. Congenital adrenal hyperplasia <ul style="list-style-type: none"> 11β-Hydroxylase deficiency <ul style="list-style-type: none"> Prevalence Clinical presentation Who should be screened? Case-detection tests 17α-Hydroxylase deficiency | <ul style="list-style-type: none"> Prevalence Clinical presentation Who should be screened? Case-detection tests B. Deoxycorticosterone-producing tumor <ul style="list-style-type: none"> Prevalence Clinical presentation Who should be screened? Case-detection tests C. Primary cortisol resistance <ul style="list-style-type: none"> Prevalence Clinical presentation Who should be screened? Case-detection tests D. Apparent mineralocorticoid excess syndrome <ul style="list-style-type: none"> Prevalence Clinical presentation Who should be screened? Case-detection tests E. Liddle syndrome: abnormal renal tubular ionic transport <ul style="list-style-type: none"> Prevalence Clinical presentation Who should be screened? |
|--|---|

ISSN Print 0163-769X ISSN Online 1945-7189

Printed in USA

Copyright © 2017 Endocrine Society

Received 24 February 2017. Accepted 27 February 2017.

Abbreviations: ACE, angiotensin-converting enzyme; ACTH, corticotropin; BP, blood pressure; ARR, aldosterone/renin ratio; CAH, congenital adrenal hyperplasia; CPAP, continuous positive airway pressure; CT, computed tomography; DOC, deoxycorticosterone; DHEA-S, dehydroepiandrosterone-sulfate; HSD11B2, 11 β -hydroxysteroid dehydrogenase type 2; OSA, obstructive sleep apnea; PA, primary aldosteronism; PPGL, pheochromocytoma and paraganglioma; RAAS, renin-angiotensin-aldosterone system; RVH, renovascular hypertension.

- Case-detection tests
- V. Secondary Aldosteronism and Renovascular Hypertension
 - A. Definition and prevalence
 - B. Clinical presentations
 - C. Who should be screened?
 - D. Case-detection tests
 - Imaging studies
 - Endocrine testing studies
- VI. Other Endocrine Disorders Associated With Hypertension
 - A. Cushing syndrome
 - Prevalence
 - Clinical presentation
 - Who should be screened?
 - Case-detection tests
 - B. Thyroid dysfunction
 - Hyperthyroidism
 - Prevalence
 - Clinical presentation
 - Who should be screened?
 - Case-detection tests
 - Hypothyroidism
- Prevalence
- Clinical presentation
- Who should be screened?
- Case-detection tests
- C. Hypercalcemia and primary hyperparathyroidism
 - Prevalence
 - Clinical presentation
 - Who should be screened?
 - Case-detection tests
- D. Acromegaly
 - Prevalence
 - Clinical presentation
 - Who should be screened?
 - Case-detection tests
- E. Obstructive sleep apnea
 - Prevalence
 - Clinical presentation
 - Who should be screened?
 - Case-detection tests
 - Polysomnography
- VII. Summary of Overall Approach to Considering and Detecting Endocrine Hypertension
 - A. Case in point

I. Introduction

Hypertension affects 28.6% of adults in United States (1–3). In most, hypertension is primary (essential or idiopathic), but a subgroup of approximately 15% has secondary hypertension (4, 5). More than 50% of children who present with hypertension have a secondary cause (6). In young adults (<40 years old), the prevalence of secondary hypertension is approximately 30% (7). The secondary causes of hypertension include renal causes (*e.g.*, renal parenchymal disease) and endocrine causes. Hypertension may be the initial clinical presentation for at least 15 endocrine disorders (Table 1). An accurate diagnosis of endocrine hypertension provides clinicians with the opportunity to render a surgical cure or to achieve an optimal clinical response with specific pharmacologic therapy (8). Primary aldosteronism (PA) is a disorder that clinicians should consider in most patients with hypertension (9). Herein, we review the different forms of endocrine hypertension with a focus on prevalence, clinical presentation, guidance on when to perform case-detection testing, and currently available case detection tests (Fig. 1).

II. Pheochromocytoma and Paraganglioma

A. Introduction and prevalence

Pheochromocytomas and paragangliomas (PPGLs) are rare neuroendocrine chromaffin-cell tumors that usually produce catecholamines and arise from the adrenal medulla (80% to 85%) or from paravertebral ganglia of the sympathetic chain (15% to 20%),

respectively (10, 11). Paragangliomas located in the neck and the skull base are usually hormonally inactive; however, some produce dopamine. The prevalence of PPGLs in the general population is extremely low (1.5 to 1.6 per 10,000 persons). The prevalence is higher in patients who present with hypertension (20 to 60 per 10,000 patients). However, it is still a rare neoplasm that a general physician will rarely encounter in clinical practice. In some patient groups, the prevalence is much higher (*e.g.*, 500 PPGLs per 10,000 of patients who also have an incidentally discovered adrenal mass) (12). The diagnosis of PPGL is frequently missed, as the prevalence of PPGLs in autopsy studies is five per 10,000 persons (13, 14).

There are two PPGL biochemical phenotypes: adrenergic and noradrenergic tumors (15, 16). Adrenergic tumors are located in the adrenal medulla and usually produce epinephrine, metanephrine (epinephrine's major metabolite), and varying amounts of norepinephrine. Noradrenergic tumors are located either in the adrenal medulla or are extra-adrenal and produce norepinephrine (predominantly or exclusively) and normetanephrine (norepinephrine's major metabolite). The lack of epinephrine secretion in these latter tumors is due to the absence of the enzyme phenylethanolamine-*N*-methyltransferase in extra-adrenal tumors; this enzyme is essential for converting norepinephrine into epinephrine (17). The biochemical phenotype is important for several reasons. First, it can predict the type of germline mutation [*e.g.*, noradrenergic tumors are more likely to be associated with the hypoxic signaling pathway mutations

Table 1. Endocrine Causes of Hypertension**Etiology**

Adrenal-dependent causes

1. Pheochromocytoma and sympathetic paraganglioma
2. Primary aldosteronism
3. Hyperdeoxycorticosteronism
 - a. Congenital adrenal hyperplasia
 - i. 11 β -Hydroxylase deficiency
 - ii. 17 α -Hydroxylase deficiency
 - b. Deoxycorticosterone-producing tumor
 - c. Primary cortisol resistance
4. Cushing syndrome

Apparent mineralocorticoid excess/11 β -hydroxysteroid dehydrogenase deficiency

1. Genetic
2. Acquired
 - a. Licorice or carbenoxolone ingestion
 - b. Cushing syndrome

Parathyroid-dependent causes

1. Hyperparathyroidism

Pituitary-dependent causes

1. Acromegaly
2. Cushing syndrome

Secondary hyperaldosteronism

1. Renovascular hypertension

Thyroid-dependent causes

1. Hypothyroidism
2. Hyperthyroidism

Complex effects

1. Obstructive sleep apnea

([cluster 1: von Hippel Lindau and succinate dehydrogenase mutations), and adrenergic tumors are more likely to be associated with kinase signaling pathway mutations (cluster 2: multiple endocrine neoplasia type 2 and neurofibromatosis type 1)]. Second, patients with adrenergic tumors may have more paroxysmal symptoms than those with noradrenergic tumors (17, 18).

B. Clinical presentation

The clinical presentation of patients with PPGLs varies widely from no symptoms, to minor discrete symptoms, to catastrophic life-threatening clinical conditions (10, 19). In general, one of every 10 patients with PPGL is completely asymptomatic; this asymptomatic subgroup is much larger in those patients with incidentally discovered adrenal masses or those tested during family screenings. When present, the classic triad of pounding headache, profuse sweating, and palpitations occurs in spells that last from several minutes to 1 hour. There is complete relief of symptoms between spells. The frequency of spells may vary from several times a day to a few times per month, occurring either spontaneously or being provoked by a variety of physical or chemical triggers, such as general anesthesia, micturition, and medications (*e.g.*, β -adrenergic inhibitors, tricyclic antidepressants, glucocorticoids) (20, 21). Enhanced blood pressure (BP) variability is

reflected by paroxysmal hypertension in about 35% of the patients. Other patients may experience severe high BP peaks, sometimes superimposed on sustained hypertension and potentially evolving into a hypertensive crisis. These high BP surges, and the underlying episodes of tumoral catecholamine release, are thought to be responsible for the high prevalence of cardiovascular emergencies, such as myocardial infarction, stroke, and heart failure (22, 23).

C. Who should be screened?

Routine biochemical screening of all new hypertensive patients for PPGL is not cost effective and, therefore, is not recommended. Clinicians should base biochemical testing for PPGL on clues that suggest the possibility of excess catecholamine secretion. In particular, clinicians should consider any paroxysmal signs or symptoms as a red flag. Since missing the diagnosis can have fatal consequences, even slight clinical suspicion of PPGL should prompt biochemical testing, irrespective of BP level (Table 2). For example, most patients with an incidentally discovered adrenal mass should have biochemical testing for PPGL (24). In addition, patients with known PPGL-causing germline mutations should have periodic biochemical testing (25). Clinicians should also consider case-detection testing in family members of newly diagnosed patients with PPGL.

D. Case-detection tests

Biochemical testing should generally precede imaging procedures because only solid evidence of excess production of catecholamines can justify performing expensive imaging procedures. Initial biochemical testing for PPGLs should include measuring plasma free or urinary fractionated metanephrines (25). Because of their very high sensitivities, normal values for these tests exclude a PPGL with high reliability. Exceptions include very small tumors that may be missed, as might be the case in asymptomatic patients tested because of a hereditary predisposition. Diagnostic accuracy is not significantly different between plasma and 24-hour urinary tests, particularly when assays use liquid chromatography with tandem mass spectrometry (26, 27). The explanation for this high sensitivity is the continuous secretion of intratumorally produced metanephrines, which contrasts with the episodic exocytotic secretion of their parent catecholamines. Conversely, elevated test results of fractionated metanephrines are not fully specific for the presence of a PPGL and, therefore, do not prove that a patient has a PPGL.

Other available biochemical tests, such as those for plasma or urinary catecholamines, urinary vanillylmandelic acid, and urinary chromogranin A, have inferior diagnostic value as compared with plasma free and urinary fractionated metanephrines.


Figure 1.

Figure 1. When to consider and how to test for endocrine hypertension. ARB, angiotensin II receptor blockers; ASO, arterial switch operation; CT, computerized tomography; CVA, cerebrovascular accident; DBP, diastolic blood pressure; DST, dexamethasone suppression test; FMD, fibromuscular dysplasia; FHx, family history; MR, magnetic resonance; SBP, systolic blood pressure; UFC, urinary free cortisol.

Preanalytical conditions

Clinicians should take several precautions regarding biochemical testing. They should assess coexisting conditions associated with increased sympathetic activity, such as occult or overt heart failure, renal failure, and

Table 2. When To Perform Biochemical Testing To Screen for Pheochromocytoma or Sympathetic Paranglioma

Indications for Biochemical Testing

Paroxysmal signs or symptoms suggesting catecholamine excess

Paradoxical BP response to drugs, surgery, or anesthesia

Resistant hypertension

Incidentally discovered adrenal mass (with or without hypertension)

Previous diagnosis of pheochromocytoma or paraganglioma (with annual biochemical testing to detect recurrent disease)

Hereditary predisposition for pheochromocytoma or paraganglioma

Syndromic feature indicating a pheochromocytoma-related hereditary syndrome

Adapted from Lenders et al. (25).

hypoglycemia. Increased sympathetic activity may result in false-positive test results, thus limiting the positive predictive value of a positive test result. A more commonly occurring example is clinicians sampling blood from patients who are in the seated position and have not had preceding rest. To minimize the risk of false-positive test results of plasma free metanephrines, clinicians should draw blood samples from an indwelling cannula after patients have had supine rest for at least 30 minutes. If taking a blood sample after supine rest is not feasible, measuring 24-hour urinary fractionated metanephrines provides an excellent alternative test.

Patients who consume biogenic amines may also exhibit false-positive test results. However, measuring plasma free metanephrines requires no special dietary precautions. The only exception is the O-methylated dopamine metabolite methoxytyramine, which can be spuriously elevated when blood is sampled from patients who are in a nonfasting state (28). Elevated levels of this metabolite have some value in predicting extra-adrenal tumor location (in particular, neck and skull-base paragangliomas) and the presence of metastatic disease (29).

When measuring plasma metanephrines, clinicians should collect blood samples in heparinized tubes or tubes with ethylenediaminetetraacetic acid and place these immediately on ice to prevent oxidative degradation (30). When measuring 24-hour urinary fractionated metanephrines, clinicians can use containers without additives. Before storage, technicians should acidify the urine sample to a pH of 4 when it arrives in the laboratory (31). Clinicians should instruct patients carefully about how to collect urine for 24 hours. Measuring urinary creatinine is a useful way to verify that urine collection was complete.

Finally, several drugs can be a source of false-positive test results due to analytical or pharmacophysiological interference (Table 3). Liquid chromatography with tandem mass spectrometry assays are more specific and experience less from analytical interference by drugs than liquid chromatography with electrochemical detection assays.

Reference values

To obtain optimal diagnostic accuracy, each laboratory should establish reference values for plasma free and 24-hour urinary fractionated metanephrines, taking into account patient-specific characteristics, such as age and sex. Specifically, the upper cutoff level of plasma free normetanephrine is higher in older patients (32, 33). There is no such effect for metanephrine or methoxytyramine.

Interpretation of test results

If a test result of plasma free or 24-hour urinary fractionated metanephrines in a symptomatic patient is within the normal range, clinicians can exclude a PPGL, provided that urine collection is complete and blood sampling was performed correctly. In asymptomatic patients, as in those who are carriers of one of the susceptibility genes, a single normal test result cannot definitely exclude a PPGL; in these cases it is preferable to wait and retest.

Plasma free metanephrine levels more than three times the upper reference limit or 24-hour urinary fractionated metanephrine levels more than two times the upper reference limit are highly reliable predictors of the presence of a PPGL. In these cases, the risk of a falsely elevated test result is very low and, in such cases, clinicians can proceed with imaging studies. However, 25% of patients have levels for the above indicators that are only slightly elevated, and these patients are more difficult to analyze. For example, patients in the intensive care unit routinely have falsely elevated test results (34). Thus, biochemical testing for PPGL should be avoided in critically ill patients. Clinicians should

Table 3. Medications That Can Cause False-Positive Biochemical Testing for Pheochromocytoma and Sympathetic Paraganglioma

Medication	Plasma		Urine	
	NMN	MN	NMN	MN
Tricyclic antidepressants ^a	↑↑	—	↑↑	—
Antipsychotics ^a	↑↑	—	↑↑	—
Buspirone ^a	↑	↑↑	↑	↑↑
MAO inhibitors ^a	↑↑	↑↑	↑↑	↑↑
Sympathomimetics ^a	↑	↑	↑	↑
Cocaine ^a	↑↑	↑	↑↑	↑
Levodopa ^b	↑	↑	↑↑	↑
Phenoxylbenzamine ^c	↑↑	—	↑↑	—
Acetaminophen ^d	↑↑	—	↑↑	—
Labetalol ^d	—	—	↑↑	↑↑
Sotalol ^d	—	—	↑↑	↑↑
α-methyl dopa ^d	↑↑	—	↑↑	—
Sulphasalazine ^d	↑↑	—	↑↑	—

Adapted from Lenders *et al.* (25).

Abbreviations: —, no change; MAO, monoamine oxidase; MN, metanephrine; NMN, normetanephrine.

^aPharmacodynamic interference affects all analytical methods.

^bThere is analytical interference with some liquid chromatography with electrochemical detection assays and pharmacodynamic interference increases measured levels of dopamine and the dopamine metabolite methoxytyramine.

^cThere is pharmacodynamic interference during the first few days of medication initiation.

^dThere is analytical interference for some but not all methods using liquid chromatography with electrochemical detection; these drugs do not cause false-positive testing in tandem mass spectroscopy assays.

first check for comorbidities associated with stress-mediated increased sympathetic activity in all patients who exhibit a slightly elevated level of plasma free metanephrine or 24-hour urinary fractionated metanephrine.

Another much more frequent condition associated with mildly elevated plasma metanephrine levels is sampling blood from patients in the seated position who have not had preceding rest. In such cases, clinicians should remeasure plasma metanephrine levels after patients have had 30 minutes' supine rest. If feasible, clinicians should discontinue medications known to cause false-positive test results (*e.g.*, tricyclic antidepressants) before blood or urine collection. If this is not possible, clinicians should administer tests irrespective of drug treatment, but take special care when interpreting test results.

When clinicians have excluded all obvious causes of false-positive test results, other less obvious possibilities remain. As a follow-up test, clinicians can measure both urinary fractionated metanephrine and plasma chromogranin A levels to exclude falsely elevated plasma metanephrine levels (35). A more laborious test is the

clonidine suppression test for distinguishing truly elevated (tumoral normetanephrine release) from falsely elevated (sympathetic activation) test results (36).

III. Primary Aldosteronism

In PA, aldosterone production exceeds the body's requirements and is relatively unchecked by its normal regulator, the renin-angiotensin II system (37, 38). Excessive unregulated production of aldosterone results in increased sodium reabsorption via amiloride-sensitive epithelial sodium channels within the distal nephron, leading to hypertension and renin-angiotensin II suppression. Urinary loss of potassium and hydrogen ions, exchanged for sodium at the distal nephron, may result in hypokalemia and metabolic alkalosis (37, 38).

A. Prevalence

Once thought to be a rare condition and not worth investigating in patients with hypertension unless hypokalemic, PA is now considered the most common, specifically treatable, and potentially curable form of hypertension, accounting for at least 5% to 10% of hypertensive patients, with most patients normokalemic (39, 40). In resistant hypertensive cohorts, the prevalence of PA is approximately 20% (41, 42). Most patients with PA are diagnosed during their third to sixth decades (43).

B. Clinical presentation

Hypertension occurs in most patients with PA and may be mild or severe but rarely malignant (44, 45). BP levels vary widely among patients with either aldosterone-producing adenoma or bilateral adrenal hyperplasia, and clinicians cannot use BP levels to distinguish these subtypes (46). In familial hyperaldosteronism type I, hypertension is often delayed, especially in female patients, but can be of early onset and severe enough to cause early death, usually from hemorrhagic stroke (47, 48). Family screening in familial hyperaldosteronism type I and familial hyperaldosteronism type II has revealed highly diverse phenotypes, with some patients being normotensive, which is consistent with PA evolving through a preclinical phase (47, 49–51).

Less than one-quarter of patients diagnosed with PA and less than one-half of those with aldosterone-producing adenoma are hypokalemic (39, 40, 52). In these patients, PA is indistinguishable from essential hypertension unless clinicians measure renin and aldosterone. When hypokalemia does occur, it may be associated with nocturia, polyuria, muscle weakness, cramps, paresthesias, and/or palpitations. The prevalence of obstructive sleep apnea (OSA) is increased in patients with PA and it improves with PA-specific therapy (53).

During pregnancy, hypertension and symptoms may improve or become worse. Improvement appears to be due to the antimineralocorticoid effects of high circulating levels of placental progesterone, which antagonize aldosterone action at the mineralocorticoid receptor (54, 55). Studies have reported increased luteinizing hormone choriogonadotropin-receptor expression in aldosterone-producing adenomas harboring β -catenin mutations in some pregnant women, and the increased pregnancy-related blood levels of human chorionic gonadotropin, in turn, increased the degree of hyperaldosteronism (56, 57).

C. Who should be screened?

Clinicians should consider PA screening for most patients with hypertension. In part, this is because hypertension in PA responds well to specific treatments directed against aldosterone excess. Unilateral laparoscopic adrenalectomy in patients with unilateral forms of PA cures hypertension in 50% to 60% of cases and results in significant improvements in the remainder (58–60). For nonoperated patients, medications that antagonize aldosterone action (*e.g.*, spironolactone, eplerenone, and amiloride) lead to substantial and often marked improvements in hypertension control (61, 62). Furthermore, it has become apparent that aldosterone excess in PA directly induces injury (*e.g.*, inflammation, remodeling, and fibrosis) in cardiovascular and renal tissues and induces adverse metabolic effects in ways that are at least partly independent of its effect on BP (63–66). As a result, rates of cardiovascular events (*e.g.*, arrhythmias, myocardial infarctions, strokes, and cardiovascular mortality) are higher in patients with PA than among those with essential hypertension matched for BP level (67–69). Importantly, the excess in cardiovascular morbidity is reversed after treatment (67), providing compelling support for early detection of individuals with PA.

The Endocrine Society clinical practice guideline on PA recommends the case detection of PA in patient groups with relatively high prevalence of PA (9). These include the following: patients with a sustained elevated BP [BP ≥ 150 mm Hg (systolic) and/or 100 mm Hg (diastolic)]; hypertension (BP $> 140/90$ mm Hg) resistant to three conventional antihypertensive drugs, including a diuretic; controlled BP (BP $< 140/90$ mm Hg) with four or more antihypertensive drugs; hypertension and spontaneous or diuretic-induced hypokalemia; hypertension and adrenal incidentaloma; hypertension and OSA; or hypertension and a family history of early-onset hypertension or cerebrovascular accident at a young age (< 40 years old). Given the known existence of familial forms, the guideline also recommends case detection for all hypertensive first-degree relatives of patients with PA (9).

D. Case-detection tests

Plasma potassium

Because only a minority (approximately 20%) of patients with PA are hypokalemic, measurements of plasma potassium lack sensitivity as a screening test (39). However, when hypokalemia is present (especially when not provoked by the use of diuretics), plasma potassium levels serve as a valuable clue of the presence of this condition.

The aldosterone/renin ratio

The aldosterone/renin ratio (ARR) is the most reliable available screening test; it is more specific than renin measurement (levels of which are almost always suppressed in patients with PA) and more sensitive than plasma potassium or aldosterone measurements. The ratio becomes elevated before aldosterone or plasma potassium leave their normal ranges (52). However, false positives and negatives are possible and can occur for the following reasons.

- Hypokalemia may be associated with false-negative ARR because potassium is a powerful chronic regulator of aldosterone secretion (44, 70).
- False-positive ARR can occur in premenopausal women during the luteal phase of the menstrual cycle and also in women receiving estrogen-containing contraceptive agents, but only when clinicians measure renin as direct active renin concentration and not as plasma renin activity (71, 72).
- False-positive results may also be seen in patients with impaired renal function (renin production is reduced, whereas any associated hyperkalemia tends to elevate aldosterone) (73), in advancing age (during which production of renin falls more quickly than that of aldosterone), and in familial hyperkalemic hypertension (also known as pseudohypoaldosteronism type II and Gordon syndrome) (74, 75).
- False-negative ARR can result when renin secretion is stimulated by the following: dietary salt restriction; concomitant malignant or renovascular hypertension (RVH); pregnancy (in which high levels of progesterone antagonize aldosterone action at the mineralocorticoid receptor); and treatment with diuretics (including spironolactone), dihydropyridine calcium channel antagonists, angiotensin-converting enzyme (ACE) inhibitors, or angiotensin receptor antagonists (70, 75–78).
- β -adrenergic blockers, α -methyldopa, clonidine, and nonsteroidal anti-inflammatory drugs can suppress renin levels and produce false-positive ARR (70, 75, 77, 79).
- Treating with antidepressants of the selective serotonin reuptake inhibitor class lowers the ARR, but whether

they can cause false-negative findings in patients with PA remains uncertain (80).

Clinicians should withdraw diuretics for at least 4 weeks and other interfering medications for at least 2 (and preferably 4) weeks before measuring ARR. To maintain BP control, clinicians should administer other medications that have a lesser effect on results, such as verapamil slow-release (with or without hydralazine) and prazosin (70, 75, 78). In cases where clinicians cannot withdraw a potentially interfering medication, they should take into account the medication's known effects when interpreting ratios. For example, an elevated ratio in patients receiving a diuretic, ACE inhibitor, angiotensin receptor blocker, or dihydropyridine calcium blocker would make PA very likely, whereas a normal ratio in the presence of β -adrenergic blocker treatment would make the diagnosis unlikely.

Clinicians should correct hypokalemia and encourage the patient to follow a liberal sodium diet before measuring ARR. Because of the effects of posture and time of day, collecting blood midmorning from seated patients who have been upright (*i.e.*, sitting, standing, or walking) for 2 to 4 hours maximizes the sensitivity of the ratio (70, 75, 78).

Clinicians should regard the ARR as a screening test only and should measure it more than once (serially if conditions of sampling, including medications, are being altered) before deciding whether to go on to a suppression test to definitively confirm or exclude the diagnosis (70, 75, 78). Although reference ranges and cutoffs differ depending on the laboratory, plasma aldosterone concentrations >10 ng/dL in concert with plasma renin activity <1 ng/mL/h should trigger confirmatory testing (9). Clinicians should consider referring those patients who have positive case detection of PA to an endocrinologist.

IV. Other Forms of Mineralocorticoid Excess or Effect

Table 1 lists medical disorders associated with mineralocorticoid excess resulting from deoxycorticosterone (DOC) or cortisol. Clinicians should consider these disorders if blood levels of aldosterone and renin are low in a patient with hypertension and hypokalemia (81).

A. Congenital adrenal hyperplasia

Congenital adrenal hyperplasia (CAH) is a group of autosomal recessive disorders caused by enzymatic defects in adrenal steroidogenesis that result in the deficient secretion of cortisol (82, 83). The lack of inhibitory feedback by cortisol on the hypothalamus and pituitary produces a corticotropin (ACTH)-driven buildup of cortisol precursors proximal to the enzymatic deficiency.

Approximately 90% of CAH cases are caused by 21-hydroxylase deficiency, which does not result in hypertension (84). Deficiencies of 11 β -hydroxylase (CYP11B1, P450c11) or 17 α -hydroxylase (CYP17, P450c17) cause hypertension and hypokalemia because of hypersecretion of the mineralocorticoid DOC. The mineralocorticoid effect of increased circulating levels of DOC also decreases renin and aldosterone secretion. These mutations are autosomal recessive in inheritance and are typically diagnosed in childhood. However, partial enzymatic defects have been shown to cause hypertension in adults.

11 β -Hydroxylase deficiency

Prevalence. 11 β -hydroxylase deficiency causes approximately 5% of all cases of CAH; the prevalence in whites is one in 100,000 (85). Research has described more than 40 mutations in *CYP11B1* (the gene encoding 11 β -hydroxylase) (86). There is an increased prevalence among Sephardic Jews from Morocco, suggestive of a founder effect.

Clinical presentation. The impaired conversion of DOC to corticosterone results in high levels of DOC and 11-deoxycortisol; the substrate mass effect results in increased levels of adrenal androgens. Girls with CAH present in infancy or childhood with hypertension, hypokalemia, acne, hirsutism, and virilization. Boys with CAH due to 11 β -hydroxylase deficiency present with hypertension, hypokalemia, and pseudoprecocious puberty. Approximately two-thirds of patients have mild to moderate hypertension.

Who should be screened? Clinicians should screen children, adolescents, and young adults who present with hypertension, spontaneous hypokalemia, and low levels of aldosterone and renin. Suspicion for this defect should be highest in girls with virilization and boys with pseudoprecocious puberty.

Case-detection tests. The initial screening tests include measuring blood levels of DOC, 11-deoxycortisol, androstenedione, testosterone, and dehydroepiandrosterone-sulfate (DHEA-S), all of which should be increased above the upper limit of their respective reference ranges (Table 4). If levels of these hormones are above their reference ranges in patients with hypertension and hypokalemia, clinicians should pursue confirmatory testing, including germline mutation testing.

17 α -Hydroxylase deficiency

Prevalence. 17 α -Hydroxylase deficiency is a very rare cause of CAH. Good prevalence data are not available, but prevalence is likely less than one in 1,000,000 live births (87).

Clinical presentation. 17 α -Hydroxylase is essential for the synthesis of cortisol and gonadal hormones, and deficiency results in decreased production of cortisol and sex steroids. Genetic 46,XY male patients present with either pseudohermaphroditism or phenotypically as female, and 46,XX female patients present with primary amenorrhea. Therefore, a person with this form of CAH may not come to medical attention until puberty.

Who should be screened? Clinicians should screen children, adolescents, and young adults who present with hypertension and spontaneous hypokalemia and low levels of aldosterone and renin. Suspicion for this mutation should be highest in the setting of primary amenorrhea and pseudohermaphroditism. Although very rare, there is an increased prevalence of 17 α -hydroxylase deficiency among Dutch Mennonites.

Case-detection tests. The initial screening tests include measuring blood levels of androstenedione, testosterone, DHEA-S, 17-hydroxyprogesterone, aldosterone, and cortisol—all of which should be either low or at the lower quartile of their respective reference ranges. The plasma concentrations of DOC and corticosterone should be above the upper limit of their respective reference ranges (Table 4). If levels of these hormones are above their reference ranges in patients with hypertension and hypokalemia, clinicians should pursue confirmatory testing, including germline mutation testing.

B. Deoxycorticosterone-producing tumor

Prevalence

Pure DOC-producing adrenal tumors are very rare and are usually large and malignant (88). Some studies have reported patients with benign DOC-producing adrenocortical adenomas (89).

Clinical presentation

Some of these adrenal neoplasms cosecrete androgens and estrogens in addition to DOC, which may cause virilization in women or feminization in men. The typical clinical presentation would be relatively rapid onset of marked hypertension associated with hypokalemia and low blood levels of aldosterone and renin. There is no specific age group that has an increased prevalence of these rare tumors.

Who should be screened?

Clinicians should screen patients who present with hypertension, spontaneous hypokalemia, and low levels of aldosterone and renin.

Table 4. Adrenal-Related Test Findings in Patients With Nondosterone-Mediated Mineralocorticoid Excess or Effect^a

Disorder	24-h UFC	Urinary Cortisol: Cortisone Ratio	DOC	11-Deoxy-cortisol	Androstenedione	DHEA-S
11 β -hydroxylase deficiency	↓	—	↑↑↑	↑↑↑	↑↑↑	↑↑↑
17 α -hydroxylase deficiency	↓	—	↑	↑	↓	↓
DOC-producing tumor	—	—	↑↑↑	—	—	—
Primary cortisol resistance	↑↑	—	↑	↑	↑	↑
Apparent mineralocorticoid excess syndrome	↑↑↑↑ ^b	↑↑↑↑	↑	—	—	—

Abbreviations: —, no change; UFC, urinary free cortisol.

^aAll of these patients have low or undetectable levels of aldosterone and renin.

^b24-Hour UFC excretion is markedly increased if the apparent mineralocorticoid excess is due to severe Cushing syndrome (e.g., ectopic ACTH syndrome).

Case-detection tests

A high level of plasma DOC or urinary tetrahydrodeoxycorticosterone and a large adrenal tumor appearing on computed tomography (CT) confirm the diagnosis. Aldosterone secretion in these patients is typically suppressed.

C. Primary cortisol resistance

Prevalence

Patients with primary cortisol (glucocorticoid) resistance, a rare familial syndrome, can have increased cortisol secretion and plasma cortisol concentrations without evidence of Cushing syndrome (90, 91). Genetic defects in the glucocorticoid receptor and the steroid-receptor complex cause primary cortisol resistance.

Clinical presentation

The syndrome is characterized by hypokalemic alkalosis, hypertension, increased plasma concentrations of DOC, and increased adrenal androgen secretion. Hypertension and hypokalemia are due to the combined effects of excess DOC and increased cortisol access to the mineralocorticoid receptor, resulting from high rates of cortisol production that overwhelm 11 β -hydroxysteroid dehydrogenase type 2 (HSD11B2) activity.

Who should be screened?

Clinicians should screen patients (primarily children) who present with hypertension, spontaneous hypokalemia, and low levels of aldosterone and renin.

Case-detection tests

The initial screening tests include measuring blood levels of cortisol, DOC, 11-deoxycortisol, androstenedione, testosterone, and DHEA-S—all of which are usually above the upper limit of their respective reference ranges (Table 4). In addition, 24-hour urinary cortisol excretion is above the upper limit of the reference range and serum ACTH is not suppressed. Confirmatory testing includes germline mutation testing.

D. Apparent mineralocorticoid excess syndrome

Apparent mineralocorticoid excess is the result of impaired activity of the microsomal enzyme HSD11B2, which normally inactivates cortisol in the kidney by converting it to cortisone (an inactive 11-keto compound) (92). Cortisol can be a potent mineralocorticoid, and when HSD11B2 is genetically deficient or its activity is blocked, high levels of cortisol are present in the kidney.

Prevalence

Decreased HSD11B2 activity may be hereditary or it may be secondary to the pharmacologic inhibition of enzyme activity by glycyrrhizic acid [the active principle of licorice root (*Glycyrrhiza glabra*)] (93). The congenital forms are rare autosomal recessive disorders; researchers have identified fewer than 50 of these cases worldwide (94).

Clinical presentation

Congenital apparent mineralocorticoid excess typically presents in childhood with hypertension, hypokalemia, low birth weight, failure to thrive, hypertension, polyuria and polydipsia, and poor growth (86). Acquired apparent mineralocorticoid excess due to licorice root ingestion presents with hypertension and hypokalemia; the cause is revealed by the patient's medical history. When massive cortisol hypersecretion associated with Cushing syndrome due to ectopic ACTH syndrome overwhelms HSD11B2, hypokalemic hypertension may be one of the outcomes (95).

Who should be screened?

Clinicians should screen patients with apparent mineralocorticoid excess due to congenital deficiency or HSD11B2 inhibition; these patients can have hypertension, hypokalemia, metabolic alkalosis, and low renin, low aldosterone, and normal plasma cortisol levels.

Case-detection tests

Clinicians can confirm a diagnosis of apparent mineralocorticoid excess by demonstrating an abnormally high

ratio of cortisol to cortisone in a 24-hour urine collection. The characteristic abnormal urinary cortisol-cortisone metabolite profile reflects decreased HSD11B2 activity; the ratio of cortisol to cortisone is typically increased 10-fold above the normal value (92). The mineralocorticoid excess state caused by ectopic ACTH secretion is related to the high rates of cortisol production that overwhelm HSD11B2 activity. DOC levels may also be increased in severe ACTH-dependent Cushing syndrome and contribute to hypertension and hypokalemia in this disorder.

E. Liddle syndrome: abnormal renal tubular ionic transport

In 1963, Grant Liddle described an autosomal dominant renal disorder with a presentation similar to PA that includes hypertension, hypokalemia, and inappropriate kaliuresis (96). However, blood levels of aldosterone and renin were very low, so researchers termed the disorder pseudoaldosteronism.

Prevalence

Autosomal dominant mutations in the β or γ subunit of the amiloride-sensitive epithelial sodium channel cause Liddle syndrome (86). It is extremely rare, with fewer than 30 families reported worldwide (97).

Clinical presentation

This mutation results in enhanced activity of the epithelial sodium channel, and patients present with increased renal sodium reabsorption, potassium wasting, hypertension, and hypokalemia. However, as mentioned, blood levels of aldosterone and renin are low.

Who should be screened?

Clinicians should screen children and adults who present with hypertension, spontaneous hypokalemia, and low levels of aldosterone and renin. A family history of hypertension associated with hypokalemia makes Liddle syndrome more likely.

Case-detection tests

A clinician who discovers low aldosterone and renin levels in a hypokalemic hypertensive patient should consider Liddle syndrome. Once the clinician excludes the other causes of this presentation (Table 4), they should consider treating the patient with amiloride or triamterene. The clinician can easily distinguish Liddle syndrome from apparent mineralocorticoid excess based on a marked improvement in hypertension when amiloride or triamterene are combined with a sodium-restricted diet. Other ways to distinguish Liddle syndrome from apparent mineralocorticoid excess include a lack of efficacy of spironolactone and dexamethasone, and a normal 24-hour urine cortisone/cortisol ratio. Clinical genetic testing is available.

V. Secondary Aldosteronism and Renovascular Hypertension

A. Definition and prevalence

Seminal studies more than 80 years ago established that reduced blood flow to the kidneys could initiate a rise in systemic BP, defined as RVH. Over the years, there has been extensive research regarding the renin-angiotensin-aldosterone system (RAAS) as a primary hormonal axis underlying this process, and renovascular disease has become one of the most widely studied “angiotensin-dependent” models of hypertension and vascular disease.

Secondary hyperaldosteronism reflects pathologically elevated aldosterone levels due to activation of the renin-angiotensin axis; RVH is one such instance. Other situations associated with secondary aldosterone excess include renal infarction, volume depletion with or without administration of diuretics, renal hypoperfusion related to cardiac or hepatic failure, or (rarely) primary overproduction of renin from a juxtaglomerular tumor. Secondary hyperaldosteronism associated with volume depletion or cardiac or hepatic failure is not typically associated with hypertension.

Most studies addressing the prevalence of renovascular disease focus on the frequency of identifying renal artery stenosis rather than establishing the actual occurrence of hypertension. This distinction is important because the kidney tolerates some degree of vascular occlusion without clinical manifestations. The prevalence of actual RVH is substantially less than that for renal artery stenosis. Some argue that establishing a diagnosis of RVH depends on reducing BP after technically successful revascularization or removal of the affected kidney.

Most renal artery stenosis in the United States is caused by atherosclerotic disease ($\approx 85\%$) or some form of fibromuscular dysplasia ($\approx 15\%$). The prevalence of atherosclerotic renal artery stenosis increases with age and other atherosclerotic manifestations. Population-based studies indicate that 6.8% of individuals older than 65 years have renal artery stenosis with more than 60% lumen occlusion.

Studies of image-based screening in patients undergoing angiography for other atherosclerotic diseases showed that 14% to 33% of these patients have renal artery stenosis, depending upon the extent of disease. Similarly, studies of image-based screening in potential kidney donors showed that 3% to 5% of normal subjects have some degree of fibromuscular dysplasia.

How often these lesions produce RVH and the true prevalence of RVH remain unclear, but estimates from referral hypertension centers suggest that 1% to 5% of hypertensive subjects may have a component of RVH.

B. Clinical presentations

Renovascular occlusive disease leading to RAAS activation can produce a spectrum of manifestations including RVH, accelerated/malignant phase hypertension, impaired cardiac function, circulatory congestion (“flash pulmonary edema”), and, ultimately, parenchymal kidney injury with irreversible loss of kidney function. Recognizing these manifestations and establishing the likelihood of improving BP control and cardiac and kidney function remain major challenges for clinicians. Table 5 details the clinical scenarios that warrant screening and diagnostic studies for RVH.

C. Who should be screened?

The utility of screening and diagnostic testing for RVH and other causes of secondary aldosteronism partly depends upon the commitment to act upon the results. The pressure to identify renal artery stenosis has diminished in recent years. One reason for this is that medical therapy is remarkably effective at blocking the RAAS, without the need for either endovascular or surgical revascularization. Several recent prospective randomized controlled trials (RCTs) have failed to identify major additional benefits from stent revascularization for atherosclerotic renal artery stenosis (98). Thus, it may be argued that clinicians should limit screening for these disorders to cases where medical therapy is failing or the risks of circulatory or renal failure truly warrant restoring the circulation, as identified in Table 5. Because these RCTs failed to enroll or excluded many high-risk subjects, clinicians must individualize treatment of many patients, particularly those with rapidly declining kidney function, accelerated hypertension, or episodes of pulmonary edema.

D. Case-detection tests

Imaging studies

Renal artery duplex ultrasound measurements of peak systolic velocity along the vessel path have a sensitivity above 85% and specificity of 92% for atherosclerotic disease with more than 60% lumen occlusion (99, 100). Renal artery duplex is relatively inexpensive and available; rarely has false-positive results in experienced laboratories; and provides information on kidney size, resistive index, and perfusion. However, it commonly has operator-dependent false-negative results and technical failures (101).

CT angiography has a sensitivity above 90% and specificity of 97%, similar to catheter angiography and high-resolution magnetic resonance angiography. CT angiography provides detailed imaging of the cortex and medulla with high-resolution of vessels. However, it is relatively expensive and has the potential for iodinated-contrast toxicity.

Magnetic resonance angiography provides detailed imaging and functional testing of oxygenation imaging.

Table 5. Clinical Scenarios Associated With Renovascular Disease

Clinical Scenario

Onset hypertension before age 30 years
Accelerated, resistant, malignant hypertension
Deterioration of renal function (rise in creatinine more than 30% of pretreatment levels) in response to ACE inhibitors or angiotensin-receptor blockers
New onset of hypertension after age 50 years (suggestive of atherosclerotic renal artery stenosis)
Asymmetric kidneys with more than 1.5-cm difference in size and otherwise unexplained loss of kidney function
Sudden unexplained pulmonary edema (“flash pulmonary edema”)

However, it is also relatively expensive and has the potential for gadolinium toxicity.

Radionuclide scintigraphy, with or without ACE inhibition (captopril renogram), rarely produces false-negative findings and provides limited information about large-vessel disease. However, it commonly has false-positive findings with impaired renal function.

Endocrine testing studies

Clinicians use peripheral plasma renin activity, direct renin mass measurements, and plasma aldosterone concentrations to define secondary aldosterone excess. However, these measurements are nonspecific and highly subject to conditions of testing, including sodium balance and drug therapy. Measuring plasma renin activity after administering an ACE inhibitor (*i.e.*, captopril)—a measurement referred to as “enhanced plasma renin activity”—is non-specific and has limited sensitivity and poor predictive value.

Clinicians commonly used renal vein renin measurements for diagnostic testing related to surgical revascularization. Although moderately invasive, these measurements have substantial validity for predicting BP response to either revascularization or nephrectomy (102) and allow localization of renin production in specific disorders (*e.g.*, juxtaglomerular cell tumors, renal infarction). However, they are subject to testing conditions (as are all hormonal measurements of the RAAS). Clinicians use these measurements less often in the endovascular stent era, in part because BP is less commonly the primary motivation for diagnostic testing (vs salvaging kidney function or treating recurrent pulmonary edema).

VI. Other Endocrine Disorders Associated With Hypertension

A. Cushing syndrome

Prevalence

Iatrogenic Cushing syndrome is relatively common. However, endogenous Cushing syndrome is rare, with an

incidence of less than one per 1 million people per year (103). Excess ACTH secretion by a pituitary tumor is the cause of endogenous Cushing syndrome in 85% of patients and is termed “Cushing disease.” Cushing disease occurs five times more frequently in women than in men, with the peak incidence occurring between 20 and 50 years of age. Ectopic ACTH-secreting neoplasms and the ACTH-independent forms of Cushing syndrome (*e.g.*, adrenal adenoma, adrenal carcinomas, and adrenal nodular hyperplasias) are responsible for 15% of the endogenous cases. Hypertension occurs in 75% to 80% of patients with Cushing syndrome (104, 105). The mechanisms of hypertension include increased production of DOC, enhanced pressor sensitivity to endogenous vasoconstrictors (*e.g.*, epinephrine, angiotensin II), increased cardiac output, activation of the RAAS by increased hepatic production of angiotensinogen, and cortisol activation of the mineralocorticoid receptor.

Clinical presentation

Typical signs and symptoms of Cushing syndrome include the following: weight gain with central obesity (thin extremities), facial rounding and plethora, supraclavicular fat pads, dorsocervical fat pads, easy bruising (“spontaneous”), fine “cigarette paper skin” and decreased skin-fold thickness, poor wound healing, purple striae (usually >1 cm in width), proximal muscle weakness, emotional and cognitive changes (*e.g.*, irritability, crying, depression, restlessness), hirsutism, hyperandrogenism (*e.g.*, acne), hypertension, osteopenia and osteoporosis, glucose intolerance and diabetes mellitus, polyuria, hyperlipidemia, opportunistic and fungal infections (*e.g.*, mucocutaneous candidiasis, tinea versicolor, pityriasis), menstrual dysfunction and altered reproductive function, and renal lithiasis. These clinical features may occur slowly over time. Therefore, comparing the patient’s current appearance with old photographs is invaluable. Many of these signs and symptoms are common and not distinguishing features (*e.g.*, obesity, hypertension, abnormal glucose tolerance, and menstrual dysfunction).

Who should be screened?

Clinicians should screen any patient with hypertension who presents with signs or symptoms consistent with the clinical presentation of Cushing syndrome (as outlined in the previous paragraph).

Case-detection tests

Standard laboratory studies may reveal fasting hyperglycemia, hyperlipidemia, hypokalemia, and leukocytosis with relative lymphopenia. Mineralocorticoid production is usually normal in endogenous Cushing syndrome; aldosterone and renin levels are usually

normal or suppressed, and DOC levels are normal or mildly increased. Aldosterone, DOC, and sex steroids may be elevated in adrenal carcinomas (106).

The case-detection tests for endogenous cortisol excess include a 1-mg overnight dexamethasone-suppression test and measuring midnight salivary cortisol and free cortisol in a 24-hour urine collection. When results of case-detection tests for Cushing syndrome are abnormal, clinicians should pursue confirmatory testing. The Endocrine Society’s clinical practice guideline on Cushing syndrome further details tests that confirm Cushing syndrome and determine the cause of cortisol excess (95).

B. Thyroid dysfunction

Hyperthyroidism

Prevalence. Hyperthyroidism is more common in women, with a prevalence between 0.5% and 1.0% (107–109). In the National Health and Nutrition Examination Survey III, two of 1,000 subjects who were not taking thyroid medication and not reporting histories of thyroid disease had “clinically significant” hyperthyroidism (110). Prevalence in the elderly population ranges between 0.4% and 2.0% (111–113). The most common causes include Graves’ disease, toxic multinodular goiter, toxic adenoma, and thyroiditis. There are no data on the prevalence of hyperthyroidism in patients who present with hypertension.

Clinical presentation. When excessive amounts of circulating thyroid hormones interact with thyroid hormone receptors on peripheral tissues, both metabolic activity and sensitivity to circulating catecholamines increase. Thyrotoxic patients usually have tachycardia, high cardiac output, increased stroke volume, decreased peripheral vascular resistance, and increased systolic BP (114). Clinical presentation may be dominated by weight loss (despite an increased appetite), heat intolerance, muscle weakness, and hyperhidrosis. In addition, in those patients with Graves’ disease, symptoms related to thyroid ophthalmopathy may dominate the presentation.

Who should be screened? Clinicians should screen all patients with hypertension suspected of also having hyperthyroidism based on the clinical presentation.

Case-detection tests. Case-detection testing includes measuring blood concentrations of thyroid-stimulating hormone and free thyroxine. In addition, these patients should undergo a comprehensive history and physical examination that includes measuring pulse rate, BP, respiratory rate, body weight, and thyroid gland palpation. After establishing clinical and biochemical

hyperthyroidism, clinicians often also perform imaging tests and assess thyroid autoantibodies; both these tests can be useful in identifying the underlying cause of hyperthyroidism. Initial management in patients with hypertension and hyperthyroidism includes using a β -adrenergic blocker to treat hypertension, tachycardia, and tremor. The definitive treatment of hyperthyroidism is cause specific (115).

Hypothyroidism

Prevalence. The prevalence of subclinical hypothyroidism ranges between 4.3% and 8.5%, and the prevalence of overt hypothyroidism ranges between 0.3% and 0.4% (110, 116). The frequency of hypertension (usually diastolic) is increased threefold in patients with hypothyroidism and may account for as many as 1% of cases of diastolic hypertension in the general population (117, 118). The mechanisms for BP elevation include increased systemic vascular resistance and extracellular volume expansion. There are no data on the prevalence of hypothyroidism in patients who present with hypertension.

Clinical presentation. The clinical presentation of hypothyroidism depends on the degree of thyroid hormone deficiency and the rapidity of the loss of the thyroid hormones thyroxine and triiodothyronine. Patients may be lethargic and exhibit slow cerebration, slow speech patterns, cold intolerance, constipation, and bradycardia. These patients typically have dry, brittle hair and edema of the face and eyelids (periorbital edema), which is associated with the subcutaneous accumulation of glycosaminoglycans. The tongue may be thickened and the voice deep and coarse. Patients with hypothyroidism generally have a slow pulse and diastolic hypertension. Cardiac output is decreased, and patients may experience dyspnea from exertion.

Who should be screened? Clinicians should screen all patients for hypothyroidism who have hypertension with any of the features listed in the previous paragraph.

Case-detection tests. Measuring serum thyrotropin and free thyroxine are the key case-detection tests. In primary hypothyroidism, the serum thyrotropin concentration is above the reference range and the blood concentration of free thyroxine is usually below the lower limit of the reference range. In central hypothyroidism due to hypothalamic or pituitary dysfunction, the serum thyrotropin concentration is inappropriately low for the low levels of free thyroxine. Treating thyroid hormone deficiency decreases BP in two-thirds of patients with hypertension and normalizes BP in the remainder. Synthetic levothyroxine is the treatment of choice for hypothyroidism (119).

C. Hypercalcemia and primary hyperparathyroidism

Prevalence

The most common cause of hypercalcemia is primary hyperparathyroidism. The prevalence of primary hyperparathyroidism is 0.20% in women and 0.09% in men, and the incidence increases with advancing age (120). Hypercalcemia is associated with an increased frequency of hypertension (121). The frequency of hypertension in patients with primary hyperparathyroidism varies from 10% to 60% (121). There are no data on the prevalence of primary hyperparathyroidism in patients who present with hypertension.

Clinical presentation

Most patients with primary hyperparathyroidism are asymptomatic. Others may present with symptoms related to chronic hypercalcemia, such as polyuria and polydipsia, constipation, osteoporosis, renal lithiasis, peptic ulcer disease, and hypertension. The mechanisms of hypertension are unclear because there is no direct correlation with the elevated parathyroid hormone or calcium levels. The hypertension associated with hyperparathyroidism can also occur as a complication of hypercalcemia-induced renal impairment.

Who should be screened?

Clinicians should screen all patients with hypertension and hypercalcemia for hyperparathyroidism.

Case-detection tests

Clinicians can confirm hyperparathyroidism as the cause of hypercalcemia by measuring serum parathyroid hormone and 24-hour urinary calcium excretion. Clinicians use urine calcium measurements to exclude familial hypocalciuric hypercalcemia (122). Clinicians should consider referral to a specialist when hypercalcemia is present. The treatment of hyperparathyroidism is surgery; hypertension may or may not remit after successful parathyroidectomy (123, 124).

D. Acromegaly

Prevalence

Acromegaly is a rare disorder, with a prevalence of 40 to 70 cases per 1 million people (125, 126).

Clinical presentation

Chronic growth hormone excess from a growth-hormone-producing pituitary tumor results in the clinical syndrome of acromegaly. The effects of chronic excess growth hormone include acral and soft tissue overgrowth, progressive dental malocclusion, degenerative arthritis related to chondral and synovial tissue overgrowth within joints, low-pitched sonorous voice,

excessive sweating and oily skin, perineural hypertrophy leading to nerve entrapment (*e.g.*, carpal tunnel syndrome), cardiac dysfunction, and hypertension (127). Hypertension occurs in 20% to 40% of the patients and is associated with sodium retention and extracellular volume expansion (128, 129). There are no data on the prevalence of acromegaly in patients who present with hypertension.

Who should be screened?

Patients with hypertension who should be screened for acromegaly include patients with incidentally discovered pituitary tumors and patients with typical clinical manifestations of acromegaly, especially those with acral and facial features or who have several associated conditions (*e.g.*, sleep apnea syndrome, type 2 diabetes mellitus, debilitating arthritis, carpal tunnel syndrome, hyperhidrosis, and hypertension).

Case-detection tests

Measuring serum insulin-like growth factor 1 is the case-detection test of choice (130). Clinicians should consider referring patients to an endocrine subspecialist when insulin-like growth factor 1 is above the upper limit of the age- and sex-adjusted reference range. Pituitary surgery is the treatment of choice; if necessary, clinicians also prescribe medical therapy or irradiation or both (130). Hypertension associated with acromegaly is treated most effectively by reducing excess growth hormone (129). If a surgical cure is not possible, the hypertension usually responds well to diuretic therapy.

E. Obstructive sleep apnea

OSA is repeated partial (hypopnea) or complete (apnea) airway closure during sleep. It is characterized by sleep disruption, intermittent hypoxemia and hypercapnea, and changes in intrathoracic pressure. The physiological effects of OSA, including those on endocrine pathways, are multiple and complex. Experimental and clinical studies indicate that OSA induces multiple adverse effects, including endothelial dysfunction, inflammation, oxidative stress, vascular stiffness, RAAS stimulation, metabolic abnormalities, cardiac structure and function alterations, and sympathetic nervous system activation (131).

OSA is strongly associated with hypertension; this is particularly true with resistant or difficult-to-treat hypertension (132–140). Several, but not all, observational studies have reported a positive correlation between aldosterone levels and the severity of OSA in patients with resistant hypertension (141–146). Studies suggest that intensifying diuretic therapy (including mineralocorticoid receptor antagonists) can reduce the severity of OSA in patients with resistant hypertension (147–150).

Prevalence

Approximately 20% of adults have at least mild sleep apnea (apnea-hypopnea index, 5 to 14 events per hour), and an estimated one in 15 has moderate to severe OSA (apnea-hypopnea index, ≥ 15 events per hour) (134). OSA is two to three times more common in men than women (151, 152), although the prevalence of OSA seemingly increases in women after menopause (153). OSA is strongly associated with being overweight or obese, with a prevalence of 60% to 80% in obese individuals (154, 155). OSA is also associated with type 2 diabetes (156). The prevalence of OSA increases with age, being two to three times more common in older persons (>65 years) compared with younger individuals (30 to 64 years) and having an estimated prevalence of 65% in a community-based sampling of persons >65 years of age (157–159). OSA appears to be more common in African Americans than whites, particularly children and adolescents, and studies of adults suggest that African Americans tend to have more severe OSA than whites (160–163).

OSA and hypertension commonly coexist. Approximately 50% of individuals with OSA are hypertensive, and an estimated 30% to 40% of patients with hypertension have OSA (131, 136–138). OSA is particularly prevalent in persons with resistant hypertension, with a prevalence of 70% to 90% (140–142, 146). However, continuous positive airway pressure (CPAP) therapy in patients with OSA and resistant hypertension, although beneficial, has not resulted in dramatic improvements in BP. For example, an RCT that blindly assessed outcomes of 117 patients with moderate to severe OSA who were randomly assigned to 6-month CPAP treatment or no therapy reported no significant difference in any BP change between CPAP and control groups (164). In an open-label, multicenter RCT of parallel groups of patients with OSA and resistant hypertension, CPAP treatment of 12 weeks compared with control groups resulted in a decrease in 24-hour mean BP and diastolic BP and an improvement in the nocturnal BP pattern (165).

Clinical presentation

Individuals with OSA often complain of a history of snoring (particularly loud snoring that is bothersome to others), episodes of witnessed apneas or chaotic breathing, and/or daytime sleepiness that is often severe. Other complaints include fragmented sleep with frequent arousals, snore arousals with a choking or gasping sensation, nocturia, dry mouth and/or headaches in the morning, daytime fatigue, cognitive dysfunction including memory impairment and/or difficulty concentrating, and symptoms of gastroesophageal reflux disease. Associated disease processes that are increased in

the setting of OSA include hypertension, heart disease, atrial fibrillation, prior stroke, gastroesophageal reflux disease, and sexual dysfunction.

The physical examination can be unremarkable, but OSA is more likely in persons who are obese (body mass index >30 kg/m²), older (>65 years), have an enlarged neck circumference (men, >43 cm; women, >37 cm), a small upper airway, and/or retrognathia or micrognathia.

Who should be screened?

Clinicians should screen all hypertensive individuals for OSA based on a history of relevant signs and symptoms. Although the decision to pursue definitive testing must be individualized, a history of loud snoring, witnessed apnea, and/or complaints of otherwise unexplained daytime sleepiness likely warrants further testing. Although resistant hypertension by itself is not yet sufficient reason alone to order definitive testing for OSA, clinicians should have a low threshold for ordering such testing, given the extraordinarily high prevalence of OSA in persons with resistant hypertension.

Case-detection tests

Polysomnography. The definitive diagnosis of OSA requires evaluation by polysomnography. Historically, sleep technicians monitor patients in a sleep laboratory. This allows for continuous recording of multiple parameters, including sleep staging using an electroencephalogram; eye movements, using an electrooculogram; body position, using video; limb movements, using an electromyogram; respiration (*i.e.*, flow, effort, oxygen saturation); and snoring. Recently, however, there is growing use of home sleep monitoring systems for diagnosing OSA. Such home-based systems are less expensive and more convenient for the individual being tested, but generally measure fewer parameters (often forgoing electroencephalogram and electrooculogram assessments). If a clinician suspects OSA in a hypertensive person, the clinician should refer the patient to a certified sleep specialist for evaluation and appropriate polysomnographic evaluation, whether via monitored or home testing.

VII. Summary of Overall Approach to Considering and Detecting Endocrine Hypertension

Clinical context is important. For example, case detection for endocrine hypertension may not be clinically important in an older patient with multiple life-limiting comorbidities. However, screening for endocrine hypertension may be key to enhancing and prolonging life in most patients with hypertension, especially younger patients.

A. Case in point

A 36-year-old woman presents with new-onset hypertension with repeated measurements of a systolic BP of 150 to 170 mm Hg and a diastolic BP of 95 to 110 mm Hg. Except for occasional headaches and palpitations, she is asymptomatic. She has been healthy and active. She does not use tobacco or alcohol. She has gained 5 kg over the past 1 year, and she has a history of recurrent renal lithiasis. Her husband said that she snores at night, but he has not observed periods of apnea. On questioning, she said that she does not feel rested in the morning. When she was in her 30s, she was told that she had a goiter. Her first menstrual period was at age 11 years and she has three biologic children. She does not eat confectionary licorice. Her family history is positive for hypertension in her father, hyperthyroidism in her maternal grandmother, and hypothyroidism in her mother. There is no family history of endocrine disorders or adrenal tumors.

The degree of hypertension in this 36-year-old woman merits an investigation for secondary causes. Because of the history of headaches and palpitations, case-detection testing for pheochromocytoma is indicated. We should check serum calcium and phosphorus levels because of her history of renal lithiasis. The weight gain over the past year could be a clue to Cushing syndrome and we should exclude this diagnosis. Her history of snoring and not feeling rested in the morning should prompt testing for OSA. We should exclude thyroid dysfunction based on her history of a goiter and family history of autoimmune thyroid disease. Because there are usually no symptoms or signs of PA, we should exclude this diagnosis in all patients with marked hypertension. In addition, she is in the age group in which we should consider fibromuscular dysplasia of the renal arteries. Depending on the degree of hypertension and urgency, we can perform case-detection testing for the potential diagnoses listed here either sequentially or simultaneously.

Acknowledgments

We thank medical writer Eric Vohr for his contribution to this Scientific Statement.

Address all correspondence and requests for reprints to: William F. Young, Jr., MD, Mayo Clinic, 200 First St SW, Rochester, Minnesota 55905. E-mail: young.william@mayo.edu.

Disclaimer Statement: The Endocrine Society develops Scientific Statements to assist clinicians by providing guidance and recommendations for particular areas of practice. One should not consider this Scientific Statement inclusive of all proper approaches or methods, or exclusive of others. It cannot guarantee any specific outcome, nor does it establish a standard of care. It is not intended to dictate the treatment of a particular patient. Treatment decisions must be made based on the independent judgment of health care providers and each patient's individual circumstances. The Endocrine Society makes no warranty, express or implied, regarding this Scientific Statement and specifically excludes any warranties of merchantability and fitness for a particular

use or purpose. The Society shall not be liable for direct, indirect, special, incidental, or consequential damages related to the use of the information contained herein.

Disclosure Summary: W.F.Y. is a consultant for Nihon Medi-Physics; D.A.C. provides consulting services for Novartis, receives industry support from Medtronic, and receives grant funding from the National Institutes of Health and the American Heart Association; the other authors have nothing to disclose.

References

1. Yoon SS, Ostchega Y, Louis T. Recent trends in the prevalence of high blood pressure and its treatment and control, 1999–2008. *NCHS Data Brief*. 2010;(48):1–8.
2. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, Lackland DT, LeFevre ML, MacKenzie TD, Oggedegbe O, Smith SC, Jr, Svetkey LP, Taler SJ, Townsend RR, Wright JT, Jr, Narva AS, Ortiz E. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014;311(5):507–520.
3. Fields LE, Burt VL, Cutler JA, Hughes J, Rocella EJ, Sorlie P. The burden of adult hypertension in the United States 1999 to 2000: a rising tide. *Hypertension*. 2004;44(4):398–404.
4. Rudnick KV, Sackett DL, Hirst S, Holmes C. Hypertension in a family practice. *Can Med Assoc J*. 1977;117(5):492–497.
5. Omura M, Saito J, Yamaguchi K, Kakuta Y, Nishikawa T. Prospective study on the prevalence of secondary hypertension among hypertensive patients visiting a general outpatient clinic in Japan. *Hypertens Res*. 2004;27(3):193–202.
6. Gupta-Malhotra M, Banker A, Shete S, Hashmi SS, Tyson JE, Barratt MS, Hecht JT, Milewicz DM, Boerwinkle E. Essential hypertension vs. secondary hypertension among children. *Am J Hypertens*. 2015;28(1):73–80.
7. Camelli S, Bobrie G, Postel-Vinay N, Azizi M, Plouin PF, Amar L. Lb01.11: Prevalence of secondary hypertension in young hypertensive adults. *J Hypertens*. 2015;33(Suppl 1):e47.
8. O'Shea PM, Griffin TP, Fitzgibbon M. Hypertension: the role of biochemistry in the diagnosis and management. *Clin Chim Acta*. 2017;465:131–143.
9. Funder JW, Carey RM, Mantero F, Murad MH, Reincke M, Shibata H, Stowasser M, Young WF, Jr. The management of primary aldosteronism: case detection, diagnosis, and treatment: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2016;101(5):1889–1916.
10. Lenders JW, Eisenhofer G, Mannelli M, Pacak K. Pheochromocytoma. *Lancet*. 2005;366(9486):665–675.
11. DeLellis R, Lloyd R, Heitz P, Eng C. *Pathology and Genetics of Tumours of Endocrine Organs*. Lyon, France: IARC Press; 2004.
12. Mansmann G, Lau J, Balk E, Rothberg M, Miyachi Y, Bornstein SR. The clinically inapparent adrenal mass: update in diagnosis and management. *Endocr Rev*. 2004;25(2):309–340.
13. Lo CY, Lam KY, Wat MS, Lam KS. Adrenal pheochromocytoma remains a frequently overlooked diagnosis. *Am J Surg*. 2000;179(3):212–215.
14. Beard CM, Sheps SG, Kurland LT, Carney JA, Lie JT. Occurrence of pheochromocytoma in Rochester, Minnesota, 1950 through 1979. *Mayo Clin Proc*. 1983;58(12):802–804.
15. Kimura N, Miura Y, Nagatsu I, Nagura H. Catecholamine synthesizing enzymes in 70 cases of functioning and non-functioning pheochromocytoma and extra-adrenal paraganglioma. *Virchows Arch A Pathol Anat Histopathol*. 1992;421(1):25–32.
16. Eisenhofer G, Lenders JW, Goldstein DS, Mannelli M, Csako G, Walther MM, Brouwers FM, Pacak K. Pheochromocytoma catecholamine phenotypes and prediction of tumor size and location by use of plasma free metanephrines. *Clin Chem*. 2005;51(4):735–744.
17. Eisenhofer G, Pacak K, Huynh TT, Qin N, Bratslavsky G, Linehan WM, Mannelli M, Friberg P, Grebe SK, Timmers HJ, Bornstein SR, Lenders JW. Catecholamine metabolomic and secretory phenotypes in pheochromocytoma. *Endocr Relat Cancer*. 2010;18(1):97–111.
18. Eisenhofer G, Huynh TT, Elkahoul A, Morris JC, Bratslavsky G, Linehan WM, Zhuang Z, Balgley BM, Lee CS, Mannelli M, Lenders JW, Bornstein SR, Pacak K. Differential expression of the regulated catecholamine secretory pathway in different hereditary forms of pheochromocytoma. *Am J Physiol Endocrinol Metab*. 2008;295(5):E1223–E1233.
19. Bravo EL, Tagle R. Pheochromocytoma: state-of-the-art and future prospects. *Endocr Rev*. 2003;24(4):539–553.
20. Eisenhofer G, Rivers G, Rosas AL, Quezado Z, Manger WM, Pacak K. Adverse drug reactions in patients with pheochromocytoma: incidence, prevention and management. *Drug Saf*. 2007;30(11):1031–1062.
21. Barrett C, van Uum SH, Lenders JW. Risk of catecholaminergic crisis following glucocorticoid administration in patients with an adrenal mass: a literature review. *Clin Endocrinol (Oxf)*. 2015;83(5):622–628.
22. Stolk RF, Bakx C, Mulder J, Timmers HJ, Lenders JW. Is the excess cardiovascular morbidity in pheochromocytoma related to blood pressure or to catecholamines? *J Clin Endocrinol Metab*. 2013;98(3):1100–1106.
23. Prejbisz A, Lenders JW, Eisenhofer G, Januszewicz A. Cardiovascular manifestations of pheochromocytoma. *J Hypertens*. 2011;29(11):2049–2060.
24. Young WF, Jr. Clinical practice. The incidentally discovered adrenal mass. *N Engl J Med*. 2007;356(6):601–610.
25. Lenders JW, Duh QY, Eisenhofer G, Gimenez-Roqueplo AP, Grebe SK, Murad MH, Naruse M, Pacak K, Young WF, Jr; Endocrine Society. Pheochromocytoma and paraganglioma: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2014;99(6):1915–1942.
26. Lenders JW, Pacak K, Walther MM, Linehan WM, Mannelli M, Friberg P, Keiser HR, Goldstein DS, Eisenhofer G. Biochemical diagnosis of pheochromocytoma: which test is best? *JAMA*. 2002;287(11):1427–1434.
27. Perry CG, Sawka AM, Singh R, Thabane L, Bajnarek J, Young WF, Jr. The diagnostic efficacy of urinary fractionated metanephrines measured by tandem mass spectrometry in detection of pheochromocytoma. *Clin Endocrinol (Oxf)*. 2007;66(5):703–708.
28. de Jong WH, Eisenhofer G, Post WJ, Muskiet FA, de Vries EG, Kema IP. Dietary influences on plasma and urinary metanephrines: implications for diagnosis of catecholamine-producing tumors. *J Clin Endocrinol Metab*. 2009;94(8):2841–2849.
29. Eisenhofer G, Lenders JW, Siegert G, Bornstein SR, Friberg P, Milosevic D, Mannelli M, Linehan WM, Adams K, Timmers HJ, Pacak K. Plasma methoxytyramine: a novel biomarker of metastatic pheochromocytoma and paraganglioma in relation to established risk factors of tumor size, location and SDHB mutation status. *Eur J Cancer*. 2012;48(11):1739–1749.
30. Willemsen JJ, Sweep CG, Lenders JW, Ross HA. Stability of plasma free metanephrines during collection and storage as assessed by an optimized HPLC method with electrochemical detection. *Clin Chem*. 2003;49(11):1951–1953.
31. Willemsen JJ, Ross HA, Lenders JW, Sweep FC. Stability of urinary fractionated metanephrines and catecholamines during collection, shipment, and storage of samples. *Clin Chem*. 2007;53(2):268–272.
32. Sawka AM, Thabane L, Gafni A, Levine M, Young WF, Jr. Measurement of fractionated plasma metanephrines for exclusion of pheochromocytoma: can specificity be improved by adjustment for age? *BMC Endocr Disord*. 2005;5(1):1.
33. Eisenhofer G, Lattke P, Herberg M, Siegert G, Qin N, Därr R, Hoyer J, Villringer A, Prejbisz A, Januszewicz A, Remaley A,

- Martucci V, Pacak K, Ross HA, Sweep FC, Lenders JW. Reference intervals for plasma free metanephrines with an age adjustment for normetanephrine for optimized laboratory testing of pheochromocytoma. *Ann Clin Biochem.* 2013;50(Pt 1):62–69.
34. Amar L, Eisenhofer G. Diagnosing pheochromocytoma/paraganglioma in a patient presenting with critical illness: biochemistry versus imaging. *Clin Endocrinol (Oxf).* 2015;83(3):298–302.
 35. Algeciras-Schimmich A, Preissner CM, Young WF, Jr, Singh RJ, Grebe SK. Plasma chromogranin A or urine fractionated metanephrines follow-up testing improves the diagnostic accuracy of plasma fractionated metanephrines for pheochromocytoma. *J Clin Endocrinol Metab.* 2008;93(1):91–95.
 36. Eisenhofer G, Goldstein DS, Walther MM, Friberg P, Lenders JW, Keiser HR, Pacak K. Biochemical diagnosis of pheochromocytoma: how to distinguish true- from false-positive test results. *J Clin Endocrinol Metab.* 2003;88(6):2656–2666.
 37. Conn JW. Presidential address. I. Painting background. II. Primary aldosteronism, a new clinical syndrome. *J Lab Clin Med.* 1955;45(1):3–17.
 38. Conn JW, Knopf RF, Nesbit RM. Clinical characteristics of primary aldosteronism from an analysis of 145 cases. *Am J Surg.* 1964;107:159–172.
 39. Mulatero P, Stowasser M, Loh KC, Fardella CE, Gordon RD, Mosso L, Gomez-Sanchez CE, Veglio F, Young WF, Jr. Increased diagnosis of primary aldosteronism, including surgically correctable forms, in centers from five continents. *J Clin Endocrinol Metab.* 2004;89(3):1045–1050.
 40. Rossi GP, Bernini G, Caliumi C, Desideri G, Fabris B, Ferri C, Ganzaroli C, Giacchetti G, Letizia C, Maccario M, Mallamaci F, Mannelli M, Mattarello MJ, Moretti A, Palumbo G, Parenti G, Porteri E, Semplicini A, Rizzoni D, Rossi E, Boscaro M, Pessina AC, Mantero F; PAPY Study Investigators. A prospective study of the prevalence of primary aldosteronism in 1,125 hypertensive patients. *J Am Coll Cardiol.* 2006;48(11):2293–2300.
 41. Calhoun DA, Nishizaka MK, Zaman MA, Thakkar RB, Weissmann P. Hyperaldosteronism among black and white subjects with resistant hypertension. *Hypertension.* 2002;40(6):892–896.
 42. Eide IK, Torjesen PA, Drolsum A, Babovic A, Lilledahl NP. Low-renin status in therapy-resistant hypertension: a clue to efficient treatment. *J Hypertens.* 2004;22(11):2217–2226.
 43. Young WF, Jr, Klee GG. Primary aldosteronism. Diagnostic evaluation. *Endocrinol Metab Clin North Am.* 1988;17(2):367–395.
 44. Gordon RD, Stowasser M, Klemm SA, Tunny TJ. Primary aldosteronism and other forms of mineralocorticoid hypertension. In: Swales J, ed. *Textbook of Hypertension*. London: Blackwell Scientific Publications; 1994:865–892.
 45. Murphy BF, Whitworth JA, Kincaid-Smith P. Malignant hypertension due to an aldosterone producing adrenal adenoma. *Clin Exp Hypertens A.* 1985;7(7):939–950.
 46. Vetter H, Siebenschein R, Studer A, Witassek F, Furrer J, Glänzer K, Siegenthaler W, Vetter W. Primary aldosteronism: inability to differentiate unilateral from bilateral adrenal lesions by various routine clinical and laboratory data and by peripheral plasma aldosterone. *Acta Endocrinol (Copenh).* 1978;89(4):710–725.
 47. Rich GM, Ulick S, Cook S, Wang JZ, Lifton RP, Dluhy RG. Glucocorticoid-remediable aldosteronism in a large kindred: clinical spectrum and diagnosis using a characteristic biochemical phenotype. *Ann Intern Med.* 1992;116(10):813–820.
 48. Stowasser M, Gartside MG, Gordon RD. A PCR-based method of screening individuals of all ages, from neonates to the elderly, for familial hyperaldosteronism type I. *Aust N Z J Med.* 1997;27(6):685–690.
 49. Gordon RD, Stowasser M. Familial forms broaden the horizons for primary aldosteronism. *Trends Endocrinol Metab.* 1998;9(6):220–227.
 50. Stowasser M, Gordon RD. Primary aldosteronism: from genesis to genetics. *Trends Endocrinol Metab.* 2003;14(7):310–317.
 51. Stowasser M, Huggard PR, Rossetti TR, Bachmann AW, Gordon RD. Biochemical evidence of aldosterone overproduction and abnormal regulation in normotensive individuals with familial hyperaldosteronism type I. *J Clin Endocrinol Metab.* 1999;84(11):4031–4036.
 52. Gordon RD, Stowasser M, Rutherford JC. Primary aldosteronism: are we diagnosing and operating on too few patients? *World J Surg.* 2001;25(7):941–947.
 53. Ono Y, Iwakura Y, Morimoto R, Kudo M, Igarashi Y, Nezu M, Tezuka Y, Ogawa H, Ito S, Stash F. Os 35-01 the prevalence of sleep apnea syndrome in primary aldosteronism. *J Hypertens.* 2016;34(Suppl 1) - ISH 2016 Abstract Book:e399.
 54. Campino C, Trejo P, Carvajal CA, Vecchiola A, Valdivia C, Fuentes CA, Delgado JF, Lagos CF, Aglony M, Carrasco C, Martinez-Aguayo A, García H, Loureiro C, Fardella CE. Pregnancy normalized familial hyperaldosteronism type I: a novel role for progesterone? *J Hum Hypertens.* 2015;29(2):138–139.
 55. Ronconi V, Turchi F, Zennaro MC, Boscaro M, Giacchetti G. Progesterone increase counteracts aldosterone action in a pregnant woman with primary aldosteronism. *Clin Endocrinol (Oxf).* 2011;74(2):278–279.
 56. Albiger NM, Sartorato P, Mariniello B, Iacobone M, Finco I, Fassina A, Mantero F. A case of primary aldosteronism in pregnancy: do LH and GNRH receptors have a potential role in regulating aldosterone secretion? *Eur J Endocrinol.* 2011;164(3):405–412.
 57. Teo AE, Garg S, Shaikh LH, Zhou J, Karet Frankl FE, Gurnell M, Happerfield L, Marker A, Bienz M, Azizan EA, Brown MJ. Pregnancy, primary aldosteronism, and adrenal CTNNB1 mutations. *N Engl J Med.* 2015;373(15):1429–1436.
 58. Celen O, O'Brien MJ, Melby JC, Beazley RM. Factors influencing outcome of surgery for primary aldosteronism. *Arch Surg.* 1996;131(6):646–650.
 59. Rutherford JC, Taylor WL, Stowasser M, Gordon RD. Success of surgery for primary aldosteronism judged by residual autonomous aldosterone production. *World J Surg.* 1998;22(12):1243–1245.
 60. Stowasser M, Gordon RD. Primary aldosteronism—careful investigation is essential and rewarding. *Mol Cell Endocrinol.* 2004;217(1-2):33–39.
 61. Karagiannis A, Tziomalos K, Papageorgiou A, Kakafika AI, Pagourelas ED, Anagnostis P, Athyros VG, Mikhailidis DP. Spironolactone versus eplerenone for the treatment of idiopathic hyperaldosteronism. *Expert Opin Pharmacother.* 2008;9(4):509–515.
 62. Lim PO, Farquharson CA, Shiels P, Jung RT, Struthers AD, MacDonald TM. Adverse cardiac effects of salt with fludrocortisone in hypertension. *Hypertension.* 2001;37(3):856–861.
 63. Fallo F, Sonino N. Metabolic syndrome and primary aldosteronism: time for reappraisal? *J Hum Hypertens.* 2010;24(10):623–624.
 64. Stowasser M. New perspectives on the role of aldosterone excess in cardiovascular disease. *Clin Exp Pharmacol Physiol.* 2001;28(10):783–791.
 65. Weber KT, Brilla CG. Pathological hypertrophy and cardiac interstitium. Fibrosis and renin-angiotensin-aldosterone system. *Circulation.* 1991;83(6):1849–1865.
 66. Young MJ, Funder JW. Mineralocorticoid receptors and pathophysiological roles for aldosterone in the cardiovascular system. *J Hypertens.* 2002;20(8):1465–1468.
 67. Catena C, Colussi G, Nadalini E, Chiuch A, Baroselli S, Lapenna R, Sechi LA. Cardiovascular outcomes in patients with primary aldosteronism after treatment. *Arch Intern Med.* 2008;168(1):80–85.
 68. Milliez P, Gierd X, Plouin PF, Blacher J, Safar ME, Mourad JJ. Evidence for an increased rate of cardiovascular events in patients with primary aldosteronism. *J Am Coll Cardiol.* 2005;45(8):1243–1248.
 69. Reincke M, Fischer E, Gerum S, Merkle K, Schulz S, Pallauf A, Quinkler M, Hanslik G, Lang K, Hahner S, Allolio B, Meisinger C,

- Holle R, Beuschlein F, Bidlingmaier M, Endres S; German Conn's Registry-Else Kröner-Fresenius-Hyperaldosteronism Registry. Observational study mortality in treated primary aldosteronism: the German Conn's registry. *Hypertension*. 2012;60(3):618–624.
70. Gordon RD. Primary aldosteronism. *J Endocrinol Invest*. 1995;18(7):495–511.
71. Ahmed AH, Gordon RD, Taylor PJ, Ward G, Pimenta E, Stowasser M. Effect of contraceptives on aldosterone/renin ratio may vary according to the components of contraceptive, renin assay method, and possibly route of administration. *J Clin Endocrinol Metab*. 2011;96(6):1797–1804.
72. Ahmed AH, Gordon RD, Taylor PJ, Ward G, Pimenta E, Stowasser M. Are women more at risk of false-positive primary aldosteronism screening and unnecessary suppression testing than men? *J Clin Endocrinol Metab*. 2011;96(2):E340–E346.
73. McKenna TJ, Sequeira SJ, Heffernan A, Chambers J, Cunningham S. Diagnosis under random conditions of all disorders of the renin-angiotensin-aldosterone axis, including primary hyperaldosteronism. *J Clin Endocrinol Metab*. 1991;73(5):952–957.
74. Gordon RD, Geddes RA, Pawsey CG, O'Halloran MW. Hypertension and severe hyperkalaemia associated with suppression of renin and aldosterone and completely reversed by dietary sodium restriction. *Australas Ann Med*. 1970;19(4):287–294.
75. Stowasser M, Gordon RD. The aldosterone-renin ratio for screening for primary aldosteronism. *Endocrinologist*. 2004;14(5):267–276.
76. Brown MJ, Hopper RV. Calcium-channel blockade can mask the diagnosis of Conn's syndrome. *Postgrad Med J*. 1999;75(882):235–236.
77. Mulatero P, Rabbia F, Milan A, Paglieri C, Morello F, Chiandussi L, Veglio F. Drug effects on aldosterone/plasma renin activity ratio in primary aldosteronism. *Hypertension*. 2002;40(6):897–902.
78. Stowasser M, Taylor PJ, Pimenta E, Ahmed AH, Gordon RD. Laboratory investigation of primary aldosteronism. *Clin Biochem Rev*. 2010;31(2):39–56.
79. Ahmed AH, Gordon RD, Taylor P, Ward G, Pimenta E, Stowasser M. Effect of atenolol on aldosterone/renin ratio calculated by both plasma renin activity and direct renin concentration in healthy male volunteers. *J Clin Endocrinol Metab*. 2010;95(7):3201–3206.
80. Ahmed AH, Calvird M, Gordon RD, Taylor PJ, Ward G, Pimenta E, Young R, Stowasser M. Effects of two selective serotonin reuptake inhibitor antidepressants, sertraline and escitalopram, on aldosterone/renin ratio in normotensive depressed male patients. *J Clin Endocrinol Metab*. 2011;96(4):1039–1045.
81. Melcescu E, Phillips J, Moll G, Subauste JS, Koch CA. 11Beta-hydroxylase deficiency and other syndromes of mineralocorticoid excess as a rare cause of endocrine hypertension. *Horm Metab Res*. 2012;44(12):867–878.
82. Speiser PW, Azziz R, Baskin LS, Ghizzoni L, Hensle TW, Merke DP, Meyer-Bahlburg HF, Miller WL, Montori VM, Oberfield SE, Ritzen M, White PC; Endocrine Society. Congenital adrenal hyperplasia due to steroid 21-hydroxylase deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2010;95(9):4133–4160.
83. Krone N, Arlt W. Genetics of congenital adrenal hyperplasia. *Best Pract Res Clin Endocrinol Metab*. 2009;23(2):181–192.
84. White PC, Dupont J, New MI, Leiber E, Hochberg Z, Rösler A. A mutation in CYP11B1 (Arg-448—His) associated with steroid 11 beta-hydroxylase deficiency in Jews of Moroccan origin. *J Clin Invest*. 1991;87(5):1664–1667.
85. Merke DP, Bornstein SR. Congenital adrenal hyperplasia. *Lancet*. 2005;365(9477):2125–2136.
86. New MI, Geller DS, Fallo F, Wilson RC. Monogenic low renin hypertension. *Trends Endocrinol Metab*. 2005;16(3):92–97.
87. Kim YM, Kang M, Choi JH, Lee BH, Kim GH, Ohn JH, Kim SY, Park MS, Yoo HW. A review of the literature on common CYP17A1 mutations in adults with 17-hydroxylase/17,20-lyase deficiency, a case series of such mutations among Koreans and functional characteristics of a novel mutation. *Metabolism*. 2014;63(1):42–49.
88. Müssig K, Wehrmann M, Horger M, Maser-Gluth C, Häring HU, Overkamp D. Adrenocortical carcinoma producing 11-deoxycorticosterone: a rare cause of mineralocorticoid hypertension. *J Endocrinol Invest*. 2005;28(1):61–65.
89. Ishikawa SE, Saito T, Kaneko K, Okada K, Fukuda S, Kuzuya T. Hypermineralocorticoidism without elevation of plasma aldosterone: deoxycorticosterone-producing adrenal adenoma and hyperplasia. *Clin Endocrinol (Oxf)*. 1988;29(4):367–375.
90. Nicolaides NC, Roberts ML, Kino T, Braatvedt G, Hurt DE, Katsantoni E, Sertedaki A, Chrousos GP, Charmandari E. A novel point mutation of the human glucocorticoid receptor gene causes primary generalized glucocorticoid resistance through impaired interaction with the LXXLL motif of the p160 coactivators: dissociation of the transactivating and transrepressive activities. *J Clin Endocrinol Metab*. 2014;99(5):E902–E907.
91. Charmandari E, Kino T, Chrousos GP. Primary generalized familial and sporadic glucocorticoid resistance (Chrousos syndrome) and hypersensitivity. *Endocr Dev*. 2013;24:67–85.
92. Chapman K, Holmes M, Seckl J. 11β-hydroxysteroid dehydrogenases: intracellular gate-keepers of tissue glucocorticoid action. *Physiol Rev*. 2013;93(3):1139–1206.
93. Robles BJ, Sandoval AR, Dardon JD, Blas CA. Lethal liquorice lollies (liquorice abuse causing pseudohyperaldosteronism). *BMJ Case Rep*. 2013;2013.
94. Stewart PM, Krozowski ZS, Gupta A, Milford DV, Howie AJ, Sheppard MC, Whorwood CB. Hypertension in the syndrome of apparent mineralocorticoid excess due to mutation of the 11 beta-hydroxysteroid dehydrogenase type 2 gene. *Lancet*. 1996;347(8994):88–91.
95. Nieman LK, Biller BM, Findling JW, Newell-Price J, Savage MO, Stewart PM, Montori VM. The diagnosis of Cushing's syndrome: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2008;93(5):1526–1540.
96. Liddle GW. A familial renal disorder simulating primary aldosteronism but with negligible aldosterone secretion. *Trans Assoc Am Physicians*. 1963;76:199–213.
97. Rossier BC, Schild L. Epithelial sodium channel: Mendelian versus essential hypertension. *Hypertension*. 2008;52(4):595–600.
98. Herrmann SM, Saad A, Textor SC. Management of atherosclerotic renovascular disease after Cardiovascular Outcomes in Renal Atherosclerotic Lesions (CORAL). *Nephrol Dial Transplant*. 2015;30(3):366–375.
99. Lewis S, Kadian-Dodov D, Bansal A, Lookstein RA. Multimodality imaging of fibromuscular dysplasia. *Abdom Radiol (NY)*. 2016;41(10):2048–2060.
100. Zachrisson K, Herlitz H, Lonn L, Falkenberg M, Eklöf H. Duplex ultrasound for identifying renal artery stenosis: direct criteria re-evaluated. *Acta Radiol*. 2017;58(2):176–182.
101. Demirpolat G, Ozbek SS, Parildar M, Oran I, Memiş A. Reliability of intrarenal Doppler sonographic parameters of renal artery stenosis. *J Clin Ultrasound*. 2003;31(7):346–351.
102. Goupil R, Cowley D, Wolley M, Ahmed AH, Gordon RD, Stowasser M. The utility of renal venous renin studies in selection of patients with renal artery stenosis for angioplasty: a retrospective study. *J Hypertens*. 2015;33(9):1931–1938; discussion 1938.
103. Lindholm J, Juul S, Jørgensen JO, Astrup J, Bjerre P, Feldt-Rasmussen U, Hagen C, Jørgensen J, Kosteljanetz M, Kristensen L, Laurberg P, Schmidt K, Weeke J. Incidence and late prognosis of Cushing's syndrome: a population-based study. *J Clin Endocrinol Metab*. 2001;86(1):117–123.
104. Sacerdote A, Weiss K, Tran T, Rokeya Noor B, McFarlane SI. Hypertension in patients with Cushing's disease: pathophysiology, diagnosis, and management. *Curr Hypertens Rep*. 2005;7(3):212–218.
105. Baid S, Nieman LK. Glucocorticoid excess and hypertension. *Curr Hypertens Rep*. 2004;6(6):493–499.

106. Kendrick ML, Curlee K, Lloyd R, Farley DR, Grant CS, Thompson GB, Rowland C, Young WF, Jr, van Heerden JA. Aldosterone-secreting adrenocortical carcinomas are associated with unique operative risks and outcomes. *Surgery*. 2002;132(6):1008–1011, discussion 1012.
107. Tunbridge WM, Evered DC, Hall R, Appleton D, Brewis M, Clark F, Evans JG, Young E, Bird T, Smith PA. The spectrum of thyroid disease in a community: the Whickham survey. *Clin Endocrinol (Oxf)*. 1977;7(6):481–493.
108. Vanderpump MP, Tunbridge WM, French JM, Appleton D, Bates D, Clark F, Grimley Evans J, Hasan DM, Rodgers H, Tunbridge F. The incidence of thyroid disorders in the community: a twenty-year follow-up of the Whickham Survey. *Clin Endocrinol (Oxf)*. 1995;43(1):55–68.
109. Knudsen N, Jorgensen T, Rasmussen S, Christiansen E, Perrild H. The prevalence of thyroid dysfunction in a population with borderline iodine deficiency. *Clin Endocrinol (Oxf)*. 1999;51(3):361–367.
110. Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, Braverman LE. Serum TSH, T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab*. 2002;87(2):489–499.
111. Parle JV, Franklyn JA, Cross KW, Jones SC, Sheppard MC. Prevalence and follow-up of abnormal thyrotrophin (TSH) concentrations in the elderly in the United Kingdom. *Clin Endocrinol (Oxf)*. 1991;34(1):77–83.
112. Gussekloo J, van Exel E, de Craen AJ, Meinders AE, Frölich M, Westendorp RG. Thyroid status, disability and cognitive function, and survival in old age. *JAMA*. 2004;292(21):2591–2599.
113. Kanaya AM, Harris F, Volpato S, Pérez-Stable EJ, Harris T, Bauer DC. Association between thyroid dysfunction and total cholesterol level in an older biracial population: the health, aging and body composition study. *Arch Intern Med*. 2002;162(7):773–779.
114. Danzi S, Klein I. Thyroid hormone and blood pressure regulation. *Curr Hypertens Rep*. 2003;5(6):513–520.
115. Bahn RS, Burch HB, Cooper DS, Garber JR, Greenlee MC, Klein I, Laurberg P, McDougall IR, Montori VM, Rivkees SA, Ross DS, Sosa JA, Stan MN. Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American Thyroid Association and American Association of Clinical Endocrinologists. *Endocr Pract*. 2011;17(3):456–520.
116. Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study. *Arch Intern Med*. 2000;160(4):526–534.
117. Streeten DH, Anderson GH, Jr, Howland T, Chiang R, Smulyan H. Effects of thyroid function on blood pressure. Recognition of hypothyroid hypertension. *Hypertension*. 1988;11(1):78–83.
118. Jian WX, Jin J, Qin L, Fang WJ, Chen XR, Chen HB, Su Q, Xing HL. Relationship between thyroid-stimulating hormone and blood pressure in the middle-aged and elderly population. *Singapore Med J*. 2013;54(7):401–405.
119. Garber JR, Cobin RH, Gharib H, Hennessey JV, Klein I, Mechanick JI, Pessah-Pollack R, Singer PA, Woebler KA. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the American Association of Clinical Endocrinologists and the American Thyroid Association. *Thyroid*. 2012;22(12):1200–1235.
120. Yeh MW, Ituarte PH, Zhou HC, Nishimoto S, Liu IL, Harari A, Haigh PI, Adams AL. Incidence and prevalence of primary hyperparathyroidism in a racially mixed population. *J Clin Endocrinol Metab*. 2013;98(3):1122–1129.
121. Richards AM, Espiner EA, Nicholls MG, Ikram H, Hamilton EJ, Maslowski AH. Hormone, calcium and blood pressure relationships in primary hyperparathyroidism. *J Hypertens*. 1988;6(9):747–752.
122. Marx SJ. Letter to the editor: Distinguishing typical primary hyperparathyroidism from familial hypocalciuric hypercalcemia by using an index of urinary calcium. *J Clin Endocrinol Metab*. 2015;100(2):L29–30.
123. Heyliger A, Tangpricha V, Weber C, Sharma J. Parathyroidectomy decreases systolic and diastolic blood pressure in hypertensive patients with primary hyperparathyroidism. *Surgery*. 2009;146(6):1042–1047.
124. Rydberg E, Birgander M, Bondeson AG, Bondeson L, Wilenheimer R. Effect of successful parathyroidectomy on 24-hour ambulatory blood pressure in patients with primary hyperparathyroidism. *Int J Cardiol*. 2010;142(1):15–21.
125. Holdaway IM, Rajasoorya C. Epidemiology of acromegaly. *Pituitary*. 1999;2(1):29–41.
126. Kwon O, Song YD, Kim SY, Lee EJ; Rare Disease Study Group, Science and Research Committee, Korean Endocrine Society. Nationwide survey of acromegaly in South Korea. *Clin Endocrinol (Oxf)*. 2013;78(4):577–585.
127. Chanson P, Salenave S. Acromegaly. *Orphanet J Rare Dis*. 2008;3:17.
128. Terzolo M, Matrella C, Boccuzzi A, Luceri S, Borriero M, Reimondo G, Pia A, Rovero E, Paccotti P, Angeli A. Twenty-four hour profile of blood pressure in patients with acromegaly. Correlation with demographic, clinical and hormonal features. *J Endocrinol Invest*. 1999;22(1):48–54.
129. Berg C, Petersenn S, Lahner H, Herrmann BL, Buchfelder M, Droste M, Stalla GK, Strasburger CJ, Roggenbuck U, Lehmann N, Moebus S, Jöckel KH, Möhlenkamp S, Erbel R, Saller B, Mann K; Investigative Group of the Heinz Nixdorf Recall Study and the German Pegvisomant Observational Study Board and Investigators. Cardiovascular risk factors in patients with uncontrolled and long-term acromegaly: comparison with matched data from the general population and the effect of disease control. *J Clin Endocrinol Metab*. 2010;95(8):3648–3656.
130. Katznelson L, Laws ER, Jr, Melmed S, Molitch ME, Murad MH, Utz A, Wass JA; Endocrine Society. Acromegaly: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2014;99(11):3933–3951.
131. Somers VK, White DP, Amin R, Abraham WT, Costa F, Culebras A, Daniels S, Floras JS, Hunt CE, Olson LJ, Pickering TG, Russell R, Woo M, Young T; American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology; American Heart Association Stroke Council; American Heart Association Council on Cardiovascular Nursing; American College of Cardiology Foundation. Sleep apnea and cardiovascular disease: an American Heart Association/American College of Cardiology Foundation Scientific Statement from the American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council on Cardiovascular Nursing. In collaboration with the National Heart, Lung, and Blood Institute National Center on Sleep Disorders Research (National Institutes of Health). *Circulation*. 2008;118(10):1080–1111.
132. Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med*. 2000;342(19):1378–1384.
133. Nieto FJ, Young TB, Lind BK, Shahar E, Samet JM, Redline S, D'Agostino RB, Newman AB, Lebowitz MD, Pickering TG. Association of sleep-disordered breathing, sleep apnea, and hypertension in a large community-based study. Sleep Heart Health Study. *JAMA*. 2000;283(14):1829–1836.
134. Young T, Peppard P, Palta M, Hla KM, Finn L, Morgan B, Skatrud J. Population-based study of sleep-disordered breathing as a risk factor for hypertension. *Arch Intern Med*. 1997;157(15):1746–1752.
135. Cano-Pumarega I, Durán-Cantolla J, Aizpuru F, Miranda-Serrano E, Rubio R, Martínez-Null C, de Miguel J, Egea C, Cancelo L, Alvarez A, Fernández-Bolaños M, Barbé F. Obstructive sleep apnea and systemic hypertension: longitudinal study in the general

- population: the Vitoria Sleep Cohort. *Am J Respir Crit Care Med*. 2011;184(11):1299–1304.
136. Fletcher EC, DeBehnke RD, Lovoi MS, Gorin AB. Undiagnosed sleep apnea in patients with essential hypertension. *Ann Intern Med*. 1985;103(2):190–195.
137. Lavie P, Ben-Yosef R, Rubin AE. Prevalence of sleep apnea syndrome among patients with essential hypertension. *Am Heart J*. 1984;108(2):373–376.
138. Worsnop CJ, Naughton MT, Barter CE, Morgan TO, Anderson AI, Pierce RJ. The prevalence of obstructive sleep apnea in hypertensives. *Am J Respir Crit Care Med*. 1998;157(1):111–115.
139. Grote L, Hedner J, Peter JH. Sleep-related breathing disorder is an independent risk factor for uncontrolled hypertension. *J Hypertens*. 2000;18(6):679–685.
140. Logan AG, Perlikowski SM, Mente A, Tisler A, Tkacova R, Niroumand M, Leung RS, Bradley TD. High prevalence of unrecognized sleep apnoea in drug-resistant hypertension. *J Hypertens*. 2001;19(12):2271–2277.
141. Gonzaga CC, Gaddam KK, Ahmed MI, Pimenta E, Thomas SJ, Harding SM, Oparil S, Cofield SS, Calhoun DA. Severity of obstructive sleep apnea is related to aldosterone status in subjects with resistant hypertension. *J Clin Sleep Med*. 2010;6(4):363–368.
142. Pratt-Ubunama MN, Nishizaka MK, Boedefeld RL, Cofield SS, Harding SM, Calhoun DA. Plasma aldosterone is related to severity of obstructive sleep apnea in subjects with resistant hypertension. *Chest*. 2007;131(2):453–459.
143. Sim JJ, Yan EH, Liu IL, Rasgon SA, Kalantar-Zadeh K, Calhoun DA, Deroose SF. Positive relationship of sleep apnea to hyperaldosteronism in an ethnically diverse population. *J Hypertens*. 2011;29(8):1553–1559.
144. Florczak E, Prejbisz A, Szwencz-Pietrasz E, Sliwiński P, Bieleń P, Klisiewicz A, Michałowska I, Warchol E, Januszewicz M, Kała M, Witkowski A, Więcek A, Narkiewicz K, Somers VK, Januszewicz A. Clinical characteristics of patients with resistant hypertension: the RESIST-POL study. *J Hum Hypertens*. 2013;27(11):678–685.
145. Lloberes P, Sampol G, Espinel E, Segarra A, Ramon MA, Romero O, Ferrer R, Martínez-García MA, Tovar JL. A randomized controlled study of CPAP effect on plasma aldosterone concentration in patients with resistant hypertension and obstructive sleep apnea. *J Hypertens*. 2014;32(8):1650–1657, discussion 1657.
146. Muxfeldt ES, Margallo VS, Guimarães GM, Salles GF. Prevalence and associated factors of obstructive sleep apnea in patients with resistant hypertension. *Am J Hypertens*. 2014;27(8):1069–1078.
147. Gaddam K, Pimenta E, Thomas SJ, Cofield SS, Oparil S, Harding SM, Calhoun DA. Spironolactone reduces severity of obstructive sleep apnoea in patients with resistant hypertension: a preliminary report. *J Hum Hypertens*. 2010;24(8):532–537.
148. Kasai T, Bradley TD, Friedman O, Logan AG. Effect of intensified diuretic therapy on overnight rostral fluid shift and obstructive sleep apnoea in patients with uncontrolled hypertension. *J Hypertens*. 2014;32(3):673–680.
149. Yang L, Zhang H, Cai M, Zou Y, Jiang X, Song L, Liang E, Bian J, Wu H, Hui R. Effect of spironolactone on patients with resistant hypertension and obstructive sleep apnea. *Clin Exp Hypertens*. 2016;38(5):464–468.
150. Krasnińska B, Miazga A, Cofta S, Szczepaniak-Chichel L, Trafas T, Krasniński Z, Pawlaczyk-Gabriel K, Tykarski A. Effect of eplerenone on the severity of obstructive sleep apnea and arterial stiffness in patients with resistant arterial hypertension. *Pol Arch Med Wewn*. 2016;126(5):330–339.
151. Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med*. 1993;328(17):1230–1235.
152. Redline S, Kump K, Tishler PV, Browner I, Ferrette V. Gender differences in sleep disordered breathing in a community-based sample. *Am J Respir Crit Care Med*. 1994;149(3 Pt 1):722–726.
153. Young T, Finn L, Austin D, Peterson A. Menopausal status and sleep-disordered breathing in the Wisconsin Sleep Cohort Study. *Am J Respir Crit Care Med*. 2003;167(9):1181–1185.
154. Punjabi NM, Sorkin JD, Katzel LI, Goldberg AP, Schwartz AR, Smith PL. Sleep-disordered breathing and insulin resistance in middle-aged and overweight men. *Am J Respir Crit Care Med*. 2002;165(5):677–682.
155. O’Keeffe T, Patterson EJ. Evidence supporting routine polysomnography before bariatric surgery. *Obes Surg*. 2004;14(1):23–26.
156. Nagayoshi M, Punjabi NM, Selvin E, Pankow JS, Shahar E, Iso H, Folsom AR, Lutsey PL. Obstructive sleep apnea and incident type 2 diabetes. *Sleep Med*. 2016;25:156–161.
157. Bixler EO, Vgontzas AN, Ten Have T, Tyson K, Kales A. Effects of age on sleep apnea in men: I. Prevalence and severity. *Am J Respir Crit Care Med*. 1998;157(1):144–148.
158. Young T, Shahar E, Nieto FJ, Redline S, Newman AB, Gottlieb DJ, Walsleben JA, Finn L, Enright P, Samet JM; Sleep Heart Health Study Research Group. Predictors of sleep-disordered breathing in community-dwelling adults: the Sleep Heart Health Study. *Arch Intern Med*. 2002;162(8):893–900.
159. Ancoli-Israel S, Kripke DF, Klauber MR, Mason WJ, Fell R, Kaplan O. Sleep-disordered breathing in community-dwelling elderly. *Sleep*. 1991;14(6):486–495.
160. Rudnick EF, Walsh JS, Hampton MC, Mitchell RB. Prevalence and ethnicity of sleep-disordered breathing and obesity in children. *Otolaryngol Head Neck Surg*. 2007;137(6):878–882.
161. Redline S, Tishler PV, Schluchter M, Aylor J, Clark K, Graham G. Risk factors for sleep-disordered breathing in children. Associations with obesity, race, and respiratory problems. *Am J Respir Crit Care Med*. 1999;159(5 Pt 1):1527–1532.
162. Pranathigieswaran S, Badr MS, Severson R, Rowley JA. The influence of race on the severity of sleep disordered breathing. *J Clin Sleep Med*. 2013;9(4):303–309.
163. Ancoli-Israel S, Klauber MR, Stepnowsky C, Estline E, Chinn A, Fell R. Sleep-disordered breathing in African-American elderly. *Am J Respir Crit Care Med*. 1995;152(6 Pt 1):1946–1949.
164. Muxfeldt ES, Margallo V, Costa LM, Guimarães G, Cavalcante AH, Azevedo JC, de Souza F, Cardoso CR, Salles GF. Effects of continuous positive airway pressure treatment on clinic and ambulatory blood pressures in patients with obstructive sleep apnea and resistant hypertension: a randomized controlled trial. *Hypertension*. 2015;65(4):736–742.
165. Martínez-García MA, Capote F, Campos-Rodríguez F, Lloberes P, Díaz de Atauri MJ, Somoza M, Masa JF, González M, Sacristán L, Barbé F, Durán-Cantolla J, Aizpuru F, Mañas E, Barreiro B, Mosteiro M, Cebrián JJ, de la Peña M, García-Río F, Maimó A, Zapater J, Hernández C, Grau SanMarti N, Montserrat JM; Spanish Sleep Network. Effect of CPAP on blood pressure in patients with obstructive sleep apnea and resistant hypertension: the HIPARCO randomized clinical trial. *JAMA*. 2013;310(22):2407–2415.