

HYPOPARATHYROIDISM AND PSEUDOHYPOPARATHYROIDISM

Geoffrey N. Hendy, Ph.D., Professor, Departments of Medicine, Physiology and Human Genetics, McGill University, Montreal, Quebec, Canada.

David E.C. Cole, M.D., Ph.D., Professor, Departments of Laboratory Medicine and Pathology, Medicine and Genetics, University of Toronto, Toronto, Ontario, Canada.

Murat Bastepe, M.D., Ph.D., Associate Professor, Harvard Medical School, Endocrine Unit, Massachusetts General Hospital, Boston MA. USA

Updated 6 February 2017

ABSTRACT

In primary hypoparathyroidism with hypocalcemia and hyperphosphatemia, deficient parathyroid hormone (PTH) secretion most commonly occurs from surgical excision of, or damage to, the parathyroid glands. The term idiopathic hypoparathyroidism describes isolated cases when a cause is not obvious and there is no family history. However, hypoparathyroidism is also a feature common to a variety of hereditary syndromes that may present *de novo*. Familial isolated hypoparathyroidism may show autosomal dominant, autosomal recessive, or X-linked inheritance. Genes involved include *PTH*, *SOX3*, *CASR*, *GNA11* and *GCM2*. Parathyroid hypoplasia is a frequent feature of 22q11.2 deletion syndrome with involvement of the *TBX1* gene. The Hypoparathyroidism, Nerve Deafness, and Renal Dysplasia syndrome is due to haploinsufficiency of the *GATA3* gene. Antibodies against parathyroid tissue are found in isolated hypoparathyroidism or combined with other endocrine deficiencies. Antibodies against the *CASR* occur in type 1 autoimmune polyglandular syndrome, due to mutations of the *AIRE* gene, or in acquired hypoparathyroidism. Disorders characterized by end-organ resistance to PTH are described collectively by the term pseudohypoparathyroidism (PHP) and PHP 1a and PHP 1b are caused by maternally-inherited changes at the complex imprinted *GNAS* locus at 20q13.32 that encodes the Gs α protein. Deleterious mutations of the *PTHR1* gene show resistance to PTH and PTHrP and present as Blomstrand lethal chondrodysplasia, Eiken syndrome, endochondromatosis, and primary failure of tooth eruption. Calcium and vitamin D are the standard therapy for management of hypoparathyroidism with hormone replacement [recombinant human PTH(1-84)] therapy recently becoming an option. Calcilytics, PTH analogs and orally-active small molecule PTHR1 agonists may, in the future, join the treatment armamentarium. For complete coverage of this and related areas in Endocrinology, visit our free web-books, www.endotext.org and www.thyroidmanager.org.

PRIMARY HYPOPARATHYROIDISM

Primary hypoparathyroidism is caused by a group of heterogeneous conditions in which hypocalcemia and hyperphosphatemia occur as a result of deficient parathyroid hormone (PTH) secretion (1). This most commonly results from surgical excision of, or damage to, the parathyroid glands. However, autoimmune disease is also a significant factor in acquired cases, and genetic forms of hypoparathyroidism due to decreased PTH secretion are not rare (Table 1).

Table 1. Forms of hypoparathyroidism having a genetic basis

1. Isolated	
1. Autosomal dominant	
1. <i>PTH</i> mutation	
2. <i>CASR</i> activating mutation (ADH1)	
1. Bartter Syndrome Type V	

3. *GCM2* mutation (dominant negative)
4. *GNA11* activating mutation (ADH2)
2. Autosomal recessive
 1. *PTH* mutation
 2. *GCM2* mutation
3. X-linked
2. Congenital multi-system syndromes*
 1. DiGeorge 1 (22q11) & 2 (10p)
 2. Barakat/HDR
 3. Kenny-Caffey 1 & 2 and Sanjad-Sakati
3. Metabolic disease
 1. Mitochondrial neuromyopathies
 2. Long-chain hydroxyacyl-CoA dehydrogenase (LCHAD) deficiency
 3. Heavy-metal storage disorders
4. Autoimmune disease
 1. Autoimmune polyendocrine syndrome type I (APS-1 / APECED)
5. Parathyroid resistance syndromes
 1. Pseudohypoparathyroidism
 2. Blomstrand chondrodysplasia and related PTH receptor defects
 3. Hypomagnesemia

* Clarke et al. (2) list other potential syndromic associations with hypoparathyroidism, including: CHARGE (**C**oloboma, **H**ear defect, **A**tresia choanae, **R**etarded growth and development, **G**enital hypoplasia, **E**ar anomalies/deafness), Dubowitz, lymphedema, nephropathy & nerve deafness

The signs and symptoms of hypoparathyroidism include evidence of latent or overt neuromuscular hyperexcitability due to hypocalcemia (Table 2). The effect may be aggravated by hyperkalemia or hypomagnesemia, but there is wide variation in the severity of symptoms. Patients may complain of circumoral numbness, paresthesias of the distal extremities or muscle cramping which can progress to carpopedal spasm or tetany. Laryngospasm or bronchospasm and seizures may also occur. Other less specific manifestations include fatigue, irritability, and personality disturbance. A comprehensive list of features associated with hypocalcemia can be found in the Endotext chapter, “Hypocalcemia: diagnosis and treatment” by Schafer & Shoback (3).

Severe hypocalcemia may be associated with a prolonged QT_c interval on electrocardiography, which reverses with treatment. More extensive cardiomyopathic changes may be seen. These include chest pain, elevated enzymes (CPK), left ventricular impairment, and T-wave inversion, suggestive of a myocardial infarction (4,5). Patients with chronic hypocalcemia may have calcification of the basal ganglia or more widespread intracranial calcification, detected by skull X-ray or CT scan. Also seen are extrapyramidal neurological symptoms (more often with intracranial calcification), subcapsular cataracts, band keratopathy, and abnormal dentition.

Table 2. Some clinical features of hypocalcemia

- | |
|--|
| <ul style="list-style-type: none"> • Neuromuscular irritability • Paresthesias • Laryngospasm |
|--|

- Bronchospasm
- Tetany
- Seizures
- Chvostek sign
- Trousseau sign
- Prolonged QT interval on ECG

Increased neuromuscular irritability may be demonstrated by eliciting a Chvostek or Trousseau sign. A positive Chvostek sign is a prolonged reflex contraction of the facial muscle in response to a digital tap on the cheek just anterior to the ear. As with other hyperreflexias, up to 20% of normal individuals may demonstrate a slight positive reaction. A positive Trousseau sign is carpopedal spasm induced by inflation of a blood pressure cuff covering the upper arm to 20 mm Hg above systolic blood pressure for three minutes. This response reflects the heightened irritability of nerves undergoing pressure ischemia.

In hypoparathyroidism, serum calcium concentrations are decreased and serum phosphate levels are increased. Serum PTH is low or undetectable. (The important exception is PTH resistance, discussed further below.) Usually, serum 1,25-dihydroxyvitamin D (1,25(OH)₂D) is low, but alkaline phosphatase activity is normal. Despite an increase in fractional excretion of calcium, intestinal calcium absorption and bone resorption are both suppressed. The renal filtered load of calcium is decreased, and the 24-h urinary calcium excretion is reduced; nephrogenous cyclic AMP excretion is low and renal tubular reabsorption of phosphate is elevated.

The terms idiopathic or isolated hypoparathyroidism have been traditionally used to describe isolated cases of glandular hypofunction when a cause is not obvious and there is no family history. However, hypoparathyroidism is a feature common to a variety of heritable syndromes that may present *de novo*. Hypoparathyroidism can occur because of a congenital hypoplasia/aplasia with or without other congenital anomalies such as dysmorphic facies, immunodeficiency, lymphedema, nephropathy, nerve deafness or cardiac malformation. Thus, in patients with hypoparathyroidism of uncertain onset, a careful examination of craniofacial features and assessment of endocrine, cardiac and renal systems should be performed to exclude a syndromic cause. Similarly, autoimmune hypoparathyroidism can occur as an isolated endocrine condition or with other glandular deficiencies in a pluriglandular autoimmune syndrome, requiring attention to multi-organ endocrine dysfunction.

A significant number of patients with idiopathic hypoparathyroidism and hypercalciuria, but no other anomalies may be found to have *de novo* activating mutations of the CASR gene. Because of the implications for treatment, CASR molecular screening of patients with this presentation is recommended (6,7).

Familial Isolated Hypoparathyroidism

Familial isolated hypoparathyroidism (FIH) may show autosomal dominant, autosomal recessive, or X-linked inheritance.

In a few instances of autosomal dominant disease, a mutation in the **PTH gene** (MIM# 168450 (8) - <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM>) has been found. In one family, a missense mutation (C18R) in the signal sequence of the preproPTH precursor has been identified (9) and the mutant shown to be defective *in vitro* in processing preproPTH to proPTH, although, as patients had one normal gene copy, the autosomal dominant mode of inheritance remained unexplained. Then, further studies in transfected cells

showed that the mutant was trapped in the endoplasmic reticulum (ER) promoting ER stress and apoptosis (10). In a family with autosomal recessive hypoparathyroidism, a different, homozygous, signal sequence mutation (S23P) segregates with affected status (11). This mutation may prevent proper cleavage of the signal peptide during processing of the nascent protein. In a girl with isolated hypoparathyroidism, a homozygous S23X signal sequence mutation was found predicting a truncated inactive PTH peptide (12). However, the circulating PTH level was not undetectable, suggesting some translational readthrough of the mutant preproPTH mRNA. A homozygous [Cys25]PTH(1-84) mutation that impairs PTHR1 activation was identified in an idiopathic hypoparathyroid family (13). Elevated circulating PTH levels were found in some (but not all) assays thus defining a novel form of hypoparathyroidism. In another family with autosomal recessive hypoparathyroidism, a donor splice site mutation at the exon 2/intron 2 junction of the PTH gene was identified (14). The mutation leads to exon skipping and loss of exon 2 containing the initiation codon and signal sequence of preproPTH mRNA. The SOX3 gene encodes a transcriptional factor likely involved in the embryonic development of the parathyroid gland (15). In two multigeneration families with X-linked recessive hypoparathyroidism exhibiting neonatal onset of hypocalcemia and parathyroid agenesis, the trait was mapped to a 906-kb region on distal Xq27 that contains three genes including SOX3 but no intragenic mutations were found (MIM# 307700). An interstitial deletion-insertion involving chromosomes 2p25.3 and Xq27.1, was found downstream of SOX3 and was speculated to exert a positional effect on SOX3 expression (16).

Gain-of-function mutations in the calcium-sensing receptor (CASR) gene (MIM#601199) have been identified in a number of families clinically diagnosed with autosomal dominant hypocalcemia type 1 (ADH1 – MIM#515361) (17,18). In the parathyroid gland, the activated CASR suppresses PTH secretion and in the kidney, it induces hypercalciuria that may contribute to the hypocalcemia. In many cases of ADH1, the family history is positive, but *de novo* mutations are quite common (19,20). Mosaicism for *de novo* mutation in an otherwise healthy parent has been described (21), and may explain some cases of apparently recessive disease. Most importantly, there are implications for counseling parents about risks of recurrence.

Almost all of the activating mutations are missense and appear almost equally divided between the amino-terminal third of the extracellular domain (ECD) and the transmembrane domain (TMD). Of special interest is the cluster of ECD mutations (A116T to C131W) that cause an increase in receptor sensitivity to extracellular calcium, suggesting that this region is critical for receptor activation. This cluster overlaps the two cysteine residues –cys-129 and cys-131– involved in the interface of the mature protein dimer (22). Further details can be found in the locus-specific database –<http://data.mch.mcgill.ca/casrdb/> (23) and (24).

Although Bartter syndrome subtype V is represented by only a handful of cases with heterozygous severe activating mutations in the CASR (MIM#601199), it provides additional insight into the functioning of the CaSR in the thick ascending limb (TAL) of the nephron (25-27). Bartter syndrome encompasses a heterogeneous group of electrolyte homeostasis disorders, the common features of which are hypokalemic alkalosis, hyperreninemia, and hyperaldosteronism. Bartter syndrome subtypes I–IV are autosomal recessive disorders due to inactivating mutations in the following ion transporters or channels active in the TAL: type I, the sodium potassium-chloride cotransporter (NKCC2); type II, the outwardly rectifying potassium channel (ROMK); type III, the voltage-gated chloride channel (CLC-Kb); type IV, Barttin, a chloride channel beta-subunit that is required for trafficking of CLC-Ka and CLC-Kb. Patients with the autosomal dominant Bartter syndrome subtype V have, in addition to the classic features of the syndrome, hypocalcemia, and may exhibit neuromuscular manifestations, seizures, and basal ganglia calcifications. NKCC2 and ROMK in the apical membrane (luminal side) of the TAL have been proposed to generate a transepithelial electrochemical gradient that drives passive paracellular transport of Na^+ , Mg^{2+} , and Ca^{2+} from the lumen to blood (28). The CASR is situated in the basolateral membrane (antiluminal side) and when activated increases 20-

hydroxyeicosatetraenoic acid and decreases cAMP concentrations, both of which would inhibit ROMK and NKCC2 activities (28,29). Thus, severe activating mutations of the CASR lead to the salt wasting of Bartter syndrome in addition to the hypercalciuric hypocalcemia of ADH1.

Heterozygous gain-of-function missense mutations of *GNA11* have been identified in ADH patients without detectable CASR activating mutations (30-33). The *GNA11* activating mutations increase the sensitivity of the parathyroid gland and renal tubule to extracellular calcium concentrations. Autosomal dominant hypocalcemia and hypoparathyroidism due to *CASR* and *GNA11* mutations are now designated as ADH type 1 (MIM#601198) and type 2 (MIM#615361) respectively. The human Gα11 protein (a Gq family member – MIM#139313) has 359 amino acids with an α-helical domain in the NH₂-terminal region, a GTPase domain in the COOH-terminal region, and three switch regions (SR1-3) in the middle portion that change conformation based on whether GTP or GDP is bound (34). The R80C, R181Q, S211W, F341L, and V304M mutations found in hypocalcemic individuals are predicted by 3D modeling to alter the normal Gα11 protein structure. Moreover, cells stably expressing the CASR and transfected with the mutants exhibit increased sensitivity to changes in extracellular calcium (30-33).

Inactivating mutations in the CASR regulator, the adaptor protein 2 sigma subunit encoded by the *AP2S1* gene, cause familial hypocalciuric hypercalcemia type 3 (35). The search for activating mutations in *AP2S1* in familial and sporadic isolated hypoparathyroid patients negative for *CASR* or *GNA11* mutations that would represent an additional genetic cause of ADH has thus far been negative (36,37).

Recessively inherited FIH may occur with mutations of the *glial cells missing-2 gene (GCM2; MIM#603716).* The *GCM2* gene localizes to chromosome 6p24.2 and encodes a transcription factor. It is expressed in the PTH-secreting cells of the developing parathyroid glands and is critical for their development in terrestrial vertebrates (38-40). A patient with neonatal hypoparathyroidism was found to be homozygous for a partial deletion acquired from both parents (41), and a pair of siblings with homozygous mutations has been reported (42). Additional studies have identified inactivating *GCM2* mutations in cases with autosomal recessive FIH (43,44). On the other hand, heterozygous mutations that cause dominant-negative *GCM2* mutants have also been identified in patients with autosomal dominant hypoparathyroidism (43,45,46). Additional recessive and dominant *GCM2* mutations have been noted in this gene that continues to be expressed in the adult parathyroid [see (47)]. Nevertheless, it appears that the prevalence of genetic defects affecting *GCM2* function is not high in isolated hypoparathyroidism, as a recent study investigating 20 unrelated cases with this disorder (10 familial and 10 sporadic) failed to identify any *GCM2* mutations segregating with the disease and/or leading to loss of function (48). Of further interest is that a genetic variant, Y282D that demonstrates significantly enhanced transcriptional activity relative to wild-type *GCM2* associates with hyperparathyroidism in some cohorts of the sporadic primary disorder (49). Most recently, novel heterozygous active *GCM2* variants that segregate with affected status in some kindreds with familial isolated **hyper**parathyroidism have been described (50). Thus, like *CASR* and *GNA11*, both gain-of-function and loss-of-function variants of *GCM2* may contribute to calcemic disorders.

Hypoparathyroidism with Syndromic Features

Hypoparathyroidism due to parathyroid hypoplasia is a frequent feature of **22q11.2 microdeletions**, the most common cause of **DiGeorge syndrome 1** (DS1; MIM#188400) (51,52). This syndrome complex arises from a failure of the third and fourth pharyngeal pouches to develop, leading to agenesis or congenital hypoplasia of the parathyroid glands, thymus, and the anterior heart field. Patients with DS1 may typically present with neonatal hypocalcemic seizures due to hypoparathyroidism, severe infections due to thymic hypoplasia, and

conotruncal heart defects (53). Because a microdeletion is involved, the identification of novel developmental genes in the 22q11 region has been keenly pursued. One of the genes is *TBX1*, encoding a DNA-binding transcription factor of the T-box family known to have important roles in vertebrate and invertebrate organogenesis and pattern formation (54,55). Mouse models with *Tbx1* haploinsufficiency established the essential contribution of this factor to conotruncal development (56), and placed it in developmental context during organogenesis (57,58). However, while the *Tbx1* null mutant mice had all the developmental anomalies of DS1 – thymic and parathyroid hypoplasia, abnormal facial structures and cleft palate, skeletal defects and cardiac outflow abnormalities – *Tbx1* haploinsufficiency in mice was associated with only defects of the fourth pharyngeal pouch responsible for the cardiac outflow abnormalities (59). cDNA microarray analyses of mice lacking *Tbx1* have identified *Gcm2* as one of the downregulated genes in the pharyngeal region, indicating that *Tbx1* is upstream of *Gcm2* (60). Furthermore, as *Tbx1* is regulated by sonic hedgehog (Shh) (61), a Shh-*Tbx1*-*Gcm2* parathyroid developmental pathway is indicated.

The basis for the phenotypic differences between DGS1 patients who are heterogeneous for *TBX1* loss and the *Tbx1*+/- mice is unclear but could reflect a species-specific gene dosage requirement together with roles of downstream genes regulated by *Tbx1*. Some patients may have late-onset DGS1 and develop symptomatic hypocalcemia in childhood or later with only subtle phenotypic abnormalities (62,63). Of note is that the age of diagnosis in rare families with DGS1 patients having *TBX1* inactivating (missense or frameshift) mutations ranged from 7 to 46 years in keeping with late-onset DGS1 (54).

The 22q11.2 deletion syndrome (22q11.2DS) encompasses a wider spectrum of clinical conditions that includes isolated congenital heart disease and velocardiofacial (VCF) syndrome (64). Associated craniofacial abnormalities include cleft palate, pharyngeal insufficiency and mildly dysmorphic facies. In the VCF syndrome, anatomical anomalies of the pharynx are prominent and hypernasal speech due to abnormal pharyngeal musculature with or without cleft palate is typical. In most patients, some degree of intellectual deficit is present and there is strong predisposition to psychiatric illness (schizophrenia or bipolar disorder) in adolescents and adults (65,66). Further information, both clinical and educational, can be found at web sites specifically devoted to this condition [see (67)].

The 22q11.2DS is due to one of the most common microdeletions (1 in 4000 live births), and it may go clinically unrecognized in its milder or variant forms. Most cases with hypoparathyroidism (~50% of cases) are the result of de novo deletion through meiotic non-allelic homologous recombination, and driven by a unique cluster of low copy repeats designated LCR22 A-H [see (67,68)]. Most commonly (~85% of cases), a deletion of ~3 Mb is found, encompassing proximal repeats A to D. Many of the others (~10% of cases) involve atypical nested deletions including those spanning LCR22 A to B. Thus LCR22 A to B, which includes the *TBX1* gene, is the primary site contributing to parathyroid dysgenesis. Detailed characterization and long-term follow-up for the hypoparathyroid component of this disorder is ongoing.

Although most cases of DiGeorge syndrome are sporadic, as mentioned above autosomal dominant inheritance is not unknown. In utero influences may be important determinants of the clinical picture, since there are instances of monozygotic twins with discordant phenotypes (69-71). Phenocopies occur with diabetic embryopathy, fetal alcohol syndrome, and retinoid embryopathy. In rare instances, it has been shown that a phenotypically normal parent can transmit a microdeletion to an offspring. Such parents have been found to carry a duplication of the 22q11 on the second chromosome, and the combination of duplication and deletion alleles in a parent generates a balanced state, termed “gene dosage compensation” (72-73).

Although the hypoparathyroidism affects about half of all carriers, it is usually not severe, and frequently treatment following neonatal hypocalcemia can be tapered or stopped in older children. However, the

hypoparathyroidism may also remain asymptomatic until adolescence or emerge at times of stress, such as corrective cardiac surgery or severe infection, suggesting that continued surveillance of parathyroid gland reserve is important. (74-76).

Traditionally, diagnosis of 22q11.2DS is established with specific cytogenetic studies -- usually with locus-specific fluorescence in-situ hybridization (FISH) testing. These tests will pick up many of the larger common deletions that involve regions of low-copy number repeats (LCRs). However, specific chromosomal array-based and MLPA analyses are now preferred, as they have been shown to have increased sensitivity for smaller deletions (67). Recently, the diagnostic power of next generation sequencing has been harnessed to identify almost all of the microdeletions underlying sporadic and inherited forms of the disorder (64). Non-invasive prenatal screening and pre-implantation genetic diagnosis are also clinically available (77). Because the clinical picture is so variable and the prevalence so high, testing for 22q11.2 microdeletion should be considered in the workup for any new hypoparathyroid case for which another cause is not found. Finally, distinct genetic defects can coexist with 22q11.2DS as exemplified by the finding of concurrence of this syndrome in an adolescent with longstanding **hypercalcemia** who had familial hypocalciuric hypercalcemia type 3 due to an *AP2S1* mutation (78).

Clinicians will also want to be aware that a small but significant minority (~10%) of patients will have associated autoimmune disease, driven in part, perhaps, by the thymus-based defect in T cell function (64,79). Among the more common (non-endocrine) conditions are arthritis, celiac disease and autoimmune hematologic disease, particularly idiopathic thrombocytopenic purpura. Autoimmune thyroid disease, with either hypo- or hyperparathyroid states, has been reported (79,80), and serum TSH assay should be measured regularly. It has been suggested that the later-onset hypoparathyroid disease may be partly autoimmune in origin, not developmental. A survey of 59 Norwegian patients showed discordance of adult onset disease with neonatal hypoparathyroidism, but a significant correlation with parathyroid autoantibodies and the presence of autoimmune disease (79).

The clinical features of DiGeorge syndrome including hypoparathyroidism also occur with other cytogenetic abnormalities, notably chromosome 10p haploinsufficiency (81,82). Deletions of two non-overlapping regions of chromosome 10p contribute to DiGeorge syndrome 2; DS2 at 10p13-14 (83), and the ***Barakat or HDR (Hypoparathyroidism, Nerve Deafness, and Renal Dysplasia) syndrome*** (MIM#146255) (84,85) at 10p14-10pter (66,87). The latter is due to haploinsufficiency of *GATA3* (MIM#131320), which encodes a dual zinc finger transcription factor (88) that is essential for normal embryonic development of the parathyroids, auditory system, and kidney. Since the original description, several additional *GATA3* loss-of-function mutations have been described in HDR patients [e.g., (89-92)]. Heterozygous *Gata3*-deficient mice develop parathyroid abnormalities as revealed by challenge with a diet low in calcium and vitamin D that are due to dysregulation of the parathyroid-specific transcription factor, *Gcm2*. *Gata3*^{-/-} embryos at E12.5 lack *Gcm2* expression and have gross defects in the fourth pharyngeal pouches, including absent parathyroid/thymus primordia (93). *GATA3* transactivates the *GCM2* promoter and with *GCM2* forms part of a transcriptional cascade essential for the differentiation and survival of parathyroid progenitor cells.

In another congenital disorder, ***Kenny-Caffey syndrome***, hypoparathyroidism is found variably associated with the typical picture of growth retardation, osteosclerosis, cortical thickening of the long bones, and delayed closure of the anterior fontanel (94-97). The original description of the syndrome was of the autosomal dominant form now identified as KCS-2 (MIM#127000) that is caused by heterozygous mutations in the *FAM111A* gene (98-100). The full functions of *FAM111A* and how mutations in it cause the disorder are unclear. *FAM111A* has some homology to peptidases, and is involved with chromatin structure during DNA replication (101). KS-2 is allelic to the lethal disorder, osteocraniostenosis (OCS, MIM#6023611).

Hypocalcemia due to hypoparathyroidism was found in some OCS patients who survived the perinatal period (97).

A recessively inherited form of Kenny-Caffey syndrome (KCS-1, MIM#244460) was noted to be similar to the recessive **Sanjad-Sakati syndrome** (MIM#241410) characterized by congenital hypoparathyroidism, seizures, growth and developmental retardation and characteristic dysmorphic features, including deep set eyes, depressed nasal bridge with beaked nose, long philtrum, thin upper lip, micrognathia and large, floppy ear lobes. Radiographs showed medullary stenosis reminiscent of Kenny-Caffey syndrome (97,102). Linkage studies localized the recessive KCS-1 and Sanjad-Sakati syndromes to 1q42-43, and causative mutations in the tubulin chaperone E, *TBCE*, gene were identified in what is now known as **Hypoparathyroidism, Retardation and Dysmorphism (HRD) syndrome** (97,103,104). This highlighted the role of TBCE that binds microtubules and proteasomes and protects against misfolded stress (105) in parathyroid development (106).

Hypoparathyroidism due to Metabolic Disease

Hypoparathyroidism is also a variable component of the neuromyopathies caused by mitochondrial gene defects (107). Among these are the **Kearns-Sayre syndrome** (ophthalmoplegia, retinal degeneration, and cardiac-conduction defects) (MIM#530000), the Pearson marrow pancreas syndrome (lactic acidosis, neutropenia, sideroblastic anemia, and pancreatic exocrine dysfunction) (108) (MIM#557000) and **mitochondrial encephalomyopathy** (MIM#540000). The molecular defects range from large deletions and duplications of the mitochondrial genomes in a large number of tissues (109,110) to single base-pair mutations in one of the transfer RNA genes found only in a restricted range of cell types (MIM#590050). The role of these mitochondrial mutations in the pathogenesis of hypoparathyroidism remains to be clarified. However, mutations in HADHB, that encodes the β -subunit of mitochondrial trifunctional protein, cause infantile onset hypoparathyroidism and peripheral polyneuropathy (111).

Long-chain hydroxyacyl-CoA dehydrogenase (LCHAD) deficiency (MIM#600890) is an inborn error of oxidative fatty acid metabolism that may be accompanied by hypoparathyroidism (112). Whether the parathyroid disease is directly related to the enzyme deficiency or secondary to the accompanying mitochondrial disease needs further study.

Parathyroid insufficiency and symptoms of hypocalcemia are occasionally seen in inherited metabolic disorders leading to excess storage of iron (**thalassemia, Diamond-Blackfan anemia, hemochromatosis**) or copper (**Wilson disease**) (113). In most instances, there is similar dysfunction in other endocrine glands, and the parathyroid disease is usually mild. Nonetheless, recognition of the hypoparathyroid state may help explain otherwise non-specific symptoms and aid in overall management of these multisystem diseases.

Autoimmune Hypoparathyroidism: Acquired and Inherited Disorders

Antibodies directed against parathyroid tissue have been detected in up to 38% of patients with isolated hypoparathyroid disease, and over 40% of patients having hypoparathyroidism combined with other endocrine deficiencies (114,115). Subsequently, a survey of a parathyroid expression library led to the identification of one protein selectively associated with the autoimmune process, the NACHT leucine-rich-repeat protein 5 (NALP5). Elevated antibody titres occur in half the patients with autoimmune hypoparathyroidism, with or without other autoimmune disease, but uncommonly in other conditions without hypoparathyroidism. (115,116).

Antibodies against the extracellular domain of the parathyroid CASR were originally reported in more than half of patients with either **type 1 autoimmune polyglandular syndrome (APS-1, also known as autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy or APECED)**, MIM# 240300, (117) or **acquired hypoparathyroidism** associated with autoimmune hypothyroidism (118). This finding was confirmed in a subsequent study of 51 cases of idiopathic hypoparathyroidism, but there was a 13% positive rate in controls (119). Other studies of APS-1 patients have also identified elevated CASR antibodies in some cases but at a lower frequency (120-122). Although some have suggested that CASR antibody assays are clinically indicated in acquired hypoparathyroidism (123), it remains to be seen whether the autoantibodies are of primary or secondary importance (115,124). There is now good evidence that autoantibodies can be functional activators of CASR and thereby could induce hypoparathyroidism. While presently there may not be a convenient clinical test for this, patient sera have been demonstrated to activate the CASR transfected into HEK cells *in vitro* (125). In some hypoparathyroid patients, both autoimmune parathyroid destruction and suppression by CASR activation may co-exist (126).

In APS-1, the most common associated manifestations are hypoparathyroidism with mucocutaneous candidiasis and Addison's disease. Additional features include pernicious anemia, chronic active hepatitis, alopecia, keratitis, gonadal failure, thyroid disease, pancreatic insufficiency and diabetes mellitus (117). The phenotype is highly variable and patients may not express all elements of the basic triad, leading to the suggestion that the criteria used for molecular screening be relaxed (126,127). The disease usually presents in infancy with chronic oral thrush, followed by hypoparathyroidism in the first decade, and then adrenocortical failure in the third. Interestingly, there is nearly 100% penetrance of hypoparathyroidism in females, but less than 60% in males, even though the adrenal hypofunction affects both sexes equally (121). Moreover, patients who develop the adrenal hypofunction first are less likely to be male and may never develop hypoparathyroidism. The responsible gene, called the autoimmune regulator (AIRE), maps to chromosome 21q22 and encodes a transcriptional regulator (128-130). In the absence of AIRE protein, tissue-specific self-antigens are not expressed in the thymus and multiorgan autoimmunity develops, because negative selection of the T cells bearing the autoantigens is disrupted (131). Many patients with APS-1 can be shown to have autosomal recessive inheritance of the AIRE defect. In families with autosomal recessive mutations of AIRE, obligate heterozygotes may also have common autoimmune disorders but APECED is not seen (132). A phenocopy leading to acquired APS-1 may occur when the AIRE gene is silenced by thymic neoplasia (133). APS-1 has been associated with more than 300 mutations of the AIRE gene, and updates can be found in the online mutation database (https://grenada.lumc.nl/LOVD2/mendelian_genes/home.php?select_db=AIRE).

PARATHYROID RESISTANCE SYNDROMES

Pseudohypoparathyroidism

Several clinical disorders characterized by end-organ resistance to PTH have been described collectively by the term pseudohypoparathyroidism (PHP) (134-137). They are associated with hypocalcemia, hyperphosphatemia, and increased circulating PTH. Target tissue unresponsiveness to the hormone manifests as a lack of increased phosphate excretion and, in some cases, cAMP excretion in response to PTH administration (138). The biochemical characteristics of the different forms of PHP are contrasted with those of hypoparathyroidism in Table 3.

Table 3. Biochemical characteristics of hypoparathyroidism and pseudohypoparathyroidism

Defects	Serum PO ₄	PTH	25(OH)D	1,25(OH) ₂ D	UcAMP*	UPO ₄ *	Multiple Endocrine Defects
Hypoparathyroidism	↑	↓	-	↓	-	-	Yes/No**

Pseudohypoparathyroidism							
Type 1a	↑	↑	-	↓	↓	↓	Yes
Type 1b	↑	↑	-	↓	↓	↓	No/Yes#
Type 1c	↑	↑	-	↓	↓	↓	Yes
Type 2	↑	↑	-	↓	-	↓	No

↑, increased;
 ↓, decreased;
 -, normal;
 *Response to PTH infusion
 **, depending upon the etiology.
 #, variable, mild defects of the thyroid axis due to TSH resistance may be seen.

Albright Hereditary Osteodystrophy

Fuller Albright first recognized that the likely cause of the hypoparathyroid state in PHP is a constitutive absence of target tissue responsiveness (139). In many patients, the end-organ resistance is accompanied by a specific pattern of physical findings, called **Albright hereditary osteodystrophy (AHO; MIM#300800)**. Typically, patients have short stature, round facies, brachydactyly, obesity, and ectopic soft tissue or dermal ossification(s) (osteoma cutis) (Figure 1). In the calvaria, this may manifest as hyperostosis frontalis interna (134). Intracranial calcification(s), cataracts and band keratopathy, subcutaneous calcifications, and dental hypoplasia are also common but are likely the consequences of longstanding hypoparathyroid hypocalcemia (Table 4, see below Figure 1). The brachydactyly may be asymmetric or not, and may involve one or both hands or feet, but the pattern is quite distinctive (140,141). The shortening tends to involve the first distal phalanx, with a thumbnail (or first toenail) that is wider than it is long. The fourth and fifth metacarpals (or metatarsals) are frequently shortened out of proportion to the others and the second metacarpal is often spared. Radiographic analysis of the hands (pattern profiling) may be helpful in assessment of the brachydactyly (Figure 1) (142).




Table 4. Incidence of signs and symptoms in PHP with AHO^a

Table 4. Incidence of signs and symptoms in PHP with AHO ^a		Percentage
Short stature		80
Obesity		50
Craniofacial		
Round face		92
^b Lenticular opacities		44
Strabismus		10
^b Dental hypoplasia		51
^b Basal ganglia calcification		50
Thickened calvaria		62
Mental deficit		75
Brachydactyly		
Brachymetacarpia		68
Brachymetatarsia		43
Brachyphalangia		50
Other connective tissue features		
Decreased bone density		15
Ectopic ossification		56
^b Subcutaneous calcification		55
^a Taken from Drezner and Neelon (1995)		
^b Features common to other forms of chronic hypoparathyroid hypocalcemia		

Although affected patients are generally short as adults, their bone age as children may be advanced and growth accelerated (142). Patients with AHO are probably predisposed to hypertension (143), conductive and sensorineural hearing loss (134,144), cord compression due to spinal anomalies (145), and movement disorders due to basal ganglia calcification (146). The features of AHO may be subtle in infancy or early childhood; in a few, there is little to see even in adulthood. The round facies, short neck, and low, flat nasal bridge are often accompanied by central obesity (147). A study showed that the obesity phenotype occurs primarily in those patients who also have multiple hormone resistance, i.e., PHP1a (see below), and according to data from mice, a hypothalamic mechanism, rather than hypothyroidism, is the primary underlying cause (148).

Patients with brachydactyly, mental retardation, and other features closely resembling AHO have been found to carry microdeletions of chromosome 2q37; **brachydactyly-mental retardation, BDMR**; MIM#600430 (149). Genes important for skeletal and neurological development lie within this region. Haploinsufficiency of *HDAC4* (MIM#605314), encoding a histone deacetylase that regulates gene expression during the development of many tissues including the bone, is responsible for the brachydactyly and the mental retardation in those patients (150). Isolated brachydactyly type E (BDE, MIM#113300) has been associated in sporadic cases with mutations in *HOX13* (MIM#168470) (151) and mutations in the *PTHLH* gene (MIM#168470) on 12p11.2 that encodes PTHrP have been implicated. In one family with autosomal BDE a *cis*-regulatory site downregulates *PTHLH* in translocation t(8;12)(q13;p11.2) and downregulates its targets *ADAMTS-7* and *ADAMTS-12* leading

to impaired chondrogenic differentiation (152). Affected individuals of one large family with BDE, short stature, and learning difficulties had an ~900 bp microdeletion encompassing *PTHLH* (153). Additional individuals with BDE and short stature from other different kindreds were found to have *PTHLH* missense, nonstop, and nonsense mutations (153). Different translocations affecting chromosome 12p have also been identified in two families with BDE, leading to increased abundance of a long noncoding RNA on chromosome 12q, which regulates the expression of *PTHLH* in cis and of the *SOX9* gene located on chromosome 17q in trans (154). BDE is associated with hypertension in some cases, in which the disease is inherited in an autosomal dominant manner (termed HTNB). Missense mutations in *PDE3A*, a gene encoding a cAMP/cGMP phosphodiesterase, have been recently found in several unrelated families with HTNB. These mutations cause increased cAMP hydrolytic activity and thus lead to diminished cAMP signaling (155). Some patients with AHO-like features have been described, who also showed platelet Gs hypofunction. Those patients were found to have *IGF2* hypermethylation and *SNURF* hypomethylation, as well as imprinting defects within *GNAS*, the gene encoding the stimulatory G protein alpha-subunit (G_{α} ; see below) (156).

PHP1a

PHP1a patients, characterized by AHO, PTH resistance, and evidence of target organ resistance to other hormones, are usually found to have a reduction in the activity of the G_{α} subunit, which is part of the membrane associated heterotrimeric G-protein complex - transducing signals between G-protein coupled receptors and adenylate cyclase (157-159). Adenylate cyclase catalyzes the synthesis of the second messenger cAMP, and therefore, PHP-1a patients tend to have a deficiency in cAMP generation, particularly in certain tissues. As explained above, this deficiency is clearly evident when measuring cAMP excretion in response to PTH administration.

The *GNAS* gene (MIM#168470) encoding the G_{α} protein maps to 20q13.2-13.3 and has at least 4 alternative transcriptional start sites (Figure 2) and an antisense transcript, *GNAS-AS1* (160). The three upstream exons and the preceding promoter regions are genetically imprinted, i.e., methylated in an allele specific manner. The promoter of the G_{α} transcript, which uses exon 1, is unmethylated. Unlike the other alternative *GNAS* products, G_{α} expression is biallelic except in a small set of tissues, where G_{α} is derived predominantly from the maternal allele (161-164). This tissue-specific monoallelic G_{α} expression affects penetrance of the PHP phenotype. The maternal transmission of the hormone resistance in PHP1a (165) can be explained by silencing of the paternal G_{α} allele, which would otherwise allow expression of 50% of G_{α} protein (166). Thus, full expression of a coding *GNAS* mutation, which occurs in maternally transmitted cases, leads to AHO plus hormone resistance (PHP1a), whereas a paternally transmitted mutation causes AHO alone (pseudopseudohypoparathyroidism; PPHP). Despite clinical evidence supporting imprinting in portions of the kidney tubule, it has been difficult to confirm this experimentally in humans (167). The imprinting of *GNAS* is complex and involves multiple differentially methylated regions (DMR) (160). Moreover, it is tissue-specific and may vary with developmental stage (168,169), although key imprinting of the 1A (also referred to as A/B) DMR is thought to be a primary event that occurs during gametogenesis and is maintained thereafter (170). Ablation of the G_{α} ortholog in mice (*Gnas*) has confirmed that maternal, but not paternal, transmission of the deleted allele results in PTH resistance. The homozygous deletion of *Gnas* is embryonic lethal (161). Comparison of G_{α} expression in mice with maternally vs paternally disrupted G_{α} expression also demonstrated that G_{α} expression is predominantly maternal in the renal cortex, but not in renal medulla (161). PTH resistance is delayed until after infancy in most PHP1a patients, and exon 1A imprinting may not be evident in the fetus (167). Indeed, a study using mice demonstrated that the silencing of the paternal G_{α} allele develops postnatally (171).




Figure 2. Simplified view of the *GNAS* region and its transcripts. The normal allele-specific methylation and expression patterns of the four alternate first exons of *GNAS* which splice onto exon 2 to produce transcripts encoding NESP55, XL α s, a transcript of unknown function (1A; also known as A/B), and Gs α (which uses exon 1). NESP55 and XL α s promoters are oppositely imprinted: NESP55 is expressed from the maternal allele and its promoter region is methylated on the paternal allele, whereas XL α s is expressed from the paternal allele and its promoter is methylated on the maternal allele. Gs α is paternally silenced in some tissues e.g., renal proximal tubule cells, indicated by the dashed arrow. NESP55 protein is unrelated to Gs α , and its entire coding region is located within its first exon. In contrast, XL α s and Gs α proteins have identical COOH-terminal domains (encoded by exons 2-13), while their unique NH₂-terminal domains are encoded within their respective first exons. Exon 1A does not have a translational start site, but is transcriptionally active. Loss of exon 1A imprinting (methylation) is associated with decreased Gs α expression in renal proximal tubules and some other hormone-responsive tissues, and is the typical cause of PHP1b. (figure from Liu et al., 2000, with permission).

A variety of inactivating mutations in the portion of the *GNAS* gene encoding Gs α have been identified in PHP1a patients (141,172-176). The spectrum includes missense mutations, point mutations impairing efficient and accurate splicing, and small insertion/deletion mutations. The 4bp deletion in exon 7 (Δ GACT 188/190) has been observed in multiple unrelated cases, suggesting that this may be a hot spot (136,176). Several other mutations have also been observed in more than one kindred, indicating that additional susceptibility regions may exist. The identification of *de novo* germline mosaicism (177) is consistent with the view that most sporadic cases harbor new mutations, but the separation of such sporadic cases from familial ones, in which there is suppression of phenotype due to imprinting, may be difficult without detailed molecular studies.

PHP1a cases have been described in which no mutations of the *GNAS* gene have been found by nucleotide sequence analysis of exons encoding Gs α . This may be because the mutation is in a regulatory region of the gene not yet examined, or it may be that a large deletion prevents amplification of the mutant allele for subsequent analyses. In cases without identified *GNAS* coding mutations, an assessment of Gs α bioactivity in erythrocytes is helpful in ruling out regulatory region mutations or large deletions. A 35 kb deletion spanning exons 1 through 5 has been identified by using comparative genome hybridization in a patient with PHP1a in whom coding mutations had been ruled out but a marked reduction of erythrocyte Gs α activity demonstrated (178,179). Typically, PHP1a is associated with multiple hormone resistance, including thyroid stimulating hormone (TSH) and gonadotropins, causing hypothyroidism and gonadal failure, respectively. Because the

hypothyroidism may express before hypocalcemia is observed (136), early surveillance of thyroid function is warranted. However, thyroid replacement from birth does not appear to prevent the mental deficit typical of PHP1a. In women, the hypogonadism is partial (180) so that oral contraceptives may help regulate the menstrual cycle. Estrogen can antagonize bone resorption leading to an exacerbation of hypocalcemia (181), but placental 1,25-dihydroxyvitamin D synthesis likely obviates this effect altogether in pregnancy so women are frequently normocalcemic at that time (182). Abnormalities of the somatotropin axis have also been reported, with documentation of subnormal growth hormone release following stimulation by L-arginine or growth hormone releasing hormone (183,184). The effect of growth hormone replacement has been investigated in a small group of pre-pubertal PHP1a patients (185). This study concluded that the treatment is potentially effective but has to be initiated as early as possible.

The tissue-specific silencing of the paternal $Gs\alpha$ allele also plays a key role in the development of the additional hormone resistance phenotypes, as monoallelic $Gs\alpha$ expression has been demonstrated in the thyroid, the ovaries, and the pituitary (162-164,186). Studies have revealed that obesity also develops primarily in patients who inherit the inactivating $Gs\alpha$ mutations from their mothers (187). $Gs\alpha$ is not imprinted in the white adipose tissue (188), but the investigations of mice in which $Gs\alpha$ is ablated conditionally in the brain showed that $Gs\alpha$ is also monoallelic in certain parts of the hypothalamus (189), thus explaining the imprinted mode of inheritance of the obesity phenotype. Likewise, it has been noted that cognitive impairment, a typical AHO feature, also develops primarily after maternal inheritance of the inactivating $Gs\alpha$ mutation (190).

PHP1b

PHP1b is typically not associated with AHO or a generalized reduction in $Gs\alpha$ expression (135). PHP1b patients show a defect in renal PTH signaling, but an apparently normal response to PTH in bone. Affected individuals are therefore functionally hypoparathyroid but show normal skeletal architecture and development. Due to unimpaired PTH responsiveness in bone, however, signs of hyperparathyroid bone disease (osteitis fibrosa cystica) are occasionally observed, complicating the picture (191,192). Biochemical abnormalities suggestive of thyroid stimulating hormone resistance are also seen in some patients (186), and abnormalities of renal uric acid handling have been documented (193,194). In fact, sometimes, PHP1b cases can present first with hypothyroidism (195,196). A recent study also demonstrated short stature and growth hormone deficiency in monozygotic twins with PHP1b (197). However, clinically significant hormone resistance is restricted to PTH in most cases. Because the hormone resistance is mostly limited to PTH, it was thought at one time that these findings could be explained by a defect in the type-1 parathyroid hormone receptor ($PTHr1$, MIM#168468), but sequencing in PHP1b patients found no mutations in protein-coding exons or gene promoter regions of the gene (198-200), and studies of PHP1b families show no linkage to $PTHr1$ (201,202).

Most cases of PHP1b are sporadic, but a familial form of PHP1b with an apparent autosomal dominant mode of inheritance also exists (AD-PHP1b). In four AD-PHP1b kindreds, linkage to chromosome 20q13.3 was established, the same region which includes the $GNAS$ locus (201). In these families, the pattern of transmission suggested paternal imprinting, and inheritance is therefore the same as for PHP1a. A further 13 PHP1b subjects were studied, some of whom had bone responsiveness to PTH (166). All lacked methylation of the alternate exon 1A, an epigenetic defect that is postulated to inhibit expression of the functional exon 1-containing $Gs\alpha$ transcript in renal tissues only (Figure 2). Thus, the loss of methylation of the maternal exon 1A allele leads to the silencing of the maternal as well as paternal $Gs\alpha$ allele, causing PTH resistance specifically in renal proximal tubule cells. Genetic analysis indicated that mutations in a regulatory region some distance from the $GNAS$ coding exons were likely to account for the unique imprinting defect(s) associated with PHP1b (203). A search for the mutation revealed the presence of a 3 kb microdeletion that segregated with the disease in 12 kindreds with AD-PHP1b and also occurred in 4 sporadic cases (204). The deletion, flanked by

direct repeats, removes 3 exons of the *STX16* gene, which encodes syntaxin-16. Two other private deletions within *STX16* have been identified in AD-PHP1b kindreds. One removes exons 2 to 4 (205) and the other exons 2 to 8 (206). In all these cases, maternal, but not paternal, inheritance of the *STX16* deletion led to PTH resistance. Because *STX16* is apparently not imprinted (205), loss of one copy of this gene is not predicted to underlie the PHP1b pathogenesis. Instead, these deletions presumably disrupt a cis-acting element that controls imprinting at *GNAS* exon 1A. Interestingly, a study identified a large deletion ablating *NESP55* without any overlap with *STX16* as the cause of PHP1b in a family in whom affected individuals showed isolated loss of 1A methylation (207). Thus, at least two distinct cis-acting regions appear to be necessary for the methylation imprints at the *GNAS* 1A region. In two other PHP1b kindreds, nearly identical deletions of the *NESP55* DMR including exons 3 and 4 of the antisense transcript segregated with the disease (208). In this instance, however, the 1A DMR was not the only region to lose the differential methylation required to allow maternal expression of *Gsα* in the kidney. Maternal methylation was also lost in the regions of the *XLαs* and *GNAS-AS1* promoters. Another kindred with these widespread epigenetic defects of *GNAS* has been described (209). The affected individuals in this kindred carried a maternally inherited deletion that removed antisense exons 3 and 4 with flanking intronic regions but not the *NESP55* exon. Additional genomic rearrangements in the chromosomal regions comprising *GNAS* have also been identified and proposed to underlie the *GNAS* methylation abnormalities in some AD-PHP-1b cases (210,211).

Sporadic PHP1b cases also show broad *GNAS* epigenetic defects that involve 1A. In some of these cases, paternal uniparental disomy of different chromosome 20 segments have been reported as the likely cause of PHP1b in several such cases (212-216). The cause of the epigenetic defects and PTH resistance, however, remains unknown for most cases of sporadic PHP1b. A few cases have deletions within *GNAS* that are *de novo* either in the patient or in the unaffected mother's paternal allele, as reported (217). Additional imprinting defects at other genomic loci have been identified in a few patients with PHP1b (218). Moreover, *GNAS* methylation defects have been identified in some cases with hypomethylation at multiple maternally methylated imprinted regions (219,220). In fact, some of those cases show both PTH resistance and the clinical features resulting from the methylation changes of the other loci, such as Beckwith-Wiedemann Syndrome (221).

A recent study revealed that, in addition to the exon 1A DMR, methylation at a new *GNAS* region close to the *GNAS-AS1* promoter (termed *GNAS-AS2*), is lost in patients who carry *STX16* deletions (222). Note that this region is also affected in those cases that display broad *GNAS* methylation changes. The effect of the loss of methylation at *GNAS-AS2* has yet to be determined at the level of gene expression, but this finding shows that the different subtypes of PHP-1b are more similar to one another at the molecular level than previously recognized. Based on one report, no clinical differences could be established according to the pattern of *GNAS* epigenetic defects, although serum PTH levels were significantly higher in females with broad *GNAS* methylation defects than females with isolated loss of 1A methylation (223). A recent study also found an intrauterine growth advantage for both AD-PHP-1b and sporadic PHP-1b cases, but the results indicate that the sporadic cases are not as markedly growth accelerated as AD-PHP-1b cases at birth (224).

In contradistinction to the classical understanding that AHO features are unique to PHP1a, some studies have identified patients with PTH resistance and AHO features who show *GNAS* epigenetic defects rather than *Gsα* coding mutations (225-227). Thus, there may be some overlap between the clinical and molecular features of PHP1a and PHP1b. It is possible that the AHO features observed in patients with *GNAS* epigenetic defects result from a genetic mechanism that is similar to the mechanism underlying the hormone resistance in PHP1a patients, i.e., due to monoallelic *Gsα* expression in additional tissues.

A PHP1b family with a novel Gs α mutation, deletion of isoleucine-382 in the carboxyl terminus (leading to uncoupling from the PTHR1 and isolated PTH resistance), shows transmission through 3 generations, consistent with paternal imprinting (228). However, such mutations within Gs α coding exons are rare (166).

PHP1c and PHP2. Patients with PHP1c have multiple hormone resistance but normal Gs α activity. The defect may be in other components of the receptor-adenylate cyclase system, such as the catalytic unit, but some PHP1c cases have been reported to carry Gs α coding mutations (229). These mutations render the Gs α protein unable to mediate cAMP generation in response to receptor activation but do not affect basal adenylate cyclase stimulating activity or the ability to be activated by non-hydrolyzable GTP analogs (230-232). Thus, some forms of PHP1c appear to be an allelic variant of PHP1a. Finally, patients with PHP2 have a normal urinary cAMP response to PTH but an impaired phosphaturic response (233). The defect could be in the cAMP-dependent protein kinase (PKA), one of its substrates or targets, or in a component of the PTH-PKC signaling pathway.

A study (234) has discovered a heterozygous mutation of the gene encoding the regulatory subunit of PKA (PRKAR1A) in three patients with **multiple hormone resistance and acrodysostosis**, a form of skeletal dysplasia that includes severe brachydactyly type-E and other skeletal findings that resemble AHO. This mutation, p.R368X, which leads to truncation of the COOH-terminal 14 residues, impairs cAMP binding to the regulatory subunit, thereby blocking the activation of PKA (234). In addition to acrodysostosis, patients carrying this mutation display evidence for target organ resistance to PTH, thyrotropin, growth hormone-releasing hormone, and gonadotropins, but these findings are accompanied by elevated basal plasma and urinary cAMP levels and with an apparently normal cAMP response to exogenous PTH administration. In certain other patients with acrodysostosis, but mostly without hormone resistance, it has been shown that the disease is caused by missense mutations in *PDE4D*, which encodes a cAMP phosphodiesterase (235-236). The type of acrodysostosis caused by *PRKAR1A* mutations has been termed **acrodysostosis-1** (MIM#101800), while the one caused by *PDE4D* mutations **acrodysostosis-2** (MIM#614613).

Other Phenotypes Associated with GNAS Mutations.

In contrast to the PHP phenotype associated with inactivating *GNAS* mutations, a different form of sporadic bone disease, (**polyostotic fibrous dysplasia**) results from *de novo* *GNAS* mutations that cause constitutive Gs α activity (237,238). A more severe form of this disease (**panostotic fibrous dysplasia**) with hyperphosphatasia and hyperphosphaturic rickets has also been described (239,240). Patients carrying these activating mutations are mosaic for mutant and wild-type cells, indicating that the mutation is acquired during postzygotic development. These mutations affect the arginine residue at position 201 (exon 8) and, rarely, the glutamine at 227 (exon 9), and inhibit the intrinsic GTP hydrolase activity of Gs α , thereby leading to constitutive activity. Such constitutively activating mutations of *GNAS* are also found in a variety of endocrine and non-endocrine tumors, such as growth hormone-secreting adenomas (241). A missense mutation in exon 13 (A366S) results in a Gs α protein that is unstable at 37°C, but constitutively active at lower temperatures (242,243). Affected patients have PHP due to PTH resistance and precocious puberty (testotoxicosis) due to hormone-independent constitutive activation of luteinizing hormone receptors at lower ambient temperatures in the testes. Another Gs α mutant carrying Ala-Val-Asp-Thr amino acid repeats in the guanine-binding domain has been described in a patient with neonatal diarrhea and PTH resistance (244). In this instance, the mutant protein is unstable and localized to the cytoplasm rather than plasma membrane, which explains the hormone resistance. On the other hand, this mutation increases the rate of GDP-GTP exchange and, thus, confers overactivity. The increased activity of Gs α seems to be evident during the neonatal period in the gut, where the mutant localizes to the plasma membrane, thus explaining the diarrhea phenotype. Undoubtedly, other patterns of hormone-receptor interaction due to a *GNAS* mutation await discovery.

Inactivating *GNAS* mutations have also been identified in patients with **congenital osteoma cutis** and **progressive osseous heteroplasia (POH)**, suggesting that these connective tissue conditions are another variant in the phenotypic spectrum of *GNAS*-related disease (245-248). No genotype-phenotype correlation has been revealed regarding these disorders, as the same mutation can be associated with either typical AHO features or severe ossifications that involve deep connective tissues and skeletal muscle (249). Nonetheless, patients with POH inherit the *GNAS* mutation from their fathers or acquire this mutation *de novo* on the paternal *GNAS* allele. This parent-of-origin specific inheritance of POH was established by analyzing 18 unrelated kindreds with this disorder (250). In a single, three generation, kindred, the inheritance of the mutation from males led to POH, while the inheritance of the same mutation from females led to typical AHO. It thus appears likely that alterations in the activity of a paternally expressed *GNAS* product, such as XLas, contribute to the pathogenesis of POH. However, POH-like features have also been seen in some patients with maternally inherited *GNAS* mutations (251). A recent study revealed that the distribution of POH lesions follow dermomyotomes and show a tendency for one-sidedness, suggesting that post-zygotic second hits may contribute to the development of these lesions on top of the inherited heterozygous mutations of *GNAS* (252).

Differential diagnosis and genetic counseling.

Patients with dysmorphic features resembling AHO may require careful endocrinologic work-up to confirm and delineate the form of PHP that is present. Similar studies of family members may also be warranted, since the biochemical and clinical features vary within families. If PHP1a with AHO is established, genetic counseling may aid in understanding the multisystemic nature of the disorder, particularly in relation to the patient's growth and development, and later-onset connective tissue complications. For either PHP1a or PHP1b, extensive counseling may be required to adequately explain the various implications of paternal imprinting for the parent-specific recurrence risks in offspring. Germline mosaicism has been reported (177), which is clearly important in assessing risks for recurrence in future sibs of a singleton family. Given the recently described complexities in the molecular, biochemical, and physical features of PHP1a and PHP1b, molecular testing is critical for achieving a clear diagnosis and validating the inheritance pattern in any given family.

THE PARATHYROID HORMONE RECEPTOR AND SKELETAL DYSPLASIAS

PTHr1 is a family B G protein-coupled receptor that signals through multiple different G proteins including Gsα (253). It responds to two ligands, PTH and the PTH-related peptide (PTHrP). It would thus be predicted that deleterious mutations might show resistance to PTH, as well as evidence for a defect of PTHrP action. Functional polymorphisms in the *PTHr1* are associated with adult height and bone mineral density (254), emphasizing the role that the receptor and its ligands play in endochondral bone formation. Inactivating or loss-of-function mutations in the *PTHr1* have been implicated in the molecular pathogenesis of **Blomstrand lethal chondrodysplasia (BLC; MIM#215045)**, and other skeletal dysplasias and dental abnormalities (255). The rare, recessive BLC is characterized by short-limbed dwarfism with craniofacial malformations, hydrops, hypoplastic lungs and aortic coarctation (256-260). The bones show accelerated endochondral ossification and deficient remodeling. The Blomstrand disease has been subdivided into **type I**, which refers to the severe (classical) form, and **type II**, which refers to a relatively milder variant, and the difference between severity is attributed to complete or incomplete inactivation of the PTHr1, respectively (261,262). A milder form of recessively inherited skeletal dysplasia, known as **Eiken syndrome** (MIM#600002), has also been linked to mutations of *PTHr1* (263). Dominantly acting *PTHr1* mutations have been identified in endochondromas of patients with **enchondromatosis (Ollier's disease - MIM#166000)**, a familial disorder with evidence of autosomal dominance characterized by multiple benign cartilage tumors, and a predisposition to malignant osteocarcinomas (264,265). As many patients with Ollier's disease do not have *PTHr1* mutations, it is likely

that the condition is genetically heterogeneous (266). Dominantly inherited ***symmetrical enchondromatosis*** is associated with duplication of 12p11.23 to 12p11.22 that includes the *PTHLH* gene encoding PTHrP suggesting that abnormal PTHR1 signaling may underlie this unusual form of endochondromatosis (267). In addition, some cases of autosomal dominant nonsyndromic ***primary failure of tooth eruption (PFE)*** are due to loss-of-function mutations in the PTHR1 that are dominantly acting, leading to haploinsufficiency of the receptor (268-272).

HYPOMAGNESEMIA

In humans, hypomagnesemia leads to a suppression of parathyroid hormone release and some degree of peripheral resistance. Although the exact molecular mechanism underlying the suppression of the parathyroid gland in hypomagnesemia is unknown, it is important to recognize that laboratory testing in cases of hypocalcemia with reduced PTH should include measurement of serum magnesium, particularly in newborns (273). Primary hypomagnesemia with secondary hypocalcemia (HSH) is an autosomal recessive disorder characterized by neuromuscular symptoms in infancy due to extremely low levels of serum magnesium and moderate to severe hypocalcemia. Homozygous mutations in the magnesium transporter gene transient receptor potential cation channel member 6 (TRPM6) cause the disease. HSH, a potentially lethal condition, can be misdiagnosed as primary hypoparathyroidism (274). Long-term prognosis after treatment with high dose of oral magnesium supplementation is good. Hypomagnesemia is also associated with long-term use of proton-pump inhibitors that decrease the luminal pH of the intestine by acting on the enterocyte apical TRPM6/7 channels (275,276).

MANAGEMENT OF HYPOPARATHYROIDISM

Calcium and Vitamin D. The goal of treatment in hypoparathyroid states is to raise the serum calcium sufficiently to alleviate acute symptoms of hypocalcemia and prevent the chronic complications (3,277,278). The calcium concentration required for this purpose is generally the low-normal range. Acute or severe symptomatic hypocalcemia is best treated with intravenous calcium infusion. Initial doses of 2 to 5 millimoles of elemental calcium as the gluconate salt can be given over a 10 to 20 minute period, followed by 2 millimoles elemental calcium per hour as a maintenance dose, to be adjusted according to symptoms and biochemical response. Care must be taken to ensure that the infusion does not extravasate, and ionized or total calcium levels should be monitored frequently on a stat basis. Doses in children 5 to 14 years of age need to be adjusted for body weight, while neonates and infants require age-specific dosing schedules. In adults, intravenous vitamin D therapy is not needed. Hyperphosphatemia, alkalosis and hypomagnesemia should be corrected concomitantly if present. Post-surgical hypocalcemia is now rarely severe and usually transient with appropriate management (279). However, the occasional patient can represent a significant problem, particularly if the indication for surgery is chronic hyperparathyroidism, and the post-operative hypoparathyroid state is permanent (280). The long-term effects of standard therapy, hypercalciuria, nephrolithiasis, nephrocalcinosis, ectopic tissue calcification and mood changes, remain a concern (270,281).

The mainstay of chronic treatment is oral calcium and vitamin D, which should be started as soon as possible to allow reduction and discontinuation of the intravenous calcium. Oral calcium comes in several forms, but **calcium carbonate** is generally the least expensive. A total of 20 to 80 millimoles elemental calcium daily (2 to 8 g calcium carbonate per day) is generally effective, but should be given in divided doses and adjusted on the basis of gastro-intestinal tolerance, relief of hypocalcemic symptoms, and appropriate biochemical response. Vitamin D is preferably administered as **calcitriol** (0.25 to 1.0 micrograms per day) but, if cost is a factor, pharmacological doses of cholecalciferol or ergocalciferol or calcidiol may be less expensive and equally efficacious (282). Cholecalciferol and ergocalciferol have the longest duration of action and can result in sustained toxicity. It is therefore appropriate to institute a starting dose of 25,000 IU/day and titrate upwards (to

100,000 IU/daily) with assessment of serum and urinary parameters afterwards with follow-up at 6 and 12 months, even if the patient is relatively asymptomatic. Some authorities recommend the use of active vitamin D (calcitriol or alphacalcidol) where possible providing the rationale that the lack of PTH along with the tendency to hyperphosphatemia impairs the renal conversion of 25-hydroxyvitamin D to active vitamin D (277,278). In any event, serum calcium and 24 hour urinary excretion should be carefully monitored when therapy is started and continued until the patient is stabilized. Hypercalciuria that occurs as treatment is initiated, even prior to the normalization of the serum calcium, may warrant an ongoing assessment of nephrocalcinosis, most sensitively detected by renal ultrasound. Consequently, only a low-normal serum calcium concentration may be attainable, but many patients feel well enough that there is no need to entirely normalize the serum calcium. That way, the risk of renal failure due to chronic hypercalciuria – especially problematic in patients with *CASR* activating mutations (6,7) – is minimized. Nevertheless, a significant number of patients report problems with easy fatigue and exhaustion, and mood disturbances (e.g., depression, anxiety, hostility, and paranoid ideation) not in keeping with the degree of hypocalcemia, suggesting that there may be non-calcitropic effects of PTH not remedied by maintenance of normocalcemia alone (281). In an epidemiological and health-related quality of life study from Norway, postsurgical hypoparathyroid patients scored worse than those with nonsurgical hypoparathyroidism or pseudohypoparathyroidism (283).

In pseudohypoparathyroidism, monitoring serum PTH levels during treatment is critical with the aim of normalizing or reducing PTH levels as much as possible. This is done to avoid the long-term elevation of circulating PTH, that would likely cause bone resorption. Also, hypercalciuria as a result of the calcitriol and calcium treatment is a lesser concern because PTH actions in the distal tubule are functional, preventing the loss of calcium in the urine. Of note, **calcitriol** (and not other forms of vitamin D) should be used for the treatment, because the PTH resistance in the proximal tubule does not allow for the efficient synthesis of 1,25(OH)₂D from 25-hydroxyvitamin D (also see comments above).

Hormone replacement therapy.

Hormone replacement has been advocated as a potentially superior form of treatment for decades but only recently have preparations of recombinant human hormone — teriparatide (PTH 1-34) and full-length parathyroid hormone (PTH 1-84) — become available. In 2015, the U.S. Food and Drug Administration (FDA) approved recombinant human (rh) PTH (1-84) for the management of hypoparathyroidism (284). This provides an additional therapeutic option for the management of those patients who demonstrate poor control with the standard calcium and active vitamin D supplemental therapy. The FDA indication is for subjects with hypoparathyroidism of any etiology, except ADH, but including postsurgical cases. The FDA approved rhPTH(-84) with a “black box” warning because of the history of rat osteosarcoma and PTH use (285), although no evidence for this in primates or in clinical use has been forthcoming (286), but did not limit the duration of use.

The use of PTH in hypoparathyroidism was demonstrated initially with the amino-terminal fragment of PTH, teriparatide [PTH (1–34)] (287). Beneficial control in children and in adults occurred when teriparatide was administered daily with better control when the peptide was administered in twice-daily dosing regimens (287-291). With a pump delivery system by which teriparatide could be administered continuously (292,293), urinary calcium excretion fell and markers of bone turnover normalized. A smaller daily dose was required with pump delivery vs multiple daily dosing regimens. An open-label trial of PTH (1–34) in adult subjects with postsurgical hypoparathyroidism showed improvement in quality of life (294). Beneficial effects on calcium homeostasis have also been demonstrated in specific ADH cases with activating *CaSR* mutations (295,296).

The full-length PTH (1-84) represents the actual secreted product of the parathyroid gland and its longer biological half-life (than PTH(1-34) makes once-daily dosing feasible (297-299). Studies by several groups have noted a substantial reduction in the requirement for calcium and active vitamin D (300-302); only transient reductions in urinary calcium excretion (299); a tendency for lumbar spine bone mineral density

(BMD) to increase and that of the distal one-third radius to fall (301); a rapid increase in bone turnover, assessed by circulating markers and dynamic histomorphometric analyses of bone that achieves a new steady state that is higher than baseline values within 2–3 years (303); and improvements in quality of life in some studies (302,304).

In a placebo-controlled 24-week clinical trial of rhPTH (1-84) in 130 hypoparathyroid patients the primary endpoints of a reduction by 50% in calcium supplements and in active vitamin D along with maintenance of the serum calcium were met in over half of the study subjects (305). There was a greater percentage of subjects in whom active vitamin D could be eliminated entirely while taking no more than 500 mg of oral calcium daily. The drug was titrated from 50 to 100 µg/d, with just over half of the subjects needing the highest dose. The rhPTH(1-84) reduced serum phosphate levels, improved the calcium-phosphate product, and maintained 1,25(OH)₂D and serum calcium levels in the normal range (306). For discussion of selection and management of patients for PTH(1-84) therapy as well as safety issues and research agenda for the future the reader is referred to (277,278).

In the future, analogs of PTH may be useful for the treatment of hypoparathyroidism. To facilitate their study, diphtheria toxin- and GFP-based mouse models of acquired hypoparathyroidism have been developed (307). In each model, a single subcutaneous injection of a long acting PTH analog increased serum calcium levels more effectively and for a longer time than a 10-fold greater dose of PTH(1-34) (307). Also of note is that orally active small-molecule PTHR1 agonists are being developed for the treatment of hypoparathyroidism. Oral administration of one such compound, PCO371, to osteopenic rats provoked a significant increase in bone turnover with some increase in bone mass. In hypocalcemic rats, the compound restored serum calcium levels without increasing urinary calcium, and the effects were stronger and longer-lasting than with PTH injections (308).

Calcilytics.

Calcilytics – small molecule allosteric modulators of the CASR that antagonize the calcium-sensing receptor and promote PTH secretion – are a promising alternative for disorders with intact but hypofunctioning parathyroid glands (309). In clinical studies in patients with ADH1, dosing with PTH(1-34) leads to better control of blood calcium levels (288). However, PTH does not correct the effects of the activated CASR in the kidney, and urinary calcium excretion may not be normalized in contrast to cases of postsurgical hypoparathyroidism (292,296). Thus while FDA approval was given for PTH treatment of hypoparathyroidism, ADH1 was excluded from the indication.

Calcilytics which inhibit the activation of the CASR in the parathyroid and renal tubule that promote not only PTH secretion but also renal calcium reabsorption are therefore of potential interest for the treatment of ADH1. In cell culture experiments studying activating CASR mutants, calcilytics normalize the left-ward shift of the calcium response curve (310,311). The utility of calcilytics was further demonstrated in studies of mice harboring activating *Casr* mutations. In one study, two knock-in mouse models of ADH1 with activating mutations in the *Casr* were generated. Daily oral administration of the calcilytic JTT-305/MK-3442 to these mice increased serum PTH and calcium levels, and reduced urinary calcium excretion (310). Intraperitoneal injection of the calcilytic NPS2143 in the *nuf* mouse model of ADH1, transiently increased circulating PTH and calcium levels without increasing urinary calcium levels (311). In a preliminary clinical study, IV administration of the calcilytic NPSP795 to five patients with ADH1 increased their plasma PTH levels and decreased their fractional urinary calcium excretion (312). Calcilytics comprise two main classes of compounds; the **amino alcohols** (e.g., NPS2143, NPSP795, JTT-305/MK-5442) and the **quinazolinones** (e.g., ATF936 and AXT914) (309). While both classes of compounds corrected the gain-of-function properties of several of the ADH1 CASR mutations tested *in vitro*, a subset of mutations involving NPS2143 binding sites within the transmembrane domain of the CASR are not fully corrected with NPS2143 but are normalized with the quinazolinone drugs (ATF936 and AXT914) (313-315). Whether this is reflected in mouse model studies and clinical situations remains to be determined.

Cases of hypoparathyroidism presenting as ADH but without *CASR* mutations have been found to have activating mutations of $G\alpha_{11}$ that couples the *CASR* to signaling pathways (316,317). The syndrome has been designated ADH2. Even though $G\alpha_{11}$ is downstream of the *CASR* the calcilytic NPS2143 has been shown to rectify the altered Ca^{2+} signaling of the overactive mutants in *in vitro* studies (318). Knock-in mice harboring an ADH2 $G\alpha_{11}$ activating mutation faithfully replicate ADH2 (319). Treatment with the calcilytic NPS2143 or a $G\alpha_{11}/q$ -specific inhibitor, YM-254890 (320), increased circulating PTH and calcium levels in the heterozygous mutant mice (319). Thus, calcilytics, by blocking the renal *CASR*, are potentially of use to treat ADH1 and ADH2, and $G\alpha_{11}$ inhibitors to treat ADH2 as well as other forms of hypoparathyroidism.

Other therapies.

If the serum calcium attainable with oral calcium and calcitriol is below the normal range and the patient remains symptomatic, then a trial of a [thiazide diuretic](#) may be considered, with the aim of reducing the hypercalciuria to raise the serum calcium further. The argument for efficacy seems greatest for responsive forms of autosomal dominant hypocalcemia due to activating *CaSR* mutations, since the thiazide-sensitive transporter, *SLC12A3* (MIM#600968), is a downstream target of and is suppressed by activated *CaSR* in the kidney. For reasons that are not clear, however, thiazides work well in some patients (321) but not others. It is critical to monitor serum potassium and magnesium levels as thiazide use can lead to renal losses of these cations with resulting hypokalemia and hypomagnesemia. Some authorities suggest thiazides should not be used in APS1 patients with adrenal insufficiency and in ADH1 patients with Bartter syndrome type V (277,278).

As the serum calcium is normalized, elevated serum phosphate concentrations generally decline but phosphate-binding gels such as aluminum hydroxide are occasionally helpful in reducing hyperphosphatemia at the beginning of therapy.

In patients with intracranial calcifications, patients may experience seizures related to chronic neuropathic changes and it may be necessary to add appropriate anti-epileptic medication(s). In all chronically hypocalcemic patients, ocular assessments should be performed periodically.

REFERENCES

1. Shoback, DM, Bilezikian JP, Costa AG, et al. (2016). Presentation of hypoparathyroidism: etiologies and clinical features. *J Clin Endocrinol Metab.* 101, 2300-2312.
2. Clarke BI, Brown EM, Collins MT, et al. (2016) Epidemiology and diagnosis of hypoparathyroidism. *J Clin Endocrinol Metab.* 101, 2284-2299.
3. Schafer AL and Shoback DM (2016) Hypocalcemia: diagnosis and treatment. In: De Groot LJ, Chrousos AG, Dungan K, Grossman A, Hershman JM, Koch C, McLachlan R, New M, Rebar R, Singer F, Vinik A, Weickert MP. Editors. *Endotext* [Internet], South Dartmouth (MA): MDText.com. Inc.; 2000-2016. Jan 3.
4. Fisher, N. G., Armitage, A., McGonigle R.J., and Gilbert, T. J. (2001). Hypocalcaemic cardiomyopathy; the relationship between myocardial damage, left ventricular function, calcium and ECG changes in a patient with idiopathic hypocalcaemia. *Eur. J. Heart Failure.* 3, 373-376.

5. Tziomalos, K., Kakavas, N., Kountana, E., Harsoulis, F., Basayannis, E. (2006). Reversible dilated hypocalcaemic cardiomyopathy in a patient with primary hypoparathyroidism. *Clin Endocrinol (Oxf)*. 64, 717-718.
6. Lienhardt, A., Bai, M., Lagarde, J. P., et al. (2001). Activating mutations of the calcium-sensing receptor: management of hypocalcemia. *J Clin. Endocrinol. Metab.* 86, 5313-5323.
7. Obermannova B, Sumnik Z, Dusatkova P, et al. (2016). Novel calcium-sensing receptor cytoplasmic tail deletion mutation causing autosomal dominant hypocalcemia: molecular and clinical study. *Eur J Endocrinol.* 174, K1-K11.
8. McKusick, V. (2001). *Online Mendelian Inheritance in Man (OMIM)*. Baltimore: Johns Hopkins University Press.
9. Arnold, A., Horst, S. A., Gardella, T. J., Baba, H., Levine, M. A., and Kronenberg, H. M. (1990). Mutation of the signal peptide-encoding region of the preproparathyroid hormone gene in familial isolated hypoparathyroidism. *J. Clin. Invest.* 86, 1084-1087.
10. Datta, R., Waheed, A., Shah, G.N., and Sly, W.S. (2007). Signal sequence mutation in autosomal dominant form of hypoparathyroidism induces apoptosis that is corrected by a chemical chaperone. *Proc. Natl. Acad. Sci. USA.* 104, 19989-19994.
11. Sunthornthepvarakul, T., Churesigaew, S., and Ngowngarmratana, S. (1999). A novel mutation of the signal peptide of the preproparathyroid hormone gene associated with autosomal recessive familial isolated hypoparathyroidism. *J. Clin. Endocrinol. Metab.* 84, 3792-3796.
12. Ertl, D-A., Stary, S., Streubel, B., Raimann, A., Haeusler, G. (2012). A novel homozygous mutation in the parathyroid hormone gene (*PTH*) in a girl with isolated hypoparathyroidism. *Bone* 51, 629-632.
13. Lee S, Mannstadt M, Guo J, et al. (2015). A homozygous [Cys25]PTH(1-84) mutation that impairs PTH/PTHrP receptor activation defines a novel form of hypoparathyroidism. *J Bone Miner Res.* 30, 1803-1813.
14. Parkinson, D. B., and Thakker, R. V. (1992). A donor splice site mutation in the parathyroid hormone gene is associated with autosomal recessive hypoparathyroidism. *Nat. Genet.* 1, 149-152.
15. Grigorieva IV, Thakker RV. (2011). Transcription factors in parathyroid development: lessons from hypoparathyroid disorders. *Ann N Y Acad Sci.* 1237, 24-38.
16. Bowl, M.R., Nesbit, M.A., Harding, B, et al. (2005). An interstitial deletion-insertion involving chromosomes 2p25.3 and Xq27.1, near SOX3, causes X-linked recessive hypoparathyroidism. *J. Clin. Invest.* 115, 2822-2831.
17. Pollak, M. R., Brown, E. M., Estep, H. L., et al. (1994). Autosomal dominant hypocalcaemia caused by a Ca²⁺-sensing receptor gene mutation. *Nat. Genet.* 8, 303-307.

18. Hendy, G. N., D'Souza-Li, L., Yang, B., Canaff, L., and Cole, D. E. (2000). Mutations of the calcium-sensing receptor (CASR) in familial hypocalciuric hypercalcemia, neonatal severe hyperparathyroidism, and autosomal dominant hypocalcemia. *Hum. Mutat.* 16, 281-296.
19. Cole, D.E.C., Yun, F.H., Wong, B.Y., et al. (2009). Calcium-sensing receptor mutations and denaturing high performance liquid chromatography. *J. Mol. Endocrinol.* 42, 331-339.
20. Hendy, G.N., Guarnieri, V., and Canaff, L. (2009). Calcium-sensing receptor and associated diseases. *Prog. Mol. Biol. Transl. Sci.* 89C, 31-95.
21. Hendy, G.N., Minutti, C., Canaff, L., et al. (2003). Recurrent familial hypocalcemia due to germline mosaicism for an activating mutation of the calcium-sensing receptor gene. *J. Clin. Endocrinol. Metab.* 88, 3674-3681.
22. Ray, K., Hauschild, B. C., Steinbach, P. J., Goldsmith, P. K., Hauache, O., and Spiegel, A. M. (1999). Identification of the cysteine residues in the amino-terminal extracellular domain of the human Ca^{2+} receptor critical for dimerization. Implications for function of monomeric Ca^{2+} receptor. *J. Biol. Chem.* 274, 27642-27650.
23. Pidasheva, S., D'Souza-Li, L., Canaff, L., Cole, D.E., and Hendy, G.N. (2004). CASRdb: calcium-sensing receptor locus-specific database for mutations causing familial (benign) hypocalciuric hypercalcemia, neonatal severe hyperparathyroidism, and autosomal dominant hypocalcemia. *Hum. Mutat.* 24, 107-111.
24. Hannan FM, Nesbit MA, Zhang C, et al. (2012) Identification of 70 calcium-sensing receptor mutations in hyper- and hypo-calcaemic patients: evidence for clustering of extracellular domain mutations at calcium-binding sites. *Hum Mol Genet.* 21, 2768-2778.
25. Vargas-Poussou R, Huang C, Hulin P, et al. (2002) Functional characterization of calcium-sensing receptor mutation in severe autosomal dominant hypocalcemia with a Bartter-like syndrome. *J Am Soc Nephrol.* 13, 2259-2266.
26. Watanabe S, Fukumoto S, Chang H, et al. (2002) Association between activating mutations of calcium-sensing receptor and Bartter's syndrome. *Lancet* 360, 692-691.
27. Vezzoli G, Arcidiacono T, Paloschi V, et al. (2006) Autosomal dominant hypocalcemia with mild type 5 Bartter syndrome. *J Am Soc Nephrol.* 19, 525-528.
28. Hebert SC. (2003) Bartter syndrome. *Curr Opin Nephrol Hypertens* 12, 527-532.
29. Carmosino M, Gerbino A, Hendy GN, et al. (2015) NKCC2 activity is inhibited by the Bartter's syndrome type 5 gain of function CaR-A843E mutant in renal cells. *Biol Cell.* 107, 98-110.
30. Nesbit MA, Hannan FM, Howles SA, et al. (2013) Mutations affecting G-protein subunit α_{11} in hypercalcemia and hypocalcemia. *N Engl J Med.* 368, 2476-2486.
31. Mannstadt M, Harris M, Bravenboer B, et al. (2013) Germline mutations affecting $\text{G}\alpha_{11}$ in hypoparathyroidism. *N Engl J Med.* 368, 2532-2534.

32. Li D, Opas EE, Tuluc F, et al. (2014) Autosomal dominant hypoparathyroidism caused by germline mutation in GNA11: phenotypic and molecular characterization. *J Clin Endocrinol Metab.* 99, E1774-E1783.
33. Piret SE, Gorvin CM, Pagnamenta AT, et al. (2016). Identification of a G-Protein Subunit- α 11 Gain-of-Function Mutation, Val340Met, in a Family With Autosomal Dominant Hypocalcemia Type 2 (ADH2). *J Bone Miner Res.* 31, 1207-1214.
34. Mizuno N, and Itoh H. (2009) Functions and regulatory mechanisms of Gq-signaling pathways. *Neurosignals* 17, 42-54
35. Hendy GN and Cole DEC. (2013) Ruling in a suspect: the role of AP2S1 mutations in familial hypocalciuric hypercalcemia type 3. *J Clin Endocrinol Metab.* 98, 4666-4669.
36. Lambert A-S, Grybek V, Francou B, et al. (2014) Analysis of AP2S1, a calcium-sensing receptor regulator, in familial and sporadic isolated hypoparathyroidism. *J Clin Endocrinol Metab.* 99, E469-E473.
37. Rogers A, Nesbit MA, Hannan FH, et al. (2014) Mutational analysis of the adaptor protein 2 sigma subunit (AP2S1) gene: search for autosomal dominant hypocalcemia type 3 (ADH3). *J Clin Endocrinol Metab.* 99, E1300-E1303.
38. Kanemura, Y., Hiraga, S., Arita, N., et al. (1999). Isolation and expression analysis of a novel human homologue of the *Drosophila* glial cells missing (gcm) gene. *FEBS Lett.* 442, 151-156.
39. Gunther, T., Chen, Z. F., Kim, J., et al. (2000). Genetic ablation of parathyroid glands reveals another source of parathyroid hormone. *Nature* 406, 199-203.
40. Okabe, M., and Graham, A. (2004). The origin of the parathyroid gland. *Proc. Natl. Acad. Sci. USA* 101, 17716-17719.
41. Ding, C., Buckingham, B., and Levine, M. A. (2001). Familial isolated hypoparathyroidism caused by a mutation in the gene for the transcription factor GCMB. *J. Clin. Invest.* 108, 1215-1220.
42. Thomee, C., Schubert, S.W., Parma, J. et al. (2005). GCMB mutation in familial isolated hypoparathyroidism with residual secretion of parathyroid hormone. *J. Clin. Endocrinol. Metab.* 90, 2487-2492.
43. Canaff, L., Zhou, X., Mosesova, I., Cole, D.E.C., and Hendy, G.N. (2009). Glial cells missing-2 (GCM2) transactivates the calcium-sensing receptor gene: effect of a dominant-negative GCM2 mutant associated with autosomal dominant hypoparathyroidism. *Hum. Mutat.* 30, 85-92.
44. Bowl, M.R., Mirczuk, S.M., Grigorieva, I.V., et al. (2010). Identification and characterization of novel parathyroid-specific transcription factor Glial Cells Missing Homolog B (GCMB) mutations in eight families with autosomal recessive hypoparathyroidism. *Hum. Mol. Genet.* 19, 2028-2038.

45. Mannstadt, M., Bertrand, G., Muresan, M., et al. (2008). Dominant-negative GCMB mutations cause an autosomal dominant form of hypoparathyroidism. *J. Clin. Endocrinol. Metab.* 93, 3568-3576.
46. Mirczuk, S.M., Bowl, M.R., Nesbit, M.A., et al. (2010). A missense glial cells missing homolog B (GCMB) mutation, Asn502His, causes autosomal dominant hypoparathyroidism. *J. Clin. Endocrinol. Metab.* 95, 3512-3516.
47. Hendy GN and Cole DEC. (2015) Chapter 16, Familial isolated hypoparathyroidism. In, *Hypoparathyroidism*. Brandi ML, Brown EM. Editors. Springer Verlag Italia. 167-175.
48. Maret, A., Ding, C., Kornfield, S.L., Levine, M.A. (2008). Analysis of the GCM2 gene in isolated hypoparathyroidism: a molecular and biochemical study. *J. Clin. Endocrinol. Metab.* 93, 1426-1432.
49. D'Agruma L, Coco M, Guarnieri V, et al. (2014) Increased prevalence of the GCM2 polymorphism, Y282D, in primary hyperparathyroidism: analysis of three Italian cohorts. *J Clin Endocrinol Metab.* 99, E2794-E2798.
50. Guan B, Welch JM, Sapp JC, et al. (2016). GCM2-activating mutations in Familial Isolated Hyperparathyroidism. *Am J Hum Genet.* 99, 1034-1044.
51. DiGeorge, A. M. (1965). Discussions on a new concept of the cellular basis of immunity. *J. Pediatr.* 67, 907-908.
52. McDonald-McGinn, D.M., and Sullivan, K.E. (2011). Chromosome 22q11.2 deletion syndrome (DiGeorge syndrome/velocardiofacial syndrome). *Medicine (Baltimore).* 90, 1-18.
53. Botto, L. D., May, K., Fernhoff, P. M., et al. (2003). A population based study of the 22q11.2 deletion: Phenotype, incidence and contribution to major birth defects in the population. *Pediatrics* 112, 101-107.
54. Yagi, H., Furutani, Y., Hamada, H., et al. (2003). Role of TBX1 in human del22q11.2 syndrome. *Lancet* 362, 1366-1373.
55. Baldini A, Fulcoli FG, and Illingworth E. (2017). Tbx1: Transcriptional and developmental functions. *Curr Top Dev Biol.* 122, 223-243.
56. Scambler PJ. (2010). 22q11 deletion syndrome: a role for TBX1 in pharyngeal and cardiovascular development. *Pediatr Cardiol.* 31, 378-390.
57. Huh, S.H., and Ornitz, D.M. (2010). Beta-catenin deficiency causes DiGeorge syndrome-like phenotypes through regulation of Tbx1. *Development.* 137, 1137-1147.
58. Guo, C., Sun, Y., Zhou, B., et al. (2011). A Tbx1-Six1/Eya1-Fgf8 genetic pathway controls mammalian cardiovascular and craniofacial morphogenesis. *J. Clin. Invest.* 121, 1585-1595.
59. Jerome, L. A., and Papaioannou, V. E. (2001) DiGeorge syndrome phenotype in mice mutant for the T-box gene, Tbx1. *Nat Genet.* 27, 286-291.

60. Ivins S, Lammerts van Beuren K, Roberts C, et al. (2005). Microarray analysis detects differentially expressed genes in the pharyngeal region of mice lacking Tbx1. *Dev Biol.* 2005 285, 554-569.
61. Garg V, Yamagishi C, Hu T, Kathiriya IS, Yamagishi H, and Srivastava D. (2001). Tbx1, a DiGeorge syndrome candidate gene, is regulated by sonic hedgehog during pharyngeal arch development. *Dev Biol.* 235, 62-73.
62. Scirè G, Dallapiccola B, Iannetti P, et al. (1994). Hypoparathyroidism as the major manifestation in two patients with 22q11 deletions. *Am J Med Genet.* 52, 478-482.
63. Sykes KS, Bachrach LK, Siegel-Bartelt J, Ipp M, Kooh SW, and Cytrynbaum C. (1997). Velocardiofacial syndrome presenting as hypocalcemia in early adolescence. *Arch Pediatr Adolesc Med.* 151, 745-747.
64. McDonald-McGinn DM, Sullivan KE, Marino B, et al. (2015). 22q11.2 deletion syndrome. *Nat Rev Dis Primers.* 1, 15071.
65. Liu, H., Abecasis, G. R., Heath, S., et al. (2002). Genetic variation in the 22q11 locus and susceptibility to schizophrenia. *Proc. Natl. Acad. Sci. USA* 99, 16859-16864.
66. Kapadia, R.K., and Bassett, A.S. (2008). Recognizing a common genetic syndrome: 22q11.2 deletion syndrome. *CMAJ.* 178, 391-393.
67. McDonald-McGinn DM, Emanuel BS, and Zackai EH. (2013). 22q11.2 deletion syndrome. In: Pagon RA, Adam MP, Ardinger HH, et al., editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle;
68. Burnside RD. (2015). 22q11.21 deletion syndromes: a review of proximal, central, and distal deletions and their associated features. *Cytogenet Genome Res.* 146, 89-99.
69. Miller, J. D., Bowker, B. M., Cole, D. E.C., and Guyda, H. J. (1983). DiGeorge's syndrome in monozygotic twins. Treatment with calcitriol. *Am. J. Dis. Child.* 137, 438-440.
70. Goodship, J., Cross, I., Scambler, P., and Burn, J. (1995). Monozygotic twins with chromosome 22q11 deletion and discordant phenotype. *J. Med. Genet.* 32, 746-748.
71. Hillebrand, G., Siebert, R., Simeoni, E., and Santer, R. (2000). DiGeorge syndrome with discordant phenotype in monozygotic twins. *J. Med. Genet.* 37, E23.
72. Carelle-Calmels, N., Saugier-veber, P., Girard-Lemaire, F., et al. (2009). Genetic compensation in a human genomic disorder. *N. Engl. J. Med.* 360, 1211-1216.
73. Alkalay, A.A., Guo, T., Montagna, C., et al. (2011). Genetic dosage compensation in a family with velo-cardio-facial/DiGeorge/22q11.2 deletion syndrome. *Am. J. Med. Genet. A* 155, 548-554.
74. Weinzimer, S.A. (2001). Endocrine aspects of the 22q11.1 deletion. *Genet. Med.* 3, 19-22.

75. Nagasaki, K., Iwasaki, Y., Ogawa, Y., Kikuchi, T., and Uchiyama, M. (2011). Evaluation of parathyroid gland function using sodium bicarbonate infusion test for 22q11.2 deletion syndrome. *Horm. Res. Paediatr.* 75, 14-18.
76. Jatana, V., Gillis, J., Webster, B.H., and Adès, L.C. (2007). Deletion 22q11.2 syndrome--implications for the intensive care physician. *Pediatr. Crit. Care Med.* 8, 459-463.
77. Jensen TJ, Dzakula Z, Deciu C, van den Boom D, and Ehrich M. (2012). Detection of microdeletion 22q11.2 in a fetus by next-generation sequencing of maternal plasma. *Clin Chem.* 58, 1148-1151.
78. Tenhola S, Hendy GN, Valta H, et al. (2015). Cinacalcet treatment in an adolescent with concurrent 22q11.2 deletion syndrome and familial hypocalciuric hypercalcemia type 3 caused by AP2S1 mutation. *J Clin Endocrinol Metab.* 100, 2515-2518.
79. Lima, K., Abrahamsen, T.G., Wolff, A.S., et al. (2011). Hypoparathyroidism and autoimmunity in the 22q11.2 deletion syndrome. *Eur. J. Endocrinol.* 165, 345-352.
80. Meek, C.L., Kaplan, F., Pereira, R.S., Viljoen, A. (2011). Hypocalcemia following treatment for hyperthyroidism. *Clin. Chem.* 57, 811-814.
81. Daw, S. C., Taylor, C., Kraman, M., et al. (1996). A common region of 10p deleted in DiGeorge and velocardiofacial syndromes. *Nat. Genet.* 13, 458-460.
82. Gottlieb, S., Driscoll, D. A., Punnett, H. H., Sellinger, B., Emanuel, B. S., and Budarf, M. L. (1998). Characterization of 10p deletions suggests two nonoverlapping regions contribute to the DiGeorge syndrome phenotype. *Am. J. Hum. Genet.* 62, 495-498.
83. Schuffenhauer, S., Lichtner, P., Peykar-Derakhshandeh, P., et al. (1998). Deletion mapping on chromosome 10p and definition of a critical region for the second DiGeorge syndrome locus (DGS2). *Eur. J. Hum. Genet.* 6, 213-225.
84. Barakat, A. Y., D'Albora, J. B., Martin, M. M., and Jose, P. A. (1977). Familial nephrosis, nerve deafness, and hypoparathyroidism. *J. Pediatr.* 91, 61-64.
85. Bilous, R. W., Murty, G., Parkinson, D. B., et al. (1992). Autosomal dominant familial hypoparathyroidism, sensorineural deafness, and renal dysplasia. *N. Engl. J. Med.* 327, 1069-1074.
86. Van Esch, H., Groenen, P., Daw, S., et al. (1999). Partial DiGeorge syndrome in two patients with a 10p rearrangement. *Clin. Genet.* 55, 269-276.
87. Lichtner, P., Konig, R., Hasegawa, T., Van Esch, H., Meitinger, T., and Schuffenhauer, S. (2000). An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14. *J. Med. Genet.* 37, 33-37.
88. Van Esch, H., Groenen, P., Nesbit, M. A., et al. (2000). GATA3 haplo-insufficiency causes human HDR syndrome. *Nature* 406, 419-422.

89. Nesbit, M. A., Bowl, M. R., Harding, B., et al. (2004). Characterization of GATA3 mutations in the hypoparathyroidism, deafness, and renal dysplasia (HDR) syndrome. *J. Biol. Chem.* 279, 22624-22634.
90. Zahirieh, A., Nesbit, M.A., Ali, A., et al. (2005). Functional analysis of a novel GATA3 mutation in a family with the hypoparathyroidism, deafness, and renal dysplasia syndrome. *J. Clin. Endocrinol. Metab.* 90, 2445-2450.
91. Ali, A., Christie, P.T., Grigorieva, I.V., et al. (2007). Functional characterization of GATA3 mutations causing the hypoparathyroidism-deafness-renal (HDR) dysplasia syndrome: insight into mechanism of DNA binding by the GATA3 transcription factor. *Hum. Mol. Genet.* 16, 265-275.
92. Nesbit, M.A. (2015) Chapter 19, Hypoparathyroidism., deafness, and renal anomaly syndrome. In, Hypoparathyroidism. Brandi ML, Brown EM. Editors. Springer Verlag Italia. 199-213.
93. Grigorieva, I.V., Mirczuk, S., Gaynor, K.U., et al. (2010). Gata3-deficient mice develop parathyroid abnormalities due to dysregulation of the parathyroid-specific transcription factor Gcm2. *J. Clin. Invest.* 120, 2144-2155.
94. Kenny, F. M., and Linarelli, L. (1966). Dwarfism and cortical thickening of tubular bones. Transient hypocalcemia in a mother and son. *Am. J. Dis. Child.* 111, 201-207.
95. Caffey, J. (1967). Congenital stenosis of medullary spaces in tubular bones and calvaria in two proportionate dwarfs--mother and son; coupled with transitory hypocalcemic tetany. *Am. J. Roentgenol. Radium. Ther. Nucl. Med.* 100, 1-11.
96. Bergada I, Schiffrin A, Abu Srair H, et al. (1988). Kenny syndrome: description of additional abnormalities and molecular studies. *Hum Genet.* 80(1):39-42.
97. HersHKovitz E, and Parvari R. (2015). Chapter 20. Hypoparathyroidism, dwarfism, medullary stenosis of long bones, and eye abnormalities (Kenny-Caffey syndrome) and hypoparathyroidism, retardation, and dysmorphism (Sanjad-Sakati syndrome). In, ML Brandi, EM Brown (eds) Hypoparathyroidism. Springer-Verlag Italia. pp. 215-224.
98. Unger S, Górna MW, Le Béhec A, et al. (2013). FAM111A mutations result in hypoparathyroidism and impaired skeletal development. *Am J Hum Genet.* 92(6):990-995.
99. Isojima T, Doi K, Mitsui J, et al. (2014). A recurrent de novo FAM111A mutation causes Kenny-Caffey syndrome type 2. *J Bone Miner Res.* 29(4):992-998.
100. Abraham MB, Li D, Tang D, et al. (2017). Short stature and hypoparathyroidism in a child with Kenny-Caffey syndrome type 2 due to a novel mutation in FAM111A gene. *Int J Pediatr Endocrinol.* 2017:1.
101. Alabert C, Bukowski-Wills JC, Lee SB, et al. (2014). Nascent chromatin capture proteomics determines chromatin dynamics during DNA replication and identifies unknown fork components. *Nat Cell Biol.* 16(3):281-293.
102. Teebi, A. S. (2000). Hypoparathyroidism, retarded growth and development, and dysmorphism or Sanjad-Sakati syndrome: an Arab disease reminiscent of Kenny-Caffey syndrome. *J. Med. Genet.* 37, 145.

103. Parvari, R., HersHKovitz, E., Kanis, A., et al. (1998). Homozygosity and linkage-disequilibrium mapping of the syndrome of congenital hypoparathyroidism, growth and mental retardation, and dysmorphism to a 1-cM interval on chromosome 1q42-43. *Am. J. Hum. Genet.* 63, 163-169.
104. HRD/Autosomal Recessive Kenny-Caffey Syndrome Consortium. (2002). Mutation of TBCE causes hypoparathyroidism-retardation-dysmorphism and autosomal recessive Kenny-Caffey syndrome. *Nat. Genet.* 32, 448-45
105. Voloshin, O., Gocheva, Y., Gutnick, M., et al. (2010). Tubulin chaperone E binds microtubules and proteasomes and protects against misfolded protein stress. *Cell. Mol. Life Sci.* 67, 2025-2038.
106. Parvari, R., Diaz, G.A., and HersHKovitz, E. (2007). Parathyroid development and the role of tubulin chaperone E. *Horm. Res.* 67, 12-21.
107. Chow J, Rahman J, Achermann JC, Dattani MT, and Rahman S. (2017). Mitochondrial disease and endocrine dysfunction. *Nat Rev Endocrinol.* 13(2):92-104.
108. Seneca, S., De Meirleir, L., De Schepper, J., et al. (1997). Pearson marrow pancreas syndrome: a molecular study and clinical management. *Clin. Genet.* 51, 338-342.
109. Tengan, C. H., Kiyomoto, B. H., Rocha, M. S., et al. (1998). Mitochondrial encephalomyopathy and hypoparathyroidism associated with a duplication and a deletion of mitochondrial deoxyribonucleic acid. *J. Clin. Endocrinol. Metab* 83, 125-129.
110. Wilichowski, E., Gruters, A., Kruse, K., et al. (1997). Hypoparathyroidism and deafness associated with pleioplasmic large scale rearrangements of the mitochondrial DNA: a clinical and molecular genetic study of four children with Kearns-Sayre syndrome. *Pediatr. Res.* 41, 193-200.
111. Naiki M, Ochi N, Kato YS, et al. (2014). Mutations in *HADHB*, which encodes the β -subunit of mitochondrial trifunctional protein, cause infantile onset hypoparathyroidism and peripheral polyneuropathy. *Am J Med Genet A.* 164A(5):1180-1187.
112. Tyni, T., Rapola, J., Palotie, A., and Pihko, H. (1997). Hypoparathyroidism in a patient with long-chain 3-hydroxyacyl-coenzyme A dehydrogenase deficiency caused by the G1528C mutation. *J. Pediatr.* 131, 766-768.
113. Carpenter TO, Carnes DL Jr, Anast CS. (1983). Hypoparathyroidism in Wilson's disease. *N Engl J Med.* 309(15), 873-877.
114. Brown, E.M. (2009). Anti-parathyroid and anti-calcium sensing receptor antibodies in autoimmune hypoparathyroidism. *Endocrinol. Metab. Clin. North. Am.* 38, 437-445.
115. Kemp EH, and Weetman AP. (2015). Chapter 17. Autoimmune hypoparathyroidism. In, ML Brandi, EM Brown (eds) *Hypoparathyroidism*. Springer-Verlag Italia. pp. 177-188.
116. Alimohammadi, M., Björklund, P., Hallgren, A., et al. (2008). Autoimmune polyendocrine syndrome type 1 and NALP5, a parathyroid autoantigen. *N. Engl. J. Med.* 358, 1018-1028.

117. Eisenbarth, G. S., and Gottlieb, P. A. (2004). Autoimmune polyendocrine syndromes. *New Engl. J. Med.* 350, 2068-2079.
118. Li, Y., Song, Y. H., Rais, N., et al. (1996). Autoantibodies to the extracellular domain of the calcium sensing receptor in patients with acquired hypoparathyroidism. *J. Clin. Invest.* 97, 910-914.
119. Goswami, R., Brown, E. M., Kochupillai, N., et al. (2004). Prevalence of calcium sensing receptor autoantibodies in patients with sporadic idiopathic hypoparathyroidism. *Eur. J. Endocrinol.* 150, 9-18.
120. Soderbergh, A., Myhre, A.G., Ekwall, O., et al. (2004). Prevalence and clinical associations in 10 defined autoantibodies in autoimmune polyendocrine syndrome type 1. *J. Clin. Endocrinol. Metab.* 89, 557-562.
121. Gylling, M., Kaariainen, E., Vaisanen, R., et al. (2003) The hypoparathyroidism of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy: protective effect of male sex. *J. Clin. Endocrinol. Metab.* 88, 4602-4608.
122. Kemp, E.H., Gavalas, N.G., Krohn, K.J., Brown, E.M., Watson, P.F., Weetman, A.P. (2009). Activating autoantibodies against the calcium-sensing receptor detected in two patients with autoimmune polyendocrine syndrome type 1. *J. Clin. Endocrinol. Metab.* 94, 4749-4756.
123. Mayer, A., Ploix, C., Orgiazzi, J., et al. (2004). Calcium-sensing receptor autoantibodies are relevant markers of acquired hypoparathyroidism. *J. Clin. Endocrinol. Metab.* 89, 4484-4488.
124. Eisenbarth, G. S., and Gottlieb, P. A. (2004). Hypoparathyroidism and autoimmune polyendocrine syndromes. *New Engl. J. Med.* 351, 1032-1033.
125. Kifor, O., McElduff, A., LeBoff, M.S., et al. (2004). Activating antibodies to the calcium-sensing receptor in two patients with autoimmune hypoparathyroidism. *J. Clin. Endocrinol. Metab.* 89, 548-556.
126. Buzi, F., Badolato, R., Mazza, C., et al. (2003). Autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy syndrome: Time to review diagnostic criteria? *J. Clin. Endocrinol. Metab.* 88, 3146-3148.
127. Ferre EM, Rose SR, Rosenzweig SD, et al. (2016). Redefined clinical features and diagnostic criteria in autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy. *JCI Insight.* 1(13). pii: e88782.
128. Finnish-German APECED Consortium (1997). An autoimmune disease, APECED, caused by mutations in a novel gene featuring two PHD-type zinc-finger domains. *Nat. Genet.* 17, 399-403.
129. Nagamine, K., Peterson, P., Scott, H. S., et al. (1997). Positional cloning of the APECED gene. *Nat. Genet.* 17, 393-398.
130. Halonen, M., Kangas, H., Ruppell, T., et al. (2004). APECED-causing mutations in AIRE reveal the functional domains of the protein. *Hum. Mutat.* 23, 245-257.
131. Akirav, E.M., Ruddle, N.H., and Herold KC. (2011). The role of AIRE in human autoimmune disease. *Nat Rev Endocrinol.* 7, 25-33.

132. Cervato, S., Mariniello, B., Lazzarotto, F., et al. (2009). Evaluation of the autoimmune regulator (AIRE) gene mutations in a cohort of Italian patients with autoimmune-polyendocrinopathy-candidiasis-ectodermal-dystrophy (APECED) and in their relatives. *Clin. Endocrinol. (Oxf)*. 70, 421-428.
133. Cheng, M.H., Fan, U., Grewal, N., et al. (2010). Acquired autoimmune polyglandular syndrome, thymoma, and an AIRE defect. *N. Engl. J. Med.* 362, 764-766.
134. Wilson, L. C., and Trembath, R. C. (1994). Albright's hereditary osteodystrophy. *J. Med. Genet.* 31, 779-784.
135. Levine, M. A. (1999). Pseudohypoparathyroidism: from bedside to bench and back. *J. Bone Miner. Res.* 14, 1255-1260.
136. Levine, M. A. (2000). The molecular basis for parathyroid hormone resistance in pseudohypoparathyroidism. In "The Genetics of Osteoporosis and Metabolic Bone Disease", (Econs, M.J. ed.), pp.179-209. Humana Press, Totawa NJ.
137. Spiegel, A. M. and Weinstein, L. S. (2001). Pseudohypoparathyroidism. In "The Metabolic and Molecular Bases of Inherited Disease", 8th ed. (Scriver, C. R., Beaudet, A. L., Sly, W. S., and Valle, D. eds.), pp.4205-4221. McGraw-Hill, New York.
138. Chase, L.R., Melson, G.L., and Aurbach, G.D. (1969) Pseudohypoparathyroidism: defective excretion of 3',5'-AMP in response to parathyroid hormone. *J. Clin. Invest.* 48, 1832-1844
139. Albright, F., Burnett, C. H., and Smith, P. H. (1942). Pseudohypoparathyroidism: An example of 'Seabright-Bantam syndrome'. *Endocrinol.* 30, 922-932.
140. De Sanctis, L., Vai, S., Andreo, M.R., et al. (2004). Brachydactyly in 14 genetically characterized pseudohypoparathyroid type 1a patients. *J. Clin. Endocrinol. Metab.* 89, 1650-1655.
141. Puzhko, S., Goodyer, C.G., Kerachian, M.A., et al. (2011) Parathyroid hormone signaling via G α s is selectively inhibited by an NH₂-terminally truncated G α s: implications for pseudohypoparathyroidism. *J. Bone Miner. Res.* 26, 2473-2485.
142. Poznanski, A.K., Werder, E.A., Giedion, A., Martin, A., Shaw, H. (1977). The pattern of shortening of the bones of the hand in PHP and PPHP--A comparison with brachydactyly E, Turner Syndrome, and acro dysostosis. *Radiology* 123, 707-718.
143. Brickman, A. S., Stern, N., and Sowers, J. R. (1988). Hypertension in pseudohypoparathyroidism type I. *Am. J. Med.* 85, 785-792.
144. Koch, T., Lehnhardt, E., Bottinger, H., et al. (1990). Sensorineural hearing loss owing to deficient G proteins in patients with pseudohypoparathyroidism: results of a multicentre study. *Eur. J. Clin. Invest.* 20, 416-421.
145. Goadsby, P. J., Lollin, Y., and Kocen, R. S. (1991). Pseudopseudohypoparathyroidism and spinal cord compression. *J Neurol. Neurosurg. Psychiatry* 54, 929-931.

146. Maeda, S.S., Fortes, E.M., Oliveira, U.M., Borba, V.C., and Lazaretti-Castro, M. (2006). Hypoparathyroidism and pseudohypoparathyroidism. *Arq. Bras. Endocrinol. Metab.* 50, 664-673.
147. Ong, K. K., Amin, R., and Dunger, D. B. (2000). Pseudohypoparathyroidism--another monogenic obesity syndrome. *Clin. Endocrinol. (Oxf)* 52, 389-391.
148. Chen, M., Wang, J., Dickerson, K.E., et al. (2009). Central nervous system imprinting of the G protein G(s) alpha and its role in metabolic regulation. *Cell Metab.* 9, 548-555.
149. Phelan, M. C., Rogers, R. C., Clarkson, K. B., et al. (1995). Albright hereditary osteodystrophy and del(2) (q37.3) in four unrelated individuals. *Am. J. Med. Genet.* 58, 1-7.
150. Williams, S.R., Aldred, M.A., Der Kaloustian, V.M., et al. (2010). Haploinsufficiency of HDAC4 causes brachydactyly mental retardation syndrome, with brachydactyly type E, developmental delays, and behavioral problems. *Am. J. Hum. Genet.* 87, 219-228.
151. Johnson, D., Kan, S.H., Oldridge, M., et al (2003). Missense mutations in the homeodomain of HOX13 are associated with brachydactyly type D and E. *Am. J. Hum. Genet.* 72, 984-997.
152. Maass, P.G., Wirth, J., Aydin, A., et al. (2010). A cis-regulatory site downregulates PTHLH in translocation t(8;12)(q13;p11.2) and leads to brachydactyly type E. *Hum. Mol. Genet.* 19, 848-860.
153. Klopocki, E., Hennig, B.P., and Dathe, K., et al. (2010). Deletion and point mutations of PTHLH cause brachydactyly type E. *Am. J. Hum. Genet.* 86, 434-439.
154. Maass PG, Rump A, Schulz H, et al. (2012). A misplaced lncRNA causes brachydactyly in humans. *J Clin Invest.* 122(11):3990-4002.
155. Maass PG, Aydin A, Luft FC, et al. (2015). PDE3A mutations cause autosomal dominant hypertension with brachydactyly. *Nat Genet.* 47(6):647-653.
156. Izzi B, Francois I, Labarque V, et al. (2012). Methylation defect in imprinted genes detected in patients with an Albright's hereditary osteodystrophy like phenotype and platelet Gs hypofunction. *PLoS One.* 7(6):e38579.
157. Farfel, Z., Brickman, A. S., Kaslow, H. R., Brothers, V. M., and Bourne, H. R. (1980). Defect of receptor-cyclase coupling protein in pseudohypoparathyroidism. *N. Engl. J. Med.* 303, 237-242.
158. Levine, M. A., Downs, R. W., Jr., Singer, M., Marx, S. J., Aurbach, G. D., and Spiegel, A. M. (1980). Deficient activity of guanine nucleotide regulatory protein in erythrocytes from patients with pseudohypoparathyroidism. *Biochem. Biophys. Res. Commun.* 94, 1319-1324.
159. Levine, M. A., Jap, T. S., Maueth, R. S., Downs, R. W., and Spiegel, A. M. (1986). Activity of the stimulatory guanine nucleotide-binding protein is reduced in erythrocytes from patients with pseudohypoparathyroidism and pseudopseudohypoparathyroidism: biochemical, endocrine, and genetic analysis of Albright's hereditary osteodystrophy in six kindreds. *J. Clin. Endocrinol. Metab* 62, 497-502.

160. Beaudet, A. L. (2004). Complex imprinting. *Nat. Genet.* 36:793-5.
161. Yu, S., Yu, D., Lee, E., et al. (1998). Variable and tissue-specific hormone resistance in heterotrimeric Gs protein α -subunit (Gsa) knockout mice is due to tissue-specific imprinting of the Gsa gene. *Proc. Natl. Acad. Sci. USA* 95, 8715-8720.
162. Hayward, B., Barlier, A., Korbonits, M., et al. (2001). Imprinting of the G(s)alpha gene GNAS1 in the pathogenesis of acromegaly. *J. Clin. Invest.* 107, R31-36
163. Mantovani, G., Ballare, E., Giammona, E., Beck-Peccoz, P., and Spada, A. (2002). The Gsalph gene: predominant maternal origin of transcription in human thyroid gland and gonads. *J. Clin. Endocrinol. Metab.* 87, 4736-4740.
164. Germain-Lee, E.L., Ding, C.L., Deng, Z., et al. (2002). Paternal imprinting of Galpha(s) in the human thyroid as the basis of TSH resistance in pseudohypoparathyroidism type 1a. *Biochem. Biophys. Res. Commun.* 296, 67-72.
165. Davies, S. J., and Hughes, H. E. (1993). Imprinting in Albright's hereditary osteodystrophy. *J Med. Genet.* 30, 101-103.
166. Liu, J., Litman, D., Rosenberg, M. J., Yu, S., Biesecker, L. G., and Weinstein, L. S. (2000). A GNAS imprinting defect in pseudohypoparathyroidism type 1B. *J Clin. Invest* 106, 1167-1174.
167. Zheng, H., Radeva, G., McCann, J. A., Hendy, G. N., and Goodyer, C. G. (2001). Galphas transcripts are biallelically expressed in the human kidney cortex: implications for pseudohypoparathyroidism type 1b. *J Clin. Endocrinol. Metab* 86, 4627-4629.
168. Weinstein, L. S., Yu, S., and Ecelbarger, C. A. (2000). Variable imprinting of the heterotrimeric G protein G(s) alpha-subunit within different segments of the nephron. *Am. J Physiol Renal Physiol* 278, F507-F514.
169. Weinstein, L. S., Yu, S., Warner, D. R., and Liu, J. (2001). Endocrine manifestations of stimulatory G-protein alpha-subunit mutations and the role of genomic imprinting. *Endocr. Rev.* 22, 675-705.
170. Weinstein, L. S., Liu, J., Sakamoto, A., Xie, T., and Chen, M. (2004). Minireview: GNAS: normal and abnormal functions. *Endocrinology.* 145, 5459-5464.
171. Turan S, Fernandez-Rebollo E, Aydin C, et al. (2014). Postnatal establishment of allelic Gas silencing as a plausible explanation for delayed onset of parathyroid hormone resistance owing to heterozygous Gas disruption. *J Bone Miner Res.* 29(3):749-760.
172. Levine, M. A., Ahn, T. G., Klupt, S. F., et al. (1988). Genetic deficiency of the alpha subunit of the guanine nucleotide- binding protein Gs as the molecular basis for Albright hereditary osteodystrophy. *Proc. Natl. Acad. Sci. U. S. A* 85, 617-621.
173. Patten, J. L., Johns, D. R., Valle, D., et al. (1990). Mutation in the gene encoding the stimulatory G protein of adenylate cyclase in Albright's hereditary osteodystrophy. *N. Engl. J Med.* 322, 1412-1419.

174. Weinstein, L. S., Gejman, P. V., Friedman, E., et al. (1990). Mutations of the Gs alpha-subunit gene in Albright hereditary osteodystrophy detected by denaturing gradient gel electrophoresis. *Proc. Natl. Acad. Sci. U. S. A* 87, 8287-8290.
175. Aldred, M. A., and Trembath, R. C. (2000). Activating and inactivating mutations in the human GNAS gene. *Hum. Mutat.* 16, 183-189.
176. Ahrens, W., Hiort, O., Staedt, P., Kirschner, T., Marschke, C., and Kruse, K. (2001). Analysis of the GNAS gene in Albright's hereditary osteodystrophy. *J Clin. Endocrinol. Metab* 86, 4630-4634.
177. Aldred, M. A., Bagshaw, R. J., Macdermot, K., et al. (2000). Germline mosaicism for a GNAS mutation and Albright hereditary osteodystrophy. *J Med. Genet.* 37, E35.
178. De Nanclares, G.P., Fernandez-Rebollo, E., Santin, I., et al. (2007) Epigenetic defects of GNAS in patients with pseudohypoparathyroidism and mild features of Albright's hereditary osteodystrophy. *J Clin Endocrinol Metab* 92:2370-2373.
179. Fernandez-Rebollo, E., Garcia-Cuartero, B., Garin, I., L., et al. (2010) Intragenic GNAS deletion involving exon A/B in pseudohypoparathyroidism type 1A resulting in an apparent loss of exon A/B methylation: potential for misdiagnosis of pseudohypoparathyroidism type 1B. *J Clin Endocrinol Metab* 95:765-771.
180. Namnoum, A. B., Merriam, G. R., Moses, A. M., and Levine, M. A. (1998). Reproductive dysfunction in women with Albright's hereditary osteodystrophy. *J Clin. Endocrinol. Metab* 83, 824-829.
181. McIlroy, J., Dryburgh, F., Hinnie, J., Dargie, R., and Al Rawi, A. (1999). Oestrogen and calcium homeostasis in women with hypoparathyroidism. *BMJ* 319, 1252-1253.
182. Breslau, N. A., and Zerwekh, J. E. (1986). Relationship of estrogen and pregnancy to calcium homeostasis in pseudohypoparathyroidism. *J Clin. Endocrinol. Metab* 62, 45-51.
183. Germain-Lee, E. L., Groman, J., Crane, J. L., Jan de Beur, S. M., and Levine, M. A. (2003). Growth hormone deficiency in pseudohypoparathyroidism type 1a: another manifestation of multihormone resistance. *J. Clin. Endocrinol. Metab.* 88, 4059-4069.
184. Mantovani, G., Maghnie, M., Weber, G., et al. (2003). Growth hormone-releasing hormone resistance in pseudohypoparathyroidism type 1a: new evidence for imprinting of the Gs alpha gene. *J. Clin. Endocrinol. Metab.* 88, 4070-4074.
185. Mantovani, G., Ferrante, E., Giavoli, C. et al. (2010) Recombinant human GH replacement therapy in children with pseudohypoparathyroidism type 1a: first study on the effect on growth. *J. Clin. Endocrinol. Metab.* 95, 5011-5017
186. Liu, J., Erlichman, B., and Weinstein, L.S. (2003) The stimulatory G protein α -subunit Gs α is imprinted in human thyroid glands: implications for thyroid function in pseudohypoparathyroidism types 1A and 1B. *J. Clin. Endocrinol. Metabol.* 88, 4336-4341

187. Long, D.N., McGuire, S., Levine, M.A., Weinstein, L.S., and Germain-Lee, E.L. (2007). Body mass index differences in pseudohypoparathyroidism type 1a versus pseudopseudohypoparathyroidism may implicate paternal imprinting of *Galpha(s)* in the development of human obesity. *J. Clin. Endocrinol. Metab.* 92, 1073-1079.
188. Mantovani, G., Bondioni, S., Locatelli, M., et al. (2004) Biallelic expression of the *Galpha* gene in human bone and adipose tissue. *J. Clin. Endocrinol. Metab.* 89:6316-6319.
189. Chen, M., Wang, J., Dickerson, K.E., et al. (2009) Central nervous system imprinting of the G protein *G(s)alpha* and its role in metabolic regulation. *Cell Metab.* 9, 548-555.
190. Mouallem, M., Shaharabany, M., Weintrob, N., et al. (2008). Cognitive impairment is prevalent in pseudohypoparathyroidism type 1a, but not in pseudopseudohypoparathyroidism: possible cerebral imprinting of *Galpha*. *Clin. Endocrinol. (Oxf)* 68, 233-239.
191. Murray, T. M., Rao, L. G., Wong, M. M., et al. (1993). Pseudohypoparathyroidism with osteitis fibrosa cystica: direct demonstration of skeletal responsiveness to parathyroid hormone in cells cultured from bone. *J. Bone Miner. Res.* 8, 83-91.
192. Ish-Shalom, S., Rao, L. G., Levine, M. A., et al. (1996). Normal parathyroid hormone responsiveness of bone-derived cells from a patient with pseudohypoparathyroidism. *J. Bone Miner. Res.* 11, 8-14.
193. Laspa, E., Bastepe, M., Jüppner, H., and Tsatsoulis, A. (2004). Phenotypic and molecular genetic aspects of pseudohypoparathyroidism type 1b in a Greek kindred: evidence for enhanced uric acid excretion due to parathyroid hormone resistance. *J Clin Endocrinol Metab.* 89, 5942-5947.
194. Unluturk, U., Harmanci, A., Babaoglu, M., et al. (2008). Molecular diagnosis and clinical characterization of pseudohypoparathyroidism type-1b in a patient with mild Albright's hereditary osteodystrophy-like features, epileptic seizures, and defective renal handling of uric acid. *Am. J. Med. Sci.* 336, 84-90.
195. Molinaro A, Tiosano D, Takatani R, et al. (2015). TSH elevations as the first laboratory evidence for pseudohypoparathyroidism type 1b (PHP-1b). *J Bone Miner Res.* 30(5):906-912.
196. Romanet P, Osei L, Netchine I, et al. (2015). Case report of *GNAS* epigenetic defect revealed by a congenital hypothyroidism. *Pediatrics.* 135(4):e1079-1083.
197. Sano S, Iwata H, Matsubara K, Fukami M, Kagami M, Ogata T. (2015). Growth hormone deficiency in monozygotic twins with autosomal dominant pseudohypoparathyroidism type 1b. *Endocr J.* 62(6):523-529.
198. Schipani, E., Weinstein, L. S., Bergwitz, C., et al. (1995). Pseudohypoparathyroidism type 1b is not caused by mutations in the coding exons of the human parathyroid hormone (PTH)/PTH-related peptide receptor gene. *J Clin. Endocrinol. Metab.* 80, 1611-1621.
199. Fukumoto, S., Suzawa, M., Takeuchi, Y., et al. (1996). Absence of mutations in parathyroid hormone (PTH)/PTH-related protein receptor complementary deoxyribonucleic acid in patients with pseudohypoparathyroidism type 1b. *J Clin. Endocrinol. Metab.* 81, 2554-2558.

200. Bettoun, J. D., Minagawa, M., Kwan, M. Y., et al. (1997). Cloning and characterization of the promoter regions of the human parathyroid hormone (PTH)/PTH-related peptide receptor gene: analysis of deoxyribonucleic acid from normal subjects and patients with pseudohypoparathyroidism type 1b. *J Clin. Endocrinol. Metab.* 82, 1031-1040.
201. Juppner, H., Schipani, E., Bastepe, M., M., et al. (1998). The gene responsible for pseudohypoparathyroidism type 1b is paternally imprinted and maps in four unrelated kindreds to chromosome 20q13.3. *Proc. Natl. Acad. Sci. U. S. A* 95, 11798-11803.
202. Jan de Beur, S. M., Ding, C. L., LaBuda, M. C., Usdin, T. B., and Levine, M. A. (2000). Pseudohypoparathyroidism 1b: exclusion of parathyroid hormone and its receptors as candidate disease genes. *J Clin. Endocrinol. Metab* 85, 2239-2246.
203. Bastepe, M., Pincus, J. E., Sugimoto, T., et al. (2001). Positional dissociation between the genetic mutation responsible for pseudohypoparathyroidism type 1b and the associated methylation defect at exon A/B: evidence for a long-range regulatory element within the imprinted GNAS locus. *Hum. Mol. Genet.* 10, 1231-1241.
204. Bastepe, M., Frohlich, L. F., Hendy, G. N., et al. (2003). Autosomal dominant pseudohypoparathyroidism type 1b is associated with a heterozygous microdeletion that likely disrupts a putative imprinting control element of GNAS. *J. Clin. Invest.* 112, 1255-1263.
205. Linglart, A., Gensure, R.C., Olney, R.C., Jüppner, H., and Bastepe, M. (2005). A novel STX16 deletion in autosomal dominant pseudohypoparathyroidism type 1b redefines the boundaries of a cis-acting imprinting control element of GNAS. *Am. J. Hum. Genet.* 76, 804-814.
206. Elli FM, de Sanctis L, Peverelli E, et al. (2014). Autosomal dominant pseudohypoparathyroidism type 1b: a novel inherited deletion ablating STX16 causes loss of imprinting at the A/B DMR. *J Clin Endocrinol Metab.* 99(4):E724-E728.
207. Richard N, Abeguilé G, Coudray N, et al. (2012). A new deletion ablating NESP55 causes loss of maternal imprint of A/B GNAS and autosomal dominant pseudohypoparathyroidism type 1b. *J Clin Endocrinol Metab.* 97(5):E863-E867.
208. Bastepe, M., Frohlich, L. F., Linglart, A., et al. (2005). Deletion of the NESP55 differentially methylated region causes loss of maternal GNAS imprints and pseudohypoparathyroidism type 1b. *Nat. Genet.* 37, 25-27.
209. Chillambhi, S., Turan, S., Hwang, D.Y., Chen, H.C., Jüppner, H., and Bastepe, M. (2010). Deletion of the noncoding GNAS antisense transcript causes pseudohypoparathyroidism type 1b and biparental defects of GNAS methylation in cis. *J. Clin. Endocrinol. Metab.* 95, 3993-4002.
210. Nakamura A, Hamaguchi E, Horikawa R, et al. (2016). Complex Genomic Rearrangement Within the GNAS Region Associated With Familial Pseudohypoparathyroidism Type 1b. *J Clin Endocrinol Metab.* 101(7):2623-2627.

211. Perez-Nanclares G, Velayos T, Vela A, Muñoz-Torres M, and Castaño L. (2015). Pseudohypoparathyroidism type 1b associated with novel duplications in the GNAS locus. *PLoS One*. 10(2):e0117691.
212. Bastepe, M., Lane, A. H., and Juppner, H. (2001). Paternal uniparental isodisomy of chromosome 20q-- and the resulting changes in GNAS methylation--as a plausible cause of pseudohypoparathyroidism. *Am. J Hum. Genet.* 68, 1283-1289.
213. Lecumberri, B., Fernandez-Rebollo, E., Sentchordi, L., et al. (2010). Coexistence of two different pseudohypoparathyroidism subtypes (1a and 1b) in the same kindred with independent Gs{alpha} coding mutations and GNAS imprinting defects. *J. Med. Genet.* 47, 276-280.
214. Fernandez-Rebollo, E., Lecumberri, B., Garin, I., et al. (2010). New mechanisms involved in paternal 20q disomy associated with pseudohypoparathyroidism. *Eur. J. Endocrinol.* 163, 953-962.
215. Bastepe, M., Altug-Teber, O., Agarwal, C., Oberfield, S.E., Bonin, M., and Jüppner, H. (2011). Paternal uniparental isodisomy of the entire chromosome 20 as a molecular cause of pseudohypoparathyroidism type 1b (PHP-1b). *Bone* 48, 659-662.
216. Takatani R, Minagawa M, Molinaro A, et al. (2015). Similar frequency of paternal uniparental disomy involving chromosome 20q (patUPD20q) in Japanese and Caucasian patients affected by sporadic pseudohypoparathyroidism type 1b (sporPHP1B). *Bone*. 79:15-20.
217. Takatani R, Molinaro A, Grigelioniene G, et al. (2016). Analysis of multiple families with single individuals affected by pseudohypoparathyroidism type 1b (PHP1B) reveals only one novel maternally inherited GNAS deletion. *J Bone Miner Res.* 31(4):796-805.
218. Maupetit-Méhouas S, Azzi S, Steunou V, et al. (2013). Simultaneous hyper- and hypomethylation at imprinted loci in a subset of patients with GNAS epimutations underlies a complex and different mechanism of multilocus methylation defect in pseudohypoparathyroidism type 1b. *Hum Mutat.* 34(8):1172-1180.
219. Blik J, Verde G, Callaway J, et al. (2009). Hypomethylation at multiple maternally methylated imprinted regions including PLAGL1 and GNAS loci in Beckwith-Wiedemann syndrome. *Eur J Hum Genet.* 17(5):611-619.
220. Mackay DJ, Callaway JL, Marks SM, et al. (2008). Hypomethylation of multiple imprinted loci in individuals with transient neonatal diabetes is associated with mutations in ZFP57. *Nat Genet.* 40(8):949-951.
221. Bakker B, Sonneveld LJ, Woltering MC, Bikker H, and Kant SG. (2015). A girl with Beckwith-Wiedemann Syndrome and Pseudohypoparathyroidism Type 1B due to multiple imprinting defects. *J Clin Endocrinol Metab.* 100(11):3963-3966.
222. Rochtus A, Martin-Trujillo A, Izzi B, et al. (2016). Genome-wide DNA methylation analysis of pseudohypoparathyroidism patients with GNAS imprinting defects. *Clin Epigenetics.* 8:10.

224. Linglart, A., Bastepe, M., and Jüppner, H. (2007). Similar clinical and laboratory findings in patients with symptomatic autosomal dominant and sporadic pseudohypoparathyroidism type Ib despite different epigenetic changes at the GNAS locus. *Clin. Endocrinol. (Oxf)* 67, 822-831.
225. Bréhin AC, Colson C, Maupetit-Méhouas S, et al. (2015). Loss of methylation at GNAS exon A/B is associated with increased intrauterine growth. *J Clin Endocrinol Metab.* 2015 100(4):E623-E631.
226. De Nanclares, G.P., Fernandez-Rebollo, E., Santin, L., et al. (2007). Epigenetic defects of GNAS in patients with pseudohypoparathyroidism and mild features of Albright's hereditary osteodystrophy. *J. Clin. Endocrinol. Metab.* 92, 2370-2373.
227. Mariot, V., Maupetit-Mehouas, S., Sinding, C., Kottler, M.L., and Linglart, A. (2008) A maternal epimutation of GNAS leads to Albright osteodystrophy and parathyroid hormone resistance. *J. Clin. Endocrinol. Metab.* 93, 661-665.
228. Mantovani, G., de Sanctis, L., Barbieri, A.M., et al. (2010). Pseudohypoparathyroidism and GNAS epigenetic defects: clinical evaluation of Albright hereditary osteodystrophy and molecular analysis in 40 patients. *J. Clin. Endocrinol. Metab.* 95, 651-658.
229. Wu, W. I., Schwindinger, W. F., Aparicio, L. F., and Levine, M. A. (2001). Selective resistance to parathyroid hormone caused by a novel uncoupling mutation in the carboxyl terminus of G alpha(s). A cause of pseudohypoparathyroidism type Ib. *J. Biol. Chem.* 276, 165-171.
230. Linglart, A., Carel, J. C., Garabedian, M., Le, T., Mallet, E., and Kottler, M. L. (2002). GNAS lesions in pseudohypoparathyroidism Ia and Ic: genotype phenotype relationship and evidence of the maternal transmission of the hormonal resistance. *J. Clin. Endocrinol. Metab.* 87, 189-197.
231. Linglart A., Mahon, M.J., Kerachian, M.A., et al. (2006). Coding GNAS mutations leading to hormone resistance impair in vitro agonist- and cholera toxin-induced adenosine cyclic 3',5'-monophosphate formation mediated by human XLas. *Endocrinology* 147, 2253-2262.
232. Linglart, A., Carel, J.C., Garabedian, M., Le, T., Mallet, E., and Kottler, M.L. (2002). GNAS1 lesions in pseudohypoparathyroidism Ia and Ic: Genotype phenotype relationship and evidence of the maternal transmission of the hormonal resistance. *J. Clin. Endocrinol. Metab.* 87, 189-97
233. Thiele, S., De Sanctis, L., Werner, R., et al. (2011). Functional characterization of GNAS mutations found in patients with pseudohypoparathyroidism type 1c defines a new subgroup of pseudohypoparathyroidism affecting selectively Gsalpha-receptor interaction. *Hum. Mutat.* 32, 653-660
234. Drezner, M., Neelon, F. A., and Lebovitz, H. E. (1973). Pseudohypoparathyroidism type II: a possible defect in the reception of the cyclic AMP signal. *N. Engl. J. Med.* 289, 1056-1060.
235. Linglart, A., Menguy, C., Couvineau, A., et al. (2011). Recurrent PRKAR1A mutation in acrodysostosis with hormone resistance. *N. Engl. J. Med.* 364, 2218-2226.
236. Lee H, Graham JM Jr, Rimoin DL, et al. et al. (2012). Exome sequencing identifies PDE4D mutations in acrodysostosis. *Am J Hum Genet.* 90(4):746-751.

237. Michot C, Le Goff C, Goldenberg A, et al. (2012). Exome sequencing identifies PDE4D mutations as another cause of acrodysostosis. *Am J Hum Genet.* 90(4):740-745.
238. Weinstein, L.S. (2000). Fibrous dysplasia and the McCune-Albright syndrome. In "The Genetics of Osteoporosis and Metabolic Bone Disease", (Econs, M. J.ed.), pp.163-177. Humana Press, Totawa NJ
239. Weinstein, L.S., and Collins, M.T. (2008). Diseases resulting from defects in the G protein Gs α . In Bilezikian, J.P., Martin, T.J., Raisz, L.G. (eds): Principles of Bone Biology. 3rd ed. Academic Press, San Diego, pp 1453-1477.
240. Cole, D.E.C., Fraser, F.C., Glorieux, F.H., et al. (1983). Panostotic fibrous dysplasia: a congenital disorder of bone with unusual facial appearance, bone fragility, hyperphosphatasemia, and hypophosphatemia. *Am. J. Med. Genet.* 14, 725-735.
241. Cole, D.E.C., and Whyte, M.P. (1997). Hyperphosphatasia Syndromes. In "Studies in Stomatology and Craniofacial Biology", (Cohen, M. M., Jr., and Baum, B. J.eds.), pp.245-272. IOS Press, Amsterdam.
242. Landis, C.A., Masters, S.B., Spada, A., Pace, A.M., Bourne, H.R., and Vallar, L. (1989). GTPase inhibiting mutations activate the alpha chain of Gs and stimulate adenylyl cyclase in human pituitary tumours. *Nature* 340, 692-696.
243. Iiri, T., Herzmark, P., Nakamoto, J. M., Van Dop, C., and Bourne, H. R. (1994). Rapid GDP release from Gs alpha in patients with gain and loss of endocrine function. *Nature* 371, 164-168.
244. Nakamoto JM, Zimmerman D, Jones EA, et al. (1996). Concurrent hormone resistance (pseudohypoparathyroidism type Ia) and hormone independence (testotoxicosis) caused by a unique mutation in the G alpha s gene. *Biochem Mol Med.* 58(1):18-24.
245. Makita, N., Sato, J., Rondard, P., et al. (2007). Human G(salpa) mutant causes pseudohypoparathyroidism type Ia/neonatal diarrhea, a potential cell-specific role of the palmitoylation cycle. *Proc. Natl. Acad. Sci. USA* 104, 17424-17429.
246. Eddy, M. C., De Beur, S. M., Yandow, S. M., et al. (2000). Deficiency of the alpha-subunit of the stimulatory G protein and severe extraskelatal ossification. *J Bone Miner. Res.* 15, 2074-2083.
247. Kaplan, F. S., and Shore, E. M. (2000). Progressive osseous heteroplasia. *J. Bone Miner. Res.* 15, 2084-2094.
248. Yeh, G. L., Mathur, S., Wivel, A., et al. (2000). GNAS mutation and Cbfa1 misexpression in a child with severe congenital platelike osteoma cutis. *J Bone Miner. Res.* 15, 2063-2073.
249. Shore, E.M., and Kaplan, F.S. (2010). Inherited human diseases of heterotopic bone formation. *Nat. Rev. Rheumatol.* 6, 518-527.
250. Adegbite, N.S., Xu, M., Kaplan, F.S., Shore, E.M., and Pignolo, R.J. (2008). Diagnostic and mutational spectrum of progressive osseous heteroplasia (POH) and other forms of GNAS-based heterotopic ossification. *Am. J. Med. Genet. A.* 146A, 1788-1796.

251. Shore, E.M., Ahn, J., Jan de Beur, S. et al. (2002). Paternally inherited inactivating mutations of the *GNAS1* gene in progressive osseous heteroplasia. *N. Engl. J. Med.* 346, 99-106.
252. Ahmed SF, Barr DG, and Bonthron DT. (2002). *GNAS1* mutations and progressive osseous heteroplasia. *N Engl J Med.* 346(21):1669-1671.
253. Cairns DM, Pignolo RJ, Uchimura T, et al. (2013). Somitic disruption of *GNAS* in chick embryos mimics progressive osseous heteroplasia. *J Clin Invest.* 123(8):3624-3633.
254. Gardella TJ, and Vilardaga JP. (2015). International Union of Basic and Clinical Pharmacology. XCIII. The parathyroid hormone receptors--family B G protein-coupled receptors. *Pharmacol Rev.* 67(2):310-37.
255. Scillitani, A., Jang, C., Wong, B.Y-L., Hendy, G.N., and Cole, D.E.C. (2006). A functional polymorphism in *PTH1R* promoter region is associated with adult height and BMD measured at the femoral neck in a large cohort of young caucasian women. *Hum. Genet.* 119, 416-421.
256. Jüppner, H., Schipani, E., and Silve, C. (2008) Genetic disorders caused by PTH/PTHrP receptor mutations. In Bilezikian, J.P., Martin, T.J., Raisz, L.G. (eds): *Principles of Bone Biology*. 3rd ed. Academic Press, San Diego, pp 1431-1452.
257. Blomstrand, S., Claesson, I., and Save-Soderbergh, J. (1985). A case of lethal congenital dwarfism with accelerated skeletal maturation. *Pediatr. Radiol.* 15, 141-143.
258. Loshkajian, A., Roume, J., Stanescu, V., Delezoide, A.-L., Stampf, F., and Maroteaux, P. (1997). Familial Blomstrand chondrodysplasia with advanced skeletal maturation: Further delineation. *Am. J. Med. Genet.* 71, 283-288.
259. Zhang, P., Jobert, A. S., Couvineau, A., and Silve, C. (1998). A homozygous inactivating mutation in the parathyroid hormone/parathyroid hormone-related peptide receptor causing Blomstrand chondrodysplasia. *J Clin. Endocrinol. Metab* 83, 3365-3368.
260. Karperien, M., van der Harten, H. J., van Schooten, R., et al. (1999). A frame-shift mutation in the type I parathyroid hormone (PTH)/PTH-related peptide receptor causing Blomstrand lethal osteochondrodysplasia. *J Clin. Endocrinol. Metab.* 84, 3713-3720.
261. Karaplis, A.C., He, B., Nguyen M.T., et al. (1998). Inactivating mutation in the human parathyroid hormone receptor type 1 gene in Blomstrand chondrodysplasia. *Endocrinology* 139, 5255-5258.
262. Oostra, R.J., van der Harten, J.J., Rijnders, W.P., Scott, R.J., Young, M.P., and Trump, D. (2000). Blomstrand osteochondrodysplasia: three novel cases and histological evidence for heterogeneity. *Virchows Arch.* 436, 28-35.
263. Hoogendam, J., Farih-Sips, H., Wynaendts, L.C., Lowik, C.W., Wit, J.M., and Karperien, M. (2007). Novel mutations in the parathyroid hormone (PTH)/PTH-related peptide receptor type 1 causing Blomstrand osteochondrodysplasia types I and II. *J. Clin. Endocrinol. Metab.* 92, 1088-1095.

264. Duchatelet, S., Ostergaard, E., Cores, D., Lemainque, A., and Julier, C., (2005). Recessive mutations in PTHR1 cause contrasting skeletal dysplasias in Eiken and Blomstrand syndromes. *Hum. Mol. Genet.* 14, 1-5.
265. Hopyan, S., Gokgoz, N., Poon, R., et al. (2002). A mutant PTH/PTHrP type I receptor in enchondromatosis. *Nat. Genet.* 30, 306-10.
266. Couvineau, A., Wouters, W., Bertrand, G., et al. (2008). PTHR1 mutations associated with Oilier disease result in receptor loss of function. *Hum. Mol. Genet.* 17, 2766-2775.
267. Rozeman, L. B., Sangiorgi, L., Briaire-de Bruijn, I. H., et al. (2004). Enchondromatosis (Ollier disease, Maffucci syndrome) is not caused by the PTHR1 mutation p.R150C. *Hum. Mutat.* 24, 466-473.
268. Collinson, M., Leonard, S.J., Charlton, J., et al. (2010). Symmetrical enchondromatosis is associated with duplication of 12p11.23 to 12p11.22 including *PTH1R*. *Am. J. Med. Genet. Part A.* 152A, 3124-3128.
269. Decker, E., Stellzig-Eisenhauer, A., Fiebig, B.S., et al. (2008) PTHR1 loss-of function mutations in familial, non-syndromic primary failure of tooth eruption. *Am. J. Hum. Genet.* 83, 781-786.
270. Frazier-Bowers, S.A., Simmons, D., Wright, J.T., Proffit, W.R., and Ackerman, J.L. (2010). Primary failure of eruption and *PTH1R*: The importance of a genetic diagnosis for orthodontic treatment planning. *Am. J. Orthod. Dentofacial. Orthop.* 137, 160.e1-160.e7.
271. Risom L, Christoffersen L, Daugaard-Jensen J, et al. (2013). Identification of six novel PTH1R mutations in families with a history of primary failure of tooth eruption. *PLoS One.* 8(9):e74601.
272. Frazier-Bowers SA, Hendricks HM, et al. (2014). Novel mutations in PTH1R associated with primary failure of eruption and osteoarthritis. *J Dent Res.* 93(2):134-139.
273. Yamaguchi T, Hosomichi K, Narita A, et al. (2011). Exome resequencing combined with linkage analysis identifies novel PTH1R variants in primary failure of tooth eruption in Japanese. *J Bone Miner Res.* 26(7):1655-1661.
274. Konrad, M., Schlingmann, K. P., and Gudermann, T. (2004). Insights into the molecular nature of magnesium homeostasis. *Am. J. Physiol. Renal Physiol.* 286, F599-F605.
275. Astor MC, Løvås K, Wolff AS, et al. (2015). Hypomagnesemia and functional hypoparathyroidism due to novel mutations in the Mg-channel TRPM6. *Endocr Connect.* 4(4):215-222.
276. Janett S, Camozzi P, Peeters GG, et al. (2015). Hypomagnesemia induced by long-term treatment with proton-pump inhibitors. *Gastroenterol Res Pract.* 2015:951768.
277. Fatuzzo P, Portale G, Scollo V, Zanolli L, Granata A. (2016). Proton pump inhibitors and symptomatic hypomagnesemic hypoparathyroidism. *J Nephrol.* [Epub ahead of print].
278. Brandi ML, Bilezikian JP, Shoback D, (2016). Management of hypoparathyroidism: summary statement and guidelines. *J Clin Endocrinol Metab.* 101(6):2273-2283.
279. Bilezikian JP, Brandi ML, Cusano NE, et al. (2016). Management of hypoparathyroidism: present and future. *J Clin Endocrinol Metab.* 01(6):2313-2324.

280. Mittendorf, E. A., Merlino, J. I., and McHenry, C. R. (2004). Post-parathyroidectomy hypocalcemia: incidence, risk factors, and management. *Am. Surg.* 70, 114-119.
281. Khan, M.I., Waguespack, S.G., and Hu, M.I. (2011). Medical management of postsurgical hypoparathyroidism. *Endocr. Pract.* 17 Suppl 1, 18-25.
282. Arlt, W., Fremerey, C., Callies, F., et al. (2002). Well-being, mood and calcium homeostasis in patients with hypoparathyroidism receiving standard treatment with calcium and vitamin D. *Eur. J. Endocrinol.* 146, 215-222.
283. Okano, K., Furukawa, Y., Morii, H., and Fujita, T. (1982). Comparative efficacy of various vitamin D metabolites in the treatment of various types of hypoparathyroidism. *J Clin. Endocrinol. Metab.* 55, 238-243.
284. Astor MC, Løvås K, Debowska A, et al. (2016). Epidemiology and health-related quality of life in hypoparathyroidism in Norway. *J Clin Endocrinol Metab.* 101(8):3045-5303.
285. Natpara [package insert]. Bedminster, NJ; NPS Pharmaceuticals, Inc; 2015.
286. Jolette, J., Wilker, C.E., Smith, S.Y., et al. (2006). Defining a noncarcinogenic dose of recombinant human parathyroid hormone 1-84 in a 2-year study in Fischer 344 rats. *Toxicol Pathol.* 34, 929-940.
287. Capriani C, Irani D, and Bilezikian JP. (2012). Safety of osteoanabolic therapy: a decade of experience. *J Bone Miner Res.* 27, 2419-2428.
288. Winer KK, Yanovski JA, Cutler GB Jr. Synthetic human parathyroid hormone 1-34 vs calcitriol and calcium in the treatment of hypoparathyroidism. *JAMA.* 1996 276(8):631-636.
289. Winer KK, Yanovski JA, Sarani B, and Cutler GB Jr. (1998). A randomized, cross-over trial of once-daily versus twice-daily parathyroid hormone 1-34 in treatment of hypoparathyroidism. *J Clin Endocrinol Metab.* 1998 83(10):3480-3486.
290. Winer, K. K., Ko, C. W., Reynolds, J. C., et al. (2003). Long-term treatment of hypoparathyroidism: a randomized controlled study comparing parathyroid hormone-(1-34) versus calcitriol and calcium. *J. Clin. Endocrinol. Metab.* 88, 4214-4220.
291. Winer, K.K., Sinaii, N., Peterson, D., Sainz, B. Jr., and Cutler, G.B. Jr. (2008). Effects of once versus twice-daily parathyroid hormone 1-34 therapy in children with hypoparathyroidism. *J. Clin. Endocrinol. Metab.* 93, 3389-3395.
292. Winer, K.K., Sinaii, N., Reynolds, J., Peterson, D., Dowdy, K., and Cutler, G.B. Jr. (2010). Long-term treatment of 12 children with chronic hypoparathyroidism: a randomized trial comparing synthetic human parathyroid hormone 1-34 versus calcitriol and calcium. *J. Clin. Endocrinol. Metab.* 95, 2680-2688.
293. Winer KK, Zhang B, Shrader JA, et al. (2012). Synthetic human parathyroid hormone 1-34 replacement therapy: a randomized crossover trial comparing pump versus injections in the treatment of chronic hypoparathyroidism. *J Clin Endocrinol Metab.* 97(2):391-9.
294. Winer KK, Fulton KA, Albert PS, Cutler GB Jr. (2014). Effects of pump versus twice-daily injection delivery of synthetic parathyroid hormone 1-34 in children with severe congenital hypoparathyroidism. *J Pediatr.* 165(3):556-563.e1.

295. Santonati A, Palermo A, Maddaloni E, et al. (2015). PTH(1-34) for surgical hypoparathyroidism: a prospective, open-label investigation of efficacy and quality of life. *J Clin Endocrinol Metab.* 100(9):3590-3597.
296. Mittelman, S.D., Hendy, G.N., Fefferman, R.A., et al. (2006). A hypocalcemic child with a novel activating mutation of the calcium-sensing receptor gene: successful treatment with recombinant human parathyroid hormone. *J. Clin. Endocrinol. Metab.* 91, 2474-2479.
297. Theman, T.A., Collins, M.T., Dempster, D.W., et al. (2009). PTH(1-34) replacement therapy in a child with hypoparathyroidism caused by a sporadic calcium receptor mutation. *J. Bone Miner. Res.* 24, 964-973.
298. Clarke BL, Kay Berg J, Fox J, Cyran JA, Lagast H. (2014). Pharmacokinetics and pharmacodynamics of subcutaneous recombinant parathyroid hormone (1-84) in patients with hypoparathyroidism: an open-label, single-dose, phase I study. *Clin Ther.* 36(5):722-736.
299. Sikjaer T, Amstrup AK, Rolighed L, et al. (2013). PTH(1-84) replacement therapy in hypoparathyroidism: a randomized controlled trial on pharmacokinetic and dynamic effects after 6 months of treatment. *J Bone Miner Res.* 28(10):2232-2243.
300. Rubin MR, Sliney J Jr, McMahon DJ, Silverberg SJ, and Bilezikian JP. (2010). Therapy of hyperparathyroidism with intact parathyroid hormone. *Osteoporos Int* 21, 1927-1934.
301. Sikjaer T, Rejnmark L, Rolighed L, Heickendorff L, and Mosenkilde L. (2011). The effect of adding PTH(1-84) to conventional treatment of hypoparathyroidism: a randomized placebo-controlled study. *J Bone Miner Res.* 26, 2358-2370.
302. Cusano NE, Rubin MR, McMahon DJ, et al. (2013). Therapy of hypoparathyroidism with PTH(1-84): a prospective four-year investigation of efficacy and safety. *J Clin Endocrinol Metab.* 98(1):137-144.
303. Cusano NE, Rubin MR, McMahon DJ, et al. (2014). PTH(1-84) is associated with improved quality of life in hypoparathyroidism through 5 years of therapy. *J Clin Endocrinol Metab.* 99(10):3694-3699
304. Rubin, M., Dempster, D., Sliney, J., E., et al. (2011). PTH(1-84) administration reverses abnormal bone remodeling dynamics and structure in hypoparathyroidism. *J Bone Miner Res.* 26, 2727-2736.
305. Cusano NE, Rubin MR, McMahon DJ, et al. (2013). The effect of PTH(1-84) on quality of life in hypoparathyroidism. *J Clin Endocrinol Metab.* 98(6):2356-2361.
306. Mannstadt M, Clarke BL, Vokes T, et al. (2013). Efficacy and safety of recombinant human parathyroid hormone (1-84) in hypoparathyroidism (REPLACE): a double-blind, placebo-controlled, randomised, phase 3 study. *Lancet Diabetes Endocrinol.* 1(4):275-283
307. Clarke BL, Vokes TJ, Bilezikian JP, Shoback DM, Lagast H, Mannstadt M. (2017). Effects of parathyroid hormone rhPTH(1-84) on phosphate homeostasis and vitamin D metabolism in hypoparathyroidism: REPLACE phase 3 study. *Endocrine.* 255(1):273-282.
308. Bi R, Fan Y, Lauter, et al. (2016). Diphtheria toxin- and GFP-based mouse models of acquired hypoparathyroidism and treatment with a long-acting parathyroid hormone analog. *J Bone Miner Res.* 31(5):975-984.

309. Tamura T, Noda H, Joyashiki E, et al. (2016). Identification of an orally active small-molecule PTHR1 agonist for the treatment of hypoparathyroidism. *Nat Commun.* 7:13384.
310. Nemeth EF, and Goodman WG. (2016). Calcimimetic and calcilytic drugs: feats, flops, and futures. *Calcif Tissue Int.* 98(4):341-358.
311. Dong B, Endo I, Ohnishi Y, et al. (2015). Calcilytic ameliorates abnormalities of mutant calcium-sensing receptor (CaSR) knock-in mice mimicking autosomal dominant hypocalcemia (ADH). *J Bone Miner Res.* 30(11):1980-1993.
312. Hannan FM, Walls GV, Babinsky VN, et al. (2015). The calcilytic agent NPS 2143 rectifies hypocalcemia in a mouse model with an activating calcium-sensing receptor (CaSR) mutation: relevance to autosomal dominant hypocalcemia type 1 (ADH1). *Endocrinology.* 156(9):3114-3121.
313. Ramnitz, M., Gafni, R., Brillante, B, et al. (2015). Treatment of autosomal dominant hypocalcemia with the calcilytic NPSP795. *J Bone Miner Res* 30 (Suppl 1). LB-SA0002.
314. Hu J, Spiegel AM. (2007). Structure and function of the human calcium-sensing receptor: insights from natural and engineered mutations and allosteric modulators. *J Cell Mol Med.* 11(5):908-922.
315. Letz S, Rus R, Haag C, et al. (2010). Novel activating mutations of the calcium-sensing receptor: the calcilytic NPS-2143 mitigates excessive signal transduction of mutant receptors. *J Clin Endocrinol Metab.* 95(10):E229-E233.
316. Letz S, Haag C, Schulze E, et al. (2014). Amino alcohol- (NPS-2143) and quinazolinone-derived calcilytics (ATF936 and AXT914) differentially mitigate excessive signalling of calcium-sensing receptor mutants causing Bartter syndrome type 5 and autosomal dominant hypocalcemia. *PLoS One.* 9(12):e115178
317. Roszko KL, Bi RD, and Mannstadt M. (2016). Autosomal dominant hypocalcemia (hypoparathyroidism) types 1 and 2. *Front Physiol.* 7:458.
318. Hannan FM, Babinsky VN, and Thakker RV. (2016). Disorders of the calcium-sensing receptor and partner proteins: insights into the molecular basis of calcium homeostasis. *J Mol Endocrinol.* 57(3):R127-R142.
319. Babinsky VN, Hannan FM, Gorvin CM, et al (2016). Allosteric modulation of the calcium-sensing receptor rectifies signaling abnormalities associated with G-protein α -11 mutations causing hypercalcemic and hypocalcemic disorders. *J Biol Chem.* 291(20):10876-10885.
320. Roszko KL, Bi R, Howles S, et al. (2016). CRISPR/Cas9-generated mouse model of autosomal dominant hypocalcemia harboring the activating G protein alpha 11 mutation Arg60Cys and use of calcilytics and a Gq/G α 11-specific Inhibitor. *J Bone Miner Res* 31 (Suppl 1). FR0077/SA0077
321. Xiong XF, Zhang H, Underwood CR, et al. (2016). Total synthesis and structure-activity relationship studies of a series of selective G protein inhibitors. *Nat Chem.* 8(11):1035-1041.
322. Sato, K., Hasegawa, Y., Nakae, J., et al. (2002). Hydrochlorothiazide effectively reduces urinary calcium excretion in two Japanese patients with gain-of-function mutations of the calcium-sensing receptor gene. *J. Clin. Endocrinol. Metab.* 87,3068-3073.