

Optimization for Data Science

Cristian Perez Jensen

April 9, 2024

Contents

1	<i>Risk minimization</i>	1
1.1	<i>Algorithms in data science</i>	1
1.2	<i>Empirical and expected risk</i>	1
1.3	<i>The map of learning</i>	3
2	<i>Theory of convex functions</i>	4
2.1	<i>Mathematical background</i>	4
2.2	<i>Convex sets</i>	4
2.3	<i>Convex functions</i>	5
2.4	<i>Minimizing convex functions</i>	9
2.5	<i>Convex programming</i>	10
3	<i>Gradient descent</i>	14
3.1	<i>Vanilla analysis</i>	14
3.2	<i>Lipschitz convex functions</i>	15
3.3	<i>Smooth functions</i>	16
3.4	<i>Smooth and strongly convex functions</i>	18
4	<i>Projected gradient descent</i>	21
4.1	<i>Lipschitz convex functions</i>	21
4.2	<i>Smooth functions</i>	22
4.3	<i>Smooth and strongly convex functions</i>	24
5	<i>Coordinate descent</i>	25
5.1	<i>Randomized coordinate descent</i>	26
5.2	<i>Importance sampling</i>	27
5.3	<i>Steepest coordinate descent</i>	28
5.4	<i>Greedy coordinate descent</i>	30
6	<i>Nonconvex functions</i>	31
6.1	<i>Trajectory analysis</i>	33

List of symbols

\doteq	Equality by definition
\mathbb{R}	Set of real numbers
$f : A \rightarrow B$	Function f that maps elements of set A to elements of set B
$\mathbf{v} \in \mathbb{R}^n$	n -dimensional vector
$\mathbf{M} \in \mathbb{R}^{m \times n}$	$m \times n$ matrix
\mathbf{M}^\top	Transpose of matrix \mathbf{M}
\mathbf{M}^{-1}	Inverse of matrix \mathbf{M}
$\det(\mathbf{M})$	Determinant of \mathbf{M}
$\frac{d}{dx}f(x)$	Ordinary derivative of $f(x)$ w.r.t. x at point $x \in \mathbb{R}$
$\frac{\partial}{\partial x}f(x)$	Partial derivative of $f(x)$ w.r.t. x at point $x \in \mathbb{R}^n$
$\nabla f(x) \in \mathbb{R}^n$	Gradient of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at point $x \in \mathbb{R}^n$
$Df(x) \in \mathbb{R}^{n \times m}$	Jacobian of $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ at point $x \in \mathbb{R}^n$
$\nabla^2 f(x) \in \mathbb{R}^{n \times n}$	Hessian of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at point $x \in \mathbb{R}^n$

1 Risk minimization

1.1 Algorithms in data science

In classical algorithm theory, an optimization problem solves a well-defined problem. For example, Kruskal's algorithm computes the minimum spanning tree of a graph. In data science, it is not as well-defined. The starting point is a learning problem, and the optimization typically happens on training data. However, even a perfect result may fail to solve the learning problem, which is a failure of the model in which the optimization algorithm was applied, rather than the optimization algorithm itself.

1.2 Empirical and expected risk

Typically, we have a data source \mathcal{X} , equipped with an unknown probability distribution. However, we do have access to independent samples $X_1, \dots, X_n \sim \mathcal{X}$. The goal is to "explain" \mathcal{X} through these samples. More specifically, we have a class \mathcal{H} of hypotheses (possible explanations of \mathcal{X}). The goal is then to select the hypothesis $H \in \mathcal{H}$ that best explains \mathcal{X} , which we measure by a *risk* (or *loss*) function $\ell : \mathcal{H} \times \mathcal{X} \rightarrow \mathbb{R}$.

The expected risk can be computed by

$$\ell(H) \doteq \mathbb{E}_{\mathcal{X}}[\ell(H, X)].$$

The best explanation is the *expected risk minimizer*,

$$H^* = \operatorname{argmin}_{H \in \mathcal{H}} \ell(H).$$

However, since we do not have access to the distribution over \mathcal{X} , we cannot compute $\ell(H)$ or H^* . Thus, we need to compute the *probably approximately correct* (PAC) hypothesis. This means that given $\delta, \epsilon > 0$, we want to produce, with probability $1 - \delta$, a hypothesis $\tilde{H} \in \mathcal{H}$ such that

$$\ell(\tilde{H}) \leq \inf_{H \in \mathcal{H}} \ell(H) + \epsilon,$$

meaning that with high probability, we approximately solve the expected risk minimization problem.

However, we can still not compute this, thus we must use our training data to compute the *empirical risk*,

$$\ell_n(H) = \frac{1}{n} \sum_{i=1}^n \ell(H, X_i).$$

This is a random variable, because it depends on the training data, which are all random variables, distributed according to the probability distribution of \mathcal{X} .

Lemma 1 (Weak law of large numbers). Let $H \in \mathcal{H}$ be a hypothesis. For any $\delta, \epsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that for $n \geq n_0$,

$$|\ell_n(H) - \ell(H)| \leq \epsilon,$$

with probability at least $1 - \delta$.

Given $n \in \mathbb{N}$ and training data X_1, \dots, X_n from \mathcal{X} , we want to produce a hypothesis \tilde{H}_n such that

$$\ell_n(\tilde{H}_n) \leq \inf_{H \in \mathcal{H}} \ell_n(H) + \epsilon.$$

In an ideal world, \tilde{H}_n is also almost the best explanation for the data source \mathcal{X} ,

$$\ell(\tilde{H}_n) \leq \inf_{H \in \mathcal{H}} \ell(H) + \epsilon.$$

Remark. Note that the weak law of large numbers can only be applied to a *fixed* hypothesis, but not to the data-dependent hypothesis \tilde{H}_n . Thus, we are not always in an ideal world scenario.

A sufficient condition for an ideal world scenario is that the weak law of large numbers uniformly holds for all hypotheses with high probability. This leads us to the following theorem.

Theorem 2. Assume that for any $\delta, \epsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that for $n \geq n_0$,

$$\sup_{H \in \mathcal{H}} |\ell_n(H) - \ell(H)| \leq \epsilon,$$

with probability at least $1 - \delta$. Then, for $n \geq n_0$, an approximate empirical risk minimizer \tilde{H}_n is PAC for expected risk minimization, meaning that it satisfies

$$\ell(\tilde{H}_n) \leq \inf_{H \in \mathcal{H}} \ell(H) + 3\epsilon,$$

with probability at least $1 - \delta$.

Proof.

$$\begin{aligned} \ell(\tilde{H}_n) &\leq \ell_n(\tilde{H}_n) + \epsilon \\ &\leq \inf_{H \in \mathcal{H}} \ell_n(H) + 2\epsilon \\ &\leq \inf_{H \in \mathcal{H}} \ell(H) + 3\epsilon, \end{aligned}$$

with probability at least $1 - \delta$. ■

It turns out that the assumption made by ?? holds in many relevant cases, but it is not always true.

Follows from $\sup_{H \in \mathcal{H}} |\ell_n(H) - \ell(H)| \leq \epsilon$.

\tilde{H}_n is an almost best explanation of the training data.

Follows from $\sup_{H \in \mathcal{H}} |\ell_n(H) - \ell(H)| \leq \epsilon$.

In this course, we will not learn how to pick the theory (\mathcal{H} and ℓ), but rather how to solve the optimization problems that arise in empirical risk minimization after the theory has been chosen.

1.3 The map of learning

The map of learning can be seen in Figure 1. It plots the empirical risk ($\ell_n(H_n)$, training data) against the expected risk ($\ell(H_n)$, estimated by a validation set). We can only be in the area denoted by “empirical risk minimization”, because

$$\begin{aligned}\ell_n(\tilde{H}_n) &\leq \inf_{H \in \mathcal{H}} \ell_n + \epsilon \\ &\leq \ell_n(\tilde{H}) + \epsilon \\ &\leq \ell(\tilde{H}) + 2\epsilon \\ &\leq \ell(\tilde{H}_n) + \epsilon\epsilon.\end{aligned}$$

A model is overfit when we have low empirical risk, while having high expected risk. This means that the explanation quality on the data source is much worse than on the training data. The main reason for this is that the theory (\mathcal{H} and ℓ) is so complex that it can explain almost anything.

A model is underfit when we have high empirical risk, while having high expected risk. This means that we neither explain the training data nor the data source. The main reason for this is that the theory is too simple to capture the nature of the data.

The model is learning when we have low empirical risk and low expected risk. This means that the training was successful. Generalization occurs when the expected risk is close to the empirical risk. Note that this does not mean that the explanation is good, since any “blind explanation” will generalize well due to the weak law of large numbers. Ideally, we want generalization and learning.



Figure 1. The map of learning.

2 Theory of convex functions

2.1 Mathematical background

Theorem 3 (Cauchy-Schwarz inequality). Let $\mathbf{u}, \mathbf{v} \in \mathbb{R}^d$, then

$$|\mathbf{u}^\top \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|.$$

For non-zero vectors, this is equivalent to

$$-1 \leq \frac{\mathbf{u}^\top \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} \leq 1.$$

Definition 4 (Spectral norm). Let A be an $m \times d$ matrix, then

$$\|A\| \doteq \max_{\mathbf{v} \in \mathbb{R}^d, \mathbf{v} \neq \mathbf{0}} \frac{\|A\mathbf{v}\|}{\|\mathbf{v}\|} = \max_{\|\mathbf{v}\|=1} \|A\mathbf{v}\|.$$

Intuitively, this means that $\|A\|$ is the largest factor by which a vector can be stretched in length under the mapping $\mathbf{v} \mapsto A\mathbf{v}$.

Theorem 5 (Mean value theorem). Let $a < b$ be real numbers, and let $h : [a, b] \rightarrow \mathbb{R}$ be a continuous function that is differentiable on (a, b) . Then, there exists $c \in (a, b)$ such that

$$h'(c) = \frac{h(b) - h(a)}{b - a}.$$

Geometrically, this means that between a and b , there is a tangent to the graph of h that has the same slope; see Figure 2.

Theorem 6 (Fundamental theorem of calculus). Let $a < b$ be real numbers, and let $h : \text{dom}(h) \rightarrow \mathbb{R}$ be a differentiable function on an open domain $\text{dom}(h) \supset [a, b]$, and such that h' is continuous on $[a, b]$. Then,

$$h(b) - h(a) = \int_a^b h'(t) dt.$$




Figure 2. Illustration of the mean value theorem.

2.2 Convex sets

Definition 7 (Convex set). A set $\mathcal{C} \subseteq \mathbb{R}^d$ is convex if the line segment between any two points of \mathcal{C} lies in \mathcal{C} . I.e., if for any $\mathbf{x}, \mathbf{y} \in \mathcal{C}$ and any λ with $0 \leq \lambda \leq 1$, we have

$$\lambda \mathbf{x} + (1 - \lambda) \mathbf{y} \in \mathcal{C}.$$

Observation. Let $\mathcal{C}_i, i \in I$ be convex sets, where I is a (possibly infinite)





Figure 3. Example of a convex set in \mathbb{R}^2 .



index set. Then,

$$\mathcal{C} = \bigcap_{i \in I} \mathcal{C}_i,$$

is a convex set.

2.3 Convex functions

Definition 8 (Convexity). A function $f : \text{dom}(f) \rightarrow \mathbb{R}$ is convex if (i) $\text{dom}(f)$ is convex and (ii) for all $x, y \in \text{dom}(f)$ and all $\lambda \in [0, 1]$, we have

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y).$$

Geometrically, this condition means that the line segment connecting the two points $(x, f(x))$ and $(y, f(y)) \in \mathbb{R}^{d+1}$ lies pointwise above the graph of f ; see Figure 5.

Definition 9 (Epigraph). The epigraph of a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is defined as

$$\text{epi}(f) \doteq \left\{ (x, \alpha) \in \mathbb{R}^{d+1} \mid x \in \text{dom}(f), \alpha \geq f(x) \right\}.$$

Observation. f is a convex function if and only if $\text{epi}(f)$ is a convex set.

Lemma 10 (Jensen's inequality). Let f be convex, $x_1, \dots, x_m \in \text{dom}(f)$, $\lambda_1, \dots, \lambda_m \in \mathbb{R}_+$ such that $\sum_{i=1}^m \lambda_i = 1$, then

$$f\left(\sum_{i=1}^m \lambda_i x_i\right) \leq \sum_{i=1}^m \lambda_i f(x_i).$$

Proof. We will prove Jensen's inequality by induction. The base case ($m = 2$) is true by definition of convexity. Let Jensen's inequality hold for $m = k$. Consider $m = k + 1$, then

$$\begin{aligned} f\left(\sum_{i=1}^{k+1} \lambda_i x_i\right) &= f\left(\sum_{i=1}^k \lambda_i x_i + \lambda_{k+1} x_{k+1}\right) \\ &= f\left((1 - \lambda_{k+1}) \sum_{i=1}^k \frac{\lambda_i}{1 - \lambda_{k+1}} x_i + \lambda_{k+1} x_{k+1}\right) \\ &\leq (1 - \lambda_{k+1}) f\left(\sum_{i=1}^k \frac{\lambda_i}{1 - \lambda_{k+1}} x_i\right) + \lambda_{k+1} f(x_{k+1}) \\ &\leq (1 - \lambda_{k+1}) \sum_{i=1}^k \frac{\lambda_i}{1 - \lambda_{k+1}} f(x_i) + \lambda_{k+1} f(x_{k+1}) \\ &= \sum_{i=1}^{k+1} \lambda_i f(x_i). \end{aligned}$$




Figure 5. Illustration of the classic definition of convexity.




Figure 6. Epigraph of a convex function.

By definition of convexity.

Induction assumption, $\sum_{i=1}^k \frac{\lambda_i}{1 - \lambda_{k+1}} = 1$.

Since it holds for the base case and the induction step, Jensen's inequality must hold for all m . ■

Lemma 11. Let f be convex and suppose that $\text{dom}(f) \subseteq \mathbb{R}^d$ is open, then f is continuous.

Definition 12 (Differentiable functions). Let $f : \text{dom}(f) \rightarrow \mathbb{R}^m$ where $\text{dom}(f) \subseteq \mathbb{R}^d$ is open. f is called differentiable at $x \in \text{dom}(f)$ if there exists an $m \times d$ matrix A and an error function $r : \mathbb{R}^d \rightarrow \mathbb{R}^m$ defined around $0 \in \mathbb{R}^d$ such that for all y in some neighborhood of x ,

$$f(y) = f(x) + A(y - x) + r(y - x),$$

where $\lim_{v \rightarrow 0} \|r(v)\|/\|v\| = 0$ (error function r is sublinear around 0). A is unique and called the Jacobian matrix of f at x .

Lemma 13 (First-order convexity). Suppose that $\text{dom}(f)$ is open and that f is differentiable. In particular, the gradient

$$\nabla f(x) \doteq \left[\frac{\partial f(x)}{\partial x_1}, \dots, \frac{\partial f(x)}{\partial x_d} \right]$$

exists at every point $x \in \text{dom}(f)$.

Then, f is convex if and only if (i) $\text{dom}(f)$ is convex and (ii)

$$\forall x, y \in \text{dom}(f) : f(y) \geq f(x) + \nabla f(x)^\top (y - x).$$

Geometrically, this means that the graph is above all tangent hyperplanes; see Figure 8.

Proof. We will first prove that convexity implies that the first-order definition holds. Then, we will prove that the first-order definition implies convexity, making the definitions equivalent for differentiable functions f .

\Rightarrow : Suppose f is convex. Then, for all $t \in (0, 1)$,

$$\begin{aligned} f((1-t)x + ty) &\leq (1-t)f(x) + tf(y) \\ \Leftrightarrow f(x + t(y-x)) &\leq f(x) + t(f(y) - f(x)) \\ \Leftrightarrow f(y) &\geq f(x) + \frac{f(x + t(y-x)) - f(x)}{t} \\ &= f(x) + \frac{\nabla f(x)^\top t(y-x) + r(t(y-x))}{t} \\ &= f(x) + \nabla f(x)^\top (y-x) + \underbrace{\frac{r(t(y-x))}{t}}_{\rightarrow 0 \text{ for } t \rightarrow 0}. \end{aligned}$$




Figure 7. Graph of the affine function $f(x) + \nabla f(x)^\top (y - x)$ is a tangent hyperplane to the graph of f at $(x, f(x))$.




Figure 8. Illustration of the first-order characterization of convexity (??).

Definition of convexity

\Leftarrow : Define $\mathbf{z} \doteq \lambda \mathbf{x} + (1 - \lambda) \mathbf{y} \in \text{dom}(f)$ for $\lambda \in [0, 1]$ by convexity of $\text{dom}(f)$. Then, we have the following inequalities,

$$\begin{aligned} f(\mathbf{x}) &\geq f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\mathbf{x} - \mathbf{z}) \\ \lambda f(\mathbf{x}) &\geq \lambda \left(f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\mathbf{x} - \mathbf{z}) \right) \\ f(\mathbf{y}) &\geq f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\mathbf{y} - \mathbf{z}) \\ (1 - \lambda) f(\mathbf{y}) &\geq (1 - \lambda) \left(f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\mathbf{y} - \mathbf{z}) \right). \end{aligned}$$

This implies the following,

$$\begin{aligned} \lambda f(\mathbf{x}) + (1 - \lambda) f(\mathbf{y}) &\geq \lambda \left(f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\mathbf{x} - \mathbf{z}) \right) + (1 - \lambda) \left(f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\mathbf{y} - \mathbf{z}) \right) \\ &= f(\mathbf{z}) + \lambda \nabla f(\mathbf{z})^\top (\mathbf{x} - \mathbf{z}) + (1 - \lambda) \nabla f(\mathbf{z})^\top (\mathbf{y} - \mathbf{z}) \\ &= f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\lambda (\mathbf{x} - \mathbf{z}) + (1 - \lambda) (\mathbf{y} - \mathbf{z})) \\ &= f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\lambda \mathbf{x} + (1 - \lambda) \mathbf{y} - (\lambda \mathbf{z} + (1 - \lambda) \mathbf{z})) \\ &= f(\mathbf{z}) + \nabla f(\mathbf{z})^\top (\mathbf{z} - \mathbf{z}) \\ &= f(\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}). \end{aligned}$$

■

Lemma 14 (First-order convexity alternative). Suppose that $\text{dom}(f)$ is open and that f is differentiable. Then, f is convex if and only if (i) $\text{dom}(f)$ is convex and (ii)

$$\forall \mathbf{x}, \mathbf{y} \in \text{dom}(f) : (\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}))^\top (\mathbf{y} - \mathbf{x}) \geq 0.$$

Geometrically, this means that the gradient is monotonic.

Proof. We will first prove that it holds from left to right, and then from right to left.

\Rightarrow : If f is convex, the first-order characterization of convexity (??) yields the following,

$$\begin{aligned} f(\mathbf{x}) &\geq f(\mathbf{y}) + \nabla f(\mathbf{y})^\top (\mathbf{x} - \mathbf{y}) \\ f(\mathbf{y}) &\geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}), \end{aligned}$$

for all $\mathbf{x}, \mathbf{y} \in \text{dom}(f)$. Adding up these two inequalities yields

$$\begin{aligned} f(\mathbf{x}) + f(\mathbf{y}) &\geq f(\mathbf{y}) + \nabla f(\mathbf{y})^\top (\mathbf{x} - \mathbf{y}) + f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) \\ \Leftrightarrow 0 &\geq \nabla f(\mathbf{y})^\top (\mathbf{x} - \mathbf{y}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) \\ &= (\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}))^\top (\mathbf{x} - \mathbf{y}) \\ \Leftrightarrow 0 &\leq (\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}))^\top (\mathbf{y} - \mathbf{x}). \end{aligned}$$

\Leftarrow : Define $\mathbf{z} \doteq (1 - t) \mathbf{x} + t \mathbf{y} \in \text{dom}(f)$ for $\mathbf{x}, \mathbf{y} \in \text{dom}(f), t \in (0, 1)$ by convexity of $\text{dom}(f)$. Observe that $\mathbf{z} = \mathbf{x} + t(\mathbf{y} - \mathbf{x})$. Then, we have the

following inequality, according to the monotonicity of the gradient,

$$\begin{aligned} (\nabla f(z) - \nabla f(x))^\top (z - x) &\geq 0 \\ (\nabla f(x + t(y - x)) - \nabla f(x))^\top (x + t(y - x) - x) &\geq 0 \\ (\nabla f(x + t(y - x)) - \nabla f(x))^\top (y - x) &\geq 0. \end{aligned}$$

Divide by t

Let $h(t) = f(x + t(y - x))$, then $h'(t) = \nabla f(x + t(y - x))^\top (y - x)$. Hence, we can rewrite the inequality as the following,

$$h'(t) \geq \nabla f(x)^\top (y - x).$$

By the mean value theorem, there exists $c \in (0, 1)$ such that $h'(c) = h(1) - h(0)$. I.e.,

$$h'(c) = f(y) - f(x).$$

Thus, we can rewrite the inequality to the following,

$$\begin{aligned} f(y) &= f(x) + h'(c) \\ &\geq f(x) + \nabla f(x)^\top (y - x), \end{aligned}$$

which is the first-order characterization of convexity (??).

■

Lemma 15 (Second-order convexity). Suppose that $\text{dom}(f)$ is open and that f is twice differentiable. In particular, the Hessian

$$\nabla^2 f(x) = \begin{bmatrix} \frac{\partial^2 f(x)}{\partial x_1^2} & \cdots & \frac{\partial^2 f(x)}{\partial x_1 \partial x_d} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f(x)}{\partial x_d \partial x_1} & \cdots & \frac{\partial^2 f(x)}{\partial x_d^2} \end{bmatrix}$$

exists at every point $x \in \text{dom}(f)$ and is symmetric.

Then, f is convex if and only if (i) $\text{dom}(f)$ is convex and (ii) for all $x \in \text{dom}(f)$, we have

I.e., $\nabla^2 f(x)$ is positive semidefinite if $x^\top \nabla^2 f(x) x \geq 0$ for all x and $x^\top \nabla^2 f(x) x > 0$ for all $x \neq 0$.

Geometrically, this means that f has non-negative curvature everywhere. I.e., the growth rate should be growing.

Observation (Operations that preserve convexity). Let f_1, \dots, f_m be convex functions, $\lambda_1, \dots, \lambda_m \in \mathbb{R}_+$, then

$$f \doteq \max_{i=1}^m f_i,$$

and

$$f \doteq \sum_{i=1}^m \lambda_i f_i.$$

are convex on $\text{dom}(f) \doteq \bigcap_{i=1}^m \text{dom}(f_i)$.

Let f be a convex function with $\text{dom}(f) \subseteq \mathbb{R}^d$, $g: \mathbb{R}^m \rightarrow \mathbb{R}^d$ an affine function, i.e. $g(x) = Ax + b$ for some $A \in \mathbb{R}^{d \times m}$, $b \in \mathbb{R}^d$. Then, the function $f \circ g$ is convex on $\text{dom}(f \circ g) \doteq \{x \in \mathbb{R}^m \mid g(x) \in \text{dom}(f)\}$.

2.4 Minimizing convex functions

Definition 16 (Local minimum). A local minimum of $f: \text{dom}(f) \rightarrow \mathbb{R}$ is a point x such that there exists $\epsilon > 0$ with

$$f(x) \leq f(y) \quad \forall y \in \text{dom}(f) \text{ s.t. } \|y - x\| < \epsilon.$$

Remark. This definition of local minima means that in some small neighborhood around x , x is the smallest point.

Lemma 17. Let x^* be a local minimum of a convex function $f: \text{dom}(f) \rightarrow \mathbb{R}$. Then, x^* is a global minimum, meaning that $f(x^*) \leq f(y) \quad \forall y \in \text{dom}(f)$.

Proof. Proof by contradiction. Suppose there exists $y \in \text{dom}(f)$ such that $f(y) < f(x^*)$. Define $y' \doteq \lambda x^* + (1 - \lambda)y$ for $\lambda \in (0, 1)$. From convexity, we get that $f(y') < f(x^*)$, because $f(y) < f(x^*)$. If we choose λ so close to 1 such that $\|y' - x^*\| < \epsilon$ gives the inequality $f(x^*) \leq f(y')$. This yields a contradiction, thus there cannot exist a $y \in \text{dom}(f)$ such that $f(y) < f(x^*)$, meaning that $f(x^*) \leq f(y)$ for all $y \in \text{dom}(f)$. ■

Lemma 18. Suppose that f is convex and differentiable over an open domain $\text{dom}(f)$. Let $x \in \text{dom}(f)$. If $\nabla f(x) = 0$ (critical point), then x is a global minimum.

Proof. Suppose that $\nabla f(x) = 0$. According to the first-order characterization of convexity, we have

$$f(y) \geq f(x) + \underbrace{\nabla f(x)^\top (y - x)}_{=0} = f(x)$$

for all $y \in \text{dom}(f)$, so x is a global minimum. ■

Definition 19 (Strict convexity). A function $f: \text{dom}(f) \rightarrow \mathbb{R}$ is strictly convex if (i) $\text{dom}(f)$ is convex and (ii) for all $x \neq y \in \text{dom}(f)$ and all $\lambda \in (0, 1)$, we have

$$f(\lambda x + (1 - \lambda)y) < \lambda f(x) + (1 - \lambda)f(y).$$




Figure 9. The maximum operator over m convex functions is a convex function. As can be seen, the epigraph of f is convex.

An example of a strictly convex function can be found in Figure 5, while an example of a function that is not strictly convex can be found in Figure 10.

Lemma 20. Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ be strictly convex. Then, f has at most one global minimum.

But, how do we know a minimizer exists? To be able to find this out, we must use the Weierstrass theorem.

Definition 21 (Sublevel set). $f : \mathbb{R}^d \rightarrow \mathbb{R}, \alpha \in \mathbb{R}$. The set $f^{\leq \alpha} \doteq \{x \in \mathbb{R}^d \mid f(x) \leq \alpha\}$ is the α -sublevel set of f .

Theorem 22 (Weierstrass). Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a continuous function, and suppose there is a non-empty and bounded sublevel set $f^{\leq \alpha}$. Then, f has a global minimum.

Proof. We know that f attains a minimum over the closed and bounded set $f^{\leq \alpha}$ at some x^* , because f is continuous. This x^* is also a global minimum as it has value $f(x^*) \leq \alpha$, while any $x \notin f^{\leq \alpha}$ has value $f(x) > \alpha \geq f(x^*)$. ■

Since convex functions are continuous by ??, the Weierstrass theorem also applies to convex functions. Thus, to figure out whether a convex function has a minimizer, we need to find any non-empty and bounded sublevel set of this function.

2.5 Convex programming

In standard form, optimization problems look like the following,

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_j(x) = 0, \quad j = 1, \dots, p. \end{aligned}$$

The problem domain is then $\mathcal{D} = \bigcap_{i=0}^m \text{dom}(f_i) \cap \bigcap_{j=1}^p \text{dom}(h_j)$.

In a convex program, all f_i are convex functions, and all h_j are affine functions with domain \mathbb{R}^d .




Figure 10. A non-strictly convex function.




Figure 11. Illustration of a sublevel set of a non-convex function.

Definition 23 (Lagrange dual function). Given an optimization problem in standard form, its Lagrangian is the function $L : \mathcal{D} \times \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$ given by

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{j=1}^p \nu_j h_j(\mathbf{x}).$$

The λ_i, ν_j are called Lagrange multipliers. The Lagrange dual function is the function $g : \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R} \cup \{-\infty\}$ defined by

$$g(\boldsymbol{\lambda}, \boldsymbol{\nu}) = \inf_{\mathbf{x} \in \mathcal{D}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}).$$

Only the $(\boldsymbol{\lambda}, \boldsymbol{\nu})$ pairs with $g(\boldsymbol{\lambda}, \boldsymbol{\nu}) > -\infty$ are interesting.

Lemma 24 (Weak Lagrange duality). Let \mathbf{x} be a feasible solution ($f_i(\mathbf{x}) \leq 0$ for $i = 1, \dots, m$ and $h_j(\mathbf{x}) = 0$ for $j = 1, \dots, p$). Let g be the Lagrange dual function and $\boldsymbol{\lambda} \in \mathbb{R}^m, \boldsymbol{\nu} \in \mathbb{R}^p$ such that $\boldsymbol{\lambda} \geq \mathbf{0}$. Then

$$g(\boldsymbol{\lambda}, \boldsymbol{\nu}) \leq f_0(\mathbf{x}).$$

Proof.

$$g(\boldsymbol{\lambda}, \boldsymbol{\nu}) \leq L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\mathbf{x}) + \underbrace{\sum_{i=1}^m \lambda_i f_i(\mathbf{x})}_{\leq 0} + \underbrace{\sum_{j=1}^p \nu_j h_j(\mathbf{x})}_{=0} \leq f_0(\mathbf{x}).$$

■

However, we want to know what the best lower bound is that we can get in this way. For this, we must choose $\boldsymbol{\lambda} \geq \mathbf{0}, \boldsymbol{\nu}$ such that $g(\boldsymbol{\lambda}, \boldsymbol{\nu})$ is maximized. This can be phrased as another optimization problem, called the Lagrange dual,

$$\begin{aligned} & \text{maximize} && g(\boldsymbol{\lambda}, \boldsymbol{\nu}) \\ & \text{subject to} && \boldsymbol{\lambda} \geq \mathbf{0}. \end{aligned}$$

Observation. The Lagrange dual is a convex program, even if the primal is not.

Theorem 25. Suppose that a convex program has a feasible solution $\tilde{\mathbf{x}}$ that in addition satisfies $f_i(\tilde{\mathbf{x}}) < 0, i = 1, \dots, m$ (Slater point). Then, the infimum value of the primal equals the supremum value of its Lagrange dual. Moreover, if the value is finite, it is attained by a feasible solution of the dual,

$$\inf f_0(\mathbf{x}) = \max g(\boldsymbol{\lambda}, \boldsymbol{\nu}).$$

A case of particular interest is that strong duality holds and the joint value is attained in both the primal and dual problem.¹ If all f_i and h_j are differentiable, then the Karush-Kuhn-Tucker (KKT) conditions provide necessary and, under convexity, also sufficient conditions for this case to occur.

¹ This would mean that $\min f_0(\mathbf{x}) = \max g(\boldsymbol{\lambda}, \boldsymbol{\nu})$.

Definition 26 (Zero duality gap). Let $\tilde{\mathbf{x}}$ be feasible for the primal and $(\tilde{\boldsymbol{\lambda}}, \tilde{\boldsymbol{\nu}})$ feasible for the Lagrange dual. The primal and dual solutions $\tilde{\mathbf{x}}$ and $(\tilde{\boldsymbol{\lambda}}, \tilde{\boldsymbol{\nu}})$ are said to have zero duality gap if $f_0(\tilde{\mathbf{x}}) = g(\tilde{\boldsymbol{\lambda}}, \tilde{\boldsymbol{\nu}})$.

If the primal and dual have zero duality gap, then $\min f_0(\mathbf{x}) = \max g(\boldsymbol{\lambda}, \boldsymbol{\nu})$.

The consequence of zero duality gap is the *master equation*,

$$\begin{aligned} f_0(\tilde{\mathbf{x}}) &= g(\tilde{\boldsymbol{\lambda}}, \tilde{\boldsymbol{\nu}}) \\ &= \inf_{\mathbf{x} \in \mathcal{D}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \tilde{\lambda}_i f_i(\mathbf{x}) + \sum_{j=1}^p \tilde{\nu}_j h_j(\mathbf{x}) \right) \\ &\leq f_0(\tilde{\mathbf{x}}) + \underbrace{\sum_{i=1}^m \tilde{\lambda}_i f_i(\tilde{\mathbf{x}})}_{\leq 0} + \underbrace{\sum_{j=1}^p \tilde{\nu}_j h_j(\tilde{\mathbf{x}})}_{=0} \\ &\leq f_0(\tilde{\mathbf{x}}), \end{aligned}$$

which means that all inequalities turn into equalities. Thus,

$$\tilde{\lambda}_i f_i(\tilde{\mathbf{x}}) = 0, \quad i = 1, \dots, m,$$

which is called complementary slackness, because if $\tilde{\lambda}_i \neq 0$, then $f_i(\tilde{\mathbf{x}}) = 0$, and vice versa. Furthermore, if all f_i and h_j are differentiable, then

$$\nabla f_0(\tilde{\mathbf{x}}) + \sum_{i=1}^m \tilde{\lambda}_i \nabla f_i(\tilde{\mathbf{x}}) + \sum_{j=1}^p \tilde{\nu}_j \nabla h_j(\tilde{\mathbf{x}}) = \mathbf{0},$$

which is called the vanishing Lagrangian gradient condition.

In summary, we get the following result.

Theorem 27 (Karush-Kuhn-Tucker necessary conditions). Let $\tilde{\mathbf{x}}$ and $(\tilde{\boldsymbol{\lambda}}, \tilde{\boldsymbol{\nu}})$ be feasible solutions of the primal optimization problem and its Lagrange dual, respectively, with zero duality gap. If all f_i and h_j are differentiable, then

$$\begin{aligned} \tilde{\lambda}_i f_i(\tilde{\mathbf{x}}) &= 0 \quad \text{complementary slackness} \\ \nabla f_0(\tilde{\mathbf{x}}) + \sum_{i=1}^m \tilde{\lambda}_i \nabla f_i(\tilde{\mathbf{x}}) + \sum_{j=1}^p \tilde{\nu}_j \nabla h_j(\tilde{\mathbf{x}}) &= \mathbf{0} \quad \text{vanishing Lagrangian gradient} \end{aligned}$$

If we have a convex program, then the KKT conditions are sufficient for zero duality gap. The motivation behind this is that they may be easier to solve for than solving the primal optimization problem. However, we cannot always count on the KKT conditions to be solvable, because they are only guaranteed if there are primal and dual solutions of zero duality

gap. But, if the primal has a Slater point², then the KKT conditions are equivalent to the existence of a primal minimizer.

² A point \tilde{x} such that $f_i(\tilde{x}) < 0$ for all $i = 1, \dots, m$.

3 Gradient descent

Gradient descent is an optimization algorithm that aims to find the global minimizer. We assume that $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex, differentiable, and has a global minimizer \mathbf{x}^* . Then, the goal of gradient descent is, for $\epsilon > 0$, to find a $\mathbf{x} \in \mathbb{R}^d$ such that

$$f(\mathbf{x}) - f(\mathbf{x}^*) \leq \epsilon.$$

It works by first choosing a $\mathbf{x}_0 \in \mathbb{R}^d$ and then iteratively updating this value by

$$\mathbf{x}_{t+1} = \mathbf{x}_t - \gamma \nabla f(\mathbf{x}_t),$$

for timesteps $t = 0, 1, \dots$ and stepsize $\gamma > 0$.

3.1 Vanilla analysis

We want to be able to bound $f(\mathbf{x}_t) - f(\mathbf{x}^*)$, which tells us how far off we are from the minimum at timestep t . For this, we can use the first-order characterization of convexity,

$$f(\mathbf{x}^*) \geq f(\mathbf{x}_t) + \nabla f(\mathbf{x}_t)^\top (\mathbf{x}^* - \mathbf{x}_t) \iff f(\mathbf{x}_t) - f(\mathbf{x}^*) \leq \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_t - \mathbf{x}^*).$$

Using the gradient descent update rule, we get

$$\nabla f(\mathbf{x}_t) = \frac{\mathbf{x}_t - \mathbf{x}_{t+1}}{\gamma},$$

which gives the following bound

$$\begin{aligned} f(\mathbf{x}_t) - f(\mathbf{x}^*) &\leq \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_t - \mathbf{x}^*) \\ &= \frac{1}{\gamma} (\mathbf{x}_t - \mathbf{x}_{t+1})^\top (\mathbf{x}_t - \mathbf{x}^*) \\ &= \frac{1}{2\gamma} \left(\|\mathbf{x}_t - \mathbf{x}_{t+1}\|^2 + \|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \right) \\ &= \frac{\gamma}{2} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2\gamma} \left(\|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \right). \end{aligned}$$

$2\mathbf{v}^\top \mathbf{w} = \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 - \|\mathbf{v} - \mathbf{w}\|^2$ (cosine theorem).

$$\mathbf{x}_t - \mathbf{x}_{t+1} = \gamma \nabla f(\mathbf{x}_t).$$

Summing this up over the first T iterations, we get an upper bound on the summed error,

$$\begin{aligned} \sum_{t=0}^{T-1} (f(\mathbf{x}_t) - f(\mathbf{x}^*)) &\leq \sum_{t=0}^{T-1} \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_t - \mathbf{x}^*) \\ &= \frac{\gamma}{2} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2\gamma} \left(\|\mathbf{x}_0 - \mathbf{x}^*\|^2 - \|\mathbf{x}_T - \mathbf{x}^*\|^2 \right) && \text{Telescoping sum.} \\ &\leq \frac{\gamma}{2} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2\gamma} \|\mathbf{x}_0 - \mathbf{x}^*\|^2. && \|\mathbf{x}_T - \mathbf{x}^*\|^2 \geq 0. \end{aligned}$$

We can also use this to get a bound on the average error,

$$\frac{1}{T} \sum_{t=0}^{T-1} (f(\mathbf{x}_t) - f(\mathbf{x}^*)) \leq \frac{1}{T} \left(\frac{\gamma}{2} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2\gamma} \|\mathbf{x}_0 - \mathbf{x}^*\|^2 \right).$$




Figure 12. Gradient descent updates.

Take a small step into the direction of the negative gradient to move toward the minimum.

3.2 Lipschitz convex functions

Theorem 28. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be convex and differentiable with a global minimum \mathbf{x}^* . Furthermore, suppose that $\|\mathbf{x}_0 - \mathbf{x}^*\| \leq R$ and $\|\nabla f(\mathbf{x})\| \leq B$ for all \mathbf{x} . Choosing the stepsize,

$$\gamma \doteq \frac{R}{B\sqrt{T}},$$

gradient descent yields

$$\frac{1}{T} \sum_{t=0}^{T-1} (f(\mathbf{x}_t) - f(\mathbf{x}^*)) \leq \frac{RB}{\sqrt{T}}.$$

A function $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is Lipschitz continuous if there exists an M , such that $\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^m$:

$$\|f(\mathbf{x}) - f(\mathbf{y})\| \leq M\|\mathbf{x} - \mathbf{y}\|.$$

This holds if and only if the gradient is bounded.

Proof. We can plug our bounds into the vanilla analysis,

$$\begin{aligned} \sum_{t=0}^{T-1} (f(\mathbf{x}_t) - f(\mathbf{x}^*)) &\leq \frac{\gamma}{2} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2\gamma} \|\mathbf{x}_0 - \mathbf{x}^*\|^2 \\ &\leq \frac{\gamma}{2} B^2 T + \frac{1}{2\gamma} R^2 \\ &\doteq q(\gamma). \end{aligned}$$

Now, we want to choose γ such that the bound $q(\gamma)$ is minimized. We can compute the minimum by computing the derivative,

$$q'(\gamma) = \frac{1}{2} B^2 T - \frac{R^2}{2\gamma^2},$$

and solving for $q'(\gamma) = 0$, which yields

$$\gamma = \frac{R}{B\sqrt{T}}.$$

Then, we can compute the bound by

$$q\left(\frac{R}{B\sqrt{T}}\right) = RB\sqrt{T}.$$

Dividing by T yields the result. ■

We want to find out how many iterations we would need to ensure that the average error is bounded by ϵ . We can use ?? to compute a lower bound on the number of iterations,

$$\frac{RB}{\sqrt{T}} \leq \epsilon \implies T \geq \frac{R^2 B^2}{\epsilon^2}.$$

So, the amount of iterations until convergence is of the order $\mathcal{O}(1/\epsilon^2)$. This means that we need at most $10000 \cdot R^2 B^2$ iterations to achieve an error of 0.01.

3.3 Smooth functions

Definition 29 (Smoothness). Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ be differentiable, $\mathcal{X} \subseteq \text{dom}(f)$ convex, $L \in \mathbb{R}_+$. f is smooth (with parameter L) over \mathcal{X} if $\forall \mathbf{x}, \mathbf{y} \in \mathcal{X}$:

$$\underbrace{f(\mathbf{y}) \leq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{L}{2} \|\mathbf{x} - \mathbf{y}\|^2}_{\text{first-order convexity}}.$$

Remark. This definition does not require convexity.

Geometrically, this definition of smoothness means that the graph f is below a not too steep tangent paraboloid at $(\mathbf{x}, f(\mathbf{x}))$; see Figure 13. Furthermore, we can easily check if f is smooth with parameter L if the following function is convex,

$$g(\mathbf{x}) \doteq \frac{L}{2} \mathbf{x}^\top \mathbf{x} - f(\mathbf{x}),$$

over $\text{dom}(g) \doteq \text{dom}(f)$.

Observation (Operations that preserve smoothness). Let f_1, \dots, f_m be smooth with parameters L_1, \dots, L_m and let $\lambda_1, \dots, \lambda_m \in \mathbb{R}_+$. Then, the function $f \doteq \sum_{i=1}^m \lambda_i f_i$ is smooth with parameter $\sum_{i=1}^m \lambda_i L_i$.

Let f be smooth with parameter L and let $g(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$. Then, the function $f \circ g$ is smooth with parameter $L\|A\|^2$, where $\|A\|$ is the spectral norm of A .

Lemma 30. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be convex and differentiable, then f is smooth with parameter L if and only if $\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^d$:

$$\|\nabla f(\mathbf{x}) - \nabla f(\mathbf{y})\| \leq L\|\mathbf{x} - \mathbf{y}\|.$$

I.e., ∇f is Lipschitz continuous.

Lemma 31 (Sufficient decrease). Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable and smooth with parameter L . With stepsize

$$\gamma \doteq \frac{1}{L},$$

gradient descent satisfies

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2L} \|\nabla f(\mathbf{x}_t)\|^2.$$

Sufficient decrease implies that, every step, we get closer to the minimum.

“Not too curved.”




Figure 13. Plot of $f(x) = x$ with the tangent paraboloids at $x = -8, 5$, showing smoothness with parameter $L = 1$. Furthermore, it shows that $g(x) = \frac{1}{2}x^2 - f(x)$ is convex.

Proof.

$$\begin{aligned}
 f(\mathbf{x}_{t+1}) &\leq f(\mathbf{x}_t) + \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_{t+1} - \mathbf{x}_t) + \frac{L}{2} \|\mathbf{x}_t - \mathbf{x}_{t+1}\|^2 \\
 &= f(\mathbf{x}_t) + \nabla f(\mathbf{x}_t)^\top \left(-\frac{\nabla f(\mathbf{x}_t)}{L} \right) + \frac{L}{2} \left\| \frac{\nabla f(\mathbf{x}_t)}{L} \right\|^2 \\
 &= f(\mathbf{x}_t) - \frac{1}{L} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2L} \|\nabla f(\mathbf{x}_t)\|^2 \\
 &= f(\mathbf{x}_t) - \frac{1}{2L} \|\nabla f(\mathbf{x}_t)\|^2.
 \end{aligned}$$

Smoothness.

$$\text{GD: } \mathbf{x}_{t+1} = \mathbf{x}_t - \frac{1}{L} \nabla f(\mathbf{x}_t).$$

Theorem 32. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be convex and differentiable with a global minimum \mathbf{x}^* . Furthermore, suppose that f is smooth with parameter L . Choosing stepsize

$$\gamma \doteq \frac{1}{L},$$

gradient descent yields

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) \leq \frac{L}{2T} \|\mathbf{x}_0 - \mathbf{x}^*\|^2, \quad T > 0.$$

Proof. Due to sufficient decrease, we can bound the sum of squared gradients (which will be useful when bounding the vanilla analysis),

$$\begin{aligned}
 \frac{1}{2L} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 &\leq \sum_{t=0}^{T-1} f(\mathbf{x}_t) - f(\mathbf{x}_{t+1}) \\
 &= f(\mathbf{x}_0) - f(\mathbf{x}_T).
 \end{aligned}$$

$$\text{SD: } f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2L} \|\nabla f(\mathbf{x}_t)\|^2.$$

Telescoping sum.

Using the vanilla analysis, we can derive a bound on the average error,

$$\begin{aligned}
 \sum_{t=0}^{T-1} (f(\mathbf{x}_t) - f(\mathbf{x}^*)) &\leq \frac{1}{2L} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{L}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2 \\
 &\leq f(\mathbf{x}_0) - f(\mathbf{x}_T) + \frac{L}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2 \\
 \sum_{t=1}^T (f(\mathbf{x}_t) - f(\mathbf{x}^*)) &\leq \frac{L}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2 \\
 \frac{1}{T} \sum_{t=1}^T (f(\mathbf{x}_t) - f(\mathbf{x}^*)) &\leq \frac{L}{2T} \|\mathbf{x}_0 - \mathbf{x}^*\|^2.
 \end{aligned}$$

$$\gamma \doteq \frac{1}{L}.$$

From sufficient decrease, we know that the last iterate must be the best, thus

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) \leq \frac{1}{T} \sum_{t=1}^T (f(\mathbf{x}_t) - f(\mathbf{x}^*)) \leq \frac{L}{2T} \|\mathbf{x}_0 - \mathbf{x}^*\|^2.$$

Using this result, we can compute a bound on T to achieve an error smaller than ϵ ,

$$f(x_T) - f(x^*) \leq \frac{L}{2T} \|x_0 - x^*\|^2 \leq \frac{R^2 L}{2T} \leq \epsilon.$$

The error then becomes,

$$T \geq \frac{R^2 L}{2\epsilon},$$

which is on the order of $\mathcal{O}(1/\epsilon)$. This means that we need at most $50 \cdot R^2 L$ iterations for an error of 0.01 (as opposed to $10000 \cdot R^2 B^2$ in the Lipschitz case).

3.4 Smooth and strongly convex functions

Definition 33 (Strong convexity). Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ be a convex and differentiable function, $\mathcal{X} \in \text{dom}(f)$ convex and $\mu \in \mathbb{R}_+, \mu > 0$. f is called *strongly convex* with parameter μ over \mathcal{X} if $\forall x, y \in \mathcal{X}$:

$$f(y) \geq f(x) + \nabla f(x)^\top (y - x) + \frac{\mu}{2} \|x - y\|^2.$$

Geometrically, this means that, for any x , the graph of f is above a not too flat tangent paraboloid at $(x, f(x))$; see Figure 14. Furthermore, we can easily check if f is strongly convex if and only if the following function is convex,

$$g(x) \doteq f(x) - \frac{\mu}{2} x^\top x.$$

By assuming that a function f is smooth and strongly convex, we can use a stronger lower bound to derive a bound on the error from the vanilla analysis.

Theorem 34. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be convex and differentiable with a global minimum x^* . Furthermore, suppose that f is smooth with parameter L and strongly convex with parameter $\mu > 0$. Choosing $\gamma \doteq 1/L$, gradient descent with arbitrary x_0 satisfies the following two properties,

1. Squared distances to x^* are geometrically decreasing,

$$\|x_{t+1} - x^*\|^2 \leq \left(1 - \frac{\mu}{L}\right) \|x_t - x^*\|^2;$$

2. The absolute error after T iterations is exponentially small in T ,

$$f(x_T) - f(x^*) \leq \frac{L}{2} \left(1 - \frac{\mu}{L}\right)^T \|x_0 - x^*\|^2.$$




Figure 14. Plot of $f(x) = x^2$ with the tangent paraboloids at $x = -6, 4$, showing strong convexity with parameter $\mu = 1$. Furthermore, it shows $g(x) = f(x) - \frac{1}{2}x^2$, which is convex.

Proof of 1. Using the vanilla analysis, we know the following,

$$\begin{aligned} f(\mathbf{x}_t) - f(\mathbf{x}^*) &\leq \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_t - \mathbf{x}^*) \\ &= \frac{\gamma}{2} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2\gamma} \left(\|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \right). \end{aligned}$$

But, using strong convexity, we have a stronger lower bound on the left hand side,

$$\begin{aligned} f(\mathbf{x}_t) - f(\mathbf{x}^*) &\leq \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_t - \mathbf{x}^*) - \frac{\mu}{2} \|\mathbf{x}_t - \mathbf{x}^*\|^2 && \text{Strong convexity.} \\ &= \frac{\gamma}{2} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{2\gamma} \left(\|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \right) \\ &\quad - \frac{\mu}{2} \|\mathbf{x}_t - \mathbf{x}^*\|^2. \end{aligned}$$

This can be rewritten to the following bound,

$$\|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \leq \underbrace{2\gamma(f(\mathbf{x}^*) - f(\mathbf{x}_t)) + \gamma^2 \|\nabla f(\mathbf{x}_t)\|^2}_{\text{"noise"}} + (1 - \mu\gamma) \|\mathbf{x}_t - \mathbf{x}^*\|^2.$$

Now we need to show that the noise is non-positive,

$$\begin{aligned} 2\gamma(f(\mathbf{x}^*) - f(\mathbf{x}_t)) + \gamma^2 \|\nabla f(\mathbf{x}_t)\|^2 &= \frac{2}{L} (f(\mathbf{x}^*) - f(\mathbf{x}_t)) + \frac{1}{L^2} \|\nabla f(\mathbf{x}_t)\|^2 && \gamma \doteq \frac{1}{L}. \\ &\leq \frac{2}{L} (f(\mathbf{x}_{t+1}) - f(\mathbf{x}_t)) + \frac{1}{L^2} \|\nabla f(\mathbf{x}_t)\|^2 && \text{Sufficient decrease.} \\ &\leq -\frac{1}{L^2} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{1}{L^2} \|\nabla f(\mathbf{x}_t)\|^2 && \text{SD: } f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2L} \|\nabla f(\mathbf{x}_t)\|^2. \\ &= 0. \end{aligned}$$

Hence, the noise is non-positive, and we get the following,

$$\|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \leq \left(1 - \frac{\mu}{L}\right) \|\mathbf{x}_t - \mathbf{x}^*\|^2.$$

■

Proof of 2. From (1), we know the following,

$$\|\mathbf{x}_T - \mathbf{x}^*\|^2 \leq \left(1 - \frac{\mu}{L}\right)^T \|\mathbf{x}_0 - \mathbf{x}^*\|^2.$$

Using smoothness, we can derive a bound on the final iterate error,

$$\begin{aligned} f(\mathbf{x}_T) - f(\mathbf{x}^*) &\leq \nabla f(\mathbf{x}^*)^\top (\mathbf{x}_T - \mathbf{x}^*) + \frac{L}{2} \|\mathbf{x}_T - \mathbf{x}^*\|^2 \\ &= \frac{L}{2} \|\mathbf{x}_T - \mathbf{x}^*\|^2 && \nabla f(\mathbf{x}^*) = 0, \text{ because it is a stationary point.} \\ &\leq \frac{L}{2} \left(1 - \frac{\mu}{L}\right)^T \|\mathbf{x}_0 - \mathbf{x}^*\|^2. && \text{Using (1).} \end{aligned}$$

■

From there, we can derive a lower bound on the number of iterations T to get an error of at most ϵ ,

$$T \geq \frac{L}{\mu} \log\left(\frac{R^2 L}{2\epsilon}\right),$$

This means that we need $\frac{L}{\mu} \log(50 \cdot R^2 L)$ iterations for an error of at most 0.01, as opposed to $50 \cdot R^2 L$ in the smooth case. This bound only depends linearly on $\frac{\mu}{L}$, which might be very high.

4 Projected gradient descent

In constrained optimization, we want to minimize $f(x)$, subject to $x \in \mathcal{X} \subseteq \mathbb{R}^d$. We will assume that $\mathcal{X} \subsetneq \mathbb{R}^d$ is a closed convex set. The idea of projected gradient descent is to project onto \mathcal{X} after every step,

$$\Pi_{\mathcal{X}}(\mathbf{y}) \doteq \operatorname{argmin}_{x \in \mathcal{X}} \|\mathbf{x} - \mathbf{y}\|^2.$$

The algorithm thus alternates between the following two steps,

$$\begin{aligned} \mathbf{y}_{t+1} &\doteq \mathbf{x}_t - \gamma \nabla f(\mathbf{x}_t) \\ \mathbf{x}_{t+1} &\doteq \Pi_{\mathcal{X}}(\mathbf{y}_{t+1}). \end{aligned}$$

Observation. Let $\mathcal{X} \subseteq \mathbb{R}^d$ be closed and convex, $\mathbf{x} \in \mathcal{X}, \mathbf{y} \in \mathbb{R}^d$, then

1. $(\mathbf{x} - \Pi_{\mathcal{X}}(\mathbf{y}))^\top (\mathbf{y} - \Pi_{\mathcal{X}}(\mathbf{y})) \leq 0$;
2. $\|\mathbf{x} - \Pi_{\mathcal{X}}(\mathbf{y})\|^2 + \|\mathbf{y} - \Pi_{\mathcal{X}}(\mathbf{y})\|^2 \leq \|\mathbf{x} - \mathbf{y}\|^2$.

The two properties equivalently say that the vectors $\mathbf{y} - \Pi_{\mathcal{X}}(\mathbf{y})$ and $\mathbf{y} - \Pi_{\mathcal{X}}(\mathbf{y})$ form an obtuse angle; see Figure 15.




Figure 15. Proof by picture of the properties of the projection step made in projected gradient descent.

4.1 Lipschitz convex functions

Since we minimize f over a closed and bounded convex set \mathcal{X} , we get the existence of a minimizer and bound R for free. Furthermore, if we assume that f is continuously differentiable, we also have a bound B for the gradient norms over \mathcal{X} . This makes the following theorem a much more useful result than in the unconstrained case, since we need to make less assumptions about f .

Theorem 35. Let $f : \operatorname{dom}(f) \rightarrow \mathbb{R}$ be convex and differentiable, $\mathcal{X} \subseteq \operatorname{dom}(f)$ closed and convex, \mathbf{x}^* a minimizer of f over \mathcal{X} . Furthermore, suppose that $\|\mathbf{x}_0 - \mathbf{x}^*\| \leq R$ and $\|\nabla f(\mathbf{x})\| \leq B$ for all $\mathbf{x} \in \mathcal{X}$. Choosing the constant stepsize,

$$\gamma \doteq \frac{R}{B\sqrt{T}},$$

with projected gradient descent yields the following bound,

$$\frac{1}{T} \sum_{t=0}^{T-1} (f(\mathbf{x}_t) - f(\mathbf{x}^*)) \leq \frac{RB}{\sqrt{T}}.$$

Proof. We only need to show that the vanilla analysis still holds for the projected \mathbf{x}_{t+1} . We know that for the non-projected iterate, we get the following equality from the vanilla analysis,

$$\nabla f(\mathbf{x}_t)^\top (\mathbf{x}_t - \mathbf{x}^*) = \frac{1}{2\gamma} \left(\gamma^2 \|\nabla f(\mathbf{x}_t)\|^2 + \|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{y}_{t+1} - \mathbf{x}^*\|^2 \right).$$

By the second property of projected gradient descent, we know $\|x_{t+1} - x^*\|^2 \leq \|y_{t+1} - x^*\|^2$, hence we get

$$\nabla f(x_t)^\top (x_t - x^*) \leq \frac{1}{2\gamma} \left(\gamma^2 \|\nabla f(x_t)\|^2 + \|x_t - x^*\|^2 - \|x_{t+1} - x^*\|^2 \right).$$

Then, we can prove the rest of the vanilla analysis as for unbounded gradient descent. ■

This is the same bound as in the unconstrained case, thus the number of necessary iteration is of the order $\mathcal{O}(1/\epsilon^2)$.

4.2 Smooth functions

Lemma 36 (Sufficient decrease of projected gradient descent). Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ be differentiable and smooth with parameter L over a closed and convex set $\mathcal{X} \subseteq \text{dom}(f)$. Choosing stepsize

$$\gamma \doteq \frac{1}{L},$$

projected gradient descent satisfies

$$f(x_{t+1}) \leq f(x_t) - \frac{1}{2L} \|\nabla f(x_t)\|^2 + \underbrace{\frac{L}{2} \|y_{t+1} - x_{t+1}\|^2}_{\text{additional term}}.$$

Proof.

$$\begin{aligned} f(x_{t+1}) &\leq f(x_t) + \nabla f(x_t)^\top (x_{t+1} - x_t) + \frac{L}{2} \|x_t - x_{t+1}\|^2 \\ &= f(x_t) - L(y_{t+1} - x_t)^\top (x_{t+1} - x_t) + \frac{L}{2} \|x_t - x_{t+1}\|^2 \\ &= f(x_t) - \frac{L}{2} \left(\|y_{t+1} - x_t\|^2 + \|x_{t+1} - x_t\|^2 - \|y_{t+1} - x_{t+1}\|^2 \right) \\ &\quad + \frac{L}{2} \|x_t - x_{t+1}\|^2 \\ &= f(x_t) - \frac{L}{2} \|y_{t+1} - x_t\|^2 + \frac{L}{2} \|y_{t+1} - x_{t+1}\|^2 \\ &= f(x_t) - \frac{1}{2L} \|\nabla f(x_t)\|^2 + \frac{L}{2} \|y_{t+1} - x_{t+1}\|^2. \end{aligned}$$

Smoothness.

$$y_{t+1} = x_t - \frac{1}{L} \nabla f(x_t) \Rightarrow \nabla f(x_t) = -L(y_{t+1} - x_t).$$

$$2v^\top w = \|v\|^2 + \|w\|^2 - \|v - w\|^2.$$

$$y_{t+1} = x_t - \frac{1}{L} \nabla f(x_t) \Rightarrow y_{t+1} - x_t = -\frac{1}{L} \nabla f(x_t).$$

Thus, we also have a sufficient decrease lemma for this version of gradient descent, which has an additional term in its bound. However, as we will see, this does not matter, because we can compensate for it in the vanilla analysis.

Theorem 37. Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ be convex and differentiable. Let $\mathcal{X} \subseteq \text{dom}(f)$ be a closed convex set, and \mathbf{x}^* the minimizer of f over \mathcal{X} . Furthermore, suppose that f is smooth over \mathcal{X} with parameter L . Choosing stepsize

$$\gamma \doteq \frac{1}{L},$$

projected gradient descent yields the following bound,

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) \leq \frac{L}{2T} \|\mathbf{x}_0 - \mathbf{x}^*\|^2.$$

Proof. From property 2 of gradient descent, we get the following inequality,

$$\|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 + \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \leq \|\mathbf{y}_{t+1} - \mathbf{x}^*\|^2.$$

Using this inequality, we get the following upper bound,

$$\begin{aligned} f(\mathbf{x}_t) - f(\mathbf{x}^*) &\leq \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_t - \mathbf{x}^*) \\ &= \frac{1}{2\gamma} \left(\gamma^2 \|\nabla f(\mathbf{x}_t)\|^2 + \|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{y}_{t+1} - \mathbf{x}^*\|^2 \right) \\ &\leq \frac{1}{2\gamma} \left(\gamma^2 \|\nabla f(\mathbf{x}_t)\|^2 + \|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \right. \\ &\quad \left. - \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \right). \end{aligned}$$

See vanilla analysis, where \mathbf{x}_{t+1} is substituted by \mathbf{y}_{t+1} , since that is the next unconstrained iterate.

We use sufficient decrease to bound the sum of gradients,

$$\begin{aligned} \frac{1}{2L} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 &\leq \sum_{t=0}^{T-1} f(\mathbf{x}_t) - f(\mathbf{x}_{t+1}) + \frac{L}{2} \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \\ &= f(\mathbf{x}_0) - f(\mathbf{x}_T) + \frac{L}{2} \sum_{t=0}^{T-1} \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2. \end{aligned}$$

Sufficient decrease.

Telescoping sum.

Now, we can bound the summed error and we will see that the additional terms cancel,

$$\begin{aligned} \sum_{t=0}^{T-1} (f(\mathbf{x}_t) - f(\mathbf{x}^*)) &\leq \frac{1}{2L} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 + \frac{L}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2 \\ &\quad - \frac{L}{2} \sum_{t=0}^{T-1} \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \\ &\leq f(\mathbf{x}_0) - f(\mathbf{x}_T) + \frac{L}{2} \sum_{t=0}^{T-1} \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \\ &\quad + \frac{L}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2 - \frac{L}{2} \sum_{t=0}^{T-1} \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \\ &= f(\mathbf{x}_0) - f(\mathbf{x}_T) + \frac{L}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2, \end{aligned}$$

which results in

$$\sum_{t=1}^T (f(\mathbf{x}_t) - f(\mathbf{x}^*)) \leq \frac{L}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2.$$

Due to sufficient decrease, we get

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) \leq \frac{L}{2T} \|\mathbf{x}_0 - \mathbf{x}^*\|^2.$$

■

This is the same bound as in the unconstrained case, thus the number of necessary iterations is of the order $\mathcal{O}(1/\epsilon)$.

4.3 Smooth and strongly convex functions

Theorem 38. Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ be convex and differentiable. Let $\mathcal{X} \subseteq \text{dom}(f)$ be a nonempty closed and convex set and suppose that f is smooth over \mathcal{X} with parameter L and strongly convex over \mathcal{X} with parameter $\mu > 0$. Choosing stepsize

$$\gamma \doteq \frac{1}{L},$$

projected gradient descent satisfies the following two properties,

1. Squared distance to \mathbf{x}^* are geometrically decreasing,

$$\|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 \leq \left(1 - \frac{\mu}{L}\right) \|\mathbf{x}_t - \mathbf{x}^*\|^2;$$

2. The absolute error after T iterations is exponentially small in T ,

$$\begin{aligned} f(\mathbf{x}_T) - f(\mathbf{x}^*) &\leq \frac{L}{2} \left(1 - \frac{\mu}{L}\right)^T \|\mathbf{x}_0 - \mathbf{x}^*\|^2 \\ &\quad + \underbrace{\|\nabla f(\mathbf{x}^*)\| \left(1 - \frac{\mu}{L}\right)^{T/2} \|\mathbf{x}_0 - \mathbf{x}^*\|}_{\text{additional term}}. \end{aligned}$$

Proof. These proofs are similar to the normal gradient descent case, starting from the constrained vanilla bound,

$$\frac{1}{2\gamma} \left(\gamma^2 \|\nabla f(\mathbf{x}_t)\|^2 + \|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 - \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \right),$$

which can be strengthened to

$$\frac{1}{2\gamma} \left(\gamma^2 \|\nabla f(\mathbf{x}_t)\|^2 + \|\mathbf{x}_t - \mathbf{x}^*\|^2 - \|\mathbf{x}_{t+1} - \mathbf{x}^*\|^2 - \|\mathbf{y}_{t+1} - \mathbf{x}_{t+1}\|^2 \right) - \frac{\mu}{2} \|\mathbf{x}_t - \mathbf{x}^*\|^2,$$

on $f(\mathbf{x}_t) - f(\mathbf{x}^*)$.

■

5 Coordinate descent

A problem with gradient descent in large-scale learning is that we need to compute the full gradient $\nabla f(x_t)$ in every iteration. The idea of *coordinate descent* is to update only one coordinate of the iterate at a time. To do this, we only need to compute one coordinate of $\nabla f(x_t)$ at a time, which we assume to be a factor of d faster to compute.

However, we also expect to pay a price for this in terms of the number of iterations required. It turns out that, in the worst case, the number of iterations will increase by a factor of d , so we only stand to gain. Under additional assumptions about f , coordinate descent can lead to provable speedups.

Definition 39 (Polyak-Łojasiewicz (PL) inequality). Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a differentiable function with a global minimum x^* . We say that f satisfies the PL inequality if the following holds for some $\mu > 0$,

$$\frac{1}{2} \|\nabla f(x)\|^2 \geq \mu(f(x) - f(x^*)), \quad \forall x \in \mathbb{R}^d.$$

Observation. The PL inequality directly implies that every critical point is a minimizer of f .

There are non-convex functions that satisfy the PL inequality.

Lemma 40. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable and strongly convex with parameter $\mu > 0$. Then f satisfies the PL inequality for the same μ .

To analyze coordinate descent, we need the notion of *coordinate-wise smoothness*.

Definition 41 (Coordinate-wise smoothness). Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable, and $\mathcal{L} = [L_1, \dots, L_d] \in \mathbb{R}_+^d$. Function f is called coordinate-wise smooth (with parameter \mathcal{L}) if for every coordinate $i = 1, \dots, d$, the following holds,

$$f(x + \lambda e_i) \leq f(x) + \lambda \nabla_i f(x) + \frac{L_i}{2} \lambda^2, \quad \forall x \in \mathbb{R}^d, \lambda \in \mathbb{R}.$$

Compare this to the definition of smoothness; ?? In our new coordinate-wise definition, we define y as $x + \lambda e_i$, since we only want to change coordinate i . Hence, $y - x$ becomes λe_i . From there, it is easy to see that $\nabla f(x)^\top (y - x)$ becomes $\lambda \nabla_i f(x)$, and $\|x - y\|$ becomes λ . Thus, smoothness with parameter L implies coordinate-wise smoothness with parameter $\mathcal{L} = [L, \dots, L]$.

Example 42. $f(x_1, x_2) = x_1^2 + 10x_2^2$ is smooth with parameter $L = 20$, but f is coordinate-wise smooth with parameter $\mathcal{L} = [2, 20]$. Such differences will become important later when showing faster convergence of coordinate descent.

In general, coordinate (gradient) descent algorithms perform the following actions,

$$\begin{aligned} &\text{choose an active coordinate } i \in [d] \\ &\mathbf{x}_{t+1} \doteq \mathbf{x}_t - \gamma_i \nabla_i f(\mathbf{x}_t) \mathbf{e}_i. \end{aligned}$$

We call the coordinate that is currently being update *active*.

Lemma 43 (Coordinate-wise sufficient decrease). Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable and coordinate-wise smooth with parameter $\mathcal{L} = [L_1, \dots, L_d]$. With active coordinate i in iteration t and stepsize

$$\gamma_i \doteq \frac{1}{L_i},$$

coordinate descent satisfies

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2L_i} |\nabla_i f(\mathbf{x}_t)|^2.$$

Proof. Let $\lambda = -\nabla_i f(\mathbf{x}_t)/L_i$, then $\mathbf{x}_{t+1} = \mathbf{x}_t + \lambda \mathbf{e}_i$. Then, we can apply coordinate-wise smoothness,

$$\begin{aligned} f(\mathbf{x}_{t+1}) &\leq f(\mathbf{x}_t) + \lambda \nabla_i f(\mathbf{x}_t) + \frac{L_i}{2} \lambda^2 \\ &= f(\mathbf{x}_t) - \frac{1}{L_i} |\nabla_i f(\mathbf{x}_t)|^2 + \frac{1}{2L_i} |\nabla_i f(\mathbf{x}_t)|^2 \\ &= f(\mathbf{x}_t) - \frac{1}{2L_i} |\nabla_i f(\mathbf{x}_t)|^2. \end{aligned}$$

Coordinate-wise smoothness. ■

5.1 Randomized coordinate descent

In *randomized gradient descent*, the active coordinate is chosen uniformly at random,

$$\begin{aligned} &\text{sample } i \in [d] \text{ uniformly at random} \\ &\mathbf{x}_{t+1} \doteq \mathbf{x}_t - \gamma_i \nabla_i f(\mathbf{x}_t) \mathbf{e}_i. \end{aligned}$$

Randomized coordinate descent is at least as fast as gradient descent on smooth functions if we assume that it is d times cheaper to update one coordinate than the full iterate [Nesterov, 2012]. If we additionally assume the PL inequality, we can obtain faster convergence.

Theorem 44. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable with global minimum \mathbf{x}^* . Suppose that f is coordinate-wise smooth with parameter L and satisfies the PL inequality with parameter $\mu > 0$. Choosing stepsize

$$\gamma_i \doteq \frac{1}{L},$$

randomized gradient descent with arbitrary \mathbf{x}_0 yields

$$\mathbb{E}[f(\mathbf{x}_T) - f(\mathbf{x}^*)] \leq \left(1 - \frac{\mu}{dL}\right)^T (f(\mathbf{x}_0) - f(\mathbf{x}^*)).$$

Proof. Sufficient decrease yields

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2L} |\nabla_i f(\mathbf{x}_t)|^2.$$

By taking the expectation of both sides with respect to i , we have

$$\begin{aligned} \mathbb{E}_i[f(\mathbf{x}_{t+1}) \mid \mathbf{x}_t] &\leq \mathbb{E}_i \left[f(\mathbf{x}_t) - \frac{1}{2L} |\nabla_i f(\mathbf{x}_t)|^2 \mid \mathbf{x}_t \right] \\ &= f(\mathbf{x}_t) - \frac{1}{2L} \sum_{i=1}^d \frac{1}{d} |\nabla_i f(\mathbf{x}_t)|^2 \\ &= f(\mathbf{x}_t) - \frac{1}{2dL} \|\nabla f(\mathbf{x}_t)\|^2 \\ &\leq f(\mathbf{x}_t) - \frac{\mu}{dL} (f(\mathbf{x}_t) - f(\mathbf{x}^*)). \end{aligned}$$

PL inequality.

Subtracting $f(\mathbf{x}^*)$ from both sides and taking the expectation over \mathbf{x}_t , we obtain

$$\mathbb{E}[f(\mathbf{x}_{t+1}) - f(\mathbf{x}^*)] \leq \left(1 - \frac{\mu}{dL}\right) (f(\mathbf{x}_t) - f(\mathbf{x}^*)).$$

$\mathbb{E}_i[f(\mathbf{x}_{t+1}) \mid \mathbf{x}_t]$ is a random variable whose expectation is $\mathbb{E}_{\mathbf{x}_t}[f(\mathbf{x}_{t+1})]$.

The statement follows. ■

5.2 Importance sampling

As seen, uniformly random selection of the active coordinate does not yield a better bound than gradient descent. However, we have not made use of the fact that the coordinate-wise smoothness parameters can differ. Intuitively, we would want to sample coordinates with high smoothness more frequently than coordinates with low smoothness, since they change more rapidly, leading to faster convergence to the optimum. This leads us to *importance sampling* [Nesterov, 2012],

$$\begin{aligned} &\text{sample } i \in [d] \text{ with probability } \frac{L_i}{\sum_{j=1}^d L_j} \\ &\mathbf{x}_{t+1} \doteq \mathbf{x}_t - \gamma_i \nabla_i f(\mathbf{x}_t) \mathbf{e}_i. \end{aligned}$$

Theorem 45. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable with a global minimum \mathbf{x}^* . Suppose that f is coordinate-wise smooth with parameter $\mathcal{L} = [L_1, \dots, L_d]$ and satisfies the PL inequality with parameter $\mu > 0$. Let

$$\bar{L} = \frac{1}{d} \sum_{i=1}^d L_i$$

be the average of coordinate-wise smoothness constants. Then, coordinate descent with importance sampling and arbitrary \mathbf{x}_0 yields

$$\mathbb{E}[f(\mathbf{x}_T) - f(\mathbf{x}^*)] \leq \left(1 - \frac{\mu}{d\bar{L}}\right)^T (f(\mathbf{x}_0) - f(\mathbf{x}^*)).$$

Proof. The proof is nearly identical to the proof of randomized coordinate descent. The difference lies in the expectation over i . Importance sampling yields

$$\begin{aligned} \mathbb{E}_i \left[f(\mathbf{x}_t) - \frac{1}{2L_i} |\nabla_i f(\mathbf{x}_t)|^2 \mid \mathbf{x}_t \right] &= f(\mathbf{x}_t) - \sum_{i=1}^d \frac{L_i}{\sum_{j=1}^d L_j} \frac{1}{2L_i} |\nabla_i f(\mathbf{x}_t)|^2 \\ &= f(\mathbf{x}_t) - \frac{1}{2d\bar{L}} \sum_{i=1}^d |\nabla_i f(\mathbf{x}_t)|^2. \end{aligned} \quad d\bar{L} = \sum_{i=1}^d L_i.$$

The rest follows identically. ■

Note that \bar{L} can be much smaller than $L = \max_{i=1}^d L_i$, so coordinate descent with important sampling is potentially faster than randomized gradient descent. In the worst-case, both algorithms are the same.

5.3 Steepest coordinate descent

In contrast to random coordinate descent, *steepest coordinate descent* chooses the active coordinate according to the coordinate with the largest gradient (*Gauss-Southwell* rule),

$$\begin{aligned} \text{choose } i &= \operatorname{argmax}_{i \in [d]} |\nabla_i f(\mathbf{x}_t)| \\ \mathbf{x}_{t+1} &\doteq \mathbf{x}_t - \gamma_i \nabla_i f(\mathbf{x}_t) \mathbf{e}_i. \end{aligned}$$

The main difference from the previous algorithms is that this algorithm is deterministic, thus we do not need to take the expectation.

Corollary. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable with a global minimum \mathbf{x}^* . Suppose that f is coordinate-wise smooth with parameter L and satisfies the PL inequality with parameter $\mu > 0$. Choosing stepsize

$$\gamma_i \doteq \frac{1}{L},$$

steepest coordinate descent with arbitrary \mathbf{x}_0 yields

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) \leq \left(1 - \frac{\mu}{dL}\right)^T (f(\mathbf{x}_0) - f(\mathbf{x}^*)).$$

This is not good. It needs the same amount of iterations as randomized coordinate descent, but each iteration takes as long as in gradient descent.³ However, this algorithm allows for a speedup in certain cases. Furthermore, it may be possible to efficiently maintain the maximum absolute gradient value throughout the iterations, so that the full evaluation of the gradient can be avoided.

Nutini et al. [2015] showed that a better convergence result can be obtained for strongly convex functions, when strong convexity is measured w.r.t. the ℓ_1 norm instead of the standard Euclidean norm, *i.e.*,

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{\mu_1}{2} \|\mathbf{y} - \mathbf{x}\|_1^2.$$

Due to $\|\mathbf{y} - \mathbf{x}\|_1 \geq \|\mathbf{y} - \mathbf{x}\|$, f is then also strongly convex with $\mu = \mu_1$. On the other hand, if f is μ -strongly convex w.r.t. the ℓ_2 norm, then f is μ/d -strongly convex w.r.t. the ℓ_1 norm, due to $\|\mathbf{y} - \mathbf{x}\| \geq \|\mathbf{y} - \mathbf{x}\|_1 / \sqrt{d}$.

Lemma 46. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable and strongly convex with parameter $\mu_1 > 0$ w.r.t. the ℓ_1 norm. Then, f is μ_1 -strongly convex w.r.t. the Euclidean norm, so a global minimum \mathbf{x}^* exists. Then, f satisfies the PL inequality w.r.t. the ℓ_∞ norm with the same μ_1 ,

$$\frac{1}{2} \|\nabla f(\mathbf{x})\|_\infty^2 \geq \mu_1 (f(\mathbf{x}) - f(\mathbf{x}^*)).$$

Theorem 47. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable with a global minimum \mathbf{x}^* . Suppose that f is coordinate-wise smooth with parameter L and satisfies the PL inequality with parameter $\mu_1 > 0$. Choosing stepsize

$$\gamma_i \doteq \frac{1}{L},$$

steepest coordinate descent with arbitrary \mathbf{x}_0 yields

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) \leq \left(1 - \frac{\mu_1}{L}\right)^T (f(\mathbf{x}_0) - f(\mathbf{x}^*)).$$

³Note that a function may be coordinate-wise smooth with an L for all coordinates that is smaller than smoothness, so it is not completely fair to compare this to gradient descent.

Proof.

$$\begin{aligned}
 f(\mathbf{x}_{t+1}) &\leq f(\mathbf{x}_t) - \frac{1}{2L} |\nabla_i f(\mathbf{x}_t)|^2 \\
 &= f(\mathbf{x}_t) - \frac{1}{2L} \|\nabla f(\mathbf{x}_t)\|_\infty^2 \\
 &\leq f(\mathbf{x}_t) - \frac{\mu_1}{L} (f(\mathbf{x}_t) - f(\mathbf{x}^*)).
 \end{aligned}$$

Subtracting $f(\mathbf{x}^*)$ from both sides yields

$$f(\mathbf{x}_{t+1}) - f(\mathbf{x}^*) \leq \left(1 - \frac{\mu_1}{L}\right) (f(\mathbf{x}_t) - f(\mathbf{x}^*)).$$

The statement follows. ■

We see that if $\mu_1 = \mu/d$, we do not gain anything. However, this is not the case in general. If, for the worst case \mathbf{x}, \mathbf{y} , which satisfy $\|\mathbf{y} - \mathbf{x}\| = \|\mathbf{y} - \mathbf{x}\|_1 / \sqrt{d}$, strong convexity holds with $\mu' > \mu$, then we can achieve $\mu_1 = \mu'/d > \mu/d$, resulting in better convergence.

5.4 Greedy coordinate descent

Greedy coordinate descent is a variant that does not even require f to be differentiable. In each iteration, we make the step that maximizes the progress in the chosen coordinate. This requires performing a *line search* by solving a one-dimensional optimization problem,

$$\begin{aligned}
 &\text{choose } i \in [d] \\
 &\mathbf{x}_{t+1} \doteq \underset{\lambda \in \mathbb{R}}{\operatorname{argmin}} f(\mathbf{x}_t + \lambda \mathbf{e}_i).
 \end{aligned}$$

However, greedy coordinate descent can get stuck in non-optimal points; see Figure 16. Thus, we need some additional conditions to make sure this does not occur.

Theorem 48. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be of the form

$$f(\mathbf{x}) \doteq g(\mathbf{x}) + h(\mathbf{x}), \quad h(\mathbf{x}) = \sum_{i=1}^d h_i(x_i), \quad \mathbf{x} \in \mathbb{R}^d,$$

with g convex and differentiable, and the h_i convex.

Let \mathbf{x} be a point such that greedy coordinate descent cannot make progress in any coordinate. Then \mathbf{x} is a global minimum of f .

In the context of machine learning, an important class of functions that satisfies the conditions of ?? is the following form,

$$f(\mathbf{x}) + \lambda \|\mathbf{x}\|_1,$$

where $\lambda \|\mathbf{x}\|_1$ is a separable ℓ_1 -regularization term used in for example LASSO [Tibshirani, 1996].

Sufficient decrease.

Active coordinate of steepest coordinate descent is the maximum gradient.

PL inequality.




Figure 16. Level set plot of $f(\mathbf{x}) = \|\mathbf{x}\|^2 + |x_1 - x_2|$. The global minimum is $[0, 0]$, but greedy coordinate descent cannot escape any point $[\mathbf{x}, \mathbf{x}]$, s.t. $|\mathbf{x}| \leq 1/2$.

6 Nonconvex functions

So far, all convergence results that we have proved have been for variants of gradient descent on convex functions. The reason for this is that, in general, we cannot expect gradient descent to come close to the global minimum x^* of nonconvex functions. Figures 17 to 19 show what can go wrong in nonconvex functions, under the assumption that we have set the γ such that we do not overshoot. These figures show points that gradient descent cannot escape; local minima and saddle points. Furthermore, it might even be that gradient descent never converges to a critical point; see Figure 19.

In practice, gradient descent works well on the nonconvex functions that we care about. But, theoretical explanations for this are mostly missing. Despite this, we will show that under favorable conditions, we can still say something useful about the behavior of gradient descent on nonconvex functions.

We can easily make an analysis of gradient descent on smooth functions. A useful property that we will use is that functions with bounded Hessians are smooth, as shown in the following lemma.

Lemma 49. Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ be twice differentiable with $\mathcal{X} \in \text{dom}(f)$ a convex set and $\|\nabla^2 f(x)\| \leq L$ for all $x \in \mathcal{X}$. Then f is smooth with parameter L over \mathcal{X} .

Proof. Bounded Hessians imply Lipschitz continuity of the gradient,

$$\|\nabla f(x) - \nabla f(y)\| \leq L\|x - y\|, \quad \forall x, y \in \mathcal{X}.$$

We will use the fundamental theorem of calculus with

$$h(t) \doteq f(x + t(y - x)), \quad t \in [0, 1].$$

The derivative can be calculated by chain rule,

$$h'(t) = \nabla f(x + t(y - x))^\top (y - x).$$




Figure 17. Gradient descent may get stuck in a local minimum $y^* \neq x^*$, in nonconvex functions.




Figure 18. Gradient may get stuck in a flat region (saddle point) in nonconvex functions.




Figure 19. Gradient descent may never even reach a critical point in nonconvex functions.

Now, we can show smoothness,

$$\begin{aligned}
& f(\mathbf{y}) - f(\mathbf{x}) - \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) \\
&= h(1) - h(0) - \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) && \text{Definition of } h. \\
&= \int_0^1 h'(t) dt - \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) && \text{Fundamental theorem of calculus.} \\
&= \int_0^1 \nabla f(\mathbf{x} + t(\mathbf{y} - \mathbf{x}))^\top (\mathbf{y} - \mathbf{x}) dt - \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) && \text{Fill in } h'(t). \\
&= \int_0^1 \nabla f(\mathbf{x} + t(\mathbf{y} - \mathbf{x}))^\top (\mathbf{y} - \mathbf{x}) - \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt && \text{Integral from 0 to 1.} \\
&= \int_0^1 (\nabla f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) - \nabla f(\mathbf{x}))^\top (\mathbf{y} - \mathbf{x}) dt \\
&\leq \int_0^1 |(\nabla f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) - \nabla f(\mathbf{x}))^\top (\mathbf{y} - \mathbf{x})| dt \\
&\leq \int_0^1 \|\nabla f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) - \nabla f(\mathbf{x})\| \|\mathbf{y} - \mathbf{x}\| dt && \text{Cauchy-Schwarz inequality.} \\
&\leq \int_0^1 L \|t(\mathbf{y} - \mathbf{x})\| \|\mathbf{y} - \mathbf{x}\| dt && \text{Lipschitz continuous gradient.} \\
&= \int_0^1 Lt \|\mathbf{x} - \mathbf{y}\|^2 dt \\
&= \frac{L}{2} \|\mathbf{x} - \mathbf{y}\|^2.
\end{aligned}$$

Thus, we have smoothness,

$$f(\mathbf{y}) \leq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{L}{2} \|\mathbf{x} - \mathbf{y}\|^2.$$

■

Now, we can use this fact and sufficient decrease⁴ to prove that the gradients of smooth functions are bounded and approach 0, as we increase the number of iterations.

⁴ Recall that sufficient decrease did not require convexity.

Theorem 50. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be differentiable with a global minimum \mathbf{x}^* . Furthermore, suppose that f is smooth with parameter L . Choosing stepsize

$$\gamma \doteq \frac{1}{L},$$

gradient descent yields

$$\frac{1}{T} \sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 \leq \frac{2L}{T} (f(\mathbf{x}_0) - f(\mathbf{x}^*)).$$

Remark. Note that concave functions are not a counter example to this theorem, despite their gradients growing, because they have no global minimum \mathbf{x}^* .

Proof. Recall that sufficient decrease does not require convexity,

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2L} \|\nabla f(\mathbf{x}_t)\|^2.$$

Rewriting this, we get

$$\|\nabla f(\mathbf{x}_t)\|^2 \leq 2L(f(\mathbf{x}_t) - f(\mathbf{x}_{t+1})).$$

Then, by telescoping sum, we get

$$\sum_{t=0}^{T-1} \|\nabla f(\mathbf{x}_t)\|^2 \leq 2L(f(\mathbf{x}_0) - f(\mathbf{x}_T)) \leq 2L(f(\mathbf{x}_0) - f(\mathbf{x}^*)).$$

The statement follows by dividing both sides by T . ■

This has the result that

$$\lim_{t \rightarrow \infty} \|\nabla f(\mathbf{x}_t)\|^2 = 0.$$

It might seem that convergence of the gradients to 0 is the same as convergence to a critical point. But, this interpretation does not hold in general; see Figure 19. In this case, the gradient converges to 0, but the iterates only move further away from the critical point. So, this is not a very strong result.

6.1 Trajectory analysis

Despite the fact that a nonconvex function may contain local minima, saddle points, and flat parts, gradient descent may avoid them and still converge to a global minimum. For this, you need a good starting point and do a trajectory analysis. As an example, we will do a trajectory analysis for a simplified deep linear neural network. It turns out that this function is smooth along the trajectories that we analysis, and this is the most important ingredient of the analysis.

Let

$$\theta = \{\mathbf{W}_1, \dots, \mathbf{W}_\ell\}$$

be the weights of the deep linear network. And, in general, we want to approximate a matrix \mathbf{Y} , given input matrix \mathbf{X} . Thus, we want to minimize

$$\|\mathbf{W}_\ell \mathbf{W}_{\ell-1} \cdots \mathbf{W}_1 \mathbf{X} - \mathbf{Y}\|_F^2.$$

Arora et al. [2018] consider this general framework, but we will only consider the case where all matrices are 1×1 , *i.e.*, scalars. Assume we have training input $x = 1$ and output $y = 1$, then we have the following function to optimize,

$$f(\mathbf{x}) \doteq \frac{1}{2} \left(\prod_{k=1}^d x_k - 1 \right)^2.$$

We can immediately see that setting $x_k = 1$ for all k minimizes the function at 0. However, we want to know whether gradient descent will also be able to find this set of weights.

Note that stacking linear layers has no benefit, since any stacking of linear layers can be represented by a single linear layer. However, the reason for doing this is that it gives us a simple playground in which we can try to understanding why training deep neural networks with gradient descent works, despite the fact that the objective function is nonconvex.

We rewrite w as x and ℓ as d to be more in line with the notation used here.

The gradient of this function is computed by

$$\nabla_i f(\mathbf{x}) = \left(\prod_{k=1}^d x_k - 1 \right) \prod_{k \neq i}^d x_k$$

Whenever at least two dimensions are zero, the gradient vanishes. Thus, any \mathbf{x} with two zero entries are critical points, despite not being global minima, since then the product of all entries must be 1, which is not possible if at least two are zero.⁵ We know that the value of all such saddle points is $1/2$.

We now want to show that for any number of layers (*i.e.* dimensionality of \mathbf{x}), anywhere in $\mathcal{X} = \{\mathbf{x} \mid \mathbf{x} > \mathbf{0}, \prod_{k=1}^d x_k \leq 1\}$, despite that f is not smooth over \mathcal{X} . However, we only need to show that f is smooth along the trajectory of gradient descent for suitable L , so that we get sufficient decrease. We will now show this by showing that the Hessians over the trajectory are bounded. The Hessian is given by

$$\nabla_{ij}^2 f(\mathbf{x}) = \begin{cases} \left(\prod_{k \neq i}^d x_k \right)^2, & j = i \\ 2 \prod_{k \neq i}^d x_k \prod_{k \neq j}^d x_k - \prod_{k \neq i, j}^d x_k, & j \neq i. \end{cases}$$

Definition 51 (*c*-balanced.). Let $\mathbf{x} > \mathbf{0}$ and $c \geq 1$. \mathbf{x} is called *c*-balanced if $x_i \leq cx_j$ for all $1 \leq i, j \leq d$.

Lemma 52. Let $\mathbf{x} > \mathbf{0}$ be *c*-balanced with $\prod_k x_k \leq 1$, then for any stepsize $\gamma > 0$, $\mathbf{x}' \doteq \mathbf{x} - \gamma \nabla f(\mathbf{x})$ satisfies $\mathbf{x}' \geq \mathbf{x}$ componentwise, and is also *c*-balanced.

Proof. Let

$$\Delta \doteq -\gamma \left(\prod_{k=1}^d x_k - 1 \right) \left(\prod_{k=1}^d x_k \right) \geq 0.$$

Then,

$$-\gamma \nabla_k f(\mathbf{x}) = \frac{\Delta}{x_k}.$$

Thus, the gradient descent update has the following form,

$$x'_k = x_k + \frac{\Delta}{x_k} \geq x_k, \quad k \in [d].$$

For all i, j , we thus get

$$x'_i = x_i + \frac{\Delta}{x_i} \leq cx_j + \frac{c\Delta}{x_j} = cx'_j.$$

■

So, we know now that all iterates are *c*-balanced if \mathbf{x}_0 is balanced with *c*. We can use this to compute a bound on the Hessian by bounding the products.

⁵ This shows that f is nonconvex, since local minima are global minima in convex functions.




Figure 20. $f(\mathbf{x}) = \frac{1}{2} (\prod_k x_k - 1)^2$ for $d = 2$, where the loss is clipped to be at most $1/2$.

$$x_j \leq cx_i \iff \frac{1}{x_i} \leq \frac{c}{x_j}.$$

Lemma 53. Suppose $\mathbf{x} > \mathbf{0}$ is c -balanced. Then, for any $I \subseteq [d]$, we have

$$\prod_{k \notin I} x_k \leq c^{|I|} \left(\prod_{k=1}^d x_k \right)^{1-|I|/d} \leq c^{|I|}.$$

Proof. For any $i \in [d]$, we have $c^d \cdot x_i^d \geq \prod_k x_k$ by c -balancedness, hence $x_i \geq \frac{1}{c} (\prod_k x_k)^{1/d}$. Thus,

$$\prod_{k \notin I} x_k = \frac{\prod_k x_k}{\prod_{i \in I} x_i} \leq \frac{\prod_k x_k}{(1/c)^{|I|} (\prod_k x_k)^{|I|/d}} = c^{|I|} \left(\prod_{k=1}^d x_k \right)^{1-|I|/d}.$$

Since $\prod_k x_k \leq 1$, we can bound the above by $c^{|I|}$. ■

Lemma 54. Let $\mathbf{x} > \mathbf{0}$ be c -balanced with $\prod_k x_k \leq 1$, then

$$\|\nabla^2 f(\mathbf{x})\| \leq \|\nabla^2 f(\mathbf{x})\|_F \leq 3dc^2.$$

Proof. The fact that $\|A\| \leq \|A\|_F$ is well known. To bound the Frobenius norm, we use the previous lemma to compute

$$|\nabla_{ii}^2 f(\mathbf{x})| = \left| \left(\prod_{k \neq i} x_k \right)^2 \right| \leq c^2,$$

and for $i \neq j$, we get

$$|\nabla_{ij}^2 f(\mathbf{x})| \leq \left| 2 \prod_{k \neq i} x_k \prod_{k \neq j} x_k \right| + \left| \prod_{k \neq i, j} x_k \right| \leq 3c^2.$$

Thus,

$$\|\nabla^2 f(\mathbf{x})\|_F^2 \leq 9d^2 c^4.$$

Taking the square root, the statement follows. ■

This lemma implies smoothness of f with parameter $L = 3dc^2$ along the whole trajectory of gradient descent, under the “smooth stepsize” $\gamma \doteq 1/L = 1/3dc^2$. Now, we can use this to prove convergence.

Theorem 55. Let $c \geq 1$ and $\delta < 0$ such that $\mathbf{x}_0 > \mathbf{0}$ is c -balanced with $\delta \leq \prod_k (\mathbf{x}_0)_k < 1$. Choosing stepsize

$$\gamma \doteq \frac{1}{3dc^2},$$

gradient descent satisfies

$$f(\mathbf{x}_T) \leq \left(1 - \frac{\delta^2}{3c^4} \right)^T f(\mathbf{x}_0).$$

Proof. For each $t \geq 0$, f is smooth over $\text{conv}(\{\mathbf{x}_t, \mathbf{x}_{t+1}\})$ with parameter $L = 3dc^2$, hence we have sufficient decrease,

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{6dc^2} \|\nabla f(\mathbf{x}_t)\|^2.$$

For every c -balanced \mathbf{x} with $\delta \leq \prod_k x_k \leq 1$, we have

$$\begin{aligned} \|\nabla f(\mathbf{x}_t)\|^2 &= 2f(\mathbf{x}) \sum_{i=1}^d \left(\prod_{k \neq i} x_k \right)^2 \\ &\geq 2f(\mathbf{x}) \frac{d}{c^2} \left(\prod_{k=1}^d x_k \right)^{2-2/d} \\ &\geq 2f(\mathbf{x}) \frac{d}{c^2} \left(\prod_{k=1}^d x_k \right)^2 \\ &\geq 2f(\mathbf{x}) \frac{d}{c^2} \delta^2. \end{aligned}$$

Hence,

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{6dc^2} 2f(\mathbf{x}_t) \frac{d}{c^2} \delta^2 = f(\mathbf{x}_t) \left(1 - \frac{\delta^2}{3c^4} \right).$$

■

Thus, we seem to have fast convergence, since the function value goes down by a constant factor in each step. However, there is a catch. Consider the $\mathbf{x}_0 = [1/2, \dots, 1/2]$, which is c -balanced with $c = 1$, and $\delta = 1/2^d$. Hence, the constant factor is

$$1 - \frac{1}{3 \cdot 4^d}.$$

This means that we would need $T \approx 4^d$ iterations to reduce the initial error by a constant factor not depending on d . Hence, for this starting value, the gradient is exponentially small. In order to get polynomial convergence, we need to start with a δ that decays at most polynomially with d . For large d , this has the consequence that we must start very close to optimality. In particular, we need to start at a distance $\mathcal{O}(1/\sqrt{d})$ from the optimal solution $[1, \dots, 1]$.

References

- Sanjeev Arora, Nadav Cohen, Noah Golowich, and Wei Hu. A convergence analysis of gradient descent for deep linear neural networks. *arXiv preprint arXiv:1810.02281*, 2018.
- Yu Nesterov. Efficiency of coordinate descent methods on huge-scale optimization problems. *SIAM Journal on Optimization*, 22(2):341–362, 2012.
- Julie Nutini, Mark Schmidt, Issam Laradji, Michael Friedlander, and Hoyt Koepke. Coordinate descent converges faster with the gauss-southwell rule than random selection. In *International Conference on Machine Learning*, pages 1632–1641. PMLR, 2015.
- Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1): 267–288, 1996.