

Accelerating role of deaggregation agents in lithium-catalysed hydrosilylation of carbonyl compounds

Cristina Ruiz,^{†,¶} Álvaro Raya-Barón,^{†,¶} Manuel A. Ortuño,^{*,†} and Ignacio Fernández^{*,‡}

[†]*Institute of Chemical Research of Catalonia (ICIQ), The Barcelona Institute of Science and Technology (BIST), Av. Països Catalans 16, 43007 Tarragona, Spain.*

[‡]*Department of Chemistry and Physics, Research centre CIAIMBITAL, Ctra. Sacramento, s/n, 04120 Almería, Spain.*

[¶]*Both authors have equally contributed*

E-mail: mortuno@iciq.es; ifernan@ual.es

Abstract

A combined computational and experimental approach demonstrates the accelerating role of deaggregation agents, especially HMPA, in the Li-catalysed hydrosilylation of acetophenone in THF solution under very mild conditions.

Introduction

The reduction of carbonyl groups into alcohols is of wide interest in synthetic chemistry and therefore, constant efforts are put into developing new efficient methodologies and perfecting the existing ones. Catalytic hydrosilylation¹ has emerged as a convenient method as it operates

under mild conditions and combines an exceptional reducing capability with a high selectivity that can be finely tuned via catalyst design.

State of the art

Many Earth-abundant first-row transition metals, specially iron,^{2,3} have been tested in catalytic hydrosilylation of carbonyl compounds, as they are usually more environmentally-friendly and less toxic than their second- and third-row counterparts. Alkali metal salts have also been explored as alternative catalysts, initially by the groups of Corriu⁴ and Hosomi,⁵ and later by Beller⁶ and Nikonov,⁷ among others. Such compounds have been employed to promote hydrosilylation due to their basic character via formation of a pentacoordinated hydridosilicate,^{6–8} usually neglecting any relevant role of the alkali cation in the reaction mechanism. We recently reported the hydrosilylation of carbonyl compounds catalysed by lithiated hydrazones.⁹ However, full understanding at atomic level of detail is still needed for the rational design of catalysts and reaction conditions. Herein we join computational and experimental efforts to understand and optimise processes catalysed by alkali-metal amides.¹⁰ Following theoretical guidance, we demonstrate how deaggregation agents (DAs) can efficiently accelerate hydrosilylation of carbonyl compounds in the presence of readily available lithium amides (Scheme 1) under very mild conditions such as room temperature and very low catalyst loading.




Figure 1: Hydrosilylation of ketones catalysed by the combination of Li-amides with deaggregation agents

Results and discussion

First, we computed the reaction mechanism for acetophenone and (MeO)₂MeSiH using lithium diisopropylamide (LDA) as catalyst.⁹ The proposed reaction mechanism entails an activation

step (Scheme 2a) prior to the catalytic cycle (Scheme 2b). We start from the LDA dimer 1 in THF solution. Dissociation of 111 via coordination of carbonyl and silane reactants yields the monomeric intermediates 2 and 3 at 17.8 kcal mol⁻¹. The Si-H bond is then activated via amide nucleophilic attack,¹² giving rise to a putative Li-hydride 5 as in Fe analogues ^{13,14} via -25 kcal mol⁻¹. Although such transient intermediate would be very difficult to detect experimentally, similar species have been proposed in related literature. ^{15,16} Insertion of acetophenone into the Li-H bond generates an alkoxide 6 which activates a silane molecule 8 through a pentacoordinated hydridosilicate. Subsequent hydride transfer releases the product and regenerates the Li-H