
# THE NEED FOR SPEED

## Finding Fast Transients with TESS


**Clarinda Montilla**[1], Ryan Ridden-Harper[1], Michael Albrow[1], Anais Moller[2,3], Armin Rest[4], Hugh Roxburgh[5], James Freeburn[6], Jefferey Cooke[2,3] & Qinan Wang[7]

For the first time ever, it is possible to obtain detailed light curves of fast optical transients – cosmic explosions with lifetimes lasting from just a few seconds to a few hours – thanks to a plethora of data provided by a range of telescopes. The unique high cadence observations of TESS allow us to observe transients during the rise, and the decay after peak brightness. Pairing these with the multi-band observations from ground-based telescopes, we can better identify transient type and constrain its physical properties.




Observations are dependent on the **integration time**, which determines how much light is received. A longer integration time will allow us to see fainter objects, but results in poorer time resolution.

If we consider integration time when fitting our models to data, we can get a more *realistic* picture of the event – often resulting in significantly higher peak fluxes and energies.




By fitting to a GRB model which has been convolved with the time integration window, we can recover the model flux and its physical parameters at an **infinite time resolution**. These precise parameters, including redshift and the peak emitted energy can give us insights into the underlying physics of these transient events.

| | 2020har | 2022eqe | 2024zuk |
|---|---|---|---|
| $log_{10}E_0$ (ergs) | $53.53^{+0.12}_{-0.19}$ | $54.79^{+2.19}_{-1.76}$ | $54.24^{+0.09}_{-0.14}$ |
| $z$ | $0.63^{+0.10}_{-0.05}$ | $0.80^{+2.29}_{-0.78}$ | $0.42^{+0.05}_{-0.04}$ |




UNIVERSITY OF CANTERBURY
Te Whare Wānanga o Waitaha
CHRISTCHURCH NEW ZEALAND


[1] University of Canterbury, [2] Swinburne University of Technology, [3] OzGrav,
[4] Space Telescope Science Institute, [5] Curtin University, [6] University of North Carolina,
[7] Massachusetts Institute of Technology

# THE NEED FOR SPEED

## Finding Fast Transients with TESS

**Clarinda Montilla**[1], Ryan Ridden-Harper[1], Michael Albrow[1], Anais Moller[2,3], Armin Rest[4], Hugh Roxburgh[5], James Freeburn[6], Jefferey Cooke[2,3] & Qinan Wang[7]

For the first time ever, it is possible to obtain detailed light curves of fast optical transients – cosmic explosions with lifetimes lasting from just a few seconds to a few hours – thanks to a plethora of data provided by a range of telescopes. The unique high cadence observations of TESS allow us to observe transients during the rise, and the decay after peak brightness. Pairing these with the multi-band observations from ground-based telescopes, we can better identify transient type and constrain its physical properties.




Observations are dependent on the **integration time**, which determines how much light is received. A longer integration time will allow us to see fainter objects, but results in poorer time resolution.

If we consider integration time when fitting our models to data, we can get a more realistic picture of the event – often resulting in significantly higher peak fluxes and energies.



By fitting to a GRB model which has been convolved with the time integration window, we can recover the model flux and its physical parameters at an **infinite** time resolution. These precise parameters, including redshift and the peak emitted energy can give us insights into the underlying physics of these transient events.

| | 2020har | 2022eqe | 2024zuk |
|---|---|---|---|
| z | 0.63 | 0.80 | 0.42 |


[1] University of Canterbury, [2] Swinburne University of Technology, [3] OzGrav, [4] Space Telescope Science Institute, [5] Curtin University, [6] University of North Carolina, [7] Massachusetts Institute of Technology

# THE NEED FOR SPEED

## Finding Fast Transients with TESS


**Clarinda Montilla**[1], Ryan Ridden-Harper[1], Michael Albrow[1], Anais Moller[2,3], Armin Rest[4], Hugh Roxburgh[5], James Freeburn[6], Jefferey Cooke[2,3] & Qinan Wang[7]

For the first time ever, it is possible to obtain detailed light curves of fast optical transients – cosmic explosions with lifetimes lasting from just a few seconds to a few hours – thanks to a plethora of data provided by a range of telescopes. The unique high cadence observations of TESS allow us to observe transients during the rise, and the decay after peak brightness. Pairing these with the multi-band observations from ground-based telescopes, we can better identify transient type and constrain its physical properties.




Observations are dependent on the **integration time**, which determines how much light is received. A longer integration time will allow us to see fainter objects, but results in poorer time resolution.

If we consider integration time when fitting our models to data, we can get a more realistic picture of the event – often resulting in significantly higher peak fluxes and energies.




26
95

# THE NEED FOR SPEED

## Finding Fast Transients with TESS


**Clarinda Montilla**[1], Ryan Ridden-Harper[1], Michael Albrow[1], Anais Moller[2,3], Armin Rest[4], Hugh Roxburgh[5], James Freeburn[6], Jefferey Cooke[2,3] & Qinan Wang[7]


For the first time ever, it is possible to obtain detailed light curves of fast optical transients – cosmic explosions with lifetimes lasting from just a few seconds to a few hours – thanks to a plethora of data provided by a range of telescopes. The unique high cadence observations of TESS allow us to observe transients during the rise, and the decay after peak brightness. Pairing these with the multi-band observations from ground-based telescopes, we can better identify transient type and constrain its physical properties.





Observations are dependent on the **integration time**, which determines how much light is received. A longer integration time will allow us to see fainter objects, but results in poorer time resolution.

If we consider integration time when fitting our models to data, we can get a more realistic picture of the event – often resulting in significantly higher peak fluxes and energies.







# THE NEED FOR SPEED

## Finding Fast Transients with TESS


**Clarinda Montilla**[1], Ryan Ridden-Harper[1], Michael Albrow[1], Anais Moller[2,3], Armin Rest[4], Hugh Roxburgh[5], James Freeburn[6], Jefferey Cooke[2,3] & Qinan Wang[7]


For the first time ever, it is possible to obtain detailed light curves of fast optical transients – cosmic explosions with lifetimes lasting from just a few seconds to a few hours – thanks to a plethora of data provided by a range of telescopes. The unique high cadence observations of TESS allow us to observe transients during the rise, and the decay after peak brightness. Pairing these with the multi-band observations from ground-based telescopes, we can better identify transient type and constrain its physical properties.





Observations are dependent on the **integration time**, which determines how much light is received. A longer integration time will allow us to see fainter objects, but results in poorer time resolution.

If we consider integration time when fitting our models to data, we can get a more realistic picture of the event – often resulting in significantly higher peak fluxes and energies.







# THE NEED FOR SPEED

## Finding Fast Transients with TESS


**Clarinda Montilla**[1], Ryan Ridden-Harper[1], Michael Albrow[1], Anais Moller[2,3], Armin Rest[4], Hugh Roxburgh[5], James Freeburn[6], Jefferey Cooke[2,3] & Qinan Wang[7]


For the first time ever, it is possible to obtain detailed light curves of fast optical transients – cosmic explosions with lifetimes lasting from just a few seconds to a few hours – thanks to a plethora of data provided by a range of telescopes. The unique high cadence observations of TESS allow us to observe transients during the rise, and the decay after peak brightness. Pairing these with the multi-band observations from ground-based telescopes, we can better identify transient type and constrain its physical properties.





Observations are dependent on the **integration time**, which determines how much light is received. A longer integration time will allow us to see fainter objects, but results in poorer time resolution.

If we consider integration time when fitting our models to data, we can get a more realistic picture of the event – often resulting in significantly higher peak fluxes and energies.







# THE NEED FOR SPEED

**Clarinda Montilla**
Dr. Ryan Ridden-Harper & Associate Professor Michael Albrow

For the first time ever, it is possible to obtain detailed light curves of fast optical transients – cosmic explosions with lifetimes lasting from just a few seconds to a few hours – thanks to a plethora of data provided by a range of telescopes. The unique high cadence observations of TESS allow us to observe transients during the rise, and the decay after peak brightness. Pairing these with the multi-band observations from ground-based telescopes, we can better identify transient type and constrain its physical properties.




Observations are dependent on the **integration time**, which determines how much light is received. A longer integration time will allow us to see fainter objects, but results in poorer time resolution.

grb fit/s

# THE NEED FOR SPEED

Clarinda Montilla
Dr. Ryan Ridden-Harper & Associate Professor Michael Albrow

Transients are elusive phenomena, living for a brief moment before disappearing forever.
To understand the physics of these explosions, we must get as many observations as possible so that we can study these events in detail – but *how* do you study an event with such a short lifetime?

# THE NEED FOR SPEED

Clarinda Montilla
Dr. Ryan Ridden-Harper & Associate Professor Michael Albrow

Transients are elusive phenomena, living for a brief moment before disappearing forever.
To understand the physics of these explosions, we must get as many observations as possible so that we can study these events in detail – but *how* do you study an event with such a short lifetime?

# THE NEED FOR SPEED

Clarinda Montilla
Dr. Ryan Ridden-Harper & Associate Professor Michael Albrow

Transient are elusive phenomena, living for a brief moment before disappearing forever.
To understand the physics of these explosions, we must get as many observations as possible so that we can study these events in detail – but *how* do you study an event with such a short lifetime?

# THE NEED FOR SPEED

Clarinda Montilla
Dr. Ryan Ridden-Harper & Associate Professor Michael Albrow

Transient are elusive phenomena, living for a brief moment before disappearing forever.
To understand the physics of these explosions, we must get as many observations as possible so that we can study these events in detail – but *how* do you study an event with such a short lifetime?

# THE NEED FOR SPEED

Clarinda Montilla
Dr. Ryan Ridden-Harper & Associate Professor Michael Albrow

Transient are elusive phenomena, living for a brief moment before disappearing forever.
To understand the physics of these explosions, we must get as many observations as possible so that we can study these events in detail – but *how* do you study an event with such a short lifetime?

# THE NEED FOR SPEED

Clarinda Montilla
Dr. Ryan Ridden-Harper & Associate Professor Michael Albrow

Transient are elusive phenomena, living for a brief moment before disappearing forever.
To understand the physics of these explosions, we must get as many observations as possible so that we can study these events in detail – but *how* do you study an event with such a short lifetime?