

ASCII MASTER FLOW – PANEL 1/12: Eastern-margin world map

CHINA -> central and north China + southern Japan: temperate monsoonal

SE UNITED STATES -> Gulf type: slight-monsoonal, Gulf of Mexico coast

NATAL + E AUSTRALIA + SE S. AMERICA -> non-monsoonal southern hemisphere

WESTERN MARGIN (same latitude) -> Mediterranean: summer drought contrast

MUST REMEMBER: Warm temperate eastern-margin China type combines humid subtropical latitude, onshore summer flow, convection/cyclones and winter continental influence, producing hot wet summers, cooler winters and mixed crops/forests in East Asia and analogous margins.

MUST REMEMBER: Warm temperate eastern-margin China type combines humid subtropical latitude, onshore summer flow, convection/cyclones and winter continental influence, producing hot wet summers, cooler winters and mixed crops/forests in East Asia and analogous margins.

ASCII MASTER FLOW – PANEL 2/12: Pressure-reversal mechanism

SUMMER -> intense heating over Asian interior -> low pressure

INFLOW -> tropical Pacific air stream drawn onshore -> summer rain

WINTER -> cooling over continental mass -> high pressure

OUTFLOW -> cold dry air moves offshore -> drier, cooler winter

ASCII MASTER FLOW – PANEL 3/12: Three sub-types comparison

CHINA TYPE -> temperate monsoonal, harshest winter (large Asian landmass)

GULF TYPE -> slight-monsoonal, SE USA, moderate winter

NATAL TYPE -> non-monsoonal, southern hemisphere, mildest winter

CAUSE -> winter severity depends on size of adjoining continental mass

ASCII MASTER FLOW – PANEL 4/12: East-west rainfall asymmetry

EASTERN MARGIN -> summer rain from onshore flow, no summer drought
WESTERN MARGIN -> summer drought under subtropical high, winter rain
MECHANISM -> same latitude, opposite pressure-wind regime
HIGHEST-VALUE COMPARISON -> in the warm temperate latitude belt

ASCII MASTER FLOW – PANEL 5/12: Typhoon track map

ORIGIN -> warm western Pacific Ocean, low-latitude genesis

TRACK -> westward toward South China Sea coasts

SEASON -> late summer, July to September peak

IMPACT -> wind, surge and flood on densely settled coastal plains

ASCII MASTER FLOW – PANEL 6/12: Agriculture-hazard duality

SUMMER INFLOW -> warm moist air gives long wet growing season

RESULT -> rice, tea, sugar-cane, maize: dense rural population

SAME MECHANISM -> brings typhoons, floods and surges

PARADOX -> richness and hazard arise from the same maritime source

ASCII MASTER FLOW – PANEL 7/12: India humid eastern-margin map

BENGAL PLAIN -> West Bengal plain, about 150 cm Bay-of-Bengal monsoon

BRAHMAPUTRA VALLEY -> Assam, hot sub-humid to perhumid

NE INDIA (R.L. Singh) -> NE except Tripura, Sikkim, NW W. Bengal

RAINFALL -> above 200 cm in Humid NE; July 25-33 C, Jan 10-25 C

ASCII MASTER FLOW – PANEL 8/12: Monsoon-branch discriminator

QUESTION: which monsoon branch serves NE India?

BAY OF BENGAL BRANCH -> correct for Bengal-Brahmaputra region

ARABIAN SEA BRANCH -> serves western coast, NOT NE India

EXAM TRAP -> this is a standard close-option Prelims discriminator

ASCII MASTER FLOW – PANEL 9/12: Flood-fertility paradox

HEAVY MONSOON RAIN -> annual flooding of Brahmaputra floodplain

POSITIVE -> alluvial fertility renews rice-tea economy each year

NEGATIVE -> bank erosion, displacement, crop loss, infrastructure damage

CONCLUSION -> flood management must preserve fertility, not exclude water

ASCII MASTER FLOW – PANEL 10/12: Tea-industry vulnerability

ASSAM -> central to India's tea output, world's 2nd-largest producer

RISK -> erratic rainfall, drought, flood: second-flush crop losses

CAUSE -> humid eastern-margin monsoon regime variability

SIGNIFICANCE -> climate change links to NE livelihoods and export crop

CLOSE DISTINCTION: China type is an eastern-margin climate label, not China's entire climate;
it differs from tropical monsoon by winter temperature and from Laurentian by warmer
conditions and different ocean-current/air-mass relations.

CLOSE DISTINCTION: China type is an eastern-margin climate label, not China's entire climate;
it differs from tropical monsoon by winter temperature and from Laurentian by warmer
conditions and different ocean-current/air-mass relations.

ASCII MASTER FLOW – PANEL 11/12: Trewartha classification note

TREWARTHA -> humid mesothermal and tropical-humid types in India

H CLASS -> undifferentiated highland for elevated regions

NE BELT -> spans several codes depending on elevation and scheme

CAUTION -> do not force entire region into one Cwg or Amw code

EVIDENCE LIMIT: East-margin similarity does not erase monsoon, current and topographic differences; Northeast India is an anchored comparison whose altitude and exceptional rainfall prevent one-code generalisation.

EVIDENCE LIMIT: East-margin similarity does not erase monsoon, current and topographic differences; Northeast India is an anchored comparison whose altitude and exceptional rainfall prevent one-code generalisation.

ASCII MASTER FLOW – PANEL 12/12: China-type answer spine

DEFINE -> warm temperate eastern margin with summer-rain regime

COMPARE -> east vs west margin at same latitude (highest-value contrast)

EXPLAIN -> pressure reversal, sub-types, agriculture-hazard duality

APPLY -> India's Bengal-NE as the China-type analogue with flood-fertility paradox