

Table S9.1 Duration of Exposure

Safety population; double-blind period only

Treatment Group	Subjects		Exposure on treatment		
	Randomized	Dosed, n (%)	Mean, weeks (SD)	Median (min-max)	Total, patient-weeks
Placebo	42	42 (100)	16.2 (4.1)	17.1 (2.9-22.4)	680
Drug X 50 mg	22	22 (100)	15.8 (4.6)	16.8 (2.1-22.2)	348
Drug X 100 mg	22	22 (100)	16.4 (4.0)	17.5 (3.4-22.6)	361
Total	86	86 (100)	16.1 (4.2)	17.1 (2.1-22.6)	1389

Time on randomized treatment from first dose to last dose or discontinuation, whichever came first; dose interruptions are included.

Table S9.2 Summary of Discontinuations

First reason recorded per subject

Reason for discontinuation	Treatment Arm			
	Placebo (N = 42)	Drug X 50 mg (N = 22)	Drug X 100 mg (N = 22)	Total (N = 86)
	n (%)	n (%)	n (%)	n (%)
Any reason	9 (21.4%)	5 (22.7%)	6 (27.3%)	20 (23.3%)
Adverse event	2 (4.8%)	3 (13.6%)	4 (18.2%)	9 (10.5%)
Withdrawal of consent	4 (9.5%)	1 (4.5%)	1 (4.5%)	6 (7.0%)
Lost to follow-up	3 (7.1%)	1 (4.5%)	0 (0.0%)	4 (4.7%)
Lack of efficacy	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Investigator decision	0 (0.0%)	0 (0.0%)	1 (4.5%)	1 (1.2%)
Protocol deviation	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Other	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)

Subjects discontinuing for an adverse event are not re-counted under subsequent reasons; the reason rows cross-foot to the 'Any reason' total.