

Table S10.1
Shifts in Clinical Laboratory Values

Safety population; values in SI units^a

Laboratory parameter	Placebo (n = 42)	Drug X 50 mg (n = 22)	Drug X 100 mg (n = 22)
Alanine aminotransferase, U/L			
Mean (SD)	34.2 (8.1)	35.1 (9.0)	37.9 (10.4)
Subjects above ULN / below LLN	1 (2.4%)	2 (9.1%)	4 (18.2%)
Aspartate aminotransferase, U/L			
Mean (SD)	31.8 (7.6)	32.9 (8.4)	36.2 (9.9)
Subjects above ULN / below LLN	0 (0.0%)	1 (4.5%)	3 (13.6%)
Gamma-glutamyl transferase, U/L			
Mean (SD)	29.4 (11.2)	33.8 (14.6)	38.1 (16.9)
Subjects above ULN / below LLN	1 (2.4%)	3 (13.6%)	5 (22.7%)
Creatinine, umol/L			
Mean (SD)	78.1 (12.4)	79.6 (13.8)	81.9 (15.1)
Subjects above ULN / below LLN	2 (4.8%)	3 (13.6%)	4 (18.2%)
Hemoglobin, g/L			
Mean (SD)	141.2 (9.8)	140.1 (10.2)	139.4 (11.0)
Subjects above ULN / below LLN	0 (0.0%)	0 (0.0%)	1 (4.5%)
Potassium, mmol/L			
Mean (SD)	4.32 (0.28)	4.28 (0.31)	4.21 (0.33)
Subjects above ULN / below LLN	1 (2.4%)	2 (9.1%)	1 (4.5%)
Glucose, mmol/L (fasting)			
Mean (SD)	5.41 (0.62)	5.62 (0.71)	5.88 (0.83)
Subjects above ULN / below LLN	3 (7.1%)	5 (22.7%)	7 (31.8%)
INR, ratio			
Mean (SD)	1.02 (0.09)	1.04 (0.11)	1.07 (0.12)
Subjects above ULN / below LLN	0 (0.0%)	0 (0.0%)	1 (4.5%)

^a Red: $\geq 13\%$ (Drug X 100 mg) or $\geq 10\%$ (Drug X 50 mg) of subjects with confirmed central-lab abnormalities in the corresponding arm. ULN/LLN per site local ranges, re-read centrally.