

MODELLING OF COVID-19 EPIDEMIC AND INTERVENTIONS WITHIN IDP CAMPS

DR. EDUARD CAMPILLO-FUNOLLET, PhD, UNIV. OF SUSSEX (UK)

JORDAN KLEIN, MPH, PRINCETON UNIV. (USA)

DR. ALBERTO PASCUAL-GARCÍA, PhD, ETH-ZÜRICH (SWITZERLAND)

DR. JENNIFER VILLERS, PhD, PRINCETON UNIV. (USA)

CHAMSY SARKIS, PAX SYRIANA FOUNDATION

JULY 16TH, 2020

PaxSyriana
Foundation

crowdfight 
COVID-19

MOTIVATION

Between December 2019 and March 2020, 1M new IDP → Informal settlements

HIGH DENSITY and no management

~700 camps with 600 person per camp on average (log-normal)

80% family-sized tents (~7 people)

Most of the informal camps have no management.

Between December 2019 and March 2020, 1M new IDP → Informal settlements

HIGH DENSITY and no management

~700 camps with 600 person per camp on average (log-normal)

80% family-sized tents (~7 people)

Most of the informal camps have no management.

NO POSSIBLE lockdown

50% no access to electricity

External water, with 25% not have enough drinking water

External letrines, and 10% households have 1 member disable/elderly.

360% drop of Syrian pound (no access to goods) → high motility to work

Between December 2019 and March 2020, 1M new IDP → Informal settlements

HIGH DENSITY and no management

~700 camps with 600 person per camp on average (log-normal)

80% family-sized tents (~7 people)

Most of the informal camps have no management.

NO POSSIBLE lockdown

50% no access to electricity

External water, with 25% not have enough drinking water

External letrines, and 10% households have 1 member disable/elderly.

360% drop of Syrian pound (no access to goods) → high motility to work

HIGH comorbidity

10% population has chronic diseases, 17% of them have no access to medicines.

of Severe Cases -Scenario One in 8 Weeks

patients need ward beds = **2,457**

of ward COVID beds in north of syria= **1,100**




Figure 6 Scenario One predicted severe cases

The health system in NW Syria would be unable to cope by the beginning of week 8; as such, severe cases could become critical and mortality could increase

Highlights

Modelling

- Compartmental models.
- Age-structured.
- Deterministic, stochastic.
- Parameters estimated for IDPCs.

Strategies


- Self-distancing
- Shielding, lockdown
- Isolation
- Combined strategies


Main conclusions



- Remark 1
- Remark 2
- Combined strategies have a greater effect.

THE MODEL


Model









Model




Model




Population Pyramid in NW of Syria




Population Pyramid in NW of Syria



Population Pyramid in NW of Syria



Model




Parameter	Description	Value	Distribution	Reference
$1/\delta_E + 1/\delta_P$	Duration of incubation period in days	5.2 (95% CI: 4.1-7.0)	Lognormal	[1]
$1/\delta_E$	Duration of latency in days	$1/\delta_E + 1/\delta_P - 1/\delta_P$		[2]
$1/\delta_P$	Duration of preclinical infectiousness in days	2.3 (95% CI: 0.8-3.0)	Gompertz	[2]
$1/\gamma_A = 1/\gamma_I$	Duration of clinical ($1/\gamma_I$) and subclinical ($1/\gamma_A$) infectiousness in days	7		[2, 3]
$1/\eta$	Delay from symptoms onset to hospitalization in days	7 (IQR: 4-8)	Gamma	[4]
$1/\alpha$	Delay from symptoms onset to ICU (here death) in days	10 (IQR: 6-12)	Gamma	[4]
$1/\gamma_H$	Delay from hospitalization to recovery in days	10 (IQR: 7-14)	Gamma	[4]
f	Fraction of infected people who develop symptoms	0.84 (95% CI: 0.8-0.88)	Binomial	[5]
h_i	Fraction of symptomatic people requiring hospitalization but not ICU	Age- and comorbidity-dependent		[6, 7]
g_i	Fraction of symptomatic people requiring ICU	Age- and comorbidity-dependent		[6, 7]


Parameter	Description	Value	Distribution	Reference
$1/\delta_E + 1/\delta_P$	Duration of incubation period in days	5.2 (95% CI: 4.1-7.0)	Lognormal	[1]
$1/\delta_E$	Duration of latency in days	$1/\delta_E + 1/\delta_P - 1/\delta_P$		[2]
$1/\delta_P$	Duration of preclinical infectiousness in days	2.3 (95% CI: 0.8-3.0)	Gompertz	[2]
$1/\gamma_A = 1/\gamma_I$	Duration of clinical ($1/\gamma_I$) and subclinical ($1/\gamma_A$) infectiousness in days	7		[2, 3]
$1/\eta$	Delay from symptoms onset to hospitalization in days	7 (IQR: 4-8)	Gamma	[4]
$1/\alpha$	Delay from symptoms onset to ICU (here death) in days	10 (IQR: 6-12)	Gamma	[4]
$1/\gamma_H$	Delay from hospitalization to recovery in days	10 (IQR: 7-14)	Gamma	[4]
f	Fraction of infected people who develop symptoms	0.84 (95% CI: 0.8-0.88)	Binomial	[5]
h_i	Fraction of symptomatic people requiring hospitalization but not ICU	Age- and comorbidity-dependent		[6, 7]
g_i	Fraction of symptomatic people requiring ICU	Age- and comorbidity-dependent		[6, 7]

Parameter	Description	Value	Distribution	Reference
$1/\delta_E + 1/\delta_P$	Duration of incubation period in days	5.2 (95% CI: 4.1-7.0)	Lognormal	[1]
$1/\delta_E$	Duration of latency in days	$1/\delta_E + 1/\delta_P - 1/\delta_P$		[2]
$1/\delta_P$	Duration of preclinical infectiousness in days	2.3 (95% CI: 0.8-3.0)	Gompertz	[2]
$1/\gamma_A = 1/\gamma_I$	Duration of clinical ($1/\gamma_I$) and subclinical ($1/\gamma_A$) infectiousness in days	7		[2, 3]
$1/\eta$	Delay from symptoms onset to hospitalization in days	7 (IQR: 4-8)	Gamma	[4]
$1/\alpha$	Delay from symptoms onset to ICU (here death) in days	10 (IQR: 6-12)	Gamma	[4]
$1/\gamma_H$	Delay from hospitalization to recovery in days	10 (IQR: 7-14)	Gamma	[4]
f	Fraction of infected people who develop symptoms	0.84 (95% CI: 0.8-0.88)	Binomial	[5]
h_i	Fraction of symptomatic people requiring hospitalization but not ICU	Age- and comorbidity-dependent	Values mapped to individuals 10 years older	[6, 7]
g_i	Fraction of symptomatic people requiring ICU	Age- and comorbidity-dependent	Values mapped to individuals 10 years older	[6, 7]


Model



Model



Model



$$\lambda_i = \sum_{j=1}^n \beta_{ij} \frac{P_j + A_j + I_j + H_j}{N_j}$$

With:

$$\beta_{ij} = \tau C_{ij}$$

τ is the probability of infection if there is a contact

C is the number of contacts between population classes i and j


$$\lambda_i = \sum_{j=1}^n \beta_{ij} \frac{P_j + A_j + I_j + H_j}{N_j}$$

With:

$$\beta_{ij} = \tau C_{ij}$$

τ is the probability of infection if there is a contact

C is the number of contacts between population classes i and j



THE RESULTS

Strategy 1: Self-distancing

Description

- Reduction of contacts between individuals.

Key-points

- Simple and rapid implementation.
- Educational-based, long-term benefit
- Starting investment then mouth-to-word

Strategy 1: Self-distancing

Description

- Reduction of contacts between individuals.

Key-points

- Simple and rapid implementation.
- Educational-based, long-term benefit
- Starting investment then mouth-to-word

Population class	Null model	Reduction 20%	Reduction 50%
Kids	25	20	12.5
Adults	15	11	7.5
Elderly	10	8	5

Strategy 1: Self-distancing

Description


- Reduction of contacts between individuals.

Key-points



- Simple and rapid implementation.
- Educational-based, long-term benefit
- Starting investment then mouth-to-word

Population class	Null model	Reduction 20%	Reduction 50%
Kids	25	20	12.5
Adults	15	11	7.5
Elderly	10	8	5



Strategy 1: Self-distancing



Strategy 1: Self-distancing



Strategy 1: Self-distancing



Up to 40% reduction in the the death tolls

Up to 40% delay in the peak of infected population

Strategy 2: Shielding and Lockdown

Description

- Shield of vulnerable population.
- Similar conditions in terms of tents occupancy and distance between tents are considered.
- Vulnerable:
 - Elderly.
 - People with co-morbidities.
 - Carers/family.
- Lockdown when first symptom is detected.

Shielding vulnerable population within a “green” zone



80% of the population


Neutral
zone

20% of the population


For a total population of 100 persons

- 50 kids (0-18 yo)
- 35 adults
- Meetings between orange and green in « **neutral zone** » (open tent), no more than 4 people at a time (mixed orange/green). No physical contact (mask and gloves, 1.5 meter distance).
- 0 kid
- 5 elderly (60+)
- 10 middle-aged NCD affected (mainly people 40-60 years old)
- Intra-green zone carers are among the 10 non-elderly NCD
- No unprotected contact with orange zone or external world
- In average, meeting in neutral zone once per week



Shielding vulnerable population within a “green” zone




Shielding vulnerable population within a “green” zone



Shielding vulnerable population within a “green” zone




Model




Strategy 2: Shielding only

Shielding
Population. size = 2000




Shielding
Population. size = 2000



Strategy 2: Shielding and Lockdown

Strategy 2: Shielding and Lockdown



Strategy 3: Isolation



Description

- Isolation of infected individuals
-

Key-points



- Remark 1.
- Remark 2.
- Requires quick identification of infected individuals

Shielding intervention and evacuation




- Assumptions

Strategy 3: Isolation



Combined strategies



Conclusions

- Remark 1
- Remark 2
- Combined strategies have a greater effect.