

MODELLING OF COVID-19 EPIDEMIC AND INTERVENTIONS WITHIN IDP CAMPS

DR. EDUARD CAMPILLO-FUNOLLET, PhD, UNIV. OF SUSSEX (UK)

JORDAN KLEIN, MPH, PRINCETON UNIV. (USA)

DR. ALBERTO PASCUAL-GARCÍA, PhD, ETH-ZÜRICH (SWITZERLAND)

DR. JENNIFER VILLERS, PhD, PRINCETON UNIV. (USA)

CHAMSY SARKIS, PAX SYRIANA FOUNDATION

JULY 16TH, 2020

PaxSyriana
Foundation

crowdfight 
COVID-19

MODEL PRELIMINARIES

Between December 2019 and March 2020, 1M new IDP → Informal settlements

HIGH DENSITY and no management

~700 camps with 600 person per camp on average (log-normal)

80% family-sized tents (~7 people)

Most of the informal camps have no management.

Between December 2019 and March 2020, 1M new IDP → Informal settlements

HIGH DENSITY and no management

~700 camps with 600 person per camp on average (log-normal)

80% family-sized tents (~7 people)

Most of the informal camps have no management.

NO POSSIBLE lockdown

50% no access to electricity

External water, with 25% not have enough drinking water

External latrines, and 10% households have 1 member disable/elderly.

360% drop of Syrian pound (no access to goods) → high motility to work

Between December 2019 and March 2020, 1M new IDP → Informal settlements

HIGH DENSITY and no management

~700 camps with 600 person per camp on average (log-normal)

80% family-sized tents (~7 people)

Most of the informal camps have no management.

NO POSSIBLE lockdown

50% no access to electricity

External water, with 25% not have enough drinking water

External letrines, and 10% households have 1 member disable/elderly.

360% drop of Syrian pound (no access to goods) → high motility to work

HIGH comorbidity

10% population has chronic diseases, 17% of them have no access to medicines.

AIM

IDENTIFY FEASIBLE INTERVENTIONS:

- IMMEDIATE APPLICABILITY
- NO NEED OF COMPLEX TECHNICAL INFRASTRUCTURE
(E.G. TESTING, PROTECTION)
- COST: AS LOW AS POSSIBLE

Outlook

Modelling


- Compartment models.
- Age-structured.
- Deterministic and stochastic simulations.
- Parameters estimated for IDPCs.
- Different camp sizes.
- ~70 different interventions modelled
- ~100K different simulations.

Strategies


- Self-distancing
- Shielding, lockdown
- Isolation
- Combined strategies

THE MODEL


Model



Model



Hospitalization capacity



Hospitalization capacity






Figure 6 Scenario One predicted severe cases

The health system in NW Syria would be unable to cope by the beginning of week 8; as such, severe cases could become critical and mortality could increase


Model




Model




Population Pyramid in NW of Syria




Population Pyramid in NW of Syria



Population Pyramid in NW of Syria



Model




Parameter	Description	Value	Distribution	Reference
$1/\delta_E + 1/\delta_P$	Duration of incubation period in days	5.2 (95% CI: 4.1-7.0)	Lognormal	[1]
$1/\delta_E$	Duration of latency in days	$1/\delta_E + 1/\delta_P - 1/\delta_P$		[2]
$1/\delta_P$	Duration of preclinical infectiousness in days	2.3 (95% CI: 0.8-3.0)	Gompertz	[2]
$1/\gamma_A = 1/\gamma_I$	Duration of clinical ($1/\gamma_I$) and subclinical ($1/\gamma_A$) infectiousness in days	7		[2, 3]
$1/\eta$	Delay from symptoms onset to hospitalization in days	7 (IQR: 4-8)	Gamma	[4]
$1/\alpha$	Delay from symptoms onset to ICU (here death) in days	10 (IQR: 6-12)	Gamma	[4]
$1/\gamma_H$	Delay from hospitalization to recovery in days	10 (IQR: 7-14)	Gamma	[4]
f	Fraction of infected people who develop symptoms	0.84 (95% CI: 0.8-0.88)	Binomial	[5]
h_i	Fraction of symptomatic people requiring hospitalization but not ICU	Age- and comorbidity-dependent		[6, 7]
g_i	Fraction of symptomatic people requiring ICU	Age- and comorbidity-dependent		[6, 7]


Parameter	Description	Value	Distribution	Reference
$1/\delta_E + 1/\delta_P$	Duration of incubation period in days	5.2 (95% CI: 4.1-7.0)	Lognormal	[1]
$1/\delta_E$	Duration of latency in days	$1/\delta_E + 1/\delta_P - 1/\delta_P$		[2]
$1/\delta_P$	Duration of preclinical infectiousness in days	2.3 (95% CI: 0.8-3.0)	Gompertz	[2]
$1/\gamma_A = 1/\gamma_I$	Duration of clinical ($1/\gamma_I$) and subclinical ($1/\gamma_A$) infectiousness in days	7		[2, 3]
$1/\eta$	Delay from symptoms onset to hospitalization in days	7 (IQR: 4-8)	Gamma	[4]
$1/\alpha$	Delay from symptoms onset to ICU (here death) in days	10 (IQR: 6-12)	Gamma	[4]
$1/\gamma_H$	Delay from hospitalization to recovery in days	10 (IQR: 7-14)	Gamma	[4]
f	Fraction of infected people who develop symptoms	0.84 (95% CI: 0.8-0.88)	Binomial	[5]
h_i	Fraction of symptomatic people requiring hospitalization but not ICU	Age- and comorbidity-dependent		[6, 7]
g_i	Fraction of symptomatic people requiring ICU	Age- and comorbidity-dependent		[6, 7]

Parameter	Description	Value	Distribution	Reference
$1/\delta_E + 1/\delta_P$	Duration of incubation period in days	5.2 (95% CI: 4.1-7.0)	Lognormal	[1]
$1/\delta_E$	Duration of latency in days	$1/\delta_E + 1/\delta_P - 1/\delta_P$		[2]
$1/\delta_P$	Duration of preclinical infectiousness in days	2.3 (95% CI: 0.8-3.0)	Gompertz	[2]
$1/\gamma_A = 1/\gamma_I$	Duration of clinical ($1/\gamma_I$) and subclinical ($1/\gamma_A$) infectiousness in days	7		[2, 3]
$1/\eta$	Delay from symptoms onset to hospitalization in days	7 (IQR: 4-8)	Gamma	[4]
$1/\alpha$	Delay from symptoms onset to ICU (here death) in days	10 (IQR: 6-12)	Gamma	[4]
$1/\gamma_H$	Delay from hospitalization to recovery in days	10 (IQR: 7-14)	Gamma	[4]
f	Fraction of infected people who develop symptoms	0.84 (95% CI: 0.8-0.88)	Binomial	[5]
h_i	Fraction of symptomatic people requiring hospitalization but not ICU	Age- and comorbidity-dependent	Values mapped to individuals 10 years older	[6, 7]
g_i	Fraction of symptomatic people requiring ICU	Age- and comorbidity-dependent	Values mapped to individuals 10 years older	[6, 7]


Model



Model



Model



$$\lambda_i = \sum_{j=1}^n \beta_{ij} \frac{P_j + A_j + I_j + H_j}{N_j}$$

With:

$$\beta_{ij} = \tau C_{ij}$$

τ is the probability of infection if there is a contact

C is the number of contacts between population classes i and j


$$\lambda_i = \sum_{j=1}^n \beta_{ij} \frac{P_j + A_j + I_j + H_j}{N_j}$$

With:


$$\beta_{ij} = \tau C_{ij}$$

τ is the probability of infection if there is a contact


C is the number of contacts between population classes i and j



Model



Model



THE RESULTS

Strategy 1: Self-distancing

Description

- Reduction of contacts between individuals.

Key-points

- Simple and rapid implementation.
- Educational-based, long-term benefit
- Starting investment then mouth-to-word

Strategy 1: Self-distancing

Description

- Reduction of contacts between individuals.

Key-points

- Simple and rapid implementation.
- Educational-based, long-term benefit
- Starting investment then mouth-to-word

Population class	Null model	Reduction 20%	Reduction 50%
Kids	25	20	12.5
Adults	15	11	7.5
Elderly	10	8	5

Strategy 1: Self-distancing

Description


- Reduction of contacts between individuals.

Key-points



- Simple and rapid implementation.
- Educational-based, long-term benefit
- Starting investment then mouth-to-word

Population class	Null model	Reduction 20%	Reduction 50%
Kids	25	20	12.5
Adults	15	11	7.5
Elderly	10	8	5



Strategy 1: Self-distancing



Strategy 1: Self-distancing



Strategy 1: Self-distancing



Up to 40% reduction in the the death tolls

Up to 40% delay in the peak of infected population

Strategy 2: Shielding and Lockdown

Description

- Shield of vulnerable population.
- Similar conditions in terms of tents occupancy and distance between tents were considered.
- Vulnerable:
 - Elderly.
 - People with co-morbidities.
 - Carers/family.
- Lockdown when first symptom is detected.

Shielding vulnerable population within a “green” zone



80% of the population

The diagram consists of two large rectangular blocks. The top block is orange and represents 80% of the population. The bottom block is green and represents 20% of the population. Between these two blocks is a blue oval labeled 'Neutral zone'.


Neutral
zone

20% of the population


For a total population of 100 persons

- 45 kids (0-18 yo)
- 35 adults
- Meetings between orange and green in « **neutral zone** » (open tent), no more than 4 people at a time (mixed orange/green). No physical contact (mask and gloves, 1.5 meter distance).
- 5 elderly (60+)
- 10 middle-aged NCD affected (mainly people 40-60 years old)
- 5 kids (<13 yrs)
- Intra-green zone carers are among the 10 non-elderly NCD
- No unprotected contact with orange zone or external world
- In average, meeting in neutral zone once per week (2 to 10 contacts per week with orange zone)



Shielding vulnerable population within a “green” zone



Shielding vulnerable population within a “green” zone




Shielding vulnerable population within a “green” zone



Strategy 2: Shielding only


Shielding

Population size = 2000



Strategy 2: Shielding only


Shielding
Population. size = 2000





Shielding
Population. size = 2000



Strategy 2: Shielding only



Lockdown of the “green” zone after first symptomatic case




Lockdown of the “green” zone after first symptomatic case


80% of the population

Neutral
zone


20% of the population




Strategy 2: Shielding and Lockdown



Strategy 3: Isolation



Strategy 3: Isolation




Isolation


Assumptions:

- Symptomatic people are isolated after they recognize their symptoms (12-48h),.
- The fate of the patients is not better nor worse than if staying in the camp → Discards isolating people together without testing.
- The model includes the presence of “carers” which cannot be neglected, even less without an adequate protection.


Model





Model



Strategy 3: Isolation




Strategy 3: Isolation




Combined strategies


Combined strategies




Combined strategies




Combined strategies



Combined strategies



Combined strategies



Conclusions

- Simple interventions can reduce the probability of outbreak and death tolls.
- **Setting safety areas** under appropriate conditions have an overall positive effect. An increase in the probability of outbreak can be mitigated by self-distancing.
- **Isolation** brings the stronger reduction in death tolls. However, it may accelerate the spread of the virus among the vulnerable population.
- The **combination of interventions** are very effective against the spread of the virus and its potential impact in the population.

Future work

- Determine the optimal population fraction that should be shielded (simulations done, analysis required).
- What can we expect in terms of herd immunity depending on the intervention, and estimate accurate times for e.g. lockdown.
- More accurate extrapolation of the spread of the infection at a regional scale.
- More realistic modelling of some interventions, e.g. introduction of carers if isolation.

Acknowledgements

Contributors:

- Jordan Klein (Princeton University)
- Jennifer Villers (Princeton University)
- Eduard Campillo-Funollet, University of Sussex
- Judith Bouman (ETH-Zürich)
- Chamsy Sarkis (Pax Syriana Foundation)

Thanks for useful discussions:

- Burcu Tepekule (ETH-Zürich)
- Juan Poyatos (Spanish Research Council)