

Empowering the crowd: Feasible strategies to minimize the spread of COVID-19 in high-density informal settlements

Alberto Pascual-García^(1,*), Jordan Klein⁽²⁾, Jennifer Villers^(2,^),
Eduard Campillo-Funollet^(3,^), Chamsy Sarkis⁽⁴⁾

August 24, 2020

(1) Institute of Integrative Biology. ETH-Zürich. Zürich, Switzerland

(2) Woodrow Wilson School. Princeton University. Princeton, NJ, USA.

(3) Genome Damage and Stability Centre, University of Sussex, Brighton, United Kingdom

(4) Pax Syriana Foundation, Valetta, Malta

(^) Equal contribution.

(*) correspondence: alberto.pascual@env.ethz.ch

Abstract

Background

More than 1 billion people live in informal settlements worldwide, in which the established paradigm to contain the spread of COVID-19 through social distancing, hygiene measures, and generalized use of personal protection equipment, present additional challenges. We investigated the impact of adapting these measures to informal settlements located in regions immerse in procrasted conflicts, which should additionally contend with the public health challenges resulting from violence, the deterioration of health-systems, and the political unstability which precludes coordinated action.

Methods

We implemented a stochastic Susceptible-Exposed-Infectious-Recovered model to simulate the spread of the virus in high-density camps of Internally Displaced Persons, with a population structure representative of the Northwest region of Syria. We chose parameters corresponding to a worst-case scenario in which no health-system is available. We evaluated interventions adapted to the living conditions in the camps, including self-distancing, self-isolation and the protection of vulnerable population in safety zones. We considered details representating social and technical challenges, which influence the micro-dynamics of the model.

Findings

All the interventions have a significant effect in reducing the probability of observing a global outbreak in the camp and the death tolls. Self-distancing brings the best results overall, if a 50% reduction in the number of contacts was considered, with mortality reduced up to a 35%. . A similar reduction in the casualties is achieved making available 1 tent for self-isolation per 200 individuals. Protecting vulnerable population in a safe area of the camp have synergistic effects with previous interventions for the whole population, and it is particularly beneficial for the vulnerable population. Complementaary mesaures such as the lockdown of the safety zone, further reduce the probability of an outbreak and the number of casualties. Importantly, the time at which the number of symptomatic individuals will peak is delayed with most of the interventions, in some cases by more than three months, and the proportion of the population recovered, near a 70%, could be compatible with herd immunity. The combination of interventions may reduce the mortality as much as 80%.

Interpretation

Our results highlight the potential of non-medical interventions to mitigate the effects of the epidemic. We demonstrate that interventions that were proved effective in other settings could be adapted to the camps. When the parameters considered have values that could be considered inefficient implementations of the interventions we still observed beneficial effects for the individual interventions and for their combination. The quantification we provide for specific details influencing the interventions such as the existence of buffer zones, daily health-checks, or the number of carers per isolated individual, suggest that the interventions are feasible and highlight the best avenues to implement them. The low cost and rapid applicability of the interventions support their idoneity, whose implementation would hence become mostly limited by the acquisition of local-communities of the necessary knowledge and means to incorporate them.

Introduction

The COVID-19 pandemic is intensifying in the developing world [1]. In Africa, SARS-CoV-2 has been spreading from urban areas to informal settlements [2]. With more than 1 billion people living in informal settlements worldwide, urgent action is needed to contain the virus in these settings, a task which necessarily involves the engagement of the communities living in them [3].

The need for action is even more pressing in regions immersed in protracted armed conflicts, where large portions of their populations have become displaced. When the displaced population exceeds official resettlement and refugee camp capacity, Internally Displaced Persons (IDPs) must live in informal settlements (hereafter named “camps”). These regions must contend with the public health challenges resulting from violence [4], the deterioration of health-systems [5], especially of critical care [6], and the breakdown of essential public infrastructure such as water and sanitation systems [7].

This study focuses on the Northwest region of Syria (NWS): a relatively small geographical area with 4.2 million people, of which 1.15 million (27.4%) are IDPs living in camps [8]. The health status of households in camps in NWS is poor; 24% have a member with a chronic disease, of whom 41% have no access to medicines [9]. As in other conflict regions, the political instability in NWS hinders coordinated public health actions, and the ongoing movements of IDPs create ample opportunity for infectious disease transmission, while making contact tracing interventions infeasible.

To investigate feasible COVID-19 prevention interventions in the camps, we considered a Susceptible-Exposed-Infectious-Recovered model in which the camps’ populations are divided into classes reflecting their estimated age-structures and comorbidity prevalence. We use this model to propose various interventions aimed at reducing the number of contacts within and between population classes in general, and with symptomatic individuals in particular. We paid special attention to how the living conditions in informal camps inform the assumptions underlying our proposed interventions. We modeled interventions previously proposed for African cities [10], such as self-distancing, isolation of symptomatic individuals and the creation of a ‘safety zone’ in which more vulnerable members of the population are protected from exposure to the virus.

Building upon the approach used to model the impact of these interventions in African cities, our model includes a parameterization of the contacts each individual has per day [10]. We further elaborate upon this approach by making a more explicit representation of contacts and other parameters in the model. We consider the micro-dynamics of contacts, the time that individuals take to recognize their symptoms before self-isolating, the effect of having carers to attend to isolated individuals, and the existence of a buffer zone in which exposed and protected population classes can interact under certain rules. We examine a potential worst-case scenario in which there is no access to any healthcare facility. Since empowering local communities in conflict regions to understand how to control COVID-19 is possibly the most (and perhaps only) effective way to minimize its spread, our models are of utmost importance for informing the implementation of realistic interventions in these regions.

Methods

The model

We considered a discrete-time stochastic model, simulating a viral outbreak in a single camp over a 12 month period. The model is divided into compartments containing individuals at different possible stages along the disease’s progression (see Supplementary Fig. 1), and splits the population into classes by age and comorbidity status. The simulation starts with a completely susceptible population where one person is exposed to the virus. The disease in exposed individuals progresses through a preclinical infectious stage, followed by either a clinical (symptomatic) or subclinical (asymptomatic) infectious stage, resolving through recovery or death. Additional susceptible individuals become infected through contact with infectious individuals. We verified that a steady state was always reached before the end of each simulation. We did not consider migration, births, nor deaths due to other causes, since they are small enough in magnitude to not significantly impact the course of an outbreak, provided additional conflict does not erupt.

Population structure

We parameterized the model with data from IDPs in NWS [11]. The population size of informal camps approximately log-normal, with a mean of 940 individuals. We simulated camps with 500, 1000 and 2000 individuals. Since interventions tend to be less effective in larger camps, the results presented refer to simulations with 2000 individuals, unless otherwise specified. We considered 3 age groups: children (age 1, 0-12 years old), working-age adults (age 2, 13-50 yrs.) and older adults (age 3, >50 yrs). For ages 2 and 3, we considered two subclasses comprising healthy individuals and individuals with comorbidities. The fraction of a simulated camp’s population in each of these 5 classes is shown in Supplementary Table 1.

Epidemiological severity assumptions

In NWS, there are 4 active and 2 in plan COVID-19 referral hospitals with 66 ventilators and current capacity of 74 ICU beds and 355 ward beds, for 4.2 million people [12, 13]. Basic estimation predicted a collapse of the health facilities after 8 weeks of the outbreak [14]. Hence, we considered a worst-case scenario in which individuals will not have access to healthcare. We consequently assumed that all critical cases, those requiring ICU care, would die. However, there is greater uncertainty about the fate of severe cases, those requiring hospitalization but not ICU care. We therefore considered a compartment for severe cases to account for a longer infectious period if they stay in the camp. This compartment also helped us model some interventions more realistically, for example by noting that the symptoms of severe cases are incompatible with self-isolation. To estimate upper and lower bounds for the outcome variables of our model, we simulated two possible scenarios for the fate of this compartment: one in which all cases recover, and another in which all die. In the simulations presented in the Main Text, we consider the worst-case scenario in which all of these cases die.

The fractions of symptomatic cases that are severe or critical are class-specific (parameters h_i and g_i , see Supplementary Table 1). We estimated these parameters using data from developed countries with superior population health [15, 16]. Following previous work [10], we reasoned that the case severity distributions of NW Syrian adult population classes would correspond with those of older age groups in developed countries.

Transmissibility assumptions

We assumed presymptomatic, asymptomatic, symptomatic and hospitalized individuals were equally infectious. We obtained the duration individuals spend in each compartment from the literature (see Supplementary Table 2). Each individual’s contact rate (see Supplementary Table 1) is class-specific, and was estimated from conversations with camp managers in NWS. The probability of random interaction with an individual from each class is proportional to this class’ fraction of the population. The product of these two values is the contact rate between two respective classes.

The probability of infection from contact with an infectious individual, τ , was estimated from a Gaussian distribution of the basic reproduction number, R_0 , with a mean of 4 (99% CI: 3–5). This distribution was a compromise between values reported in the literature from regions with high-density informal settlements: $R_0=2.77$ in Abuja and 3.44 in Lagos, Nigeria [17], 3.3 in Buenos Aires [18], and 5 in Rohingya refugee camps in Bangladesh [19]. The probability distribution of τ was estimated by randomly generating a value for R_0 , and dividing this value by the real part of the main eigenvalue of the Next Generation Matrix (see Supplementary Material).

Interventions




Figure 1: **Diagram of interventions.**

Self-distancing

We considered a situation in which the whole population reduces their mean number of contacts per day by a certain magnitude, 20% or 50% (see Fig. 1-1). Since the mean number of people per tent in a camp is 5.5 and sanitation facilities are shared [11] we inferred that the number of contacts per day cannot be reduced by more than 50%. For an adult, this would mean 7.5 contacts per day.

Buffer zone

Since only moderate self-distancing can be achieved, additional control measures involve splitting the population into subgroups occupying different zones of the camp, limiting cross-group contacts to specified locations, or “buffer zones”. We envision these zones as open spaces, with guidelines in place to limit occupancy to 4 individuals wearing masks, with 2 meters separating individuals from the two separate zones. Based on discussions with local organizations, we assumed a level of non-compliance such that the probability of transmission from contacts in these buffer zones will be reduced by 80%, rather than 100%, compared to the baseline (see Fig. 1).

Self-isolation

Self-isolation is a challenge in informal settlements, where households consist of a single (often small) space, water is collected at designated locations, sanitation facilities are communal and food supplies are scarce. We considered the possibility of those showing symptoms self-isolating in individual tents in dedicated parts of the camps, or next to the tents of their relatives. We simulated this intervention with various numbers of isolation tents per camp, ranging from 10 to 2000 for a camp of 2000 people (see Fig. 1-2). In addition, we modeled the role of carers dedicated to supplying isolated individuals, who interact with them via a buffer zone (see Fig. 1-2a). In considering one carer per individual with one contact per day, we do not neglect their probability of infecting the rest of the camp. We further considered that severe cases, or those in the hospitalized compartment, are fully infectious since they require more intensive care that is not possible to deliver while adhering to the guidelines of the buffer zone. We also considered minimum time intervals for individuals to recognize their symptoms: with means of 12, 24 and 48 hours (see Fig. 1-2b). Safety zone

In this intervention, the camp is divided in two areas: a safety zone, in which more vulnerable people live (hereby referred to as a “green” zone following previous studies [10]), and an exposed (“orange”) zone with the remaining population. In our simulations, the first exposed individual always belongs to the orange zone. We avoided the use of the term “shielding” to describe this intervention, since it may erroneously suggest that the vulnerable population is isolated in a closed space, such as a separated building. Such an intervention would require additional assumptions on how contacts occur in closed spaces. The living conditions within both zones remain the same, so the overall contact rate does not change unless self-distancing is also implemented. In practice, reducing the contact rate with individuals living in a different zone implies an increase in the contact rate with individuals in the same zone (see Supplementary Material). This allows us to investigate undesired side-effects of this intervention. Since proposals for partitioning the population may be received differently across camps, we considered several scenarios for allocating a camp’s population to the two zones (see Fig. 1-3, Supplementary Table 3). We consider the scenario in which all older adults, working-age adults with comorbidities and their family members up to 20% of the camp’s population live in the green zone in the Main Text, unless otherwise specified.

Interaction between the two zones is limited to a buffer zone. Individuals in the green zone cannot leave and thus need to be provided with supplies by individuals in the orange zone. Delivery of supplies will take place in the buffer zone. In our simulations, we considered limiting individuals in the green zone to 10 or 2 contacts with individuals from the orange zone per week (see Fig. 1-3a). Other variations of this intervention we explored include preventing symptomatic individuals from entering the buffer zone (see Fig. 1-3b) and a “lockdown” of the green zone, where the number of weekly contacts in the buffer zone is reduced by 50% or 90% (see Fig. 1-3c).

Evacuation

The last intervention we simulated is the evacuation of individuals in the hospitalization compartment. We assume the evacuees will be taken to isolation centers, not hospitals, so this measure does not change their fate; the only effect this intervention has is reducing the infectivity of evacuees to zero (see Fig. 1-4).

Analysis of the interventions

For each implementation of the interventions, we ran 500 simulations and compared results between them. The main variables considered are the fraction of simulations in which at least one death is observed, a proxy for the probability of an outbreak, the fraction of the population that dies and the time until the symptomatic population peaks, as well as the case fatality rate (CFR) and fraction of the population that recovers. For consistency, we only considered simulations in which there was an outbreak when comparing the outcome of a variable between interventions. We used the Shapiro-Wilk test [20] to verify that our results do not exhibit normally distributed residuals, Kruskal-Wallis test for pairwise comparisons [21], and Conover-Iman test for multiple comparisons [22]. The model and all statistical analysis were implemented in R [23]; we used the package PMCMRplus [24]. All the code and results are freely available at the url: <https://github.com/crowdfightcovid19/req-550-Syria>.

Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the article.

Results

Without any interventions, the mean CFR is ~2% in simulations where all severe cases requiring hospitalization recover (see Supplementary Fig. 2), and ~11% in simulations where all severe cases die. We consider the latter scenario to evaluate the interventions. In this scenario, the probability of observing an outbreak is close to 0.85, in which ~10% of the camp dies, the number of symptomatic cases peaks after 55 days and ~84% of the population recovers.




Figure 2: **Effect of interventions on outbreak probability, fatalities and time until symptomatic cases peak.** A: Self-distancing, probability of an outbreak. B: Self-distancing, fraction of the population dying. C: Self-distancing, time until peak symptomatic cases. D: Self-isolation, probability of an outbreak. E: Self-isolation, fraction of the population dying. F: Self-isolation, time until peak symptomatic cases. G: Safety zone, probability of an outbreak. H: Safety zone, fraction of the population dying. I: Safety zone, time until peak symptomatic cases. Note that, for the safety zone, the mean of the whole is not the mean of the exposed and safety zones, since means are computed considering simulations in which at least one death was observed in the population classes inhabiting the zone. This fact explains why in panel I there is a reduction in the mean time for the whole population despite of being a strong increase in the safety zone: in $\sim 35\%$ of the simulations there is an outbreak in the orange zone but not in the green zone (panel G).

Self-distancing has a notable effect on reducing the probability of an outbreak; even a 20% reduction in daily contacts is associated with a $\sim 10\%$ decrease in outbreak probability (see Fig. 2A). A greater reduction in daily contacts, of 50%, is required to observe a significant decrease, of as much as 35%, in the fraction of the population dying (see Fig. 2B). A reduction in contacts of this magnitude also significantly extends the time until the number of symptomatic cases peaks to 110 days, roughly double the time when no interventions are in place (see Fig. 2C). However, the proportion of the population recovered after 12 months is reduced by $\sim 30\%$ (see Supplementary Fig. 3)

With only 10 tents for a camp of 2000 people (i.e. 1 tent for every 200 people), self-isolation yields a modest decrease in the probability of observing an outbreak (see Fig. 2D), but a stronger reduction in mortality ($\sim 30\%$) (see Fig. 2E). However, further increasing the number of tents significantly augment this reduction up to 1 per every 10 people (Kruskal-Conover posthoc-test, $p\text{-val} < 3 \times 10^{-5}$) and, above this number, mortality even

begin to slightly increase again. This effect is a consequence of having one carer per individual isolated (see Supplementary Material), so when the isolated (infected) population increases, the number of healthy working-age adults in contact with them increases in tandem. Nevertheless, we observe a continuous reduction in CFR, albeit small, with an increasing number of tents (see Supplementary Fig. 4).

Importantly, the potential reductions in overall fatalities and CFR from self-isolation are realized whether the time required for individuals to recognize their symptoms is 12h or 24h on average, but the intervention becomes less effective when this time increases to 48h (see Supplementary Fig. 5). We observe no significant effects when severe cases requiring hospitalization are evacuated (see Supplementary Fig. 6). Since we assume that evacuation does not change their fate, this intervention only affects their infectivity. Although the period between developing more severe symptoms and dying is relatively long (~10 days), the number of individuals under these conditions is only a small fraction of the total infectious population at any given time.

Creating a green zone improves the effect of previous interventions overall, but with sometimes opposite outcomes for the exposed and protected populations. For example, the probability of an outbreak sharply decreases for the protected population, by almost 60%, if only two contacts are allowed per week in the buffer zone (see Fig. 2G). Notably, most of this reduction is only achieved when health-checks excluding symptomatic individuals from the buffer zone are in place (see Supplementary Fig. 7). On the other hand, the probability of an outbreak may slightly increase for the exposed population, a consequence of the relative increase in intra-zone contacts. Despite this side-effect, by shifting the burden of an outbreak towards the less vulnerable population in the orange zone, this intervention not only reduces fatalities among the more vulnerable population in the green zone (Kruskal-Wallis test, $p\text{-val} < 10^{-15}$; see Supplementary Fig. 8), but also reduces the overall CFR (see Supplementary Fig. 9) and thus the number of fatalities globally (see Fig. 2H). Another important outcome of this intervention is the notable increase in time until the number of symptomatic cases peaks for the vulnerable population. This effect is not observed globally, when there is an outbreak in the orange zone but not in the green zone (see Fig. 2I).

Considering different scenarios for allocating people to the green zone, the lowest probability of an outbreak is achieved when only older adults or at most older adults and working-age adults with comorbidities move there (see Supplementary Fig. 10). Positive effects of the green zone intervention are even more marked in camps with smaller populations, for every outcome of interest except time until symptomatic cases peak (see Supplementary Fig. 11). The incorporation of a lockdown has the greatest effect on reducing the probability of an outbreak in the green zone, to under 0.10. While lockdowns show no positive effect on green zone fatalities in the few instances where an outbreak does reach there, they decrease CFR and overall fatalities by further concentrating outbreaks in the less vulnerable population (see Supplementary Fig. 12).

The effects of the interventions observed when we examine them individually build upon each other when multiple interventions are implemented in tandem (see Fig. 3 and Supplementary Fig. 13). The protective effects of the safety zone intervention especially are most fully realized when paired with other interventions, less so when implemented on its own. When all interventions are implemented together: strict self-distancing (50% reduction in contacts), self-isolation of symptomatic cases (1 tent for every 40 people), a safety zone with 2 contacts per week in the buffer zone, health checks and a moderate lockdown (50%), and evacuation of severe cases, mortality is reduced by ~80%.




Figure 3: **Combinations of interventions.** Probability of an outbreak (top), fraction of the population dying (middle) and time until peak symptomatic cases (bottom) for different combination of interventions. Evac = evacuation of severely symptomatic, self = self-distancing, tents = number of available self-isolation tents, safety = safety zone, lock = lockdown of the buffer zone. For combinations of interventions including a safety zone, we distinguish between the population living in the green zone, in the orange zone and the whole population.

Discussion

In this study, we propose a number of interventions of immediate applicability to informal settlements. We focused on IDP settlements in NW Syria, taking into account the interventions' feasibility, cultural acceptance and their need for low-cost. When confronted with different possible scenarios, we generally considered the worst-

cases, highlighting the interventions that are most effective in the direst conditions, but possibly resulting in an overestimate of mortality. Our results align with previous simulation studies of potential COVID-19 interventions in similarly densely populated, low resource settings where informal settlements are present, such as urban areas of sub-Saharan Africa. In these settings, social-distancing is demonstrated to be an effective intervention, and even small changes are estimated to have large effects on outbreaks [25], in some cases determining whether or not already inadequate healthcare systems become overwhelmed [26]. Zandvoort et al. show that similar measures to the ones we consider: self-isolation, physical distancing and “shielding” the vulnerable, may reduce mortality by 60%-75% in African cities [10].

Self-distancing proves to be an effective measure in our models as well; reducing contacts by 50% has the greatest effect across the most outcomes of interest of any of the interventions we examined. However, the difficulty of achieving a reduction of this magnitude cannot be overlooked, especially considering the large proportion of the population composed of children, a group with an already high contact rate that may prove difficult to control [11].

We also propose self-isolation using individual tents which can be located in a dedicated zone or next to the tents of relatives, where contact with non-isolated individuals is mediated by a buffer zone. This intervention is effective with even a small number of isolation tents, as low as 5-10 tents per 1000 camp residents. After conversations with camp managers, we found that this intervention is more likely to be accepted in NW Syria than evacuation to community-based isolation centers. Community-based isolation not only poses cultural challenges; the capacity required to implement it has hardly been met [27].

One of the key parameters we assessed for the implementation of self-isolation is the need for carers. In considering one carer per isolated individual, with daily contact in a buffer zone, once a certain threshold of isolated cases (~ 200 per camp of 2000) is surpassed, the benefits of isolation begin to be outweighed by an increase in infectivity resulting from a growing number of exposed carers. This pitfall could be circumvented through the creation of a more organized group of carers, thus reducing the number of healthy working-age adults in contact with isolated (infectious) individuals.

Much of the success or failure of the safety zone intervention hinges on the functioning of the buffer zone. The number of inter-zone contacts per week, the implementation of health checks, and potential lockdowns all have notable effects. Also important is the portion of the population that is protected; protecting only the vulnerable may have the most beneficial effects, but it is precisely these vulnerable individuals, older adults and people with comorbidities, who may most need family members to care for them. While safety zone scenarios that allow greater numbers of family members to accompany their vulnerable relatives to the green zone may confer greater epidemiological risk, they may also engender greater well-being and social cohesion.

Although setting up a safety zone sharply reduces the probability of an outbreak in the population classes with the highest CFRs, thus reducing the CFR of the entire population, it is possible that our model may overestimate mortality from an outbreak in the green zone in the few instances when there is one. Since total numbers of contacts are conserved in our modelization, individuals do not reduce their contacts when moved to the green zone, which implies an increase in the number of contacts among vulnerable individuals. Despite of this increase, we observed a significant reduction in the casualties in the vulnerable population, when the safety zone is implemented. This result answers concerns raised around this type of intervention due to the large number of casualties registered in nursing-homes in wealthy countries [28]. While in wealthy countries individuals in a nursing-home will increase their exposure with respect to living confined in their household, in the camps individuals moving to the green zone will become less exposed.

An instrumental consideration for our models is the fraction of the population recovered from COVID-19 after a steady state is reached. Although the duration for which SARS-CoV-2 infection confers immunity is uncertain, the proportion of the population recovered after an outbreak should play a role in its protection against future ones. For every intervention except self-distancing of 50%, we observed that the fraction of the population recovered meets or exceeds 75%. This is quite promising to prevent future outbreaks.

Unaccounted for social and cultural dynamics will undeniably complicate the feasibility of our proposed interventions. One example we have not addressed here is the unlikelihood of children under 13 self-isolating. Although the number of exceptions to the cases proposed are possibly endless, the community-based nature of our approach may contribute to circumvent these challenges much faster than health-based interventions, which often depend on complex political decisions and may take years to reach the needed level of response. If the dynamics of the virus are well understood by local communities and at least some of the interventions we propose are implemented, the impacts of COVID-19 can be mitigated even in an environment as challenging as NW Syria.

Given a rapidly changing environment and slow responses of local and international authorities, empowering

local communities themselves is perhaps the best, if not the only way, to help them avoid the worst consequences of the pandemic. This not only applies to IDP camps in NW Syria, but to conflict-torn regions, informal settlements and vulnerable communities around the world: from the urban slums of India, townships of South Africa and favelas of Brazil, to the refugee camps of Kenya, Bangladesh and South Sudan; the low-cost, effective interventions we present are feasible, needed and urgent.

Acknowledgements

This collaboration was organized by crowdfightCOVID19 (www.crowdfightcovid19.org) upon request from CS. We thank Judith Boumann for valuable contributions. We thank Peter Ashcroft, Juan Poyatos and members of Sebastian Bonhoeffer's and Bryan Grenfell's groups for useful discussions. ECF research is supported by Wellcome Trust grant 204833/Z/16/Z.

Conflicts of interests

APG is a Board Member of crowdfightCOVID19, an initiative from the scientific community to put all available resources at the service of the fight against COVID-19. CS is a Board Member of the Pax Syria Foundation, a non-profit organization set up for social and philanthropic purposes including promoting and providing support and assistance to civilian aid projects in the fields of education, health, emergency assistance, psychological assistance and humanitarian aid for people affected by wars or humanitarian crisis. These organizations had no role in study design, data collection, data analysis, data interpretation, or writing of the article.

References

- [1] The Guardian. Total confirmed coronavirus cases in africa pass 1 million. url: <https://www.theguardian.com/world/2020/aug/06/total-confirmed-coronavirus-cases-in-africa-pass-1-million>. Accessed: 2020-08-17.
- [2] World Health Organization. Africa marks six months of covid-19. url: <https://www.afro.who.int/news/africa-marks-six-months-covid-19>. Accessed: 2020-08-17.
- [3] Annie Wilkinson. Local response in health emergencies: key considerations for addressing the COVID-19 pandemic in informal urban settlements. *Environment and Urbanization*, page 0956247820922843, May 2020.
- [4] Mohamed Abbas, Tammam Aloudat, Javier Bartolomei, Manuel Carballo, Sophie Durieux-Paillard, Laure Gabus, Alexandra Jablonka, Yves Jackson, Kanokporn Kaojaroen, Daniel Koch, Esperanza Martinez, Marc Mendelson, Roumyana Petrova-Benedict, Sotirios Tsiodras, Derek Christie, Mirko Saam, Sally Hargreaves, and Didier Pittet. Migrant and refugee populations: a public health and policy perspective on a continuing global crisis. *Antimicrobial Resistance and Infection Control*, 7, September 2018.
- [5] Peter S. Hill, Ghulam Farooq Mansoor, and Fernanda Claudio. Conflict in least-developed countries: challenging the Millennium Development Goals. *Bulletin of the World Health Organization*, 88:562–562, August 2010.
- [6] Mohammed Z. Sahloul, Jaber Monla-Hassan, Abdulghani Sankari, Mazen Kherallah, Bassel Atassi, Safwan Badr, Aula Abbara, and Annie Sparrow. War is the enemy of health. pulmonary, critical care, and sleep medicine in war-torn Syria. *Annals of the American Thoracic Society*, 13(2):147–155, January 2016.
- [7] Mustafa Sikder, Umar Daraz, Daniele Lantagne, and Roberto Saltori. Water, sanitation, and hygiene access in southern Syria: analysis of survey data and recommendations for response. *Conflict and Health*, 12(1):17, April 2018.
- [8] Health Information Central Unit and Health Directorates. Capacity of health sector in north of syria, mar 2020. Accessed: 2020-08-18.
- [9] REACH initiative. Syria: Camps & sites needs assessment - Northwest Syria sub-district profiles & comparative dashboard, January - February 2020. url: <https://reliefweb.int/report/syrian-arab-republic/syria-camps-sites-needs-assessment-northwest-syria-sub-district-profiles>.

- [10] Kevin Van Zandvoort, Christopher I Jarvis, Carl Pearson, Nicholas G Davies, Timothy W Russell, Adam J Kucharski, Mark J Jit, Stefan Flasche, Rosalind M Eggo, Francesco Checchi, et al. Response strategies for covid-19 epidemics in african settings: a mathematical modelling study. *MedRxiv*, 2020.
- [11] Assistance Coordinator Unit. The Syrian IDP camps monitoring study - Northern Syria camps - Humanitarian Data Exchange. url: <https://data.humdata.org/dataset/idp-camps-monitoring-november-of-2018>.
- [12] REACH initiative. North West Syria situation overview, (16 march 2020). url: https://www.impact-repository.org/document/reach/0169bf6a/REACH_SYR_Situation-Overview_Northwest-Syria_16-March-2020-1.pdf.
- [13] United Nations Office for the Coordination of Humanitarian Affairs. Situation report no. 18 - as of 25 july 2020. url: https://www.humanitarianresponse.info/sites/www.humanitarianresponse.info/files/documents/files/nw_syria_sitrep18_25july2020.pdf. Accessed: 2020-08-24.
- [14] Mahmoud Hariri, Hazem Rihawi, Salah Safadi, Mary Ana McGlasson, and Wael Obaid. The COVID-19 forecast in northwest syria: The imperative of global action to avoid catastrophe. *medRxiv*, page 2020.05.07.20085365, May 2020.
- [15] Yuanyuan Dong, Xi Mo, Yabin Hu, Xin Qi, Fang Jiang, Zhongyi Jiang, and Shilu Tong. Epidemiological characteristics of 2143 pediatric patients with 2019 coronavirus disease in china. *Pediatrics*, 2020.
- [16] CDC Covid, CDC COVID, CDC COVID, Nancy Chow, Katherine Fleming-Dutra, Ryan Gierke, Aron Hall, Michelle Hughes, Tamara Pilishvili, Matthew Ritchey, et al. Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019 in usa, february 12–march 28, 2020. *Morbidity and Mortality Weekly Report*, 69(13):382, 2020.
- [17] Mutiu Abimbola Oyinlola, Tolulope Osayomi, and Oluwatosin Adeniyi. Empirical modelling of confirmed COVID-19 cases in Nigeria: Forecasts and implications. SSRN Scholarly Paper ID 3596095, Social Science Research Network, Rochester, NY, May 2020.
- [18] Juan E. Santos, Jose’ M. Carcione, Gabriela B. Savioli, Patricia M. Gauzellino, Alejandro Ravecca, and Alfredo Moras. A numerical simulation of the COVID-19 epidemic in Argentina using the SEIR model. *arXiv:2005.06297 [physics, q-bio]*, July 2020. arXiv: 2005.06297.
- [19] Shaun A. Truelove, Orit Abraham, Chiara Altare, Stephen A. Lauer, Andrew Azman, and Paul B. Spiegel. The potential impact of COVID-19 in refugee camps in Bangladesh and beyond: a modeling study. *medRxiv*, page 2020.03.27.20045500, May 2020.
- [20] Samuel Sanford Shapiro and Martin B Wilk. An analysis of variance test for normality (complete samples). *Biometrika*, 52(3/4):591–611, 1965.
- [21] William H Kruskal and W Allen Wallis. Use of ranks in one-criterion variance analysis. *Journal of the American statistical Association*, 47(260):583–621, 1952.
- [22] William Jay Conover and Ronald L Iman. On multiple-comparisons procedures. *Los Alamos Sci. Lab. Tech. Rep. LA-7677-MS*, 1:14, 1979.
- [23] R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2020.
- [24] Thorsten Pohlert. *The Pairwise Multiple Comparison of Mean Ranks Package Extended (PMCMRplus)*, 2020. R package.
- [25] Farai Nyabadza, Faraimunashe Chirove, Williams Chidozie Chukwu, and Maria Vivien Visaya. Modelling the potential impact of social distancing on the covid-19 epidemic in south africa. *medRxiv*, 2020.
- [26] Amir S. Siraj, Alemayehu Worku, Kiros Berhane, Yemane Berhane, and Dawd S. Siraj. Early estimates of COVID-19 infections in small, medium and large population clusters. *medRxiv*, page 2020.04.07.20053421, May 2020.

- [27] United Nations Office for the Coordination of Humanitarian Affairs. Situation report no. 19 - as of 21 august 2020. url: https://www.humanitarianresponse.info/sites/www.humanitarianresponse.info/files/documents/files/nw_-syria_sitrep19_21aug2020.pdf. Accessed: 2020-08-24.
- [28] Maysoon Dahab, Kevin van Zandvoort, Stefan Flasche, Abdihamid Warsame, Ruwan Ratnayake, Caroline Favas, Paul B Spiegel, Ronald J Waldman, and Francesco Checchi. Covid-19 control in low-income settings and displaced populations: what can realistically be done? *Conflict and Health*, 14(1):1–6, 2020.