



## Hamburger-Siegel Theorem

The **Hamburger-Siegel theorem** (more commonly known as **Hamburger's theorem** for the moment problem) characterizes when a sequence of real numbers is a moment sequence of a positive measure on the real line. <sup>[1] [2] [3] [4]</sup>

### Statement of the Theorem

Given a sequence  $(s_n)_{n \geq 0}$  of real numbers, the **Hamburger moment problem** asks whether there exists a positive Borel measure  $\mu$  on  $\mathbb{R}$  such that

$$s_n = \int_{-\infty}^{\infty} x^n d\mu(x), \quad n = 0, 1, 2, \dots$$

**Hamburger's Theorem** (Existence): The sequence  $(s_n)_{n \geq 0}$  is a moment sequence if and only if the **Hankel matrix**

$$H = \begin{pmatrix} s_0 & s_1 & s_2 & \cdots \\ s_1 & s_2 & s_3 & \cdots \\ s_2 & s_3 & s_4 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

is **positive semidefinite**. That is, for every finite sequence  $(c_j)_{j \geq 0}$  of complex numbers with only finitely many nonzero terms, <sup>[3] [4] [1]</sup>

$$\sum_{j,k \geq 0} s_{j+k} c_j \overline{c_k} \geq 0.$$

Equivalently, all principal minors (finite Hankel determinants) must be nonnegative:

$$\det \begin{pmatrix} s_0 & s_1 & \cdots & s_{n-1} \\ s_1 & s_2 & \cdots & s_n \\ \vdots & \vdots & \ddots & \vdots \\ s_{n-1} & s_n & \cdots & s_{2n-2} \end{pmatrix} \geq 0$$

for all  $n \geq 1$ . <sup>[1]</sup>

## Uniqueness (Determinacy)

Hamburger also addressed the uniqueness question. The moment problem is **determinate** (the measure  $\mu$  is unique) if and only if

$$\lim_{n \rightarrow \infty} \frac{\det(H_n)}{\det(H'_n)} = 0,$$

where  $H_n$  and  $H'_n$  are certain finite sections of the Hankel matrix. A more practical sufficient condition for determinacy is **Carleman's criterion**: if<sup>[1]</sup>

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt[2n]{s_{2n}}} = \infty,$$

then the moment problem is determinate.<sup>[2] [1]</sup>

## Hamburger's Converse Theorem for the Riemann Zeta Function

A different but related result, also due to Hamburger, characterizes the Riemann zeta function via its functional equation.<sup>[5] [6] [7] [8]</sup>

**Hamburger's Converse Theorem** (1921): Let  $h(s) = \sum_{n=1}^{\infty} a_n n^{-s}$  and  $g(s) = \sum_{n=1}^{\infty} b_n n^{-s}$  be absolutely convergent for  $\Re(s) > 1$ , and suppose both  $(s-1)h(s)$  and  $(s-1)g(s)$  are entire functions of finite order. If the functional equation<sup>[6] [5]</sup>

$$\pi^{-s/2} \Gamma\left(\frac{s}{2}\right) h(s) = \pi^{-(1-s)/2} \Gamma\left(\frac{1-s}{2}\right) g(1-s)$$

holds, then  $h(s) = g(s) = a_1 \zeta(s)$ .

This says the Riemann zeta function is **uniquely determined** by its functional equation (up to a constant factor), provided suitable regularity conditions hold.<sup>[5] [6]</sup>

## Detailed Proof of Hamburger's Theorem

The proof uses orthogonal polynomials, functional analysis, and the theory of linear functionals on polynomial algebras.

### Step 1: Reformulation via Linear Functionals

For a sequence  $s = (s_n)_{n \geq 0}$ , define a linear functional  $L_s$  on the polynomial algebra  $\mathbb{R}[x]$  by

$$L_s(x^n) = s_n, \quad n = 0, 1, 2, \dots$$

By linearity of integration,  $(s_n)$  is a moment sequence for measure  $\mu$  if and only if

$$L_s(p) = \int_{-\infty}^{\infty} p(x) d\mu(x)$$

for all polynomials  $p \in \mathbb{R}[x]$ .<sup>[4]</sup>

## Step 2: Positive Definiteness

The measure  $\mu$  is positive if and only if

$$L_s(p^2) = \int_{-\infty}^{\infty} p(x)^2 d\mu(x) \geq 0$$

for all  $p \in \mathbb{R}[x]$ . Writing  $p(x) = \sum_{j=0}^m c_j x^j$ , we have<sup>[4]</sup>

$$L_s(p^2) = L_s \left( \left( \sum_j c_j x^j \right)^2 \right) = \sum_{j,k} c_j c_k s_{j+k}.$$

Thus,  $\mu$  is positive if and only if the associated Hankel form is positive semidefinite.<sup>[3] [4]</sup>

## Step 3: Construction via Orthogonal Polynomials

Given a positive definite sequence  $(s_n)$ , construct orthonormal polynomials  $(P_n)$  via Gram-Schmidt orthogonalization with respect to the inner product

$$\langle p, q \rangle = L_s(pq).$$

These polynomials satisfy a three-term recurrence relation

$$xP_n(x) = b_n P_{n+1}(x) + a_n P_n(x) + b_{n-1} P_{n-1}(x),$$

with  $b_n > 0$ .<sup>[2] [1]</sup>

By **Favard's theorem**, such a recurrence relation determines a positive measure  $\mu$  for which  $(P_n)$  are orthonormal.<sup>[2] [1]</sup>

## Step 4: Spectral Theorem

The multiplication operator  $Mf(x) = xf(x)$  on  $L^2(\mu)$  is self-adjoint. By the spectral theorem,  $\mu$  is the spectral measure of this operator, uniquely determined by the moments.<sup>[4] [1]</sup>

## Step 5: Uniqueness (Determinacy)

Hamburger proved that the measure  $\mu$  is unique if and only if the polynomials are dense in  $L^2(\mu)$ , or equivalently,

$$\sum_{n=0}^{\infty} (P_n^2(0) + Q_n^2(0)) < \infty,$$

where  $Q_n$  are the polynomials of the second kind. Carleman's sufficient condition follows from analyzing growth rates of moments.<sup>[1] [2]</sup>

## Conclusion

Hamburger's theorem provides a complete solution to the existence and uniqueness questions for the moment problem on  $\mathbb{R}$ . The positive definiteness criterion is both necessary and sufficient for existence, and various criteria (Carleman, Krein) determine uniqueness. <sup>[3] [2] [4] [1]</sup>

\*\*

1. [https://web.williams.edu/Mathematics/sjmiller/public\\_html/book/papers/jcmp.pdf](https://web.williams.edu/Mathematics/sjmiller/public_html/book/papers/jcmp.pdf)
2. [https://en.wikipedia.org/wiki/Stieltjes\\_moment\\_problem](https://en.wikipedia.org/wiki/Stieltjes_moment_problem)
3. [https://en.wikipedia.org/wiki/Hamburger\\_moment\\_problem](https://en.wikipedia.org/wiki/Hamburger_moment_problem)
4. <https://arxiv.org/pdf/2008.12698.pdf>
5. <https://arxiv.org/pdf/1910.09837.pdf>
6. [https://www.clas.kitasato-u.ac.jp/~miyazaki/rims2022/RIMS\\_Jan23.pdf](https://www.clas.kitasato-u.ac.jp/~miyazaki/rims2022/RIMS_Jan23.pdf)
7. <https://sites.math.rutgers.edu/~zeilberg/EM18/TitchmarshZeta.pdf>
8. <https://sites.math.rutgers.edu/~sdmiller/gelbart-miller-zeta.pdf>
9. [https://en.wikipedia.org/wiki/Siegel's\\_theorem\\_on\\_integral\\_points](https://en.wikipedia.org/wiki/Siegel's_theorem_on_integral_points)
10. <https://surya-teja.com/2010/11/24/riemann-functional-equation-and-hamburgers-theorem/>
11. <https://www.fandm.edu/directory/wendell-ressler.html>
12. <https://www.sciencedirect.com/science/article/pii/S0096300396003050>
13. <https://gorogoro.cis.ibaraki.ac.jp/web/papers/kano2018-a.pdf>
14. <https://projecteuclid.org/journals/pacific-journal-of-mathematics/volume-27/issue-3/On-determinate-Hamburger-moment-problems/pjm/1102983768.pdf>
15. <https://arxiv.org/abs/2407.08511>
16. [https://womengovtcollegevisakha.ac.in/departments/Number theory \[p211-226\].pdf](https://womengovtcollegevisakha.ac.in/departments/Number%20theory%20[p211-226].pdf)
17. <https://harvest.usask.ca/bitstreams/7c81b396-34c2-47cc-9b9c-be6d20ddc34c/download>
18. [https://kam.mff.cuni.cz/~kyncl/presentations/40\\_hamburger\\_theorem/burger.pdf](https://kam.mff.cuni.cz/~kyncl/presentations/40_hamburger_theorem/burger.pdf)
19. <https://mathworld.wolfram.com/PositiveDefiniteSequence.html>
20. <https://math.iisc.ac.in/~khare/papers/hankel.pdf>
21. [https://www.cambridge.org/core/services/aop-cambridge-core/content/view/517508F83FF5A35EF04D6F0F7C1483FA/S0013091500022628a.pdf/an\\_extended\\_hamburger\\_moment\\_problem.pdf](https://www.cambridge.org/core/services/aop-cambridge-core/content/view/517508F83FF5A35EF04D6F0F7C1483FA/S0013091500022628a.pdf/an_extended_hamburger_moment_problem.pdf)
22. <https://www.sciencedirect.com/science/article/pii/S0925772117300640>
23. <https://www.msand.dk/article/download/112091/163320/236397>
24. [https://en.wikipedia.org/wiki/Moment\\_problem](https://en.wikipedia.org/wiki/Moment_problem)
25. <https://www.sciencedirect.com/science/article/pii/S0377042787901750>
26. <https://ems.press/journals/jems/articles/2579441>
27. <https://www.sciencedirect.com/science/article/pii/S0024379515006035>
28. [https://onlinelibrary.wiley.com/doi/pdf/10.1002/\(SICI\)1522-2616\(200002\)210:1<67::AID-MANA67>3.0.CO;2-N](https://onlinelibrary.wiley.com/doi/pdf/10.1002/(SICI)1522-2616(200002)210:1<67::AID-MANA67>3.0.CO;2-N)

29. <https://projecteuclid.org/journals/tokyo-journal-of-mathematics/volume-5/issue-1/On-an-Analogoue-to-Hecke-Correspondence/10.3836/tjm/1270215036.pdf>