

# Testing for Harmonizability

BY H. L. HURD

*IEEE Transactions on Information Theory, Vol. IT-19, No. 3, May 1973*

## Abstract

Let  $R(s, t)$  be a covariance function having the representation

$$R(s, t) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp(i s x - i t y) d^2 G(x, y) \quad (1)$$

where  $G(x, y)$  is continuous to the right in both variables and is of bounded variation in the plane; then  $R(s, t)$  is harmonizable in that  $G(x, y)$  is also a covariance. Examples are given in which this result is used to determine the harmonizability of new processes and covariances that are formed by operations on old processes and covariances. Specifically, if  $X(t)$  is a real Gaussian harmonizable process, then  $X^n(t)$  is harmonizable. If  $X(t)$  is harmonizable,  $G(x, y)$  has bounded support and  $g(t)$  is a Fourier-Stieltjes transform, then  $X(g(t))$  and  $X(t + g(t))$  are harmonizable. If

$$X(t) = \int_{-\infty}^{\infty} f(t, u) dZ(u) \quad (2)$$

where  $f(t, u) = f(t - u)$  is a Fourier-Stieltjes transform and  $G(u, v) = \mathbb{E}\{Z(u)Z^*(v)\}$  has finite total variation, then  $X(t)$  is harmonizable. A sufficient condition for Priestley's oscillatory processes to be harmonizable is also obtained. The Bochner-Eberlein characterization of Fourier-Stieltjes transforms is particularly convenient for determining harmonizability in these cases.

## 1 Introduction

Let  $\{X(t, \omega), -\infty < t < \infty, \omega \in \Omega\}$  be a second-order continuous-parameter stochastic process defined on some probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . The process  $X(t, \omega)$  is said to be harmonizable [loeve1955probability, p. 474] if it has the quadratic mean representation

$$X(t, \omega) = \int_{-\infty}^{\infty} \exp(i t x) dZ(x, \omega) \quad (3)$$

where  $\{Z(x, \omega), -\infty < x < \infty\}$  is a process whose covariance is of bounded variation (BV) in the plane. Harmonizable processes are of engineering interest because decomposition relative to  $\exp(i t x)$  admits the usual frequency interpretation of linear filtering. If  $H(x)$  is the frequency response of a stable, linear time-invariant system, then the system output process  $Y(t, \omega)$  is given by the quadratic mean integral

$$Y(t, \omega) = \int_{-\infty}^{\infty} \exp(i t x) H(x) dZ(x, \omega) \quad (4)$$

A detailed account may be found in [blanc-lapierre1968random, Ch. 8]. For recent results on harmonizable processes in engineering, see [cambanis1970harmonizable, donati1971spectra, ogura1971spectral].

The covariance functions for  $X(t)$  and  $Y(t)$  are

$$R(s, t) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp(i s x - i t y) d^2 G(x, y) \quad (5)$$

$$R_Y(s, t) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp(i s x - i t y) H(x) H^*(y) d^2 G(x, y) \quad (6)$$

where  $d^2 G(x, y) = \mathbb{E} \{dZ(x) dZ^*(y)\}$ . Any covariance with representation (5), with  $G(x, y)$  a covariance of bounded variation, is called harmonizable; harmonizable processes have harmonizable covariances. Conversely, processes with harmonizable covariances are themselves harmonizable [loeve1955probability, p. 476]. For brevity, we call those corresponding to  $G$  with finite total variation simply “harmonizable.”

This paper addresses the determination of harmonizability for new processes or covariances constructed from old ones. The main results are as follows:

- If  $X(t)$  is a real Gaussian harmonizable process, then  $X^n(t)$  is harmonizable.
- If  $X(t)$  is harmonizable with spectral decomposition of bounded support and  $g(t)$  is a Fourier-Stieltjes transform, then  $X(t + g(t))$  and  $X(g(t))$  are harmonizable.
- If  $R_1$  and  $R_2$  are harmonizable covariances, then for  $T$  of finite Lebesgue measure,

$$R_3(s, t) = \int_T R_1(s, u) R_2(u, t) du \quad (7)$$

is harmonizable.

- If  $X(t)$  is a moving average as in (2) with  $f(t, u) = f(t - u)$  a Fourier-Stieltjes transform and  $G(u, v)$  of bounded variation, then  $X(t)$  is harmonizable.
- If  $X(t)$  is as above,  $Z(u)$  has orthogonal increments with  $dF(u) = \mathbb{E} |dZ(u)|^2$  and

$$f(t, u) = \exp(i u t) \int_{-\infty}^{\infty} \exp(i t x) dH_u(x) \quad (8)$$

with  $H_u(x)$  of bounded variation for every  $u$ , the resulting processes include Priestley’s oscillatory processes [priestley1965evolutionary], which are harmonizable under suitable conditions.

The method is to use the following result.

**Theorem 1.** *If  $R(s, t)$  is simultaneously a covariance and a Fourier-Stieltjes (FS) transform with respect to some  $G(x, y)$  of bounded variation, then  $R(s, t)$  is harmonizable in that  $G$  is necessarily a covariance.*

**Proof.** Sufficiency is immediate: one can find a process  $Z(x, \omega)$  whose covariance is  $G$  and whose FS transform  $X(t, \omega)$  as in (3) has covariance  $R(s, t)$ . Conversely, suppose  $R$  is both a covariance and an FS transform with respect to some  $G$  that is BV. Define

$$G_a(x, y) = G(x, y) - G(a, y) - G(x, a) + G(a, a) \quad (9)$$

For any  $n$  and sequence  $\{x_j \geq a, j = 1, \dots, n\}$  and complex  $\{c_j\}$ ,

$$\sum_{j,k=1}^n c_j c_k^* G_a(x_j, x_k) \geq 0$$

This follows by constructing  $g_a(s) = \sum_{j=1}^n c_j [1 - \exp(-i s x_j)] \exp(-i s a)$  and applying the inversion theorem [loeve1955probability, p. 475]. Letting  $a \rightarrow -\infty$ ,  $G_a(x, y) \rightarrow G(x, y)$ . Thus  $G(x, y)$  is non-negative definite.  $\square$

Thus, any characterization of FS transforms, such as the Bochner-Eberlein theorem, also provides a characterization for harmonizable covariances [bochner1934fst, eberlein1955fst, rudin1962groups]. Cramér [cramer1939representation] and Dominguez [dominguez1940fst] provide alternative characterizations.

## 2 Mathematical Preliminaries

Assume  $G(x, y)$  is normalized, e.g.,

$$G(x, y) = \frac{1}{4} [G(x+0, y+0) + G(x+0, y-0) + G(x-0, y+0) + G(x-0, y-0)] \quad (10)$$

and satisfies

$$\lim_{x \rightarrow -\infty} G(x, y) = \lim_{y \rightarrow -\infty} G(x, y) = 0.$$

We now state the key characterizations.

**Theorem 2.** [Bochner] A necessary and sufficient condition that  $f(t)$ ,  $-\infty < t < \infty$ , has the representation

$$f(t) = \int_{-\infty}^{\infty} \exp(i x t) dG(x) \quad (11)$$

for a complex measure  $G$  of bounded variation is that, for any  $n$ , any sequence  $\{t_j\}_{j=1}^n$  and any complex  $\{c_j\}_{j=1}^n$ ,

$$\left| \sum_{j=1}^n c_j f(t_j) \right| \leq M \left[ \sum_{j=1}^n \sum_{k=1}^n c_j c_k^* \exp(i x (t_j - t_k)) \right]^{1/2} \quad (12)$$

for some  $M > 0$ .

**Theorem 3.** [Bochner-Eberlein] A necessary and sufficient condition for a function  $R(s, t)$  to have representation (1) with

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |d^2 G(x, y)| \leq M \quad (13)$$

is that for any  $n$ , any sequence of pairs  $\{(s_j, t_j)\}_{j=1}^n$  and any complex  $\{c_j\}_{j=1}^n$ ,

$$\left| \sum_{j=1}^n c_j R(s_j, t_j) \right| \leq M \left[ \sum_{j=1}^n \sum_{k=1}^n c_j c_k^* \exp(i (x s_j - y t_j - x s_k + y t_k)) \right]^{1/2} \quad (14)$$

for some  $M > 0$ .

### 3 Examples and Application

**Example 4.** Let  $X(t)$  be a zero-mean real Gaussian harmonizable process with covariance  $R(s, t)$ . Then, for any  $n \geq 1$ ,  $X^n(t)$  is harmonizable.

**Proof.** An exercise with characteristic functions shows that

$$\mathbb{E}[X^n(s) X^n(t)] = \sum_{p, q, r \geq 0} c_z(p, q, r, n) R^p(s, s) R^q(s, t) R^r(t, s) R^p(t, t) \quad (15)$$

where the sum is over all  $p, q, r \geq 0$  with  $n = 2p + q + r$  and  $c_z(p, q, r, n)$  are combinatorial coefficients.

Since both FS transforms [bochner1934fst, p.151] and covariances [loeve1955probability, p.468] are closed under products,  $R^q(s, t) R^r(t, s) = R^{q+r}(s, t)$  is an FS transform and a covariance. The product  $R^p(s, s) R^p(t, t)$  is also an FS transform, as for  $f(s) = R(s, s)$

$$f^p(s) f^p(t)$$

is NND and an FS transform by Theorem 2. The sum in (15) is harmonizable since FS transforms and covariances are closed under positive sums.  $\square$

**Example 5.** Suppose  $X(t)$  is harmonizable with spectral decomposition supported by a bounded set  $A$  and  $g(t)$  is the FS transform of some  $G(x)$  of finite variation. Then  $X(t + g(t))$  and  $X(g(t))$  are harmonizable.

**Proof.** Set  $Y(t) = X(t + g(t))$ , so

$$R_Y(s, t) = R_X(s + g(s), t + g(t)) = \langle \text{int} \rangle_A \exp(i x(s + g(s)) - i y(t + g(t))) d^2 G(x, y).$$

Let  $M_A$  denote the variation over  $A \times A$  of  $G(x, y)$ . For any complex  $\{c_j\}$  and parameter pairs,

$$Q = \left| \sum_{j=1}^n c_j R_Y(s_j, t_j) \right| \leq M_A \left| \sum_{j=1}^n c_j \exp(i x' s_j + i x' g(s_j) - i y' t_j - i y' g(t_j)) \right|$$

where  $x', y'$  in closure of  $A$ . The mappings  $t \mapsto \exp[i x' g(t)]$  and  $t \mapsto \exp[-i y' g(t)]$  are FS transforms, so by repeated application of Bochner's condition this is bounded, and  $R_Y(s, t)$  is an FS transform.  $\square$

**Example 6.** Suppose  $R_1$  and  $R_2$  are harmonizable covariances, and for  $T$  of finite Lebesgue measure define  $R_3(s, t)$  as in (7). Then  $R_3$  is harmonizable.

**Proof.** By the Bochner-Eberlein condition, for any  $\{c_j\}$ ,

$$Q = \left| \sum_{j=1}^n c_j \int_T R_1(s_j, u) R_2(u, t_j) du \right| \leq M_1 \sup_{x, y} \left| \sum_{j=1}^n c_j \int_T R_2(u, t_j) \exp(i s_j x - i u y) du \right|.$$

With  $M_2$  the variation bound for  $R_2$ ,

$$Q \leq M_1 M_2 m(T) \sup_{x, y} \left| \sum_{j=1}^n c_j \exp(i s_j x - i t_j y) \right|$$

where  $m(T)$  is Lebesgue measure. Thus  $R_3$  is an FS transform.  $\square$

**Example 7.** Suppose  $X(t)$  is a moving average as in (2) with  $f(t, u) = f(t - u)$  a Fourier-Stieltjes transform and  $G(u, v)$  of bounded variation. Then  $X(t)$  is harmonizable.

**Proof.** From (2) and the bounded variation of  $H(x)$  and  $G(u, v)$ ,

$$R(s, t) = \langle \text{iiiiint} \rangle \exp [i x (s - u) - i y (t - v)] d^2 G(u, v) d H(x) d H^*(y).$$

This is an FS transform with variation bound  $M_G M_H^2$ , where  $M_G$  bounds  $G(u, v)$  and  $M_H$  bounds  $H(x)$ .  $\square$

**Example 8.** Suppose  $X(t)$  is as above,  $Z(u)$  has orthogonal increments with  $d F(u) = \mathbb{E}[|d Z(u)|^2]$  and  $f(t, u)$  as in (8). Then  $X(t)$  is a Priestley oscillatory process, and is harmonizable provided

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |d H_u(x)| |d H_u^*(y)| d F(u) < \infty. \quad (16)$$

**Proof.** The covariance is

$$R(s, t) = \int_{-\infty}^{\infty} \exp [i u (s - t)] f(u, s) f^*(u, t) d F(u).$$

By repeated application of Bochner's and Fubini's theorems, provided (16) holds,  $R(s, t)$  is an FS transform and thus harmonizable.  $\square$

## Acknowledgment

The author is indebted to H. J. Landau for remarks on an early version of this paper and to a referee for several helpful suggestions.

## Bibliography

- [loeve1955probability] M. Loève, *Probability Theory*. New York: Van Nostrand, 1955.
- [blanc-lapierre1968random] A. Blanc-Lapierre and R. Fortet, *Theory of Random Functions*. New York: Gordon and Breach, 1968.
- [cambanis1970harmonizable] S. Cambanis and B. Liu, "On harmonizable stochastic processes," *Inform. Contr.*, vol. 17, pp. 183–202, 1970.
- [donati1971spectra] F. Donati, "Finite-time averaged power spectra", *IEEE Trans. Inform. Theory*, vol. IT-17, pp. 7–16, Jan. 1971.
- [ogura1971spectral] H. Ogura, "Spectral representation of a periodic nonstationary random process," *IEEE Trans. Inform. Theory*, vol. IT-17, pp. 143–149, Mar. 1971.
- [gladyshev1963almost] E. G. Gladyshev, "Periodically and almost-periodically correlated random processes with continuous time parameter," *Theory Prob. Appl.*, vol. 8, pp. 173–177, 1963.
- [priestley1965evolutionary] M. B. Priestley, "Evolutionary spectra and nonstationary processes," *J. Roy. Statist. Soc. B*, vol. 27, pp. 204–237, 1965.
- [bochner1934fst] S. Bochner, "A theorem on Fourier-Stieltjes integrals," *Bull. Amer. Math. Soc.*, vol. 40, pp. 271–276, 1934.

- [eberlein1955fst] W. F. Eberlein, “Characterization of Fourier-Stieltjes transforms,” *Duke Math. J.*, vol. 22, pp. 465–468, 1955.
- [rudin1962groups] W. Rudin, *Fourier Analysis on Groups*. New York: Interscience, 1962.
- [cramer1939representation] H. Cramér, “On the representation of a function by certain Fourier integrals,” *Trans. Amer. Math. Soc.*, vol. 46, pp. 191–201, 1939.
- [dominguez1940fst] A. G. Dominguez, “The representation of functions by Fourier integrals,” *Duke Math. J.*, vol. 6, pp. 246–255, 1940.