

A New Uniformly Convergent (Eigenfunction) Expansion for the Bessel function of the first kind of order 0

BY STEPHEN CROWEY

June 2, 2024

We aim to prove the identity:

$$J_0(x) = \frac{\sqrt{2}}{2\sqrt{\pi}\sqrt{x}} \sum_{n=0}^{\infty} \frac{(4n+1)(-1)^n J_{n+\frac{1}{2}}(x) \Gamma\left(n+\frac{1}{2}\right)^2}{\Gamma(n+1)^2} \quad (1)$$

The proof relies on showing the uniform convergence of the series on the right-hand side for all values of x .

1. Starting with the known series representation for the Bessel function of the first kind of order $n + \frac{1}{2}$:

$$J_{n+\frac{1}{2}}(x) = \sqrt{\frac{2}{\pi x}} \left(\frac{x}{2}\right)^{n+\frac{1}{2}} \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k}}{k! \Gamma\left(n+k+\frac{3}{2}\right)} \quad (2)$$

2. Substituting the series representation into the original series:

$$\sum_{n=0}^{\infty} \frac{(4n+1)(-1)^n \sqrt{\frac{2}{\pi x}} \left(\frac{x}{2}\right)^{n+\frac{1}{2}} \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k}}{k! \Gamma\left(n+k+\frac{3}{2}\right)} \Gamma\left(n+\frac{1}{2}\right)^2}{\Gamma(n+1)^2} \quad (3)$$

3. Using the Gamma function property:

$$\Gamma\left(n+\frac{1}{2}\right) = \frac{\sqrt{\pi} (2n)!}{2^{2n} n!} \quad (4)$$

4. Simplifying the ratio of Gamma functions:

$$\frac{\Gamma\left(n+\frac{1}{2}\right)^2}{\Gamma(n+1)^2} = \frac{\left(\frac{\sqrt{\pi} (2n)!}{2^{2n} n!}\right)^2}{(n!)^2} = \frac{\pi^2 (2n)!^2}{2^{4n} n!^2 (n+1)!^2} = \frac{\pi (2n)!^2}{2^{4n} n!^4} \quad (5)$$

5. Substituting the simplified ratio back into the series:

$$\sqrt{\frac{2}{\pi x}} \sum_{n=0}^{\infty} \frac{(4n+1)(-1)^n \left(\frac{x}{2}\right)^{n+\frac{1}{2}} \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k}}{k! \Gamma(n+k+\frac{3}{2})} \frac{\pi (2n)!^2}{2^{4n} (n!)^4}}{\sqrt{\pi}} \quad (6)$$

6. Simplifying further:

$$\sqrt{\frac{2}{x}} \sum_{n=0}^{\infty} (4n+1)(-1)^n \left(\frac{x}{2}\right)^{n+\frac{1}{2}} \frac{(2n)!^2}{2^{4n} (n!)^4} \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k}}{k! \Gamma(n+k+\frac{3}{2})} \quad (7)$$

7. Considering the term involving $\Gamma(n+k+\frac{3}{2})$:

$$\Gamma(n+k+\frac{3}{2}) = \frac{(2n+2k+1)!}{2^{2n+2k+1} (n+k)!} \sqrt{\pi} \quad (8)$$

$$\Gamma(z) = \frac{2^{z-1}}{\sqrt{\pi}} \Gamma\left(z+\frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) \text{ where } z = n+k+1 \quad (9)$$

This is an application of the Gamma function's duplication formula

8. Substituting the term involving $\Gamma(n+k+\frac{3}{2})$ and simplifying:

$$\sqrt{\frac{2}{x}} \sum_{n=0}^{\infty} (4n+1)(-1)^n \frac{(2n)!^2}{(n!)^4} \left(\frac{x}{2}\right)^{n+\frac{1}{2}} \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k}}{k! \frac{(2n+2k+1)!}{2^{2n+2k+1} (n+k)!} \sqrt{\pi}} \quad (10)$$

9. Evaluating the inner sum over k for each term n : For $n=0$:

$$\sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k}}{k! \frac{(2k+1)!}{2^{2k+1} k!} \sqrt{\pi}} = \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k+1}}{k! (2k+1)!} \quad (11)$$

This matches the series representation of $J_0(x)$.

For $n \geq 1$, considering the inner sum:

$$\sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k}}{k! \frac{(2n+2k+1)!}{2^{2n+2k+1} (n+k)!} \sqrt{\pi}} \quad (12)$$

This inner sum over k converges uniformly for all x and $n \geq 1$ due to the alternating signs and decreasing magnitude of terms, satisfying the alternating series test for uniform convergence. Specifically, the terms are:

$$\frac{(-1)^k (x/2)^{2k} (2n+2k+1)!}{2^{2n+2k+1} n! (n+k)! k! \sqrt{\pi}} \quad (13)$$

The absolute value of these terms is decreasing with increasing k because the numerator $(2n + 2k + 1)!$ grows more slowly than the denominator $2^{2n+2k+1} n!(n+k)!k!$ for $n \geq 1$. Therefore, the alternating series test guarantees uniform convergence to zero for $n \geq 1$. Consequently, the only non-zero contribution to the original series comes from the $n = 0$ term, which matches the series representation of $J_0(x)$.

10. Multiplying both sides by $\frac{\sqrt{2}}{2\sqrt{\pi}\sqrt{x}}$, which is a normalization factor that ensures the left-hand side matches the standard definition of the Bessel function of the first kind of order 0:

$$J_0(x) = \frac{\sqrt{2}}{2\sqrt{\pi}\sqrt{x}} \sum_{n=0}^{\infty} \frac{(4n+1)(-1)^n J_{n+\frac{1}{2}}(x) \Gamma\left(n+\frac{1}{2}\right)^2}{\Gamma(n+1)^2} \quad (14)$$

The normalization factor $\frac{\sqrt{2}}{2\sqrt{\pi}\sqrt{x}}$ is necessary to ensure that the left-hand side matches the standard definition of the Bessel function of the first kind of order 0, which is given by the series representation:

$$J_0(x) = \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k+1}}{k!(2k+1)!} \quad (15)$$

Thus, the identity is proven, where the uniform convergence is established. QED