

so assume it is and show me the fucking inverse for F u fuck

Certainly. Let's be precise and explicit.

We assume that the random measure \$ Z(\omega) \$ is absolutely continuous with respect to Lebesgue measure, so that

$$dZ(\omega) = F(\omega) d\omega$$

where \$ F(\omega) \$ is a (complex-valued) random process, the "spectral process density."

The oscillatory process is then

$$X(t) = \int_{-\infty}^{\infty} \phi_t(\omega) \, F(\omega) \, d\omega,$$

Explicit Inversion Formula for \$ F(\omega) \$

Since $(\phi_t(\omega))_{t\in\mathbb{R}}\$ form an orthonormal basis, you invert by integrating $X(t)\$ against the conjugate basis. Specifically,

$$F(\omega) = \int_{-\infty}^{\infty} X(t) \, \overline{\phi_t(\omega)} \, dt.$$

Proof:

Plug this back into the forward formula:

$$\begin{split} \int_{-\infty}^{\infty} X(t) \overline{\phi_t(\omega)} dt &= \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \phi_t(\lambda) F(\lambda) d\lambda \right) \overline{\phi_t(\omega)} dt \\ &= \int_{-\infty}^{\infty} F(\lambda) \left(\int_{-\infty}^{\infty} \phi_t(\lambda) \overline{\phi_t(\omega)} dt \right) d\lambda \\ &= \int_{-\infty}^{\infty} F(\lambda) \delta(\lambda - \omega) d\lambda \\ &= F(\omega). \end{split}$$

Where we used Fubini's theorem and the orthonormality:

$$\int_{-\infty}^{\infty}\phi_t(\lambda)\overline{\phi_t(\omega)}dt=\delta(\lambda-\omega).$$

Summary Table

Forward (synthesis)	Inverse (analysis)
\$ X(t) = \int \phi_t(\omega) F(\omega) d\omega \$	\$ F(\omega) = \int X(t) \overline{\phi_t(\omega)} dt \$

Special Cases

• **Stationary case:** \$\phi_t(\omega) = e^{i\omega t} \$ gives the usual Fourier inversion:

$$X(t)=\int_{-\infty}^{\infty}e^{i\omega t}F(\omega)d\omega,\quad F(\omega)=\int_{-\infty}^{\infty}X(t)e^{-i\omega t}dt.$$

• Oscillatory case: all above formulas hold for any orthonormal time–frequency family.

Final Answer:

If \$ dZ(\omega) \$ is absolutely continuous, the inverse for \$ F(\omega) \$ is

$$F(\omega) = \int_{-\infty}^{\infty} X(t) \, \overline{\phi_t(\omega)} \, dt$$

where \$ \phi_t(\omega) \$ is the oscillatory orthonormal family used in the forward representation.