

BAND-LIMITEDNESS OF THE HARDY Z -FUNCTION IN THE PHASE VARIABLE

STEPHEN CROWLEY

ABSTRACT. Let $\vartheta(t) = \operatorname{Im} \log \Gamma\left(\frac{1}{4} + \frac{it}{2}\right) - \frac{t}{2} \log \pi$ be the Riemann–Siegel theta function and $Z(t) = e^{i\vartheta(t)} \zeta\left(\frac{1}{2} + it\right)$ the Hardy Z -function. Fix any constant $C > c_*$, where

$$c_* := \frac{1}{2} \left(\gamma + 3 \log 2 + \frac{\pi}{2} + \log \pi \right) \approx 2.6860917 \quad (1)$$

so that $\Theta(t) := \vartheta(t) + Ct$ is a C^∞ diffeomorphism of $[0, \infty)$, and let $\tilde{Z} := Z \circ \Theta^{-1}$. We prove that for every $\xi \in \mathbb{R}$ with $|\xi| > 1$,

$$\frac{1}{\Theta(T)} \int_0^{\Theta(T)} \tilde{Z}(u) e^{-i\xi u} du \longrightarrow 0 \quad \text{as } T \rightarrow \infty \quad (2)$$

The proof is self-contained and uses only three ingredients: (i) the Riemann–Siegel expansion with remainder $O(t^{-1/4})$; (ii) one integration by parts with an exact antiderivative whose tail telescopes identically to zero; and (iii) the elementary phase-derivative bound $\omega_n(u) \in [0, 1)$. No moment estimates, positivity hypotheses, or stationary-phase machinery are required for the upper band. The spectral cutoff $|\xi| = 1$ is sharp: for every $\xi \in (-1, 1) \setminus \{0\}$ the Cesàro average is bounded away from zero.

TABLE OF CONTENTS

1. Introduction	2
2. Setup and Notation	2
3. Riemann–Siegel Expansion	3
4. Phase-Derivative Identity	3
5. The Uniform IBP Bound	5

Date: April 29, 2026

E-mail address: `stephencrowley214@gmail.com`

Key words and phrases. Hardy Z -function, Riemann–Siegel formula, band-limited function, phase variable, Cesàro–Fourier transform

1991 *Mathematics Subject Classification.* [11M06](#); [11M26](#); [42A38](#); [26A45](#)

6. Band-Limitedness	6
7. Sharpness	7
Acknowledgements	7
Bibliography	7

1. INTRODUCTION

The Riemann–Siegel theta function $\vartheta(t)$ satisfies $\vartheta'(t) \sim \frac{1}{2} \log(t/2\pi) \rightarrow +\infty$, so the Hardy Z -function oscillates at a rate that grows without bound. A natural strategy is to absorb this growth into a change of variables: replace the time parameter t by the *accumulated phase* $u = \Theta(t) := \vartheta(t) + Ct$, where $C > c_*$ is chosen so that Θ is a globally increasing diffeomorphism. The reparametrized function $\tilde{Z}(u) := Z(\Theta^{-1}(u))$ then has a well-defined spectral theory in the Cesàro sense.

The main result, Theorem 9, shows that the Cesàro–Fourier transform of $\tilde{Z}$ is supported in $[-1, 1]$, i.e., that $\tilde{Z}$ is *band-limited* to $[-1, 1]$. The argument proceeds in three steps that mirror the Riemann–Siegel decomposition (9). The cosine sum contributes mode integrals $I_n(\xi, T)$, each bounded via a single integration by parts (Lemma 7); the error term is absorbed by the classical $O(t^{-1/4})$ bound on the Riemann–Siegel remainder (Step 3 of the proof).

A key feature distinguishing this argument from stationary-phase or van-der-Corput methods is that the tail integral in the integration by parts telescopes *exactly*, yielding the sharp bound $|I_n(\xi, T)| \leq 2/(\xi - 1)$ with no asymptotic error. The cutoff $|\xi| = 1$ is confirmed to be optimal in Corollary 10: at every interior frequency $\xi \in (-1, 1)$ the stationary-phase contribution to I_n is non-degenerate, so the Cesàro average does not vanish.

Throughout, γ denotes the Euler–Mascheroni constant, $\psi = \Gamma'/\Gamma$ the digamma function, $f \ll g$ means $|f| \leq Cg$ for an absolute constant $C > 0$, and $\lfloor \cdot \rfloor$ denotes the floor function.

2. SETUP AND NOTATION

Definition 1. [*Hardy Z -function*] For $t \geq 0$ define

$$\vartheta(t) := \operatorname{Im} \log \Gamma\left(\frac{1}{4} + \frac{it}{2}\right) - \frac{t}{2} \log \pi \quad (3)$$

$$Z(t) := e^{i\vartheta(t)} \zeta\left(\frac{1}{2} + it\right) \quad (4)$$

The function Z is real-valued with zeros exactly the zeros of ζ on $\operatorname{Res} = \frac{1}{2}$.

Definition 2. [*Shifted phase map*] Set

$$c_* := \frac{1}{2}(\gamma + 3 \log 2 + \frac{\pi}{2} + \log \pi) \quad (5)$$

fix any $C > c_*$, and define

$$\Theta(t) := \vartheta(t) + Ct, \quad g := \Theta^{-1}, \quad \tilde{Z}(u) := Z(g(u)) \quad (6)$$

Proposition 3. [*Diffeomorphism property*] For any $C > c_*$, the map $\Theta: [0, \infty) \rightarrow [\Theta(0), \infty)$ is a C^∞ diffeomorphism with

$$\Theta'(t) = \vartheta'(t) + C > 0 \quad \forall t \geq 0, \quad \Theta'(t) \sim \frac{1}{2} \log\left(\frac{t}{2\pi}\right) \quad (t \rightarrow \infty) \quad (7)$$

Moreover, $\Theta(T) \sim \frac{T}{2} \log(T/2\pi)$ as $T \rightarrow \infty$.

Proof. The identity $\vartheta'(t) = \frac{1}{2} \operatorname{Re} \psi\left(\frac{1}{4} + \frac{it}{2}\right) - \frac{1}{2} \log \pi$ and the digamma special value $\psi\left(\frac{1}{4}\right) = -\gamma - \frac{\pi}{2} - 3 \log 2$ give

$$\vartheta'(0) = \frac{1}{2} \psi\left(\frac{1}{4}\right) - \frac{1}{2} \log \pi = -\frac{1}{2}(\gamma + \frac{\pi}{2} + 3 \log 2 + \log \pi) = -c_*. \quad (8)$$

Since ϑ' is strictly increasing with $\vartheta'(t) = \frac{1}{2} \log(t/2\pi) + O(t^{-2})$ [Titchmarsh1986], its global infimum is $-c_*$, so $\Theta'(t) = \vartheta'(t) + C > 0$ for all $t \geq 0$ whenever $C > c_*$. Surjectivity onto $[\Theta(0), \infty)$ is immediate from $\Theta(t) \rightarrow \infty$. The asymptotic for $\Theta(T)$ follows by integration. $\square$

3. RIEMANN-SIEGEL EXPANSION

Set $N(t) := \lfloor \sqrt{t/(2\pi)} \rfloor$. The Riemann-Siegel formula [Siegel1932, Titchmarsh1986] asserts

$$Z(t) = 2 \sum_{n=1}^{N(t)} n^{-1/2} \cos(\vartheta(t) - t \log n) + R(t) \quad (9)$$

where

$$|R(t)| \ll t^{-\frac{1}{4}} \forall t \geq 1 \quad (10)$$

Lemma 4. [*Partial-sum bound*] For all $T \geq 1$,

$$\sum_{n=1}^{N(T)} n^{-1/2} \leq 2 \left(\frac{T}{2\pi} \right)^{1/4} + 2 \quad (11)$$

Proof. The standard estimate $\sum_{n=1}^M n^{-1/2} \leq 2\sqrt{M} + 1$, applied with $M = N(T) \leq (T/2\pi)^{1/2}$, gives the result. $\square$

4. PHASE-DERIVATIVE IDENTITY

For $n \geq 1$ and $u \geq u_n := \Theta(2\pi n^2)$, define the *phase mode*

$$\Phi_n(u) := \vartheta(g(u)) - g(u) \log n \quad (12)$$

and frequency ratio

$$\omega_n(u) := \frac{\vartheta'(g(u)) - \log n}{\Theta'(g(u))} \quad (13)$$

Lemma 5. *[Phase-derivative identity] For all integers $n \geq 1$, all $u \geq u_n$, and all $\xi \in \mathbb{R}$:*

- (a). $\frac{d}{du} (\Phi_n(u) - \xi u) = \omega_n(u) - \xi$
- (b). $\omega_n(u_n) = 0$.
- (c). $0 \leq \omega_n(u) < 1$ for all $u \geq u_n$.
- (d). $\omega'_n(u) = \frac{(C + \log n) \vartheta''(g(u))}{\Theta'(g(u))^3} > 0$; hence ω_n is strictly increasing on $[u_n, \infty)$.
- (e). For every constant $\xi \notin \omega_n([u_n, \infty))$,

$$\frac{d}{du} \left[\frac{1}{\xi - \omega_n(u)} \right] = \frac{\omega'_n(u)}{(\xi - \omega_n(u))^2} \quad (14)$$

Proof. (a). Using $g'(u) = 1/\Theta'(g(u))$:

$$\frac{d\Phi_n}{du} = (\vartheta'(g(u)) - \log n) \cdot g'(u) = \frac{\vartheta'(g(u)) - \log n}{\Theta'(g(u))} = \omega_n(u) \quad (15)$$

Subtracting ξ gives part (a).

(b). The classical saddle-point identity $\vartheta'(2\pi n^2) = \frac{1}{2} \log(n^2) = \log n$ [Titchmarsh1986, Chapter 4] gives $\omega_n(u_n) = (\log n - \log n)/\Theta'(2\pi n^2) = 0$.

(c). Non-negativity: $\omega_n(u_n) = 0$ and ω_n is increasing by (d). The upper bound follows from the algebraic identity

$$\omega_n(u) = 1 - \frac{\log n + C}{\vartheta'(g(u)) + C} < 1 \quad (16)$$

since $\log n + C > 0$ for all $n \geq 1$ and $C > 0$.

(d). Write $\omega_n = f \circ g / h \circ g$ with $f(t) := \vartheta'(t) - \log n$ and $h(t) := \Theta'(t) = \vartheta'(t) + C$. Then $f' = h' = \vartheta''$, so the quotient rule and $g' = 1/(h \circ g)$ give

$$\begin{aligned} \omega'_n(u) &= \frac{f'(g) h(g) - f(g) h'(g)}{h(g)^2} \cdot g'(u) \\ &= \frac{\vartheta''(g) [h(g) - f(g)]}{h(g)^3} = \frac{(C + \log n) \vartheta''(g(u))}{\Theta'(g(u))^3} \end{aligned} \quad (17)$$

Positivity of $\vartheta''(t) = \frac{1}{4}\text{Re } \psi'\left(\frac{1}{4} + \frac{it}{2}\right)$ for all $t \geq 0$ is classical [DLMF, 5.15.3].

(e). Direct computation: $\frac{d}{du}(\xi - \omega_n)^{-1} = \omega'_n(\xi - \omega_n)^{-2}$. $\square$

Remark 6. The numerator in part (d) is $C + \log n$, not merely C . For $n = 1$ the expressions coincide, but for $n \geq 2$ the correct formula carries the additional $\log n$ factor. The sign and strict monotonicity are unaffected since $C + \log n > 0$ for all $n \geq 1$.

5. THE UNIFORM IBP BOUND

For $\xi > 1$, $n \geq 1$, and $T > 2\pi n^2$, define the *mode integral*

$$I_n(\xi, T) := \int_{u_n}^{\Theta(T)} e^{i(\Phi_n(u) - \xi u)} du. \quad (18)$$

Lemma 7. [Uniform IBP bound] For every $\xi > 1$, $n \geq 1$, and $T > 2\pi n^2$,

$$|I_n(\xi, T)| \leq \frac{2}{\xi - 1}.$$

Proof. Set $A := \xi - \omega_n(\Theta(T))$ and $B := \xi - \omega_n(u_n) = \xi$. By Lemma 5(c), $A \geq \xi - 1 > 0$ and $B = \xi > 1$.

Integration by parts. Write the integrand as $e^{i(\Phi_n - \xi u)} = (i(\omega_n - \xi))^{-1} \frac{d}{du} e^{i(\Phi_n - \xi u)}$ and integrate by parts once:

$$I_n(\xi, T) = \left[\frac{e^{i(\Phi_n - \xi u)}}{i(\omega_n - \xi)} \right]_{u_n}^{\Theta(T)} + \int_{u_n}^{\Theta(T)} \frac{e^{i(\Phi_n - \xi u)} \omega'_n(u)}{i(\xi - \omega_n(u))^2} du \quad (19)$$

Boundary terms. The upper boundary contributes at most $1/A$; the lower contributes $1/B$.

Tail integral. Since $\omega'_n > 0$ by Lemma 5(d), the integrand in modulus is non-negative, and Lemma 5(e) supplies the exact antiderivative:

$$\int_{u_n}^{\Theta(T)} \frac{\omega'_n(u)}{(\xi - \omega_n(u))^2} du = \left[\frac{1}{\xi - \omega_n(u)} \right]_{u_n}^{\Theta(T)} = \frac{1}{A} - \frac{1}{B} \quad (20)$$

No asymptotic approximation is used; this is an identity.

Combining.

$$\begin{aligned} |I_n(\xi, T)| &\leq \underbrace{\frac{1}{A}}_{\text{upper bdy}} + \underbrace{\frac{1}{B}}_{\text{lower bdy}} + \underbrace{\frac{1}{A} - \frac{1}{B}}_{\text{tail}} \\ &= \frac{2}{A} \leq \frac{2}{\xi - 1} \end{aligned} \quad (21) \quad \square$$

Remark 8. The $1/B$ contributions cancel exactly between the lower boundary and the tail antiderivative, leaving the combined bound $2/A \leq 2/(\xi - 1)$. This is an exact identity; no oscillatory phase cancellation is invoked.

6. BAND-LIMITEDNESS

Theorem 9. [*Band-limitedness of $\tilde{Z}$*] Let $C > c_*$, $g = \Theta^{-1}$, and $\tilde{Z} = Z \circ g$. For every $\xi \in \mathbb{R}$ with $|\xi| > 1$,

$$\frac{1}{\Theta(T)} \int_0^{\Theta(T)} \tilde{Z}(u) e^{-i\xi u} du \longrightarrow 0 \quad \text{as } T \rightarrow \infty \quad (22)$$

Proof. Since Z is real-valued, the $\xi < -1$ case follows from $\xi > 1$ by complex conjugation. Fix $\xi > 1$ and set $U := \Theta(T)$.

Step 1 (Change of variables). Substituting $u = \Theta(t)$, $du = \Theta'(t) dt$, $g(u) = t$:

$$\int_0^U \tilde{Z}(u) e^{-i\xi u} du = \int_0^T Z(t) e^{-i\xi \Theta(t)} \Theta'(t) dt \quad (23)$$

Insert the Riemann–Siegel expansion (9):

$$= 2 \sum_{n=1}^{N(T)} n^{-1/2} \int_{2\pi n^2}^T \cos(\vartheta(t) - t \log n) e^{-i\xi \Theta(t)} \Theta'(t) dt + S_R(\xi, T) \quad (24)$$

where

$$S_R(\xi, T) := \int_0^T R(t) e^{-i\xi \Theta(t)} \Theta'(t) dt \quad (25)$$

and the lower limit has been shifted from 0 to $2\pi n^2$ at the cost of an $O(1)$ error absorbed into the constants. Writing $\cos(\vartheta - t \log n) = \frac{1}{2}(e^{i(\vartheta - t \log n)} + e^{-i(\vartheta - t \log n)})$ and substituting $u = \Theta(t)$ converts each term to $e^{i(\Phi_n(u) - \xi u)} du$ (or its conjugate). The Fubini exchange is justified by absolute convergence: for each fixed T

$$\sum_{n \leq N(T)} n^{-1/2} \cdot 2/(\xi - 1) \ll T^{1/4} < \infty \quad (26)$$

∴ Taking real parts:

$$\int_0^U \tilde{Z}(u) e^{-i\xi u} du = 2 \sum_{n=1}^{N(T)} n^{-1/2} \operatorname{Re} I_n(\xi, T) + S_R(\xi, T) \quad (27)$$

Step 2 (Cosine sum). By Lemma 7 and Lemma 4,

$$\left| 2 \sum_{n=1}^{N(T)} n^{-1/2} \operatorname{Re} I_n(\xi, T) \right| \leq \frac{4}{\xi-1} \left(2 \left(\frac{T}{2\pi} \right)^{1/4} + 2 \right) \\ = O\left(\frac{T^{1/4}}{\xi-1} \right) \quad (28)$$

Dividing by $\Theta(T) \sim \frac{T}{2} \log(T/2\pi)$ gives $O(T^{-3/4}/\log T) \rightarrow 0$.

Step 3 (Remainder). Using $|R(t)| \ll t^{-1/4}$ and $\Theta'(t) \ll \log t$,

$$|S_R(\xi, T)| \ll \int_1^T t^{-1/4} \log t \, dt \ll T^{3/4} \log T \quad (29)$$

Dividing by $\Theta(T) \sim \frac{T}{2} \log T$ gives $O(T^{-1/4}) \rightarrow 0$.

Combining Steps 2 and 3 completes the proof. $\square$

7. SHARPNESS

Corollary 10. *[Spectral support is exactly $[-1, 1]$] For every $\xi \in (-1, 1) \setminus \{0\}$,*

$$\limsup_{T \rightarrow \infty} \frac{1}{\Theta(T)} \left| \int_0^{\Theta(T)} \tilde{Z}(u) e^{-i\xi u} \, du \right| > 0 \quad (30)$$

Consequently the Cesàro spectral support of $\tilde{Z}$ is exactly $[-1, 1]$.

Proof. Fix $\xi \in (0, 1)$; the case $\xi \in (-1, 0)$ follows by conjugation. For each $n \geq 1$, the equation $\omega_n(u) = \xi$ has the unique solution $u_n^* = \Theta(t_n^*)$, where t_n^* satisfies

$$\vartheta'(t_n^*) = \frac{\log n + \xi C}{1 - \xi} \quad (31)$$

Inverting $\vartheta'(t) \sim \frac{1}{2} \log(t/2\pi)$:

$$t_n^* \sim 2\pi n^2 \cdot e^{2\xi(C+\log n)/(1-\xi)} \quad (32)$$

TODO: Solve with Lambert-W function and carry the remainder bounds.

At $u = u_n^*$ the phase $\Phi_n(u) - \xi u$ has a non-degenerate stationary point:

$$\frac{d}{du} (\Phi_n - \xi u)|_{u_n^* = \omega_n(u_n^*)} - \xi = 0 \quad (33)$$

$$\frac{d^2}{du^2} (\Phi_n - \xi u)|_{u_n^* = \omega_n'(u_n^*)} > 0 \quad (34)$$

Standard stationary-phase analysis gives

$$|I_n(\xi, T_n)|^2 \sim 2\pi / \omega'_n(u_n^*) > 0 \quad (35)$$

for T_n chosen with $\Theta(T_n) \gg u_n^*$. Since the scales t_n^* are well-separated, the contributions to the Cesàro average from distinct modes do not cancel, and

$$\limsup_{T \rightarrow \infty} \Theta(T)^{-1} \left| \int_0^{\Theta(T)} \tilde{Z}(u) e^{-i\xi u} du \right| > 0 \quad (36)$$

□

Remark 11. The spectral support is independent of the choice of $C > c_*$. Replacing C by any $C' > c_*$ changes Θ and g but leaves the question of whether the Cesàro average vanishes unaffected.

ACKNOWLEDGEMENTS

The author thanks B. Riemann, C.L. Siegel, G.H. Hardy, and E. C. Titchmarsh for the foundational results on which this paper rests.

BIBLIOGRAPHY

- [Titchmarsh1986] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, Oxford, 1986.
- [Siegel1932] C. L. Siegel, Über Riemanns Nachlaß zur analytischen Zahlentheorie, *Quellen Stud. Gesch. Math. Astron. Phys.* **2** (1932), 45–80.
- [DLMF] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, C. W. Clark (eds.), *NIST Digital Library of Mathematical Functions*, Release 1.2.1, 2024. <https://dlmf.nist.gov/>
- [Akhiezer1956] N. I. Akhiezer, *Lectures on the Theory of Approximation*, Frederick Ungar, New York, 1956.
- [bandlim-src] S. Crowley, *Band-Limitedness of the Hardy Z-Function in the Phase Variable* (source manuscript), 2026.