

The Riemann–Hilbert View of Padé Approximation and Global Analytic Continuation, with Application to the Fractional Riccati Equation

Abstract

The diagonal Padé approximant of a function analytic at the origin admits a representation as the solution to a 2×2 matrix Riemann–Hilbert problem. This formulation replaces the local derivative-matching of Taylor approximation with the global reconstruction of a spectral measure from its moments. The convergence domain is determined not by the distance to the nearest singularity, but by the maximal region on which the associated Riemann–Hilbert factorization remains solvable, a region governed by the equilibrium measure of an extremal contour selected by the S -property. The poles of the approximant are the eigenvalues of a truncated Jacobi operator, and their condensation along the equilibrium contour is the signature of the underlying Riemann surface. We present the general theory in the form of theorems, corollaries, and limit statements, identify the precise hypotheses under which each conclusion holds, and apply the framework to the constant-coefficient fractional Riccati equation that governs the rough Heston characteristic function in mathematical finance. The application section develops the Müntz spectral solution, derives a Gamma-ratio convolution recurrence for the Puiseux coefficients, establishes a strictly positive radius of convergence by a Banach fixed-point argument, constructs the diagonal Padé resummation in the variable $z = t^\mu$, proves meromorphy of the transformed solution on $\mathbb{C}$ by a classical Riccati majorization with a sharp Gautschi-type constant and a z_* -dependent residue formula, and proves global convergence on the positive real axis, with a computable a-posteriori error bound.

Contents

1	Introduction	1
2	The Riemann–Hilbert formulation	2
3	The contour is determined, not given	3
4	Froissart doublets and convergence in capacity	4
5	The Jacobi operator and the Toda lattice	5
6	Pole condensation: orthogonal polynomials and Fekete points	6

7	The Borel plane and resummation	7
8	Application to rough Heston: the constant-coefficient fractional Riccati equation	8
8.1	Fractional calculus and the Müntz basis	8
8.2	The series solution and Volterra representation	9
8.3	Radius of convergence and generic finiteness	11
8.4	Padé resummation in the variable $z = t^\mu$	12
8.5	Remainder representation and the computable error bound	13
8.6	Meromorphy on $\mathbb{C}$ via classical Riccati majorization	14
8.7	Global convergence on the positive real axis	18
8.8	The bordered rank-one update and incremental Hankel solver	19
8.9	Spectral interpretation and operator-theoretic synthesis	20
8.10	Algorithmic summary	21
9	The essential difference	22

1 Introduction

Taylor approximation asks for the polynomial that matches the derivatives of a function at a single point. It is local, and its radius of convergence is the distance to the nearest singularity. Padé approximation asks for the spectral measure that generates the given moment sequence, and for the Stieltjes transform of that measure. It is global, and its convergence domain is the maximal region on which the Riemann–Hilbert factorization exists.

The distinction is not merely one of degree. A Taylor polynomial of degree $2n$ and a diagonal Padé approximant $[n/n]$ both consume the same $2n + 1$ coefficients, yet they encode fundamentally different information. The Taylor polynomial encodes derivatives at a point; the Padé approximant encodes moments of a measure. The first is governed by local analyticity; the second by global potential theory. The convergence theory of the two paradigms reflects this distinction at every level.

The document proceeds as follows. Section 2 presents the Fokas–Its–Kitaev Riemann–Hilbert formulation for diagonal Padé approximants. Section 3 describes Stahl’s geometric theorem and the selection of the equilibrium contour by the S -property, together with the Deift–Zhou steepest descent asymptotic. Section 4 explains why convergence holds in capacity rather than uniformly, through the mechanism of Froissart doublets, and gives the Gončar condition under which uniform convergence is recovered. Section 5 identifies the Padé table with the Toda lattice and the Jacobi operator resolvent, including the non-Hermitian extension via pseudo-spectrum. Section 6 contrasts orthogonal polynomial zeros with Fekete points and gives bulk and edge residue estimates. Section 7 discusses the Borel plane and Padé–Borel resummation for factorially divergent series. Section 8 develops the application to the rough Heston model in detail, treating the constant-coefficient fractional Riccati equation through the Müntz spectral method and Padé resummation in $z = t^\mu$.

Section 9 summarizes the essential difference between the two paradigms.

2 The Riemann–Hilbert formulation

We begin with the foundational equivalence between diagonal Padé approximants and matrix Riemann–Hilbert problems. This equivalence, due to Fokas, Its, and Kitaev, recasts a problem of rational approximation as a problem of matrix factorization on a contour.

Definition 2.1 (Cauchy transform). Let $\Sigma \subset \mathbb{C}$ be a finite union of smooth oriented arcs, and let $w : \Sigma \rightarrow \mathbb{C}$ be a measurable weight with $\int_{\Sigma} |w(s)| |ds| < \infty$. The *Cauchy transform* of w on Σ is the function

$$f(z) = \int_{\Sigma} \frac{w(s) ds}{z - s}, \quad z \in \mathbb{C} \setminus \Sigma. \quad (1)$$

Definition 2.2 (Diagonal Padé approximant). Let f be analytic at infinity with formal expansion $f(z) = \sum_{k=0}^{\infty} a_k z^{-k-1}$. The *diagonal Padé approximant* $[n/n]_f$ of order n is the unique rational function P_n/Q_n with $\deg P_n < n$, $\deg Q_n = n$, and Q_n monic, satisfying

$$f(z) - \frac{P_n(z)}{Q_n(z)} = O(z^{-2n-1}) \quad \text{as } z \rightarrow \infty, \quad (2)$$

whenever the underlying Hankel determinant $\tau_n = \det(a_{i+j})_{0 \leq i, j \leq n-1}$ is nonzero.

Theorem 2.3 (Fokas–Its–Kitaev representation). *Let f be the Cauchy transform (1) of a weight w on a fixed contour Σ , and assume the Hankel determinants $\tau_1, \dots, \tau_n$ are nonzero. Then there exists a unique 2×2 matrix-valued function $Y : \mathbb{C} \setminus \Sigma \rightarrow GL_2(\mathbb{C})$ satisfying:*

(i) Y is analytic on $\mathbb{C} \setminus \Sigma$;

(ii) Y has continuous boundary values $Y_{\pm}$ on Σ related by the jump

$$Y_+(s) = Y_-(s) \begin{pmatrix} 1 & 2\pi i w(s) \\ 0 & 1 \end{pmatrix}, \quad s \in \Sigma; \quad (3)$$

(iii) Y has the asymptotic behavior at infinity

$$Y(z) = \left(I + \frac{Y_1}{z} + \frac{Y_2}{z^2} + \dots \right) \begin{pmatrix} z^n & 0 \\ 0 & z^{-n} \end{pmatrix}, \quad z \rightarrow \infty. \quad (4)$$

The entries of Y encode the diagonal Padé approximant via

$$Y_{11}(z) = Q_n(z), \quad Y_{12}(z) = \int_{\Sigma} \frac{Q_n(s) w(s) ds}{s - z}, \quad (5)$$

and the approximant itself is

$$[n/n]_f(z) = \frac{P_n(z)}{Q_n(z)} = \frac{Y_{12}(z)}{Y_{11}(z)}. \quad (6)$$

Corollary 2.4 (Orthogonality). *The denominator Q_n is the monic orthogonal polynomial of degree n for the moment functional $\mathcal{L}[p] = \int_{\Sigma} p(s)w(s) ds$:*

$$\int_{\Sigma} Q_n(s)s^k w(s) ds = 0, \quad k = 0, 1, \dots, n-1. \quad (7)$$

Remark 2.5 (Limitation: the contour must be given). Theorem 2.3 requires a fixed contour Σ . For functions specified only by a power series, with branch cuts whose location is unknown, the contour is not given; it must be determined as part of the problem. The Riemann–Hilbert formulation in the form stated above becomes exact only after the equilibrium contour has been selected by the procedure of Section 3.

3 The contour is determined, not given

For functions whose singular structure is not specified in advance, the appropriate contour is selected by an extremal principle in logarithmic potential theory. This is the content of Stahl’s geometric theorem.

Definition 3.1 (Stahl class). A function f analytic at infinity belongs to the *Stahl class* $\mathcal{S}$ if it admits multi-valued meromorphic continuation to a domain $\overline{\mathbb{C}} \setminus E$, where E is a compact set of logarithmic capacity zero.

Definition 3.2 (Admissible compact). A compact set $K \subset \mathbb{C}$ is *admissible* for $f \in \mathcal{S}$ if $\overline{\mathbb{C}} \setminus K$ is a domain on which f admits single-valued meromorphic continuation.

Theorem 3.3 (Stahl’s geometric theorem). *For every $f \in \mathcal{S}$, there exists a unique admissible compact Δ_f with the following properties:*

- (i) $\overline{\mathbb{C}} \setminus \Delta_f$ is a connected domain D_f ;
- (ii) f has single-valued meromorphic continuation to D_f ;
- (iii) $\text{cap}(\Delta_f) \leq \text{cap}(K)$ for every other admissible compact K .

The set Δ_f is called the Stahl compact or S -compact of f .

Theorem 3.4 (The S -property). *Let $g(z, \infty)$ denote the Green’s function of D_f with pole at infinity. The arcs of Δ_f satisfy the symmetry condition*

$$\frac{\partial g}{\partial n_+}(z) = \frac{\partial g}{\partial n_-}(z), \quad z \in \Delta_f \setminus \{\text{endpoints}\}, \quad (8)$$

where $n_{\pm}$ are the two unit normals to the arc.

Remark 3.5 (Variational interpretation). The S -property is the Euler–Lagrange equation of a min-max variational problem. Among admissible compacts in a fixed homotopy class, the minimum of $\text{cap}(K)$ is attained at Δ_f , and the harmonic measure exerts equal pull on both sides of each arc.

Theorem 3.6 (Stahl's convergence theorem). *Let $f \in \mathcal{S}$ with Stahl compact Δ_f and complementary domain D_f . The diagonal Padé approximants converge to f in capacity on compact subsets of $D_f \setminus \mathcal{P}_f$, where $\mathcal{P}_f$ is the pole set of f . For every compact $K \subset D_f \setminus \mathcal{P}_f$ and every $\varepsilon > 0$,*

$$\lim_{n \rightarrow \infty} \text{cap}\{z \in K : |f(z) - [n/n]_f(z)| > \varepsilon\} = 0. \quad (9)$$

Theorem 3.7 (Deift–Zhou asymptotic). *Under the hypotheses of Theorem 3.6, with the additional regularity that w is analytic in a neighborhood of Δ_f and the endpoints exhibit square-root behavior of the equilibrium density, nonlinear steepest descent yields*

$$[n/n]_f(z) = f(z)(1 + O(e^{-ng(z, \infty)})) \quad (10)$$

uniformly on compact subsets of D_f bounded away from $\Delta_f \cup \mathcal{P}_f$.

Corollary 3.8 (Exponential rate of convergence).

$$\limsup_{n \rightarrow \infty} |f(z) - [n/n]_f(z)|^{1/n} \leq e^{-g(z, \infty)}, \quad z \in D_f \setminus \mathcal{P}_f. \quad (11)$$

The level sets $\{g(z, \infty) = c\}$ are loci of equal convergence rate, and $\Delta_f = \{g = 0\}$.

Remark 3.9 (Comparison with Taylor). The Taylor polynomial of degree $2n$ converges geometrically only inside the disk of convergence, with rate $|z|/r$ where r is the distance to the nearest singularity. The Padé rate $e^{-g(z, \infty)}$ extends to all of D_f , generally a much larger domain. The two rates agree only inside the disk of convergence, where $g(z, \infty) \sim -\log|z| + \text{const}$.

4 Froissart doublets and convergence in capacity

The convergence statement of Theorem 3.6 is in capacity, not uniformly. The reason is the appearance of spurious poles, called Froissart doublets, that wander through the domain of holomorphy without condensing on the Stahl compact.

Definition 4.1 (Froissart doublet). A *Froissart doublet* of $[n/n]_f$ is a pole $z_j^{(n)} \in D_f \setminus \mathcal{P}_f$ accompanied by a zero $\zeta_j^{(n)}$ at distance $|z_j^{(n)} - \zeta_j^{(n)}| = O(e^{-cn})$ for some $c > 0$, with residue

$$\text{Res}([n/n]_f, z_j^{(n)}) = O(e^{-cn}) \quad \text{as } n \rightarrow \infty. \quad (12)$$

Proposition 4.2 (Boundedness of doublet count). *For $f \in \mathcal{S}$, the number of Froissart doublets of $[n/n]_f$ in any compact $K \subset D_f \setminus \mathcal{P}_f$ is $O(1)$ as $n \rightarrow \infty$, but their location may oscillate erratically with n .*

Definition 4.3 (Gončar condition). A function $f \in \mathcal{S}$ satisfies the *Gončar condition* on a compact $K \subset D_f \setminus \mathcal{P}_f$ if the number of poles of $[n/n]_f$ in K is uniformly bounded in n .

Theorem 4.4 (Gončar's uniform convergence theorem). *If $f \in \mathcal{S}$ satisfies the Gončar condition on a compact $K \subset D_f \setminus \mathcal{P}_f$, then $[n/n]_f \rightarrow f$ uniformly on K . The exceptional set of spurious poles is empty for sufficiently large n .*

Remark 4.5. The Gončar condition is restrictive for generic $f \in \mathcal{S}$ because Froissart doublets, though $O(1)$ in number, could in principle visit every compact infinitely often. For meromorphic functions with well-separated poles and a regular Padé table, the condition holds on every compact disjoint from the pole set; this is the situation of Section 8.7.

5 The Jacobi operator and the Toda lattice

The Padé problem admits a third equivalent formulation, in addition to rational approximation and Riemann–Hilbert factorization. This third formulation is operator-theoretic: the Padé approximant is the resolvent of a truncated Jacobi operator.

Definition 5.1 (Hankel matrix and tau function). For a sequence $\{a_k\}_{k \geq 0}$, the *Hankel matrix* of order n is

$$H^{(n)} = (a_{i+j})_{0 \leq i, j \leq n-1}, \quad (13)$$

and its determinant $\tau_n = \det H^{(n)}$ is the *tau function*.

Theorem 5.2 (Toda lattice structure). Assume $\tau_n \neq 0$ for all $n \geq 1$. The monic orthogonal polynomials Q_n for the moment functional defined by $\{a_k\}$ satisfy the three-term recurrence

$$Q_{n+1}(z) = (z - b_n)Q_n(z) - a_n^2 Q_{n-1}(z), \quad Q_{-1} = 0, \quad Q_0 = 1. \quad (14)$$

Under the spectral evolution $d\mu_t(x) = e^{tx} d\mu(x)$, the recurrence coefficients evolve according to the Toda equations

$$\dot{b}_n = a_n^2 - a_{n-1}^2, \quad \dot{a}_n = a_n(b_{n+1} - b_n). \quad (15)$$

The flow is isospectral.

Definition 5.3 (Jacobi operator). The *infinite Jacobi operator* associated to $\{a_n, b_n\}$ is the tridiagonal matrix

$$J_\infty = \begin{pmatrix} b_0 & a_0 & 0 & \cdots \\ a_0 & b_1 & a_1 & \ddots \\ 0 & a_1 & b_2 & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix} \quad (16)$$

acting on $\ell^2(\mathbb{N}_0)$. Its $n \times n$ *principal truncation* is $J_n = [J_\infty]_{0 \leq i, j \leq n-1}$.

Theorem 5.4 (Resolvent representation, self-adjoint case). Assume the moment functional is positive, so that J_∞ is self-adjoint and bounded with spectral measure μ supported on a compact subset of $\mathbb{R}$. Then

$$f(z) = \langle e_0, (J_\infty - zI)^{-1} e_0 \rangle = \int \frac{d\mu(x)}{x - z}, \quad (17)$$

and

$$[n/n]_f(z) = \langle e_0, (J_n - zI)^{-1} e_0 \rangle = \sum_{j=1}^n \frac{|\langle e_0, \psi_j^{(n)} \rangle|^2}{\lambda_j^{(n)} - z}, \quad (18)$$

where $\{\lambda_j^{(n)}, \psi_j^{(n)}\}_{j=1}^n$ are the eigenvalue-eigenvector pairs of J_n .

Corollary 5.5 (Poles as eigenvalues). The poles of $[n/n]_f$ are precisely the eigenvalues of J_n , and the residues are the squared first components of the corresponding normalized eigenvectors.

Theorem 5.6 (Strong resolvent convergence). *In the setting of Theorem 5.4, $J_n \rightarrow J_\infty$ in the strong resolvent sense:*

$$\lim_{n \rightarrow \infty} (J_n - zI)^{-1} \xi = (J_\infty - zI)^{-1} \xi \quad \forall \xi \in \ell^2, z \notin \text{spec}(J_\infty). \quad (19)$$

Consequently, $[n/n]_f(z) \rightarrow f(z)$ for every $z \in \mathbb{C} \setminus \text{spec}(J_\infty)$, locally uniformly.

Corollary 5.7 (Self-adjoint convergence domain). *For real measures, the convergence domain of the diagonal Padé approximation coincides with $\mathbb{C} \setminus \text{spec}(J_\infty)$, and $\Delta_f = \text{spec}(J_\infty)$.*

Remark 5.8 (Non-Hermitian extension). For complex measures and functions with branch points off the real axis, J_∞ is non-Hermitian and Theorem 5.6 does not apply directly. The relevant convergence theory uses the pseudo-spectrum

$$\text{spec}_\varepsilon(J_\infty) = \{z \in \mathbb{C} : \|(J_\infty - zI)^{-1}\| \geq \varepsilon^{-1}\}. \quad (20)$$

Theorem 5.9 (Pseudo-spectral limit, regular case). *Suppose the recurrence coefficients $\{a_n, b_n\}$ admit a regular asymptotic expansion in $1/n$ and the limiting density is supported on a finite union of analytic arcs. Then*

$$\lim_{\varepsilon \rightarrow 0} \partial \text{spec}_\varepsilon(J_\infty) = \Delta_f. \quad (21)$$

Remark 5.10 (Beyond the regular case). In less regular non-Hermitian settings, the limiting object of (21) is a quadratic differential trajectory in the sense of Strebel; its identification with Δ_f requires additional symmetry conditions and is established case-by-case.

6 Pole condensation: orthogonal polynomials and Fekete points

Definition 6.1 (Fekete points). The n -th Fekete points of a compact $K \subset \mathbb{C}$ are the maximizers

$$\{z_1^{F,n}, \dots, z_n^{F,n}\} = \arg \max_{(z_1, \dots, z_n) \in K^n} \prod_{i < j} |z_i - z_j|. \quad (22)$$

Theorem 6.2 (Equilibrium limit of Fekete points). *The normalized counting measures of the Fekete points of K converge weak-* to the equilibrium measure μ_K :*

$$\frac{1}{n} \sum_{j=1}^n \delta_{z_j^{F,n}} \xrightarrow{w^*} \mu_K \quad \text{as } n \rightarrow \infty. \quad (23)$$

Theorem 6.3 (Equilibrium limit of orthogonal polynomial zeros). *For $f \in \mathcal{S}$ with Stahl compact Δ_f , the normalized counting measures of the zeros of Q_n converge weak-* to the equilibrium measure μ_{Δ_f} :*

$$\frac{1}{n} \sum_{j=1}^n \delta_{\zeta_j^{(n)}} \xrightarrow{w^*} \mu_{\Delta_f}. \quad (24)$$

Remark 6.4 (Distinct objects, common limit). Theorems 6.2 and 6.3 share a conclusion but describe different sequences. Fekete points are extremal for discrete logarithmic energy and have no recursive relationship across n . Orthogonal polynomial zeros are determined by the spectral truncation of the Jacobi operator and are linked across n by the three-term recurrence (14). Only their normalized counting measures agree in the limit.

Theorem 6.5 (Bulk residue estimate). *Let $f \in \mathcal{S}$ satisfy the regularity conditions of Theorem 3.7. For poles $z_j^{(n)}$ approaching the interior of Δ_f ,*

$$\text{Res}([n/n]_f, z_j^{(n)}) = \frac{P_n(z_j^{(n)})}{Q'_n(z_j^{(n)})} = O\left(\frac{1}{n}\right), \quad (25)$$

uniformly for poles bounded away from the endpoints.

Theorem 6.6 (Edge residue scaling). *For poles within an $O(n^{-2/3})$ neighborhood of an endpoint of Δ_f , the rescaled residues satisfy*

$$n^{1/3} \text{Res}([n/n]_f, z_j^{(n)}) \rightarrow \rho_{\text{Ai}}(\xi_j), \quad \xi_j = n^{2/3}(z_j^{(n)} - \text{endpoint}), \quad (26)$$

where ρ_{Ai} is determined by the Airy kernel. The bulk $1/n$ law is replaced by an Airy-type $n^{-1/3}$ decay at the edge, the universal Painlevé II scaling of random matrix theory.

Corollary 6.7 (Pole-zero cancellation on contours avoiding the cut). *For any rectifiable contour $\gamma \subset D_f \setminus \mathcal{P}_f$ enclosing a region disjoint from Δ_f ,*

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi i} \oint_{\gamma} [n/n]_f(z) dz = \frac{1}{2\pi i} \oint_{\gamma} f(z) dz. \quad (27)$$

7 The Borel plane and resummation

Definition 7.1 (Borel transform). *For a formal power series $\hat{f}(z) = \sum_{n=0}^{\infty} a_n z^n$, the Borel transform is*

$$B(t) = \sum_{n=0}^{\infty} \frac{a_n}{n!} t^n. \quad (28)$$

Theorem 7.2 (Inverse Borel transform). *If $B(t)$ admits analytic continuation to a sector $|\arg t| < \theta$ with at most exponential growth, the Laplace integral*

$$f_{\text{Borel}}(z) = \int_0^{\infty} e^{-t/z} B(t) dt \quad (29)$$

defines a holomorphic function in a sector of the z -plane.

Theorem 7.3 (Padé–Borel resummation). *Let $\hat{f}$ have factorial coefficient growth $a_n \sim n!$, and assume $B(t)$ is in Stahl’s class with no positive real singularities. The composition*

$$f_{\text{P-B}}(z) = \int_0^{\infty} e^{-t/z} [n/n]_B(t) dt \quad (30)$$

converges in capacity to $f_{\text{Borel}}(z)$ on the Borel domain of regularity, and assigns a finite value to a series whose original radius of convergence was zero.

Remark 7.4 (Resummation versus continuation). Padé–Borel resummation is distinct from analytic continuation: the latter extends a function across a singularity from a region where it already exists, while the former assigns a value to a divergent series. The Padé approximant in (30) approximates the measure in the Borel plane whose Laplace transform is the physical quantity.

8 Application to rough Heston: the constant-coefficient fractional Riccati equation

The rough Heston characteristic function reduces, after standard transformations, to the constant-coefficient fractional Riccati equation

$$D^\mu y(t) = c_0 + c_1 y(t) + c_2 y(t)^2, \quad I^{1-\mu} y(0) = 0, \quad (31)$$

where D^μ is the Caputo derivative of order $\mu \in (0, 1]$, $H = \mu - \frac{1}{2}$ is the Hurst parameter, and the coefficients $c_0, c_1, c_2 \in \mathbb{C}$ depend on the model parameters and the Fourier variable. We develop the Müntz spectral solution, the Padé resummation in the variable $z = t^\mu$, and the global convergence theorem with computable error bound.

8.1 Fractional calculus and the Müntz basis

Definition 8.1 (Riemann–Liouville integral and Caputo derivative). For $r > 0$ and $f \in L^1_{\text{loc}}[0, T]$, the *Riemann–Liouville integral* is

$$I^r f(t) = \frac{1}{\Gamma(r)} \int_0^t (t-s)^{r-1} f(s) ds. \quad (32)$$

For $\mu \in (0, 1)$ and f such that $I^{1-\mu} f \in C^1[0, T]$, the *Caputo derivative* is

$$D^\mu f(t) = \frac{d}{dt}(I^{1-\mu} f(t)) - \frac{f(0)}{\Gamma(1-\mu)} t^{-\mu}. \quad (33)$$

The fundamental relations are $I^\mu D^\mu f(t) = f(t) - f(0)$ and $D^\mu I^\mu f(t) = f(t)$.

Proposition 8.2 (Action on power functions). For $s > -1$ and $r > 0$,

$$I^r t^s = \frac{\Gamma(s+1)}{\Gamma(s+r+1)} t^{s+r}. \quad (34)$$

For $\mu \in (0, 1)$ and $s > 0$,

$$D^\mu t^s = \frac{\Gamma(s+1)}{\Gamma(s+1-\mu)} t^{s-\mu}. \quad (35)$$

Proof. For (34), the substitution $u = tx$ in the integral defining $I^r t^s$ yields a Beta function evaluation. For (35), observe that $s > 0$ implies $f(0) = 0$, so the correction term in (33) vanishes; differentiating $I^{1-\mu} t^s = \Gamma(s+1)/\Gamma(s+2-\mu) \cdot t^{s+1-\mu}$ and applying $\Gamma(s+2-\mu) = (s+1-\mu)\Gamma(s+1-\mu)$ gives the result. $\square$

Remark 8.3 (The Caputo derivative of a constant). The hypothesis $s > 0$ in (35) cannot be relaxed. For $s = 0$, the correction term in (33) cancels the formal power $t^{-\mu}/\Gamma(1-\mu)$ and yields $D^\mu(1) = 0$, consistent with the convention that constants have zero Caputo derivative.

Definition 8.4 (Müntz lattice). Fix $\mu \in (0, 1]$ and $T > 0$. The *Müntz lattice* is

$$\mathcal{M} = \text{span}\{1, t^\mu, t^{2\mu}, t^{3\mu}, \dots\}. \quad (36)$$

Theorem 8.5 (Closure of the Müntz lattice). *For every $k \geq 1$ and $r \in \mu\mathbb{N}$:*

$$D^\mu t^{k\mu} = \frac{\Gamma(k\mu + 1)}{\Gamma((k-1)\mu + 1)} t^{(k-1)\mu}, \quad (37)$$

$$I^r t^{k\mu} = \frac{\Gamma(k\mu + 1)}{\Gamma(k\mu + r + 1)} t^{k\mu+r}, \quad (38)$$

$$t^{j\mu} \cdot t^{k\mu} = t^{(j+k)\mu}. \quad (39)$$

Consequently $D^\mu \mathcal{M} \subset \mathcal{M}$, $I^r \mathcal{M} \subset \mathcal{M}$ for $r \in \mu\mathbb{N}$, and $\mathcal{M}$ is closed under multiplication.

Remark 8.6 (Why the Müntz lattice is the natural basis). The three closure properties of Theorem 8.5, under fractional differentiation, fractional integration of order μ , and pointwise multiplication, make the Müntz lattice the canonical basis for any constant-coefficient fractional ODE with polynomial nonlinearity. The Müntz–Tau substitution preserves the algebraic structure of the equation. Ordinary Taylor series in t fail because they are not closed under D^μ for non-integer μ . Note that $I^{1-\mu}$ does *not* preserve $\mathcal{M}$ for generic μ ; however, the Volterra reformulation (40) requires only I^μ , which does.

8.2 The series solution and Volterra representation

Applying I^μ to (31) and using $I^\mu D^\mu y = y - y(0) = y$ yields the equivalent Volterra integral equation

$$y(t) = \frac{1}{\Gamma(\mu)} \int_0^t (t-s)^{\mu-1} (c_0 + c_1 y(s) + c_2 y(s)^2) ds. \quad (40)$$

Remark 8.7 (Hölder regularity at the origin). Evaluating (40) at $t = 0$ gives $y(0) = 0$. Near the origin,

$$y(t) = \frac{c_0}{\Gamma(\mu + 1)} t^\mu + O(t^{2\mu}), \quad (41)$$

so y is Hölder continuous of order μ but not differentiable at 0. The Hölder exponent μ is the natural regularity scale of the problem.

Theorem 8.8 (Müntz–Tau Puiseux series solution). *There exist $R > 0$ and a sequence $(a_k)_{k \geq 1} \subset \mathbb{C}$ such that*

$$y(t) = \sum_{k=1}^{\infty} a_k t^{k\mu}, \quad |t| < R^{1/\mu}, \quad (42)$$

where the coefficients satisfy the Gamma-ratio convolution recurrence

$$a_1 = \frac{c_0}{\Gamma(\mu + 1)}, \quad (43)$$

$$a_{k+1} = \frac{\Gamma(k\mu + 1)}{\Gamma((k+1)\mu + 1)} \left(c_1 a_k + c_2 \sum_{\substack{j+\ell=k \\ 1 \leq j, \ell \leq k-1}} a_j a_\ell \right), \quad k \geq 1. \quad (44)$$

Proof. Existence of $R > 0$ is established in Theorem 8.10 below by a Banach fixed-point argument. Granting analyticity of y in $z = t^\mu$ on $|z| < R$, write $y(t) = \sum_{k \geq 1} a_k t^{k\mu}$. By (39),

$$y(t)^2 = \sum_{m=2}^{\infty} b_m t^{m\mu}, \quad b_m = \sum_{\substack{j+\ell=m \\ 1 \leq j, \ell \leq m-1}} a_j a_\ell. \quad (45)$$

Substituting into (40), splitting the integral, and applying (34) with $s = k\mu$, $r = \mu$:

$$\int_0^t (t-s)^{\mu-1} s^{k\mu} ds = \Gamma(\mu) \frac{\Gamma(k\mu+1)}{\Gamma((k+1)\mu+1)} t^{(k+1)\mu}. \quad (46)$$

Matching the coefficient of $t^{n\mu}$ for each $n \geq 1$ yields (43) for $n = 1$ and (44) after reindexing $n \mapsto k+1$. $\square$

Remark 8.9 (Quadratic convolution at low orders). For $k = 1$, the convolution sum in (44) is empty: there are no pairs (j, ℓ) with $j, \ell \geq 1$ and $j + \ell = 1$. Hence

$$a_2 = \frac{\Gamma(\mu+1)}{\Gamma(2\mu+1)} c_1 a_1. \quad (47)$$

The quadratic term first contributes at $k = 2$:

$$a_3 = \frac{\Gamma(2\mu+1)}{\Gamma(3\mu+1)} (c_1 a_2 + c_2 a_1^2). \quad (48)$$

8.3 Radius of convergence and generic finiteness

Theorem 8.10 (Strictly positive Puiseux radius and Cauchy bound). *Let*

$$A = |c_0| + |c_1| + |c_2|, \quad B = |c_1| + 2|c_2|, \quad (49)$$

and define

$$T_0^\mu = \min\left(\frac{\Gamma(\mu+1)}{2A}, \frac{\Gamma(\mu+1)}{4B}\right) > 0. \quad (50)$$

(i) *There exists a unique solution $y \in C[0, T_0]$ of (40) with $\|y\|_\infty \leq 1$. This solution is analytic in $z = t^\mu$ on $\{|z| < T_0^\mu\}$, so the Puiseux series (42) has radius of convergence $R \geq T_0^\mu > 0$.*

(ii) *For every $0 < r < R$, with $g(z) = y(z^{1/\mu})$ and $M(r) = \sup_{|z|=r} |g(z)|$,*

$$|a_k| \leq \frac{M(r)}{r^k} \quad \forall k \geq 1. \quad (51)$$

Proof. Define $B_1 = \{y \in C[0, T_0] : \|y\|_\infty \leq 1\}$ with the supremum norm and the operator

$$\Phi(y)(t) = \frac{1}{\Gamma(\mu)} \int_0^t (t-s)^{\mu-1} (c_0 + c_1 y(s) + c_2 y(s)^2) ds = I^\mu(c_0 + c_1 y + c_2 y^2)(t). \quad (52)$$

For $y \in B_1$, the integrand is bounded by A , and (34) gives

$$|\Phi(y)(t)| \leq A \cdot \frac{T_0^\mu}{\Gamma(\mu+1)} \leq \frac{1}{2} < 1 \quad (53)$$

by the first term in (50). For $y_1, y_2 \in B_1$,

$$|c_1(y_1 - y_2) + c_2(y_1^2 - y_2^2)| \leq B \|y_1 - y_2\|_\infty, \quad (54)$$

so

$$\|\Phi(y_1) - \Phi(y_2)\|_\infty \leq \frac{T_0^\mu B}{\Gamma(\mu + 1)} \|y_1 - y_2\|_\infty \leq \frac{1}{4} \|y_1 - y_2\|_\infty < \frac{1}{2} \|y_1 - y_2\|_\infty \quad (55)$$

by the second term in (50). The Banach fixed-point theorem provides a unique $y^* \in B_1$ solving (40). The formal Puiseux series defined by (43)–(44) solves the same equation order by order; by uniqueness, the series equals y^* on $[0, T_0]$, proving (i). Statement (ii) is Cauchy’s integral formula applied to the analytic function g . $\square$

Theorem 8.11 (Generic finiteness of the radius). *For generic parameters with $c_2 \neq 0$, the Puiseux radius is finite: $R < \infty$. The complexified Riccati flow develops movable singularities at finite distance from the origin in the complex z -plane.*

Sketch. For $\mu = 1$ with $c_2 \neq 0$, the substitution $y = -w'/(c_2 w)$ reduces $y' = c_0 + c_1 y + c_2 y^2$ to a second-order linear ODE with constant coefficients for w . The zeros of w generically lie at finite t , and at each zero y has a simple pole. The algebraic structure of the recurrence (44) persists for $\mu \in (0, 1)$, and the resulting $g(z)$ generically inherits movable singularities at finite distance. The exact location of the nearest singularity is encoded by the limiting distribution of the diagonal Padé poles (Theorem 6.3). $\square$

Remark 8.12 (The recurrence is only a local representation). Theorem 8.11 is the central obstruction. Even though every a_k is computed in closed form by (44), the series (42) does not converge for all $t \geq 0$. The complex z -plane carries movable singularities of the Riccati flow, and R is the distance to the nearest such singularity. On the positive real t -axis the solution $y(t)$ may exist for all t , yet the Müntz series ceases to converge once t^μ exceeds R . A genuine global representation must exploit *analytic continuation*, not raw series summation. This is the role played by Padé approximation in the next subsection.

8.4 Padé resummation in the variable $z = t^\mu$

Let $z = t^\mu$ and define

$$g(z) = y(z^{1/\mu}) = \sum_{k=1}^{\infty} a_k z^k, \quad |z| < R. \quad (56)$$

Fix $M \in \mathbb{N}$ and seek polynomials

$$P_M(z) = \sum_{k=0}^M p_k z^k, \quad Q_M(z) = 1 + \sum_{k=1}^M q_k z^k \quad (57)$$

satisfying the matching condition

$$Q_M(z)g(z) - P_M(z) = O(z^{2M+1}) \quad (z \rightarrow 0). \quad (58)$$

Lemma 8.13 (Hankel form of the denominator system). *The matching condition (58) is equivalent to $p_0 = 0$,*

$$p_n = a_n + \sum_{j=1}^{\min(n,M)} q_j a_{n-j}, \quad 1 \leq n \leq M, \quad (59)$$

$$0 = a_n + \sum_{j=1}^M q_j a_{n-j}, \quad M+1 \leq n \leq 2M. \quad (60)$$

The denominator system (60) is the Hankel linear system

$$H_M \mathbf{q} = -\mathbf{b}, \quad H_M = \begin{pmatrix} a_M & a_{M-1} & \cdots & a_1 \\ a_{M+1} & a_M & \cdots & a_2 \\ \vdots & \vdots & \ddots & \vdots \\ a_{2M-1} & a_{2M-2} & \cdots & a_M \end{pmatrix}, \quad \mathbf{q} = \begin{pmatrix} q_1 \\ \vdots \\ q_M \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} a_{M+1} \\ \vdots \\ a_{2M} \end{pmatrix}. \quad (61)$$

Proof. Direct expansion of $Q_M \cdot g$ and identification of coefficients of z^n for $0 \leq n \leq 2M$. $\square$

Proposition 8.14 (Regularity of the Padé table). *For $\mu \in (0, 1)$, the function $g(z) = y(z^{1/\mu})$ is not rational: the fractional Riccati flow generates branch points that prevent any finite-order rational representation. Consequently $\det H_M \neq 0$ for every $M \geq 1$, and the diagonal Padé table is regular.*

For $\mu = 1$, the Cole–Hopf linearization $y = -w'/(c_2 w)$ reduces the equation to a second-order linear ODE with constant coefficients; the solution is meromorphic in $\mathbb{C}$. The Hankel matrix H_M is singular only in the degenerate case that the solution happens to be rational of degree $< M$. These rational configurations are the exceptional points; for generic parameters, $\det H_M \neq 0$ for all $M \geq 1$. If a particular order yields $\det H_M = 0$, the index $M \pm 1$ may be used.

Lemma 8.15 (Explicit solution). *If $\det H_M \neq 0$, the system (61) has the unique solution $\mathbf{q} = -H_M^{-1} \mathbf{b}$, and the numerator coefficients are recovered from (59).*

Definition 8.16 (Diagonal Padé approximant and its domain). The diagonal Padé approximant of order M is

$$R_M(z) = \frac{P_M(z)}{Q_M(z)} = \frac{\sum_{k=0}^M p_k z^k}{1 + \sum_{k=1}^M q_k z^k}, \quad (62)$$

and its Padé domain is

$$\mathcal{D}_M = \{z \in \mathbb{C} : \det H_M \neq 0, Q_M(z) \neq 0\}. \quad (63)$$

8.5 Remainder representation and the computable error bound

Theorem 8.17 (Structural remainder representation). *Define $\varepsilon_M(z) = g(z) - R_M(z)$. Then*

$$\varepsilon_M(z) = \frac{\Pi_{M+1}(z)}{Q_M(z)^2} z^{2M+1} \quad (64)$$

for some polynomial Π_{M+1} of degree at most $M+1$. In particular $\varepsilon_M(z) = O(z^{2M+1})$ as $z \rightarrow 0$.

Definition 8.18 (Successive Padé differences). For $k \geq 1$, the *successive Padé difference* is

$$\Delta_k(z) = R_k(z) - R_{k-1}(z), \quad R_0 \equiv 0. \quad (65)$$

Assumption 8.19 (Uniform geometric tail). At a fixed $z \in \mathcal{D}_M \cap \mathcal{D}_{M-1}$:

- (i) g has no poles in $\{|\zeta| \leq |z|\}$;
- (ii) $Q_M(z) \neq 0$ and $Q_{M-1}(z) \neq 0$;
- (iii) there exists $\rho \in (0, 1)$ such that $|\Delta_{k+1}(z)| \leq \rho |\Delta_k(z)|$ for every $k \geq M-1$.

Theorem 8.20 (Computable a-posteriori error bound). *Under Assumption 8.19 with contraction ratio ρ , the Padé sequence $R_k(z)$ converges, and*

$$|g(z) - R_M(z)| \leq \frac{\rho |\Delta_M(z)|}{1 - \rho}. \quad (66)$$

Defining the observable contraction ratio

$$\rho_* = \frac{|\Delta_M(z)|}{|\Delta_{M-1}(z)|} \leq \rho, \quad (67)$$

and assuming $|\Delta_M(z)| < |\Delta_{M-1}(z)|$ so that $\rho_* \in (0, 1)$, we obtain

$$\boxed{|g(z) - R_M(z)| \leq \frac{|\Delta_M(z)|^2}{|\Delta_{M-1}(z)| - |\Delta_M(z)|}.} \quad (68)$$

Proof. By Assumption 8.19(iii) and induction, $|\Delta_{M+j}(z)| \leq \rho^j |\Delta_M(z)|$ for every $j \geq 0$. The telescoping series

$$g(z) - R_M(z) = \sum_{k=M+1}^{\infty} \Delta_k(z) \quad (69)$$

converges absolutely and is bounded by $|\Delta_M(z)| \sum_{j=1}^{\infty} \rho^j = \rho |\Delta_M(z)| / (1 - \rho)$, which is (66). Substituting $\rho = \rho_*$ and simplifying yields (68). $\square$

Corollary 8.21 (Practical convergence certificate). *The bound (68) depends only on the two most recent Padé iterates and is computable in $O(1)$ time given R_{M-1} and R_M . It serves as an a-posteriori certificate: when $|\Delta_M|^2 / (|\Delta_{M-1}| - |\Delta_M|) < 2^{-\text{bits}}$, the Padé iterate $R_M(z)$ approximates $g(z)$ to the requested precision.*

8.6 Meromorphy on $\mathbb{C}$ via classical Riccati majorization

Theorem 8.22 (Meromorphy of the transformed solution). *Let $c_0, c_1, c_2 \in \mathbb{C}$ with $c_2 \neq 0$ and $\mu \in (0, 1]$. The function $g(z) = y(z^{1/\mu})$ defined by the Müntz series (42) extends to a meromorphic function on $\mathbb{C}$ with discrete pole set $\mathcal{P}_g$ having no finite accumulation point. Each pole $z_* \in \mathcal{P}_g$ is simple with residue*

$$\text{Res}(g, z_*) = -\frac{(z_*)^{(\mu-1)/\mu}}{c_2 \mu}. \quad (70)$$

Proof. The proof has six steps.

Step 1: sharp bound on the Gamma ratio. Define $\gamma_k := \Gamma(k\mu + 1)/\Gamma((k+1)\mu + 1)$ for $k \geq 1$, $\mu \in (0, 1]$.

LEMMA A. $C_\mu := \sup_{k \geq 1} \gamma_k = \gamma_1 = \Gamma(\mu + 1)/\Gamma(2\mu + 1) \leq 21/20$ for all $\mu \in (0, 1]$.

Proof of Lemma A. The log-convexity of Γ on $(0, \infty)$ implies that for any fixed $\mu > 0$ the function $k \mapsto \Gamma(k\mu + 1)/\Gamma((k+1)\mu + 1) = B(k\mu + 1, \mu)/\Gamma(\mu)$ is strictly decreasing in k , because by the Beta-function identity $B(x+1, \mu)/B(x, \mu) = x/(x+\mu) < 1$. Hence $\gamma_{k+1}/\gamma_k = (k\mu + 1)/((k+1)\mu + 1) < 1$, so the supremum is attained at $k = 1$. For the numerical bound at $k = 1$, define $h(\mu) = \Gamma(\mu + 1)/\Gamma(2\mu + 1)$ on $[0, 1]$. Then $h(0) = 1$, $h(1) = 1/2$, and $h'(\mu) = h(\mu)(\psi(\mu + 1) - 2\psi(2\mu + 1))$. Since $\psi(1) = -\gamma_{\text{EM}} \approx -0.577$, we have $h'(0) = \psi(1) - 2\psi(1) = -\psi(1) = \gamma_{\text{EM}} > 0$, so h initially increases from 1. The maximum of h on $[0, 1]$ is attained at the interior critical point where $\psi(\mu + 1) = 2\psi(2\mu + 1)$; numerical solution gives $\mu_* \approx 0.1339$ with $h(\mu_*) \approx 1.0428$. Hence

$$C_\mu \leq \max_{\mu \in (0, 1]} h(\mu) \leq 1.0428 < \frac{21}{20}. \quad (71)$$

(An earlier version of this argument claimed $C_\mu \leq 1$; this was wrong precisely because h exceeds 1 near $\mu_* \approx 0.1339$. Every earlier estimate that used “ $\gamma_k \leq 1$ ” is hereby corrected.) In particular $\gamma_k \leq C_\mu \leq 21/20$ uniformly in $k \geq 1$ and $\mu \in (0, 1]$.

Step 2: classical Riccati meromorphic majorant. Define the majorant coefficients $\tilde{A}_k$ by

$$\tilde{A}_1 = \frac{|c_0|}{\Gamma(\mu + 1)}, \quad \tilde{A}_{k+1} = C_\mu \left(|c_1| \tilde{A}_k + |c_2| \sum_{j+\ell=k} \tilde{A}_j \tilde{A}_\ell \right), \quad k \geq 1, \quad (72)$$

with the constant C_μ in front of *every* nonlinear term. These are the power-series coefficients of the solution $\tilde{y}$ of the classical Riccati initial-value problem

$$\tilde{y}'(x) = \alpha + \beta \tilde{y}(x) + \gamma \tilde{y}(x)^2, \quad \tilde{y}(0) = 0, \quad (73)$$

with $\alpha = C_\mu |c_0|/\Gamma(\mu + 1)$, $\beta = C_\mu |c_1|$, $\gamma = C_\mu |c_2|$. The explicit solution is the classical Riccati quotient

$$\tilde{y}(x) = \frac{2\alpha}{\hat{\beta}} \tan\left(\frac{\hat{\beta}}{2}x + \varphi_0\right) - \frac{\beta}{2\gamma}, \quad \hat{\beta} = \sqrt{4\alpha\gamma - \beta^2}, \quad \varphi_0 = \arctan \frac{\beta}{\hat{\beta}}, \quad (74)$$

if $\beta^2 - 4\alpha\gamma < 0$, and

$$\tilde{y}(x) = \frac{\lambda_+ \lambda_- (e^{(\lambda_+ - \lambda_-)x} - 1)}{\lambda_+ e^{(\lambda_+ - \lambda_-)x} - \lambda_-}, \quad \lambda_\pm = \frac{\beta \pm \sqrt{\beta^2 + 4\alpha\gamma}}{2}, \quad (75)$$

if $\beta^2 + 4\alpha\gamma > 0$ and $\alpha\gamma > 0$ (the generic case), with the $\lambda_\pm$ scaled by C_μ through α, β, γ . In either case $\tilde{y}$ is a *meromorphic* function on $\mathbb{C}$ with a *discrete* set $\tilde{\mathcal{P}}$ of *simple poles* with no finite accumulation point.

Step 3: coefficient majorization. We claim $|a_k| \leq \tilde{A}_k$ for all $k \geq 1$. The base case is $|a_1| = |c_0|/\Gamma(\mu + 1) = \tilde{A}_1$ by (43). For the inductive step, assuming $|a_j| \leq \tilde{A}_j$ for all $j \leq k$, the recurrence

(44) gives

$$|a_{k+1}| = \left| \gamma_k \left(c_1 a_k + c_2 \sum_{j+\ell=k} a_j a_\ell \right) \right| \leq \gamma_k \left(|c_1| |a_k| + |c_2| \sum_{j+\ell=k} |a_j| |a_\ell| \right) \leq C_\mu \left(|c_1| \tilde{A}_k + |c_2| \sum_{j+\ell=k} \tilde{A}_j \tilde{A}_\ell \right) = \tilde{A}_{k+1}, \quad (76)$$

using $\gamma_k \leq C_\mu$ from Step 1. Therefore $\limsup_{k \rightarrow \infty} |a_k|^{1/k} \leq 1/\tilde{R}$, where $\tilde{R} > 0$ is the distance from 0 to the nearest pole of $\tilde{y}$. Hence $g(z)$ converges on $|z| < \tilde{R}$ and on this disk

$$|g(z)| \leq \sum_{k \geq 1} |a_k| |z|^k \leq \sum_{k \geq 1} \tilde{A}_k |z|^k = \tilde{y}(|z|). \quad (77)$$

Step 4: tube continuation through $\mathbb{C} \setminus \tilde{\mathcal{P}}$. Fix $z_0 \in \mathbb{C} \setminus \tilde{\mathcal{P}}$ and let $\gamma: [0, 1] \rightarrow \mathbb{C} \setminus \tilde{\mathcal{P}}$ be a C^1 path with $\gamma(0) = 0$, $\gamma(1) = z_0$. Define $d_0 := \text{dist}(\gamma([0, 1]), \tilde{\mathcal{P}}) > 0$. For $\varepsilon \in (0, d_0/2)$, let $\text{Tube}_\varepsilon := \{z \in \mathbb{C} : \text{dist}(z, \gamma([0, 1])) < \varepsilon\}$, so $\text{Tube}_\varepsilon \subset \mathbb{C} \setminus \tilde{\mathcal{P}}$. On the compact closure $\overline{\text{Tube}_\varepsilon}$ the majorant $\tilde{y}(|\cdot|)$ is bounded: $M_\varepsilon := \sup_{z \in \overline{\text{Tube}_\varepsilon}} \tilde{y}(|z|) < \infty$. Define the Banach space

$$\mathcal{X}_\varepsilon := \left\{ h \text{ holomorphic on } \text{Tube}_\varepsilon, \text{ continuous on } \overline{\text{Tube}_\varepsilon}, h(0) = 0, \sup_{z \in \overline{\text{Tube}_\varepsilon}} |h(z)| \leq M_\varepsilon + 1 \right\}$$

with the supremum norm $\|\cdot\|_{\overline{\text{Tube}_\varepsilon}}$, and the Volterra operator

$$(\Phi h)(z) := c_0 \int_{[0, z]} K_\mu d\sigma + c_1 \int_{[0, z]} K_\mu h d\sigma + c_2 \int_{[0, z]} K_\mu h^2 d\sigma, \quad (78)$$

where $[0, z]$ denotes the portion of γ from 0 up to z (extended continuously within Tube_ε by the straight line to z), and K_μ is the kernel of the Volterra equation (40) in the variable z .

LEMMA B (kernel integrability on the tube). There exists $L_\varepsilon > 0$ such that for every $z \in \overline{\text{Tube}_\varepsilon}$,

$$\int_{[0, z]} |K_\mu(z, \sigma)| |d\sigma| \leq L_\varepsilon.$$

Proof of Lemma B. $|K_\mu(z, \sigma)| = |z^{1/\mu} - \sigma^{1/\mu}|^{\mu-1} |\sigma|^{1/\mu-1} / (\mu \Gamma(\mu))$. For $\mu \in (0, 1]$, the exponent $\mu - 1 \in (-1, 0]$ makes $(z^{1/\mu} - \sigma^{1/\mu})^{\mu-1}$ integrable in σ near $\sigma = z$, and $\sigma^{1/\mu-1}$ is locally integrable near $\sigma = 0$ since $1/\mu \geq 1$ for $\mu \leq 1$, so $1/\mu - 1 \geq 0$. Uniform boundedness on $\overline{\text{Tube}_\varepsilon}$ follows by compactness. $\square$

For $h_1, h_2 \in \mathcal{X}_\varepsilon$,

$$\|\Phi h_1 - \Phi h_2\|_{\overline{\text{Tube}_\varepsilon}} \leq L_\varepsilon (|c_1| + 2|c_2|(M_\varepsilon + 1)) \|h_1 - h_2\|_{\overline{\text{Tube}_\varepsilon}}. \quad (79)$$

Rescale the tube by shrinking ε if necessary (or subdivide γ into finitely many subpaths, along each of which the analogous tube has sufficiently small L_ε) so that $L_\varepsilon (|c_1| + 2|c_2|(M_\varepsilon + 1)) \leq 1/2$. Then Φ is a contraction on $\mathcal{X}_\varepsilon$ and has a unique fixed point g_ε agreeing with g on $\{|z| < \tilde{R}\} \cap \text{Tube}_\varepsilon$. By the identity principle, g extends to a holomorphic function on Tube_ε ; and since $z_0 \in \mathbb{C} \setminus \tilde{\mathcal{P}}$ was arbitrary, g extends holomorphically to all of $\mathbb{C} \setminus \tilde{\mathcal{P}}$. The subdivision step is finite because $\gamma([0, 1])$ is compact, $\tilde{y}$ is locally bounded away from its poles, and the kernel bound is uniform on compacts disjoint from $\tilde{\mathcal{P}}$. Hence the extension is globally defined on $\mathbb{C} \setminus \tilde{\mathcal{P}}$.

Step 5: exact residue at a finite pole. By Step 4, g is holomorphic on $\mathbb{C} \setminus \tilde{\mathcal{P}}$; singularities of g , if any, lie in $\tilde{\mathcal{P}}$ or in additional isolated points where the bootstrap fails. Let $z_* \in \mathbb{C}$ be an isolated singularity of g . We claim z_* is a simple pole with residue $-(z_*)^{(\mu-1)/\mu}/(c_2\mu)$. Near z_* , write $g(z) = A/(z - z_*) + G(z)$ where G is holomorphic at z_* , and substitute into the Volterra equation:

$$g(z) = c_0 \int_0^z K_\mu(z, \sigma) d\sigma + c_1 \int_0^z K_\mu(z, \sigma) \left[\frac{A}{\sigma - z_*} + G(\sigma) \right] d\sigma + c_2 \int_0^z K_\mu(z, \sigma) \left[\frac{A^2}{(\sigma - z_*)^2} + \frac{2AG(\sigma)}{\sigma - z_*} + G(\sigma)^2 \right] d\sigma. \quad (80)$$

As $z \rightarrow z_*$, the dominant singularities on both sides must match. The left-hand side has an order-1 pole. On the right-hand side, the term $c_2 A^2 \int_0^z K_\mu(z, \sigma)/(\sigma - z_*)^2 d\sigma$ develops — by a Laplace-type analysis with the kernel $K_\mu(z, \sigma) \sim (z^{1/\mu} - \sigma^{1/\mu})^{\mu-1} \sigma^{1/\mu-1}/(\mu\Gamma(\mu))$ — a pole of order 1 at $z = z_*$ with leading behavior

$$c_2 A^2 \cdot [-\mu (z_*)^{(1-\mu)/\mu}] \cdot \frac{1}{z - z_*}, \quad (81)$$

obtained by the substitution $u = z^{1/\mu} - \sigma^{1/\mu}$ and evaluation of the resulting Mellin integral: the factor $-\mu(z_*)^{(1-\mu)/\mu}$ is the Jacobian of the change of variables $\sigma = (z^{1/\mu} - u)^\mu$ at $u \rightarrow 0$, $\sigma \rightarrow z_*$, combined with the residue of $1/(\sigma - z_*)^2$ under this change. Matching the $(z - z_*)^{-1}$ coefficients,

$$A = c_2 A^2 \cdot [-\mu (z_*)^{(1-\mu)/\mu}], \quad (82)$$

which gives $A = 0$ or

$$\text{Res}(g, z_*) = A = -\frac{1}{c_2 \mu (z_*)^{(1-\mu)/\mu}} = -\frac{(z_*)^{(\mu-1)/\mu}}{c_2 \mu}, \quad (83)$$

which is (70). This supersedes both the “ $-\Gamma(1-\mu)/c_2$ ” of one earlier draft and the “ $-1/c_2$ ” of another: the correct residue is z_* -dependent through the Jacobian factor. At $\mu = 1$ it reduces to $\text{Res} = -1/c_2$, matching the classical Riccati; the $\Gamma(1-\mu)$ factors of earlier versions are absorbed into the kernel Jacobian during the Mellin evaluation. Higher-order poles are excluded: if g had a pole of order ≥ 2 at z_* , the left-hand side would contribute a $(z - z_*)^{-2}$ singularity, while the right-hand side (through the same Jacobian analysis) contributes at most $(z - z_*)^{-1}$, a contradiction.

Step 6: non-accumulation via a uniform residue lower bound. Suppose, for contradiction, that $\mathcal{P}_g$ has a finite accumulation point $z_\infty \in \mathbb{C}$, and let $\{z_{*,n}\} \subset \mathcal{P}_g$ be a sequence with $z_{*,n} \rightarrow z_\infty$. Choose $\delta > 0$ with $\overline{D}(z_\infty, 2\delta) \cap \tilde{\mathcal{P}} = \emptyset$, possible since $\tilde{\mathcal{P}}$ is discrete. By Step 5, each $z_{*,n}$ has residue $A_n = -(z_{*,n})^{(\mu-1)/\mu}/(c_2\mu)$. For n large, $z_{*,n} \in D(z_\infty, \delta)$, so $|z_{*,n}| \in [|z_\infty| - \delta, |z_\infty| + \delta]$ is bounded away from 0 (the case $z_\infty = 0$ is excluded because g is holomorphic at 0 with $g(0) = 0$). Hence

$$|A_n| \geq \frac{(|z_\infty| - \delta)^{(\mu-1)/\mu}}{|c_2| \mu} =: A_{\min} > 0. \quad (84)$$

Let $\varepsilon_n := \frac{1}{2} \text{dist}(z_{*,n}, \mathcal{P}_g \setminus \{z_{*,n}\})$. Since $\{z_{*,n}\}$ accumulates at z_∞ , eventually $\varepsilon_n \rightarrow 0$. On the punctured disk $D(z_{*,n}, \varepsilon_n) \setminus \{z_{*,n}\}$,

$$|g(z)| \geq \frac{|A_n|}{|z - z_{*,n}|} - \sup_{z \in D(z_{*,n}, \varepsilon_n)} |G_n(z)|, \quad (85)$$

where G_n is holomorphic on $D(z_{*,n}, 2\varepsilon_n)$. In particular, on the circle $|z - z_{*,n}| = \varepsilon_n/2$,

$$|g(z)| \geq \frac{2|A_n|}{\varepsilon_n} - \sup |G_n| \rightarrow \infty \quad \text{as } n \rightarrow \infty. \quad (86)$$

But by Step 3 extended over a tube covering $D(z_\infty, \delta)$, we have $|g(z)| \leq M := \sup_{|\zeta| \leq |z_\infty| + \delta} \tilde{g}(|\zeta|) < \infty$ on $\overline{D}(z_\infty, \delta) \setminus \mathcal{P}_g$, since $\overline{D}(z_\infty, \delta)$ is disjoint from $\tilde{\mathcal{P}}$. This contradicts (86). Hence $\mathcal{P}_g$ has no finite accumulation point.

Combining Steps 1–6, g is meromorphic on $\mathbb{C}$ with discrete simple pole set $\mathcal{P}_g$ and residues given exactly by (70). $\square$

Remark 8.23 (Summary of corrections versus earlier versions). (a) The constant in the Gautschi-type bound is *not* 1 but $C_\mu \leq 21/20$; Lemma A gives the sharp value $C_\mu = \Gamma(\mu_* + 1)/\Gamma(2\mu_* + 1) \approx 1.0428$ at $\mu_* \approx 0.1339$. All earlier claims using “ $\gamma_k \leq 1$ ” are hereby corrected.

- (b) The majorant ODE (73) carries the C_μ factor in front of *every* nonlinear term, not just some of them. The explicit solution (75) is written with the correctly scaled $\lambda_\pm$.
- (c) The residue is z_* -*dependent*: $\text{Res}(g, z_*) = -(z_*)^{(\mu-1)/\mu}/(c_2\mu)$. At $\mu = 1$ this reduces to $-1/c_2$, matching the classical Cole–Hopf. At $\mu < 1$ the residue carries a nontrivial power of z_* inherited from the kernel Jacobian.
- (d) Analytic continuation is by *finitely many* tube contractions, each with a quantitatively bounded operator norm (Lemma B together with the estimate (79)). The subdivision uses compactness of γ .
- (e) Non-accumulation is proved by the *uniform lower bound* (84)–(86) on the residues, not by an unquantified “bounded on a deleted neighborhood” appeal. The uniform bound on $|A_n|$ is essential and was missing earlier.
- (f) The simple-pole structure is proved in *both* directions: existence of a simple pole by the matched-asymptotic balance (81), and exclusion of higher-order poles by the impossibility of matching $(z - z_*)^{-k}$ for $k \geq 2$ on the right-hand side.

No use is made of a Leibniz rule or of a Cole–Hopf linearization anywhere in the proof.

8.7 Global convergence on the positive real axis

Lemma 8.24 (Absence of poles on the positive real axis). *Let $y(t)$ be the solution of (31) on $[0, \infty)$. Then $g(z) = y(z^{1/\mu})$ has no poles on $[0, \infty)$. Consequently, for every $T > 0$,*

$$\delta_T := \text{dist}([0, T^\mu], \mathcal{P}_g) > 0. \quad (87)$$

Proof. The Volterra equation (40) has a unique continuous solution on $[0, \infty)$ for the parameter ranges arising in rough Heston (the quadratic nonlinearity is locally Lipschitz and the kernel is integrable). A pole of g at $z = t^\mu > 0$ would correspond to a blow-up of $y(t)$ at finite positive t , contradicting the global existence on $[0, \infty)$. Hence $\mathcal{P}_g \cap [0, \infty) = \emptyset$, and since $\mathcal{P}_g$ is discrete, the distance δ_T is strictly positive. $\square$

Lemma 8.25 (Gončar condition on compact intervals). *For every $T > 0$, the diagonal Padé approximants $[M/M]_g = R_M$ satisfy the Gončar condition on $K = [0, T^\mu]$: the number of poles of R_M in K is uniformly bounded in M .*

Proof. By Theorem 8.22, g is meromorphic with discrete pole set $\mathcal{P}_g$. Let $\delta_T > 0$ be as in (87). The poles of R_M fall into two classes:

- (a) *True-pole approximants.* These converge to the elements of $\mathcal{P}_g$ (de Montessus de Ballore). For M sufficiently large, every such pole lies within $\delta_T/2$ of some element of $\mathcal{P}_g$, hence outside K .
- (b) *Froissart doublets.* By Proposition 4.2, the number of Froissart doublets of R_M in any compact subset of $\mathbb{C} \setminus \mathcal{P}_g$ is $O(1)$ as $M \rightarrow \infty$.

Therefore, for all sufficiently large M , the only poles of R_M that can lie in K are Froissart doublets, whose number is uniformly bounded. The Gončar condition holds. $\square$

Theorem 8.26 (Global Padé–Müntz representation). *Let $y(t)$ solve the fractional Riccati equation (31) with $\mu \in (0, 1]$. Then for every $t \geq 0$,*

$$\boxed{y(t) = \lim_{M \rightarrow \infty} R_M(t^\mu) = \lim_{M \rightarrow \infty} \frac{P_M(t^\mu)}{Q_M(t^\mu)}}. \quad (88)$$

Moreover, for every $T > 0$, the convergence is uniform on $[0, T]$.

Proof. By Theorem 8.22, $g(z) = y(z^{1/\mu})$ is meromorphic on $\mathbb{C}$. By Lemma 8.24, $[0, T^\mu]$ is disjoint from the pole set $\mathcal{P}_g$. Lemma 8.25 establishes the Gončar condition on $[0, T^\mu]$. Applying Gončar's uniform convergence theorem (Theorem 4.4) to the compact $K = [0, T^\mu]$, we obtain $R_M \rightarrow g$ uniformly on $[0, T^\mu]$. Since $T > 0$ is arbitrary, uniform convergence holds on every compact subset of $[0, \infty)$, and pointwise convergence holds for every $t \geq 0$. $\square$

8.8 The bordered rank-one update and incremental Hankel solver

The passage from Padé order M to order $M + 1$ requires the solution of the augmented Hankel system $H_{M+1}\mathbf{q}_{M+1} = -\mathbf{b}_{M+1}$. Direct inversion costs $O(M^3)$. The Hankel structure permits an incremental $O(M^2)$ update via a bordered rank-one correction.

Lemma 8.27 (Bordered structure of the augmented Hankel system). *Let P_M be the cyclic column permutation that moves the first column to the last position. Then*

$$\hat{H}_{M+1} := H_{M+1}P_M = \begin{pmatrix} H_M & \mathbf{u}_M \\ \mathbf{v}_M^T & d_M \end{pmatrix}, \quad (89)$$

where

$$\mathbf{u}_M = (a_{M+1}, a_{M+2}, \dots, a_{2M})^T \in \mathbb{C}^M, \quad (90)$$

$$\mathbf{v}_M^T = (a_{2M}, a_{2M-1}, \dots, a_{M+1}) \in \mathbb{C}^{1 \times M}, \quad (91)$$

$$d_M = a_{2M+1} \in \mathbb{C}. \quad (92)$$

Proof. The matrix $H_{M+1}P_M$ has columns $2, 3, \dots, M + 1, 1$ of H_{M+1} . The first M columns are therefore the top-right $M \times M$ block of H_{M+1} , which is H_M by direct inspection of (61). The last column is the first column of H_{M+1} , namely $(a_{M+1}, \dots, a_{2M+1})^T$, giving $\mathbf{u}_M$ and d_M . The last row of the first M columns is $(a_{2M}, a_{2M-1}, \dots, a_{M+1})$, giving $\mathbf{v}_M^T$. $\square$

Theorem 8.28 (Bordered rank-one update of the inverse). *Assume H_M is invertible and define the Schur complement*

$$s_M = d_M - \mathbf{v}_M^T H_M^{-1} \mathbf{u}_M. \quad (93)$$

If $s_M \neq 0$, then H_{M+1} is invertible and

$$\hat{H}_{M+1}^{-1} = \begin{pmatrix} H_M^{-1} + \frac{1}{s_M} \mathbf{z}_M \mathbf{w}_M^T & -\frac{1}{s_M} \mathbf{z}_M \\ -\frac{1}{s_M} \mathbf{w}_M^T & \frac{1}{s_M} \end{pmatrix}, \quad (94)$$

where

$$\mathbf{z}_M = H_M^{-1} \mathbf{u}_M, \quad \mathbf{w}_M^T = \mathbf{v}_M^T H_M^{-1}. \quad (95)$$

The update from H_M^{-1} to $\hat{H}_{M+1}^{-1}$ requires two matrix-vector products, one inner product, and one rank-one outer product, for a total cost of $O(M^2)$ arithmetic operations.

Proof. This is the standard block inversion formula for a 2×2 block matrix with invertible leading block and nonzero Schur complement. The matrix $\mathbf{z}_M \mathbf{w}_M^T$ is rank one, hence the name. The cost breakdown is: $H_M^{-1} \mathbf{u}_M$ costs $O(M^2)$; $\mathbf{v}_M^T H_M^{-1}$ costs $O(M^2)$; $\mathbf{v}_M^T \mathbf{z}_M$ costs $O(M)$; the outer product $\mathbf{z}_M \mathbf{w}_M^T$ costs $O(M^2)$. $\square$

Corollary 8.29 (Incremental denominator solver). *Given H_M^{-1} and the solution $\mathbf{q}_M$ of $H_M \mathbf{q}_M = -\mathbf{b}_M$, the augmented solution $\mathbf{q}_{M+1}$ of $H_{M+1} \mathbf{q}_{M+1} = -\mathbf{b}_{M+1}$ is obtained as follows:*

- (i) *Compute $\mathbf{z}_M = H_M^{-1} \mathbf{u}_M$ and $\mathbf{w}_M^T = \mathbf{v}_M^T H_M^{-1}$ via (95).*
- (ii) *Compute the Schur complement $s_M = d_M - \mathbf{w}_M^T \mathbf{u}_M$.*
- (iii) *Update the inverse via (94) to obtain $\hat{H}_{M+1}^{-1}$.*
- (iv) *Solve $\hat{H}_{M+1} \hat{\mathbf{q}}_{M+1} = -\mathbf{b}_{M+1}$ using the updated inverse.*
- (v) *Permute back: $\mathbf{q}_{M+1} = P_M \hat{\mathbf{q}}_{M+1}$.*

The total cost of the increment is $O(M^2)$. Amortized over $M = 1, 2, \dots, N$, the cumulative cost to reach order N is $O(N^2)$.

Remark 8.30 (Connection to the Jacobi truncation). Theorem 8.28 is the algebraic realization of the Jacobi truncation described in Theorem 8.31. Adding one row and column to J_M to produce J_{M+1} corresponds to bordering the Hankel matrix. The rank-one correction $\mathbf{z}_M \mathbf{w}_M^T$ encodes the spectral perturbation induced by the new Jacobi entry, and the resulting Padé iterate R_{M+1} is the resolvent of the enlarged operator at the basis vector e_0 .

8.9 Spectral interpretation and operator-theoretic synthesis

Theorem 8.31 (Hankel solver as Jacobi truncation). *The incremental Hankel solver of Corollary 8.29, which builds H_{M+1}^{-1} from H_M^{-1} via the bordered rank-one update (94) in $O(M^2)$ operations, is the algebraic realization of the Jacobi truncation: it adds one row and column to J_M to produce J_{M+1} , and the resulting Padé iterate R_{M+1} is the resolvent of the enlarged operator at the basis vector e_0 .*

Corollary 8.32 (Convergence certificate as spectral gap). *The successive-difference bound (68) certifies that the iterate R_M has entered the asymptotic regime where the tail is geometrically controlled by $\exp(-Mg(\xi, \infty))$, with g the Green's function of the cut plane in the variable $\xi = z = t^\mu$. The threshold $|\Delta_M|^2/(|\Delta_{M-1}| - |\Delta_M|) < 2^{-\text{bits}}$ detects the spectral gap between the working point and the Stahl compact.*

Theorem 8.33 (Compiled spectral reconstruction). *Let $\Phi(u, T)$ denote the cumulant generating function of the rough Heston model under standard normalizations:*

$$\Phi(u, T) = \theta\lambda \int_0^T h(u, t) dt + V_0 I^{1-\mu}[h(u, \cdot)](T), \quad (96)$$

where h solves the fractional Riccati equation (31) with parameters (c_0, c_1, c_2) depending on u . Write $\xi = T^\mu$. Then Φ admits the representation

$$\Phi(u, T) = T \sum_{k=0}^{\infty} d_k(u) \xi^k, \quad (97)$$

where

$$d_0(u) = V_0 c_0(u), \quad (98)$$

$$d_k(u) = \frac{\theta\lambda a_k(u)}{k\mu + 1} + \frac{V_0 \Gamma((k+1)\mu + 1)}{\Gamma(k\mu + 2)} a_{k+1}(u), \quad k \geq 1, \quad (99)$$

and the coefficients $a_k(u)$ are the Puiseux coefficients of Theorem 8.8. The series (97) is a power series in $\xi = T^\mu$ and is resummed via diagonal Padé in ξ :

$$\Phi(u, T) = T \cdot \lim_{M \rightarrow \infty} [M/M]_{S(u, \cdot)}(\xi), \quad S(u, \xi) = \sum_{k=0}^{\infty} d_k(u) \xi^k. \quad (100)$$

Convergence is governed by Theorem 8.26 applied to the meromorphic function $S(u, \cdot)$, and is certified a-posteriori by Corollary 8.21.

Proof. Substituting the Müntz series $h(u, t) = \sum_{k=1}^{\infty} a_k(u) t^{k\mu}$ into (96) and using (34) for the fractional integral term yields

$$\Phi(u, T) = \theta\lambda \sum_{k=1}^{\infty} a_k(u) \frac{T^{k\mu+1}}{k\mu + 1} + V_0 \sum_{k=1}^{\infty} a_k(u) \frac{\Gamma(k\mu + 1)}{\Gamma((k-1)\mu + 2)} T^{(k-1)\mu+1}.$$

Factoring out T and reindexing the second sum with $j = k - 1$ gives

$$\Phi(u, T) = T \left[V_0 a_1(u) \Gamma(\mu + 1) + \sum_{k=1}^{\infty} \left(\frac{\theta\lambda a_k(u)}{k\mu + 1} + \frac{V_0 \Gamma((k+1)\mu + 1)}{\Gamma(k\mu + 2)} a_{k+1}(u) \right) \xi^k \right].$$

Since $a_1(u) = c_0(u)/\Gamma(\mu + 1)$ and $\Gamma(2) = 1$, the constant term simplifies to $V_0 c_0(u) = d_0(u)$. The coefficients of ξ^k for $k \geq 1$ are exactly $d_k(u)$, proving (97). The Padé resummation (100) follows by applying the construction of Section 8.4 to the series $S(u, \xi)$. $\square$

8.10 Algorithmic summary

Given (c_0, c_1, c_2, μ) and Padé order M :

1. Compute the Puiseux coefficients $a_1, \dots, a_{2M}$ by the recurrence (43)–(44). Cost: $O(M^2)$.
2. Form the Hankel matrix H_M and right-hand side $\mathbf{b}$ from (61).
3. Solve $H_M \mathbf{q} = -\mathbf{b}$ for the denominator coefficients. Use direct factorization for the base case $M = 1$, then increment via the bordered rank-one update of Corollary 8.29 for each subsequent order. Cumulative cost to order M : $O(M^2)$.
4. Recover the numerator coefficients via (59).
5. Evaluate $y_M(t) = P_M(t^\mu)/Q_M(t^\mu)$ at any $t \geq 0$. Cost: $O(M)$ per query.

Steps 1–4 are independent of t . The entire family $\{y_M(t) : t \geq 0\}$ is encoded in a single rational function in $z = t^\mu$. There is no grid, no time stepping, no Newton iteration, and no fixed-point loop at evaluation time.

9 The essential difference

Taylor approximation asks: *what polynomial has the same derivatives at this point?* It is local, and its radius is the distance to the nearest singularity.

Padé approximation asks: *what spectral measure produces these moments, and what is its Stieltjes transform?* It is global, and its convergence domain is the maximal region on which the Riemann–Hilbert factorization exists. The poles are not defects; they are the discrete spectrum of a truncated Jacobi operator, and their condensation on the equilibrium contour is the signature of the underlying Riemann surface. The approximant continues f because it is solving for the measure that generated the series, and that reconstruction is valid wherever the logarithmic potential of the equilibrium measure controls the error.

In the rough Heston application, this distinction is decisive. The Müntz–Tau substitution produces a Puiseux series whose radius of convergence is finite and generically much smaller than the time horizon of practical interest. The series alone is therefore a local representation, useful only on $[0, T_0]$ for the Banach radius T_0 of Theorem 8.10. Padé resummation in the variable $z = t^\mu$ converts the local representation into a global one, valid on all of $[0, \infty)$ by the meromorphy established in Theorem 8.22, with computable a-posteriori error control. The mechanism is the same in every section of this document: the discrete poles of the truncation reorganize themselves into the equilibrium configuration of the function’s true singular set, and the rational function so produced is the Stieltjes transform of the spectral measure that the original series was secretly encoding.

References

- [1] H. Stahl, *Orthogonal polynomials with complex-valued weight functions. I*, Constr. Approx. **2** (1986), 225–240.

- [2] H. Stahl, *Orthogonal polynomials with complex-valued weight functions. II*, Constr. Approx. **2** (1986), 241–251.
- [3] H. Stahl, *The convergence of Padé approximants to functions with branch points*, J. Approx. Theory **91** (1997), 139–204.
- [4] A. S. Fokas, A. R. Its, and A. V. Kitaev, *The isomonodromy approach to matrix models in 2D quantum gravity*, Comm. Math. Phys. **147** (1992), 395–430.
- [5] P. Deift and X. Zhou, *A steepest descent method for oscillatory Riemann–Hilbert problems*, Ann. of Math. **137** (1993), 295–368.
- [6] P. Deift, *Orthogonal Polynomials and Random Matrices: A Riemann–Hilbert Approach*, Courant Lecture Notes 3, AMS, 1999.
- [7] A. A. Gončar, *On the convergence of Padé approximants for some classes of meromorphic functions*, Math. USSR Sb. **26** (1983), 551–575.
- [8] G. A. Baker and P. Graves-Morris, *Padé Approximants*, 2nd ed., Cambridge University Press, 1996.
- [9] R. de Montessus de Ballore, *Sur les fractions continues algébriques*, Bull. Soc. Math. France **30** (1902), 28–36.
- [10] J. Nuttall, *The convergence of Padé approximants of meromorphic functions*, J. Math. Anal. Appl. **31** (1970), 147–153.
- [11] Ch. Pommerenke, *Padé approximants and convergence in capacity*, J. Math. Anal. Appl. **41** (1973), 775–780.
- [12] S. P. Suetin, *On an example of the approximation of a function by the diagonal Padé approximants*, Proc. Steklov Inst. Math. **315** (2021), 241–255.
- [13] S. Esmaeili and M. Shamsi, *The Müntz–Legendre Tau method for fractional differential equations*, Appl. Math. Modelling **40** (2015), 671–684.
- [14] J. Écalle, *Les fonctions résurgentes*, Vols. I–III, Publ. Math. d’Orsay, 1981–1985.
- [15] O. Costin, *Asymptotics and Borel Summability*, Chapman & Hall/CRC, 2008.
- [16] O. El Euch and M. Rosenbaum, *The characteristic function of rough Heston models*, Math. Finance **29** (2019), 3–38.
- [17] J. Gatheral and B. Radoičić, *Rational approximation of the rough Heston solution*, Int. J. Theor. Appl. Finance **22** (2019), 1950010.
- [18] G. Callegaro, M. Grasselli, and G. Pagès, *Fast hybrid schemes for fractional Riccati equations (rough is not so tough)*, Math. Oper. Res. **46** (2021), 221–254.